Spontaneous scalarization of Gauss-Bonnet black holes surrounded by massive scalar fields

Yan $Peng^{1*}$

¹ School of Mathematical Sciences, Qufu Normal University, Qufu, Shandong 273165, China

Abstract

For massless scalar fields, a relation $\Delta_n = \frac{\sqrt{3}}{2}\pi$ for $n \to \infty$ was observed in the scalar-Gauss-Bonnet theory. In the present paper, we extend the discussion by including a nonzero scalar field mass. For massive scalar fields, we show that the relation $\Delta_n = \frac{\sqrt{3}}{2}\pi$ for $n \to \infty$ still holds. We demonstrate this relation with both analytical and numerical methods. The analytical analysis implies that this relation may be a very universal behavior.

PACS numbers: 11.25.Tq, 04.70.Bw, 74.20.-z

^{*} yanpengphy@163.com

I. INTRODUCTION

The recent gravitational wave observation confirmed the existence of astrophysical black holes in nature [1–3]. Future more precise detection of gravitational waves may also can be used to examine various black holes theories. A well known property of black holes is the no hair theorem, which suggests that asymptotically flat black holes cannot support the formation of scalar field hairs, see references [4]-[11] and reviews [12, 13]. This no hair theorem is a very general property and it still holds even including the non-minimally coupling between scalar fields and the Ricci curvature [14–16].

Interestingly, it was recently found that the no hair theorem is violated when considering a non-minimally coupling between scalar fields and the Gauss-Bonnet invariant [17–20]. In particular, this scalar-Gauss-Bonnet coupling also provides a mechanism that a thermodynamically unstable pure black hole evolves into a thermodynamically more stable hairy black hole [18, 19]. This intriguing mechanism is usually called spontaneous scalarization, which was firstly found in the background of neutron stars [21]. As a further step, spontaneous scalarization of Kerr black holes was also observed in the scalar-Gauss-Bonnet theory [22]. In fact, spontaneous scalarization is a very general property and it also can be triggered by non-minimal couplings between scalar fields and Maxwell fields [23, 24]. At present, spontaneous scalarization phenomenons were observed in various completed gravity models [25–37]. In addition, analytical properties of black hole spontaneous scalarization were explored in [38–40]. In particular, for massless scalar fields, an interesting relation $\Delta_n = \sqrt{\eta_{n+1}} - \sqrt{\eta_n} = \frac{\sqrt{3}}{2}\pi$ was obtained through WKB approach in [38] and this relation is also precisely supported by numerical data in [19]. Whether this relation is a universal behavior is still a question to be answered. In this work, we plan to examine whether this relation holds for scalar fields with nonzero scalar field mass.

The rest of this paper is as follows. We start by introducing a model with a massive scalar field coupled to the Gauss-Bonnet invariant in the black hole spacetime. For massive scalar fields, we obtain a relation, which was firstly found in the case of massless scalar fields. We demonstrate this relation with both analytical and numerical methods. The conclusions are given in the last section.

II. INVESTIGATIONS ON THE COUPLING PARAMETER THROUGH THE WKB METHOD

We study the system with a real scalar field non-minimally coupled to the Einstein gravity. The model is described by the Lagrange density [17–20]

$$\mathcal{L} = R - \nabla^{\nu} \nabla_{\nu} \Psi - \mu^2 \Psi^2 + f(\Psi) \mathcal{R}_{GB}^2.$$
⁽¹⁾

Here R is the Ricci scalar of the metric. Ψ is the scalar field with mass μ . The term \mathcal{R}_{GB}^2 is the Gauss-Bonnet invariant expressed as $\mathcal{R}_{GB}^2 = R_{\alpha\beta\rho\sigma}R^{\alpha\beta\rho\sigma} - 4R_{\alpha\beta}R^{\alpha\beta} + R^2$. In the Schwarzschild black hole background, it is given by $\mathcal{R}_{GB}^2 = \frac{48M^2}{r^6}$ [38, 39, 41, 42]. The function $f(\Psi)$ describes the coupling between the scalar field and the Gauss-Bonnet invariant. In the linearized regime, without loss of generality, the coupling function can be taken in the form $f(\Psi) = \eta \Psi^2$, with η as the coupling parameter [38, 39, 41, 42].

The scalar field equation of motions is

$$\nabla^{\nu}\nabla_{\nu}\Psi - \mu^{2}\Psi + \frac{f'_{\Psi}\mathcal{R}^{2}_{GB}}{2} = 0.$$
⁽²⁾

The line element describing a spherically symmetric black hole solution is

$$ds^{2} = -g(r)dt^{2} + \frac{dr^{2}}{g(r)} + r^{2}(d\theta^{2} + \sin^{2}\theta d\phi^{2}).$$
(3)

The metric function of a scalar free Schwarzschild black hole is given by $g(r) = 1 - \frac{2M}{r}$ with M as the black hole mass. The black hole horizon $r_h = 2M$ is determined by the relation $g(r_h) = 0$.

According to relations (2) and (3), the scalar field equation can be expressed in the form [41, 42]

$$\Psi'' + \left(\frac{2}{r} + \frac{g'}{g}\right)\Psi' + \left(\frac{\eta \mathcal{R}_{GB}^2}{g} - \frac{\mu^2}{g}\right)\Psi = 0 \tag{4}$$

with $g = 1 - \frac{2M}{r}$ and $\mathcal{R}_{GB}^2 = \frac{48M^2}{r^6}$.

At the horizon $r_h = 2M$, we take finite boundary conditions [38]

$$\Psi(r_h) < \infty. \tag{5}$$

In the far region, the general asymptotical behavior of massive static scalar fields are $\Psi(r \to \infty) \sim \frac{A_1}{r}e^{-\mu r} + \frac{B_1}{r}e^{\mu r}$ with A_1 and B_1 as constants. For bounded scalar fields, we impose the relation $B_1 = 0$. So the infinity boundary condition is [39, 41, 43, 44]

$$\Psi(r \to \infty) \propto \frac{1}{r} e^{-\mu r}.$$
(6)

4

We introduce a new function $\psi = r\Psi$. Then there is $\psi(r \to \infty) \to 0$. We also take the tortoise radial coordinate y defined by the relation

$$\frac{dr}{dy} = g(r). \tag{7}$$

With relations (4), (7) and $\mathcal{R}_{GB}^2 = \frac{48M^2}{r^6}$, we derive the Schrödinger-like differential equation

$$\frac{d^2\psi}{dy^2} - V\psi = 0\tag{8}$$

with the effective potential expressed as

$$V(r) = (1 - \frac{2M}{r})(\frac{2M}{r^3} + \mu^2 - \frac{12\eta M^2}{r^6}).$$
(9)

The Schrödinger-like radial differential equation (8) can be studied through a standard WKB analysis. For a bounded scalar field, there is a well-known quantization condition [38]

$$\int_{y_{-}}^{y_{+}} dy \sqrt{-V(y,\eta)} = (n - \frac{1}{4})\pi; \quad n = 1, 2, 3, \dots,$$
(10)

where y_{\pm} are the classical turning points derived from $V(y_{\pm}) = 0$. And n is a positive integer.

With the relation (7), we can transfer the resonance condition (10) into

$$\int_{r_{-}}^{r_{+}} dr \frac{\sqrt{-V(r,\eta)}}{g(r)} = (n - \frac{1}{4})\pi; \quad n = 1, 2, 3, \dots,$$
(11)

where the two turning points r_{\pm} are determined by $V(r_{\pm}) = 0$ from equations

$$1 - \frac{2M}{r_{-}} = 0 \tag{12}$$

and

$$\frac{2M}{r_+^3} + \mu^2 - \frac{12\eta M^2}{r_+^6} = 0.$$
(13)

Then the turning points are

$$r_{-} = 2M, \qquad r_{+} = \left(\frac{12\eta}{1 + \sqrt{1 + 12\eta\mu^{2}}}\right)^{1/3}.$$
 (14)

Substituting (9) into equation (11), we obtain the relation

$$\int_{r_{-}}^{r_{+}} dr \sqrt{\frac{12\eta M^2 - \mu^2 r^6 - 2Mr^3}{r^6(1 - \frac{2M}{r})}} = (n - \frac{1}{4})\pi; \quad n = 1, 2, 3, \dots$$
(15)

For fixed values of η , μ and M, the integration on the left side of (15) is finite. From the relation (15), for fixed μ and M, there is $\eta \to \infty$ in the limit of $n \to \infty$. Also considering (14), it yields $r_+ \to \infty$ in the limit of $n \to \infty$. In the large n limit, the left side of (15) can be approximated by

$$\int_{r_{-}}^{r_{+}} dr \sqrt{\frac{12\eta M^2 - \mu^2 r^6 - 2Mr^3}{r^6(1 - \frac{2M}{r})}} = \int_{2M}^{\infty} dr \sqrt{\frac{12\eta M^2}{r^6(1 - \frac{2M}{r})}}.$$
(16)

According to (16), the relation (15) can be expressed as

$$\int_{2M}^{\infty} dr \sqrt{\frac{12\bar{\eta}M^4}{r^6(1-\frac{2M}{r})}} = (n-\frac{1}{4})\pi \quad for \quad n \to \infty,$$
(17)

where we define a new dimensionless parameter $\bar{\eta} = \frac{\eta}{M^2}$ using the symmetry $r \to \gamma r$, $\mu \to \mu/\gamma$, $M \to \gamma M$, $\eta \to \gamma^2 \eta$ of equation (4). According to the integration $\int_{2M}^{\infty} dr \sqrt{\frac{12M^4}{r^6(1-\frac{2M}{r})}} = \frac{2}{\sqrt{3}}$, one deduces the expression

$$\sqrt{\bar{\eta}_n} = \frac{\sqrt{3}}{2}(n-\frac{1}{4})\pi.$$
(18)

The discrete resonant spectrum (18) yields a relation

$$\Delta_n = \sqrt{\bar{\eta}_{n+1}} - \sqrt{\bar{\eta}_n} = \frac{\sqrt{3}}{2}\pi.$$
(19)

We mention that the relation (19) is the same as the case of massless scalar fields in [38]. Our analysis implies that this relation may hold for very general models with the property that $r_+ \to \infty$ as $n \to \infty$. Since the WKB approach is an approximate method, we still have to examine the relation with numerical data.

III. NUMERICAL EXAMINATION OF THE UNIVERSAL BEHAVIOR

In this part, we examine the relation (19) with numerical solutions by integrating the equation (4) from the black hole horizon to the infinity. Using the symmetry $\psi \to \kappa \psi$ of equation (4), we can take $\psi(r_h) = 1$ without loss of generality. From mathematical aspects, it is a boundary value problem. For fixed parameters μ^2 and M, we find discrete η , which leads to bounded scalar field solutions satisfying the boundary conditions $\psi(r_h) = 1$ and $\psi(\infty) = 0$.

With $\mu^2 = 0.002$ and M = 1, we choose different values of η to try to search for the proper η corresponding to a scalar field decaying at the infinity. We numerically obtain a discrete value $\eta \approx 3.110$, which leads to the solution with $\Psi(\infty) = 0$. We plot the physical solution with blue curves in the left panel of Fig. 1. With a larger discrete value $\eta \approx 20.263$, we get a solution with one node as shown in the right panel of Fig. 1. In fact, we can also get physical solutions with many nodes.

With numerical solutions, we can examine whether the relation (19) holds. We study effects of scalar field mass μ^2 and node number n on the discrete dimensionless coupling parameter $\bar{\eta}(\mu^2, n)$. With M = 1, we



FIG. 1: (Color online) We show the scalar field $\Psi(r)$ as a function of the coordinate r with $\mu^2 = 0.002$ and M = 1. The left panel corresponds to the case of $\eta = 3.110$ and the right panel is with $\eta = 20.263$.

choose $\mu^2 = 0.002$ in Table I and $\mu^2 = 0.004$ in Table II. The data shows that, for nonzero scalar field mass μ^2 , $\Delta_n = \sqrt{\bar{\eta}_{n+1}} - \sqrt{\bar{\eta}_n} \rightarrow 2.72$ for very large n. Since $\frac{\sqrt{3}}{2}\pi \approx 2.72 \times 1.0003$, we have numerically demonstrated the analytical relation (19). For larger mass μ^2 , we usually need a larger n to obtain the limit value $\bar{\eta} \approx 2.72$, as can be seen from Table I and Table II.

BLE I: The parameter η with $\mu = 0.002$ and M											
	n	1	2	3	$4\leqslant n\leqslant 28$	$n \geqslant 29$					
	$\bar{\eta}$	2.74	2.74	2.74	2.73	2.72					

TABLE I: The parameter $\bar{\eta}$ with $\mu^2 = 0.002$ and M = 1

TABLE II: The parameter $\bar{\eta}$ with $\mu^2 = 0.004$ and M = 1

n	1	2	$3\leqslant n\leqslant 5$	$6\leqslant n\leqslant 41$	$n \geqslant 42$
$\bar{\eta}$	2.75	2.75	2.74	2.73	2.72

IV. CONCLUSIONS

We investigated formations of scalar clouds outside asymptotically flat spherical black holes. We considered massive scalar fields non-minimally coupled to the Gauss-Bonnet invariant. For massive scalar fields, with WKB methods, we analytically obtained a relation $\Delta_n = \sqrt{\overline{\eta}_{n+1}} - \sqrt{\overline{\eta}_n} = \frac{\sqrt{3}}{2}\pi$ in the limit of $n \to \infty$. We numerically found $\Delta_n = \sqrt{\overline{\eta}_{n+1}} - \sqrt{\overline{\eta}_n} \to 2.72$ in the limit of $n \to \infty$. Considering the fact $\frac{\sqrt{3}}{2}\pi \approx 2.72 \times 1.0003$, we numerically demonstrated the relation $\Delta_n = \sqrt{\overline{\eta}_{n+1}} - \sqrt{\overline{\eta}_n} = \frac{\sqrt{3}}{2}\pi$ in the large n limit. In fact, this relation was firstly obtained in the case of massless scalar fields in [38] and supported by numerical data in [19]. We also pointed out that the WKB analysis could be generalized to more general models, which implies that the relation may be a very universal behavior.

Acknowledgments

This work was supported by the Shandong Provincial Natural Science Foundation of China under Grant

No. ZR2018QA008. This work was also supported by a grant from Qufu Normal University of China under

Grant No. xkjjc201906.

- [1] B. P. Abbott et al. (Virgo and LIGO Scientific Collaborations), Phys. Rev. Lett. 116(2016)061102.
- [2] B. P. Abbott et al. (Virgo and LIGO Scientific Collaborations), Phys. Rev. Lett. 116(2016)241103.
- [3] B. P. Abbott et al. (Virgo and LIGO Scientific Collabora- tions), Phys. Rev. X 6(2016)041015.
- [4] J. D. Bekenstein, Transcendence of the law of baryon-number conservation in black hole physics, Phys. Rev. Lett. 28, 452 (1972).
- [5] J. E. Chase, Event horizons in Static Scalar-Vacuum Space-Times, Commun. Math. Phys. 19, 276 (1970).
- [6] C. Teitelboim, Nonmeasurability of the baryon number of a black-hole, Lett. Nuovo Cimento 3, 326 (1972).
- [7] R. Ruffini and J. A. Wheeler, Introducing the black hole, Phys. Today 24, 30 (1971).
- [8] J.D. Bekenstein, Novel "no-scalar-hair" theorem for black holes, Phys. Rev. D 51(1995)no.12,R6608.
- [9] D. Núñez, H. Quevedo, and D. Sudarsky, Black Holes Have No Short Hair, Phys. Rev. Lett. 76, 571 (1996).
- [10] S. Hod, Hairy Black Holes and Null Circular Geodesics, Phys. Rev. D 84, 124030(2011).
- [11] Yan Peng, Hair mass bound in the black hole with non-zero cosmological constants, Physical Review D 98,104041,(2018).
- [12] J. D. Bekenstein, Black hole hair: 25-years after, arXiv:gr-qc/9605059.
- [13] Carlos A. R. Herdeiro, Eugen Radu, Asymptotically flat black holes with scalar hair: a review, Int.J.Mod.Phys.D 24(2015)09,1542014.
- [14] Avraham E. Mayo, Jacob D. Bekenstein, No hair for spherical black holes: Charged and nonminimally coupled scalar field with selfinteraction, Phys. Rev. D 54(1996)5059-5069.
- [15] S. Hod, No nonminimally coupled massless scalar hair for spherically symmetric neutral black holes, Phys. Lett. B 771(2017)521-523.
- [16] S. Hod, No hair for spherically symmetric neutral black holes: Nonminimally coupled massive scalar fields, Phys. Rev. D 96(2017) no.12,124037.
- [17] Thomas P. Sotiriou and Shuang-Yong Zhou, Black Hole Hair in Generalized Scalar-Tensor Gravity, Phys. Rev. Lett. 112(2014)251102.
- [18] Daniela D. Doneva and Stoytcho S. Yazadjiev, New Gauss-Bonnet black holes with curvature induced scalarization in the extended scalar-tensor theories, Phys. Rev. Lett. 120(2018)131103.
- [19] Hector O. Silva, Jeremy Sakstein, Leonardo Gualtieri, Thomas P. Sotiriou, Emanuele Berti, Spontaneous scalarization of black holes and compact stars from a Gauss-Bonnet coupling, Phys. Rev. Lett. 120(2018)131104.
- [20] G. Antoniou, A. Bakopoulos, P. Kanti, Evasion of No-Hair Theorems and Novel Black-Hole Solutions in Gauss-Bonnet Theories, Phys. Rev. Lett. 120(2018)131102.
- [21] T. Damour and G. Esposito-Farese, Nonperturbative strong-field effects in tensor-scalar theories of gravitation, Phys. Rev. Lett. 70(1993)2220.
- [22] Pedro V. P. Cunha, Carlos A. R. Herdeiro, Eugen Radu, Spontaneously scalarised Kerr black holes, Phys. Rev. Lett. 123(2019)011101.
- [23] Carlos A.R. Herdeiro, Eugen Radu, Nicolas Sanchis-Gual, José A. Font, Spontaneous Scalarization of Charged Black Holes, Phys. Rev. Lett. 121(2018)101102.
- [24] Pedro G.S. Fernandes, Carlos A.R. Herdeiro, Alexandre M. Pombo, Eugen Radu, Nicolas Sanchis-Gual, Spontaneous Scalarisation of Charged Black Holes: Coupling Dependence and Dynamical Features, Class. Quant. Grav. 36(2019)no.13,134002.
- [25] Yves Brihaye, Carlos Herdeiro, Eugen Radu, The scalarised Schwarzschild-NUT spacetime, Phys. Lett. B 788(2019)295-301.
- [26] Yves Brihaye, Betti Hartmann, Charged scalar-tensor solitons and black holes with (approximate) Anti-de Sitter asymptotics, JHEP 1901(2019)142.
- [27] Carlos A. R. Herdeiro, Eugen Radu, Black hole scalarisation from the breakdown of scale-invariance, Phys. Rev. D 99(2019)084039.
- [28] Daniela D. Doneva, Stella Kiorpelidi, Petya G. Nedkova, Eleftherios Papantonopoulos, Stoytcho S. Yazadjiev, Charged Gauss-Bonnet black holes with curvature induced scalarization in the extended scalar-tensor theories, Phys. Rev. D 98(2018)104056.
- [29] Hayato Motohashi, Shinji Mukohyama, Shape dependence of spontaneous scalarization, Phys. Rev. D 99(2019)044030.
- [30] Masato Minamitsuji, Taishi Ikeda Scalarized black holes in the presence of the coupling to Gauss-Bonnet gravity,

Phys. Rev. D 99(2019)044017.

- [31] De-Cheng Zou, Yun Soo Myung, Scalarized charged black holes with scalar mass term, arXiv:1909.11859[gr-qc].
- [32] Caio F. B. Macedo, Jeremy Sakstein, Emanuele Berti, Leonardo Gualtieri, Hector O. Silva, Thomas P. Sotiriou, Self-interactions and Spontaneous Black Hole Scalarization, Phys. Rev. D 99(2019)104041.
- [33] Jose Luis Blzquez-Salcedo, Daniela D. Doneva, Jutta Kunz, Stoytcho S. Yazadjiev Radial perturbations of the scalarized EGB black holes, Phys. Rev. D 98(2018)084011.
- [34] Yun Soo Myung, De-Cheng Zou, Instability of Reissner-Nordström black hole in Einstein-Maxwell-scalar theory, Eur. Phys. J. C 79(2019)no.3,273.
- [35] Yun Soo Myung, De-Cheng Zou, Stability of scalarized charged black holes in the Einstein-Maxwell-Scalar theory, Eur. Phys. J. C 79(2019)no.8,641.
- [36] Yan Peng, Scalarization of horizonless reflecting stars: neutral scalar fields non-minimally coupled to Maxwell fields Phys.Lett.B 804(2020)135372.
- [37] De-Cheng Zou, Yun Soo Myung, Scalar hairy black holes in Einstein-Maxwell-conformally coupled scalar theory, arXiv:1911.08062[gr-qc].
- [38] S. Hod, Spontaneous scalarization of Gauss-Bonnet black holes: Analytic treatment in the linearized regime, Phys. Rev. D 100(2019)064039.
- [39] S. Hod, Gauss-Bonnet black holes supporting massive scalar field configurations: the large-mass regime, Eur. Phys. J. C 79(2019)966.
- [40] S. Hod, Spontaneous scalarization of charged Reissner-Nordström black holes: Analytic treatment along the existence line, Physics Letters B 798(2019)135025.
- [41] Yan Peng, Scalarization of compact stars in the scalar-Gauss-Bonnet gravity, JHEP 12(2019)064.
- [42] Yan Peng, Analytical investigations on formations of hairy neutral reflecting shells in the scalar-Gauss-Bonnet gravity, Eur.Phys.J.C 80(2020)3,202.
- [43] S. Hod, Stationary bound-state scalar configurations supported by rapidly-spinning exotic compact objects, Phys.Lett.B 770(2017)186-192.
- [44] S. Hod, Marginally bound resonances of charged massive scalar fields in the background of a charged reflecting shell, Physics Letters B 768(2017)97-102.