

GALOIS-THEORETIC FEATURES FOR 1-SMOOTH PRO- p GROUPS

CLAUDIO QUADRELLI

ABSTRACT. Let p be a prime. A pro- p group G is said to be 1-smooth if it can be endowed with a continuous representation $\theta: G \rightarrow \mathrm{GL}_1(\mathbb{Z}_p)$ such that every open subgroup H of G , together with the restriction $\theta|_H$, satisfies a formal version of Hilbert 90. We prove that every 1-smooth pro- p group contains a unique maximal closed abelian normal subgroup, in analogy with a result by Engler and Koenigsmann on maximal pro- p Galois groups of fields, and that if a 1-smooth pro- p group is solvable, then it is locally uniformly powerful, in analogy with a result by Ware on maximal pro- p Galois groups of fields. Finally we ask whether 1-smooth pro- p groups satisfy a “Tits’ alternative”.

1. INTRODUCTION

Throughout the paper p will denote a prime number, and \mathbb{K} a field containing a root of unity of order p . Let $\mathbb{K}(p)$ denote the compositum of all finite Galois p -extensions of \mathbb{K} . The maximal pro- p Galois group of \mathbb{K} , denoted by $G_{\mathbb{K}}(p)$, is the Galois group $\mathrm{Gal}(\mathbb{K}(p)/\mathbb{K})$, and it coincides with the maximal pro- p quotient of the absolute Galois group of \mathbb{K} . Characterising maximal pro- p Galois groups of fields among pro- p groups is one of the most important — and challenging — problems in Galois theory. One of the obstructions for the realization of a pro- p group as maximal pro- p Galois group for some field \mathbb{K} is given by the Artin-Schrier theorem: the only finite group realizable as $G_{\mathbb{K}}(p)$ is the cyclic group of order 2 (cf. [1]).

The proof of the celebrated *Bloch-Kato conjecture*, completed by M. Rost and V. Voevodsky with Ch. Weibel’s “patch” (cf. [12, 27, 29]) provided new tools to study absolute Galois groups of field and their maximal pro- p quotients (see, e.g., [2, 3, 17, 21]). In particular, the now-called Norm Residue Theorem implies that the \mathbb{Z}/p -cohomology algebra of a maximal pro- p Galois group $G_{\mathbb{K}}(p)$

$$H^\bullet(G_{\mathbb{K}}(p), \mathbb{Z}/p) := \bigoplus_{n \geq 0} H^n(G_{\mathbb{K}}(p), \mathbb{Z}/p),$$

with \mathbb{Z}/p a trivial $G_{\mathbb{K}}(p)$ -module and endowed with the cup-product, is a quadratic algebra: i.e., all its elements of positive degree are combinations of products of elements of degree 1, and its defining relations are homogeneous relations of degree 2 (see § 2.3 below). For instance, from this property one may recover the Artin-Schreier obstruction (see, e.g., [17, § 2]).

Date: March 14, 2024.

2010 *Mathematics Subject Classification.* Primary 12G05; Secondary 20E18, 20J06, 12F10.

Key words and phrases. Galois cohomology, Maximal pro- p Galois groups, Bloch-Kato conjecture, Kummerian pro- p pairs, Tits’ alternative.

More recently, a formal version of *Hilbert 90* for pro- p groups was employed to find further results on the structure of maximal pro- p Galois groups (see [9, 19, 21]). A pair $\mathcal{G} = (G, \theta)$ consisting of a pro- p group G endowed with a continuous representation $\theta: G \rightarrow \mathrm{GL}_1(\mathbb{Z}_p)$ is called a *pro- p pair*. For a pro- p pair $\mathcal{G} = (G, \theta)$ let $\mathbb{Z}_p(1)$ denote the continuous left G -module isomorphic to \mathbb{Z}_p as an abelian pro- p group, with G -action induced by θ (namely, $g.v = \theta(g) \cdot v$ for every $v \in \mathbb{Z}_p(1)$). The pair \mathcal{G} is called a *Kummerian pro- p pair* if the canonical map

$$H^1(G, \mathbb{Z}_p(1)/p^n) \longrightarrow H^1(G, \mathbb{Z}_p(1)/p),$$

is surjective for every $n \geq 1$. Moreover the pair \mathcal{G} is said to be a *1-smooth pro- p pair* if every closed subgroup H , endowed with the restriction $\theta|_H$, gives rise to a Kummerian pro- p pair (see Definition 2.1). By Kummer theory, the maximal pro- p Galois group $G_{\mathbb{K}}(p)$ of a field \mathbb{K} , together with the pro- p cyclotomic character $\theta_{\mathbb{K}}: G_{\mathbb{K}}(p) \rightarrow \mathrm{GL}_1(\mathbb{Z}_p)$ (induced by the action of $G_{\mathbb{K}}(p)$ on the roots of unity of order a p -power lying in $\mathbb{K}(p)$) gives rise to a 1-smooth pro- p pair $\mathcal{G}_{\mathbb{K}}$ (see Theorem 2.9).

In [5] — driven by the pursuit of an “explicit” proof of the Bloch-Kato conjecture as an alternative to the proof by Voevodsky — C. De Clerq and M. Florence introduced the 1-smoothness property, and formulated the so-called “Smoothness Conjecture”: namely, that it is possible to deduce the surjectivity of the norm residue homomorphism (which is acknowledged to be the “hard part” of the Bloch-Kato conjecture) from the fact that $G_{\mathbb{K}}(p)$ together with the pro- p cyclotomic character is a 1-smooth pro- p pair (see [5, Conj. 14.25] and [15, § 3.1.6], and Question 2.12 below).

In view of the Smoothness Conjecture, it is natural to ask which properties of maximal pro- p Galois groups of fields arise also for 1-smooth pro- p pairs. For example, the Artin-Schrier obstruction does: the only finite p -group which may complete into a 1-smooth pro- p pair is the cyclic group C_2 of order 2, together with the non-trivial representation $\theta: C_2 \rightarrow \{\pm 1\} \subseteq \mathrm{GL}_1(\mathbb{Z}_2)$ (see Example 2.10 below).

A pro- p pair $\mathcal{G} = (G, \theta)$ comes endowed with a distinguished closed subgroup: the θ -center $Z(\mathcal{G})$ of \mathcal{G} , defined by

$$Z(\mathcal{G}) = \left\langle h \in \mathrm{Ker}(\theta) \mid ghg^{-1} = h^{\theta(g)} \forall g \in G \right\rangle.$$

This subgroup is abelian, and normal in G . In [10], A. Engler and J. Koenigsmann showed that if the maximal pro- p Galois group $G_{\mathbb{K}}(p)$ of a field \mathbb{K} is not cyclic then it has a unique maximal normal abelian closed subgroup (i.e., one containing all normal abelian closed subgroups of $G_{\mathbb{K}}(p)$), which coincides with the $\theta_{\mathbb{K}}$ -center $Z(\mathcal{G}_{\mathbb{K}})$, and the short exact sequence of pro- p groups

$$\{1\} \longrightarrow Z(\mathcal{G}_{\mathbb{K}}) \longrightarrow G_{\mathbb{K}}(p) \longrightarrow G_{\mathbb{K}}(p)/Z(\mathcal{G}_{\mathbb{K}}) \longrightarrow \{1\}$$

splits. We prove a group-theoretic analogue of Engler-Koenigsmann’s result for 1-smooth pro- p groups.

Theorem 1.1. *Let G be a torsion-free pro- p group, $G \not\cong \mathbb{Z}_p$, endowed with a representation $\theta: G \rightarrow \mathrm{GL}_1(\mathbb{Z}_p)$ such that $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair. Then $Z(\mathcal{G})$ is the unique maximal normal abelian closed subgroup of G , and the quotient $G/Z(\mathcal{G})$ is a torsion-free pro- p group.*

In [28], R. Ware proved the following result on maximal pro- p Galois groups of fields: if $G_{\mathbb{K}}(p)$ is solvable, then it is locally uniformly powerful, i.e., $G_{\mathbb{K}}(p) \simeq A \rtimes \mathbb{Z}_p$, where A is a free abelian pro- p group, and the right-side factor acts by scalar multiplication by a unit of \mathbb{Z}_p (see § 3.1). We prove that the same property holds also for 1-smooth pro- p groups.

Theorem 1.2. *Let G be a solvable torsion-free pro- p group, endowed with a representation $\theta: G \rightarrow \mathrm{GL}_1(\mathbb{Z}_p)$ such that $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair. Then G is locally uniformly powerful.*

This gives a complete description of solvable torsion-free pro- p groups which may be completed into a 1-smooth pro- p pair. Moreover, Theorem 1.2 settles the Smoothness Conjecture positively for the class of solvable pro- p groups.

Corollary 1.3. *If $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair with G solvable, then G is a Bloch-Kato pro- p group, i.e., the \mathbb{Z}/p -cohomology algebra of every closed subgroup of G is quadratic.*

Remark 1.4. After the submission of this paper, I. Snopce and S. Tanushevski showed in [24] that Theorems 1.2–1.1 hold for a wider class of pro- p groups. A pro- p group is said to be *Frattini-injective* if distinct finitely generated closed subgroups have distinct Frattini subgroups (cf. [24, Def. 1.1]). By [24, Thm. 1.11 and Cor. 4.3], a pro- p group which may complete into a 1-smooth pro- p pair is Frattini-injective. By [24, Thm. 1.4] a Frattini-injective pro- p group has a unique maximal normal abelian closed subgroup, and by [24, Thm. 1.3] a Frattini-injective pro- p group is solvable if, and only if, it is locally uniformly powerful.

A solvable pro- p group does not contain a free non-abelian closed subgroup. For Bloch-Kato pro- p groups — and thus in particular for maximal pro- p Galois groups of fields containing a root of unity of order p — Ware proved the following Tits' alternative: either such a pro- p group contains a free non-abelian closed subgroup; or it is locally uniformly powerful (see [28, Cor. 1] and [17, Thm. B]). We conjecture that the same phenomenon occurs for 1-smooth pro- p groups.

Conjecture 1.5. *Let G be a torsion-free pro- p group which may be endowed with a representation $\theta: G \rightarrow \mathrm{GL}_1(\mathbb{Z}_p)$ such that $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair. Then either G is locally uniformly powerful, or G contains a closed non-abelian free pro- p group.*

2. CYCLOTOMIC PRO- p PAIRS

Henceforth, every subgroup of a pro- p group will be tacitly assumed to be closed, and the generators of a subgroup will be intended in the topological sense.

In particular, for a pro- p group G and a positive integer n , G^{p^n} will denote the closed subgroup of G generated by the p^n -th powers of all elements of G . Moreover, for two elements $g, h \in G$, we set

$$h^g = g^{-1}hg, \quad \text{and} \quad [h, g] = h^{-1} \cdot h^g,$$

and for two subgroups H_1, H_2 of G , $[H_1, H_2]$ will denote the closed subgroup of G generated by all commutators $[h, g]$ with $h \in H_1$ and $g \in H_2$. In particular, G' will

denote the commutator subgroup $[G, G]$ of G , and the Frattini subgroup $G^p \cdot G'$ of G is denoted by $\Phi(G)$. Finally, $d(G)$ will denote the minimal number of generators of G , i.e., $d(G) = \dim(G/\Phi(G))$ as a \mathbb{Z}/p -vector space.

2.1. Kummerian pro- p pairs. Let $1 + p\mathbb{Z}_p = \{1 + p\lambda \mid \lambda \in \mathbb{Z}_p\} \subseteq \mathrm{GL}_1(\mathbb{Z}_p)$ denote the pro- p Sylow subgroup of the group of units of the ring of p -adic integers \mathbb{Z}_p . A pair $\mathcal{G} = (G, \theta)$ consisting of a pro- p group G and a continuous homomorphism

$$\theta: G \longrightarrow 1 + p\mathbb{Z}_p$$

is called a *cyclotomic pro- p pair*, and the morphism θ is called an *orientation* of G (cf. [7, § 3] and [21]).

A cyclotomic pro- p pair $\mathcal{G} = (G, \theta)$ is said to be *torsion-free* if $\mathrm{Im}(\theta)$ is torsion-free: this is the case if p is odd; or if $p = 2$ and $\mathrm{Im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$. Observe that a cyclotomic pro- p pair $\mathcal{G} = (G, \theta)$ may be torsion-free even if G has non-trivial torsion — e.g., if G is the cyclic group of order p and θ is constantly equal to 1. Given a cyclotomic pro- p pair $\mathcal{G} = (G, \theta)$ one has the following constructions:

- (a) if H is a subgroup of G , $\mathrm{Res}_H(\mathcal{G}) = (H, \theta|_H)$;
- (b) if N is a normal subgroup of G contained in $\mathrm{Ker}(\theta)$, then θ induces an orientation $\bar{\theta}: G/N \rightarrow 1 + p\mathbb{Z}_p$, and we set $\mathcal{G}/N = (G/N, \bar{\theta})$;
- (c) if A is an abelian pro- p group, we set $A \rtimes \mathcal{G} = (A \rtimes G, \theta \circ \pi)$, with $a^g = a^{\theta(g)^{-1}}$ for all $a \in A$, $g \in G$, and π the canonical projection $A \rtimes G \rightarrow G$.

Given a cyclotomic pro- p pair $\mathcal{G} = (G, \theta)$, the pro- p group G has two distinguished subgroups:

- (a) the subgroup

$$(2.1) \quad K(\mathcal{G}) = \left\langle h^{-\theta(g)} \cdot h^{g^{-1}} \mid g \in G, h \in \mathrm{Ker}(\theta) \right\rangle$$

introduced in [9, § 3];

- (b) the θ -center

$$(2.2) \quad Z(\mathcal{G}) = \left\langle h \in \mathrm{Ker}(\theta) \mid ghg^{-1} = h^{\theta(g)} \forall g \in G \right\rangle$$

introduced in [17, § 1].

Both $Z(\mathcal{G})$ and $K(\mathcal{G})$ are normal subgroups of G , and they are contained in $\mathrm{Ker}(\theta)$. Moreover, $Z(\mathcal{G})$ is abelian, while

$$K(\mathcal{G}) \supseteq \mathrm{Ker}(\theta)', \quad \text{and} \quad K(\mathcal{G}) \subseteq \Phi(G).$$

Thus, the quotient $\mathrm{Ker}(\theta)/K(\mathcal{G})$ is abelian, and if \mathcal{G} is torsion-free one has an isomorphism of pro- p pairs

$$(2.3) \quad \mathcal{G}/K(\mathcal{G}) \simeq (\mathrm{Ker}(\theta)/K(\mathcal{G})) \rtimes (\mathcal{G}/\mathrm{Ker}(\theta)),$$

namely, $G/K(\mathcal{G}) \simeq (\mathrm{Ker}(\theta)/K(\mathcal{G})) \rtimes (G/\mathrm{Ker}(\theta))$ (where the action is induced by θ , in the latter), and both pro- p groups are endowed with the orientation induced by θ (cf. [18, Prop. 3.1]).

Definition 2.1. Given a cyclotomic pro- p pair $\mathcal{G} = (G, \theta)$, let $\mathbb{Z}_p(1)$ denote the continuous G -module of rank 1 induced by θ , i.e., $\mathbb{Z}_p(1) \simeq \mathbb{Z}_p$ as abelian pro- p groups, and

$g \cdot \lambda = \theta(g) \cdot \lambda$ for every $\lambda \in \mathbb{Z}_p(1)$. The pair \mathcal{G} is said to be *Kummerian* if for every $n \geq 1$ the map

$$(2.4) \quad H^1(G, \mathbb{Z}_p(1)/p^n) \longrightarrow H^1(G, \mathbb{Z}_p(1)/p),$$

induced by the epimorphism of G -modules $\mathbb{Z}_p(1)/p^n \rightarrow \mathbb{Z}_p(1)/p$, is surjective. Moreover, \mathcal{G} is *1-smooth* if $\text{Res}_H(\mathcal{G})$ is Kummerian for every subgroup $H \subseteq G$.

Observe that the action of G on $\mathbb{Z}_p(1)/p$ is trivial, as $\text{Im}(\theta) \subseteq 1 + p\mathbb{Z}_p$. We say that a pro- p group G may complete into a Kummerian, or 1-smooth, pro- p pair if there exists an orientation $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ such that the pair (G, θ) is Kummerian, or 1-smooth.

Kummerian pro- p pairs and 1-smooth pro- p pairs were introduced in [9] and in [5, § 14] respectively. In [21], if $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair, the orientation θ is said to be *1-cyclotomic*. Note that in [5, § 14.1], a pro- p pair is defined to be 1-smooth if the maps (2.4) are surjective for every *open* subgroup of G , yet by a limit argument this implies also that the maps (2.4) are surjective also for every *closed* subgroup of G (cf. [21, Cor. 3.2]).

Remark 2.2. Let $\mathcal{G} = (G, \theta)$ be a cyclotomic pro- p pair. Then \mathcal{G} is Kummerian if, and only if, the map

$$H_{\text{cts}}^1(G, \mathbb{Z}_p(1)) \longrightarrow H^1(G, \mathbb{Z}_p(1)/p),$$

induced by the epimorphism of continuous left G -modules $\mathbb{Z}_p(1) \rightarrow \mathbb{Z}_p(1)/p$, is surjective (cf. [21, Prop. 2.1]) — here H_{cts}^* denotes continuous cochain cohomology as introduced by J. Tate in [26].

One has the following group-theoretic characterization of Kummerian torsion-free pro- p pairs (cf. [9, Thm. 5.6 and Thm. 7.1] and [20, Thm. 1.2]).

Proposition 2.3. *A torsion-free cyclotomic pro- p pair $\mathcal{G} = (G, \theta)$ is Kummerian if and only if $\text{Ker}(\theta)/K(\mathcal{G})$ is a free abelian pro- p group.*

Remark 2.4. Let $\mathcal{G} = (G, \theta)$ be a cyclotomic pro- p pair with $\theta \equiv \mathbf{1}$, i.e., θ is constantly equal to 1. Since $K(\mathcal{G}) = G'$ in this case, \mathcal{G} is Kummerian if and only if the quotient G/G' is torsion-free. Hence, by Proposition 2.3, \mathcal{G} is 1-smooth if and only if H/H' is torsion-free for every subgroup $H \subseteq G$. Pro- p groups with such property are called *absolutely torsion-free*, and they were introduced by T. Würfel in [30]. In particular, if $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair (with θ non-trivial), then $\text{Res}_{\text{Ker}(\theta)}(\mathcal{G}) = (\text{Ker}(\theta), \mathbf{1})$ is again 1-smooth, and thus $\text{Ker}(\theta)$ is absolutely torsion-free. Hence, a pro- p group which may complete into a 1-smooth pro- p pair is an absolutely torsion-free-by-cyclic pro- p group.

Example 2.5. (a) A cyclotomic pro- p pair (G, θ) with G a free pro- p group is 1-smooth for any orientation $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ (cf. [21, § 2.2]).

(b) A cyclotomic pro- p pair (G, θ) with G an infinite Demushkin pro- p group is 1-smooth if and only if $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ is defined as in [14, Thm. 4] (cf. [9, Thm. 7.6]). E.g., if G has a minimal presentation

$$G = \left\langle x_1, \dots, x_d \mid x_1^{p^f} [x_1, x_2] \cdots [x_{d-1}, x_d] = 1 \right\rangle$$

with $f \geq 1$ (and $f \geq 2$ if $p = 2$), then $\theta(x_2) = (1 - p^f)^{-1}$, while $\theta(x_i) = 1$ for $i \neq 2$.

- (c) For $p \neq 2$ let G be the pro- p group with minimal presentation

$$G = \langle x, y, z \mid [x, y] = z^p \rangle.$$

Then the pro- p pair (G, θ) is not Kummerian for any orientation $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ (cf. [9, Thm. 8.1]).

- (d) Let

$$H = \left\{ \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \mid a, b, c \in \mathbb{Z}_p \right\}$$

be the Heisenberg pro- p group. The pair $(H, \mathbf{1})$ is Kummerian, as $H/H' \simeq \mathbb{Z}_p^2$, but H is not absolutely torsion-free. In particular, H can not complete into a 1-smooth pro- p pair (cf. [18, Ex. 5.4]).

- (e) The only 1-smooth pro- p pair (G, θ) with G a finite p -group is the cyclic group of order 2 $G \simeq \mathbb{Z}/2$, endowed with the only non-trivial orientation $\theta: G \rightarrow \{\pm 1\} \subseteq 1 + 2\mathbb{Z}_2$ (cf. [9, Ex. 3.5]).

Remark 2.6. By Example 2.5–(e), if $\mathcal{G} = (G, \theta)$ is a torsion-free 1-smooth pro- p pair, then G is torsion-free.

A torsion-free pro- p pair $\mathcal{G} = (G, \theta)$ is said to be θ -abelian if the following equivalent conditions hold:

- (i) $\text{Ker}(\theta)$ is a free abelian pro- p group, and $\mathcal{G} \simeq \text{Ker}(\theta) \rtimes (\mathcal{G}/\text{Ker}(\theta))$;
- (ii) $Z(\mathcal{G})$ is a free abelian pro- p group, and $Z(\mathcal{G}) = \text{Ker}(\theta)$;
- (iii) \mathcal{G} is Kummerian and $K(\mathcal{G}) = \{1\}$

(cf. [17, Prop. 3.4] and [20, § 2.3]). Explicitly, a torsion-free pro- p pair $\mathcal{G} = (G, \theta)$ is θ -abelian if and only if G has a minimal presentation

$$(2.5) \quad G = \langle x_0, x_i, i \in I \mid [x_0, x_i] = x_i^q, [x_i, x_j] = 1 \forall i, j \in I \rangle \simeq \mathbb{Z}_p^I \rtimes \mathbb{Z}_p$$

for some set I and some p -power q (possibly $q = p^\infty = 0$), and in this case $\text{Im}(\theta) = 1 + q\mathbb{Z}_p$. In particular, a θ -abelian pro- p pair is also 1-smooth, as every open subgroup U of G is again isomorphic to $\mathbb{Z}_p^I \rtimes \mathbb{Z}_p$, with action induced by $\theta|_U$, and therefore $\text{Res}_U(\mathcal{G})$ is $\theta|_U$ -abelian.

Remark 2.7. From [9, Thm. 5.6], one may deduce also the following group-theoretic characterization of Kummerian pro- p pairs: a pro- p group G may complete into a Kummerian oriented pro- p group if, and only if, there exists an epimorphism of pro- p groups $\varphi: G \rightarrow \bar{G}$ such that \bar{G} has a minimal presentation (2.5), and $\text{Ker}(\varphi)$ is contained in the Frattini subgroup of G (cf., e.g., [22, Prop. 3.11]).

Remark 2.8. If $G \simeq \mathbb{Z}_p$, then the pair (G, θ) is θ -abelian, and thus also 1-smooth, for any orientation $\theta: G \rightarrow 1 + p\mathbb{Z}_p$.

On the other hand, if $\mathcal{G} = (G, \theta)$ is a θ -abelian pro- p pair with $d(G) \geq 2$, then θ is the only orientation which may complete G into a 1-smooth pro- p pair. Indeed, let $\mathcal{G}' = (G, \theta')$ be a cyclotomic pro- p pair, with $\theta': G \rightarrow 1 + p\mathbb{Z}_p$ different to θ , and let $\{x_0, x_i, i \in I\}$ be a minimal generating set of G as in the presentation (2.5) — thus, $\theta(x_i) = 1$ for all $i \in I$, and $\theta(x_0) \in 1 + q\mathbb{Z}_p$. Then for some $i \in I$ one has $\theta'|_H \neq \theta|_H$, with H the subgroup of G generated by the two elements x_0 and x_i . In particular, one has $\theta([x_0, x_i]) = \theta'([x_0, x_i]) = 1$.

Suppose that \mathcal{G}' is 1-smooth. If $\theta'(x_i) \neq 1$, then

$$x_i^q = x_i \cdot x_i^q \cdot x_i^{-1} = (x_i^q)^{\theta'(x_i)} = x_i^{q\theta'(x_i)},$$

hence $x_i^{q(1-\theta'(x_i))} = 1$, a contradiction as G is torsion-free by Remark 2.6. If $\theta'(x_i) = 1$ then necessarily $\theta'(x_0) \neq \theta(x_0)$, and thus

$$x_i^{\theta(x_0)} = x_0 \cdot x_i \cdot x_0^{-1} = x_i^{\theta'(x_0)},$$

hence $x_i^{\theta(x_0)-\theta'(x_0)} = 1$, again a contradiction as G is torsion-free. (See also [21, Cor. 3.4].)

2.2. The Galois case. Let \mathbb{K} be a field containing a root of 1 of order p , and let μ_{p^∞} denote the group of roots of 1 of order a p -power contained in the separable closure of \mathbb{K} . Then $\mu_{p^\infty} \subseteq \mathbb{K}(p)$, and the action of the maximal pro- p Galois group $G_{\mathbb{K}}(p) = \text{Gal}(\mathbb{K}(p)/\mathbb{K})$ on μ_{p^∞} induces a continuous homomorphism

$$\theta_{\mathbb{K}}: G_{\mathbb{K}}(p) \longrightarrow 1 + p\mathbb{Z}_p$$

— called the *pro- p cyclotomic character of $G_{\mathbb{K}}(p)$* —, as the group of the automorphisms of μ_{p^∞} which fix the roots of order p is isomorphic to $1 + p\mathbb{Z}_p$ (see, e.g., [8, p. 202] and [9, § 4]). In particular, if \mathbb{K} contains a root of 1 of order p^k for $k \geq 1$, then $\text{Im}(\theta_{\mathbb{K}}) \subseteq 1 + p^k\mathbb{Z}_p$.

Set $\mathcal{G}_{\mathbb{K}} = (G_{\mathbb{K}}(p), \theta_{\mathbb{K}})$. Then by Kummer theory one has the following (see, e.g., [9, Thm. 4.2]).

Theorem 2.9. *Let \mathbb{K} be a field containing a root of 1 of order p . Then $\mathcal{G}_{\mathbb{K}} = (G_{\mathbb{K}}(p), \theta_{\mathbb{K}})$ is 1-smooth.*

1-smooth pro- p pairs share the following properties with maximal pro- p Galois groups of fields.

- Example 2.10.** (a) The only finite p -group which occurs as maximal pro- p Galois group for some field \mathbb{K} is the cyclic group of order 2, and this follows from the pro- p version of the Artin-Schreier Theorem (cf. [1]). Likewise, the only finite p -group which may complete into a 1-smooth pro- p pair, is the cyclic group of order 2 (endowed with the only non-trivial orientation onto $\{\pm 1\}$), as it follows from Example 2.5–(e) and Remark 2.6.
- (b) If x is an element of $G_{\mathbb{K}}(2)$ for some field \mathbb{K} and x has order 2, then x self-centralizes (cf. [4, Prop. 2.3]). Likewise, if x is an element of a pro-2 group G which may complete into a 1-smooth pro-2 pair, then x self-centralizes (cf. [21, § 6.1]).

2.3. Bloch-Kato and the Smoothness Conjecture. A non-negatively graded algebra $A_{\bullet} = \bigoplus_{n \geq 0} A_n$ over a field \mathbb{F} , with $A_0 = \mathbb{F}$, is called a *quadratic algebra* if it is 1-generated — i.e., every element is a combination of products of elements of degree 1 —, and its relations are generated by homogeneous relations of degree 2. One has the following definitions (cf. [5, Def. 14.21] and [17, § 1]).

Definition 2.11. Let G be a pro- p group, and let $n \geq 1$. Cohomology classes in the image of the natural cup-product

$$H^1(G, \mathbb{Z}/p) \times \dots \times H^1(G, \mathbb{Z}/p) \xrightarrow{\cup} H^n(G, \mathbb{Z}/p)$$

are called *symbols* (relative to \mathbb{Z}/p , viewed as trivial G -module).

- (i) If for every open subgroup $U \subseteq G$ every element $\alpha \in H^n(U, \mathbb{Z}/p)$, for every $n \geq 1$, can be written as

$$\alpha = \text{cor}_{V_1, U}^n(\alpha_1) + \dots + \text{cor}_{V_r, U}^n(\alpha_r),$$

with $r \geq 1$, where $\alpha_i \in H^n(V_i, \mathbb{Z}/p)$ is a symbol and

$$\text{cor}_{V_i, U}^n: H^n(V_i, \mathbb{Z}/p) \longrightarrow H^n(U, \mathbb{Z}/p)$$

is the *corestriction map* (cf. [16, Ch. I, § 5]), for some open subgroups $V_i \subseteq U$, then G is called a *weakly Bloch-Kato pro- p group*.

- (ii) If for every closed subgroup $H \subseteq G$ the \mathbb{Z}/p -cohomology algebra

$$H^\bullet(H, \mathbb{Z}/p) = \bigoplus_{n \geq 0} H^n(H, \mathbb{Z}/p),$$

endowed with the cup-product, is a quadratic algebra over \mathbb{Z}/p , then G is called a *Bloch-Kato pro- p group*. As the name suggests, a Bloch-Kato pro- p group is also weakly Bloch-Kato.

By the Norm Residue Theorem, if \mathbb{K} contains a root of unity of order p , then the maximal pro- p Galois group $G_{\mathbb{K}}(p)$ is Bloch-Kato. The pro- p version of the “Smoothness Conjecture”, formulated by De Clerq and Florence, states that being 1-smooth is a sufficient condition for a pro- p group to be weakly Bloch-Kato (cf. [5, Conj. 14.25]).

Conjecture 2.12. *Let $\mathcal{G} = (G, \theta)$ be a 1-smooth pro- p pair. Then G is weakly Bloch-Kato.*

In the case of $\mathcal{G} = \mathcal{G}_{\mathbb{K}}$ for some field \mathbb{K} containing a root of 1 of order p , using Milnor K -theory one may show that the weak Bloch-Kato condition implies that $H^\bullet(G, \mathbb{Z}/p)$ is 1-generated (cf. [5, Rem. 14.26]). In view of Theorem 2.9, a positive answer to the Smoothness Conjecture would provide a new proof of the surjectivity of the norm residue isomorphism, i.e., the “surjectivity” half of the Bloch-Kato conjecture (cf. [5, § 1.1]).

Conjecture 2.12 has been settled positively for the following classes of pro- p groups.

- (a) Finite p -groups: indeed, if $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair with G a finite (non-trivial) p -group, then by Example 2.5–(e) $p = 2$, G is a cyclic group of order two and $\theta: G \twoheadrightarrow \{\pm 1\}$, so that $\mathcal{G} \simeq (\text{Gal}(\mathbb{C}/\mathbb{R}), \theta_{\mathbb{R}})$, and G is Bloch-Kato.
- (b) Analytic pro- p groups: indeed if $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair with G a p -adic analytic pro- p group, then by [18, Thm. 1.1] G is locally uniformly powerful and thus Bloch-Kato (see § 3.1 below).
- (c) Pro- p completions of right-angled Artin groups: indeed, in [25] it is shown that if $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair with G the pro- p completion of a right-angled Artin group induced by a simplicial graph Γ , then necessarily θ is trivial and Γ has the diagonal property — namely, G may be constructed starting from free pro- p groups by iterating the following two operations: free pro- p products, and direct products with \mathbb{Z}_p —, and thus G is Bloch-Kato (cf. [25, Thm. 1.2]).

3. NORMAL ABELIAN SUBGROUPS

3.1. Powerful pro- p groups.

Definition 3.1. A finitely generated pro- p group G is said to be *powerful* if one has $G' \subseteq G^p$, and also $G' \subseteq G^4$ if $p = 2$. A powerful pro- p group which is also torsion-free and finitely generated is called a *uniformly powerful* pro- p group.

For the properties of powerful and uniformly powerful pro- p groups we refer to [6, Ch. 4].

A pro- p group whose finitely generated subgroups are uniformly powerful, is said to be *locally uniformly powerful*. As mentioned in the Introduction, a pro- p group G is locally uniformly powerful if, and only if, G has a minimal presentation (2.5) — i.e., G is locally powerful if, and only if, there exists an orientation $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ such that (G, θ) is a torsion-free θ -abelian pro- p pair (cf. [17, Thm. A] and [3, Prop. 3.5]).

Therefore, a locally uniformly powerful pro- p group G comes endowed automatically with an orientation $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ such that $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair. In fact, finitely generated locally uniformly powerful pro- p groups are precisely those uniformly powerful pro- p groups which may complete into a 1-smooth pro- p pair (cf. [18, Prop. 4.3]).

Proposition 3.2. *Let $\mathcal{G} = (G, \theta)$ be a 1-smooth torsion-free pro- p pair. If G is locally powerful, then \mathcal{G} is θ -abelian, and thus G is locally uniformly powerful.*

It is well-known that the \mathbb{Z}/p -cohomology algebra of a pro- p group G with minimal presentation (2.5) is the exterior \mathbb{Z}/p -algebra

$$H^\bullet(H, \mathbb{Z}/p) \simeq \bigwedge_{n \geq 0} H^1(H, \mathbb{Z}/p)$$

— if $p = 2$ then $\bigwedge_{n \geq 0} V$ is defined to be the quotient of the tensor algebra over \mathbb{Z}/p generated by V by the two-sided ideal generated by the elements $v \otimes v$, $v \in V$ —, so that $H^\bullet(G, \mathbb{Z}/p)$ is quadratic. Moreover, every subgroup $H \subseteq G$ is again locally uniformly powerful, and thus also $H^\bullet(H, \mathbb{Z}/p)$ is quadratic. Hence, a locally uniformly powerful pro- p group is Bloch-Kato.

3.2. Normal abelian subgroups of maximal pro- p Galois groups. Let \mathbb{K} be a field containing a root of 1 of order p (and also $\sqrt{-1}$ if $p = 2$). In Galois theory one has the following result, due to A. Engler, J. Koenigsmann and J. Nogueira (cf. [11] and [10]).

Theorem 3.3. *Let \mathbb{K} be a field containing a root of 1 of order p (and also $\sqrt{-1}$ if $p = 2$), and suppose that the maximal pro- p Galois group $G_{\mathbb{K}}(p)$ of \mathbb{K} is not isomorphic to \mathbb{Z}_p . Then $G_{\mathbb{K}}(p)$ contains a unique maximal abelian normal subgroup.*

By [21, Thm. 7.7], such a maximal abelian normal subgroup coincides with the $\theta_{\mathbb{K}}$ -center $Z(\mathcal{G}_{\mathbb{K}})$ of the pro- p pair $\mathcal{G}_{\mathbb{K}} = (G_{\mathbb{K}}(p), \theta_{\mathbb{K}})$ induced by the pro- p cyclotomic character $\theta_{\mathbb{K}}$ (cf. § 2.2). Moreover, the field \mathbb{K} admits a *p -Henselian valuation* with residue characteristic not p and non- p -divisible value group, such that the *residue field* κ of such a valuation gives rise to the cyclotomic pro- p pair \mathcal{G}_{κ} isomorphic to $\mathcal{G}_{\mathbb{K}}/Z(\mathcal{G}_{\mathbb{K}})$, and the

induced short exact sequence of pro- p groups

$$(3.1) \quad \{1\} \longrightarrow Z(\mathcal{G}_{\mathbb{K}}) \longrightarrow G_{\mathbb{K}}(p) \longrightarrow G_{\kappa}(p) \longrightarrow \{1\}$$

splits (cf. [10, § 1] and [8, Ex. 22.1.6] — for the definitions related to p -henselian valuations of fields we direct the reader to [8, § 15.3]). In particular, $G_{\mathbb{K}}(p)/Z(\mathcal{G}_{\mathbb{K}})$ is torsion-free.

Remark 3.4. By [21, Thm. 1.2 and Thm. 7.7], Theorem 3.3 and the splitting of (3.1) generalize to 1-smooth pro- p pairs whose underlying pro- p group is Bloch-Kato. Namely, if $\mathcal{G} = (G, \theta)$ is a 1-smooth pro- p pair with G a Bloch-Kato pro- p group, then $Z(\mathcal{G})$ is the unique maximal abelian normal subgroup of G , and it has a complement in G .

3.3. Proof of Theorem 1.1. In order to prove Theorem 1.1 (and also Theorem 1.2 later on), we need the following result.

Proposition 3.5. *Let $\mathcal{G} = (G, \theta)$ be a torsion-free 1-smooth pro- p pair, with $d(G) = 2$ and $G = \langle x, y \rangle$. If $[[x, y], y] = 1$, then $\text{Ker}(\theta) = \langle y \rangle$ and*

$$xyx^{-1} = y^{\theta(x)}.$$

Proof. Let H be the subgroup of G generated by y and $[x, y]$. Recall that by Remark 2.6, G (and hence also H) is torsion-free.

If $d(H) = 1$ then $H \simeq \mathbb{Z}_p$, as H is torsion-free. Moreover, H is generated by y and $x^{-1}yx$, and thus $xHx^{-1} \subseteq H$. Therefore, x acts on $H \simeq \mathbb{Z}_p$ by multiplication by $1 + p\lambda$ for some $\lambda \in \mathbb{Z}_p$. If $\lambda = 0$ then G is abelian, and thus $G \simeq \mathbb{Z}_p^2$ as it is absolutely torsion-free, and $\theta \equiv \mathbf{1}$ by Remark 2.8. If $\lambda \neq 0$ then x acts non-trivially on the elements of H , and thus $\langle x \rangle \cap H = \{1\}$ and $G = H \rtimes \langle x \rangle$: by (2.5), (G, θ') is a θ' -abelian pro- p pair, with $\theta': G \rightarrow 1 + p\mathbb{Z}_p$ defined by $\theta'(x) = 1 + p\lambda$ and $\theta'(y) = 1$. By Remark 2.8, one has $\theta' \equiv \theta$, and thus $\theta(x) = 1 + p\lambda$ and $\theta(y) = 1$.

If $d(H) = 2$, then H is abelian by hypothesis, and torsion-free, and thus (H, θ') is θ' -abelian, with $\theta' \equiv \mathbf{1}: H \rightarrow 1 + p\mathbb{Z}_p$ trivial. By Remark 2.8, one has $\theta' = \theta|_H$, and thus $y, [x, y] \in \text{Ker}(\theta)$. Now put $z = [x, y]$ and $t = y^p$, and let U be the open subgroup of G generated by x, z, t . Clearly, $\text{Res}_U(\mathcal{G})$ is again 1-smooth. By hypothesis one has $z^y = z$, and hence commutator calculus yields

$$(3.2) \quad [x, t] = [x, y^p] = z \cdot z^y \cdots z^{y^{p-1}} = z^p.$$

Put $\lambda = 1 - \theta(x)^{-1} \in p\mathbb{Z}_p$. Since $t \in \text{Ker}(\theta)$, by (2.1) $[x, t] \cdot t^{-\lambda}$ lies in $K(\text{Res}_U(\mathcal{G}))$. Since t and z commute, from (3.2) one deduces

$$(3.3) \quad [x, t]t^{-\lambda} = z^p t^{-\lambda} = z^p t^{-\frac{\lambda}{p}p} = \left(z t^{-\lambda/p} \right)^p \in K(\text{Res}_U(\mathcal{G})).$$

Moreover, $z t^{-\lambda/p} \in \text{Ker}(\theta|_U)$. Since $\text{Res}_U(\mathcal{G})$ is 1-smooth, by Proposition 2.3 the quotient $\text{Ker}(\theta|_U)/K(\text{Res}_U(\mathcal{G}))$ is a free abelian pro- p group, and therefore (3.3) implies that also $z t^{-\lambda/p}$ is an element of $K(\text{Res}_U(\mathcal{G}))$.

Since $K(\text{Res}_U(\mathcal{G})) \subseteq \Phi(U)$, one has $z \equiv t^{\lambda/p} \pmod{\Phi(U)}$. Then by [6, Prop. 1.9] $d(U) = 2$ and U is generated by x and t . Since $[x, t] \in U^p$ by (3.2), the pro- p group U is powerful. Therefore, $\text{Res}_U(\mathcal{G})$ is $\theta|_U$ -abelian by Proposition 3.2. In particular, the subgroup $K(\text{Res}_U(\mathcal{G}))$ is trivial, and thus

$$[x, y] = z = t^{\lambda/p} = y^{1 - \theta(x)^{-1}},$$

and the claim follows. \square

Proposition 3.5 is a generalization of [18, Prop. 5.6].

Theorem 3.6. *Let $\mathcal{G} = (G, \theta)$ be a torsion-free 1-smooth pro- p pair, with $d(G) \geq 2$.*

- (i) *The θ -center $Z(\mathcal{G})$ is the unique maximal abelian normal subgroup of G .*
- (ii) *The quotient $G/Z(\mathcal{G})$ is a torsion-free pro- p group.*

Proof. Recall that G is torsion-free by Remark 2.6. Since $Z(\mathcal{G})$ is an abelian normal subgroup of G by definition, in order to prove (i) we need to show that if A is an abelian normal subgroup of G , then $A \subseteq Z(\mathcal{G})$.

First, we show that $A \subseteq \text{Ker}(\theta)$. If $A \simeq \mathbb{Z}_p$, let y be a generator of A . For every $x \in G$ one has $xyx^{-1} \in A$, and thus $xyx^{-1} = y^\lambda$, for some $\lambda \in 1 + p\mathbb{Z}_p$. Let H be the subgroup of G generated by x and y , for some $x \in G$ such that $d(H) = 2$. Then the pair (H, θ') is θ' -abelian for some orientation $\theta': H \rightarrow 1 + p\mathbb{Z}_p$ such that $y \in \text{Ker}(\theta')$, as H has a presentation as in (2.5). Since both $\text{Res}_H(\mathcal{G})$ and (H, θ') are 1-smooth pro- p pairs, by Remark 2.8 one has $\theta' = \theta|_H$, and thus $A \subseteq \text{Ker}(\theta)$.

If $A \not\simeq \mathbb{Z}_p$, then A is a free abelian pro- p group with $d(A) \geq 2$, as G is torsion-free. Therefore, by Remark 2.4 the pro- p pair $(A, 1)$ is 1-smooth. Since also $\text{Res}_A(\mathcal{G})$ is 1-smooth, Remark 2.8 implies that $\theta|_A = 1$, and hence $A \subseteq \text{Ker}(\theta)$.

Now, for arbitrary elements $x \in G$ and $y \in A$, put $z = [x, y]$. Since A is normal in G , one has $z \in A$, and since A is abelian, one has $[z, y] = 1$. Then Proposition 3.5 applied to the subgroup of G generated by $\{x, y\}$ yields $xyx^{-1} = x^{\theta(x)}$, and this completes the proof of statement (i).

In order to prove statement (ii), suppose that $y^p \in Z(\mathcal{G})$ for some $y \in G$. Then $y^p \in \text{Ker}(\theta)$, and since $\text{Im}(\theta)$ has no non-trivial torsion, also y lies in $\text{Ker}(\theta)$. Since G is torsion-free by Remark 2.6, $y^p \neq 1$. Let H be the subgroup of G generated by y and x , for some $x \in G$ such that $d(H) \geq 2$. Since $xy^px^{-1} = (y^p)^{\theta(x)}$, commutator calculus yields

$$(3.4) \quad y^{p(1-\theta(x)^{-1})} = [x, y^p] = [x, y] \cdot [x, y]^y \cdots [x, y]^{y^{p-1}}.$$

Put $z = [x, y]$, and let S be the subgroup of H generated by y, z . Clearly, $\text{Res}_S(\mathcal{G})$ is 1-smooth, and since $y, z \in \text{Ker}(\theta)$, one has $\theta|_S = 1$, and thus S/S' is a free abelian pro- p group by Remark 2.4. From (3.4) one deduces

$$(3.5) \quad y^{p(1-\theta(x)^{-1})} \cdot z^{-p} \equiv \left(y^{1-\theta(x)^{-1}} \cdot z^{-1} \right)^p \equiv 1 \pmod{S'}.$$

Since S/S' is torsion-free, (3.5) implies that $z \equiv y^{1-\theta(x)^{-1}} \pmod{\Phi(S)}$, so that S is generated by y , and $S \simeq \mathbb{Z}_p$, as G is torsion-free. Therefore, $S' = \{1\}$, and (3.5) yields $[x, y] = y^{1-\theta(x)^{-1}}$, and this completes the proof of statement (ii). \square

Remark 3.7. Let G be a pro- p group isomorphic to \mathbb{Z}_p , and let $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ be a non-trivial orientation. Then by Example 2.5-(a), $\mathcal{G} = (G, \theta)$ is 1-smooth. Since G is abelian and $\theta(x) \neq 1$ for every $x \in G$, $x \neq 1$, $Z(\mathcal{G}) = \{1\}$, still every subgroup of G is normal and abelian.

In view of the splitting of (3.1) (and in view of Remark 3.4), it seems natural to ask the following question.

Question 3.8. *Let $\mathcal{G} = (G, \theta)$ be a torsion-free 1-smooth pro- p pair, with $d(G) \geq 2$. Is the pro- p pair $\mathcal{G}/Z(\mathcal{G}) = (G/Z(\mathcal{G}), \bar{\theta})$ 1-smooth? Does the short exact sequence of pro- p groups*

$$\{1\} \longrightarrow Z(\mathcal{G}) \longrightarrow G \longrightarrow G/Z(\mathcal{G}) \longrightarrow \{1\}$$

split?

If $\mathcal{G} = (G, \theta)$ is a torsion-free pro- p pair, then either $\text{Ker}(\theta) = G$, or $\text{Im}(\theta) \simeq \mathbb{Z}_p$, hence in the former case one has $G \simeq \text{Ker}(\theta) \rtimes (G/\text{Ker}(\theta))$, as the right-side factor is isomorphic to \mathbb{Z}_p , and thus p -projective (cf. [16, Ch. III, § 5]). Since $Z(\mathcal{G}) \subseteq Z(\text{Ker}(\theta))$ (and $Z(\mathcal{G}) = Z(G)$ if $\text{Ker}(\theta) = G$), and since $\text{Ker}(\theta)$ is absolutely torsion-free if \mathcal{G} is 1-smooth, Question 3.8 is equivalent to the following question (of its own group-theoretic interest): if G is an absolutely torsion-free pro- p group, does G split as direct product

$$G \simeq Z(G) \times (G/Z(G)) ?$$

One has the following partial answer (cf. [30, Prop. 5]): if G is absolutely torsion-free, and $Z(G)$ is finitely generated, then $\Phi_n(G) = Z(\Phi_n(G)) \times H$, for some $n \geq 1$ and some subgroup $H \subseteq \Phi_n(G)$ (here $\Phi_n(G)$ denotes the iterated Frattini series of G , i.e., $\Phi_1(G) = G$ and $\Phi_{n+1}(G) = \Phi(\Phi_n(G))$ for $n \geq 1$).

4. SOLVABLE PRO- p GROUPS

4.1. Solvable pro- p groups and maximal pro- p Galois groups. Recall that a (pro- p) group G is said to be meta-abelian if there is a short exact sequence

$$\{1\} \longrightarrow N \longrightarrow G \longrightarrow \bar{G} \longrightarrow \{1\}$$

such that both N and \bar{G} are abelian; or, equivalently, if the commutator subgroup G' is abelian. Moreover, a pro- p group G is solvable if the derived series $(G^{(n)})_{n \geq 1}$ of G — i.e., $G^{(1)} = G$ and $G^{(n+1)} = [G^{(n)}, G^{(n)}]$ — is finite, namely $G^{(N+1)} = \{1\}$ for some finite N .

Example 4.1. A non-abelian locally uniformly powerful pro- p group G is meta-abelian: if $\theta: G \rightarrow 1+p\mathbb{Z}_p$ is the associated orientation, then $G' \subseteq \text{Ker}(\theta)^p$, and thus G' is abelian.

In Galois theory one has the following result by R. Ware (cf. [28, Thm. 3], see also [13] and [17, Thm. 4.6]).

Theorem 4.2. *Let \mathbb{K} be a field containing a root of 1 of order p (and also $\sqrt{-1}$ if $p = 2$). If the maximal pro- p Galois group $G_{\mathbb{K}}(p)$ is solvable, then $\mathcal{G}_{\mathbb{K}}$ is $\theta_{\mathbb{K}}$ -abelian.*

4.2. Proof of Theorem 1.2 and Corollary 1.3. In order to prove Theorem 1.2, we prove first the following intermediate results — a consequence of Würfel's result [30, Prop. 2] and of [18, Prop. 6.11] —, which may be seen as the “1-smooth analogue” of [28, Thm. 2].

Proposition 4.3. *Let $\mathcal{G} = (G, \theta)$ be a torsion-free 1-smooth pro- p pair. If G is meta-abelian, then \mathcal{G} is θ -abelian.*

Proof. Assume first that $\theta \equiv 1$ — i.e., G is absolutely torsion-free (cf. Remark 2.4). Then G is a free abelian pro- p group by [30, Prop. 2].

Assume now that $\theta \neq 1$. Since \mathcal{G} is 1-smooth, also $\text{Res}_{\text{Ker}(\theta)}(\mathcal{G})$ and $\text{Res}_{\text{Ker}(\theta)'}(\mathcal{G})$ are 1-smooth pro- p pairs, and thus $\text{Ker}(\theta)$ and $\text{Ker}(\theta)'$ are absolutely torsion-free. Moreover, $\text{Ker}(\theta)' \subseteq G'$, and since the latter is abelian, also $\text{Ker}(\theta)'$ is abelian, i.e., $\text{Ker}(\theta)$ is meta-abelian. Thus $\text{Ker}(\theta)$ is a free abelian pro- p group by [30, Prop. 2]. Consequently, for arbitrary $y \in \text{Ker}(\theta)$ and $x \in G$, the commutator $[x, y]$ lies in $\text{Ker}(\theta)$ and $[[x, y], y] = 1$. Therefore, Proposition 3.5 implies that $xyx^{-1} = y^{\theta(y)}$ for every $x \in G$ and $y \in \text{Ker}(\theta)$, namely, \mathcal{G} is θ -abelian. \square

Note that Proposition 4.3 generalizes [30, Prop. 2] from absolutely torsion-free pro- p groups to 1-smooth pro- p groups. From Proposition 4.3, we may deduce Theorem 1.2.

Proposition 4.4. *Let $\mathcal{G} = (G, \theta)$ be a torsion-free 1-smooth pro- p pair. If G is solvable, then G is locally uniformly powerful.*

Proof. Let N be the positive integer such that $G^{(N)} \neq \{1\}$ and $G^{(N+1)} = \{1\}$. Then for every $1 \leq n \leq N$, the pro- p pair $\text{Res}_{G^{(n)}}(\mathcal{G})$ is 1-smooth, and $G^{(n)}$ is solvable, and moreover $\theta|_{G^{(n)}} \equiv 1$ if $n \geq 2$.

Suppose that $N \geq 3$. Since $G^{(N-1)}$ is metabelian and $\theta|_{G^{(N-1)}} \equiv 1$, Proposition 4.3 implies that $G^{(N-1)}$ is a free abelian pro- p group, and therefore $G^{(N)} = \{1\}$, a contradiction. Thus, $N \leq 2$, and G is meta-abelian. Therefore, Proposition 4.3 implies that the pro- p pair \mathcal{G} is θ -abelian, and hence G is locally uniformly powerful (cf. § 3.1). \square

Proposition 4.4 may be seen as the 1-smooth analogue of Ware's Theorem 4.2. Corollary 1.3 follows from Proposition 4.4 and from the fact that a locally uniformly powerful pro- p group is Bloch-Kato (cf. § 3.1).

Corollary 4.5. *Let $\mathcal{G} = (G, \theta)$ be a torsion-free 1-smooth pro- p pair. If G is solvable, then G is Bloch-Kato.*

This settles the Smoothness Conjecture for the class of solvable pro- p groups.

4.3. A Tits' alternative for 1-smooth pro- p groups. For maximal pro- p Galois groups of fields one has the following Tits' alternative (cf. [28, Cor. 1]).

Theorem 4.6. *Let \mathbb{K} be a field containing a root of 1 of order p (and also $\sqrt{-1}$ if $p = 2$). Then either $G_{\mathbb{K}}$ is $\theta_{\mathbb{K}}$ -abelian, or $G_{\mathbb{K}}(p)$ contains a closed non-abelian free pro- p group.*

Actually, the above Tits' alternative holds also for the class of Bloch-Kato pro- p groups, with p odd: if a Bloch-Kato pro- p group G does not contain any free non-abelian subgroups, then it can complete into a θ -abelian pro- p pair $\mathcal{G} = (G, \theta)$ (cf. [17, Thm. B]), this Tits' alternative holds also for $p = 2$ under the further assumption that the Bockstein morphism $\beta: H^1(G, \mathbb{Z}/2) \rightarrow H^2(G, \mathbb{Z}/2)$ is trivial, see [17, Thm. 4.11]).

Clearly, a solvable pro- p group contains no free non-abelian subgroups.

A pro- p group is *p -adic analytic* if it is a p -adic analytic manifold and the map $(x, y) \mapsto x^{-1}y$ is analytic, or, equivalently, if it contains an open uniformly powerful subgroup (cf. [6, Thm. 8.32]) — e.g., the Heisenberg pro- p group is analytic. Similarly to solvable pro- p groups, a p -adic analytic pro- p group does not contain a free non-abelian subgroup (cf. [6, Cor. 8.34]).

Even if there are several p -adic analytic pro- p groups which are solvable (e.g., finitely generated locally uniformly powerful pro- p groups), none of these two classes of pro- p groups contains the other one: e.g.,

- (a) the wreath product $\mathbb{Z}_p \wr \mathbb{Z}_p \simeq \mathbb{Z}_p^{\mathbb{Z}_p} \rtimes \mathbb{Z}_p$ is a meta-abelian pro- p group, but it is not p -adic analytic (cf. [23]);
- (b) if G is a pro- p -Sylow subgroup of $\mathrm{SL}_2(\mathbb{Z}_p)$, then G is a p -adic analytic pro- p group, but it is not solvable.

In addition, it is well-known that also for the class of pro- p completions of right-angled Artin pro- p groups one has a Tits' alternative: the pro- p completion of a right-angled Artin pro- p group contains a free non-abelian subgroup unless it is a free abelian pro- p group (i.e., unless the associated graph is complete) — and thus it is locally uniformly powerful.

In [18], it is shown that analytic pro- p groups which may complete into a 1-smooth pro- p pair are locally uniformly powerful. Therefore, after the results in [18] and [25], and Theorem 1.2, it is natural to ask whether a Tits' alternative, analogous to Theorem 4.6 (and its generalization to Bloch-Kato pro- p groups), holds also for all torsion-free 1-smooth pro- p pairs.

Question 4.7. *Let $\mathcal{G} = (G, \theta)$ be a torsion-free 1-smooth pro- p pair, and suppose that \mathcal{G} is not θ -abelian. Does G contain a closed non-abelian free pro- p group?*

In other words, we are asking whether there exists torsion-free 1-smooth pro- p pairs $\mathcal{G} = (G, \theta)$ such that G is not analytic nor solvable, and yet it contains no free non-abelian subgroups. In view of Theorem 4.6 and of the Tits' alternative for Bloch-Kato pro- p groups [17, Thm. B], a positive answer to Question 4.7 would corroborate the Smoothness Conjecture.

Observe that — analogously to Question 3.8 — Question 4.7 is equivalent to asking whether an absolutely torsion-free pro- p group which is not abelian contains a closed non-abelian free subgroup. Indeed, by Proposition 3.5 (in fact, just by [18, Prop. 5.6]), if $\mathcal{G} = (G, \theta)$ is a torsion-free 1-smooth pro- p pair and $\mathrm{Ker}(\theta)$ is abelian, then \mathcal{G} is θ -abelian.

Acknowledgement. The autor wishes to thank I. Efrat, J. Mináč, N.D. Tân and Th. Weigel for working together on maximal pro- p Galois groups and their cohomology; and P. Guillot and I. Snopce for the discussions on 1-smooth pro- p groups. Also, the author wishes to thank the editors of CMB-BMC, for their helpfulness, and the anonymous referee.

REFERENCES

- [1] E. Becker, *Euklidische Körper und euklidische Hüllen von Körpern*, J. Reine Angew. Math. **268/269** (1974), 41–52 (German). Collection of articles dedicated to Helmut Hasse on his seventy-fifth birthday, II.
- [2] S. K. Chebolu, I. Efrat, and J. Minač, *Quotients of absolute Galois groups which determine the entire Galois cohomology*, Math. Ann. **352** (2012), no. 1, 205–221.
- [3] S. K. Chebolu, J. Mináč, and C. Quadrelli, *Detecting fast solvability of equations via small powerful Galois groups*, Trans. Amer. Math. Soc. **367** (2015), no. 12, 8439–8464.
- [4] Th. C. Craven and T. L. Smith, *Formally real fields from a Galois-theoretic perspective*, J. Pure Appl. Algebra **145** (2000), no. 1, 19–36.

- [5] C. De Clercq and M. Florence, *Lifting theorems and smooth profinite groups*, 2017. Preprint, available at [arXiv:1711.06585](https://arxiv.org/abs/1711.06585).
- [6] J. D. Dixon, M. P. F. du Sautoy, A. Mann, and D. Segal, *Analytic pro- p groups*, 2nd ed., Cambridge Studies in Advanced Mathematics, vol. 61, Cambridge University Press, Cambridge, 1999.
- [7] I. Efrat, *Small maximal pro- p Galois groups*, Manuscripta Math. **95** (1998), no. 2, 237–249.
- [8] ———, *Valuations, orderings, and Milnor K -theory*, Mathematical Surveys and Monographs, vol. 124, American Mathematical Society, Providence, RI, 2006.
- [9] I. Efrat and C. Quadrelli, *The Kummerian property and maximal pro- p Galois groups*, J. Algebra **525** (2019), 284–310.
- [10] A. J. Engler and J. Koenigsmann, *Abelian subgroups of pro- p Galois groups*, Trans. Amer. Math. Soc. **350** (1998), no. 6, 2473–2485.
- [11] A. J. Engler and J. Nogueira, *Maximal abelian normal subgroups of Galois pro-2-groups*, J. Algebra **166** (1994), no. 3, 481–505.
- [12] C. Haesemeyer and Ch. Weibel, *The norm residue theorem in motivic cohomology*, Annals of Mathematics Studies, vol. 200, Princeton University Press, Princeton, NJ, 2019.
- [13] J. Koenigsmann, *Solvable absolute Galois groups are metabelian*, Invent. Math. **144** (2001), no. 1, 1–22.
- [14] J. P. Labute, *Classification of Demushkin groups*, Canad. J. Math. **19** (1967), 106–132.
- [15] J. Minač, F. Pop, A. Topaz, and K. Wickelgren, *Nilpotent Fundamental Groups*, BIRS for Mathematical Innovation and Discovery, 2017, <https://www.birs.ca/workshops/2017/17w5112/report17w5112.pdf>. Report of the workshop “Nilpotent Fundamental Groups”, Banff AB, Canada, June 2017.
- [16] J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of number fields*, 2nd ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 323, Springer-Verlag, Berlin, 2008.
- [17] C. Quadrelli, *Bloch-Kato pro- p groups and locally powerful groups*, Forum Math. **26** (2014), no. 3, 793–814.
- [18] ———, *1-smooth pro- p groups and Bloch-Kato pro- p groups*, 2019. Preprint, available at [arXiv:19040.0667](https://arxiv.org/abs/19040.0667).
- [19] ———, *Two families of pro- p groups that are not absolute Galois groups*, J. Group Theory (2021). In press.
- [20] ———, *Chasing maximal pro- p Galois groups with 1-cyclotomicity*, 2021. Preprint, available at [arXiv:2106.00335](https://arxiv.org/abs/2106.00335).
- [21] C. Quadrelli and Th. Weigel, *Profinite groups with a cyclotomic p -orientation*, Doc. Math. **25** (2020), 1881–1917.
- [22] ———, *Pro- ℓ groups with the Bogomolov property* (2021). Preprint, available at [arXiv:2103.12438](https://arxiv.org/abs/2103.12438).
- [23] A. Shalev, *Characterization of p -adic analytic groups in terms of wreath products*, J. Algebra **145** (1992), no. 1, 204–208.
- [24] I. Snopce and S. Tanushevski, *Frattini-injectivity and maximal pro- p Galois groups*, 2020. Preprint, available at [arXiv:2009.09297](https://arxiv.org/abs/2009.09297).
- [25] I. Snopce and P. A. Zalesskii, *Right-angled Artin pro- p -groups*, 2020. Preprint, available at [arXiv:2005.01685](https://arxiv.org/abs/2005.01685).
- [26] J. Tate, *Relations between K_2 and Galois cohomology*, Invent. Math. **36** (1976), 257–274.
- [27] V. Voevodsky, *On motivic cohomology with \mathbf{Z}/l -coefficients*, Ann. of Math. (2) **174** (2011), no. 1, 401–438.
- [28] R. Ware, *Galois groups of maximal p -extensions*, Trans. Amer. Math. Soc. **333** (1992), no. 2, 721–728.
- [29] Charles Weibel, *2007 Trieste lectures on the proof of the Bloch-Kato conjecture*, Some recent developments in algebraic K -theory, ICTP Lect. Notes, vol. 23, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2008, pp. 277–305. MR2509183
- [30] T. Würfel, *On a class of pro- p groups occurring in Galois theory*, J. Pure Appl. Algebra **36** (1985), no. 1, 95–103.

DEPARTMENT OF MATHEMATICS AND APPLICATIONS, UNIVERSITY OF MILANO BICOCCA, 20125 MILAN,
ITALY EU

Email address: `claudio.quadrelli@unimib.it`