

# Variational problems of splitting-type with mixed linear- superlinear growth conditions

Michael Bildhauer & Martin Fuchs

## Abstract

Variational problems of splitting-type with mixed linear- superlinear growth conditions are considered. In the twodimensional case the minimizing problem is given by

$$J[w] = \int_{\Omega} \left[ f_1(\partial_1 w) + f_2(\partial_2 w) \right] dx \rightarrow \min$$

w.r.t. a suitable class of comparison functions. Here  $f_1$  is supposed to be a convex energy density with linear growth,  $f_2$  is supposed to be of superlinear growth, for instance to be given by a  $N$ -function or just bounded from below by a  $N$ -function. One motivation for this kind of problem located between the well known splitting-type problems of superlinear growth and the splitting-type problems with linear growth (recently considered in [1]) is the link to mathematical problems in plasticity (compare [2]). Here we prove results on the appropriate way of relaxation including approximation procedures, duality, existence and uniqueness of solutions as well as some new higher integrability results.<sup>1</sup>

## 1 Introduction

In the last decades the study of variational problems with nonstandard growth conditions developed to one of the main topics in the calculus of variations and related areas. We do not want to go into details and will not present the historical line of this development. The reader will find this background information, for instance, in the recent paper [3].

Let us just mention a few aspects, which serve as a motivation for the manuscript at hand.

---

<sup>1</sup>AMS-Classification: 49J45, 49N60

As one of the first main contributions Giaquinta considered the most common prototype of energies with  $(p, q)$ -growth in the sense of minimizing a splitting functional

$$\int_{\Omega} f(\nabla u) \, dx = \int_{\Omega} \left[ f_1(\partial_1 u) + f_2(\partial_2 u) \right] \, dx \rightarrow \min \quad (1.1)$$

in a suitable class of comparison functions, where  $f_1$  and  $f_2$  are supposed to have different growth rates larger than 1. In [4] he presented a famous counterexample which shows that in general and even in the scalar case we cannot expect the smoothness of solutions to this variational problem.

Of course (1.1) also serves as a motivation to study variational problems with non-uniform ellipticity conditions. As one variant we may consider energy densities of class  $C^2$  satisfying with different exponents  $1 < p < q$  ( $c_1, c_2 > 0$ ,  $\xi, \eta \in \mathbb{R}^{nN}$ )

$$c_1(1 + |\xi|^2)^{\frac{p-2}{2}}|\eta|^2 \leq D^2 f(\xi)(\eta, \eta) \leq c_2(1 + |\xi|^2)^{\frac{q-2}{2}}|\eta|^2. \quad (1.2)$$

Here a lot of important contributions in the scalar and also in the vectorial setting can be found, we just mention [5] as one central reference in the long series of papers in this direction.

Related to (1.2), Frehse and Seregin ([6]) considered plastic materials with logarithmic hardening, i.e. the energy density  $f(\xi) = |\xi| \ln(1 + |\xi|)$  of nearly linear growth. Due to [7] we have full regularity for this particular kind of model.

We finally pass to the case of linear growth problems for which the energy density is of (uniform) linear growth w.r.t. the gradient but just satisfies a non-uniform ellipticity condition in the sense of (1.2). Of course the minimal surface case is the most prominent representative for this kind of problems satisfying with suitable constants  $a_1, b_1, c_1, c_2 > 0$ ,  $a_2, b_2 \geq 0$ , for all  $\xi, \eta \in \mathbb{R}^{nN}$  and with some exponent  $\mu > 1$

$$\begin{aligned} a_1|\xi| - a_2 &\leq f(\xi) \leq b_1|\xi| + b_2, \\ c_1(1 + |\xi|^2)^{-\frac{\mu}{2}}|\eta|^2 &\leq D^2 f(\xi)(\eta, \eta) \leq c_2(1 + |\xi|^2)^{-\frac{1}{2}}|\eta|^2. \end{aligned} \quad (1.3)$$

In the minimal surface case we have  $\mu = 3$  and we like to mention the pioneering work of Giaquinta, Modica and Souček [8], [9] in the list of outstanding contributions. In [8] and [9] a suitable relaxation is discussed together with a subsequent proof of apriori estimates. We note that the uniqueness of solutions in general is lost by passing to the relaxed problem.

In [10], condition (1.3) was introduced defining a class of  $\mu$ -elliptic energy densities. The regularity theory for minimizers was studied in a series of subsequent papers, compare, e.g., [11]. We also like to mention the Lipschitz estimates of Marcellini and Papi [12], which cover a broad class of the functionals we discussed up to now.

Very recently, the authors [1] considered variational problems of splitting-type as given in (1.1) but now with two energy parts being of linear growth. Here it turns out that the right-hand side of the ellipticity condition in (1.3) is no longer valid and we just have for all  $\xi, \eta \in \mathbb{R}^{nN}$  with a positive constant  $c$

$$D^2 f(\xi)(\eta, \eta) \leq c|\eta|^2.$$

Nevertheless, some natural assumptions still imply regularity and uniqueness properties of solutions to the relaxed problem.

In the manuscript at hand we follow this line of studying variational problems of splitting-type by now considering variational problems with mixed linear- superlinear growth conditions.

A first step in this direction was already made in Chapter 6 of [13]. The results given there follow from suitable apriori estimates which are available if the non-uniform ellipticity is not too bad. This leads to the analysis of the set of cluster points of minimizing sequences and to some kind of local interpretation for the stress tensor, although the existence and the uniqueness of dual solutions were not established (compare Remark 6.15 of [13]).

We like to finish this introductory remarks by mentioning a prominent application of a mixed linear- superlinear growth problem, which in [2] is discussed as the Hencky plasticity model. This problem takes the form (compare (4.17), Chapter I, of [2])

$$\inf_{v \in \mathcal{C}_a} \left\{ \int_{\Omega} (\operatorname{div} v)^2 dx + \int_{\Omega} \psi(\varepsilon^D(v)) dx - L(v) \right\}$$

with a suitable class  $\mathcal{C}_a \subset W^{1,2}(\Omega; \mathbb{R}^3)$ , some volume force  $L$  and the deviatoric part  $\varepsilon^D(v)$  of the symmetric gradient  $\varepsilon(v)$ . Here  $\operatorname{div} v$  enters with quadratic growth while the function  $\psi$  is of linear growth w.r.t. the tensor  $\varepsilon^D(v)$ . Let us note that this plasticity model is based on the dual point of view, i.e. the so-called sur-potential  $\psi$  is introduced through the conjugate function  $\psi^*$ . In general a more explicit expression cannot be given (see Remark 4.1, p. 75 of [2]). One particular example is of the form  $\psi(\xi^D) = \Phi(|\xi^D|)$

with (for some positive constants  $k, \nu$ )

$$\Phi(s) = \begin{cases} \nu s^2 & \text{if } |s| \leq \frac{k}{\sqrt{2\nu}}, \\ \sqrt{2k}|s| - \frac{k^2}{2\nu} & \text{if } |s| \geq \frac{k}{\sqrt{2\nu}}. \end{cases}$$

However, at this stage our main difficulty in comparison to the known results for the Hencky model is quite hidden. We postpone a refined discussion to Remark 2.1.

Now let us introduce the general framework of our considerations in a more precise way. For the sake of notational simplicity we restrict our considerations to the case that a linear growth condition is satisfied in only one coordinate direction.

Suppose that  $\Omega \subset \mathbb{R}^n$  is a bounded Lipschitz domain and let  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  be an energy density of class  $C^2(\mathbb{R}^n)$  which is decomposed in the form

$$f(\xi) = f_1(\xi_1) + f_2(\xi_2), \quad \xi = (\xi_1, \xi_2) \in \mathbb{R} \times \mathbb{R}^{n-1}. \quad (1.4)$$

Here we assume that  $f_1: \mathbb{R} \rightarrow \mathbb{R}$  and  $f_2: \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  are convex functions of class  $C^2(\mathbb{R})$  and  $C^2(\mathbb{R}^{n-1})$ , respectively, satisfying with  $a_i, b_i \geq 0$ ,  $a_1, a_3, b_1 > 0$  and with a  $N$ -function  $A: \mathbb{R} \rightarrow \mathbb{R}$ :

$$\begin{aligned} a_1|\xi_1| - a_2 &\leq f_1(\xi_1) \leq a_3|\xi_1| + a_4, \quad \xi_1 \in \mathbb{R}, \\ b_1A(|\xi_2|) - b_2 &\leq f_2(\xi_2), \quad \xi_2 \in \mathbb{R}^{n-1}. \end{aligned} \quad (1.5)$$

We wish to note that we work with energy densities of class  $C^2$  just for notational simplicity. This hypothesis just enters Section 4. It is easy to check that for example no differentiability assumptions are needed in Section 3.1, whereas the results of Section 3.2 hold for densities of class  $C^1$ .

Having the decomposition (1.4) in mind, we now suppose  $n = 2$  throughout the rest of this manuscript. This essentially clarifies the notation while the main ideas and results remain unchanged.

For the definition and the properties of  $N$ -functions and Orlicz-Sobolev space we refer to the monographs [14] or [15]. The basics needed here are summarized in [16] or [17]. We suppose that  $A: [0, \infty) \rightarrow [0, \infty)$  satisfies

(N1)  $A$  is continuous, strictly increasing and convex.

$$(N2) \quad \lim_{t \downarrow 0} \frac{A(t)}{t} = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \frac{A(t)}{t} = \infty.$$

(N3) There exist constants  $k, t_0 > 0$  such that  $A(2t) \leq kA(t)$  for all  $t \geq t_0$ ,

where (N3) is called a  $\Delta_2$ -condition near infinity. We note that there exists an exponent  $p \geq 1$  such that with some constant  $c > 0$

$$c|t|^p \leq A(t) \quad \text{for all } t \gg 1. \quad (1.6)$$

Moreover we suppose some kind of triangle inequality for  $f_2$ : there exists a real number  $c_3 > 0$  such that for all  $t, \hat{t} \in \mathbb{R}$

$$f_2(t + \hat{t}) \leq c_3 \left[ f_2(t) + f_2(\hat{t}) \right]. \quad (1.7)$$

This condition, for instance, follows from the convexity of  $f_2$  together with some  $\Delta_2$ -condition.

For the definition of the Sobolev spaces  $W_p^k$  and their local variants we refer to the textbook of Adams ([18]), the notation needed later in the case of functions of bounded variation can be found, e.g., in the monographs [19] and [20]. For the sake of completeness we recall the definition of the Orlicz-Sobolev space generated by a  $N$ -function  $A$  satisfying (N1)-(N3) (see [14], [15]).

In the following we suppose that the bounded Lipschitz domain  $\Omega$  is normal w.r.t. the  $x_2$ -axis (compare the approximation arguments presented in Section 2), i.e. there exist Lipschitz functions  $\kappa_1, \kappa_2: (a, b) \rightarrow \mathbb{R}$  such that

$$\Omega = \{x \in \mathbb{R}^2 : x_1 \in (a, b), \kappa_1(x_1) < x_2 < \kappa_2(x_1)\}. \quad (1.8)$$

The space

$$L_A(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{R} : u \text{ is a measurable function such that} \right. \\ \left. \text{there exists } \lambda > 0 \text{ with } \int_{\Omega} A(\lambda|u|) \, dx < +\infty \right\}$$

is called Orlicz space equipped with the Luxemburg norm

$$\|u\|_{L_A(\Omega)} = \inf \left\{ l > 0 : \int_{\Omega} A\left(\frac{|u|}{l}\right) \, dx \leq 1 \right\}.$$

The Orlicz-Sobolev space is given by

$$W_A^1(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} : u \text{ is a measurable function, } u, |\nabla u| \in L_A(\Omega) \right\}$$

with norm

$$\|u\|_{W_A^1(\Omega)} = \|u\|_{L_A(\Omega)} + \|\nabla u\|_{L_A(\Omega)}.$$

The closure in  $W_A^1(\Omega)$  of  $C_0^\infty(\Omega)$ -functions w.r.t. this norm is according to Theorem 2.1 of [16] (recall that we suppose (N3))

$$\mathring{W}_A^1(\Omega) = W_A^1(\Omega) \cap \mathring{W}_1^1(\Omega) .0 \quad (1.9)$$

We additionally use the notation

$$W_{1,A}^1(\Omega) := \left\{ w \in W^{1,1}(\Omega) : \partial_2 w \in L_A(\Omega) \right\},$$

$$E[v] := \int_{\Omega} f_2(v) \, dx, \quad v \in L_A(\Omega).$$

In formal accordance with (1.9) we define the class

$$\mathring{W}_{1,A}^1(\Omega) := W_{1,A}^1(\Omega) \cap \mathring{W}_1^1(\Omega).$$

observing that this set is just the completion of  $C_0^\infty(\Omega)$  in  $W_{1,A}^1(\Omega)$  w.r.t. the natural norm of  $W_{1,A}^1(\Omega)$ .

The main classes of functions under consideration are:

$$\mathcal{C}_{\text{Sob}} := u_0 + \mathring{W}_{1,A}^1(\Omega),$$

$$\mathcal{C}_{\text{BV}} := \left\{ w \in \text{BV}(\Omega) : \|\partial_2 w\|_{L_A(\Omega)} < \infty, (w - u_0)\nu_2 = 0 \text{ } \mathcal{H}^1\text{-a.e. on } \partial\Omega \right\},$$

where the boundary values  $u_0$  are always supposed to be of class  $W_\infty^1(\Omega)$  (see also Remark 6.12 in [13]). Note that in the definition of the space  $\mathcal{C}_{\text{BV}}$  we require that the distributional derivative  $\partial_2 w$  is generated by a function from the space  $L_A(\Omega)$ . Moreover, we consider the BV-trace of  $w$  and denote by  $\nu = (\nu_1, \nu_2)$  the outward unit normal to  $\partial\Omega$ .

With respect to these classes we consider the minimization problem

$$J[w] := \int_{\Omega} f(\nabla w) \, dx \rightarrow \min \quad \text{in the class } \mathcal{C}_{\text{Sob}} \quad (1.10)$$

and its relaxed version

$$\begin{aligned} K[w] &:= \int_{\Omega} f_1(\partial_1^a w) \, dx + \int_{\Omega} f_1^\infty \left( \frac{\partial_1^s w}{|\partial_1^s w|} \right) d|\partial_1^s w| \\ &\quad + \int_{\partial\Omega} f_1^\infty((u_0 - w)\nu_1) d\mathcal{H}^1 + E[\partial_2 w] \\ &=: K_1[w] + E[\partial_2 w] \rightarrow \min \quad \text{in the class } \mathcal{C}_{\text{BV}}. \end{aligned} \quad (1.11)$$

Here  $\nabla^a w$  denotes the absolutely continuous part of  $\nabla w$  w.r.t. the Lebesgue measure,  $\nabla^s w$  represents the singular part.

As a matter of fact, problem (1.10) in general is not solvable and one has to pass to the relaxed version in order to have the existence of at least generalized minimizers. The approach via relaxation in the case of linear growth is well-known and outlined, e.g., in the monographs [19] or [20].

It will turn out in Section 2 and in Section 3 that the functional  $K$  together with the class  $\mathcal{C}_{BV}$  is the suitable choice in the setting at hand: in Section 3.1 we show that there exists a solution  $\bar{u}$  of problem (1.11). Moreover, with the help of the geometric approximation procedure of Section 2, we show in Corollary 3.1 that the infima of (1.10) and (1.11) are equal. In the case that the superlinear part is given by a  $N$ -function, we obtain in addition a complete dual point of view.

In Section 4 the apriori higher integrability and regularity results of the recent paper [1] on splitting-type variational problems with linear growth are essentially refined and carried over to the mixed linear- superlinear setting.

Concerning the function  $f_1$  we suppose that there exist real numbers  $\mu > 1$  and  $\gamma \geq 0$  such that

$$c_1(1 + |t|)^{-\mu} \leq f_1''(t) \leq \bar{c}_1(1 + |t|)^\gamma, \quad t \in \mathbb{R}, \quad (1.12)$$

holds with constants  $c_1, \bar{c}_2 > 0$ . We note that  $\mu > 1$  is in accordance with the required linear growth of  $f_1$ .

For  $f_2$  we suppose that there exist real numbers  $\hat{\mu} < 2$  and  $q \geq 1$  such that

$$c_2(1 + |t|)^{-\hat{\mu}} \leq f_2''(t) \leq \bar{c}_2(1 + |t|)^{q-2}, \quad t \in \mathbb{R}, \quad (1.13)$$

holds with constants  $c_2, \bar{c}_2 > 0$ . Since  $f_2$  is of superlinear growth, the condition  $\hat{\mu} < 2$  is a quite mild assumption. In the case of  $(p, q)$ -growth of  $f_2$  we have  $-\hat{\mu} = p - 2$  for some  $p > 1$ . We also note that for  $n > 2$  the condition (1.13) is replaced by considering a function  $f_2: \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  such that with constants  $\lambda, \Lambda > 0$  and for all  $\xi_2, \eta \in \mathbb{R}^{n-1}$

$$\lambda(1 + |\xi_2|)^{-\hat{\mu}}|\eta|^2 \leq D^2 f_2(\xi_2)(\eta, \eta) \leq \Lambda(1 + |\xi_2|)^{q-2}|\eta|^2. \quad (1.14)$$

A Caccioppoli-type inequality w.r.t.  $\partial_2 \nabla u$  and  $\partial_1 \nabla u$ , respectively, using in addition negative exponents gives different variants of regularity results depending on the properties of  $f_{1,2}$ .

In Section 5 we finally turn our attention to the question of uniqueness of solutions. First results were already given in Theorem 4.1 and Corollary 4.1 by quoting [13].

It remains to discuss the  $N$ -function case. If the ellipticity parameter from (1.12) satisfies  $\mu < 2$ , then the smoothness properties of  $\sigma$  together with the uniqueness of  $\sigma$  imply the uniqueness of generalized solutions. In order to make this argument precise we prove a generalization of [21], Theorem 7, to the situation at hand.

## 2 Approximation procedure

In this section we present an approximation procedure which is adapted to the particular linear- superlinear setting. Although the arguments seem to be quite technical, the principle idea is a geometric one.

We have to take care of various aspects:

- A retracting and smoothing procedure of the form  $u_0 + \eta_\varepsilon * [(u - u_0)(x + \delta e_2)]$  is compatible with Lebesgue spaces. However it does not work w.r.t. the “BV”-direction  $e_1$  which is due to the possible concentration of masses on the boundary.
- In the linear growth situation the methods of local approximation (compare, e.g., [20], Theorem 1.17, p. 14) together with some extension by  $u_0$  outside of  $\Omega$  (see, e.g., [8], [9]) serve as a powerful tool. However, a partition of the unity  $\{\varphi_i\}$  is involved in this kind of argument. This causes serious difficulties proving the convergence of  $f_2(\partial_2 w_m)$  for the approximating sequence  $w_m$  since the derivatives of  $\varphi_i$  do not cancel when calculating the integral of  $f_2$  evaluated at the corresponding expression.
- Combining and adjusting both methods and using the geometric structure of the domain we obtain a partition of the unity such that the derivatives w.r.t. the relevant direction vanish.

We start with a generalization of Lemma B.1 of [13] including strong  $L^p$ -convergence of  $\partial_2 w_m$ . The main new feature is the way of constructing the sequence  $\{w_m\}$  which is crucial for proving Lemma 2.2.

**Lemma 2.1.** *Let  $w \in \text{BV}(\Omega)$  such that  $\partial_2 w \in L^p(\Omega)$  for some  $1 \leq p < \infty$  and such that  $(w - u_0)\nu_2 = 0$  a.e. on  $\partial\Omega$ .*

Then there exists a sequence  $\{w_m\}$  such that for all  $m \in \mathbb{N}$  we have  $w_m \in W_1^1(\Omega) \cap C^\infty(\Omega)$ ,  $\partial_2 w_m \in L^p(\Omega)$ ,  $\text{trace } w_m = \text{trace } w$  and such that we have the convergences

$$\begin{aligned} \lim_{m \rightarrow \infty} \int_{\Omega} |w_m - w| \, dx &= 0, \\ \lim_{m \rightarrow \infty} \int_{\Omega} \sqrt{1 + |\nabla w_m|^2} \, dx &= \int_{\Omega} \sqrt{1 + |\nabla w|^2}, \\ \lim_{m \rightarrow \infty} \int_{\Omega} |\partial_2 w_m - \partial_2 w|^p \, dx &= 0. \end{aligned}$$

*Proof.* Recalling our assumption (1.8) imposed on the domain  $\Omega$  we may consider w.l.o.g. the case

$$\Omega = (-1, 1) \times (-1, 1). \quad (2.1)$$

We fix a function  $w \in BV(\Omega)$  and proceed in five steps.

**Step 1.** In the following we suppose that  $u_0 = 0$ . The general case is obtained by considering  $w - u_0$  and adding  $u_0$  at the end of the proof.

We then reduce the problem by choosing two smooth functions  $\psi_1, \psi_2: [-1, 1] \rightarrow [0, 1]$  such that  $\psi_1 + \psi_2 \equiv 1$ ,  $\psi_1(t) = 0$  on  $[-1, -1/2]$ ,  $\psi_1(t) = 1$  on  $[1/2, 1]$  and  $\psi_2(t) = 1$  on  $[-1, -1/2]$ ,  $\psi_2(t) = 0$  on  $[1/2, 1]$ .

We consider  $\psi_1(x_2)w$  and  $\psi_2(x_2)w$  separately, hence w.l.o.g.  $w \equiv 0$  in a neighborhood of  $[x_2 = -1]$ .

**Step 2.** Fix some  $\varepsilon_0 > 0$  and let (w.r.t. the  $x_2$ -direction)

$$w_{\varepsilon_0}(x) = w(x + \varepsilon_0 e_2), \quad (2.2)$$

where  $w$  is extended by 0 on  $(-1, 1) \times [1, \infty)$ . At the end of our proof we pass to the limit  $\varepsilon_0 \rightarrow 0$ .

Thus we may suppose w.l.o.g. that

$$w \equiv 0 \quad \text{on} \quad [(-1, 1) \times (1 - \varepsilon_0, 1)] \cap [(-1, 1) \times (-1, -1 + \varepsilon_0)]. \quad (2.3)$$

**Step 3.** We now take [20], proof of Theorem 1.17, as a reference (compare also [13]), Lemma B.1), fix  $\varepsilon > 0$ , (recalling Step 1 and Step 2) and for  $l \in \mathbb{N}$  we let

$$\Omega_k = \Omega_k^l := \left\{ x \in \Omega : -1 + \frac{1}{l+k} < x_1 < 1 - \frac{1}{l+k} \right\}, \quad k \in \mathbb{N}_0,$$

where  $l$  is chosen sufficiently large such that

$$\int_{\Omega - \Omega_0} |\nabla w| \, dx < \varepsilon. \quad (2.4)$$

With this notation we define  $A_1 := \Omega_2$  and

$$\begin{aligned} A_i = \Omega_{i+1} - \overline{\Omega}_{i-1} &:= \left\{ x \in \Omega : -1 + \frac{1}{l+i+1} < x_1 < -1 + \frac{1}{l+i-1} \right. \\ &\quad \left. \text{and } 1 - \frac{1}{l+i-1} < x_1 < 1 - \frac{1}{l+i+1} \right\} \\ &=: \{x \in \Omega : x_1 \in I_i^- \cup I_i^+\}. \end{aligned}$$

A partition  $\{\varphi_i\}$  of the unity is defined w.r.t. these sets by

$$\varphi_i \in C_0^\infty(A_i), \quad 0 \leq \varphi_i \leq 1, \quad \sum_{i=1}^{\infty} \varphi_i = 1 \text{ on } \Omega.$$

For proving Lemma 2.2 below it will be crucial to observe that the functions  $\varphi_i$  may be chosen respecting the structure of the stripes, i.e. for all  $i \in \mathbb{N}$

$$\varphi_i(x_1, x_2) = \tilde{\varphi}_i(x_1), \quad \tilde{\varphi}_i \in C_0^\infty(I_i^- \cup I_i^+). \quad (2.5)$$

**Step 4.** Now we proceed essentially as described in Lemma B.1 of [13]: let  $\Omega_{-1} = \emptyset$  and denote by  $\eta$  a smoothing kernel. On account of (2.3) we select  $\varepsilon_i$  small enough such that the smoothing procedure is well defined and such that we have

$$\begin{aligned} \text{spt } \eta_{\varepsilon_i} * (\varphi_i w) &\subset \Omega_{i+2} - \overline{\Omega}_{i-2}, \\ \int_{\Omega} |\eta_{\varepsilon_i} * (\varphi_i w) - \varphi_i w| \, dx &< 2^{-i} \varepsilon, \\ \int_{\Omega} |\eta_{\varepsilon_i} * (w \nabla \varphi_i) - w \nabla \varphi_i| \, dx &< 2^{-i} \varepsilon, \\ \int_{\Omega} |\eta_{\varepsilon_i} * \partial_2(\varphi_i w) - \partial_2(\varphi_i w)|^p \, dx &< 2^{-i} \varepsilon. \end{aligned} \quad (2.6)$$

Moreover, the analogue to (2.3) holds for  $w_m$  with some  $\tilde{\varepsilon}_0 < \varepsilon_0$ . Here with the choice  $\varepsilon = 1/m$  we have set

$$w_m = \sum_{i=1}^{\infty} \eta_{\varepsilon_i} * (\varphi_i w).$$

By the above remarks we suppose with a slight abuse of notation (relabeling  $\varepsilon_0$ ) that we have in addition to (2.3) for all  $m \in \mathbb{N}$

$$w_m \equiv 0 \quad \text{on} \quad \left[(-1, 1) \times (1 - \varepsilon_0, 1)\right] \cap \left[(-1, 1) \times (-1, -1 + \varepsilon_0)\right]. \quad (2.7)$$

Given (2.7) we follow exactly the proof of Lemma B.1, where in particular the notion of a convex function  $g$  of a measure (see [22]) is exploited via the representation

$$\int_U g(\nabla w) := \sup_{\kappa \in C_0^\infty(U; \mathbb{R}^n), |\kappa| \leq 1} \left\{ - \int_U w \operatorname{div} \kappa \, dx - \int_U g^*(\kappa) \, dx \right\}$$

and where  $g$  is of linear growth and  $g^*$  denotes the conjugate function (see the definition given in Section 3.2).

**Step 5.** With  $\varepsilon \ll \varepsilon_0$  (i.e. choosing  $m = m(\varepsilon_0)$  sufficiently large) we pass to the limit  $\varepsilon_0 \rightarrow 0$ , which finally proves the lemma.  $\square$

Following the lines of Lemma 2.1 we obtain the convergence of the superlinear part of the energy under consideration.

**Lemma 2.2.** *Given the notation of Lemma 2.1 we suppose that we have (1.5) and (1.6). Moreover, we now assume (1.7).*

*Then the sequence  $\{w_m\}$  of Lemma 2.1 satisfies*

$$\int_{\Omega} f_2(\partial_2 w_m) \, dx \rightarrow \int_{\Omega} f_2(\partial w) \, dx \quad \text{as } m \rightarrow \infty.$$

*Proof.* We start with the first three steps of the proof of Lemma 2.1, in particular we have (2.3), (2.7) and (2.5).

If  $p$  is the exponent given in (1.6), then the strong  $L^p$ -convergence of the sequence  $\{\partial_2 w_m\}$  yields (after passing to a subsequence)

$$\partial_2 w_m \rightarrow \partial_2 w \quad \text{a.e. in } \Omega. \quad (2.8)$$

The first ingredient of the proof follows from our assumption (1.7) and Jensen's inequality, where we recall that in fact only finite sums are considered: for all  $x \in \Omega$  and for all  $m \in \mathbb{N}$  we have

$$\begin{aligned} f_2(\partial_2 w_m) &= f_2\left(\partial_2 \sum_{i=1}^{\infty} \eta_{\varepsilon_i} * (\varphi_i w)\right) = f_2\left(\sum_{i=1}^{\infty} \eta_{\varepsilon_i} * \partial_2(\varphi_i w)\right) \\ &\leq c \sum_{i=1}^{\infty} f_2\left(\eta_{\varepsilon_i} * \partial_2(\varphi_i w)\right) \leq c \sum_{i=1}^{\infty} \eta_{\varepsilon_i} * f_2(\partial_2(\varphi_i w)). \end{aligned} \quad (2.9)$$

We also recall (2.5) which means  $\partial_2 \varphi_i = 0$ . In conclusion, (2.9) shows

$$f_2(\partial_2 w_m) \leq c \sum_{i=1}^{\infty} \eta_{\varepsilon_i} * f_2(\varphi_i \partial_2 w). \quad (2.10)$$

Now we benefit from (2.8) and Egoroff's theorem: for any  $\bar{\varepsilon} > 0$  and for any  $i \in \mathbb{N}$  there exists a measurable set  $A_{i,\bar{\varepsilon}}$  such that

$$|A_i - A_{i,\bar{\varepsilon}}| < \bar{\varepsilon}_i \ll \bar{\varepsilon} \quad \text{and} \quad \partial_2 w_m \rightrightarrows \partial_2 w \quad \text{on } A_{i,\bar{\varepsilon}}. \quad (2.11)$$

A suitable choice of  $\bar{\varepsilon}_i$  is made in (2.13).

With the help of (2.10) one obtains for fixed  $i \in \mathbb{N}$  (note that by the first condition of (2.6), there exist at most three different numbers  $k \in \mathbb{N}$  such that the function  $\eta_{\varepsilon_k} * f_2(\varphi_k \partial_2 w) \not\equiv 0$  on  $A_i$ )

$$\begin{aligned} & \int_{A_i - A_{i,\bar{\varepsilon}}} f_2(\partial_2 w_m) \, dx \\ & \leq c \sum_{k=1}^{\infty} \int_{A_i - A_{i,\bar{\varepsilon}}} \eta_{\varepsilon_k} * f_2(\varphi_k \partial_2 w) \, dx \\ & = c \sum_{k=1}^{\infty} \int_{A_i - A_{i,\bar{\varepsilon}}} \left[ \int \eta_{\varepsilon_k}(x - y) f_2((\varphi_k \partial_2 w)(y)) \, dy \right] \, dx \\ & = c \sum_{k=1}^{\infty} \int_{A_i - A_{i,\bar{\varepsilon}}} \left[ \int_{B_1} \eta(z) f_2((\varphi_k \partial_2 w)(x - \varepsilon_k z)) \, dz \right] \, dx \\ & \leq c \sum_{k=1}^{\infty} \int_{B_1} \eta(z) \left[ \int_{T_{i,\bar{\varepsilon}}^k} f_2((\partial_2 w)(y)) \, dy \right] \, dz, \end{aligned} \quad (2.12)$$

where it is abbreviated ( $|U|$  denoting the Lebesgue measure of  $U \subset \Omega$ )

$$T_{i,\bar{\varepsilon}}^k := \{y = x - \varepsilon_k z : x \in A_i - A_{i,\bar{\varepsilon}}\}, \quad \text{in particular } |T_{i,\bar{\varepsilon}}^k| = |A_i - A_{i,\bar{\varepsilon}}|.$$

Now, since for fixed  $i$  the sum is just taken over three indices, we may choose  $\bar{\varepsilon}_i$  sufficiently small and finally obtain from (2.12) (recalling  $\int_{\Omega} f_2(\partial_2 w) \, dx < \infty$ )

$$\int_{A_i - A_{i,\bar{\varepsilon}}} f_2(\partial_2 w_m) \, dx \leq 2^{-i} \bar{\varepsilon}. \quad (2.13)$$

Decreasing  $\bar{\varepsilon}_i$ , if necessary, it may also be assumed that

$$\int_{A_i - A_{i,\bar{\varepsilon}}} f_2(\partial_2 w) \, dx \leq 2^{-i} \bar{\varepsilon}. \quad (2.14)$$

By (2.3) and (2.7) we note once more that only finite sums have to be considered and recalling (2.11), (2.13) and (2.14) we obtain

$$\begin{aligned}
& \left| \int_{\Omega} f_2(\partial_2 w_m) \, dx - \int_{\Omega} f_2(\partial_2 w) \, dx \right| \\
& \leq \sum_{i=1}^{N_0} \int_{A_i} \left| f_2(\partial_2 w_m) - f_2(\partial_2 w) \right| \, dx \\
& \leq \sum_{i=0}^{N_0} \int_{A_{i,\bar{\varepsilon}}} \left| f_2(\partial_2 w_m) - f_2(\partial_2 w) \right| \, dx + 2\bar{\varepsilon} \\
& \leq \sum_{i=0}^{N_0} \sup_{A_{i,\bar{\varepsilon}}} \left| f_2(\partial_2 w_m) - f_2(\partial_2 w) \right| |A_i| + 2\bar{\varepsilon} \leq 3\bar{\varepsilon}
\end{aligned}$$

provided that  $m > m_0$  with  $m_0$  sufficiently large. This finishes the proof of the lemma.  $\square$

**Remark 2.1.** *Now we can shortly discuss one main difference to the model of Hencky plasticity investigated in [2]. There an approximation lemma is formulated as Theorem 5.3 in Chapter II. The convergence of the deviatoric part in terms of the density  $f$  considered there corresponds to the convergence of the square root in Lemma 2.1 which follows from the linear growth of  $f$  and the notion of a convex function of a measure.*

*Our main difficulty is proving the convergence of the  $f_2$ -energy. In the case of the Hencky plasticity the analogue is just a consequence of considering the intermediate topology (defined in formula (3.37), Chapter II, of [2]) which respects the linear operator  $\operatorname{div} v$ , see also Theorem 3.4 and formula (5.53), Chapter II, of [2].*

Now we define

$$\hat{\Omega} := (-2, 2) \times (-1, 1)$$

and for  $w \in \operatorname{BV}(\Omega)$  we let

$$\hat{w} := \begin{cases} w & \text{on } \Omega, \\ u_0 & \text{on } \hat{\Omega} - \Omega, \end{cases}$$

where  $u_0$  represents a Lipschitz extension of our fixed boundary datum from the space  $W_{\infty}^1(\Omega)$ .

We then have the validity of an approximation result corresponding to Lemma B.2 of [13]. It can be seen as a kind of generalization of Lemma 2.1 and Lemma 2.2 where now  $C^\infty$  is replaced by  $C_0^\infty$ .

**Lemma 2.3.** *Using the above notation suppose that  $w \in \text{BV}(\Omega)$ ,  $\|\partial_2 w\|_{L_A(\Omega)} < \infty$  and that we have (1.4), (1.5), (1.6) and (1.7). Then there exists a sequence  $\{w_m\}$  in  $u_0 + C_0^\infty(\Omega)$  such that passing to the limit  $m \rightarrow \infty$  we have*

$$\begin{aligned} i) \quad & \hat{w}_m \rightarrow \hat{w} \quad \text{in } L^1(\hat{\Omega}), \\ ii) \quad & \int_{\hat{\Omega}} \sqrt{1 + |\nabla \hat{w}_m|^2} \, dx \rightarrow \int_{\hat{\Omega}} \sqrt{1 + |\nabla \hat{w}|^2} \, dx, \\ iii) \quad & \int_{\Omega} f_2(\partial_2 w_m) \, dx \rightarrow \int_{\Omega} f_2(\partial_2 w) \, dx. \end{aligned}$$

### 3 Existence of solutions

There are two approaches towards the existence of generalized solutions to problem (1.10). The first one follows the direct method and leads to the existence of solutions to problem (1.11). This works under quite weak assumptions, for example, the densities  $f_1$  and  $f_2$  need not to be of class  $C^2$ .

The second approach yields the stress tensor as the unique solution of the dual problem and by the stress-strain relation a complete picture of the situation is drawn. However, following the duality approach, we have to suppose that  $f_2$  is given in terms of a  $N$ -function.

#### 3.1 Generalized solutions

**Theorem 3.1.** *Suppose that we have (1.4) - (1.7). Then the relaxed problem (1.11) admits a solution  $\bar{u} \in \mathcal{C}_{\text{BV}}$ .*

*Proof of Theorem 3.1.* We recall that w.l.o.g. we suppose  $n = 2$  and consider a  $K$ -minimizing sequence  $\{u^{(n)}\}$  in the admissible class  $\mathcal{C}_{\text{BV}}$  of comparison functions. After passing to a subsequence we may assume that there exists a function  $\bar{u} \in \text{BV}(\Omega)$  and a function  $v \in L^p(\Omega)$  such that as  $n \rightarrow \infty$

$$u^{(n)} \rightarrow \bar{u} \quad \text{in } L^1(\Omega), \quad \partial_2 u^{(n)} \rightharpoonup v \quad \text{in } L^p(\Omega). \quad (3.1)$$

Here, in the case  $p = 1$ , we refer to the Theorem of De LaValee-Poussin (see, e.g., [15]).

We have for any  $\varphi \in C_0^\infty(\Omega)$

$$\int_{\Omega} u^{(n)} \partial_2 \varphi \, dx = - \int_{\Omega} \partial_2 u^{(n)} \varphi \, dx,$$

hence  $v = \partial_2 \bar{u}$  and since we have for any  $\psi \in C^\infty(\Omega)$

$$\begin{aligned} \int_{\Omega} u^{(n)} \partial_2 \psi \, dx &= - \int_{\Omega} \partial_2 u^{(n)} \psi \, dx + \int_{\partial\Omega} u_0 \psi \nu_2 \, d\mathcal{H}^1, \\ \int_{\Omega} \bar{u} \partial_2 \psi \, dx &= - \int_{\Omega} \partial_2 \bar{u} \psi \, dx + \int_{\partial\Omega} \bar{u} \psi \nu_2 \, d\mathcal{H}^1, \end{aligned}$$

the convergences stated in (3.1) prove  $\bar{u} \in \mathcal{C}_{\text{BV}}$ .

We note that

$$\liminf_{n \rightarrow \infty} K[u^{(n)}] \geq \liminf_{n \rightarrow \infty} K_1[u^{(n)}] + \liminf_{n \rightarrow \infty} E[\partial_2 u^{(n)}].$$

By [23], see also [19], Theorem 5.47, p. 304, we have the lower semicontinuity

$$K_1[\bar{u}] \leq \liminf_{n \rightarrow \infty} K_1[u^{(n)}].$$

Discussing  $E$  we cite Theorem 2.3, p. 18, of [24], hence

$$E[\partial_2 \bar{u}] \leq \liminf_{n \rightarrow \infty} E[\partial_2 u^{(n)}].$$

Since  $\{u^{(n)}\}$  was chosen as a  $K$ -minimizing sequence, the proof of Theorem 3.1 is complete.  $\square$

Now, on account of our approximation Lemma 2.3, we have

**Corollary 3.1.** *With the notation and under the hypotheses of Theorem 3.1 we have*

$$\inf_{w \in \mathcal{C}_{\text{Sob}}} J[w] = \inf_{v \in \mathcal{C}_{\text{BV}}} K[v] = K[\bar{u}].$$

## 3.2 The dual solution

Another approach leading to an analogue of the stress tensor, occurring as basic quantity in problems from mechanics, is to consider the dual problem. As the main references on convex analysis we mention [25] and [26].

Let us assume that we have (1.5) with  $A(|\xi_2|) = f_2(\xi_2)$  for all  $\xi \in \mathbb{R}^{n-1}$ , with  $A$  being of class  $C^1([0, \infty))$  and with  $A$  satisfying (N1)-(N3). In this case we suppose for notational simplicity that  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ ,

$$f(\xi) = f_1(\xi_1) + A(|\xi_2|), \quad \xi \in \mathbb{R}^2. \quad (3.2)$$

As usual we define the conjugate function  $A^*: [0, \infty) \rightarrow [0, \infty)$  by

$$A^*(s) := \max_{t \geq 0} \{st - A(t)\}. \quad (3.3)$$

and note that we have for all  $t \in [0, \infty)$

$$A(t) + A^*(A'(t)) = tA'(t). \quad (3.4)$$

In order to obtain a well-posed dual problem we additionally require

$$A^*(A'(t)) \leq c[A(t) + 1] \quad \text{for all } t \in \mathbb{R}. \quad (3.5)$$

Since

$$f_1^*(s) := \sup_{\xi_1 \in \mathbb{R}} \left\{ s\xi_1 - f_1(\xi_1) \right\},$$

we obtain from the decomposition of  $f$

$$f^*(\xi) = f_1^*(\xi_1) + A^*(|\xi_2|) \quad (3.6)$$

as formula for the conjugate function  $f^*: \mathbb{R}^2 \rightarrow \mathbb{R}$ . The conjugate function  $f^*$  satisfies in correspondence to (3.4)

$$f(\xi) + f^*(Df(\xi)) = \xi \cdot Df(\xi), \quad \xi \in \mathbb{R}^2. \quad (3.7)$$

Given these preliminaries we define the Lagrangian

$$\begin{aligned} l(v, \tau) &:= \int_{\Omega} \tau \cdot \nabla v \, dx - \int_{\Omega} f_1^*(\tau_1) \, dx - \int_{\Omega} A^*(|\tau_2|) \, dx, \\ v &\in u_0 + \overset{\circ}{W}_{1,A}^1(\Omega), \quad \tau \in L^{\infty, A^*}(\Omega; \mathbb{R}^2). \end{aligned} \quad (3.8)$$

In (3.8) we have set

$$L^{\infty, A^*}(\Omega; \mathbb{R}^2) := L^{\infty}(\Omega) \times L_{A^*}(\Omega).$$

With the help of the formula for the conjugate function given in (3.6) we have the representation for the energy  $J$  defined in (1.10)

$$\begin{aligned} J[w] &= \sup_{\varkappa \in L^{\infty, A^*}(\Omega; \mathbb{R}^2)} \left\{ \int_{\Omega} \varkappa \cdot \nabla w \, dx - \int_{\Omega} f_1^*(\varkappa_1) \, dx \right. \\ &\quad \left. - \int_{\Omega} A^*(|\varkappa_2|) \, dx \right\}, \\ &= \sup_{\varkappa \in L^{\infty, A^*}(\Omega; \mathbb{R}^2)} l(w, \varkappa), \quad w \in u_0 + \overset{\circ}{W}_{1,A}^1(\Omega). \end{aligned} \quad (3.9)$$

The dual functional finally is defined via

$$R[\tau] := \inf_{w \in u_0 + \overset{\circ}{W}_{1,A}^1(\Omega)} l(w, \tau), \quad \tau \in L^{\infty, A^*}(\Omega; \mathbb{R}^2). \quad (3.10)$$

This functional leads to the dual problem as the maximizing problem

$$R[\tau] \rightarrow \max \quad \text{in } \tau \in L^{\infty, A^*}(\Omega; \mathbb{R}^2). \quad (3.11)$$

Then we have recalling Theorem 3.1

**Theorem 3.2.** *Suppose that we have our general assumptions (1.4) - (1.7). Moreover, suppose that  $f$  is given in (3.2) with  $A$  satisfying (3.5). Let  $\bar{u}$  denote a solution of the problem (1.11).*

*Then the “stress tensor” defined by*

$$\sigma(x) := Df(\nabla^a \bar{u}) = \left( f'_1(\partial_1^a \bar{u}), A'(|\partial_2 \bar{u}|) \right) \quad (3.12)$$

*is of class  $L^{\infty, A^*}(\Omega; \mathbb{R}^2)$  and maximizes the dual variational problem (3.11) with  $R$  given in (3.10).*

*Proof.* We first note that the boundedness of  $|f'_1|$  and condition (3.5) imply  $\sigma \in L^{\infty, A^*}(\Omega; \mathbb{R}^2)$ .

We then follow an Ansatz similar to Lemma 5.1 of [27]. For any  $v \in u_0 + \overset{\circ}{W}_{1,A}^1(\Omega)$  we have recalling (3.8) and using (3.7)

$$\begin{aligned} l(v, \sigma) &= \int_{\Omega} \nabla v \cdot Df(\nabla^a \bar{u}) \, dx - \int_{\Omega} f^* \left( Df(\nabla^a \bar{u}) \right) \, dx \\ &= \int_{\Omega} Df(\nabla^a \bar{u}) \cdot (\nabla v - \nabla^a \bar{u}) \, dx + \int_{\Omega} f(\nabla^a \bar{u}) \, dx. \end{aligned} \quad (3.13)$$

Now given  $|t| \ll 1$  let  $\bar{u}_t := \bar{u} + t(v - \bar{u}) \in u_0 + \overset{\circ}{W}_{1,A}^1(\Omega)$ . The  $K$ -minimality of  $\bar{u}$  obviously implies

$$\frac{d}{dt} \Big|_{t=0} K[\bar{u}_t] = 0,$$

hence by  $\nabla^s v = 0$

$$\begin{aligned} 0 &= \int_{\Omega} Df(\nabla^a \bar{u}) \cdot (\nabla v - \nabla^a \bar{u}) \, dx + \frac{d}{dt} \Big|_{t=0} \int_{\Omega} f_1^{\infty} \left( \frac{\partial_1^s \bar{u}_t}{|\partial_1^s \bar{u}_t|} \right) d|\partial_1^s \bar{u}_t| \\ &\quad + \frac{d}{dt} \Big|_{t=0} \int_{\partial\Omega} f_1^{\infty} \left( (u_0 - \bar{u}_t) \nu_1 \right) d\mathcal{H}^1, \end{aligned} \quad (3.14)$$

where  $\partial_1^s \bar{u}_t = (1-t)\partial_1^s \bar{u}$ . Now we note that

$$\frac{d}{dt} \Big|_{t=0} \int_{\Omega} f_1^{\infty} \left( \frac{\partial_1^s \bar{u}}{|\partial_1^s \bar{u}|} \right) d((1-t)|\partial_1^s \bar{u}|) = - \int_{\Omega} f_1^{\infty} \left( \frac{\partial_1^s \bar{u}}{|\partial_1^s \bar{u}|} \right) d|\partial_1^s \bar{u}|,$$

and since  $v$  takes the boundary data  $u_0$  on  $\partial\Omega$  we have

$$\frac{d}{dt}\bigg|_{t=0} \int_{\partial\Omega} f_1^\infty((u_0 - \bar{u}_t)\nu_1) d\mathcal{H}^1 = - \int_{\partial\Omega} f_1^\infty((u_0 - \bar{u})\nu_1) d\mathcal{H}^1.$$

Hence, inserting (3.13) in (3.14) we have shown

$$l(v, \sigma) = K[\bar{u}] \quad \text{for any } v \in u_0 + \mathring{W}_{1,A}^1(\Omega)$$

and taking the infimum w.r.t. the comparison function  $v$  we have

$$R[\sigma] \geq K[\bar{u}]. \quad (3.15)$$

We already know from Section 2 that  $\inf J = K[\bar{u}]$  and the representation (3.9) finally yields ( $w \in u_0 + \mathring{W}_{1,A}^1(\Omega)$ )

$$\begin{aligned} J[w] &= \sup_{\varkappa \in L^{\infty, A^*}(\Omega; \mathbb{R}^2)} l(w, \varkappa) \\ &\geq \sup_{\varkappa \in L^{\infty, A^*}(\Omega; \mathbb{R}^2)} \left\{ \inf_{v \in u_0 + \mathring{W}_{1,A}^1(\Omega)} l(v, \varkappa) \right\} \\ &= \sup_{\varkappa \in L^{\infty, A^*}(\Omega; \mathbb{R}^2)} R[\varkappa], \quad \text{i.e.} \\ \inf_{w \in u_0 + \mathring{W}_{1,A}^1(\Omega)} J[w] &\geq \sup_{\varkappa \in L^{\infty, A^*}(\Omega; \mathbb{R}^2)} R[\varkappa]. \end{aligned}$$

This together with (3.15) and Corollary 3.1 proves the theorem.  $\square$

## 4 Higher integrability

### 4.1 Anisotropic behaviour of the superlinear part with $q = 2$

Let us start by recalling Theorem 6.5 of [13] together with our general assumption  $\hat{\mu} < 2$  in (1.13).

**Theorem 4.1.** *Suppose that we are given the assumptions (1.4), (1.5), (1.12) with  $\mu < 2$ ,  $\gamma = 0$  and (1.13) with  $q = 2$ . Let*

$$u^* \in \mathcal{M} := \left\{ u \in \text{BV}(\Omega) : u \text{ is the } L^1\text{-limit of a } J\text{-minimizing sequence from } u_0 + \mathring{W}_1^1(\Omega) \right\}.$$

*Then  $u^*$  is of class  $C^{1,\alpha}(\Omega)$  for any  $0 < \alpha < 1$ . Moreover, the elements of  $\mathcal{M}$  are uniquely determined up to constants.*

**Remark 4.1.** *We again emphasize that the theorem holds without the restriction  $n = 2$ , i.e. the general case  $f_2: \mathbb{R}^{n-1} \rightarrow \mathbb{R}$*

$$\lambda(1 + |\xi_2|)^{-\hat{\mu}}|\eta|^2 \leq D^2 f_2(\xi_2)(\eta, \eta) \leq \Lambda|\eta|^2, \quad \xi_2, \eta \in \mathbb{R}^{n-1},$$

$\lambda, \Lambda > 0$  is included without being explicitly mentioned.

*Main idea of the proof of Theorem 4.1.* The proof of the theorem is based on uniform apriori estimates for the minimizers  $u_\delta$  of the standard quadratic regularization. However, is not immediate that  $\{u_\delta\}$  is a  $J$ -minimizing sequence, if the superlinear part is not generated by a  $N$ -function.

One way to overcome this difficulty is to introduce some kind of local regularized stress tensor  $\sigma_\delta$  without knowing that a solution to a dual problem exists (see Remark 6.15 of [13]).

This, together with the equation  $\operatorname{div} \sigma_\delta = 0$ , leads to the minimality of weak  $L^1$ -cluster points of  $\{u_\delta\}$  via a generalized inf – sup relation. In conclusion, a variational inequality is derived for any  $u^* \in \mathcal{M}$  which provides the regularity of  $u^*$  by Corollary 6.13 of [13] in the  $C^{0,\alpha}$ -regularity of the stress tensor.  $\square$

With our approximation result Corollary 3.1 we identify the elements of  $\mathcal{M}$  with  $K$ -minimizers and observe that the class  $\mathcal{C}_{\text{BV}}$  is defined respecting the condition  $(w - u_0)\nu_2 = 0$   $\mathcal{H}^1$  a.e. on  $\partial\Omega$ .

**Corollary 4.1.** *Given the assumptions of Theorem 4.1, the problem (1.11) is uniquely solvable.*

With the help of Theorem 4.1 we now establish the first regularity result of this paper which is very much in the spirit of [1], i.e.: if we drop the ellipticity condition  $\mu < 2$ , then we still have higher integrability of  $\partial_2 u$ .

**Theorem 4.2.** *Suppose that we have the assumptions of Theorem 4.1 without the requirement  $\mu < 2$ .*

*Then there exists a generalized minimizer  $u \in \mathcal{M}$  such that*

$$\partial_2 u \in L_{\text{loc}}^\chi(\Omega) \quad \text{for any finite } \chi.$$

*Proof of Theorem 4.2.* With the ideas presented in [1] in the linear growth case, the main point is to introduce in the mixed linear-superlinear case a suitable regularization procedure to obtain a sufficiently smooth minimizing sequence.

We fix  $1 < \nu < 2$  and define (compare [10] and related papers)

$$\begin{aligned}\Phi_\nu(t) &:= (\nu - 1) \int_0^t \int_0^s (1 + r)^{-\nu} dr ds, \\ &= t - \frac{1}{2 - \nu} (1 + t)^{2 - \nu} - \frac{1}{\nu - 2}.\end{aligned}$$

Then  $\Phi_\nu$  satisfies (1.12) with  $\mu = \nu$  and  $\gamma = 0$  (in fact we may even choose  $\gamma = -1$ ).

With

$$f_\delta(\xi) = \delta \Phi_\nu(|\xi_1|) + f_1(\xi_1) + f_2(\xi_2)$$

we consider with the obvious meaning of notation the regularized minimization problem ( $\delta \ll 1$ )

$$K_\delta[w] = K_{1,\delta}[w] + E[\partial_2 w] \rightarrow \min \quad \text{in the class } \mathcal{C}_{\text{BV}}. \quad (4.1)$$

By Corollary 4.1 there exists a unique solution denoted by  $u_\delta$  and by Theorem 4.1  $u_\delta$  is of class  $C^{1,\alpha}(\Omega)$ , hence arguing with the Euler equation we also have  $u_\delta \in W_{2,\text{loc}}^2(\Omega)$ .

For the minimizing property of the sequence  $\{u_\delta\}$  we observe for any fixed  $w \in \mathcal{C}_{\text{Sob}}$

$$J[u_\delta] \leq J_\delta[u_\delta] \leq J_\delta[w],$$

hence

$$\liminf_{\delta \rightarrow 0} J[u_\delta] \leq \limsup_{\delta \rightarrow 0} J_\delta[w] = J[w].$$

Passing to a subsequence, if necessary,  $\{u_\delta\}$  obviously is a  $J$ -minimizing sequence.

Now we argue exactly as in [1], proof of Theorem 1, and complete the proof of Theorem 4.2.  $\square$

## 4.2 Higher integrability results in case of superlinear parts of isotropic $p$ -growth

In this subsection we restrict our considerations to the case that the superlinear part is of  $p$ -power growth. This is done in order to simplify our exposition which otherwise would be based on additional parameters. Generalizations are left to the reader.

The main issue is that linear growth conditions are also compatible with  $p$ -growth conditions,  $p > 2$ . Higher integrability results for  $\partial_2 u$  are established in Theorem 4.3 under quite general assumptions.

In Theorem 4.4 we suppose in addition that  $\mu < 2$  and obtain higher integrability of the full gradient, again without restrictions on  $p$ .

Throughout this section we concentrate on the following model case:

Given the general splitting hypothesis (1.4), we suppose that  $f_1$  is of linear growth,

$$a_1|t| - a_2 \leq f_1(t) \leq a_3|t| + a_4, \quad t \in \mathbb{R},$$

$a_2, a_4 \geq 0$ ,  $a_1, a_3 > 0$ , satisfying for some  $\mu > 1$ ,  $\gamma \geq 0$

$$c_1(1 + |t|)^{-\mu} \leq f_1''(t) \leq \bar{c}_1(1 + |t|)^\gamma, \quad t \in \mathbb{R}, \quad (4.2)$$

with constants  $c_1, \bar{c}_1 > 0$ .

We recall the examples discussed in [1] having linear growth and with unbounded second derivative, i.e. the case  $\gamma > 0$  is included in our considerations.

The function  $f_2$  is supposed to be of  $p$ -power growth,  $p > 1$ , in the sense that

$$b_1|t|^p - b_2 \leq f_2(t) \leq b_3|t|^p + b_4, \quad t \in \mathbb{R},$$

$b_2, b_4 \geq 0$ ,  $b_1, b_3 > 0$ , satisfying in addition

$$c_2(1 + |t|)^{p-2} \leq f_2''(t) \leq \bar{c}_2(1 + |t|)^{p-2}, \quad t \in \mathbb{R}, \quad (4.3)$$

with constants  $c_2, \bar{c}_2 > 0$ .

**Theorem 4.3.** *Suppose that the hypotheses of this section are valid.*

i) *If  $\gamma = 0$ , then there exists a generalized minimizer  $u \in \mathcal{M}$  such that*

$$\partial_2 u \in L_{\text{loc}}^\chi(\Omega) \quad \text{for all finite } \chi > p.$$

ii) *If*

$$0 \leq \gamma < \frac{p}{p+1}, \quad (4.4)$$

*then there exists a generalized minimizer  $u \in \mathcal{M}$  such that*

$$\partial_2 u \in L_{\text{loc}}^\chi(\Omega) \quad \text{for some } \chi > p+1.$$

*Proof of Theorem 4.3.* The proof of Theorem 4.3 is carried out in four steps.

*Step 1. Regularization.* We fix some  $0 < \delta < 1$  and define an appropriate variant of regularization: let

$$f_\delta(\xi) := \delta(1 + |\xi_1|^2)^{\frac{p}{2}} + f_1(\xi_1) + f_2(\xi_2), \quad \xi = (\xi_1, \xi_2) \in \mathbb{R}^2.$$

We note that this Ansatz respects the splitting structure of the energy density under consideration.

We then consider the minimization problem

$$J_\delta[w] := \int_{\Omega} f_\delta(\nabla w) \, dx \rightarrow \min \quad \text{in } u_0 + W_0^{1,p}(\Omega) \quad (1.1_\delta)$$

with  $u_\delta$  denoting the unique solution of  $(1.1_\delta)$  satisfying in addition (see the monographs [28], Theorem 8.1, p. 267, and [29], Theorem 5.2, p. 277)

$$u_\delta \in W_{\text{loc}}^{2,2}(\Omega) \cap W_{\text{loc}}^{1,\infty}(\Omega). \quad (4.5)$$

In the situation at hand one may use the dual problem in order to show that the sequence  $\{u_\delta\}$  is  $J$ -minimizing. To this purpose we let

$$\begin{aligned} \tau_\delta &:= \nabla f(\nabla u_\delta), \\ \sigma_\delta &:= \delta X_\delta + \tau_\delta = \nabla f_\delta(\nabla u_\delta), \quad X_\delta := p(1 + |\partial_1 u_\delta|^2)^{\frac{p-2}{2}} \partial_1 u_\delta, \end{aligned}$$

and adapt the arguments of [13], Section 4.1.2: we note that  $\sigma_\delta$  is of class  $W_{\text{loc}}^{1,2}$  (recall (4.5)) satisfying  $\text{div } \sigma_\delta = 0$  and by  $J_\delta[u_\delta] \leq J_\delta[u_0] \leq J_1[u_0]$  we have with a finite constant  $c$  independent of  $\delta$  and not relabelled in the following

$$\delta \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p}{2}} \, dx \leq c, \quad \text{i.e.} \quad \|\delta^{\frac{p-1}{p}} X_\delta\|_{p/(p-1)} \leq c \quad (4.6)$$

$$\int_{\Omega} f(\nabla u_\delta) \, dx \leq c, \quad \|\tau_\delta\|_{p/(p-1)} \leq c. \quad (4.7)$$

Observe that (4.6) implies as  $\delta \rightarrow 0$

$$\delta X_\delta \rightarrow 0 \quad \text{in } L^{\frac{p}{p-1}}(\Omega). \quad (4.8)$$

After passing to subsequences we obtain from (4.7) and (4.8) as  $\delta \rightarrow 0$

$$\tau_\delta, \sigma_\delta \rightharpoonup \sigma \quad \text{in } L^{\frac{p}{p-1}}(\Omega). \quad (4.9)$$

We recall the equation

$$\tau_\delta : \nabla u_\delta - f^*(\tau_\delta) = f(\nabla u_\delta)$$

and arrive at

$$\begin{aligned}
J_\delta[u_\delta] &= \delta \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p}{2}} dx + \int_{\Omega} [\tau_\delta : \nabla u_\delta - f^*(\tau_\delta)] dx \\
&= \delta \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p}{2}} dx + \int_{\Omega} [\sigma_\delta : \nabla u_\delta - f^*(\tau_\delta)] dx \\
&\quad - \delta p \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p-2}{2}} |\nabla u_\delta|^2 dx
\end{aligned}$$

Using  $\operatorname{div} \sigma_\delta = 0$  we obtain

$$\begin{aligned}
J_\delta[u_\delta] &= \delta \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p}{2}} dx + \int_{\Omega} [\sigma_\delta : \nabla u_0 - f^*(\tau_\delta)] dx \\
&\quad - \delta p \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p-2}{2}} |\nabla u_\delta|^2 dx \\
&= \int_{\Omega} [\tau_\delta : \nabla u_0 - f^*(\tau_\delta)] dx + \delta \int_{\Omega} X_\delta : \nabla u_0 dx \\
&\quad + (1-p)\delta \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p}{2}} dx + \delta p \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p-2}{2}} dx.
\end{aligned}$$

Here, as  $\delta \rightarrow 0$ , the second integral (recall (4.8)) and the last integral on the right-hand side converge to 0 and we obtain as  $\delta \rightarrow 0$  (recall (4.9) and upper semicontinuity of  $-f^*$ )

$$\begin{aligned}
R[\mathcal{K}] &\leq \inf_{u \in \mathcal{C}_{\text{Sob}}} J[u] \leq J[u_\delta] \leq J_\delta[u_\delta] \\
&\rightarrow R[\sigma] + (1-p)\delta \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p}{2}} dx.
\end{aligned}$$

Hence we have the minimizing property of the sequence  $\{u_\delta\}$  and additionally

$$\delta \int_{\Omega} (1 + |\partial_1 u_\delta|^2)^{\frac{p}{2}} dx \rightarrow 0 \quad \text{as } \delta \rightarrow 0.$$

*Step 2. Caccioppoli-type inequality.* Proceeding with the proof of Theorem 4.3 we note that by  $u_0 \in W_\infty^1(\Omega)$

$$\sup_{\delta} \|u_\delta\|_{L^\infty(\Omega)} < \infty.$$

As usual we let

$$\Gamma_{i,\delta} := 1 + |\partial_i u_\delta|^2, \quad i = 1, 2.$$

We note that for proving *ii*) of Theorem 4.3, we also need an iterated Caccioppoli-type inequality with negative exponents.

**Proposition 4.1.** *Fix  $l \in \mathbb{N}$  and suppose that  $\eta \in C_0^\infty(\Omega)$ ,  $0 \leq \eta \leq 1$ . Then, given the assumptions at the beginning of Section 4.2, the inequality*

$$\begin{aligned} & \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \partial_2 u_{\delta}) \eta^{2l} \Gamma_{2,\delta}^{\alpha} dx \\ & \leq c \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \eta, \nabla \eta) \eta^{2l-2} \Gamma_{2,\delta}^{\alpha+1} dx \end{aligned} \quad (4.10)$$

*holds for any  $\alpha > -1/2$ , which in particular implies (again for all  $\alpha > -1/2$ )*

$$\begin{aligned} & \int_{\Omega} \eta^{2l} \Gamma_{2,\delta}^{\alpha + \frac{p-2}{2}} |\partial_{22} u_{\delta}|^2 dx \\ & \leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\frac{\alpha+1}{1-\gamma}} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\alpha+1 + \frac{p-2}{2}} dx \right]. \end{aligned} \quad (4.11)$$

*Here and in what follows  $c$  is a finite constant independent of  $\delta$  ( $c = c(l, \alpha)$ ).*

*Proof of Proposition 4.1.* Let us first suppose that  $-1/2 < \alpha \leq 0$ .

We differentiate the Euler equation

$$0 = \int_{\Omega} D f_{\delta}(\nabla u_{\delta}) \cdot \nabla \varphi dx \quad \text{for all } \varphi \in C_0^\infty(\Omega)$$

by inserting  $\varphi = \partial_2 \psi$  as test function, hence

$$0 = \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \psi) dx \quad \text{for all } \psi \in C_0^\infty(\Omega).$$

With the choice

$$\psi := \eta^{2l} \partial_2 u_{\delta} \Gamma_{2,\delta}^{\alpha}$$

we obtain

$$\begin{aligned} & \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \partial_2 u_{\delta}) \eta^{2l} \Gamma_{2,\delta}^{\alpha} dx \\ & = - \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \Gamma_{2,\delta}^{\alpha}) \partial_2 u_{\delta} \eta^{2l} dx \\ & \quad - \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla (\eta^{2l})) \partial_2 u_{\delta} \Gamma_{2,\delta}^{\alpha} dx =: S_1 + S_2, \end{aligned} \quad (4.12)$$

where we note

$$\begin{aligned} |S_1| &= 2|\alpha| \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \partial_2 u_{\delta}) |\partial_2 u_{\delta}|^2 \Gamma_{2,\delta}^{\alpha-1} \eta^{2l} dx \\ &\leq 2|\alpha| \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \partial_2 u_{\delta}) \Gamma_{2,\delta}^{\alpha} \eta^{2l} dx. \end{aligned}$$

Since  $2|\alpha| < 1$ , we may absorb  $|S_1|$  on the left-hand side of (4.12) with the result

$$\int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \partial_2 u_{\delta}) \eta^{2l} \Gamma_{2,\delta}^{\alpha} dx \leq c|S_2|. \quad (4.13)$$

In the case  $\alpha > 0$  we immediately have (4.13) on account of the observation

$$\int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \Gamma_{2,\delta}^{\alpha}) \partial_2 u_{\delta} \eta^{2l} dx \geq 0.$$

The right-hand side of (4.13) is estimated with the help of the Cauchy-Schwarz inequality which gives for any  $\varepsilon > 0$

$$\begin{aligned} & \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \eta) \eta^{2l-1} \Gamma_{2,\delta}^{\alpha} \partial_2 u_{\delta} dx \\ & \leq \varepsilon \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \partial_2 u_{\delta}, \nabla \partial_2 u_{\delta}) \eta^{2l} \Gamma_{2,\delta}^{\alpha} dx \\ & \quad + c(\varepsilon) \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \eta, \nabla \eta) \eta^{2l-2} \Gamma_{2,\delta}^{\alpha} |\partial_2 u_{\delta}|^2 dx \end{aligned}$$

and we have (4.10) by choosing  $\varepsilon$  sufficiently small.

We estimate (recall (4.2) and (4.3))

$$\begin{aligned} & \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \eta, \nabla \eta) \eta^{2l-2} \Gamma_{2,\delta}^{\alpha+1} dx \\ & \leq c \left[ \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{\frac{\gamma}{2}} \Gamma_{2,\delta}^{\alpha+1} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\alpha+1+\frac{p-2}{2}} dx \right]. \quad (4.14) \end{aligned}$$

Let  $\gamma > 0$  and define the numbers  $p_1, p_2$  via

$$1 < p_1 = \frac{1}{\gamma}, \quad p_2 = \frac{1}{1-\gamma}.$$

Using Young's inequality we obtain for the first integral on the right-hand side of (4.14)

$$\begin{aligned} \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{\frac{\gamma}{2}} \Gamma_{2,\delta}^{\alpha+1} dx &= \int_{\Omega} \left[ |\nabla \eta|^2 \eta^{2l-2} \right]^{\frac{1}{p_1}} \Gamma_{1,\delta}^{\frac{\gamma}{2}} \left[ |\nabla \eta|^2 \eta^{2l-2} \right]^{\frac{1}{p_2}} \Gamma_{2,\delta}^{\alpha+1} dx \\ &\leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\frac{\alpha+1}{1-\gamma}} dx \right]. \quad (4.15) \end{aligned}$$

With (4.14) and (4.15) the proof of Proposition 4.1 is finished by observing

$$\begin{aligned}
& \int_{\Omega} \eta^{2l} \Gamma_{2,\delta}^{\alpha + \frac{p-2}{2}} |\partial_{22} u_{\delta}|^2 dx \\
& \leq c \int_{\Omega} f_2''(\partial_2 u_{\delta}) |\partial_{22} u_{\delta}|^2 \eta^{2l} \Gamma_{2,\delta}^{\alpha} dx \\
& \leq c \int_{\Omega} \left[ f_{1,\delta}''(\partial_1 u_{\delta}) |\partial_{12} u_{\delta}|^2 + f_{2,\delta}''(\partial_2 u_{\delta}) |\partial_{22} u_{\delta}|^2 \right] \eta^{2l} \Gamma_{2,\delta}^{\alpha} dx \\
& \leq c \int_{\Omega} D^2 f_{\delta}(\nabla \partial_2 u_{\delta}, \nabla \partial_2 u_{\delta}) \eta^{2l} \Gamma_{2,\delta}^{\alpha} dx. \quad \square
\end{aligned}$$

*Step 3. Main inequality.*

**Proposition 4.2.** *Given the above hypotheses let for some  $\tau_s > 0$ ,  $\tau_{\alpha} > 0$*

$$0 \leq s := \frac{p-2}{2} + \tau_s, \quad \alpha := -\frac{1}{2} + \tau_{\alpha}.$$

*Then for  $l$  sufficiently large and a local constant  $c(\eta, l)$  independent of  $\delta$ , it holds*

$$\begin{aligned}
\int_{\Omega} \eta^{2l} \Gamma_{2,\delta}^{s+1} dx & \leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\frac{\alpha+1}{1-\gamma}} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\alpha+1+\frac{p-2}{2}} dx \right. \\
& \quad \left. + \int_{\Omega} \Gamma_{2,\delta}^{2s-\alpha-\frac{p-2}{2}} \eta^{2l} dx \right]. \quad (4.16)
\end{aligned}$$

*Proof of Proposition 4.2.* Let us first note that on the left-hand side of (4.16) we have  $s+1 = (p/2) + \tau_s$ . Moreover, since  $\|u_{\delta}\|_{L^{\infty}(\Omega)} \leq c$ , we estimate

$$\begin{aligned}
\int_{\Omega} |\partial_2 u_{\delta}|^2 \Gamma_{2,\delta}^s \eta^{2l} dx & = \int_{\Omega} \partial_2 u_{\delta} \partial_2 u_{\delta} \Gamma_{2,\delta}^s \eta^{2l} dx = - \int_{\Omega} u_{\delta} \partial_2 \left[ \partial_2 u_{\delta} \Gamma_{2,\delta}^s \eta^{2l} \right] dx \\
& \leq c \left[ \int_{\Omega} |\partial_{22} u_{\delta}| \Gamma_{2,\delta}^s \eta^{2l} dx + \int_{\Omega} |\partial_2 u_{\delta}| \eta^{2l-1} |\nabla \eta| \Gamma_{2,\delta}^s dx \right. \\
& \quad \left. + \int_{\Omega} \Gamma_{2,\delta}^{s-1} |\partial_2 u_{\delta}|^2 |\partial_{22} u_{\delta}| \eta^{2l} dx \right] \\
& \leq c \left[ \int_{\Omega} |\partial_{22} u_{\delta}| \Gamma_{2,\delta}^s \eta^{2l} dx \right. \\
& \quad \left. + \varepsilon \int_{\Omega} |\partial_2 u_{\delta}|^2 \Gamma_{2,\delta}^s \eta^{2l} dx + c(\varepsilon) \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^s dx \right],
\end{aligned}$$

leading to ( $\varepsilon > 0$  sufficiently small)

$$\int_{\Omega} |\partial_2 u_{\delta}|^2 \Gamma_{2,\delta}^s \eta^{2l} dx \leq c \left[ \int_{\Omega} |\partial_{22} u_{\delta}| \Gamma_{2,\delta}^s \eta^{2l} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^s dx \right]. \quad (4.17)$$

The first term on the right-hand side of (4.17) is estimated with the help of Young's inequality

$$\begin{aligned} \int_{\Omega} |\partial_{22} u_{\delta}| \Gamma_{2,\delta}^s \eta^{2l} dx &= \int_{\Omega} |\partial_{22} u_{\delta}| \Gamma_{2,\delta}^{\frac{\alpha}{2} + \frac{p-2}{4}} \Gamma_{2,\delta}^{s - \frac{\alpha}{2} - \frac{p-2}{4}} \eta^{2l} dx \\ &\leq c \left[ \int_{\Omega} |\partial_{22} u_{\delta}|^2 \Gamma_{2,\delta}^{\alpha + \frac{p-2}{2}} \eta^{2l} dx + \int_{\Omega} \Gamma_{2,\delta}^{2s - \alpha - \frac{p-2}{2}} \eta^{2l} dx \right]. \end{aligned} \quad (4.18)$$

Now (4.11) is applied to the first term on the right-hand side of (4.18) which gives

$$\begin{aligned} \int_{\Omega} |\partial_{22} u_{\delta}| \Gamma_{2,\delta}^s \eta^{2l} dx &\leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\frac{\alpha+1}{1-\gamma}} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\alpha+1 + \frac{p-2}{2}} dx \right. \\ &\quad \left. + \int_{\Omega} \Gamma_{2,\delta}^{2s - \alpha - \frac{p-2}{2}} \eta^{2l} dx \right]. \end{aligned} \quad (4.19)$$

Combining (4.17) and (4.19) we finally obtain

$$\begin{aligned} \int_{\Omega} \Gamma_{2,\delta}^{s+1} \eta^{2l} dx &\leq c \left[ 1 + \int_{\Omega} (\eta^{2l} + |\nabla \eta|^2 \eta^{2l-2}) \Gamma_{2,\delta}^s dx \right. \\ &\quad + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\frac{\alpha+1}{1-\gamma}} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{2,\delta}^{\alpha+1 + \frac{p-2}{2}} dx \\ &\quad \left. + \int_{\Omega} \Gamma_{2,\delta}^{2s - \alpha - \frac{p-2}{2}} \eta^{2l} dx \right]. \end{aligned} \quad (4.20)$$

The first integral on the right-hand side of (4.20) can be absorbed in the left-hand side whenever  $l$  is sufficiently large (compare, e.g., [1], Proof of Proposition 2.2) which completes the proof of the proposition.  $\square$

*Step 4. Conclusion.* To finish the proof of Theorem 4.3 we observe that we may exactly follow the lines of [1], Theorem 1.3, provided that

$$\alpha + 1 + \frac{p-2}{2} = \alpha + \frac{p}{2} < s + 1, \quad 2s - \alpha - \frac{p-2}{2} < s + 1, \quad (4.21)$$

and provided that

$$\alpha + 1 < (s + 1)(1 - \gamma) = s + 1 - (s + 1)\gamma. \quad (4.22)$$

We note that (4.21) is equivalent to

$$\tau_{\alpha} < \tau_s + \frac{1}{2}, \quad \tau_s < \tau_{\alpha} + \frac{1}{2}, \quad \text{i.e. } |\tau_s - \tau_{\alpha}| < \frac{1}{2}. \quad (4.23)$$

Condition (4.22) turns into

$$-\frac{1}{2} + \tau_\alpha < \frac{p-2}{2} + \tau_s - \frac{p+2\tau_s}{2}\gamma,$$

which can be written as

$$\gamma < \frac{p-1+2(\tau_s-\tau_\alpha)}{p+2\tau_s}. \quad (4.24)$$

- i) If  $\gamma = 0$ , then (4.23) and (4.24) are satisfied for any  $\tau_s > 1/4$  and with the choice  $\tau_\alpha = \tau_s - 1/4$ , which implies the first claim of the theorem.
- ii) If we have (4.4) with  $\gamma > 0$ , then we choose  $\tau_\alpha > 0$  sufficiently small and  $\tau_s = \frac{1}{2} + \frac{\tau_\alpha}{2}$ . Then we have (4.23) and (4.24) for  $\tau_\alpha$  sufficiently small on account of our assumption (4.4). We note that with the notation introduced above we have

$$\chi = 2(s+1) = 2 \left[ \frac{p-2}{2} + \tau_s + 1 \right] = p + 2\tau_s.$$

The choice  $\tau_s > 1/2$  completes the proof of Theorem 4.3.  $\square$

In our final theorem on regularity we pose the question, whether an improved ellipticity for the linear part is compatible with  $p$ -growth for arbitrary  $p$  in the following sense: do we have higher integrability of  $\partial_2 u$  and simultaneously of  $\partial_1 u$ ? In fact, in addition to Theorem 4.3 we have

**Theorem 4.4.** *Suppose that we have (1.4) and the first condition of (1.5), i.e. a linear growth condition for  $f_1$ . Moreover, suppose that we have (1.12) with  $2 < \mu$  and  $\gamma = 0$  and let assumption (4.3) hold.*

*Then there exists  $u \in \mathcal{M}$  such that for any  $\kappa < \infty$*

$$\partial_1 u \in L^\kappa(\Omega).$$

*Proof of Theorem 4.4.* Since we already discussed Theorem 4.1 covering the case  $p = 2$  we may suppose that  $p > 2$ . Given the regularization of Theorem 4.3 we need a counterpart for Proposition 4.1 with  $\partial_{22} u_\delta$  replaced by  $\partial_{11} u_\delta$  and  $\Gamma_{2,\delta}$  by  $\Gamma_{1,\delta}$ .

**Proposition 4.3.** *Given the hypotheses of Theorem 4.4 we have for any real numbers  $\chi > 1$ ,  $\alpha > -1/2$*

$$\begin{aligned} & \int_{\Omega} \eta^{2l} \Gamma_{1,\delta}^{\alpha-\frac{\mu}{2}} |\partial_{11} u_\delta|^2 dx \\ & \leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{\alpha+1} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{(\alpha+1)\frac{\chi}{\chi-(p-2)}} dx \right] \end{aligned} \quad (4.25)$$

*Proof of Proposition 4.3.* The counterpart of (4.14) reads as

$$\begin{aligned} & \int_{\Omega} D^2 f_{\delta}(\nabla u_{\delta})(\nabla \eta, \nabla \eta) \eta^{2l-2} \Gamma_{1,\delta}^{\alpha+1} dx \\ & \leq c \left[ \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{\alpha+1} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{\alpha+1} \Gamma_{2,\delta}^{\frac{p-2}{2}} dx \right]. \end{aligned} \quad (4.26)$$

Now we let (recall  $p > 2$ )

$$p_1 = \frac{\chi}{\chi - (p-2)}, \quad 1 < p_2 = \frac{\chi}{p-2}.$$

For the choice of  $\chi$  we have Theorem 4.3, *i*), in mind. Now Young's inequality implies for the second integral on the right-hand side of (4.26)

$$\begin{aligned} \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{\alpha+1} \Gamma_{2,\delta}^{\frac{p-2}{2}} dx &= \int_{\Omega} \left[ |\nabla \eta|^2 \eta^{2l-2} \right]^{\frac{1}{p_1}} \Gamma_{1,\delta}^{\alpha+1} \left[ |\nabla \eta|^2 \eta^{2l-2} \right]^{\frac{1}{p_2}} \Gamma_{2,\delta}^{\frac{p-2}{2}} dx \\ &\leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{(\alpha+1) \frac{\chi}{\chi - (p-2)}} dx \right]. \end{aligned} \quad (4.27)$$

By (4.26) and (4.27) we have (4.25), i.e. the proposition is proved.  $\square$

Now a variant of Proposition 4.1 is needed.

**Proposition 4.4.** *Given the hypotheses of Theorem 4.4 let for some  $\tau_s > 0$ ,  $\tau_{\alpha} > 0$*

$$0 \leq s := -\frac{1}{2} + \tau_s, \quad \alpha := -\frac{1}{2} + \tau_{\alpha}.$$

*Then for  $l$  sufficiently large and a local constant  $c(\eta, l)$  independent of  $\delta$  it holds*

$$\begin{aligned} \int_{\Omega} \eta^{2l} \Gamma_{1,\delta}^{s+1} dx &\leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{\alpha+1} dx \right. \\ &\quad + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{(\alpha+1) \frac{\chi}{\chi - (p-2)}} dx \\ &\quad \left. + \int_{\Omega} \Gamma_{1,\delta}^{2s-\alpha+\frac{\mu}{2}} \eta^{2l} dx \right]. \end{aligned} \quad (4.28)$$

*Proof of Proposition 4.4.* Here we have instead of (4.17), (4.18)

$$\int_{\Omega} |\partial_1 u_{\delta}|^2 \Gamma_{1,\delta}^s \eta^{2l} dx \leq c \left[ \int_{\Omega} |\partial_{11} u_{\delta}| \Gamma_{1,\delta}^s \eta^{2l} dx + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^s dx \right] \quad (4.29)$$

and

$$\begin{aligned} \int_{\Omega} |\partial_{11} u_{\delta}| \Gamma_{1,\delta}^s \eta^{2l} dx &= \int_{\Omega} |\partial_{11} u_{\delta}| \Gamma_{1,\delta}^{\frac{\alpha}{2} - \frac{\mu}{4}} \Gamma_{1,\delta}^{s - \frac{\alpha}{2} + \frac{\mu}{4}} \eta^{2l} dx \\ &\leq c \left[ \int_{\Omega} |\partial_{11} u_{\delta}|^2 \Gamma_{1,\delta}^{\alpha - \frac{\mu}{2}} \eta^{2l} dx + \int_{\Omega} \Gamma_{1,\delta}^{2s - \alpha + \frac{\mu}{2}} \eta^{2l} dx \right]. \end{aligned} \quad (4.30)$$

We then obtain

$$\begin{aligned} \int_{\Omega} \Gamma_{1,\delta}^{s+1} \eta^{2l} dx &\leq c \left[ 1 + \int_{\Omega} (\eta^{2l} + |\nabla \eta|^2 \eta^{2l-2}) \Gamma_{1,\delta}^s dx \right. \\ &\quad + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{\alpha+1} + \int_{\Omega} |\nabla \eta|^2 \eta^{2l-2} \Gamma_{1,\delta}^{(\alpha+1) \frac{\chi}{\chi - (p-2)}} dx \\ &\quad \left. + \int_{\Omega} \Gamma_{1,\delta}^{2s - \alpha + \frac{\mu}{2}} \eta^{2l} dx \right], \end{aligned} \quad (4.31)$$

and (4.31) gives the proposition.  $\square$

Finally we arrive at the following choice of parameters.

$$\alpha + 1 < s + 1 \Leftrightarrow \tau_{\alpha} < \tau_s, \quad (4.32)$$

$$2s - \alpha + \frac{\mu}{2} < s + 1 \Leftrightarrow \tau_s - \tau_{\alpha} < 1 - \frac{\mu}{2}. \quad (4.33)$$

Moreover, the inequality

$$\left[ \frac{1}{2} + \tau_{\alpha} \right] \left[ \frac{\chi}{\chi - (p-2)} \right] < \frac{1}{2} + \tau_s \quad (4.34)$$

needs to be true.

For any given  $\tau_s > 0$  we may choose  $\tau_{\alpha} > 0$  such that (4.32) and (4.33) hold (recalling  $\mu < 2$ ). On account of Theorem 4.3 we then choose  $\chi$  sufficiently large such that (4.34) is satisfied as well. This proves the theorem.  $\square$

**Remark 4.2.** We note that even higher integrability of  $\partial_2 u$  with some  $\chi > p + 1$  (compare Theorem 4.3, ii)) would give some higher integrability of  $\partial_1 u$  by choosing

$$\tau_{\alpha} := \frac{3(2 - \mu) - (p - 2)}{2(p - 2)}$$

whenever  $3(2 - \mu) - (p - 2) > 0$ . This indicates that we also may prove variants of Theorem 4.4 with some weaker hypotheses. This is left to the reader.

Finally we observe that with the higher integrability of the gradient w.r.t. any exponent we have using the idea, e.g., of [13], Section 4.2.4 we obtain as a corollary:

**Corollary 4.2.** *Suppose that we have the hypotheses of Theorem 4.4. Then any weak  $L^1$ -cluster-point of the regularizing sequence  $\{u_\delta\}$  is of class  $C^{1,\alpha}(\Omega)$ .*

## 5 Uniqueness of solutions

The first uniqueness result is already established in Theorem 4.1 and Corollary 4.1.

We now finally discuss the case that the superlinear part of the energy density under consideration is given in terms of a  $N$ -function.

In this case the uniqueness of solutions to the generalized problems can be established without using the results of the previous section.

The reason is the existence of an unique dual solution which, by the duality relation, can be carried over to establish the uniqueness of generalized minimizers.

So let us suppose that we have the situation as described in the hypotheses of Theorem 3.2, in particular

$$\sigma := Df(\nabla^a \bar{u}) = \left( f'_1(\partial_1^a \bar{u}), A'(|\partial_2 \bar{u}|) \right)$$

is a solution of the dual variational problem (3.11) whenever  $\bar{u}$  is a generalized minimizer in the sense of (1.11).

Now, if we have a closer look at the arguments from measure theory leading to Theorem 7 of [21], then we may adapt the proof to the situation at hand and obtain:

**Theorem 5.1.** *Suppose that we have the hypotheses stated in the beginning of Section 4.2 in front of Theorem 4.3. Moreover, suppose that  $u^* \in \mathcal{M}$  is a  $L^1$ -cluster-point of a sequence  $\{u_\delta\}$  from  $u_0 + \overset{\circ}{W}_1^1(\Omega)$  such that for some  $\varkappa \in \mathbb{R}$  the functions*

$$h_\delta := h_\delta^1 + h_\delta^2 := (1 + |\partial_1 u_\delta|^2)^{\frac{\varkappa}{2}} + (1 + |\partial_2 u_\delta|^2)^{\frac{\varkappa}{2}} \quad (5.1)$$

*are of class  $W_{1,\text{loc}}^1(\Omega)$  uniform w.r.t.  $\delta \in (0, 1)$ .*

*Then the dual problem (3.11) is uniquely solvable.*

We emphasize that Theorem 5.1 is valid without any restriction on the exponent  $\mu$  in (1.12). This is just needed for Corollary 5.1 below.

*Main idea of the proof of Theorem 5.1.* Using the notation of [13], Section 2.2, we first note that the unboundedness of  $U := \text{Im}(\nabla f)$  w.r.t. the direction  $\xi_2$  causes no essential changes in the previous arguments.

Given the regularization  $\{u_\delta\}$  of Step 1 in the proof of Theorem 4.3, the main changes concern the proof of (compare p. 22 of [13])

$$\sigma_\delta(x) \rightarrow \sigma(x), \quad \delta |\partial_1 u_\delta(x)|^{p-1} \rightarrow 0, \quad (5.2)$$

on  $\Sigma$  as  $\delta \rightarrow 0$ . In [13], these convergences follow from the uniform local  $W_2^1$ -regularity of  $\sigma_\delta$  which in [13] is a consequence of

$$D^2 f(\xi)(\eta, \eta) \leq (1 + |\xi|)^{-1} |\eta|^2, \quad \xi, \eta \in \mathbb{R}^2. \quad (5.3)$$

Since (5.3) in general is no longer valid, we use (5.1) to get the a.e. convergence of  $\nabla u_\delta$ .  $\square$

We finally have

**Corollary 5.1.** *Suppose that we have the assumptions of Theorem 4.4 with  $p > 2$ . Then we have the uniqueness of generalized solutions of problem (1.11).*

*Proof of Corollary 5.1.* We first establish the uniqueness of the dual solution  $\sigma$ . To this purpose we recall (4.11) and (4.25), i.e. given the regularization introduced in Step 1 of the proof of Theorem 4.3 we have choosing  $\alpha = 0$

$$\begin{aligned} & \int_{\Omega} D^2 f_\delta(\nabla u_\delta)(\nabla \partial_1 u_\delta, \nabla \partial_1 u_\delta) \eta^2 dx \\ & \leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \Gamma_{1,\delta} dx + \int_{\Omega} |\nabla \eta|^2 \Gamma_{1,\delta}^{\frac{x}{x-(p-2)}} dx \right] \end{aligned} \quad (5.4)$$

as well as

$$\begin{aligned} & \int_{\Omega} D^2 f_\delta(\nabla u_\delta)(\nabla \partial_2 u_\delta, \nabla \partial_2 u_\delta) \eta^2 dx \\ & \leq c \left[ 1 + \int_{\Omega} |\nabla \eta|^2 \Gamma_{2,\delta} dx + \int_{\Omega} |\nabla \eta|^2 \Gamma_{2,\delta}^{\frac{p}{2}} dx \right]. \end{aligned} \quad (5.5)$$

Now we compute

$$\begin{aligned}
\int_{\Omega} \eta^2 |\nabla h_{\delta}| \, dx &\leq c \int_{\Omega} \eta^{2l} (1 + |\partial_1 u|^2)^{\frac{\kappa-1}{2}} |\nabla \partial_1 u_{\delta}| \, dx \\
&\quad + c \int_{\Omega} \eta^2 (1 + |\partial_2 u|^2)^{\frac{\kappa-1}{2}} |\nabla \partial_2 u_{\delta}| \, dx \\
&\leq c \int_{\Omega} \eta^2 \left[ \Gamma_{1,\delta}^{-\frac{\mu}{4}} |\partial_1 \partial_1 u_{\delta}| + |\partial_2 \partial_1 u_{\delta}| \right] \Gamma_{1,\delta}^{\frac{\mu}{4} + \frac{\kappa-1}{2}} \, dx \\
&\quad + c \int_{\Omega} \eta^2 \left[ \Gamma_{1,\delta}^{-\frac{\mu}{4}} |\partial_1 \partial_2 u_{\delta}| + |\partial_2 \partial_2 u_{\delta}| \right] \Gamma_{1,\delta}^{\frac{\mu}{4}} \Gamma_{2,\delta}^{\frac{\kappa-1}{2}} \, dx
\end{aligned}$$

which leads to

$$\begin{aligned}
\int_{\Omega} \eta^2 |\nabla h_{\delta}| \, dx &\leq \int_{\Omega} \eta^2 D^2 f_{\delta}(\nabla u_{\delta}) (\nabla \partial_1 u_{\delta}, \nabla \partial_1 u_{\delta}) \, dx \\
&\quad + \int_{\Omega} \eta^2 D^2 f_{\delta}(\nabla u_{\delta}) (\nabla \partial_2 u_{\delta}, \nabla \partial_2 u_{\delta}) \, dx \\
&\quad + \int_{\Omega} \eta^2 \Gamma_{1,\delta}^{t_1} \, dx + \int_{\Omega} \eta^2 \Gamma_{2,\delta}^{t_2} \, dx \tag{5.6}
\end{aligned}$$

with some finite exponents  $t_1$  and  $t_2$ . By (5.4) - (5.6) and Theorem 4.3, *i*), Theorem 4.4 we have that  $h_{\delta}$  is uniformly of class  $W_{1,\text{loc}}^1$  which by Theorem 5.1 implies the uniqueness of the dual solution.

The next step for proving Corollary 5.1 is to use Corollary 4.2 to obtain  $C^{0,\alpha}$ -regularity of the dual solution. Then, as outlined in the proof of Theorem A.9 in [13], a suitable comparison argument w.r.t.  $\sigma$  can be carried out to obtain for any generalized minimizer  $\bar{u} \in \mathcal{C}_{\text{BV}}$

$$\nabla \bar{u} = \nabla f^*(\sigma). \tag{5.7}$$

We note that (5.4) in particular gives

$$\nabla^s \bar{u} = 0.$$

By the uniqueness of  $\sigma$  the uniqueness of generalized minimizers up to an additive constant is established. Finally, on account of  $\bar{u} \in \mathcal{C}_{\text{BV}}$  we have  $(\bar{u} - u_0)\nu_2 = 0$   $\mathcal{H}^1$  a.e. on  $\partial\Omega$  and in conclusion the uniqueness of generalized minimizers.  $\square$

## References

- [1] Bildhauer, M.; Fuchs, M. Splitting type variational problems with linear growth conditions. *arXiv:2004.08169*, 2020.

- [2] Temam, R. *Mathematical Problems in Plasticity*. Dover republication of the edition originally published by Gauthier-Villars, Paris, 1983, Mineola, New York, 2018.
- [3] Beck, L.; Mingione, G. Lipschitz bounds and nonuniform ellipticity. *Comm. Pure Appl. Math.*, LXXIII:944–1034, 2020.
- [4] Giaquinta, M. Growth conditions and regularity, a counterexample. *Manuscripta Math.*, 59(2):245–248, 1987.
- [5] Marcellini, P. Regularity and existence of solutions of elliptic equations with  $p, q$ -growth conditions. *J. Differential Equations*, 90(1):1–30, 1991.
- [6] Frehse, J.; Seregin, G. Regularity of solutions to variational problems of the deformation theory of plasticity with logarithmic hardening. *Amer. Math. Soc. Transl. Ser. 2 (Translation from Proceedings of the St. Petersburg Mathematical Society (1998))*, 193:127–152, 1999.
- [7] Mingione, G.; Siepe, F. Full  $C^{1,\alpha}$ -regularity for minimizers of integral functionals with  $L \log L$ -growth. *Z. Anal. Anwendungen*, 18(4):1083–1100, 1999.
- [8] Giaquinta, M.; Modica, G.; Souček, J. Functionals with linear growth in the calculus of variations. I. *Comm. Math. Univ. Carolinae*, 20:143–156, 1979.
- [9] Giaquinta, M.; Modica, G.; Souček, J. Functionals with linear growth in the calculus of variations. II. *Comm. Math. Univ. Carolinae*, 20:157–172, 1979.
- [10] Bildhauer, M.; Fuchs, M. On a class of variational integrals with linear growth satisfying the condition of  $\mu$ -ellipticity. *Rend. Mat. Appl.*, 22 (2002)(7):249–274, 2003.
- [11] Bildhauer, M. A priori gradient estimates for bounded generalized solutions of a class of variational problems with linear growth. *J. Convex Anal.*, 9(1):117–137, 2002.
- [12] Marcellini, P.; Papi, G. Nonlinear elliptic systems with general growth. *J. Differential Equations*, 221(2):412–443, 2006.
- [13] Bildhauer, M. *Convex variational problems. Linear, nearly linear and anisotropic growth conditions*, volume 1818 of *Lecture Notes in Mathematics*. Springer, Berlin, 2003.
- [14] Rao, M.M.; Ren, Z.D. *Theory of Orlicz spaces.*, volume 146 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, New York, 1991.

- [15] Krasnoselskii, M.A.; Rutickii, J.B. *Convex functions and Orlicz spaces. (Translated from the first Russian edition by Leo F. Boron. P.)*. Noordhoff Ltd., Groningen, 1961.
- [16] Fuchs, M.; Osmolovski, V. Variational integrals on Orlicz-Sobolev spaces. *Z. Anal. Anwendungen*, 17(2):393–415, 1998.
- [17] Fuchs, M.; Li, G. Variational inequalities for energy functionals with nonstandard growth conditions. *Abstr. Appl. Anal.*, 3(1-2):41–64, 1998.
- [18] Adams, R.A. *Sobolev spaces*, volume 65 of *Pure and Applied Mathematics*. Academic Press, New York-London, 1975.
- [19] Ambrosio, L.; Fusco, N.; Pallara, D. *Functions of bounded variation and free discontinuity problems*, volume 224 of *Oxford Mathematical Monographs*. The Clarendon Press, Oxford University Press, New York, 2000.
- [20] Giusti, E. *Minimal surfaces and functions of bounded variation*, volume 80 of *Monographs in Mathematics*. Birkhäuser, Basel, 1984.
- [21] Bildhauer, M. A uniqueness theorem for the dual problem associated to a variational problem with linear growth. *Zap. Nauchn. Sem. St.-Petersburg Otdel. Math. Inst. Steklov (POMI)*, 271:83–91, 2000.
- [22] Demengel, F.; Temam, R. Convex functions of a measure and Applications. *Indiana Univ. Math. J.*, 33(5):673–709, 1984.
- [23] Reshetnyak, Y. Weak convergence of completely additive vector functions on a set. *Sibirsk. Maz. Ž. – English translation: Sib. Math. J.*, 9:1386–1394 – English translation: 1039–1045, 1968.
- [24] Giaquinta, M. *Multiple intergals in the calculus of variations and non-linear elliptic systems*, volume 105 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1983.
- [25] Rockafellar, R.T. *Convex analysis*, volume 28 of *Princeton Mathematical Series*. Princeton University Press, Princeton, N.J., 1970.
- [26] Ekeland, I.; Temam, R. *Convex analysis and variational problems (translated from the French)*, volume 1 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam-Oxford, 1976.
- [27] Fuchs, M.; Tietz, C. Existence of generalized minimizers and of dual solutions for a class of variational problems with linear growth related to image recovery. *J. Math. Sci. (N.Y.)*, 210(4):458–475, 2015.
- [28] Giusti, E. *Direct methods in the calculus of variations*. World Scientific Publishing Co., River Edge, NJ, 2003.

- [29] Ladyzhenskaya, O.A.; Uraltseva, N.N. *Linear and quasilinear elliptic equations. (Translated from the Russian by Scripta Technica, Translation editor: Leon Ehrenpreis)*. Academic Press, New York-London, 1968.

Michael Bildhauer

bibi@math.uni-sb.de

Martin Fuchs

fuchs@math.uni-sb.de

Department of Mathematics

Saarland University

P.O. Box 15 11 50

66041 Saarbrücken

Germany