

Carleman estimates and some inverse problems for the coupled quantitative thermoacoustic equations by boundary data. Part I: Carleman estimates

Yunxia Shang^{*†} and Shumin Li[‡]

Abstract. In this paper, we consider Carleman estimates and inverse problems for the coupled quantitative thermoacoustic equations. In Part I, we establish Carleman estimates for the coupled quantitative thermoacoustic equations by assuming that the coefficients satisfy suitable conditions and taking the usual weight function $\varphi(x, t) = e^{\lambda\psi(x, t)}$, $\psi(x, t) = |x - x_0|^2 - \beta(t - t_0)^2 + \beta t_0^2$ for x in a bounded domain in $\mathbb{R}^n$ with C^3 -boundary and $t \in (0, T)$, where $t_0 = T/2$. We will discuss applications of the Carleman estimates to some inverse problems for the coupled quantitative thermoacoustic equations in the succeeding Part II paper [6].

Keywords. Carleman estimates, thermoacoustic equations, coupled, inverse problems.

2010 Mathematics Subject Classification. 35M10, 35B45, 35R30.

1 Introduction and main results

We investigate the so called quantitative thermoacoustic tomography process (e.g. [1, 5, 23, 25] and their references). According to [5], assuming that the variations in temperature and pressure are weak and neglecting the nonlinear effects, we

^{*}Mathematics & Science college, Shanghai Normal University, 100 Guilin Road, Shanghai, 200234, P.R. China (syx@shnu.edu.cn).

[†]School of Mathematical Sciences, University of Science and Technology of China, 96 Jinzhai Road Baohe, Hefei, Anhui, 230026, P.R. China.

[‡]CAS Wu Wen-Tsun Key Laboratory of Mathematics, University of Science and Technology of China, 96 Jinzhai Road Baohe, Hefei, Anhui, 230026, P.R. China (shuminli@ustc.edu.cn).

obtain the system

$$\begin{cases} \partial_t^2 p - \rho v_s^2 \operatorname{div} \left(\frac{1}{\rho} \nabla p \right) - \Gamma \partial_t \{ \operatorname{div} (\kappa \nabla \theta) \} = \Gamma \partial_t \Pi_a, \\ \partial_t \theta - \frac{1}{\rho C_p} \operatorname{div} (\kappa \nabla \theta) - \frac{\theta_0 \varsigma}{\rho C_p} \partial_t p = \frac{\Pi_a}{\rho C_p}, \end{cases} \quad \text{in } Q, \quad (1.1)$$

for the temperature rise θ and the pressure perturbation p from the equilibrium steady state depending on $(x, t) = (x_1, \dots, x_n, t) \in Q \triangleq \Omega \times (0, T)$. Here $n \in \mathbb{N}^*$ and Ω is a bounded domain in $\mathbb{R}^n$ with the boundary $\partial\Omega \in C^3$. Throughout this paper, we set $\partial_j = \frac{\partial}{\partial x_j}$, $\partial_t = \frac{\partial}{\partial t}$, $\partial_j^2 = \frac{\partial^2}{\partial x_j^2}$, $\partial_j \partial_k = \frac{\partial^2}{\partial x_j \partial x_k}$, $\Delta = \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2}$, $\partial_t^2 = \frac{\partial^2}{\partial t^2}$, $1 \leq j, k \leq n$. We assume that the mass density at steady state ρ , the thermal conductivity κ , the acoustic wave velocity v_s and the isobar specific heat capacity C_p are given strictly positive functions of x and independent of t , the Grüneisen parameter Γ , the background temperature θ_0 and the volume thermal expansivity ς are given non-negative functions of x and independent of t , and the absorbed energy Π_a is an unknown function of x and t .

We set

$$\Theta(x, t) = \partial_t \theta(x, t). \quad (1.2)$$

Differentiating the second equation in (1.1) with respect to t , we obtain

$$\begin{cases} \partial_t^2 p - \rho_1(x) \operatorname{div} (q(x) \nabla p) - \rho_2(x) \operatorname{div} (\kappa(x) \nabla \Theta) = \Gamma(x) \partial_t \Pi_a, \\ \partial_t \Theta - \rho_3(x) \operatorname{div} (\kappa(x) \nabla \Theta) - \rho_4(x) \partial_t^2 p = \rho_3(x) \partial_t \Pi_a, \end{cases} \quad \text{in } Q, \quad (1.3)$$

where $\rho_1(x) = \rho(x) v_s^2(x)$, $q(x) = \frac{1}{\rho(x)}$, $\rho_2(x) = \Gamma(x)$, $\rho_3(x) = \frac{1}{\rho(x) C_p(x)}$, and $\rho_4(x) = \frac{\theta_0(x) \varsigma(x)}{\rho(x) C_p(x)}$. Let $a_1(x) = \rho_1(x) q(x)$, $a_2(x) = \rho_2(x) \kappa(x)$, $a_3(x) = \rho_3(x) \kappa(x)$, and $a_4(x) = \rho_4(x)$.

Therefore, in this paper, we consider Carleman estimates for the following strongly coupled hyperbolic-parabolic system

$$\begin{cases} \partial_t^2 p(x, t) - a_1(x) \Delta p(x, t) - a_2(x) \Delta \Theta(x, t) = f(x, t), \\ \partial_t \Theta(x, t) - a_3(x) \Delta \Theta(x, t) - a_4(x) \partial_t^2 p(x, t) = g(x, t), \end{cases} \quad \text{in } Q, \quad (1.4)$$

where $f(x, t), g(x, t) \in L^2(Q)$, $a_j(x) \in C^2(\overline{\Omega})$ ($j = 1, 2, 3, 4$) are real-valued functions.

In order to state our main results, we introduce more notations. Let $(x \cdot x')$ denote the scalar product in $\mathbb{R}^n$. Let $\nu = \nu(x) = (\nu_1(x), \dots, \nu_n(x))$ denote the outward unit normal vector to $\partial\Omega$ at x . We assume that $\omega \subset \Omega$ is a subdomain of Ω satisfy

$$\overline{\partial\Omega \setminus \partial\omega} \subset \{x \in \partial\Omega \mid ((x - x_0) \cdot \nu(x)) < 0\} \quad (1.5)$$

with some $x_0 = (x_0^1, x_0^2, \dots, x_0^n) \in \mathbb{R}^n \setminus \overline{\Omega}$. Let $T > 0$ be given. We shall establish a Carleman estimate with the measurement in a subboundary layer

$$\Theta(x, t), \quad (x, t) \in Q_\omega \triangleq \omega \times (0, T).$$

We introduce two sets which are concerned with the coefficients $a_j(x)$, $j = 1, 2, 3, 4$:

$$\begin{aligned} \mathcal{U} = \mathcal{U}_{\sigma_0, \sigma_1, M_0, M_1, M_2} = & \left\{ (a_1(x), a_2(x), a_3(x), a_4(x)) \in (C^2(\overline{\Omega}))^4 \mid \right. \\ & a_1(x) \geq \sigma_1, \ a_3(x) \geq \sigma_1, \ a_2(x) \geq 0, \ a_4(x) \geq 0, \ \forall x \in \overline{\Omega}, \\ & \|a_j\|_{C(\overline{\Omega})} \leq M_0, \ \|a_j\|_{C^1(\overline{\Omega})} \leq M_1, \ \|a_j\|_{C^2(\overline{\Omega})} \leq M_2, \ j = 1, 2, 3, 4, \\ & \left. 3a_1 - 2((x - x_0) \cdot \nabla a_1) + \frac{a_1}{(1 + \frac{a_2 a_4}{a_3})} \left(2(x - x_0) \cdot \nabla \left(\frac{a_2 a_4}{a_3} \right) \right) \geq \sigma_0 \right\} \end{aligned} \quad (1.6)$$

where the constants $M_0 > 1$, $M_1 > 0$, $M_2 > 0$, $\sigma_0 > 0$, $M_0 > \sigma_1 > 0$ are given. Denote

$$m = \inf_{x \in \overline{\Omega}} |x - x_0|^2, \quad M = \sup_{x \in \overline{\Omega}} |x - x_0|^2, \quad \text{and} \quad \mathcal{D} = \sqrt{M - m}. \quad (1.7)$$

We assume that $(a_1(x), a_2(x), a_3(x), a_4(x)) \in \mathcal{U} = \mathcal{U}_{\sigma_0, \sigma_1, M_0, M_1, M_2}$. Denote

$$\begin{aligned} \alpha_1 &= 228nM_0^3M_1M^{\frac{1}{2}} + 20M_0^4 - 8\sigma_1^2, \quad \alpha_2 = \left(132nM_1M^{\frac{1}{2}} + 16M_0 \right) M_0^3, \\ \alpha_3 &= \min \left\{ \frac{\sigma_0\sigma_1}{\left(\frac{2M_0^4\alpha_1}{\sigma_1^2} + \alpha_2 \right)}, \frac{\sigma_1}{8M}, \frac{\sigma_1^4}{16M_0^8} \right\}, \\ \alpha_4 &= 6M_0^2\alpha_3 + \frac{2M_0^2}{\sigma_1}, \quad \alpha_5 = 2M_0^2\alpha_3 + 2 \left(1 + \frac{M_0^2}{\sigma_1} + \frac{3M_0^3}{\sigma_1^2} \right) M_1 + \frac{2M_0^2}{\sigma_1}, \\ \alpha_6 &= \frac{4M_0^3\alpha_3}{\sigma_1} + 2 \left(1 + \frac{M_0^2}{\sigma_1} + \frac{3M_0^3}{\sigma_1^2} \right) nM_1, \\ \alpha_7 &= \frac{16M_0^8}{\sigma_1^7} \left(1 + \frac{M_0^2}{\sigma_1} \right)^2, \quad \alpha_8 = 2 \left(1 + \frac{M_0^2}{\sigma_1} \right)^2, \\ \alpha_9 &= \frac{-\mathcal{D}\alpha_6 + \sqrt{\mathcal{D}^2\alpha_6^2 + \sigma_1(\mathcal{D}^2\alpha_7 + \alpha_8)}}{4(\mathcal{D}^2\alpha_7 + \alpha_8)}. \end{aligned} \quad (1.8)$$

We choose $\beta > 0$ such that

$$0 < \beta < \min \left\{ \alpha_9^2, \frac{\alpha_1^2\alpha_3^2}{16\alpha_4^2\mathcal{D}^2}, \frac{\sigma_0^2\sigma_1^2}{16\alpha_5^2\mathcal{D}^2}, \frac{m^2\sigma_1^3}{2M_0(\sigma_1 + M_0^2)\mathcal{D}^2} \right\}. \quad (1.9)$$

Denote $t_0 = \frac{T}{2}$. We will prove a Carleman estimate for (1.4) with the exponential weight function $e^{2s\varphi}$ where

$$\varphi(x, t) = e^{\lambda\psi(x, t)}, \quad \psi(x, t) = |x - x_0|^2 - \beta(t - t_0)^2 + \beta t_0^2, \quad \forall (x, t) \in Q, \quad (1.10)$$

and $\lambda > 0$ is a suitably large constant.

We set $\nabla = (\partial_1, \dots, \partial_n)$, $\nabla_{x,t} = (\partial_1, \dots, \partial_n, \partial_t)$, $|\nabla w|^2 = \sum_{k=1}^n |\partial_k w|^2$, $|\nabla_{x,t} w|^2 = |\nabla w|^2 + |\partial_t w|^2$, and so on. $L^2(Q)$, $H^2(\Omega)$, etc. denote usual Sobolev spaces. We further set

$$\begin{aligned} \mathcal{H}^{2,1}(Q) &= \{u \in L^2(Q); \partial_j u, \partial_j^2 u, \partial_j \partial_k u, \partial_t u \in L^2(Q), j, k = 1, \dots, n\}, \\ \mathcal{H}^{2,2}(Q) &= \{u \in L^2(Q); \partial_j u, \partial_j^2 u, \partial_j \partial_k u, \partial_t u, \partial_t^2 u \in L^2(Q), j, k = 1, \dots, n\}, \end{aligned}$$

and $\mathcal{W} = \mathcal{H}^{2,1}(Q) \times \mathcal{H}^{2,2}(Q)$.

Theorem 1.1. *Let $(\Theta, p) \in \mathcal{W}$ satisfy (1.4) and*

$$\Theta(x, t) = 0, \quad p(x, t) = 0, \quad (x, t) \in \Sigma \triangleq \partial\Omega \times (0, T) \quad (1.11)$$

$$\Theta(x, 0) = \Theta(x, T) = 0, \quad \partial_t^j p(x, 0) = \partial_t^j p(x, T) = 0, \quad x \in \overline{\Omega}, \quad j = 0, 1. \quad (1.12)$$

We assume that $(a_1, a_2, a_3, a_4) \in \mathcal{U}$, and that (1.9) holds. Then there exists a constant $\eta(\beta) > 0$ such that for any $T \in \left(0, \frac{2(\mathcal{D}+\eta)}{\sqrt{\beta}}\right)$, there exists a constant $\lambda_0 > 0$ such that for all $\lambda > \lambda_0$, there exist constants $s_0(\lambda) > 0$ and $K_1 = K_1(s_0, \lambda_0, \beta, \Omega, T, m, M, M_0, M_1, M_2, \sigma_0, \sigma_1) > 0$ such that

$$\begin{aligned} & \int_Q \left(s^3 \lambda^4 \varphi^3 \Theta^2 + \frac{1}{s\varphi} |\partial_t \Theta|^2 + s\lambda\varphi |\nabla \Theta|^2 + s^3 \lambda^4 \varphi^3 p^2 + s\lambda\varphi |\nabla_{x,t} p|^2 \right) e^{2s\varphi} dx dt \\ & \leq K_1 \int_Q (f^2 + g^2) e^{2s\varphi} dx dt \\ & + K_1 \int_{Q_\omega} \left\{ s^3 \lambda^4 \varphi^3 (\Theta^2 + p^2) + \frac{1}{s\varphi} |\partial_t \Theta|^2 + s^3 \lambda^3 \varphi^3 |\partial_t p|^2 \right\} e^{2s\varphi} dx dt, \end{aligned} \quad (1.13)$$

for all $s \geq s_0$.

Let $\Gamma_0 \subset \partial\Omega$ satisfy

$$\overline{\partial\Omega \setminus \Gamma_0} \subset \{x \in \partial\Omega \mid ((x - x_0) \cdot \nu(x)) < 0\} \quad (1.14)$$

Let $\frac{\partial u}{\partial \nu}$ denote the normal derivative: $\frac{\partial u}{\partial \nu}(x) = \sum_{j=1}^n \nu_j(x) \partial_j u(x)$, $x \in \partial\Omega$. The following Carleman estimate plays an important role in the proof of Theorem 1.1:

Theorem 1.2. *Let $(\Theta, p) \in \mathcal{W}$ satisfy (1.4), (1.11) and (1.12). We assume that $(a_1, a_2, a_3, a_4) \in \mathcal{U}$, and that (1.9) holds. Then there exists a constant $\eta(\beta) > 0$ such that for any $T \in \left(0, \frac{2(\mathcal{D}+\eta)}{\sqrt{\beta}}\right)$, there exists a constant $\lambda_0 > 0$ such that for all $\lambda > \lambda_0$, there exist constants $s_0(\lambda) > 0$ and $K_2 = K_2(s_0, \lambda_0, \beta, \Omega, T, m, M, M_0, M_1, M_2, \sigma_0, \sigma_1, \epsilon) > 0$ such that*

$$\begin{aligned} & \int_Q \left(s^3 \lambda^4 \varphi^3 \Theta^2 + \frac{1}{s\varphi} |\partial_t \Theta|^2 + s\lambda \varphi |\nabla \Theta|^2 + s^3 \lambda^4 \varphi^3 p^2 + s\lambda \varphi |\nabla_{x,t} p|^2 \right) e^{2s\varphi} dx dt \\ & \leq K_2 \int_Q (f^2 + g^2) e^{2s\varphi} dx dt + K_2 s \lambda \int_{\Gamma_0 \times (0, T)} \varphi \left(\left| \frac{\partial \Theta}{\partial \nu} \right|^2 + \left| \frac{\partial p}{\partial \nu} \right|^2 \right) d\sigma dt, \quad (1.15) \end{aligned}$$

for all $s \geq s_0$.

The Carleman estimate was first introduced by Carleman [3] in 1939 for the purpose of proving uniqueness in the general Cauchy problem of first-order systems in two dimensions with simple characteristics. After that the Carleman estimate and applications were greatly developed. We refer to Bellassoued and Yamamoto [2], Chae, Imanuvilov and Kim [4], Egorov [7], Eller and Isakov [8], Fursikov and Imanuvilov [9], Hörmander [10, 11], Imanuvilov [12], Imanuvilov and Yamamoto [13], Isakov [14]–[18], Klibanov [19], Klibanov and Timonov [20], Lavrent'ev, Romanov and Shishatskii [21], Li [22], Romanov [24], Tataru [26], Trèves [27], Yamamoto [28], Yuan and Yamamoto [29], and related references in them. In this references, Carleman estimates are proved by general methods or direct methods.

In this paper, we concentrate on the proof of Carleman estimates for (1.4), that is, Theorem 1.1 and 1.2, by a direct method. We will discuss applications of Theorem 1.1 to several inverse problems in the domain of thermoacoustic tomography (TAT) in the succeeding Part II paper [6]. To our best knowledge, inverse problems in the domain of TAT are very few addressed until now because the models of the literature do not consider the heating process.

The paper consists of three sections. In Section 2, we will give a direct derivation of Theorem 1.2 by apply the argument in e.g. [20, 21, 24, 28]. In Section 3, we will derive Theorem 1.1 by applying Theorem 1.2 and the argument in [12].

2 Proof of Theorem 1.2

By a usual density argument, we can assume that $(\Theta, p) \in C^\infty(Q) \times C^\infty(Q)$. Let $\lambda > 1$ and $s > 1$. Eliminating $\partial_t^2 p$ from the two equations in (1.4), we have

$$\partial_t \Theta - a_5 \Delta \Theta - a_6 \Delta p = a_4 f + g, \quad (2.1)$$

where

$$a_5 = a_3 + a_2 a_4 \quad \text{and} \quad a_6 = a_1 a_4. \quad (2.2)$$

Denote

$$L_0(\Theta, p) = \partial_t \Theta - a_5 \Delta \Theta - a_6 \Delta p. \quad (2.3)$$

We set $(\vartheta, w) = e^{s\varphi}(\Theta, p)$, $\mathcal{L}(\vartheta, w) = e^{s\varphi} L_0(e^{-s\varphi}(\vartheta, w))$. By (1.10), we have

$$\begin{aligned} \partial_t \psi &= -2\beta(t - t_0), \quad \partial_t^2 \psi = -2\beta, \quad \partial_j \psi = 2(x_j - x_0^j), \quad \partial_j \partial_k \psi = 2\delta_{jk}, \\ \partial_t \varphi &= \lambda \varphi(\partial_t \psi), \quad \partial_t^2 \varphi = \lambda \varphi(\partial_t^2 \psi) + \lambda^2 \varphi(\partial_t \psi)^2, \quad \partial_j \varphi = \lambda \varphi(\partial_j \psi), \\ \partial_j \partial_k \varphi &= \lambda \varphi(\partial_j \partial_k \psi) + \lambda^2 \varphi(\partial_j \psi)(\partial_k \psi), \quad \text{for all } j, k = 1, \dots, n. \end{aligned} \quad (2.4)$$

By direct calculations, we have

$$\begin{aligned} e^{s\varphi}(a_4 f + g) &= \mathcal{L}(\vartheta, w) = e^{s\varphi} \{ \partial_t(e^{-s\varphi} \vartheta) - a_5 \Delta(e^{-s\varphi} \vartheta) - a_6 \Delta(e^{-s\varphi} w) \} \\ &= -a_5 \Delta \vartheta - a_6 \Delta w - s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 a_5 \vartheta - s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 a_6 w + \partial_t \vartheta \\ &\quad + 2s \lambda \varphi (\nabla \psi \cdot (a_5 \nabla \vartheta + a_6 \nabla w)) + A(\vartheta, w), \quad \text{in } Q, \end{aligned}$$

where

$$A(\vartheta, w) = s \lambda^2 \varphi |\nabla \psi|^2 (a_5 \vartheta + a_6 w) + 2ns \lambda \varphi (a_5 \vartheta + a_6 w) - s \lambda \varphi (\partial_t \psi) \vartheta. \quad (2.5)$$

Then taking into consideration the orders of (s, λ, φ) , we divide $\mathcal{L}(\vartheta, w)$ as follows:

$$\mathcal{L}(\vartheta, w) = L_1(\vartheta, w) + L_2(\vartheta, w), \quad (2.6)$$

where

$$\begin{aligned} L_1(\vartheta, w) &= -a_5 \Delta \vartheta - a_6 \Delta w - s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 a_5 \vartheta - s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 a_6 w, \\ L_2(\vartheta, w) &= \partial_t \vartheta + 2s \lambda \varphi (\nabla \psi \cdot (a_5 \nabla \vartheta + a_6 \nabla w)) + A(\vartheta, w). \end{aligned} \quad (2.7)$$

Therefore we have

$$\begin{aligned} \|(a_4 f + g) e^{s\varphi}\|_{L^2(Q)}^2 &= \|\mathcal{L}(\vartheta, w)\|_{L^2(Q)}^2 = \|L_1(\vartheta, w) + L_2(\vartheta, w)\|_{L^2(Q)}^2 \\ &= \|L_1(\vartheta, w)\|_{L^2(Q)}^2 + 2 \int_Q L_1(\vartheta, w) L_2(\vartheta, w) dx dt + \|L_2(\vartheta, w)\|_{L^2(Q)}^2. \end{aligned} \quad (2.8)$$

Direct calculations yeild

$$\begin{aligned} L_1(\vartheta, w) L_2(\vartheta, w) &= -a_5 (\Delta \vartheta) (\partial_t \vartheta) - 2a_5 s \lambda \varphi (\Delta \vartheta) (\nabla \psi \cdot (a_5 \nabla \vartheta + a_6 \nabla w)) \\ &\quad - a_5 (\Delta \vartheta) A(\vartheta, w) - a_6 (\Delta w) (\partial_t \vartheta) - 2s \lambda \varphi a_6 (\Delta w) (\nabla \psi \cdot (a_5 \nabla \vartheta + a_6 \nabla w)) \\ &\quad - a_6 (\Delta w) A(\vartheta, w) - s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 a_5 \vartheta (\partial_t \vartheta) \\ &\quad - 2s^3 \lambda^3 \varphi^3 |\nabla \psi|^2 a_5 (\nabla \psi \cdot (a_5 \nabla \vartheta + a_6 \nabla w)) \vartheta - s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 a_5 \vartheta A(\vartheta, w) \end{aligned}$$

$$\begin{aligned}
& -s^2\lambda^2\varphi^2|\nabla\psi|^2a_6w(\partial_t\vartheta) - 2s^3\lambda^3\varphi^3|\nabla\psi|^2a_6(\nabla\psi\cdot(a_5\nabla\vartheta+a_6\nabla w))w \\
& -s^2\lambda^2\varphi^2|\nabla\psi|^2a_6wA(\vartheta,w) \triangleq \sum_{k=1}^{12} J_k.
\end{aligned} \tag{2.9}$$

We integrate J_k , $k = 1, 2, \dots, 12$ over Q . We calculate them by applying the integration by parts and apply the conditions (1.11), (1.12), and $(a_1, a_2, a_3, a_4) \in \mathcal{U}$.

$$\begin{aligned}
\int_Q J_1 dx dt &= - \int_Q a_5 (\Delta\vartheta) (\partial_t\vartheta) dx dt \\
&= \int_Q (\nabla a_5 \cdot \nabla\vartheta) (\partial_t\vartheta) dx dt + \int_Q a_5 \partial_t \left(\frac{1}{2} |\nabla\vartheta|^2 \right) dx dt \\
&= \int_Q (\nabla a_5 \cdot \nabla\vartheta) (\partial_t\vartheta) dx dt.
\end{aligned} \tag{2.10}$$

$$\begin{aligned}
\int_Q J_2 dx dt &= -2s\lambda \int_Q \varphi (\Delta\vartheta) (\nabla\psi \cdot (a_5^2\nabla\vartheta + a_5a_6\nabla w)) dx dt \\
&= -2s\lambda \int_Q \varphi a_5^2 (\Delta\vartheta) (\nabla\psi \cdot \nabla\vartheta) dx dt - 2s\lambda \int_Q \varphi a_5a_6 (\Delta\vartheta) (\nabla\psi \cdot \nabla w) dx dt \\
&= -2s\lambda \int_{\Sigma} \varphi a_5^2 (\nabla\psi \cdot \nabla\vartheta) (\nabla\vartheta \cdot \nu) d\sigma dt + 2s\lambda^2 \int_Q \varphi a_5^2 (\nabla\psi \cdot \nabla\vartheta)^2 dx dt \\
&\quad + 4s\lambda \int_Q \varphi a_5 (\nabla a_5 \cdot \nabla\vartheta) (\nabla\psi \cdot \nabla\vartheta) dx dt + 4s\lambda \int_Q \varphi a_5^2 |\nabla\vartheta|^2 dx dt \\
&\quad + s\lambda \int_Q \varphi a_5^2 (\nabla\psi \cdot \nabla (|\nabla\vartheta|^2)) dx dt - 2s\lambda \int_{\Sigma} \varphi a_5a_6 (\nabla\psi \cdot \nabla w) (\nabla\vartheta \cdot \nu) d\sigma dt \\
&\quad + 2s\lambda^2 \int_Q \varphi a_5a_6 (\nabla\psi \cdot \nabla\vartheta) (\nabla\psi \cdot \nabla w) dx dt \\
&\quad + 2s\lambda \int_Q \varphi (\nabla (a_5a_6) \cdot \nabla\vartheta) (\nabla\psi \cdot \nabla w) dx dt + 4s\lambda \int_Q \varphi a_5a_6 (\nabla\vartheta \cdot \nabla w) dx dt \\
&\quad + 2s\lambda \int_Q \varphi a_5a_6 \sum_{j,k=1}^n (\partial_j\psi) (\partial_k\vartheta) (\partial_j\partial_kw) dx dt.
\end{aligned}$$

Calculating the fifth term by the integration by parts, we have

$$\begin{aligned}
(\text{the fifth term}) &= s\lambda \int_{\Sigma} \varphi a_5^2 (\nabla\psi \cdot \nu) |\nabla\vartheta|^2 d\sigma dt - s\lambda^2 \int_Q \varphi |\nabla\psi|^2 a_5^2 |\nabla\vartheta|^2 dx dt \\
&\quad - 2s\lambda \int_Q \varphi a_5 (\nabla\psi \cdot \nabla a_5) |\nabla\vartheta|^2 dx dt - s\lambda \int_Q \varphi (\Delta\psi) a_5^2 |\nabla\vartheta|^2 dx dt.
\end{aligned}$$

We note that $\Delta\psi = 2n$ on $\bar{\Omega}$. By (1.11), we have $\vartheta = 0$ and $w = 0$ on Σ , so that we have $\nabla\vartheta = \frac{\partial\vartheta}{\partial\nu}\nu$ and $\nabla w = \frac{\partial w}{\partial\nu}\nu$ on Σ . Therefore we have

$$\begin{aligned}
\int_Q J_2 dxdt &= -s\lambda \int_{\Sigma} \varphi (\nabla\psi \cdot \nu) a_5^2 \left| \frac{\partial\vartheta}{\partial\nu} \right|^2 d\sigma dt - 2s\lambda \int_{\Sigma} \varphi (\nabla\psi \cdot \nu) a_5 a_6 \frac{\partial\vartheta}{\partial\nu} \frac{\partial w}{\partial\nu} d\sigma dt \\
&+ 2s\lambda^2 \int_Q \varphi a_5^2 (\nabla\psi \cdot \nabla\vartheta)^2 dxdt + 4s\lambda \int_Q \varphi a_5 (\nabla a_5 \cdot \nabla\vartheta) (\nabla\psi \cdot \nabla\vartheta) dxdt \\
&+ 4s\lambda \int_Q \varphi a_5^2 |\nabla\vartheta|^2 dxdt - s\lambda^2 \int_Q \varphi |\nabla\psi|^2 a_5^2 |\nabla\vartheta|^2 dxdt \\
&- 2s\lambda \int_Q \varphi a_5 (\nabla\psi \cdot \nabla a_5) |\nabla\vartheta|^2 dxdt - 2ns\lambda \int_Q \varphi a_5^2 |\nabla\vartheta|^2 dxdt \\
&+ 2s\lambda^2 \int_Q \varphi a_5 a_6 (\nabla\psi \cdot \nabla\vartheta) (\nabla\psi \cdot \nabla w) dxdt \\
&+ 2s\lambda \int_Q \varphi (\nabla(a_5 a_6) \cdot \nabla\vartheta) (\nabla\psi \cdot \nabla w) dxdt + 4s\lambda \int_Q \varphi a_5 a_6 (\nabla\vartheta \cdot \nabla w) dxdt \\
&+ 2s\lambda \int_Q \varphi a_5 a_6 \sum_{j,k=1}^n (\partial_j\psi) (\partial_k\vartheta) (\partial_j\partial_k w) dxdt. \tag{2.11}
\end{aligned}$$

$$\begin{aligned}
\int_Q J_3 dxdt &= - \int_Q a_5 (\Delta\vartheta) A(\vartheta, w) dxdt \\
&= -s\lambda^2 \int_Q \varphi |\nabla\psi|^2 a_5^2 (\Delta\vartheta) \vartheta dxdt - s\lambda^2 \int_Q \varphi |\nabla\psi|^2 a_5 a_6 (\Delta\vartheta) w dxdt \\
&- 2ns\lambda \int_Q \varphi a_5^2 (\Delta\vartheta) \vartheta dxdt - 2ns\lambda \int_Q \varphi a_5 a_6 (\Delta\vartheta) w dxdt \\
&+ s\lambda \int_Q \varphi (\partial_t\psi) a_5 (\Delta\vartheta) \vartheta dxdt \\
&= s\lambda^3 \int_Q \varphi |\nabla\psi|^2 a_5^2 (\nabla\psi \cdot \nabla\vartheta) \vartheta dxdt + s\lambda^2 \int_Q \varphi (\nabla(a_5^2 |\nabla\psi|^2) \cdot \nabla\vartheta) \vartheta dxdt \\
&+ s\lambda^2 \int_Q \varphi |\nabla\psi|^2 a_5^2 |\nabla\vartheta|^2 dxdt + s\lambda^3 \int_Q \varphi |\nabla\psi|^2 a_5 a_6 (\nabla\psi \cdot \nabla\vartheta) w dxdt \\
&+ s\lambda^2 \int_Q \varphi (\nabla(|\nabla\psi|^2 a_5 a_6) \cdot \nabla\vartheta) w dxdt \\
&+ s\lambda^2 \int_Q \varphi |\nabla\psi|^2 a_5 a_6 (\nabla\vartheta \cdot \nabla w) dxdt + 2ns\lambda^2 \int_Q \varphi a_5^2 (\nabla\psi \cdot \nabla\vartheta) \vartheta dxdt \\
&+ 4ns\lambda \int_Q \varphi a_5 (\nabla a_5 \cdot \nabla\vartheta) \vartheta dxdt + 2ns\lambda \int_Q \varphi a_5^2 |\nabla\vartheta|^2 dxdt
\end{aligned}$$

$$\begin{aligned}
& + 2ns\lambda^2 \int_Q \varphi a_5 a_6 (\nabla \psi \cdot \nabla \vartheta) w dx dt + 2ns\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla \vartheta) w dx dt \\
& + 2ns\lambda \int_Q \varphi a_5 a_6 (\nabla \vartheta \cdot \nabla w) dx dt - s\lambda^2 \int_Q \varphi (\partial_t \psi) a_5 (\nabla \psi \cdot \nabla \vartheta) \vartheta dx dt \\
& - s\lambda \int_Q \varphi (\partial_t \psi) (\nabla a_5 \cdot \nabla \vartheta) \vartheta dx dt - s\lambda \int_Q \varphi (\partial_t \psi) a_5 |\nabla \vartheta|^2 dx dt \\
& \geq s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_5^2 |\nabla \vartheta|^2 dx dt + s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_5 a_6 (\nabla \vartheta \cdot \nabla w) dx dt \\
& + 2ns\lambda \int_Q \varphi a_5^2 |\nabla \vartheta|^2 dx dt + 2ns\lambda \int_Q \varphi a_5 a_6 (\nabla \vartheta \cdot \nabla w) dx dt \\
& - s\lambda \int_Q \varphi (\partial_t \psi) a_5 |\nabla \vartheta|^2 dx dt - C_1 s \lambda^3 \int_Q \varphi |\nabla \vartheta| (|\vartheta| + |w|) dx dt. \tag{2.12}
\end{aligned}$$

Here and henceforth, C_k ($k = 1, 2, \dots$) denote generic positive constants which may depend on $n, \Omega, T, \beta, m, M, M_j$ ($j = 0, 1, 2$), $\sigma_0, \sigma_1, s_0, \lambda_0$, but are independent of s and λ . Integrating by parts several times, we have

$$\begin{aligned}
\int_Q J_4 dx dt & = - \int_Q a_6 (\Delta w) (\partial_t \vartheta) dx dt \\
& = \int_Q (\partial_t \vartheta) (\nabla a_6 \cdot \nabla w) dx dt + \int_Q a_6 (\nabla w \cdot \nabla (\partial_t \vartheta)) dx dt \\
& = - \int_Q (\nabla a_6 \cdot \nabla (\partial_t w)) \vartheta dx dt - \int_Q a_6 (\nabla (\partial_t w) \cdot \nabla \vartheta) dx dt \\
& = \int_Q (\Delta a_6) (\partial_t w) \vartheta dx dt + 2 \int_Q (\nabla a_6 \cdot \nabla \vartheta) (\partial_t w) dx dt \\
& \quad + \int_Q a_6 (\partial_t w) (\Delta \vartheta) dx dt. \tag{2.13}
\end{aligned}$$

We calculate the third term by applying the second equation in (1.4). Since

$$\begin{aligned}
\Delta \vartheta & = \Delta (e^{s\varphi} \Theta) = (s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 + s \lambda^2 \varphi |\nabla \psi|^2 + 2ns\lambda \varphi) \Theta e^{s\varphi} \\
& \quad + (\Delta \Theta) e^{s\varphi} + 2s\lambda \varphi (\nabla \psi \cdot \nabla \Theta) e^{s\varphi}, \\
\partial_t w & = \partial_t (e^{s\varphi} p) = s\lambda \varphi (\partial_t \psi) p e^{s\varphi} + (\partial_t p) e^{s\varphi},
\end{aligned}$$

we have

$$\begin{aligned}
& a_6 (\partial_t w) (\Delta \vartheta) \\
& = s\lambda \varphi (\partial_t \psi) (s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 + s \lambda^2 \varphi |\nabla \psi|^2 + 2ns\lambda \varphi) a_6 p \Theta e^{2s\varphi} \\
& \quad + s\lambda \varphi (\partial_t \psi) a_6 (\Delta \Theta) p e^{2s\varphi} + 2s^2 \lambda^2 \varphi^2 (\partial_t \psi) a_6 (\nabla \psi \cdot \nabla \Theta) p e^{2s\varphi}
\end{aligned}$$

$$\begin{aligned}
& + (s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 + s \lambda^2 \varphi |\nabla \psi|^2 + 2ns \lambda \varphi) a_6 \Theta (\partial_t p) e^{2s\varphi} \\
& + a_6 (\partial_t p) (\Delta \Theta) e^{2s\varphi} + 2s \lambda \varphi a_6 (\nabla \psi \cdot \nabla \Theta) (\partial_t p) e^{2s\varphi}.
\end{aligned} \tag{2.14}$$

Integrating by parts, we have

$$\begin{aligned}
& \int_Q s \lambda \varphi (\partial_t \psi) a_6 (\Delta \Theta) p e^{2s\varphi} dx dt \\
& = -s \lambda \int_Q \varphi (\partial_t \psi) (\nabla a_6 \cdot \nabla \Theta) p e^{2s\varphi} dx dt \\
& \quad - s \lambda^2 \int_Q \varphi (\partial_t \psi) a_6 (\nabla \psi \cdot \nabla \Theta) p e^{2s\varphi} dx dt - s \lambda \int_Q \varphi (\partial_t \psi) a_6 (\nabla \Theta \cdot \nabla p) e^{2s\varphi} dx dt \\
& \quad - 2s^2 \lambda^2 \int_Q \varphi^2 (\partial_t \psi) a_6 (\nabla \psi \cdot \nabla \Theta) p e^{2s\varphi} dx dt.
\end{aligned} \tag{2.15}$$

By the second equation in (1.4), we have

$$\begin{aligned}
a_6 (\partial_t p) (\Delta \Theta) e^{2s\varphi} & = \frac{a_6}{a_3} (\partial_t p) (\partial_t \Theta - a_4 \partial_t^2 p - g) e^{2s\varphi} \\
& = \frac{a_6}{a_3} (\partial_t p) (\partial_t \Theta) e^{2s\varphi} - \frac{a_4 a_6}{2a_3} \partial_t (|\partial_t p|^2) e^{2s\varphi} - \frac{a_6}{a_3} (\partial_t p) g e^{2s\varphi}.
\end{aligned}$$

Therefore we have

$$\begin{aligned}
\int_Q a_6 (\partial_t p) (\Delta \Theta) e^{2s\varphi} dx dt & = \int_Q \frac{a_6}{a_3} (\partial_t p) (\partial_t \Theta) e^{2s\varphi} dx dt \\
& \quad + s \lambda \int_Q \varphi (\partial_t \psi) \frac{a_4 a_6}{a_3} |\partial_t p|^2 e^{2s\varphi} dx dt - \int_Q \frac{a_6}{a_3} (\partial_t p) g e^{2s\varphi} dx dt.
\end{aligned} \tag{2.16}$$

We note that $s > 1$, $\lambda > 1$ and $\varphi > 1$ on $\overline{Q}$. Combining of (2.13)-(2.16) and noting that the integration of the third term in (2.14) over Q plus the last term in (2.15) equals to 0, we obtain

$$\begin{aligned}
\int_Q J_4 dx dt & \geq -s \lambda \int_Q \varphi (\partial_t \psi) a_6 (\nabla \Theta \cdot \nabla p) e^{2s\varphi} dx dt \\
& \quad + \int_Q \frac{a_6}{a_3} (\partial_t p) (\partial_t \Theta) e^{2s\varphi} dx dt + s \lambda \int_Q \varphi (\partial_t \psi) \frac{a_4 a_6}{a_3} |\partial_t p|^2 e^{2s\varphi} dx dt \\
& \quad + 2s \lambda \int_Q \varphi a_6 (\nabla \psi \cdot \nabla \Theta) (\partial_t p) e^{2s\varphi} dx dt \\
& \quad - C_2 \int_Q |\partial_t w| |\vartheta| dx dt - C_2 \int_Q |\partial_t w| |\nabla \vartheta| dx dt - C_2 s^3 \lambda^3 \int_Q \varphi^3 |p| |\Theta| e^{2s\varphi} dx dt \\
& \quad - C_2 s \lambda^2 \int_Q \varphi |p| |\nabla \Theta| e^{2s\varphi} dx dt - C_2 s^2 \lambda^2 \int_Q \varphi^2 |\partial_t p| |\Theta| e^{2s\varphi} dx dt
\end{aligned}$$

$$-C_2 \int_Q |\partial_t p| |g| e^{2s\varphi} dx dt. \quad (2.17)$$

Similarly to J_2 , we have

$$\begin{aligned} \int_Q J_5 dx dt &= -2s\lambda \int_Q \varphi a_6 (\Delta w) (\nabla \psi \cdot (a_5 \nabla \vartheta + a_6 \nabla w)) dx dt \\ &= -2s\lambda \int_Q \varphi a_5 a_6 (\nabla \psi \cdot \nabla \vartheta) (\Delta w) dx dt - 2s\lambda \int_Q \varphi a_6^2 (\nabla \psi \cdot \nabla w) (\Delta w) dx dt \\ &= -2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_5 a_6 \frac{\partial \vartheta}{\partial \nu} \frac{\partial w}{\partial \nu} d\sigma dt + 4s\lambda \int_Q \varphi a_5 a_6 (\nabla \vartheta \cdot \nabla w) dx dt \\ &\quad + 2s\lambda^2 \int_Q \varphi a_5 a_6 (\nabla \psi \cdot \nabla \vartheta) (\nabla \psi \cdot \nabla w) dx dt \\ &\quad + 2s\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla w) (\nabla \psi \cdot \nabla \vartheta) dx dt \\ &\quad + 2s\lambda \int_Q \varphi a_5 a_6 \sum_{j,k=1}^n (\partial_j \psi) (\partial_j \partial_k \vartheta) (\partial_k w) dx dt \\ &\quad - 2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_6^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt + 2s\lambda^2 \int_Q \varphi a_6^2 (\nabla \psi \cdot \nabla w)^2 dx dt \\ &\quad + 4s\lambda \int_Q \varphi a_6 (\nabla a_6 \cdot \nabla w) (\nabla \psi \cdot \nabla w) dx dt + 4s\lambda \int_Q \varphi a_6^2 |\nabla w|^2 dx dt \\ &\quad + s\lambda \int_Q \varphi a_6^2 (\nabla \psi \cdot \nabla (|\nabla w|^2)) dx dt. \end{aligned}$$

Calculating the fifth term by the integration by parts, we have

$$\begin{aligned} (\text{the fifth term}) &= 2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_5 a_6 \frac{\partial \vartheta}{\partial \nu} \frac{\partial w}{\partial \nu} d\sigma dt \\ &\quad - 2s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_5 a_6 (\nabla \vartheta \cdot \nabla w) dx dt \\ &\quad - 2s\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla \psi) (\nabla \vartheta \cdot \nabla w) dx dt - 4ns\lambda \int_Q \varphi a_5 a_6 (\nabla \vartheta \cdot \nabla w) dx dt \\ &\quad - 2s\lambda \int_Q \varphi a_5 a_6 \sum_{j,k=1}^n (\partial_j \psi) (\partial_k \vartheta) (\partial_j \partial_k w) dx dt. \end{aligned}$$

Calculating the tenth term by the integration by parts, we have

$$(\text{the tenth term}) = s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_6^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt - s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_6^2 |\nabla w|^2 dx dt$$

$$- 2s\lambda \int_Q \varphi a_6 (\nabla a_6 \cdot \nabla \psi) |\nabla w|^2 dxdt - 2ns\lambda \int_Q \varphi a_6^2 |\nabla w|^2 dxdt.$$

Therefore we obtain

$$\begin{aligned} \int_Q J_5 dxdt &= 4(1-n)s\lambda \int_Q \varphi a_5 a_6 (\nabla \vartheta \cdot \nabla w) dxdt \\ &+ 2s\lambda^2 \int_Q \varphi a_5 a_6 (\nabla \psi \cdot \nabla \vartheta) (\nabla \psi \cdot \nabla w) dxdt \\ &+ 2s\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla w) (\nabla \psi \cdot \nabla \vartheta) dxdt \\ &- 2s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_5 a_6 (\nabla \vartheta \cdot \nabla w) dxdt \\ &- 2s\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla \psi) (\nabla \vartheta \cdot \nabla w) dxdt \\ &- 2s\lambda \int_Q \varphi a_5 a_6 \sum_{j,k=1}^n (\partial_j \psi) (\partial_k \vartheta) (\partial_j \partial_k w) dxdt \\ &- s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_6^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt + 2s\lambda^2 \int_Q \varphi a_6^2 (\nabla \psi \cdot \nabla w)^2 dxdt \\ &+ 4s\lambda \int_Q \varphi a_6 (\nabla a_6 \cdot \nabla w) (\nabla \psi \cdot \nabla w) dxdt + 2(2-n)s\lambda \int_Q \varphi a_6^2 |\nabla w|^2 dxdt \\ &- s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_6^2 |\nabla w|^2 dxdt - 2s\lambda \int_Q \varphi a_6 (\nabla \psi \cdot \nabla a_6) |\nabla w|^2 dxdt. \quad (2.18) \end{aligned}$$

$$\begin{aligned} \int_Q J_6 dxdt &= - \int_Q a_6 (\Delta w) A(\vartheta, w) dxdt \\ &= -s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_5 a_6 (\Delta w) \vartheta dxdt - s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_6^2 (\Delta w) w dxdt \\ &- 2ns\lambda \int_Q \varphi a_5 a_6 (\Delta w) \vartheta dxdt - 2ns\lambda \int_Q \varphi a_6^2 (\Delta w) w dxdt \\ &+ s\lambda \int_Q \varphi (\partial_t \psi) a_6 (\Delta w) \vartheta dxdt \\ &= s\lambda^3 \int_Q \varphi |\nabla \psi|^2 a_5 a_6 (\nabla \psi \cdot \nabla w) \vartheta dxdt \\ &+ s\lambda^2 \int_Q \varphi (\nabla (|\nabla \psi|^2 a_5 a_6) \cdot \nabla w) \vartheta dxdt + s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_5 a_6 (\nabla w \cdot \nabla \vartheta) dxdt \\ &+ s\lambda^3 \int_Q \varphi |\nabla \psi|^2 a_6^2 (\nabla \psi \cdot \nabla w) w dxdt + s\lambda^2 \int_Q \varphi (\nabla (|\nabla \psi|^2 a_6^2) \cdot \nabla w) w dxdt \end{aligned}$$

$$\begin{aligned}
& + s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_6^2 |\nabla w|^2 dxdt + 2ns\lambda^2 \int_Q \varphi a_5 a_6 (\nabla \psi \cdot \nabla w) \vartheta dxdt \\
& + 2ns\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla w) \vartheta dxdt + 2ns\lambda \int_Q \varphi a_5 a_6 (\nabla w \cdot \nabla \vartheta) dxdt \\
& + 2ns\lambda^2 \int_Q \varphi a_6^2 (\nabla \psi \cdot \nabla w) w dxdt + 4ns\lambda \int_Q \varphi a_6 (\nabla a_6 \cdot \nabla w) w dxdt \\
& + 2ns\lambda \int_Q \varphi a_6^2 |\nabla w|^2 dxdt - s\lambda^2 \int_Q \varphi (\partial_t \psi) a_6 (\nabla \psi \cdot \nabla w) \vartheta dxdt \\
& - s\lambda \int_Q \varphi (\partial_t \psi) (\nabla a_6 \cdot \nabla w) \vartheta dxdt - s\lambda \int_Q \varphi (\partial_t \psi) a_6 (\nabla w \cdot \nabla \vartheta) dxdt \\
& \geq s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_6^2 |\nabla w|^2 dxdt + 2ns\lambda \int_Q \varphi a_6^2 |\nabla w|^2 dxdt \\
& + s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_5 a_6 (\nabla w \cdot \nabla \vartheta) dxdt + 2ns\lambda \int_Q \varphi a_5 a_6 (\nabla w \cdot \nabla \vartheta) dxdt \\
& - s\lambda \int_Q \varphi (\partial_t \psi) a_6 (\nabla w \cdot \nabla \vartheta) dxdt - C_3 s \lambda^3 \int_Q \varphi |\nabla w| (|\vartheta| + |w|) dxdt. \quad (2.19)
\end{aligned}$$

$$\begin{aligned}
\int_Q J_7 dxdt & = -\frac{1}{2} s^2 \lambda^2 \int_Q \varphi^2 |\nabla \psi|^2 a_5 \{ \partial_t (\vartheta^2) \} dxdt \\
& = s^2 \lambda^3 \int_Q \varphi^2 |\nabla \psi|^2 (\partial_t \psi) a_5 \vartheta^2 dxdt \geq -C_4 s^2 \lambda^3 \int_Q \varphi^2 \vartheta^2 dxdt. \quad (2.20)
\end{aligned}$$

$$\begin{aligned}
\int_Q J_8 dxdt & = -2s^3 \lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 a_5 (\nabla \psi \cdot (a_5 \nabla \vartheta + a_6 \nabla w)) \vartheta dxdt \\
& = 3s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 a_5^2 \vartheta^2 dxdt + s^3 \lambda^3 \int_Q \varphi^3 \sum_{j=1}^n \partial_j \{ (\partial_j \psi) |\nabla \psi|^2 a_5^2 \} \vartheta^2 dxdt \\
& \quad - 2s^3 \lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 a_5 a_6 (\nabla \psi \cdot \nabla w) \vartheta dxdt \\
& \geq 3s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 a_5^2 \vartheta^2 dxdt - 2s^3 \lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 a_5 a_6 (\nabla \psi \cdot \nabla w) \vartheta dxdt \\
& \quad - C_5 s^3 \lambda^3 \int_Q \varphi^3 \vartheta^2 dxdt. \quad (2.21)
\end{aligned}$$

$$\int_Q J_9 dxdt = -s^2 \lambda^2 \int_Q \varphi^2 |\nabla \psi|^2 a_5 \vartheta A(\vartheta, w) dxdt$$

$$\begin{aligned}
&= -s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_5^2 \vartheta^2 dxdt - s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_5 a_6 \vartheta w dxdt \\
&\quad - C_6 s^3 \lambda^3 \int_Q \varphi^3 \vartheta^2 dxdt - C_6 s^3 \lambda^3 \int_Q \varphi^3 |\vartheta| |w| dxdt.
\end{aligned} \tag{2.22}$$

$$\begin{aligned}
\int_Q J_{10} dxdt &= -s^2\lambda^2 \int_Q \varphi^2 |\nabla\psi|^2 a_6 w (\partial_t \vartheta) dxdt \\
&= s^2\lambda^2 \int_Q \varphi^2 |\nabla\psi|^2 a_6 \vartheta (\partial_t w) dxdt + 2s^2\lambda^3 \int_Q \varphi^2 (\partial_t \psi) |\nabla\psi|^2 a_6 \vartheta w dxdt \\
&\geq -C_7 s^2 \lambda^2 \int_Q \varphi^2 |\vartheta| |\partial_t w| dxdt - C_7 s^2 \lambda^3 \int_Q \varphi^2 |\vartheta| |w| dxdt.
\end{aligned} \tag{2.23}$$

$$\begin{aligned}
\int_Q J_{11} dxdt &= -2s^3\lambda^3 \int_Q \varphi^3 |\nabla\psi|^2 a_6 (\nabla\psi \cdot (a_5 \nabla \vartheta + a_6 \nabla w)) w dxdt \\
&= 2s^3\lambda^3 \int_Q \varphi^3 \sum_{j=1}^n \partial_j \{ |\nabla\psi|^2 (\partial_j \psi) a_5 a_6 \} \vartheta w dxdt \\
&\quad + 6s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_5 a_6 \vartheta w dxdt + 2s^3\lambda^3 \int_Q \varphi^3 |\nabla\psi|^2 a_5 a_6 (\nabla\psi \cdot \nabla w) \vartheta dxdt \\
&\quad + 3s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_6^2 w^2 dxdt + s^3\lambda^3 \int_Q \varphi^3 \sum_{j=1}^n \partial_j \{ |\nabla\psi|^2 (\partial_j \psi) a_6^2 \} w^2 dxdt \\
&\geq 6s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_5 a_6 \vartheta w dxdt + 2s^3\lambda^3 \int_Q \varphi^3 |\nabla\psi|^2 a_5 a_6 (\nabla\psi \cdot \nabla w) \vartheta dxdt \\
&\quad + 3s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_6^2 w^2 dxdt \\
&\quad - C_8 s^3 \lambda^3 \int_Q \varphi^3 |\vartheta| |w| dxdt - C_8 s^3 \lambda^3 \int_Q \varphi^3 w^2 dxdt.
\end{aligned} \tag{2.24}$$

$$\begin{aligned}
\int_Q J_{12} dxdt &= -s^2\lambda^2 \int_Q \varphi^2 |\nabla\psi|^2 a_6 w A(\vartheta, w) dxdt \\
&\geq -s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_5 a_6 \vartheta w dxdt - s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_6^2 w^2 dxdt \\
&\quad - C_9 s^3 \lambda^3 \int_Q \varphi^3 |\vartheta| |w| dxdt - C_9 s^3 \lambda^3 \int_Q \varphi^3 w^2 dxdt.
\end{aligned} \tag{2.25}$$

We combine (2.9)-(2.12) and (2.17)-(2.25). The sum of the third and ninth terms in the right hand side of (2.11) and the second and eighth terms in the right hand

side of (2.18) equals to

$$2s\lambda^2 \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dxdt.$$

The sum of the sixth term in the right hand side of (2.11) and the first term in the right hand side of (2.12) equals to 0. The sum of the eighth term in the right hand side of (2.11) and the third term in the right hand side of (2.12) equals to 0. The sum of the twelfth term in the right hand side of (2.11) and the sixth term in the right hand side of (2.18) equals to 0. The sum of the second term in the right hand side of (2.12), the fourth term in the right hand side of (2.18), and the third term in the right hand side of (2.19) equals to 0. The sum of the eleventh term in the right hand side of (2.18) and the first term in the right hand side of (2.19) equals to 0. The sum of the second term in the right hand side of (2.21) and the second term in the right hand side of (2.24) equals to 0. Furthermore, we combine the eleventh term in right hand side of (2.11), the fourth term in the right term in the right hand of (2.12), the first term in the right hand side of (2.18), and the fourth term in the right hand side of (2.19). We combine the tenth term in the right hand side of (2.18) and the second term in the right hand side of (2.19). We combine the first term in the right hand side of (2.21) and the first term in the right hand side of (2.22). We combine the second term in the right hand side of (2.22), the first term in the right hand side of (2.24), and the first term in the right hand side of (2.25). We combine the third term in the right hand side of (2.24) and the second term in the right hand side of (2.25). Therefore, we can obtain

$$\begin{aligned} \int_Q L_1(\vartheta, w) L_2(\vartheta, w) dxdt &\geq -s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) \left(a_5 \frac{\partial \vartheta}{\partial \nu} + a_6 \frac{\partial w}{\partial \nu} \right)^2 d\sigma dt \\ &+ 2s\lambda^2 \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dxdt + \int_Q (\nabla a_5 \cdot \nabla \vartheta) (\partial_t \vartheta) dxdt \\ &+ 4s\lambda \int_Q \varphi a_5 (\nabla a_5 \cdot \nabla \vartheta) (\nabla \psi \cdot \nabla \vartheta) dxdt + 4s\lambda \int_Q \varphi a_5^2 |\nabla \vartheta|^2 dxdt \\ &- 2s\lambda \int_Q \varphi a_5 (\nabla \psi \cdot \nabla a_5) |\nabla \vartheta|^2 dxdt + 2s\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla \vartheta) (\nabla \psi \cdot \nabla w) dxdt \\ &+ 8s\lambda \int_Q \varphi a_5 a_6 (\nabla \vartheta \cdot \nabla w) dxdt - s\lambda \int_Q \varphi (\partial_t \psi) a_5 |\nabla \vartheta|^2 dxdt \\ &- s\lambda \int_Q \varphi (\partial_t \psi) a_6 (\nabla \Theta \cdot \nabla p) e^{2s\varphi} dxdt + \int_Q \frac{a_6}{a_3} (\partial_t p) (\partial_t \Theta) e^{2s\varphi} dxdt \\ &+ s\lambda \int_Q \varphi (\partial_t \psi) \frac{a_4 a_6}{a_3} |\partial_t p|^2 e^{2s\varphi} dxdt + 2s\lambda \int_Q \varphi a_6 (\nabla \psi \cdot \nabla \Theta) (\partial_t p) e^{2s\varphi} dxdt \end{aligned}$$

$$\begin{aligned}
& + 2s\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla w) (\nabla \psi \cdot \nabla \vartheta) \, dxdt \\
& - 2s\lambda \int_Q \varphi (\nabla (a_5 a_6) \cdot \nabla \psi) (\nabla \vartheta \cdot \nabla w) \, dxdt \\
& + 4s\lambda \int_Q \varphi a_6 (\nabla a_6 \cdot \nabla w) (\nabla \psi \cdot \nabla w) \, dxdt + 4s\lambda \int_Q \varphi a_6^2 |\nabla w|^2 \, dxdt \\
& - 2s\lambda \int_Q \varphi a_6 (\nabla \psi \cdot \nabla a_6) |\nabla w|^2 \, dxdt - s\lambda \int_Q \varphi (\partial_t \psi) a_6 (\nabla w \cdot \nabla \vartheta) \, dxdt \\
& + 2s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 a_5^2 \vartheta^2 \, dxdt + 4s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 a_5 a_6 \vartheta w \, dxdt \\
& + 2s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 a_6^2 w^2 \, dxdt - C_{10} \int_Q |\partial_t w| (|\vartheta| + |\nabla \vartheta|) \, dxdt \\
& - C_{10} s^3 \lambda^3 \int_Q \varphi^3 |p| |\Theta| e^{2s\varphi} \, dxdt - C_{10} s^2 \lambda^2 \int_Q \varphi^2 |\partial_t p| |\Theta| e^{2s\varphi} \, dxdt \\
& - C_{10} s \lambda^2 \int_Q \varphi |p| |\nabla \Theta| e^{2s\varphi} \, dxdt - C_{10} \int_Q |\partial_t p| |g| e^{2s\varphi} \, dxdt \\
& - C_{10} s \lambda^3 \int_Q \varphi (|\nabla \vartheta| + |\nabla w|) (|\vartheta| + |w|) \, dxdt \\
& - C_{10} s^3 \lambda^3 \int_Q \varphi^3 (\vartheta^2 + w^2) \, dxdt - C_{10} s^2 \lambda^2 \int_Q \varphi^2 |\vartheta| |\partial_t w| \, dxdt. \tag{2.26}
\end{aligned}$$

For $s > 1$, we have, $(\Theta, p) = e^{-s\varphi} (\vartheta, w)$,

$$\begin{aligned}
\partial_t \Theta &= \partial_t (e^{-s\varphi} \vartheta) = -s\lambda \varphi (\partial_t \psi) e^{-s\varphi} \vartheta + e^{-s\varphi} \partial_t \vartheta \\
&= 2s\lambda \varphi \beta (t - t_0) e^{-s\varphi} \vartheta + e^{-s\varphi} \partial_t \vartheta, \\
\partial_j \Theta &= \partial_j (e^{-s\varphi} \vartheta) = -s\lambda \varphi (\partial_j \psi) e^{-s\varphi} \vartheta + e^{-s\varphi} \partial_j \vartheta \\
&= -2s\lambda \varphi (x_j - x_0^j) e^{-s\varphi} \vartheta + e^{-s\varphi} \partial_j \vartheta, \\
\partial_t p &= \partial_t (e^{-s\varphi} w) = -s\lambda \varphi (\partial_t \psi) e^{-s\varphi} w + e^{-s\varphi} \partial_t w \\
&= 2s\lambda \varphi \beta (t - t_0) e^{-s\varphi} w + e^{-s\varphi} \partial_t w, \\
\partial_j p &= \partial_j (e^{-s\varphi} w) = -s\lambda \varphi (\partial_j \psi) e^{-s\varphi} w + e^{-s\varphi} \partial_j w \\
&= -2s\lambda \varphi (x_j - x_0^j) e^{-s\varphi} w + e^{-s\varphi} \partial_j w, \quad \text{in } Q. \tag{2.27}
\end{aligned}$$

By the Cauchy-Schwarz inequality, $(a_1, a_2, a_3, a_4) \in \mathcal{U}$, and (2.27), we estimate every term of (2.26). We note that $\|a_5\|_{C(\overline{\Omega})} \leq 2M_0^2$, $\|a_6\|_{C(\overline{\Omega})} \leq M_0^2$, $\|\partial_j a_5\|_{C(\overline{\Omega})} \leq 3M_0 M_1$, $\|\partial_j a_6\|_{C(\overline{\Omega})} \leq 2M_0 M_1$, and $\|\partial_j (a_5 a_6)\|_{C(\overline{\Omega})} = \|a_5 \partial_j a_6 + a_6 \partial_j a_5\|_{C(\overline{\Omega})} \leq 7M_0^3 M_1$, $j = 1, 2, \dots, n$. By the Cauchy-Schwarz inequality, we can get, for any

$\varepsilon_1 > 0$,

$$|(\nabla a_5 \cdot \nabla \vartheta)(\partial_t \vartheta)| \leq \frac{9M_0^2 M_1^2}{2} \varepsilon_1^{-1} s \varphi |\nabla \vartheta|^2 + \frac{n \varepsilon_1}{2s \varphi} |\partial_t \vartheta|^2, \quad \text{in } Q. \quad (2.28)$$

$$\begin{aligned} |4s \lambda \varphi a_5 (\nabla a_5 \cdot \nabla \vartheta)(\nabla \psi \cdot \nabla \vartheta)| &= |8s \lambda \varphi a_5 (\nabla a_5 \cdot \nabla \vartheta)((x - x_0) \cdot \nabla \vartheta)| \\ &\leq 48n M_0^3 M_1 M^{\frac{1}{2}} s \lambda \varphi |\nabla \vartheta|^2, \quad \text{in } Q. \end{aligned} \quad (2.29)$$

$$4s \lambda \varphi a_5^2 |\nabla \vartheta|^2 \geq 4\sigma_1^2 s \lambda \varphi |\nabla \vartheta|^2, \quad \text{in } Q. \quad (2.30)$$

$$\begin{aligned} |-2s \lambda \varphi a_5 (\nabla \psi \cdot \nabla a_5) |\nabla \vartheta|^2| &= |4s \lambda \varphi a_5 ((x - x_0) \cdot \nabla a_5)| |\nabla \vartheta|^2 \\ &\leq 24n M_0^3 M_1 M^{\frac{1}{2}} s \lambda \varphi |\nabla \vartheta|^2, \quad \text{in } Q. \end{aligned} \quad (2.31)$$

Similarly,

$$|-2s \lambda \varphi a_6 (\nabla \psi \cdot \nabla a_6) |\nabla w|^2| \leq 8n M_0^3 M_1 M^{\frac{1}{2}} s \lambda \varphi |\nabla w|^2, \quad \text{in } Q. \quad (2.32)$$

$$|4s \lambda \varphi a_6 (\nabla a_6 \cdot \nabla w)(\nabla \psi \cdot \nabla w)| \leq 16n M_0^3 M_1 M^{\frac{1}{2}} s \lambda \varphi |\nabla w|^2, \quad \text{in } Q. \quad (2.33)$$

Similarly, we have

$$\begin{aligned} &|2s \lambda \varphi (\nabla (a_5 a_6) \cdot \nabla \vartheta)(\nabla \psi \cdot \nabla w)| + |2s \lambda \varphi (\nabla (a_5 a_6) \cdot \nabla w)(\nabla \psi \cdot \nabla \vartheta)| \\ &\quad + |-2s \lambda \varphi (\nabla (a_5 a_6) \cdot \nabla \psi)(\nabla \vartheta \cdot \nabla w)| \\ &\leq 42n M_0^3 M_1 M^{\frac{1}{2}} s \lambda \varphi (|\nabla \vartheta|^2 + |\nabla w|^2), \quad \text{in } Q. \end{aligned} \quad (2.34)$$

$$|8s \lambda \varphi a_5 a_6 (\nabla \vartheta \cdot \nabla w)| \leq 8M_0^4 s \lambda \varphi (|\nabla \vartheta|^2 + |\nabla w|^2), \quad \text{in } Q. \quad (2.35)$$

$$|-s \lambda \varphi (\partial_t \psi) a_5 |\nabla \vartheta|^2| = |2s \lambda \varphi \beta a_5 (t - t_0)| |\nabla \vartheta|^2 \leq 2M_0^2 s \lambda \varphi \beta T |\nabla \vartheta|^2, \quad (2.36)$$

in Q . By (2.27), we have

$$\begin{aligned} &\left| 2s \lambda \varphi a_6 (\nabla \psi \cdot \nabla \Theta)(\partial_t p) e^{2s \varphi} \right| \\ &= \left| 2s \lambda \varphi a_6 (\nabla \psi \cdot \nabla \vartheta)(\partial_t w) - 2s^2 \lambda^2 \varphi^2 (\partial_t \psi) a_6 (\nabla \psi \cdot \nabla \vartheta) w \right. \\ &\quad \left. - 2s^2 \lambda^2 \varphi^2 |\nabla \psi|^2 a_6 (\partial_t w) \vartheta + 2s^3 \lambda^3 \varphi^3 |\nabla \psi|^2 (\partial_t \psi) a_6 w \vartheta \right| \\ &\leq |4s \lambda \varphi ((\partial_t w)(x - x_0) \cdot a_6 \nabla \vartheta)| \end{aligned}$$

$$\begin{aligned}
& + \left| 8\beta (t - t_0) a_6 \left((x - x_0) \cdot \left(s^{\frac{1}{2}} \lambda^{\frac{1}{4}} \varphi^{\frac{1}{2}} \nabla \vartheta \right) \right) s^{\frac{3}{2}} \lambda^{\frac{7}{4}} \varphi^{\frac{3}{2}} w \right| \\
& + |\nabla \psi|^2 \left| 2a_6 \left(s^{\frac{1}{2}} \lambda^{\frac{1}{4}} \varphi^{\frac{1}{2}} \partial_t w \right) s^{\frac{3}{2}} \lambda^{\frac{7}{4}} \varphi^{\frac{3}{2}} \vartheta \right| + s^3 \lambda^3 \varphi^3 |\nabla \psi|^2 |\partial_t \psi| |2a_6 w \vartheta| \\
& \leq 2s\lambda\varphi (M |\partial_t w|^2 + M_0^4 |\nabla \vartheta|^2) + C_{11} s \lambda^{\frac{1}{2}} \varphi (|\nabla \vartheta|^2 + |\partial_t w|^2) \\
& \quad + C_{11} s^3 \lambda^{\frac{7}{2}} \varphi^3 (\vartheta^2 + w^2), \quad \text{in } Q.
\end{aligned} \tag{2.37}$$

Similarly to (2.37), we have

$$\begin{aligned}
& \left| -s\lambda\varphi (\partial_t \psi) a_6 (\nabla \Theta \cdot \nabla p) e^{2s\varphi} \right| \\
& = \left| -s\lambda\varphi (\partial_t \psi) a_6 (\nabla \vartheta \cdot \nabla w) + s^2 \lambda^2 \varphi^2 (\partial_t \psi) a_6 (\nabla \psi \cdot \nabla \vartheta) w \right. \\
& \quad \left. + s^2 \lambda^2 \varphi^2 (\partial_t \psi) a_6 (\nabla \psi \cdot \nabla w) \vartheta - s^3 \lambda^3 \varphi^3 |\nabla \psi|^2 (\partial_t \psi) a_6 \vartheta w \right| \\
& \leq \frac{1}{2} M_0^2 s \lambda \varphi \beta T (|\nabla \vartheta|^2 + |\nabla w|^2) + C_{12} s \lambda^{\frac{1}{2}} \varphi (|\nabla \vartheta|^2 + |\nabla w|^2) \\
& \quad + C_{12} s^3 \lambda^{\frac{7}{2}} \varphi^3 (\vartheta^2 + w^2), \quad \text{in } Q.
\end{aligned} \tag{2.38}$$

Similarly, we can obtain

$$s^2 \lambda^2 \varphi^2 |\vartheta| |\partial_t w| \leq C_{13} s \lambda^{\frac{1}{2}} \varphi |\partial_t w|^2 + C_{13} s^3 \lambda^{\frac{7}{2}} \varphi^3 \vartheta^2, \tag{2.39}$$

$$\begin{aligned}
s^2 \lambda^2 \varphi^2 |\partial_t p| |\Theta| e^{2s\varphi} & \leq s^2 \lambda^2 \varphi^2 (|\partial_t w| |\vartheta| + |-s\lambda\varphi (\partial_t \psi) w| |\vartheta|) \\
& \leq C_{14} s \lambda^{\frac{1}{2}} \varphi |\partial_t w|^2 + C_{14} s^3 \lambda^{\frac{7}{2}} \varphi^3 (\vartheta^2 + w^2), \quad \text{in } Q.
\end{aligned} \tag{2.40}$$

$$\begin{aligned}
& \left| \frac{a_6}{a_3} (\partial_t \Theta) (\partial_t p) e^{2s\varphi} \right| = \left| \frac{a_6}{a_3} (\partial_t \vartheta) (\partial_t w) - s\lambda\varphi (\partial_t \psi) \frac{a_6}{a_3} (\partial_t \vartheta) w \right. \\
& \quad \left. - s\lambda\varphi (\partial_t \psi) \frac{a_6}{a_3} (\partial_t w) \vartheta + s^2 \lambda^2 \varphi^2 (\partial_t \psi)^2 \frac{a_6}{a_3} \vartheta w \right| \\
& \leq \left| \left(\frac{1}{\sqrt{6s\varphi M}} \partial_t \vartheta \right) \left(\frac{M_0^2}{\sigma_1} \sqrt{6s\varphi M} \partial_t w \right) \right| \\
& \quad + \frac{M_0^2 \beta T}{\sigma_1} \left| \left(\sqrt{\frac{1}{s\lambda\varphi}} \partial_t \vartheta \right) \left(s\lambda\varphi \sqrt{s\lambda\varphi} w \right) \right| + \frac{M_0^2 \beta T}{\sigma_1} |(\partial_t w) (s\lambda\varphi \vartheta)| \\
& \quad + \frac{M_0^2 \beta^2 T^2}{\sigma_1} |s^2 \lambda^2 \varphi^2 \vartheta w| \\
& \leq \frac{1}{12s\varphi M} |\partial_t \vartheta|^2 + \frac{3M_0^4 M}{\sigma_1^2} s\varphi |\partial_t w|^2 \\
& \quad + C_{15} \left(\frac{1}{s\lambda\varphi} |\partial_t \vartheta|^2 + s^3 \lambda^3 \varphi^3 w^2 + |\partial_t w|^2 + s^2 \lambda^2 \varphi^2 \vartheta^2 \right), \quad \text{in } Q.
\end{aligned} \tag{2.41}$$

$$\begin{aligned}
& \left| s\lambda\varphi(\partial_t\psi) \frac{a_4a_6}{a_3} |\partial_t p|^2 e^{2s\varphi} \right| = \left| 2s\lambda\varphi\beta(t-t_0) \frac{a_4a_6}{a_3} |\partial_t w + 2s\lambda\varphi\beta(t-t_0)w|^2 \right| \\
& \leq \frac{2M_0^3\beta T}{\sigma_1} s\lambda\varphi |\partial_t w|^2 + C_{16}s^3\lambda^3\beta^3\varphi^3w^2, \quad \text{in } Q.
\end{aligned} \tag{2.42}$$

$$4s\lambda\varphi a_6^2 |\nabla w|^2 \geq 0, \quad \text{in } Q. \tag{2.43}$$

$$|-s\lambda\varphi(\partial_t\psi) a_6 (\nabla w \cdot \nabla \vartheta)| \leq \frac{1}{2} M_0^2 s\lambda\beta T\varphi (|\nabla \vartheta|^2 + |\nabla w|^2), \quad \text{in } Q. \tag{2.44}$$

$$2s^3\lambda^4\varphi^3 |\nabla\psi|^4 a_5^2\vartheta^2 \geq 2\sigma_1^2 s^3\lambda^4\varphi^3 |\nabla\psi|^4 \vartheta^2, \quad \text{in } Q. \tag{2.45}$$

By the Cauchy-Schwarz inequality, we have, for any $\varepsilon_2 > 0$,

$$|4s^3\lambda^4\varphi^3 |\nabla\psi|^4 a_5a_6\vartheta w| \leq 4M_0^4 s^3\lambda^4\varphi^3 |\nabla\psi|^4 (\varepsilon_2\vartheta^2 + \varepsilon_2^{-1}w^2), \quad \text{in } Q. \tag{2.46}$$

$$2s^3\lambda^4\varphi^3 |\nabla\psi|^4 a_6^2w^2 \geq 0, \quad \text{in } Q. \tag{2.47}$$

Moreover, by (2.27), we have

$$\begin{aligned}
s\lambda^2\varphi|p| |\nabla\Theta| e^{2s\varphi} & \leq s\lambda^2\varphi (|\nabla\vartheta| + |-s\lambda\varphi(\nabla\psi)\vartheta|) |w| \\
& \leq C_{17} |\nabla\vartheta|^2 + C_{17}s^2\lambda^4\varphi^2 (\vartheta^2 + w^2), \quad \text{in } Q.
\end{aligned} \tag{2.48}$$

$$\begin{aligned}
|\partial_t p| |g| e^{2s\varphi} & \leq (|\partial_t w| + |-s\lambda\varphi(\partial_t\psi)w|) (|g|e^{s\varphi}) \\
& \leq C_{18} |\partial_t w|^2 + C_{18}s^2\lambda^2\varphi^2w^2 + C_{18}g^2e^{2s\varphi}, \quad \text{in } Q.
\end{aligned} \tag{2.49}$$

$$\begin{aligned}
& s\lambda^3\varphi (|\nabla\vartheta| + |\nabla w|) (|\vartheta| + |w|) \\
& \leq C_{19} (|\nabla\vartheta|^2 + |\nabla w|^2) + C_{19}s^2\lambda^6\varphi^2 (\vartheta^2 + w^2), \quad \text{in } Q.
\end{aligned} \tag{2.50}$$

Moreover, by the second equality in (2.7) and the Cauchy-Schwarz inequality, we have, for any $\varepsilon_3 > 0$,

$$\begin{aligned}
|L_2(\vartheta, w)|^2 & = |\partial_t\vartheta + \{2s\lambda\varphi(\nabla\psi \cdot (a_5\nabla\vartheta + a_6\nabla w)) + A(\vartheta, w)\}|^2 \\
& = |\partial_t\vartheta|^2 + b^2 + 2(\sqrt{\varepsilon_3}\partial_t\vartheta) \frac{b}{\sqrt{\varepsilon_3}} \geq (1 - \varepsilon_3) |\partial_t\vartheta|^2 + (1 - \varepsilon_3^{-1}) b^2,
\end{aligned}$$

where $b = 2s\lambda\varphi(\nabla\psi \cdot (a_5\nabla\vartheta + a_6\nabla w)) + A(\vartheta, w)$. Noting

$$b^2 \leq 8s^2\lambda^2\varphi^2 |a_5(\nabla\psi \cdot \nabla\vartheta) + a_6(\nabla\psi \cdot \nabla w)|^2 + C_{20}s^2\lambda^4\varphi^2 (\vartheta^2 + w^2),$$

and choosing $\varepsilon_3 = \frac{4s\varphi M}{4s\varphi M + 1}$, we have

$$\begin{aligned}
|L_2(\vartheta, w)|^2 &\geq \frac{1}{4s\varphi M + 1} |\partial_t \vartheta|^2 - 2s\lambda^2 \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 \\
&\quad - C_{21} s \lambda^4 \varphi (\vartheta^2 + w^2) \\
&\geq \frac{1}{5s\varphi M} |\partial_t \vartheta|^2 - 2s\lambda^2 \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 \\
&\quad - C_{21} s \lambda^4 \varphi (\vartheta^2 + w^2), \tag{2.51}
\end{aligned}$$

for all $s > s_1 \triangleq \max\{\frac{1}{M}, 1\}$. By (2.8), (2.26), (2.28)–(2.51), we obtain

$$\begin{aligned}
&\int_Q |\mathcal{L}(\vartheta, w)|^2 dx dt \\
&= \|L_1(\vartheta, w)\|_{L^2(Q)}^2 + 2 \int_Q L_1(\vartheta, w) L_2(\vartheta, w) dx dt + \|L_2(\vartheta, w)\|_{L^2(Q)}^2 \\
&\geq -2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) \left(a_5 \frac{\partial \vartheta}{\partial \nu} + a_6 \frac{\partial w}{\partial \nu} \right)^2 d\sigma dt + \|L_1(\vartheta, w)\|_{L^2(Q)}^2 \\
&\quad + 2s\lambda^2 \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dx dt \\
&\quad + (4\sigma_1^2 - 8M_0^4 \varepsilon_2) s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 \vartheta^2 dx dt - \frac{8M_0^4 s^3 \lambda^4}{\varepsilon_2} \int_Q \varphi^3 |\nabla \psi|^4 w^2 dx dt \\
&\quad + \left(\frac{1}{30M} - n\varepsilon_1 \right) \int_Q \frac{1}{s\varphi} |\partial_t \vartheta|^2 dx dt - 4Ms\lambda \int_Q \varphi |\partial_t w|^2 dx dt \\
&\quad - \left(228nM_0^3 M_1 M^{\frac{1}{2}} + 20M_0^4 - 8\sigma_1^2 \right) s\lambda \int_Q \varphi |\nabla \vartheta|^2 dx dt \\
&\quad - \left(132nM_1 M^{\frac{1}{2}} + 16M_0 \right) M_0^3 s\lambda \int_Q \varphi |\nabla w|^2 dx dt \\
&\quad - 6M_0^2 s\lambda \beta T \int_Q \varphi |\nabla \vartheta|^2 dx dt - 2M_0^2 s\lambda \beta T \int_Q \varphi |\nabla w|^2 dx dt \\
&\quad - \frac{4M_0^3}{\sigma_1} s\lambda \beta T \int_Q \varphi |\partial_t w|^2 dx dt - C_{22} \int_Q s\lambda^{\frac{1}{2}} \varphi (|\nabla \vartheta|^2 + |\nabla w|^2 + |\partial_t w|^2) dx dt \\
&\quad - C_{22} \int_Q \left(s^3 \lambda^{\frac{7}{2}} \varphi^3 + s^2 \lambda^6 \varphi^2 \right) (\vartheta^2 + w^2) dx dt - C_{22} \int_Q \frac{1}{s\lambda \varphi} |\partial_t \vartheta|^2 dx dt \\
&\quad - C_{22} \int_Q g^2 e^{2s\varphi} dx dt,
\end{aligned}$$

for all $s > s_1$. Choosing $\varepsilon_1 > 0$ and $\varepsilon_2 > 0$ sufficiently small such that

$$4\sigma_1^2 - 8M_0^4 \varepsilon_2 = 2\sigma_1^2 \quad \text{and} \quad \frac{1}{30M} - n\varepsilon_1 = \frac{1}{60M},$$

that is, $\varepsilon_1 = \frac{1}{60nM}$ and $\varepsilon_2 = \frac{\sigma_1^2}{4M_0^4}$, we arrive at

$$\begin{aligned}
& 2\sigma_1^2 s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 \vartheta^2 dx dt + \frac{1}{60M} \int_Q \frac{1}{s\varphi} |\partial_t \vartheta|^2 dx dt \\
& + 2s\lambda^2 \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dx dt \\
& - 2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) \left(a_5 \frac{\partial \vartheta}{\partial \nu} + a_6 \frac{\partial w}{\partial \nu} \right)^2 d\sigma dt + \|L_1(\vartheta, w)\|_{L^2(Q)}^2 \\
& \leq C_{23} \int_Q |\mathcal{L}(\vartheta, w)|^2 dx dt + \alpha_1 s\lambda \int_Q \varphi |\nabla \vartheta|^2 dx dt + \alpha_2 s\lambda \int_Q \varphi |\nabla w|^2 dx dt \\
& + 4Ms\lambda \int_Q \varphi |\partial_t w|^2 dx dt + \frac{32M_0^8}{\sigma_1^2} s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 w^2 dx dt \\
& + 6M_0^2 s\lambda \beta T \int_Q \varphi |\nabla \vartheta|^2 dx dt + 2M_0^2 s\lambda \beta T \int_Q \varphi |\nabla w|^2 dx dt \\
& + \frac{4M_0^3}{\sigma_1} s\lambda \beta T \int_Q \varphi |\partial_t w|^2 dx dt + C_{23} s\lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2 + |\partial_t w|^2) dx dt \\
& + C_{23} \int_Q \left(s^3 \lambda^{\frac{7}{2}} \varphi^3 + s^2 \lambda^6 \varphi^2 \right) (\vartheta^2 + w^2) dx dt + C_{23} \int_Q \frac{1}{s\lambda\varphi} |\partial_t \vartheta|^2 dx dt \\
& + C_{23} \int_Q g^2 e^{2s\varphi} dx dt, \tag{2.52}
\end{aligned}$$

for all $s > s_1$, where α_1 and α_2 are given by the first and second equality in (1.8).

Next we will estimate $s\lambda \int_Q \varphi |\nabla \vartheta|^2 e^{2s\varphi} dx dt$. Noting (2.6) and (2.7) and applying the integration by parts, similarly to (2.38), we can obtain

$$\begin{aligned}
& \int_Q \mathcal{L}(\vartheta, w) (s\lambda\varphi\vartheta) dx dt \\
& = \int_Q \left\{ -s\lambda\varphi a_5 (\Delta \vartheta) \vartheta - s\lambda\varphi a_6 (\Delta w) \vartheta + \frac{1}{2} s\lambda\varphi (\partial_t (\vartheta^2)) \right\} dx dt \\
& + \int_Q \left\{ -s^3 \lambda^3 \varphi^3 |\nabla \psi|^2 a_5 \vartheta^2 - s^3 \lambda^3 \varphi^3 |\nabla \psi|^2 a_6 \vartheta w \right. \\
& + 2s^2 \lambda^2 \varphi^2 a_5 (\nabla \psi \cdot \nabla \vartheta) \vartheta \\
& + 2s^2 \lambda^2 \varphi^2 a_6 (\nabla \psi \cdot \nabla w) \vartheta + s^2 \lambda^3 \varphi^2 |\nabla \psi|^2 (a_5 \vartheta + a_6 w) \vartheta \\
& \left. + 2ns^2 \lambda^2 \varphi^2 (a_5 \vartheta + a_6 w) \vartheta - s^2 \lambda^2 \varphi^2 (\partial_t \psi) \vartheta^2 \right\} dx dt \\
& \geq s\lambda \int_Q \varphi (\nabla a_5 \cdot \nabla \vartheta) \vartheta dx dt + s\lambda^2 \int_Q \varphi a_5 (\nabla \psi \cdot \nabla \vartheta) \vartheta dx dt
\end{aligned}$$

$$\begin{aligned}
& + s\lambda \int_Q \varphi a_5 |\nabla \vartheta|^2 dxdt + s\lambda \int_Q \varphi (\nabla a_6 \cdot \nabla w) \vartheta dxdt \\
& + s\lambda^2 \int_Q \varphi a_6 (\nabla \psi \cdot \nabla w) \vartheta dxdt + s\lambda \int_Q \varphi a_6 (\nabla w \cdot \nabla \vartheta) dxdt \\
& - \frac{1}{2} s\lambda^2 \int_Q \varphi (\partial_t \psi) \vartheta^2 dxdt - C_{24} s^3 \lambda^3 \int_Q \varphi^3 w^2 dxdt \\
& - C_{24} s^3 \lambda^{\frac{7}{2}} \int_Q \varphi^3 \vartheta^2 dxdt - C_{24} s \lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2) dxdt \\
& \geq \sigma_1 s\lambda \int_Q \varphi |\nabla \vartheta|^2 dxdt - C_{25} \int_Q (|\nabla \vartheta|^2 + |\nabla w|^2) dxdt \\
& - C_{25} s^2 \lambda^2 \int_Q \varphi^2 \vartheta^2 dxdt - C_{25} s \lambda^{\frac{7}{2}} \int_Q \varphi \vartheta^2 dxdt \\
& - C_{25} s \lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2) dxdt \\
& - s\lambda \int_Q \varphi (\sqrt{\sigma_1} |\nabla \vartheta|) \left(\frac{M_0^2}{\sqrt{\sigma_1}} |\nabla w| \right) dxdt \\
& - C_{25} s^3 \lambda^3 \int_Q \varphi^3 w^2 dxdt - C_{25} s^3 \lambda^{\frac{7}{2}} \int_Q \varphi^3 \vartheta^2 dxdt \\
& \geq \frac{1}{2} \sigma_1 s\lambda \int_Q \varphi |\nabla \vartheta|^2 dxdt - \frac{M_0^4}{2\sigma_1} s\lambda \int_Q \varphi |\nabla w|^2 dxdt \\
& - C_{26} s \lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2) dxdt - C_{26} s^3 \lambda^{\frac{7}{2}} \int_Q \varphi^3 \vartheta^2 dxdt \\
& - C_{26} s^3 \lambda^3 \int_Q \varphi^3 w^2 dxdt.
\end{aligned}$$

Since

$$\int_Q |\mathcal{L}(\vartheta, w)(s\lambda\varphi\vartheta)| dxdt \leq \frac{1}{2} \int_Q |\mathcal{L}(\vartheta, w)|^2 dxdt + \frac{1}{2} s^2 \lambda^2 \int_Q \varphi^2 \vartheta^2 dxdt,$$

we obtain

$$\begin{aligned}
& \sigma_1 \int_Q s\lambda\varphi |\nabla \vartheta|^2 dxdt \\
& \leq \frac{M_0^4}{\sigma_1} \int_Q s\lambda\varphi |\nabla w|^2 dxdt + C_{27} \int_Q |\mathcal{L}(\vartheta, w)|^2 dxdt \\
& + C_{27} s \lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2) dxdt + C_{27} s^3 \lambda^{\frac{7}{2}} \int_Q \varphi^3 \vartheta^2 dxdt
\end{aligned}$$

$$+ C_{27}s^3\lambda^3 \int_Q \varphi^3 w^2 dx dt. \quad (2.53)$$

Multiplying (2.53) by $\frac{2\alpha_1}{\sigma_1}$, adding it to (2.52), and noting the equality in (2.8), we obtain

$$\begin{aligned} & 2\sigma_1^2 s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 \vartheta^2 dx dt + \frac{1}{60M} \int_Q \frac{1}{s\varphi} |\partial_t \vartheta|^2 dx dt \\ & + \alpha_1 s \lambda \int_Q \varphi |\nabla \vartheta|^2 dx dt + 2s\lambda^2 \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dx dt \\ & - 2s\lambda \int_\Sigma \varphi (\nabla \psi \cdot \nu) \left(a_5 \frac{\partial \vartheta}{\partial \nu} + a_6 \frac{\partial w}{\partial \nu} \right)^2 d\sigma dt + \|L_1(\vartheta, w)\|_{L^2(Q)}^2 \\ & \leq C_{28} \int_Q (f^2 + g^2) e^{2s\varphi} dx dt + \left(\frac{2M_0^4 \alpha_1}{\sigma_1^2} + \alpha_2 \right) s \lambda \int_Q \varphi |\nabla w|^2 dx dt \\ & + 4Ms\lambda \int_Q \varphi |\partial_t w|^2 dx dt + \frac{32M_0^8}{\sigma_1^2} s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 w^2 dx dt \\ & + 6M_0^2 s \lambda \beta T \int_Q \varphi |\nabla \vartheta|^2 dx dt + 2M_0^2 s \lambda \beta T \int_Q \varphi |\nabla w|^2 dx dt \\ & + \frac{4M_0^3}{\sigma_1} s \lambda \beta T \int_Q \varphi |\partial_t w|^2 dx dt + C_{28} s \lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2 + |\partial_t w|^2) dx dt \\ & + C_{28} \int_Q \left(s^3 \lambda^{\frac{7}{2}} \varphi^3 + s^2 \lambda^6 \varphi^2 \right) (\vartheta^2 + w^2) dx dt + C_{28} \int_Q \frac{1}{s\lambda\varphi} |\partial_t \vartheta|^2 dx dt, \quad (2.54) \end{aligned}$$

for all $s > s_1$.

Next we estimate $s\lambda \int_Q \varphi |\nabla_{x,t} w|^2 dx dt$ and $s^3 \lambda^4 \int_Q \varphi^3 w^2 dx dt$. By (1.4), we can get the following equation

$$a_7 \partial_t^2 p - a_1 \Delta p - a_8 \partial_t \Theta = f - a_8 g, \quad (2.55)$$

where

$$a_7 = 1 + \frac{a_2 a_4}{a_3} \quad \text{and} \quad a_8 = \frac{a_2}{a_3}. \quad (2.56)$$

Denote

$$L_3(\Theta, p) = a_7 \partial_t^2 p - a_1 \Delta p - a_8 \partial_t \Theta. \quad (2.57)$$

We recall $(\vartheta, w) = e^{s\varphi}(\Theta, p)$. We set

$$\mathcal{L}_*(\vartheta, w) = e^{s\varphi} L_3(e^{-s\varphi}(\vartheta, w)). \quad (2.58)$$

Then we have

$$(f - a_8 g) e^{s\varphi} = \mathcal{L}_*(\vartheta, w) = e^{s\varphi} \{ a_7 \partial_t^2 (e^{-s\varphi} w) - a_1 \Delta (e^{-s\varphi} w) - a_8 \partial_t (e^{-s\varphi} \vartheta) \}$$

$$\begin{aligned}
&= \{s^2\lambda^2\varphi^2 (\partial_t\psi)^2 - s\lambda^2\varphi (\partial_t\psi)^2 - s\lambda\varphi\partial_t^2\psi\} a_7w + a_7\partial_t^2w \\
&\quad - 2s\lambda\varphi (\partial_t\psi) a_7\partial_tw - (s^2\lambda^2\varphi^2 |\nabla\psi|^2 - s\lambda^2\varphi |\nabla\psi|^2 - 2ns\lambda\varphi) a_1w \\
&\quad - a_1\Delta w + 2s\lambda\varphi a_1 (\nabla\psi \cdot \nabla w) - a_8\partial_t\vartheta + s\lambda\varphi (\partial_t\psi) a_8\vartheta.
\end{aligned}$$

We set

$$J_0 = s\lambda\varphi\gamma_0w + \sum_{j=1}^n \gamma_j\partial_jw, \quad (2.59)$$

where $\gamma_0 = \gamma_0(x)$ and $\gamma_j = \gamma_j(x) \in C^2(\overline{\Omega})$, ($j = 1, \dots, n$) are functions of x , but are independent of t, s, λ, φ , and will be suitably chosen later. We divide $\mathcal{L}_*(\vartheta, w)$ as follows:

$$\mathcal{L}_*(\vartheta, w) = L_{1*}(\vartheta, w) + L_{2*}(w), \quad (2.60)$$

where

$$\begin{aligned}
L_{1*}(\vartheta, w) &= a_7\partial_t^2w - a_1\Delta w + s^2\lambda^2\varphi^2 (\partial_t\psi)^2 a_7w \\
&\quad - s^2\lambda^2\varphi^2 |\nabla\psi|^2 a_1w - a_8\partial_t\vartheta + s\lambda\varphi (\partial_t\psi) a_8\vartheta + J_0,
\end{aligned} \quad (2.61)$$

$$\begin{aligned}
L_{2*}(w) &= -2s\lambda\varphi (\partial_t\psi) a_7\partial_tw + 2s\lambda\varphi a_1 (\nabla\psi \cdot \nabla w) \\
&\quad - s\lambda^2\varphi (\partial_t\psi)^2 a_7w - s\lambda\varphi (\partial_t^2\psi) a_7w \\
&\quad + s\lambda^2\varphi |\nabla\psi|^2 a_1w + 2ns\lambda\varphi a_1w - J_0.
\end{aligned} \quad (2.62)$$

Then we have

$$\begin{aligned}
&\|(f - a_8g) e^{s\varphi}\|_{L^2(Q)}^2 = \|\mathcal{L}_*(\vartheta, w)\|_{L^2(Q)}^2 \\
&= \|\mathcal{L}_{1*}(\vartheta, w)\|_{L^2(Q)}^2 + 2 \int_Q L_{1*}(\vartheta, w) L_{2*}(w) dxdt + \|\mathcal{L}_{2*}(w)\|_{L^2(Q)}^2.
\end{aligned} \quad (2.63)$$

We have

$$\begin{aligned}
L_{1*}(\vartheta, w) L_{2*}(w) &= a_7 (\partial_t^2w) L_{2*}(w) - a_1 (\Delta w) L_{2*}(w) \\
&\quad + s^2\lambda^2\varphi^2 (\partial_t\psi)^2 a_7w L_{2*}(w) - s^2\lambda^2\varphi^2 |\nabla\psi|^2 a_1w L_{2*}(w) \\
&\quad - a_8 (\partial_t\vartheta) L_{2*}(w) + s\lambda\varphi (\partial_t\psi) a_8\vartheta L_{2*}(w) + J_0 L_{2*}(w) \triangleq \sum_{k=1}^7 I_k.
\end{aligned} \quad (2.64)$$

We integrate I_k , $k = 1, 2, \dots, 6$ over Q . We calculate them by applying the integration by parts and $(a_1, a_2, a_3, a_4) \in \mathcal{U}$. By (1.12) and $\partial_t w = \partial_t(e^{s\varphi}p) = e^{s\varphi}\partial_t p + s(\partial_t\varphi)p$, we have

$$\vartheta(x, 0) = \vartheta(x, T) = 0, \quad \partial_t^j w(x, 0) = \partial_t^j w(x, T) = 0, \quad x \in \overline{\Omega}, \quad j = 0, 1.$$

We further recall that, by (1.11), we have $\vartheta = 0$ and $w = 0$ on Σ , so that we have $\nabla \vartheta = \frac{\partial \vartheta}{\partial \nu} \nu$ and $\nabla w = \frac{\partial w}{\partial \nu} \nu$ on Σ .

$$\begin{aligned}
\int_Q I_1 dx dt &= \int_Q a_7 (\partial_t^2 w) L_{2*}(w) dx dt = - \int_Q a_7 (\partial_t w) \partial_t \{L_{2*}(w)\} dx dt \\
&= 2s\lambda^2 \int_Q \varphi (\partial_t \psi)^2 a_7^2 |\partial_t w|^2 dx dt + 2s\lambda \int_Q \varphi (\partial_t^2 \psi) a_7^2 |\partial_t w|^2 dx dt \\
&\quad + s\lambda \int_Q \varphi (\partial_t \psi) a_7^2 \partial_t \{(\partial_t w)^2\} dx dt \\
&\quad - 2s\lambda^2 \int_Q \varphi (\partial_t \psi) a_1 a_7 (\nabla \psi \cdot \nabla w) (\partial_t w) dx dt \\
&\quad - s\lambda \int_Q \varphi a_1 a_7 (\nabla \psi \cdot \nabla \{(\partial_t w)^2\}) dx dt \\
&\quad + \int_Q \left\{ s\lambda^2 \varphi (\partial_t \psi)^2 a_7^2 |\partial_t w|^2 + s\lambda \varphi (\partial_t^2 \psi) a_7^2 |\partial_t w|^2 \right. \\
&\quad \left. - s\lambda^2 \varphi |\nabla \psi|^2 a_1 a_7 |\partial_t w|^2 - 2ns\lambda \varphi a_1 a_7 |\partial_t w|^2 + s\lambda \varphi a_7 \gamma_0 |\partial_t w|^2 \right\} dx dt \\
&\quad + \frac{1}{2} \int_Q a_7 \sum_{j=1}^n \gamma_j \partial_j \{(\partial_t w)^2\} dx dt \\
&\quad + \int_Q \left\{ s\lambda^3 \varphi (\partial_t \psi)^3 a_7^2 (\partial_t w) w + 2s\lambda^2 \varphi (\partial_t \psi) (\partial_t^2 \psi) a_7^2 (\partial_t w) w \right. \\
&\quad \left. + s\lambda^2 \varphi (\partial_t \psi) (\partial_t^2 \psi) a_7^2 (\partial_t w) w - s\lambda^3 \varphi (\partial_t \psi) |\nabla \psi|^2 a_1 a_7 (\partial_t w) w \right. \\
&\quad \left. - 2ns\lambda^2 \varphi (\partial_t \psi) a_1 a_7 (\partial_t w) w + s\lambda^2 \varphi (\partial_t \psi) a_7 \gamma_0 (\partial_t w) w \right\} dx dt.
\end{aligned}$$

We integrate the third and fifth terms by parts and then combine them with the sixth term. We integrate the seventh term by parts. We further estimate the eighth term by the Cauchy-Schwarz inequality. Then we obtain

$$\begin{aligned}
\int_Q I_1 dx dt &\geq 2s\lambda^2 \int_Q \varphi (\partial_t \psi)^2 a_7^2 |\partial_t w|^2 dx dt + 2s\lambda \int_Q \varphi (\partial_t^2 \psi) a_7^2 |\partial_t w|^2 dx dt \\
&\quad - 2s\lambda^2 \int_Q \varphi (\partial_t \psi) a_1 a_7 (\nabla \psi \cdot \nabla w) (\partial_t w) dx dt \\
&\quad + s\lambda \int_Q \varphi (\nabla \psi \cdot \nabla (a_1 a_7)) |\partial_t w|^2 dx dt + s\lambda \int_Q \varphi a_7 \gamma_0 |\partial_t w|^2 dx dt \\
&\quad - \frac{1}{2} \int_Q \left\{ \sum_{j=1}^n \partial_j (a_7 \gamma_j) \right\} |\partial_t w|^2 dx dt
\end{aligned}$$

$$- C_{29} \int_Q |\partial_t w|^2 dx dt - C_{29} s^2 \lambda^6 \int_Q \varphi^2 w^2 dx dt. \quad (2.65)$$

$$\begin{aligned}
\int_Q I_2 dx dt &= - \int_Q a_1 (\Delta w) L_{2*}(w) dx dt \\
&= -2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_1^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt + \int_Q a_1 (\nabla w \cdot \nabla \{L_{2*}(w)\}) dx dt \\
&\quad + \int_Q (\nabla a_1 \cdot \nabla w) L_{2*}(w) dx dt \\
&\geq -2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_1^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt - 2s\lambda^2 \int_Q \varphi (\partial_t \psi) a_1 a_7 (\nabla \psi \cdot \nabla w) (\partial_t w) dx dt \\
&\quad - 2s\lambda \int_Q \varphi (\partial_t \psi) a_1 (\nabla a_7 \cdot \nabla w) (\partial_t w) dx dt - s\lambda \int_Q \varphi (\partial_t \psi) a_1 a_7 \partial_t (|\nabla w|^2) dx dt \\
&\quad + 2s\lambda^2 \int_Q \varphi a_1^2 (\nabla \psi \cdot \nabla w)^2 dx dt + 2s\lambda \int_Q \varphi a_1 (\nabla a_1 \cdot \nabla w) (\nabla \psi \cdot \nabla w) dx dt \\
&\quad + 4s\lambda \int_Q \varphi a_1^2 |\nabla w|^2 dx dt + s\lambda \int_Q \varphi a_1^2 (\nabla \psi \cdot \nabla \{|\nabla w|^2\}) dx dt \\
&\quad - s\lambda^2 \int_Q \varphi (\partial_t \psi)^2 a_1 a_7 |\nabla w|^2 dx dt - s\lambda \int_Q \varphi (\partial_t^2 \psi) a_1 a_7 |\nabla w|^2 dx dt \\
&\quad + s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_1^2 |\nabla w|^2 dx dt + 2ns\lambda \int_Q \varphi a_1^2 |\nabla w|^2 dx dt \\
&\quad - s\lambda \int_Q \varphi \gamma_0 a_1 |\nabla w|^2 dx dt - \frac{1}{2} \int_Q a_1 \sum_{j=1}^n \gamma_j \partial_j (|\nabla w|^2) dx dt \\
&\quad - 2s\lambda \int_Q \varphi (\partial_t \psi) a_7 (\nabla a_1 \cdot \nabla w) (\partial_t w) dx dt \\
&\quad + 2s\lambda \int_Q \varphi a_1 (\nabla a_1 \cdot \nabla w) (\nabla \psi \cdot \nabla w) dx dt \\
&\quad - C_{30} \int_Q |\nabla w|^2 dx dt - C_{30} s^2 \lambda^6 \int_Q \varphi^2 |w|^2 dx dt.
\end{aligned}$$

We combine the third and fifteenth terms. We integrate the fourth term by parts and then combine them with the ninth and tenth terms. We combine the sixth and sixteenth terms. We integrate the eighth term by parts and then combine them with the first, eleventh and twelfth terms. We further integrate the fourteenth

term by parts. Thus we obtain

$$\begin{aligned}
\int_Q I_2 dx dt &\geq -s\lambda \int_{\Sigma} \varphi(\nabla\psi \cdot \nu) a_1^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt - \frac{1}{2} \int_{\Sigma} a_1 \left(\sum_{j=1}^n \gamma_j \nu_j \right) \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt \\
&\quad - 2s\lambda^2 \int_Q \varphi(\partial_t \psi) a_1 a_7 (\nabla\psi \cdot \nabla w) (\partial_t w) dx dt \\
&\quad - 2s\lambda \int_Q \varphi(\partial_t \psi) (\nabla(a_1 a_7) \cdot \nabla w) (\partial_t w) dx dt \\
&\quad + 2s\lambda^2 \int_Q \varphi a_1^2 (\nabla\psi \cdot \nabla w)^2 dx dt + 4s\lambda \int_Q \varphi a_1 (\nabla a_1 \cdot \nabla w) (\nabla\psi \cdot \nabla w) dx dt \\
&\quad + 4s\lambda \int_Q \varphi a_1^2 |\nabla w|^2 dx dt - 2s\lambda \int_Q \varphi a_1 (\nabla\psi \cdot \nabla a_1) |\nabla w|^2 dx dt \\
&\quad - s\lambda \int_Q \varphi \gamma_0 a_1 |\nabla w|^2 dx dt + \frac{1}{2} \int_Q \left\{ \sum_j^n \partial_j (a_1 \gamma_j) \right\} |\nabla w|^2 dx dt \\
&\quad - C_{31} \int_Q |\nabla w|^2 dx dt - C_{31} s^2 \lambda^6 \int_Q \varphi^2 |w|^2 dx dt. \tag{2.66}
\end{aligned}$$

$$\begin{aligned}
\int_Q I_3 dx dt &= s^2 \lambda^2 \int_Q \varphi^2 (\partial_t \psi)^2 a_7 w L_{2*}(w) dx dt \\
&\geq -s^3 \lambda^3 \int_Q \varphi^3 (\partial_t \psi)^3 a_7^2 \{ \partial_t (w^2) \} dx dt \\
&\quad + s^3 \lambda^3 \int_Q \varphi^3 (\partial_t \psi)^2 a_1 a_7 (\nabla\psi \cdot \nabla (w^2)) dx dt \\
&\quad - s^3 \lambda^4 \int_Q \varphi^3 (\partial_t \psi)^4 a_7^2 w^2 dx dt + s^3 \lambda^4 \int_Q \varphi^3 (\partial_t \psi)^2 |\nabla\psi|^2 a_1 a_7 w^2 dx dt \\
&\quad - C_{32} s^3 \lambda^3 \int_Q \varphi^3 w^2 dx dt - \frac{1}{2} s^2 \lambda^2 \int_Q \varphi^2 (\partial_t \psi)^2 a_7 \sum_{j=1}^n \gamma_j \{ \partial_j (w^2) \} dx dt.
\end{aligned}$$

We integrate the first term by parts and then combine it with the third term. We integrate the second term by parts and then combine it with the fourth term. We further integrate the sixth term by parts and estimate it. Therefore we can obtain

$$\begin{aligned}
\int_Q I_3 dx dt &\geq 2s^3 \lambda^4 \int_Q \varphi^3 (\partial_t \psi)^4 a_7^2 w^2 dx dt + 3s^3 \lambda^3 \int_Q \varphi^3 (\partial_t \psi)^2 (\partial_t^2 \psi) a_7^2 w^2 dx dt \\
&\quad - 2s^3 \lambda^4 \int_Q \varphi^3 (\partial_t \psi)^2 |\nabla\psi|^2 a_1 a_7 w^2 dx dt
\end{aligned}$$

$$\begin{aligned}
& -s^3\lambda^3 \int_Q \varphi^3 (\partial_t \psi)^2 (\nabla \psi \cdot \nabla (a_1 a_7)) w^2 dx dt \\
& -C_{33}s^3\lambda^3 \int_Q \varphi^3 w^2 dx dt + s^2\lambda^3 \int_Q \varphi^2 (\partial_t \psi)^2 a_7 \left\{ \sum_{j=1}^n \gamma_j (\partial_j \psi) \right\} w^2 dx dt \\
& + \frac{1}{2}s^2\lambda^2 \int_Q \varphi^2 (\partial_t \psi)^2 \left\{ \sum_{j=1}^n \partial_j (a_7 \gamma_j) \right\} w^2 dx dt.
\end{aligned}$$

Noting the first term is nonnegative, we obtain

$$\int_Q I_3 dx dt \geq -2s^3\lambda^4 \int_Q \varphi^3 (\partial_t \psi)^2 |\nabla \psi|^2 a_1 a_7 w^2 dx dt - C_{34}s^3\lambda^3 \int_Q \varphi^3 w^2 dx dt. \quad (2.67)$$

$$\begin{aligned}
\int_Q I_4 dx dt &= -s^2\lambda^2 \int_Q \varphi^2 |\nabla \psi|^2 a_1 w L_{2*}(w) dx dt \\
&= s^3\lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 (\partial_t \psi) a_1 a_7 \{ \partial_t (w^2) \} dx dt \\
&\quad - s^3\lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 a_1^2 (\nabla \psi \cdot \nabla (w^2)) dx dt \\
&\quad + s^3\lambda^4 \int_Q \varphi^3 |\nabla \psi|^2 (\partial_t \psi)^2 a_1 a_7 w^2 dx dt + s^3\lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 (\partial_t^2 \psi) a_1 a_7 w^2 dx dt \\
&\quad - s^3\lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 a_1^2 w^2 dx dt - 2ns^3\lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 a_1^2 w^2 dx dt \\
&\quad + s^3\lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 a_1 \gamma_0 w^2 dx dt + \frac{1}{2}s^2\lambda^2 \int_Q \varphi^2 |\nabla \psi|^2 a_1 \sum_{j=1}^n \gamma_j \{ \partial_j (w^2) \} dx dt.
\end{aligned}$$

We integrate the first term by parts and combine it with the third and fourth term. We integrate the second term by parts and combine it with the fifth and sixth term. We integrate the eighth term by parts. Then we obtain

$$\begin{aligned}
\int_Q I_4 dx dt &\geq -2s^3\lambda^4 \int_Q \varphi^3 |\nabla \psi|^2 (\partial_t \psi)^2 a_1 a_7 w^2 dx dt \\
&\quad + 2s^3\lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 a_1^2 w^2 dx dt + s^3\lambda^3 \int_Q \varphi^3 (\nabla \psi \cdot \nabla (|\nabla \psi|^2 a_1^2)) w^2 dx dt \\
&\quad + s^3\lambda^3 \int_Q \varphi^3 |\nabla \psi|^2 a_1 \gamma_0 w^2 dx dt - s^2\lambda^3 \int_Q \varphi^2 |\nabla \psi|^2 a_1 \left(\sum_{j=1}^n \gamma_j \partial_j \psi \right) w^2 dx dt
\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}s^2\lambda^2 \int_Q \varphi^2 \left\{ \sum_{j=1}^n \partial_j (|\nabla\psi|^2 a_1 \gamma_j) \right\} w^2 dx dt \\
& \geq -2s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^2 (\partial_t\psi)^2 a_1 a_7 w^2 dx dt + 2s^3\lambda^4 \int_Q \varphi^3 |\nabla\psi|^4 a_1^2 w^2 dx dt \\
& \quad - C_{35}s^3\lambda^3 \int_Q \varphi^3 w^2 dx dt.
\end{aligned} \tag{2.68}$$

$$\begin{aligned}
I_5 & = -a_8 (\partial_t\vartheta) L_{2*}(w) \\
& = 2s\lambda\varphi (\partial_t\psi) a_7 a_8 (\partial_t\vartheta) (\partial_t w) - 2s\lambda\varphi a_1 a_8 (\nabla\psi \cdot \nabla w) (\partial_t\vartheta) \\
& \quad + s\lambda^2\varphi (\partial_t\psi)^2 a_7 a_8 (\partial_t\vartheta) w + \{s\lambda\varphi (\partial_t^2\psi) a_7 a_8 (\partial_t\vartheta) w \\
& \quad - s\lambda^2\varphi |\nabla\psi|^2 a_1 a_8 (\partial_t\vartheta) w - 2ns\lambda\varphi a_1 a_8 (\partial_t\vartheta) w \\
& \quad + s\lambda\varphi a_8 \gamma_0 (\partial_t\vartheta) w\} + a_8 (\partial_t\vartheta) \sum_{j=1}^n \gamma_j \partial_j w \triangleq \sum_{k=1}^5 I_{5k}.
\end{aligned} \tag{2.69}$$

Integrating by parts and using the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
\int_Q I_{51} dx dt & = 2s\lambda \int_Q \varphi (\partial_t\psi) a_7 a_8 (\partial_t\vartheta) (\partial_t w) dx dt \\
& = -2s\lambda \int_Q \varphi (\partial_t\psi) a_7 a_8 \vartheta (\partial_t^2 w) dx dt \\
& \quad - 2s\lambda^2 \int_Q \varphi (\partial_t\psi)^2 a_7 a_8 \vartheta (\partial_t w) dx dt - 2s\lambda \int_Q \varphi (\partial_t^2\psi) a_7 a_8 \vartheta (\partial_t w) dx dt \\
& \geq -2s\lambda \int_Q \varphi (\partial_t\psi) a_7 a_8 \vartheta (\partial_t^2 w) dx dt \\
& \quad - C_{36}s^2\lambda^4 \int_Q \varphi^2 \vartheta^2 dx dt - C_{36} \int_Q |\partial_t w|^2 dx dt.
\end{aligned}$$

We calculate the first term as follows. By (2.61), we have

$$\begin{aligned}
a_7 \partial_t^2 w & = L_{1*}(\vartheta, w) + a_1 \Delta w - s^2\lambda^2\varphi^2 (\partial_t\psi)^2 a_7 w \\
& \quad + s^2\lambda^2\varphi^2 |\nabla\psi|^2 a_1 w + a_8 \partial_t\vartheta - s\lambda\varphi (\partial_t\psi) a_8 \vartheta - J_0, \text{ in } Q.
\end{aligned}$$

Therefore we have

$$\begin{aligned}
\int_Q I_{51} dx dt & \geq -2s\lambda \int_Q \varphi (\partial_t\psi) a_1 a_8 \vartheta (\Delta w) dx dt \\
& \quad + 2s^3\lambda^3 \int_Q \varphi^3 (\partial_t\psi) a_8 \{(\partial_t\psi)^2 a_7 - |\nabla\psi|^2 a_1\} \vartheta w dx dt
\end{aligned}$$

$$\begin{aligned}
& -s\lambda \int_Q \varphi (\partial_t \psi) a_8^2 \{ \partial_t (\vartheta^2) \} dxdt + 2s^2\lambda^2 \int_Q \varphi^2 (\partial_t \psi)^2 a_8^2 \vartheta^2 dxdt \\
& - 2s\lambda \int_Q \varphi (\partial_t \psi) a_8 L_{1*} (\vartheta, w) \vartheta dxdt \\
& + 2s^2\lambda^2 \int_Q \varphi^2 (\partial_t \psi) a_8 \gamma_0 w \vartheta dxdt + 2s\lambda \int_Q \varphi (\partial_t \psi) a_8 \vartheta \sum_{j=1}^n \gamma_j \partial_j w dxdt \\
& - C_{37} s^2 \lambda^4 \int_Q \varphi^2 \vartheta^2 dxdt - C_{37} \int_Q |\partial_t w|^2 dxdt.
\end{aligned}$$

Integrating the first and third terms by parts and estimating the second and fifth terms by the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
\int_Q I_{51} dxdt & \geq 2s\lambda \int_Q \varphi (\partial_t \psi) a_1 a_8 (\nabla \vartheta \cdot \nabla w) dxdt \\
& + 2s\lambda^2 \int_Q \varphi (\partial_t \psi) a_1 a_8 (\nabla \psi \cdot \nabla w) \vartheta dxdt \\
& + 2s\lambda \int_Q \varphi (\partial_t \psi) (\nabla (a_1 a_8) \cdot \nabla w) \vartheta dxdt \\
& + s\lambda^2 \int_Q \varphi (\partial_t \psi)^2 a_8^2 \vartheta^2 dxdt + s\lambda \int_Q \varphi (\partial_t^2 \psi) a_8^2 \vartheta^2 dxdt \\
& + 2s^2\lambda^2 \int_Q \varphi^2 (\partial_t \psi)^2 a_8^2 \vartheta^2 dxdt - 5s^2\lambda^2 \int_Q \varphi^2 (\partial_t \psi)^2 a_8^2 \vartheta^2 dxdt \\
& - \frac{1}{4} \int_Q |L_{1*} (\vartheta, w)|^2 dxdt - C_{38} s^3 \lambda^3 \int_Q \varphi^3 (\vartheta^2 + w^2) dxdt \\
& - C_{38} s^2 \lambda^4 \int_Q \varphi^2 \vartheta^2 dxdt - C_{38} \int_Q (|\nabla w|^2 + |\partial_t w|^2) dxdt \\
& \geq 2s\lambda \int_Q \varphi (\partial_t \psi) a_1 a_8 (\nabla \vartheta \cdot \nabla w) dxdt - \frac{1}{4} \int_Q |L_{1*} (\vartheta, w)|^2 dxdt \\
& - C_{39} s^3 \lambda^3 \int_Q \varphi^3 (\vartheta^2 + w^2) dxdt - C_{39} s^2 \lambda^4 \int_Q \varphi^2 \vartheta^2 dxdt \\
& - C_{39} \int_Q (|\nabla w|^2 + |\partial_t w|^2) dxdt. \tag{2.70}
\end{aligned}$$

Integrating by parts two times, we have

$$\begin{aligned}
\int_Q I_{52} dxdt & = -2s\lambda \int_Q \varphi a_1 a_8 (\nabla \psi \cdot \nabla w) (\partial_t \vartheta) dxdt \\
& = 2s\lambda \int_Q \varphi a_1 a_8 (\nabla \psi \cdot \nabla (\partial_t w)) \vartheta dxdt
\end{aligned}$$

$$\begin{aligned}
& + 2s\lambda^2 \int_Q \varphi (\partial_t \psi) a_1 a_8 (\nabla \psi \cdot \nabla w) \vartheta dx dt \\
& = -2s\lambda \int_Q \varphi a_1 a_8 (\nabla \psi \cdot \nabla \vartheta) (\partial_t w) dx dt \\
& \quad - 2s\lambda \int_Q \varphi \{ \operatorname{div} (a_1 a_8 \nabla \psi) \} (\partial_t w) \vartheta dx dt \\
& \quad - 2s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_1 a_8 (\partial_t w) \vartheta dx dt \\
& \quad + 2s\lambda^2 \int_Q \varphi (\partial_t \psi) a_1 a_8 (\nabla \psi \cdot \nabla w) \vartheta dx dt \\
& \geq -2s\lambda \int_Q \varphi a_1 a_8 (\nabla \psi \cdot \nabla \vartheta) (\partial_t w) dx dt \\
& \quad - C_{40} s^2 \lambda^4 \int_Q \varphi^2 \vartheta^2 dx dt - C_{40} \int_Q (|\nabla w|^2 + |\partial_t w|^2) dx dt. \tag{2.71}
\end{aligned}$$

Integrating by parts, we have

$$\begin{aligned}
\int_Q I_{53} dx dt & = s\lambda^2 \int_Q \varphi (\partial_t \psi)^2 a_7 a_8 (\partial_t \vartheta) w dx dt \\
& = -s\lambda^2 \int_Q \varphi (\partial_t \psi)^2 a_7 a_8 \vartheta (\partial_t w) dx dt \\
& \quad - 2s\lambda^2 \int_Q \varphi (\partial_t \psi) (\partial_t^2 \psi) a_7 a_8 \vartheta w dx dt - s\lambda^3 \int_Q \varphi (\partial_t \psi)^3 a_7 a_8 \vartheta w dx dt \\
& \geq -C_{41} s^2 \lambda^4 \int_Q \varphi^2 \vartheta^2 dx dt - C_{41} \int_Q |\partial_t w|^2 dx dt - C_{41} s\lambda^3 \int_Q \varphi w^2 dx dt. \tag{2.72}
\end{aligned}$$

Similarly, integrating by parts, we can obtain

$$\int_Q I_{54} dx dt \geq -C_{42} s^2 \lambda^4 \int_Q \varphi^2 \vartheta^2 dx dt - C_{42} \int_Q |\partial_t w|^2 dx dt - C_{42} s\lambda^3 \int_Q \varphi w^2 dx dt. \tag{2.73}$$

Integrating by parts two times, we have

$$\begin{aligned}
\int_Q I_{55} dx dt & = \int_Q a_8 \sum_{j=1}^n \gamma_j (\partial_j w) (\partial_t \vartheta) dx dt = - \int_Q a_8 \sum_{j=1}^n \gamma_j (\partial_t \partial_j w) \vartheta dx dt \\
& = \int_Q a_8 \sum_{j=1}^n \gamma_j (\partial_j \vartheta) (\partial_t w) dx dt + \int_Q \left\{ \sum_{j=1}^n \partial_j (a_8 \gamma_j) \right\} (\partial_t w) \vartheta dx dt
\end{aligned}$$

$$\geq -C_{43} \int_Q (|\partial_t w|^2 + |\nabla \vartheta|^2 + |\vartheta|^2) dx dt. \quad (2.74)$$

Combining (2.69)–(2.74), we arrive at

$$\begin{aligned} \int_Q I_5 dx dt &\geq 2s\lambda \int_Q \varphi (\partial_t \psi) a_1 a_8 (\nabla \vartheta \cdot \nabla w) dx dt - \frac{1}{4} \int_Q |L_{1*}(\vartheta, w)|^2 dx dt \\ &\quad - 2s\lambda \int_Q \varphi a_1 a_8 (\nabla \psi \cdot \nabla \vartheta) (\partial_t w) dx dt \\ &\quad - C_{44} s^3 \lambda^3 \int_Q \varphi^3 (\vartheta^2 + w^2) dx dt - C_{44} s^2 \lambda^4 \int_Q \varphi^2 \vartheta^2 dx dt \\ &\quad - C_{44} \int_Q (|\nabla w|^2 + |\partial_t w|^2 + |\nabla \vartheta|^2) dx dt. \end{aligned} \quad (2.75)$$

$$\begin{aligned} \int_Q I_6 dx dt &= s\lambda \int_Q \varphi (\partial_t \psi) a_8 \vartheta L_{2*}(w) dx dt \\ &= -2s^2 \lambda^2 \int_Q \varphi^2 (\partial_t \psi)^2 a_7 a_8 \vartheta (\partial_t w) dx dt \\ &\quad + 2s^2 \lambda^2 \int_Q \varphi^2 (\partial_t \psi) a_1 a_8 (\nabla \psi \cdot \nabla w) \vartheta dx dt - s^2 \lambda^3 \int_Q \varphi^2 (\partial_t \psi)^3 a_7 a_8 w \vartheta dx dt \\ &\quad - s^2 \lambda^2 \int_Q \varphi^2 (\partial_t \psi) (\partial_t^2 \psi) a_7 a_8 \vartheta w dx dt + s^2 \lambda^3 \int_Q \varphi^2 (\partial_t \psi) |\nabla \psi|^2 a_1 a_8 \vartheta w dx dt \\ &\quad + 2ns^2 \lambda^2 \int_Q \varphi^2 (\partial_t \psi) a_1 a_8 \vartheta w dx dt - s^2 \lambda^2 \int_Q \varphi^2 (\partial_t \psi) a_8 \gamma_0 \vartheta w dx dt \\ &\quad - s\lambda \int_Q \varphi (\partial_t \psi) a_8 \vartheta \sum_{j=1}^n \gamma_j \partial_j w dx dt. \end{aligned} \quad (2.76)$$

Similarly to (2.37), we can estimate the first and second terms. Therefore we have

$$\begin{aligned} \int_Q I_6 dx dt &\geq -C_{45} s \lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla w|^2 + |\partial_t w|^2) dx dt - C_{45} s^3 \lambda^{\frac{7}{2}} \int_Q \varphi^3 \vartheta^2 dx dt \\ &\quad - C_{45} s^2 \lambda^3 \int_Q \varphi^2 (\vartheta^2 + w^2) dx dt - C_{45} \int_Q |\nabla w|^2 dx dt. \end{aligned} \quad (2.77)$$

By (2.59), we have

$$\int_Q I_7 dx dt = \int_Q J_0 L_{2*}(w) dx dt$$

$$= s\lambda \int_Q \varphi \gamma_0 w L_{2*}(w) \, dxdt + \int_Q \left(\sum_{j=1}^n \gamma_j \partial_j w \right) L_{2*}(w) \, dxdt.$$

Similarly to (2.76)-(2.77), we can estimate the first term.

$$\begin{aligned} (\text{second term}) &= -2s\lambda \int_Q \varphi (\partial_t \psi) a_7 (\partial_t w) \sum_{j=1}^n \gamma_j (\partial_j w) \, dxdt \\ &\quad + 2s\lambda \int_Q \varphi a_1 (\nabla \psi \cdot \nabla w) \left(\sum_{j=1}^n \gamma_j \partial_j w \right) \, dxdt \\ &\quad - s\lambda^2 \int_Q \varphi (\partial_t \psi)^2 a_7 \sum_{j=1}^n \gamma_j (\partial_j w) w \, dxdt \\ &\quad - s\lambda \int_Q \varphi (\partial_t^2 \psi) a_7 \sum_{j=1}^n \gamma_j (\partial_j w) w \, dxdt \\ &\quad + s\lambda^2 \int_Q \varphi |\nabla \psi|^2 a_1 \sum_{j=1}^n \gamma_j (\partial_j w) w + 2ns\lambda \int_Q \varphi a_1 \sum_{j=1}^n \gamma_j (\partial_j w) w \, dxdt \\ &\quad - s\lambda \int_Q \varphi \gamma_0 \sum_{j=1}^n \gamma_j (\partial_j w) w \, dxdt - \int_Q \left(\sum_{j=1}^n \gamma_j \partial_j w \right)^2 \, dxdt \\ &\geq -2s\lambda \int_Q \varphi (\partial_t \psi) a_7 (\partial_t w) \sum_{j=1}^n \gamma_j (\partial_j w) \, dxdt \\ &\quad + 2s\lambda \int_Q \varphi a_1 (\nabla \psi \cdot \nabla w) \left(\sum_{j=1}^n \gamma_j \partial_j w \right) \, dxdt \\ &\quad - C_{46}s^2\lambda^4 \int_Q \varphi^2 w^2 \, dxdt - C_{46} \int_Q |\nabla w|^2 \, dxdt. \end{aligned}$$

Therefore we can obtain

$$\begin{aligned} \int_Q I_7 \, dxdt &\geq -2s\lambda \int_Q \varphi (\partial_t \psi) a_7 (\partial_t w) \sum_{j=1}^n \gamma_j (\partial_j w) \, dxdt \\ &\quad + 2s\lambda \int_Q \varphi a_1 (\nabla \psi \cdot \nabla w) \left(\sum_{j=1}^n \gamma_j \partial_j w \right) \, dxdt \\ &\quad - C_{47}s\lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla w|^2 + |\partial_t w|^2) \, dxdt - C_{47}s^3\lambda^{\frac{7}{2}} \int_Q \varphi^3 w^2 \, dxdt \\ &\quad - C_{47}s^2\lambda^4 \int_Q \varphi^2 w^2 \, dxdt - C_{47} \int_Q |\nabla w|^2 \, dxdt. \end{aligned} \tag{2.78}$$

We combine of (2.63), (2.64), (2.65)–(2.68), (2.75), (2.77), and (2.78). We note that the sum of the first and third terms in (2.65) and the third and fifth terms in (2.66) equals

$$2s\lambda^2 \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 dx dt.$$

Moreover, we take

$$\gamma_0 = a_1 - \frac{(\nabla(a_1 a_7) \cdot \nabla \psi)}{a_7}, \quad \gamma_j = -2(\partial_j a_1), \quad j = 1, 2, \dots, n. \quad (2.79)$$

Then the sum of the fourth and fifth terms in (2.65) equals to $s\lambda \int_Q \varphi a_1 a_7 |\partial_t w|^2 dx dt$. The sum of the sixth term in (2.66) and the second term in (2.78) equals 0. The sum of the fourth term in (2.66) and the first term in (2.78) equals to

$$2s\lambda \int_Q \varphi (\partial_t \psi) a_7 (\nabla a_1 \cdot \nabla w) (\partial_t w) dx dt - 2s\lambda \int_Q \varphi (\partial_t \psi) a_1 (\nabla a_7 \cdot \nabla w) (\partial_t w) dx dt.$$

We combine the first term in (2.67) and the first term in (2.68). Thus we arrive at

$$\begin{aligned} & \| (f - a_8 g) e^{s\varphi} \|_{L^2(Q)}^2 = \| \mathcal{L}_* (\vartheta, w) \|_{L^2(Q)}^2 \\ & = \| \mathcal{L}_{1*} (\vartheta, w) \|_{L^2(Q)}^2 + 2 \int_Q L_{1*} (\vartheta, w) L_{2*} (w) dx dt + \| \mathcal{L}_{2*} (w) \|_{L^2(Q)}^2 \\ & \geq \frac{1}{2} \| \mathcal{L}_* (\vartheta, w) \|_{L^2(Q)}^2 + \| \mathcal{L}_{2*} (w) \|_{L^2(Q)}^2 \\ & \quad - 2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_1^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt + 2 \int_{\Sigma} a_1 (\nabla a_1 \cdot \nu) \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt \\ & \quad + 4s\lambda^2 \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 dx dt + 2s\lambda \int_Q \varphi a_1 a_7 |\partial_t w|^2 dx dt \\ & \quad + 2s\lambda \int_Q \varphi a_1 \left\{ 3a_1 - (\nabla a_1 \cdot \nabla \psi) + \frac{a_1 (\nabla a_7 \cdot \nabla \psi)}{a_7} \right\} |\nabla w|^2 dx dt \\ & \quad + 4s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 a_1^2 w^2 dx dt - 4s\lambda \int_Q \varphi a_1 a_8 (\nabla \psi \cdot \nabla \vartheta) (\partial_t w) dx dt \\ & \quad + 4s\lambda \int_Q \varphi (\partial_t^2 \psi) a_7^2 |\partial_t w|^2 dx dt + 4s\lambda \int_Q \varphi (\partial_t \psi) a_7 (\nabla a_1 \cdot \nabla w) (\partial_t w) dx dt \\ & \quad - 4s\lambda \int_Q \varphi (\partial_t \psi) a_1 (\nabla a_7 \cdot \nabla w) (\partial_t w) dx dt \\ & \quad + 4s\lambda \int_Q \varphi (\partial_t \psi) a_1 a_8 (\nabla \vartheta \cdot \nabla w) dx dt \end{aligned}$$

$$\begin{aligned}
& - 8s^3\lambda^4 \int_Q \varphi^3 (\partial_t \psi)^2 |\nabla \psi|^2 a_1 a_7 w^2 dx dt \\
& - C_{48} s \lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla w|^2 + |\partial_t w|^2) dx dt - C_{48} s^3 \lambda^{\frac{7}{2}} \int_Q \varphi^3 (\vartheta^2 + w^2) dx dt \\
& - C_{48} \int_Q (|\nabla \vartheta|^2 + s^2 \lambda^4 \varphi^2 \vartheta^2 + s^2 \lambda^6 \varphi^2 w^2) dx dt.
\end{aligned} \tag{2.80}$$

By $(a_1(x), a_2(x), a_3(x), a_4(x)) \in \mathcal{U}$ and the Cauchy-Schwarz inequality, we estimate every term of (2.80). We note that $\|a_7\|_{C(\bar{\Omega})} \leq 1 + \frac{M_0^2}{\sigma_1}$, $\|a_8\|_{C(\bar{\Omega})} \leq \frac{M_0}{\sigma_1}$, $\|\partial_j a_7\|_{C(\bar{\Omega})} \leq \frac{3M_0^2 M_1}{\sigma_1^2}$, and $\|\partial_j a_8\|_{C(\bar{\Omega})} \leq \frac{2M_1 M_0}{\sigma_1^2}$, $j = 1, 2, \dots, n$.

$$2a_1 a_7 s \lambda \varphi |\partial_t w|^2 \geq 2\sigma_1 s \lambda \varphi |\partial_t w|^2, \tag{2.81}$$

$$2s \lambda \varphi a_1 \left\{ 3a_1 - (\nabla a_1 \cdot \nabla \psi) + \frac{a_1 (\nabla a_7 \cdot \nabla \psi)}{a_7} \right\} |\nabla w|^2 \geq 2\sigma_0 \sigma_1 s \lambda \varphi |\nabla w|^2, \tag{2.82}$$

and

$$4s^3 \lambda^4 \varphi^3 |\nabla \psi|^4 a_1^2 w^2 \geq 4\sigma_1^2 s^3 \lambda^4 \varphi^3 |\nabla \psi|^4 w^2, \text{ in } Q. \tag{2.83}$$

By the Cauchy-Schwarz inequality, we have, for any $\varepsilon_4 > 0$,

$$|-4s \lambda \varphi a_1 a_8 (\nabla \psi \cdot \nabla \vartheta) \partial_t w| \leq \frac{2M_0^2}{\sigma_1} s \lambda \varphi (\varepsilon_4 |\partial_t w|^2 + \varepsilon_4^{-1} |(\nabla \psi \cdot \nabla \vartheta)|^2), \text{ in } Q.$$

Choosing $\varepsilon_4 > 0$ small such that $\frac{2M_0^2}{\sigma_1} \varepsilon_4 = \sigma_1$, that is, $\varepsilon_4 = \frac{\sigma_1^2}{2M_0^2}$, we have

$$|-4s \lambda \varphi a_1 a_8 (\nabla \psi \cdot \nabla \vartheta) \partial_t w| \leq \sigma_1 s \lambda \varphi |\partial_t w|^2 + \frac{4M_0^4}{\sigma_1^3} s \lambda \varphi |(\nabla \psi \cdot \nabla \vartheta)|^2, \text{ in } Q. \tag{2.84}$$

Next we estimate the second term of (2.84).

$$\begin{aligned}
s \lambda \varphi |(\nabla \psi \cdot \nabla \vartheta)|^2 &= \frac{1}{a_5^2} s \lambda \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w) - a_6 (\nabla \psi \cdot \nabla w)|^2 \\
&\leq \frac{2}{a_5^2} s \lambda \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 + \frac{2a_6^2}{a_5^2} s \lambda \varphi |(\nabla \psi \cdot \nabla w)|^2 \\
&= \frac{2}{a_5^2} s \lambda \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 \\
&\quad + \frac{2a_6^2}{a_1^2 a_5^2} s \lambda \varphi |-a_1 (\nabla \psi \cdot \nabla w) + a_7 (\partial_t \psi) (\partial_t w) - a_7 (\partial_t \psi) (\partial_t w)|^2
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{2}{a_5^2} s \lambda \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 + \frac{4a_6^2 a_7^2}{a_1^2 a_5^2} s \lambda \varphi (\partial_t \psi)^2 |\partial_t w|^2 \\
&\quad + \frac{4a_6^2}{a_1^2 a_5^2} s \lambda \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 \\
&\leq \frac{4M_0^4 \beta^2 T^2}{\sigma_1^4} \left(1 + \frac{M_0^2}{\sigma_1}\right)^2 s \lambda \varphi |\partial_t w|^2 + C_{49} s \lambda \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 \\
&\quad + C_{49} s \lambda \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2. \tag{2.85}
\end{aligned}$$

By (2.84) and (2.85), we obtain

$$\begin{aligned}
&|-4s\lambda\varphi a_1 a_8 (\nabla \psi \cdot \nabla \vartheta) \partial_t w| \\
&\leq \sigma_1 s \lambda \varphi |\partial_t w|^2 + \frac{16M_0^8 \beta^2 T^2}{\sigma_1^7} \left(1 + \frac{M_0^2}{\sigma_1}\right)^2 s \lambda \varphi |\partial_t w|^2 \\
&\quad + C_{50} s \lambda \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 \\
&\quad + C_{50} s \lambda \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2, \text{ in } Q. \tag{2.86}
\end{aligned}$$

Furthermore, we have

$$|4s\lambda\varphi (\partial_t^2 \psi) a_7^2 |\partial_t w|^2| \leq 8 \left(1 + \frac{M_0^2}{\sigma_1}\right)^2 \beta s \lambda \varphi |\partial_t w|^2, \tag{2.87}$$

$$\begin{aligned}
&|4s\lambda\varphi (\partial_t \psi) a_7 (\nabla a_1 \cdot \nabla w) (\partial_t w)| \leq 4 \left(1 + \frac{M_0^2}{\sigma_1}\right) M_1 \beta T s \lambda \varphi \sum_{j=1}^n |\partial_t w| |\partial_j w| \\
&\leq 2 \left(1 + \frac{M_0^2}{\sigma_1}\right) M_1 \beta T s \lambda \varphi (n |\partial_t w|^2 + |\nabla w|^2), \tag{2.88}
\end{aligned}$$

$$|-4s\lambda\varphi (\partial_t \psi) a_1 (\nabla a_7 \cdot \nabla w) (\partial_t w)| \leq \frac{6M_0^3 M_1}{\sigma_1^2} \beta T s \lambda \varphi (n |\partial_t w|^2 + |\nabla w|^2), \tag{2.89}$$

$$\begin{aligned}
&|4s\lambda\varphi (\partial_t \psi) a_1 a_8 (\nabla \vartheta \cdot \nabla w)| \leq 4\beta T s \lambda \varphi |a_1 a_8| |\nabla \vartheta| |\nabla w| \\
&\leq \frac{2M_0^2}{\sigma_1} \beta T s \lambda \varphi (|\nabla \vartheta|^2 + |\nabla w|^2), \tag{2.90}
\end{aligned}$$

and

$$|-8s^3 \lambda^4 \varphi^3 (\partial_t \psi)^2 |\nabla \psi|^2 a_1 a_7 w^2| \leq 8M_0 \left(1 + \frac{M_0^2}{\sigma_1}\right) \beta^2 T^2 s^3 \lambda^4 \varphi^3 |\nabla \psi|^2 w^2, \tag{2.91}$$

in Q . Combining (2.80)–(2.83) and (2.86)–(2.91), we can obtain

$$\begin{aligned}
& \frac{1}{2} \|\mathcal{L}_*(\vartheta, w)\|_{L^2(Q)}^2 + \|\mathcal{L}_{2*}(w)\|_{L^2(Q)}^2 \\
& - 2s\lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_1^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt + 2 \int_{\Sigma} a_1 (\nabla a_1 \cdot \nu) \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt \\
& + 4s\lambda^2 \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 dx dt + \sigma_1 s\lambda \int_Q \varphi |\partial_t w|^2 dx dt \\
& + 2\sigma_0 \sigma_1 s\lambda \int_Q \varphi |\nabla w|^2 dx dt + 4\sigma_1^2 s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 w^2 dx dt \\
& \leq C_{51} \int_Q (f^2 + g^2) e^{2s\varphi} dx dt \\
& + C_{51} s\lambda \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dx dt \\
& + C_{51} s\lambda \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 dx dt \\
& + \frac{16M_0^8 \beta^2 T^2}{\sigma_1^7} \left(1 + \frac{M_0^2}{\sigma_1}\right)^2 s\lambda \int_Q \varphi |\partial_t w|^2 dx dt \\
& + 8 \left(1 + \frac{M_0^2}{\sigma_1}\right)^2 \beta s\lambda \int_Q \varphi |\partial_t w|^2 dx dt \\
& + 2 \left(1 + \frac{M_0^2}{\sigma_1} + \frac{3M_0^3}{\sigma_1^2}\right) M_1 \beta T s\lambda \int_Q \varphi (n |\partial_t w|^2 + |\nabla w|^2) dx dt \\
& + \frac{2M_0^2}{\sigma_1} \beta T s\lambda \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2) dx dt \\
& + 8M_0 \left(1 + \frac{M_0^2}{\sigma_1}\right) \beta^2 T^2 s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^2 w^2 dx dt \\
& + C_{51} s\lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla w|^2 + |\partial_t w|^2) dx dt + C_{51} s^3 \lambda^{\frac{7}{2}} \int_Q \varphi^3 (\vartheta^2 + w^2) dx dt \\
& + C_{51} \int_Q (|\nabla \vartheta|^2 + s^2 \lambda^4 \varphi^2 \vartheta^2 + s^2 \lambda^6 \varphi^2 w^2) dx dt. \tag{2.92}
\end{aligned}$$

We multiply (2.54) by α_3 and then add it to (2.92). By the definition of α_3 in (1.8), the second, third and fourth terms in the right hand side of (2.54) can be absorbed by the left hand side of (2.92), so that we arrive at

$$2\alpha_3 \sigma_1^2 s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 \vartheta^2 dx dt + \frac{\alpha_3}{60M} \int_Q \frac{1}{s\varphi} |\partial_t \vartheta|^2 dx dt$$

$$\begin{aligned}
& + \alpha_1 \alpha_3 s \lambda \int_Q \varphi |\nabla \vartheta|^2 dx dt \\
& + 2 \alpha_3 s \lambda^2 \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dx dt \\
& + 4 s \lambda^2 \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 dx dt \\
& + \frac{1}{2} \sigma_1 s \lambda \int_Q \varphi |\partial_t w|^2 dx dt + \sigma_0 \sigma_1 s \lambda \int_Q \varphi |\nabla w|^2 dx dt \\
& + 2 \sigma_1^2 s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^4 w^2 dx dt + \frac{1}{2} \|\mathcal{L}_*(\vartheta, w)\|_{L^2(Q)}^2 + \|\mathcal{L}_{2*}(w)\|_{L^2(Q)}^2 \\
& + \alpha_3 \|L_1(\vartheta, w)\|_{L^2(Q)}^2 - 2 \alpha_3 s \lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) \left(a_5 \frac{\partial \vartheta}{\partial \nu} + a_6 \frac{\partial w}{\partial \nu} \right)^2 d\sigma dt \\
& - 2 s \lambda \int_{\Sigma} \varphi (\nabla \psi \cdot \nu) a_1^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt + 2 \int_{\Sigma} a_1 (\nabla a_1 \cdot \nu) \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt \\
& \leq C_{52} \int_Q (f^2 + g^2) e^{2s\varphi} dx dt + C_{52} s \lambda \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dx dt \\
& + C_{52} s \lambda \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 dx dt \\
& + \beta T s \lambda \int_Q \varphi (\alpha_4 |\nabla \vartheta|^2 + \alpha_5 |\nabla w|^2 + \alpha_6 |\partial_t w|^2) dx dt \\
& + \alpha_7 \beta^2 T^2 s \lambda \int_Q \varphi |\partial_t w|^2 dx dt + 4 \alpha_8 \beta s \lambda \int_Q \varphi |\partial_t w|^2 dx dt \\
& + \frac{8 M_0 (\sigma_1 + M_0^2)}{\sigma_1} \beta^2 T^2 s^3 \lambda^4 \int_Q \varphi^3 |\nabla \psi|^2 w^2 dx dt \\
& + C_{52} s \lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2 + |\partial_t w|^2) dx dt + C_{52} s^3 \lambda^{\frac{7}{2}} \int_Q \varphi^3 (\vartheta^2 + w^2) dx dt \\
& + C_{52} \int_Q \left(\frac{1}{s \lambda \varphi} |\partial_t \vartheta|^2 + s^2 \lambda^6 \varphi^2 \vartheta^2 + s^2 \lambda^6 \varphi^2 w^2 \right) dx dt, \tag{2.93}
\end{aligned}$$

for all $s > s_1$. We note that $4m \leq |\nabla \psi|^2 \leq 4M$ on $\overline{\Omega}$. By (1.8) and (1.9), we have

$$2\mathcal{D}\alpha_6\sqrt{\beta} + 4\mathcal{D}^2\alpha_7\beta + 4\alpha_8\beta < \frac{1}{4}\sigma_1. \tag{2.94}$$

By (1.9) and (2.94), there exists a sufficiently small constant $\eta = \eta(\beta) > 0$ such that

$$0 < \sqrt{\beta}(\mathcal{D} + \eta) < \frac{\alpha_1\alpha_3}{4\alpha_4}, \quad 0 < \sqrt{\beta}(\mathcal{D} + \eta) < \frac{\sigma_0\sigma_1}{4\alpha_5},$$

$$2(\mathcal{D} + \eta) \alpha_6 \sqrt{\beta} + 4(\mathcal{D} + \eta)^2 \alpha_7 \beta + 4\alpha_8 \beta < \frac{1}{4} \sigma_1,$$

$$0 < \beta(\mathcal{D} + \eta)^2 < \frac{m^2 \sigma_1^3}{2M_0(\sigma_1 + M_0^2)}.$$

For any $T \in \left(0, \frac{2(\mathcal{D} + \eta)}{\sqrt{\beta}}\right)$, we have $0 < \beta T < 2\sqrt{\beta}(\mathcal{D} + \eta)$. Therefore, the fourth, fifth, sixth and seventh terms in the right hand side of (2.93) can be absorbed by the left hand side of it. Hence we arrive at

$$\begin{aligned} & 32\alpha_3 m^2 \sigma_1^2 s^3 \lambda^4 \int_Q \varphi^3 \vartheta^2 dx dt + \frac{\alpha_3}{60M} \int_Q \frac{1}{s\varphi} |\partial_t \vartheta|^2 dx dt \\ & + \frac{1}{2} \alpha_1 \alpha_3 s \lambda \int_Q \varphi |\nabla \vartheta|^2 dx dt + 2\alpha_3 s \lambda^2 \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dx dt \\ & + 4s \lambda^2 \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 dx dt \\ & + \frac{1}{4} \sigma_1 s \lambda \int_Q \varphi |\partial_t w|^2 dx dt + \frac{1}{2} \sigma_0 \sigma_1 s \lambda \int_Q \varphi |\nabla w|^2 dx dt \\ & + 16m^2 \sigma_1^2 s^3 \lambda^4 \int_Q \varphi^3 w^2 dx dt + \frac{1}{2} \|\mathcal{L}_*(\vartheta, w)\|_{L^2(Q)}^2 + \|\mathcal{L}_{2*}(w)\|_{L^2(Q)}^2 \\ & + \alpha_3 \|L_1(\vartheta, w)\|_{L^2(Q)}^2 \\ & - 2\alpha_3 s \lambda \int_{(\partial\Omega \setminus \Gamma_0) \times (0, T)} \varphi (\nabla \psi \cdot \nu) \left(a_5 \frac{\partial \vartheta}{\partial \nu} + a_6 \frac{\partial w}{\partial \nu} \right)^2 d\sigma dt \\ & - 2s \lambda \int_{(\partial\Omega \setminus \Gamma_0) \times (0, T)} \varphi (\nabla \psi \cdot \nu) a_1^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt \\ & - 2\sqrt{n} M_0 M_1 \int_{(\partial\Omega \setminus \Gamma_0) \times (0, T)} \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt \\ & \leq C_{53} \int_Q (f^2 + g^2) e^{2s\varphi} dx dt \\ & + 2\alpha_3 s \lambda \int_{\Gamma_0 \times (0, T)} \varphi (\nabla \psi \cdot \nu) \left(a_5 \frac{\partial \vartheta}{\partial \nu} + a_6 \frac{\partial w}{\partial \nu} \right)^2 d\sigma dt \\ & + 2s \lambda \int_{\Gamma_0 \times (0, T)} \varphi (\nabla \psi \cdot \nu) a_1^2 \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt + 2\sqrt{n} M_0 M_1 \int_{\Gamma_0 \times (0, T)} \left| \frac{\partial w}{\partial \nu} \right|^2 d\sigma dt \\ & + C_{53} s \lambda \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 dx dt \\ & + C_{53} s \lambda \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 dx dt \end{aligned}$$

$$\begin{aligned}
& + C_{53}s\lambda^{\frac{1}{2}} \int_Q \varphi (|\nabla \vartheta|^2 + |\nabla w|^2 + |\partial_t w|^2) \, dxdt + C_{53}s^3\lambda^{\frac{7}{2}} \int_Q \varphi^3 (\vartheta^2 + w^2) \, dxdt \\
& + C_{53} \int_Q \frac{1}{s\lambda\varphi} |\partial_t \vartheta|^2 \, dxdt + C_{53}s^2\lambda^6 \int_Q \varphi^2 (\vartheta^2 + w^2) \, dxdt,
\end{aligned}$$

for all $s > s_1$. By (1.14), there exists a constant $\epsilon > 0$ such that $-(\nabla \psi \cdot \nu) \geq \epsilon$ on $\partial\Omega \setminus \Gamma_0$. Therefore, there exists a constant $\lambda_0 > 1$ such that for any $\lambda \geq \lambda_0$, the fourteenth term in the left hand side can be absorbed by the thirteenth term in the left hand side, and the terms from the fifth to the ninth in the right hand side can be absorbed by the left hand side. Moreover, for any fixed $\lambda \geq \lambda_0$, there exists a constant $s_0 = s_0(\lambda) > s_1$ such that for any $s \geq s_0(\lambda)$, the tenth term in the right hand side can be absorbed by the left hand side. We further note that the twelfth term in the left hand side is positive. Thus there exists a constant $K_0 = K_0(s_0, \lambda_0, \beta, \Omega, T, m, M, M_0, M_1, M_2, \sigma_0, \sigma_1, \epsilon) > 0$ such that

$$\begin{aligned}
& \int_Q \left(s^3\lambda^4\varphi^3\vartheta^2 + \frac{1}{s\varphi} |\partial_t \vartheta|^2 + s\lambda\varphi |\nabla \vartheta|^2 + s^3\lambda^4\varphi^3w^2 + s\lambda\varphi |\nabla_{x,t} w|^2 \right) \, dxdt \\
& + s\lambda^2 \int_Q \varphi |a_5 (\nabla \psi \cdot \nabla \vartheta) + a_6 (\nabla \psi \cdot \nabla w)|^2 \, dxdt \\
& + s\lambda^2 \int_Q \varphi |a_7 (\partial_t \psi) (\partial_t w) - a_1 (\nabla \psi \cdot \nabla w)|^2 \, dxdt + \|\mathcal{L}_*(\vartheta, w)\|_{L^2(Q)}^2 \\
& + \|\mathcal{L}_{2*}(w)\|_{L^2(Q)}^2 + \|L_1(\vartheta, w)\|_{L^2(Q)}^2 + \epsilon s\lambda \int_{(\partial\Omega \setminus \Gamma_0) \times (0, T)} \varphi \left| \frac{\partial w}{\partial \nu} \right|^2 \, d\sigma dt \\
& \leq K_0 \int_Q (f^2 + g^2) e^{2s\varphi} \, dxdt + K_0 s\lambda \int_{\Gamma_0 \times (0, T)} \varphi \left(\left| \frac{\partial \vartheta}{\partial \nu} \right|^2 + \left| \frac{\partial w}{\partial \nu} \right|^2 \right) \, d\sigma dt,
\end{aligned}$$

for all $\lambda \geq \lambda_0$ and $s \geq s_0(\lambda)$. Noting $(\vartheta, w) = e^{s\varphi}(\Theta, p)$, we complete the proof of Theorem 1.2.

3 Proof of Theorem 1.1

We prove Theorem 1.1 by applying Theorem 1.2 and the argument in [12]. By a usual density argument, it is sufficient to prove (1.15) for $(\Theta, p) \in C^\infty(Q) \times C^\infty(Q)$. Let $G_\varepsilon = \{x \in \omega \mid \text{dist}(x, \partial\omega \cap \Omega) \leq \varepsilon\}$ and $\omega_\varepsilon = \omega \setminus G_\varepsilon$. Let $Q_{\omega_\varepsilon} = \omega_\varepsilon \times (0, T)$. By (1.5), there exists $\varepsilon > 0$ such that

$$\overline{\partial\Omega \setminus \partial\omega_{4\varepsilon}} \subset \{x \in \partial\Omega \mid ((x - x_0) \cdot \nu(x)) < 0\}. \quad (3.1)$$

We take $\Gamma_0 = \partial\omega_{4\varepsilon} \cap \partial\Omega$. By (3.1), Γ_0 satisfies (1.14). In order to apply Theorem 1.2, we take a cut-off function $\chi_1(x) \in C^\infty(\overline{\Omega})$ such that $0 \leq \chi_1(x) \leq 1$ for

all $x \in \overline{\Omega}$, $\chi_1(x) = 1$ for all $x \in \Omega \setminus \omega_{2\varepsilon}$, and $\chi_1(x) = 0$ for all $x \in \omega_{3\varepsilon}$. Let $u(x, t) = \chi_1(x)p(x, t)$ and $v(x, t) = \chi_1(x)\Theta(x, t)$. By (1.4), (1.11) and (1.12), we have

$$\begin{cases} \partial_t^2 u - a_1 \Delta u - a_2 \Delta v = \chi_1 f - a_1 (\Delta \chi_1) p - 2a_1 (\nabla \chi_1 \cdot \nabla p) \\ \quad - a_2 (\Delta \chi_1) \Theta - 2a_2 (\nabla \chi_1 \cdot \nabla \Theta), \\ \partial_t v - a_3 \Delta v - a_4 \partial_t^2 u = \chi_1 g - a_3 (\Delta \chi_1) \Theta - 2a_3 (\nabla \chi_1 \cdot \nabla \Theta), \end{cases} \text{ in } Q,$$

$v(x, t) = u(x, t) = 0$ for $(x, t) \in \Sigma$, and $v(x, 0) = v(x, T) = \partial_t^j u(x, 0) = \partial_t^j u(x, T) = 0$ for $x \in \overline{\Omega}$ and $j = 0, 1$. Furthermore, by the definition of χ_1 , we have $\nabla v = \nabla u = 0$ in some neighborhood of $\Gamma_0 \times (0, T)$. By Theorem 1.2, there exists a constant $\eta(\beta) > 0$ such that for any $T \in \left(0, \frac{2(\mathcal{D}+\eta)}{\sqrt{\beta}}\right)$, there exists a constant $\lambda_0 > 0$ such that for all $\lambda > \lambda_0$, there exist constants $s_0(\lambda) > 0$ and $K = K(s_0, \lambda_0, \beta, \Omega, T, m, M, M_0, M_1, M_2, \sigma_0, \sigma_1) > 0$ such that

$$\begin{aligned} & \int_Q \left(s^3 \lambda^4 \varphi^3 v^2 + \frac{1}{s\varphi} |\partial_t v|^2 + s\lambda\varphi |\nabla v|^2 + s^3 \lambda^4 \varphi^3 u^2 + s\lambda\varphi |\nabla_{x,t} u|^2 \right) e^{2s\varphi} dx dt \\ & \leq K \int_Q (f^2 + g^2) e^{2s\varphi} dx dt + K \int_{(\omega_{2\varepsilon} \setminus \omega_{3\varepsilon}) \times (0, T)} \left\{ |(\Delta \chi_1) p|^2 + |(\nabla \chi_1 \cdot \nabla p)|^2 \right. \\ & \quad \left. + |(\Delta \chi_1) \Theta|^2 + |(\nabla \chi_1 \cdot \nabla \Theta)|^2 \right\} e^{2s\varphi} dx dt, \quad \text{for all } s \geq s_0. \end{aligned}$$

Therefore we have

$$\begin{aligned} & \int_Q \left\{ s^3 \lambda^4 \varphi^3 (\Theta^2 + p^2) + \frac{1}{s\varphi} |\partial_t \Theta|^2 + s\lambda\varphi (|\nabla \Theta|^2 + |\nabla_{x,t} p|^2) \right\} e^{2s\varphi} dx dt \\ & = \int_{Q \setminus Q_{\omega_{2\varepsilon}}} \left\{ s^3 \lambda^4 \varphi^3 (v^2 + u^2) + \frac{1}{s\varphi} |\partial_t v|^2 + s\lambda\varphi (|\nabla v|^2 + |\nabla_{x,t} u|^2) \right\} e^{2s\varphi} dx dt \\ & \quad + \int_{Q_{\omega_{2\varepsilon}}} \left\{ s^3 \lambda^4 \varphi^3 (\Theta^2 + p^2) + \frac{1}{s\varphi} |\partial_t \Theta|^2 + s\lambda\varphi (|\nabla \Theta|^2 + |\nabla_{x,t} p|^2) \right\} e^{2s\varphi} dx dt \\ & \leq K \int_Q (f^2 + g^2) e^{2s\varphi} dx dt + K \int_{(\omega_{2\varepsilon} \setminus \omega_{3\varepsilon}) \times (0, T)} \left\{ |(\Delta \chi_1) p|^2 + |(\nabla \chi_1 \cdot \nabla p)|^2 \right. \\ & \quad \left. + |(\Delta \chi_1) \Theta|^2 + |(\nabla \chi_1 \cdot \nabla \Theta)|^2 \right\} e^{2s\varphi} dx dt \\ & \quad + \int_{Q_{\omega_{2\varepsilon}}} \left\{ s^3 \lambda^4 \varphi^3 (\Theta^2 + p^2) + \frac{1}{s\varphi} |\partial_t \Theta|^2 + s\lambda\varphi (|\nabla \Theta|^2 + |\nabla_{x,t} p|^2) \right\} e^{2s\varphi} dx dt \\ & \leq K \int_Q (f^2 + g^2) e^{2s\varphi} dx dt + C_{54} \int_{Q_{\omega_{2\varepsilon}}} \left\{ s^3 \lambda^4 \varphi^3 (\Theta^2 + p^2) + \frac{1}{s\lambda\varphi} |\partial_t \Theta|^2 \right. \\ & \quad \left. + s\lambda\varphi (|\nabla \Theta|^2 + |\nabla_{x,t} p|^2) \right\} e^{2s\varphi} dx dt, \quad \text{for all } s \geq s_0. \end{aligned} \tag{3.2}$$

Next we estimate $s\lambda \int_{Q_{\omega_{2\varepsilon}}} \varphi (|\nabla\Theta|^2 + |\nabla p|^2) e^{2s\varphi} dxdt$ in (3.2). We take a cut-off function $\chi_2(x) \in C^\infty(\overline{\Omega})$ such that $0 \leq \chi_2(x) \leq 1$ for all $x \in \overline{\Omega}$, $\chi_2(x) = 0$ for all $x \in \Omega \setminus \omega_\varepsilon$, and $\chi_2(x) = 1$ for all $x \in \overline{\omega_{2\varepsilon}}$. We multiply the second equation in (1.4) by $s\lambda\varphi\chi_2\Theta e^{2s\varphi}$ and integrate it over Q . Integrating by parts and using (1.11) and (1.12), we have

$$\begin{aligned}
s\lambda \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 g \Theta e^{2s\varphi} dxdt &= \frac{1}{2} s\lambda \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 \{ \partial_t (\Theta^2) \} e^{2s\varphi} dxdt \\
&\quad - s\lambda \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 a_3 (\Delta \Theta) \Theta e^{2s\varphi} dxdt - s\lambda \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 a_4 (\partial_t^2 p) \Theta e^{2s\varphi} dxdt \\
&= -\frac{1}{2} s\lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 (\partial_t \psi) \Theta^2 e^{2s\varphi} dxdt - s^2 \lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi^2 (\partial_t \psi) \chi_2 \Theta^2 e^{2s\varphi} dxdt \\
&\quad + s\lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 a_3 (\nabla \psi \cdot \nabla \Theta) \Theta e^{2s\varphi} dxdt \\
&\quad + s\lambda \int_{Q_{\omega_\varepsilon}} \varphi (\nabla (\chi_2 a_3) \cdot \nabla \Theta) \Theta e^{2s\varphi} dxdt \\
&\quad + s\lambda \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 a_3 |\nabla \Theta|^2 e^{2s\varphi} dxdt + 2s^2 \lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi^2 \chi_2 a_3 (\nabla \psi \cdot \nabla \Theta) \Theta e^{2s\varphi} dxdt \\
&\quad + s\lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi (\partial_t \psi) \chi_2 a_4 (\partial_t p) \Theta e^{2s\varphi} dxdt + s\lambda \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 a_4 (\partial_t p) (\partial_t \Theta) e^{2s\varphi} dxdt \\
&\quad + 2s^2 \lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi^2 (\partial_t \psi) \chi_2 a_4 (\partial_t p) \Theta e^{2s\varphi} dxdt \triangleq \sum_{k=1}^{10} \tilde{J}_k, \quad \text{for all } s \geq s_0. \quad (3.3)
\end{aligned}$$

We recall $s > s_0 \geq 1$ and $\lambda > 1$. Thus we have

$$\tilde{J}_1 + \tilde{J}_2 \geq -C_{55} s^2 \lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi^2 \Theta^2 e^{2s\varphi} dxdt, \quad \text{for all } s \geq s_0.$$

By the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
\tilde{J}_3 &\geq -C_{56} \int_{Q_{\omega_\varepsilon}} \chi_2 \left(\frac{1}{2} s^{\frac{1}{2}} \lambda^{\frac{1}{2}} \sigma_1^{\frac{1}{2}} \varphi^{\frac{1}{2}} |\nabla \Theta| \right) \left(2s^{\frac{1}{2}} \lambda^{\frac{3}{2}} \sigma_1^{-\frac{1}{2}} \varphi^{\frac{1}{2}} |\Theta| \right) e^{2s\varphi} dxdt \\
&\geq -\frac{1}{4} s\lambda \sigma_1 \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 |\nabla \Theta|^2 e^{2s\varphi} dxdt - C_{57} s\lambda^3 \int_{Q_{\omega_\varepsilon}} \varphi \chi_2 \Theta^2 e^{2s\varphi} dxdt,
\end{aligned}$$

for all $s \geq s_0$. Integrating by parts, we have

$$\tilde{J}_4 = \frac{1}{2} s\lambda \int_{Q_{\omega_\varepsilon}} \varphi (\nabla (\chi_2 a_3) \cdot \nabla (\Theta^2)) e^{2s\varphi} dxdt$$

$$\begin{aligned}
&= -\frac{1}{2}s\lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi (\nabla (\chi_2 a_3) \cdot \nabla \psi) \Theta^2 e^{2s\varphi} dxdt \\
&\quad - \frac{1}{2}s\lambda \int_{Q_{\omega_\varepsilon}} \varphi \{\Delta (\chi_2 a_3)\} \Theta^2 e^{2s\varphi} dxdt \\
&\quad - s^2\lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi^2 (\nabla (\chi_2 a_3) \cdot \nabla \psi) \Theta^2 e^{2s\varphi} dxdt \\
&\geq -C_{58}s^2\lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi^2 \Theta^2 e^{2s\varphi} dxdt.
\end{aligned}$$

$$\tilde{J}_5 \geq s\lambda\sigma_1 \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 |\nabla\Theta|^2 e^{2s\varphi} dxdt, \text{ for all } s \geq s_0.$$

By the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
\tilde{J}_6 &\geq -C_{59} \int_{Q_{\omega_\varepsilon}} \chi_2 \left(\frac{1}{2}s^{\frac{1}{2}}\lambda^{\frac{1}{2}}\sigma_1^{\frac{1}{2}}\varphi^{\frac{1}{2}}|\nabla\Theta| \right) \left(2s^{\frac{3}{2}}\lambda^{\frac{3}{2}}\sigma_1^{-\frac{1}{2}}\varphi^{\frac{3}{2}}|\Theta| \right) e^{2s\varphi} dxdt \\
&\geq -\frac{1}{4}s\lambda\sigma_1 \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 |\nabla\Theta|^2 e^{2s\varphi} dxdt - C_{60}s^3\lambda^3 \int_{Q_{\omega_\varepsilon}} \varphi^3\chi_2\Theta^2 e^{2s\varphi} dxdt,
\end{aligned}$$

$$\tilde{J}_7 + \tilde{J}_9 \geq -C_{61}s^2\lambda^2 \int_{Q_{\omega_\varepsilon}} \varphi^2\chi_2 (|\partial_t p|^2 + \Theta^2) e^{2s\varphi} dxdt,$$

$$\begin{aligned}
\tilde{J}_8 &\geq -C_{62} \int_{Q_{\omega_\varepsilon}} \chi_2 \left(s^{\frac{3}{2}}\lambda^{\frac{3}{2}}\varphi^{\frac{3}{2}}|\partial_t p| \right) \left(\frac{1}{s^{\frac{1}{2}}\lambda^{\frac{1}{2}}\varphi^{\frac{1}{2}}}|\partial_t\Theta| \right) e^{2s\varphi} dxdt \\
&\geq -C_{63}s^3\lambda^3 \int_{Q_{\omega_\varepsilon}} \varphi^3\chi_2 |\partial_t p|^2 e^{2s\varphi} dxdt - \int_{Q_{\omega_\varepsilon}} \frac{\chi_2}{s\lambda\varphi} |\partial_t\Theta|^2 e^{2s\varphi} dxdt,
\end{aligned}$$

$$\begin{aligned}
s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 g\Theta e^{2s\varphi} dxdt &\leq C_{64} \int_{Q_{\omega_\varepsilon}} \chi_2 \left(\frac{1}{s^{\frac{1}{2}}\lambda^{\frac{1}{2}}\varphi^{\frac{1}{2}}}|g| \right) \left(s^{\frac{3}{2}}\lambda^{\frac{3}{2}}\varphi^{\frac{3}{2}}|\Theta| \right) e^{2s\varphi} dxdt \\
&\leq \int_{Q_{\omega_\varepsilon}} \frac{\chi_2}{s\lambda\varphi} g^2 e^{2s\varphi} dxdt + C_{65}s^3\lambda^3 \int_{Q_{\omega_\varepsilon}} \varphi^3\chi_2\Theta^2 e^{2s\varphi} dxdt,
\end{aligned}$$

for all $s \geq s_0$. Therefore, we obtain

$$\begin{aligned}
&\frac{1}{2}s\lambda\sigma_1 \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 |\nabla\Theta|^2 e^{2s\varphi} dxdt \\
&\leq \int_{Q_{\omega_\varepsilon}} \frac{\chi_2}{s\lambda\varphi} |\partial_t\Theta|^2 e^{2s\varphi} dxdt + \int_{Q_{\omega_\varepsilon}} \frac{\chi_2}{s\lambda\varphi} g^2 e^{2s\varphi} dxdt \\
&\quad + C_{66}s^3\lambda^3 \int_{Q_{\omega_\varepsilon}} \varphi^3 (|\partial_t p|^2 + \Theta^2) e^{2s\varphi} dxdt, \text{ for all } s \geq s_0. \tag{3.4}
\end{aligned}$$

By (1.4), we can get (2.55) where a_7 and a_8 are given by (2.56). Similarly to (3.3)–(3.4), we multiply (2.55) by $s\lambda\varphi\chi_2pe^{2s\varphi}$ and integrate it over Q , so that, by the integration by parts and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 (f - a_8g) pe^{2s\varphi} dxdt &= s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 a_7 (\partial_t^2 p) pe^{2s\varphi} dxdt \\ &\quad - s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 a_1 (\Delta p) pe^{2s\varphi} dxdt - s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 a_8 (\partial_t \Theta) pe^{2s\varphi} dxdt \\ &\geq -C_{67}s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 |\partial_t p|^2 e^{2s\varphi} dxdt - C_{67}s^3\lambda^3 \int_{Q_{\omega_\varepsilon}} \varphi^3 p^2 e^{2s\varphi} dxdt \\ &\quad + \frac{1}{2}s\lambda\sigma_1 \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 |\nabla p|^2 e^{2s\varphi} dxdt - C_{67}s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 \Theta^2 e^{2s\varphi} dxdt, \end{aligned}$$

for all $s \geq s_0$. Then we have

$$\begin{aligned} &\frac{1}{2}s\lambda\sigma_1 \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 |\nabla p|^2 e^{2s\varphi} dxdt \\ &\leq \int_{Q_{\omega_\varepsilon}} \frac{\chi_2}{s\lambda\varphi} (f^2 + g^2) e^{2s\varphi} dxdt + C_{68}s^3\lambda^3 \int_{Q_{\omega_\varepsilon}} \varphi^3 p^2 e^{2s\varphi} dxdt \\ &\quad + C_{68}s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 |\partial_t p|^2 e^{2s\varphi} dxdt + C_{68}s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 \Theta^2 e^{2s\varphi} dxdt, \end{aligned} \quad (3.5)$$

for all $s \geq s_0$. Adding (3.4) and (3.5) and noting $\omega_{2\varepsilon} \subseteq \omega_\varepsilon \subseteq \omega$ and $\chi_2(x) = 1$ for all $x \in \overline{\omega_{2\varepsilon}}$, we obtain

$$\begin{aligned} s\lambda \int_{Q_{\omega_{2\varepsilon}}} \varphi (|\nabla \Theta|^2 + |\nabla p|^2) e^{2s\varphi} dxdt &\leq s\lambda \int_{Q_{\omega_\varepsilon}} \varphi\chi_2 (|\nabla \Theta|^2 + |\nabla p|^2) e^{2s\varphi} dxdt \\ &\leq \frac{2}{\sigma_1} \int_{Q_{\omega_\varepsilon}} \frac{\chi_2}{s\lambda\varphi} |\partial_t \Theta|^2 e^{2s\varphi} dxdt + \frac{4}{\sigma_1} \int_{Q_{\omega_\varepsilon}} \frac{\chi_2}{s\lambda\varphi} (f^2 + g^2) e^{2s\varphi} dxdt \\ &\quad + C_{69}s^3\lambda^3 \int_{Q_{\omega_\varepsilon}} \varphi^3 (|\partial_t p|^2 + p^2 + \Theta^2) e^{2s\varphi} dxdt, \text{ for all } s \geq s_0. \end{aligned} \quad (3.6)$$

Substituting (3.6) into (3.2), we obtain

$$\begin{aligned} &\int_Q \left\{ s^3\lambda^4\varphi^3 (\Theta^2 + p^2) + \frac{1}{s\varphi} |\partial_t \Theta|^2 + s\lambda\varphi (|\nabla \Theta|^2 + |\nabla_{x,t} p|^2) \right\} e^{2s\varphi} dxdt \\ &\leq C_{70} \int_Q (f^2 + g^2) e^{2s\varphi} dxdt \\ &\quad + C_{70} \int_{Q_{\omega_\varepsilon}} \left\{ s^3\lambda^4\varphi^3 (\Theta^2 + p^2) + \frac{1}{s\varphi} |\partial_t \Theta|^2 + s^3\lambda^3\varphi^3 |\partial_t p|^2 \right\} e^{2s\varphi} dxdt, \end{aligned} \quad (3.7)$$

for all $s \geq s_0$. Noting $\omega_\varepsilon \subseteq \omega$, we obtain (1.13). The proof of Theorem 1.1 is complete.

Acknowledgements

This paper has been partially done during the stay of Li S at the Institut de Mathématiques de Marseille, Aix-Marseille University, France, in July 2019, and the stay was supported by Aix-Marseille University.

References

- [1] H. Akhouayri, M. Bergounioux, A. D. Silva, P. Elbau, A. Litman and L. Min-drinos, Quantitative thermoacoustic tomography with microwaves sources, *J. Inverse Ill-Posed Probl.* 25 (2016), 703-717.
- [2] M. Bellassoued and M. Yamamoto, *Carleman Estimates and Applications to Inverse Problems for Hyperbolic Systems*, Springer, Tokyo, 2018.
- [3] T. Carleman, Sur un problème d'unicité pour les systmes d'équations aux dérivées partielles à deux variables indépendentes [On a problem of the uniqueness for systems of partial differential equations with two independent variables], *Ark. Mat.Astr. Fys.* 26B (1939), 1-9.
- [4] D. Chae, O. Y. Imanuvilov and S. M. Kim, Exact controllability for semilinear parabolic equations with Neumann boundary conditions, *J. Dyn. Control Syst.* 2 (1996), 449-483.
- [5] B. Cox and P. C. Beard, Modeling photoacoustic propagation in tissue using k-space techniques, In: Wang L V, editor, *Photoacoustic Imaging and Spectroscopy*, chapter 3, pp. 25-34, CRC Press, 2009.
- [6] M. Cristofol, S. Li and Y. Shang, Carleman estimates and inverse problems for the coupled quantitative thermoacoustic equations. Part II: inverse problems, preprint, 2020.
- [7] Y. V. Egorov, *Linear Differential Equations of Principal Type*, Consultants Bureau, New York, 1986.
- [8] M. M. Eller and V. Isakov, Carleman estimates with two large parameters and applications, *Contemp. Math.* 268 (2000), 117-136.
- [9] A. V. Fursikov and O. Y. Imanuvilov, *Controllability of Evolution Equations*, Lecture Notes Series vol 34, Seoul National University, Korea, 1996.
- [10] L. Hörmander, *Linear Partial Differential Operators*, Springer, Berlin, 1963.

- [11] L. Hörmander, The Analysis of Linear Partial Differential Operators I-IV, Springer, Berlin, 1985.
- [12] O. Y. Imanuvilov, On Carleman estimates for hyperbolic equations, *Asymptot. Anal.* 32 (2002), 185-220.
- [13] O. Y. Imanuvilov and M. Yamamoto, Carleman estimate for a stationary isotropic Lamé system and the applications, *Appl. Anal.* 83 (2004), 243-270.
- [14] V. Isakov, A nonhyperbolic Cauchy problem for $\square_b \square_c$ and its applications to elasticity theory, *Comm. Pure Appl. Math.* 39 (1986), 747-767.
- [15] V. Isakov, Inverse Source Problems, American Mathematical Society, Providence, RI, 1990.
- [16] V. Isakov, Carleman type estimates in an anisotropic case and applications, *J. Differential Equations* 105 (1993), 217-238.
- [17] V. Isakov, Carleman estimates and applications to inverse problems, *Milan J. Math.* 72 (2004), 249-271.
- [18] V. Isakov, Inverse problems for partial differential equations, Springer-Verlag, Berlin, 2006.
- [19] M. V. Klibanov, Inverse problems and Carleman estimates, *Inverse Problems* 8 (1992), 575-596.
- [20] M. V. Klibanov and A. Timonov, Carleman estimates for coefficient inverse problems and numerical applications, VSP, Utrecht, 2004.
- [21] M. M. Lavrent'ev, V. G. Romanov and S. P. Shishatskii, Ill-posed problems of mathematical physics and analysis, American Mathematical Society, Providence, RI, 1986.
- [22] S. Li, Carleman estimates for second-order hyperbolic systems in anisotropic cases and applications. Part I: Carleman estimates, *Appl. Anal.* 94(2015), 2261-2286.
- [23] S. K. Patch and O. Scherzer, Guest editors' introduction: Photo-and thermo-acoustic imaging, *Inverse Problems* 23 (2007), S1-S10.
- [24] V. G. Romanov, Carleman estimates for second-order hyperbolic equations, *Sib. Math. J.* 47 (2006) 135-151.

- [25] P. Stefanov and G. Uhlmann, Thermoacoustic tomography with variable sound speed, *Inverse Problems* 25 (2009), 075011.
- [26] D. Tataru, Carleman estimates and unique continuation for solutions to boundary value problems, *J. Math. Pures Appl.* 75 (1996), 367-408.
- [27] F. Trèves, *Linear Partial Differential Equations*, Gordon and Breach, New York, 1970.
- [28] M. Yamamoto, Carleman estimates for parabolic equations and applications, *Inverse Problems* 25 (2009), 123013.
- [29] G. Yuan and M. Yamamoto, Lipschitz stability in the determination of the principal part of a parabolic equation, *ESAIM Control Optim. Calc. Var.* 15 (2009), 525-554.