# CONVERGENCE ANALYSIS OF THE SCALED BOUNDARY FINITE ELEMENT METHOD FOR THE LAPLACE EQUATION

#### FLEURIANNE BERTRAND, DANIELE BOFFI, AND GONZALO G. DE DIEGO

ABSTRACT. The scaled boundary finite element method (SBFEM) is a relatively recent boundary element method that allows the approximation of solutions to PDEs without the need of a fundamental solution. A theoretical framework for the convergence analysis of SBFEM is proposed here. This is achieved by defining a space of semi-discrete functions and constructing an interpolation operator onto this space. We prove error estimates for this interpolation operator and show that optimal convergence to the solution can be obtained in SBFEM. These theoretical results are backed by a numerical example.

#### 1. INTRODUCTION

The scaled boundary finite element method (SBFEM), first proposed by Song and Wolf, is a boundary element method that does not require a fundamental solution. It has proven to be particularly effective for problems with singularities or posed over unbounded media, see [6, 7, 9]. In SBFEM, a semi-analytical (or semidiscrete, as we shall call it here) solution to a PDE is constructed by transforming the weak formulation of the PDE into an ODE. Essentially, given a star-shaped domain  $\Omega \subset \mathbb{R}^n$ , a coordinate transformation is performed (the scaled boundary transformation) in terms of a radial variable and n-1 circumferential variables. Then, an approximate solution is sought in a space of functions discretized solely in the circumferential direction. The resulting weak formulation posed over this space is then transformed into an ODE which, under certain circumstances, can be solved exactly, yielding a semi-analytical approximation of the solution to the PDE.

The SBFEM has been applied to a wide range of problems that arise in science and engineering, such as crack propagation [8] and acoustic-structure interactions [3]. Moreover, the limitation to star-shaped domains has been overcome with the development of scaled-boundary polygon elements, in which the domain is broken into arbitrarily shaped polygons and shape functions are constructed over these polygons based on SBFEM [5, 4, 1].

The first author gratefully acknowledges support by the German Research Foundation (DFG) in the Priority Programme SPP 1748 Reliable simulation techniques in solid mechanics under grant number BE6511/1-1. The second author is a member of the INdAM Research group GNCS and his research is partially supported by IMATI/CNR and by PRIN/MIUR. The first and third author gratefully acknowledge the support by Mercator Research Center Ruhr (MERCUR) under grant Pr-2017-0017. We would also like to thank the project partners Prof. Carolin Birk (Universität Duisburg-Essen, Germany) and Prof. Christian Meyer (TU Dortmund, Germany) as well as Professor Gerhard Starke for the fruitful discussions.



FIGURE 1. Sector of a disk of angle  $\Theta$ .

The objective of this paper is to introduce a rigorous framework in which the error of the approximate solution obtained by SBFEM can be estimated. In particular, the notion of a semi-discrete solution to a PDE is formalized by defining a space of semi-discrete functions and constructing an interpolation operator onto this space. Then, given a semi-analytical solution obtained in the framework of SBFEM, estimates of its error can be obtained by bounding the interpolation operator's error using Céa's lemma. We limit the analysis to Poisson's equation posed on a circular domain for simplicity; this setting is appropriate to highlight the main features of our theoretical setting.

The overview of this paper is as follows: in Section 2 we describe the continuous problem together with the polar coordinate change of variables. In Section 3 we introduce a semi-discretization of our problem, where the domain is discretized only in the angular coordinate. It is shown that the semi-discrete solution converges optimally to the continuous solution. Section 4, making use of the semidiscretization, transforms the original problem into an ODE. Finally, two numerical results reported in Section 5 show that the method is performing optimally also in the presence of singularities.

# 2. Setting of the problem

Given an angle  $\Theta$  in  $(0, 2\pi)$ , we are considering the Poisson problem in the following circular sector (see Figure 1):

$$\Omega := \left\{ (x,y) \in \mathbb{R}^2 : 0 < x^2 + y^2 < 1, \ 0 < \arctan\left(\frac{y}{x}\right) < \Theta \right\}.$$

Since we are going to consider a change of variables when defining the scaled boundary method, we denote with  $\hat{\bullet}$  (with the hat symbol) quantities defined on  $\Omega$ that correspond to quantities  $\bullet$  defined on the reference domain. Hence our problem reads: find  $\hat{u} : \Omega \to \mathbb{R}$  such that

(1) 
$$\begin{aligned} & -\hat{\Delta}\hat{u} = \hat{f} \quad \text{in } \Omega\\ & \hat{u} = 0 \qquad \text{on } \partial\Omega. \end{aligned}$$

where  $\hat{f} \in L^2(\Omega)$  and  $\hat{\Delta} = \partial_x^2 + \partial_y^2$  is the Laplace operator in the Cartesian coordinates (x, y).

Let the curved part of the boundary of  $\Omega$  be parametrized by the graph

$$(x_b(\theta), y_b(\theta)) = (\cos \theta, \sin \theta) \qquad \theta \in (0, \Theta)$$

 $\mathbf{2}$ 

and define the open rectangle  $Q := (0,1) \times (0,\Theta)$ . We consider the mapping  $F: Q \to \mathbb{R}^2$  given by

(2) 
$$F(r,\theta) = r\left(x_b(\theta), y_b(\theta)\right).$$

In this particular case, the scaled boundary transformation is given by the change of variables  $(r, \theta) = F^{-1}(x, y)$  for  $(x, y) \in \Omega$ , i.e., by the polar coordinate transformation. The Jacobian of F is given by

$$DF(r,\theta) = \begin{pmatrix} \partial_r x & \partial_r y \\ \partial_\theta x & \partial_\theta y \end{pmatrix} = \begin{pmatrix} \cos\theta & \sin\theta \\ -r\sin\theta & r\cos\theta \end{pmatrix}$$

and its determinant is  $|DF(r, \theta)| = r$ . Since F is differentiable and  $|DF(r, \theta)|$  is invertible in the open set Q we have

$$DF^{-1}(F(r,\theta)) = \begin{pmatrix} \partial_x r & \partial_x \theta \\ \partial_y r & \partial_y \theta \end{pmatrix} = \frac{1}{r} \begin{pmatrix} r\cos\theta & -\sin\theta \\ r\sin\theta & \cos\theta \end{pmatrix}.$$

Let  $u(r,\theta) = \hat{u}(F(r,\theta))$ , then the relation between the gradient in Cartesian coordinates  $\hat{\nabla} = (\partial_x, \partial_y)^\top$  and the gradient in polar coordinates  $\nabla = (\partial_r, \partial_\theta)^\top$  is given by

$$\hat{\nabla}\hat{u}(x,y) = DF^{-1}(x,y)\nabla u(F^{-1}(x,y)) \quad \text{in } \Omega.$$

Moreover, the solution  $\hat{u}$  of (1) satisfies

(3) 
$$||\hat{u}||^{2}_{H^{1}(\Omega)} = \int_{0}^{1} \int_{0}^{\Theta} \left( ru^{2} + r(\partial_{r}u)^{2} + \frac{1}{r}(\partial_{\theta}u)^{2} \right) dr d\theta < \infty.$$

In order to consider the variational formulation of (1) in polar coordinates we have to consider appropriate weighted functional spaces. While this is pretty straightforward and well understood, we present the procedure in detail since the notation will be useful for the analysis of the numerical approximation.

Given a weight function  $w(r, \theta)$  in Q, we define the weighted Lebesgue space

$$L^2_w(Q) = \left\{ v : Q \to \mathbb{R} \text{ measurable} : \int_0^1 \int_0^\Theta v^2 w \, \mathrm{d}r \, \mathrm{d}\theta < \infty \right\}$$

with inner product

$$(u,v)_{L^2_w(Q)} := \int_0^1 \int_0^\Theta uvw \,\mathrm{d}r \,\mathrm{d}\theta.$$

We will use in particular w = r and w = 1/r; it is not difficult to see that we have  $||u||_{L^2(Q)} \leq ||u||_{L^2(Q)} \leq ||u||_{L^2_{1/r}(Q)}$  for all  $u \in L^2_{1/r}(Q)$ . Furthermore, these spaces are complete [2].

The bound (3) motivates the definition of the following weighted Sobolev space

$$\tilde{H}^{1}(Q) = \left\{ v \in L^{2}_{r}(Q) : ||v||_{L^{2}_{r}(Q)} + ||\partial_{r}v||_{L^{2}_{r}(Q)} + ||\partial_{\theta}u||_{L^{2}_{1/r}(Q)} < \infty \right\}$$

with inner product

$$(u,v)_{\tilde{H}^{1}(Q)} := (u,v)_{L^{2}_{r}(Q)} + (\partial_{r}u,\partial_{r}v)_{L^{2}_{r}(Q)} + (\partial_{\theta}u,\partial_{\theta}v)_{L^{2}_{1/r}(Q)}.$$

The following lemma shows that  $H^1(\Omega)$  and  $\tilde{H}^1(Q)$  are isometric.

**Lemma 1.** Let  $\Phi : L^2(Q) \to L^2_r(Q)$  be defined by  $\hat{u} \mapsto \hat{u} \circ F$ . Then the spaces  $H^1(\Omega)$  and  $\tilde{H}^1(Q)$  are isometric via  $\Phi$ .

*Proof.* Let  $\hat{u} \in H^1(\Omega)$  and, for  $0 < \rho < 1$ , let  $B_\rho$  be the ball of radius  $\rho$  centred at the origin and  $B_\rho^c$  its complement. For  $\Omega_\rho = \Omega \cap B_\rho^c$ , the map  $F : Q_\rho \to \Omega_\rho$  with  $Q_\rho := (\rho, 1) \times (0, \Theta)$  is a bi-Lipschitz map, i.e. there exist two constants  $C_1, C_2 > 0$  such that

$$C_1 \left| (r_1, \theta_1) - (r_2, \theta_2) \right| \le \left| F(r_1, \theta_1) - F(r_2, \theta_2) \right| \le C_2 \left| (r_1, \theta_1) - (r_2, \theta_2) \right|$$

holds for all  $(r_1, \theta_1), (r_2, \theta_2) \in Q_{\rho}$ . Indeed, by the mean value theorem we have

$$F(r_1, \theta_1) - F(r_2, \theta_2) \le \|\nabla F\| |(r_1, \theta_1) - (r_2, \theta_2)|$$

and clearly  $\|\nabla F\|_{\infty} \leq 1$ . In the same way,  $F^{-1}: \Omega_{\rho} \to Q_{\rho}$  is a smooth, bijective map and  $\|F^{-1}\|_{\infty} \leq 1/\rho$ ; hence it is Lipschitz continuous and it follows that F and  $F^{-1}$  are bi-Lipschitz when restricted to  $Q_{\rho}$  and  $\Omega_{\rho}$  respectively. As a result of [10, Theorem 2.2.2.],  $u = \Phi(\hat{u})$  is weakly differentiable on  $Q_{\rho}$  and the chain rule holds. For  $n \in \mathbb{N}$ , define  $u_n = \Phi|_{\Omega_{1/n}}(\hat{u})$  on Q by extending  $\hat{u}$  by zero outside  $\Omega_{\rho}$ . For any  $0 < \rho < 1$  one has that

(4) 
$$\|u\|_{\tilde{H}^{1}(Q_{\rho})} = \|\hat{u}\|_{H^{1}(\Omega_{\rho})} \le \|\hat{u}\|_{H^{1}(\Omega)},$$

so  $u_n$  and its derivatives belong to the associated weighted Lebesgue spaces. As a result of the monotone convergence theorem we have that  $u \in \tilde{H}^1(Q)$  and

$$||u||_{\tilde{H}^1(Q)} = ||\hat{u}||_{H^1(\Omega)}$$

Repeating the steps above, we can also show that for  $u \in \tilde{H}^1(Q)$  one has  $\Phi^{-1}(u) \in H^1(\Omega)$ .

It is then natural to define the following space in order to take into account the boundary conditions

$$\tilde{H}^1_0(Q) := \Phi(H^1_0(\Omega)).$$

We are now ready to state the variational formulation of (1) in both coordinate systems.

**Definition 2** (Weak form of the Poisson problem in Cartesian coordinates). Find  $\hat{u} \in H_0^1(\Omega)$  such that

(5) 
$$\hat{a}(\hat{u},\hat{v}) = \hat{b}(\hat{u}) \text{ for all } \hat{v} \in H_0^1(\Omega),$$

with

$$\hat{a}(\hat{u},\hat{v}) = \left(\hat{\nabla}\hat{u},\hat{\nabla}\hat{v}\right)_{L^{2}(\Omega)}, \quad \hat{b}(\hat{u}) = \left(\hat{f},\hat{v}\right)_{L^{2}(\Omega)}$$

**Definition 3** (Weak form of the Poisson problem in polar coordinates). Find  $u \in \tilde{H}^1_0(Q)$  such that

(6) 
$$a(u,v) = b(v) \quad for \ all \ v \in \tilde{H}^1_0(Q),$$

with

$$a(u,v) = \int_0^1 \int_0^\Theta \left( \partial_r u \partial_r v + \frac{1}{r^2} \partial_\theta u \partial_\theta v \right) r \, \mathrm{d}r \, \mathrm{d}\theta$$
$$b(v) = \int_0^1 \int_0^\Theta f v \, r \, \mathrm{d}r \, \mathrm{d}\theta.$$

It is well-known that (5) is well posed and so is (6) thanks to the properties of the map  $\Phi$  and of the isometry shown above.

# 3. The semi-discrete Poisson equation

The discretization of (1) with the scaled boundary finite element method is based on a spatial semi-discretization that is described in this section.

We introduce a partition of the parametrized boundary  $\theta \mapsto (\cos \theta, \sin \theta)$  given by

$$\mathcal{T}_{\Gamma} = \{\theta_1, \dots, \theta_N\}$$

and consider a finite dimensional approximation of  $H^1(0,\Theta)$  generated by a basis  $\{e_i(\theta)\}_{i=1}^N$  with the property that

$$e_i(\theta_j) = \delta_{ij}.$$

**Remark 4.** The choice of  $\{e_i(\theta)\}_{i=1}^N$  at this point is arbitrary. It could be based on finite elements, splines, global Lagrange polynomials, etc.

Due to our choice of boundary conditions, we could also have defined the basis  $\{e_i(\theta)\}_{i=1}^N$  in  $H_0^1(0,\Theta)$ , but we prefer to avoid this in order to allow our analysis to be extended more easily to more general boundary conditions or to a situation where  $\Theta = 2\pi$ .

The main idea behind the semi-discretization is to consider families of functions where the variables r and  $\theta$  are separated formally as follows:

$$u_s(r,\theta) = \sum_{i=1}^N u_i(r)e_i(\theta).$$

Ideally, we would like to have  $u_i(r) = u(r, \theta_i)$  and this choice will be used later in Subsection 3.1 for the error analysis; it will lead to the analogous of the interpolation operator for standard finite elements. In order to do so, we need to give sense to the radial trace  $u(r, \theta_i)$ . For the sake of readability, we now introduce an abstract setting and we postpone the actual definition of the involved functional spaces to Subsection 3.1. Ultimately, we want to define a semi-discrete space

$$U^s := \left\{ v_s \in \tilde{H}^1(Q) : v_s = \sum_{i=1}^N v_i(r)e_i(\theta) \text{ with } v_i \in \tilde{U} \text{ for } 1 \le i \le N \right\},$$

where  $\tilde{U}$  is a suitable functional space on the interval (0, 1). We will then consider its subspace  $U_0^s = U^s \cap \tilde{H}_0^1(Q)$ , so that the semi-discretization of problem (6) will read: find  $u_s \in U_0^s$  such that

(7) 
$$a(u_s, v_s) = b(v_s)$$
 for all  $v_s \in U_0^s$ .

We will prove (see Theorem 6) that  $U_0^s$  is a closed subspace of  $H_0^1(Q)$  so that problem (7) is uniquely solvable and the error between u and  $u_s$  is bounded as usual by the best approximation by Céa's lemma:

(8) 
$$\|u - u_s\|_{\tilde{H}^1(Q)} \le C \inf_{v \in U_0^s} \|u - v\|_{\tilde{H}^1(Q)}.$$

The solution of problem (7) is actually computed by solving a system of ordinary differential equations where the unknowns are the coefficients  $u_i(r)$  of  $u_s(r,\theta)$ . This procedure is detailed in Section 4.

In order to show the convergence of this procedure, we need to estimate the right-hand side of (8).

3.1. Error estimates for the interpolation operator. We plan to construct an interpolation operator  $\Pi$  with values in  $U^s$  that, if applied to smooth functions, would act as follows

$$(\Pi u)(r,\theta) = \sum_{i=1}^{N} u(r,\theta_i) e_i(\theta).$$

Since we will work with Sobolev functions, it is useful to define an adequate trace-like operator that we are going to call the "radial trace operator". To this end, the following bound is required.

**Lemma 5.** For all  $u \in C^{\infty}(\overline{Q})$  and  $0 \leq \vartheta \leq \Theta$  we have

(9) 
$$\int_{0}^{1} r u^{2}(r, \vartheta) \mathrm{d}r \leq C \left( \left\| u \right\|_{L^{2}_{r}(Q)}^{2} + \left\| \partial_{\theta} u \right\|_{L^{2}_{r}(Q)}^{2} \right)$$

where C > 0 only depends on  $\Theta$ .

*Proof.* Let  $u \in C^{\infty}(\overline{Q})$  and assume, without loss of generality, that  $0 \leq \zeta < \Theta$ . Then we have

$$\int_{\vartheta}^{\vartheta} \partial_{\zeta} \left( u^2(r,\zeta) \right) r \, \mathrm{d}\zeta = r u^2(r,\theta) - r u^2(r,\vartheta) \quad \text{for all } \theta \in (\vartheta,\Theta].$$

Reordering and integrating over r, we have

$$\int_0^1 r u^2(r,\vartheta) \,\mathrm{d}r = \int_0^1 r u^2(r,\theta) \,\mathrm{d}r - 2 \int_0^1 \int_\vartheta^\theta u(r,\zeta) \partial_\zeta u(r,\zeta) r \,\mathrm{d}\zeta \,\mathrm{d}r.$$

For the last term, we can apply Hölder's inequality, so that

$$-2\int_{0}^{1}\int_{\vartheta}^{\theta}u(r,\zeta)\partial_{\zeta}u(r,\zeta)r\,\mathrm{d}\zeta\,\mathrm{d}r \leq 2\int_{0}^{1}\int_{0}^{\Theta}|u(r,\zeta)\partial_{\zeta}u(r,\zeta)|\,r\,\mathrm{d}\zeta\,\mathrm{d}r$$
$$\leq 2\|u\|_{L^{2}_{r}(Q)}\|\partial_{\theta}u\|_{L^{2}_{r}(Q)}$$
$$\leq \|u\|_{L^{2}_{\pi}(Q)}^{2}+\|\partial_{\theta}u\|_{L^{2}_{\pi}(Q)}^{2}.$$

Finally, integrating over  $\theta$ , we have

$$\Theta \int_0^1 r u^2(r, \vartheta) \, \mathrm{d}r \le 2 \, \|u\|_{L^2_r(Q)}^2 + \|\partial_\theta u\|_{L^2_r(Q)}^2 \,,$$

so that (9) holds for smooth functions.

The following space on the interval (0,1) will be used for the definition of  $\tilde{U}$ 

$$H_r^1(0,1) = \left\{ u \in L_r^2(0,1) : \int_0^1 \left( u'(r) \right)^2 r \, \mathrm{d}r < \infty \right\}.$$

Given an angle  $0 \le \vartheta \le \Theta$ , inequality (9) shows that the natural norm for a space U where the radial trace operator can be defined, is

$$||u||_U := \left( ||u||^2_{\tilde{H}^1(Q)} + ||\partial_{r\theta} u||^2_{L^2_r(Q)} \right)^{\frac{1}{2}}.$$

It is apparent that not all functions in  $C^{\infty}(\overline{Q})$  have a bounded U-norm because in general  $C^{\infty}(\overline{Q})$  is not included in  $\tilde{H}^1(Q)$ . This is due to the fact that  $\|\partial_{\theta} u\|_{L^2_{1/r}(Q)}$  might not be bounded for some  $u \in C^{\infty}(\overline{Q})$ . Hence, we define

$$\tilde{\gamma}_{\vartheta}: C^{\infty}(\overline{Q}) \cap H^1(Q) \to H^1_r(0,1) \qquad u \mapsto u(\cdot,\vartheta)$$

and extend it to the closure of  $C^{\infty}(\overline{Q}) \cap \tilde{H}^1(Q)$  with respect to the *U*-norm. We denote by  $U \subset \tilde{H}^1(Q)$  this space and by  $\gamma_{\vartheta}$  the extension of the trace operator, so that we have a bounded radial trace operator

$$\gamma_{\vartheta}: U \to H^1_r(0,1)$$

that extends the restriction operator  $\tilde{\gamma}_\vartheta$  defined on smooth enough functions.

It is then natural to choose  $\tilde{U} = H^1_r(0,1)$ , so that the definition of  $U^s$  reads as follows

$$U^{s} := \left\{ v_{s} \in \tilde{H}^{1}(Q) : v_{s} = \sum_{i=1}^{N} v_{i}(r)e_{i}(\theta) \text{ with } v_{i} \in H^{1}_{r}(0,1) \text{ for } 1 \le i \le N \right\}.$$

**Theorem 6.** The space of semi-discrete functions  $U^s$  is a closed subspace of  $\tilde{H}^1(Q)$ .

*Proof.* Let  $(u_n)$  be a Cauchy sequence in  $U^s$ . By completeness, there is a function  $\tilde{u}$  such that  $u_n \to \tilde{u}$  in  $\tilde{H}^1(Q)$ . By Lemma 5 we have

$$\int_0^1 r |u_m(r,\theta_i) - u_n(r,\theta_i)|^2 \, \mathrm{d}r \le C ||u_m - u_n||^2_{\tilde{H}^1(Q)} \to 0$$

as  $n, m \to \infty$ , so  $(u_n(\cdot, \theta_i))$  is a Cauchy sequence in  $L^2_r(0, 1)$  and by completeness there is a limit  $u_n(\cdot, \theta_i) \to u_i$  for each *i*. Define  $u = \sum_{i=1}^N u_i e_i$  in  $U^s$  and note that

$$\begin{aligned} \|u - \tilde{u}\|_{L^{2}_{r}(Q)} &\leq \|u - u_{n}\|_{L^{2}_{r}(Q)} + \|\tilde{u} - u_{n}\|_{L^{2}_{r}(Q)} \\ &\leq C \left(\sum_{i=1}^{N} \|u_{i}(\cdot) - u_{n}(\cdot, \theta_{i})\|_{L^{2}_{r}(0, 1)}^{2}\right)^{\frac{1}{2}} + \|\tilde{u} - u_{n}\|_{L^{2}_{r}(Q)} \to 0 \end{aligned}$$

as  $n \to \infty$ , so  $u = \tilde{u}$  and therefore  $u_n \to u$  in  $\tilde{H}^1(Q)$ .

Given  $u \in U$  we can then define the interpolant as

$$(\Pi u)(r,\theta) = \sum_{i=1}^{N} u_i(r)e_i(\theta),$$

where  $u_i(r)$  is defined as  $\gamma_{\theta_i}(u)$  in  $H_r^1(0, 1)$ . In order to see that the interpolant is well defined, we need to show that  $\Pi u$  belongs to  $\tilde{H}^1(Q)$ . To limit the technicalities, from now on in this section we are assuming that  $\{e_i\}$  is the basis of continuous piecewise linear finite elements on  $(0, \Theta)$ . The general case can be handled with similar arguments.

**Lemma 7.** For  $u \in U$ , we have  $\Pi u \in \tilde{H}^1(Q)$ .

*Proof.* We have that  $\Pi u \in \tilde{H}^1(Q)$  if and only if

(10) 
$$\int_0^1 \int_0^{2\pi} \left( r(\Pi u)^2 + r(\partial_r \Pi u)^2 + \frac{1}{r} (\partial_\theta \Pi u)^2 \right) \, \mathrm{d}\theta \, \mathrm{d}r < \infty.$$

We apply Lemma 5 and obtain (11)

$$\begin{split} \|\Pi u\|_{L^{2}_{r}(Q)}^{2} + \|\partial_{r}\Pi u\|_{L^{2}_{r}(Q)}^{2} &\leq N \sum_{i=1}^{N} \left( \int_{0}^{1} r u_{i}^{2}(r) \,\mathrm{d}r \int_{0}^{2\pi} e_{i}^{2}(\theta) \,\mathrm{d}\theta \right. \\ &+ \int_{0}^{1} r \left( \partial_{r} u_{i}(r) \right)^{2} \,\mathrm{d}r \int_{0}^{2\pi} e_{i}^{2}(\theta) \,\mathrm{d}\theta \Big) \\ &\leq CN \left( \|u\|_{\tilde{H}^{1}(Q)}^{2} + \|\partial_{r\theta} u\|_{L^{2}_{r}(Q)}^{2} \right) \sum_{i=1}^{N} \int_{0}^{2\pi} e_{i}^{2}(\theta) \,\mathrm{d}\theta \\ &\leq \frac{4\pi^{2}Ch}{3h_{\min}^{2}} \left( \|u\|_{\tilde{H}^{1}(Q)}^{2} + \|\partial_{r\theta} u\|_{L^{2}_{r}(Q)}^{2} \right), \end{split}$$

where  $h = \max_i(\theta_{i+1} - \theta_i)$  and  $h_{\min} = \min_i(\theta_{i+1} - \theta_i)$ .

For the third term in (10), we fix  $r \in (0, 1)$  and observe that

 $u(r,\theta_i)=(\Pi u)(r,\theta_i),\quad u(r,\theta_{i+1})=(\Pi u)(r,\theta_{i+1})$ 

for i = 1, 2, ..., N - 1. Taking into account that  $\partial_{\theta} \Pi u$  is well defined in  $(\theta_i, \theta_{i+1})$ , we apply the mean value theorem and

$$u(r,\theta_{i+1}) - u(r,\theta_i) = (\theta_{i+1} - \theta_i) \left( \partial_{\theta} \Pi u(r,\tilde{\theta}) \right)$$

holds for some  $\tilde{\theta} \in (\theta_i, \theta_{i+1})$ . Since  $\Pi u$  is linear in this interval, the following equality holds

$$\left|\partial_{\theta}\Pi u(r,\theta)\right|^{2}\left(\theta_{i+1}-\theta_{i}\right)^{2} = \left|\int_{\theta_{i}}^{\theta_{i+1}}\partial_{\theta}u(r,\zeta)\,\mathrm{d}\zeta\right|^{2}$$

for all  $\theta \in (\theta_i, \theta_{i+1})$  and  $r \in (0, 1)$ . After multiplying by 1/r, integrating, and applying Hölder's inequality, we have

$$\int_0^1 \int_{\theta_i}^{\theta_{i+1}} \frac{1}{r} \left| \partial_\theta \Pi u(r,\theta) \right|^2 \, \mathrm{d}\theta \, \mathrm{d}r \le \int_0^1 \int_{\theta_i}^{\theta_{i+1}} \frac{1}{r} \left| \partial_\theta u(r,\theta) \right|^2 \, \mathrm{d}\theta \, \mathrm{d}r.$$

By integrating over each interval and summing up the terms, we have

(12) 
$$\|\partial_{\theta}\Pi u\|_{L^{2}_{1/r}(Q)} \leq \|\partial_{\theta} u\|_{L^{2}_{1/r}(Q)}.$$

Finally, putting (11) and (12) together, we get

$$\|\Pi u\|_{\tilde{H}^{1}(Q)}^{2} \leq \max\left\{\frac{4\pi^{2}Ch}{3h_{\min}^{2}}, 1\right\} \left(2\|u\|_{\tilde{H}^{1}(Q)}^{2} + \|\partial_{r\theta}u\|_{L_{r}^{2}(Q)}^{2}\right) < \infty.$$

In the next theorem we prove the approximation properties of  $U^s$ . As usual, we need to assume suitable regularity that will be characterized by the following space

$$U' = \left\{ u \in U : \left\| \partial_{\theta \theta} u \right\|_{L^2_{1/r}(Q)} < \infty \right\}.$$

**Remark 8.** The space U' requires extra regularity only in the angular variable  $\theta$ . We will see in Section 5 that singular solutions (with respect to the Cartesian coordinates) can be in U' and be approximated optimally by SBFEM.

**Theorem 9.** Let u be in U'. Then there exists C > 0 independent of  $\mathcal{T}_{\Gamma}$  such that

$$\|u - \Pi u\|_{L^2_r(Q)} \le h^2 \|\partial_{\theta\theta} u\|_{L^2_{1/r}(Q)}$$

and

$$\|u - \Pi u\|_{\tilde{H}^{1}(Q)} \le Ch\left(\|\partial_{r\theta} u\|_{L^{2}_{r}(Q)}^{2} + \|\partial_{\theta\theta} u\|_{L^{2}_{1/r}(Q)}^{2}\right).$$

*Proof.* Let  $u \in C^{\infty}(\overline{Q})$  and define  $\varepsilon(r, \theta) = (u - \Pi u)(r, \theta)$ . Due to the properties of the interpolation operator, we have  $\varepsilon(r, \theta_i) = 0$  for i = 1, ..., N. As a result, there is a  $\tilde{\theta}_i \in (\theta_i, \theta_{i+1})$  such that  $\partial_{\theta} \varepsilon(r, \tilde{\theta}_i) = 0$ . It follows that

$$\partial_{\theta}\varepsilon(r,\vartheta) = \int_{\tilde{\theta}_{i}}^{\vartheta} \partial_{\theta\theta}\varepsilon(r,\zeta) \,\mathrm{d}\zeta = \int_{\tilde{\theta}_{i}}^{\vartheta} \partial_{\theta\theta}u(r,\zeta) \,\mathrm{d}\zeta \quad \text{for } \tilde{\theta}_{i} < \vartheta \leq \theta_{i+1},$$

since  $\Pi u$  is linear in  $(\theta_i, \theta_{i+1})$  in the  $\theta$  direction. Applying Hölder's inequality we have

$$\left|\partial_{\theta}\varepsilon(r,\vartheta)\right|^{2} \leq \left(\theta_{i+1} - \theta_{i}\right) \int_{\theta_{i}}^{\theta_{i+1}} \left|\partial_{\theta\theta}u(r,\theta)\right|^{2} \,\mathrm{d}\theta.$$

Integrating over the domain and summing up the different terms corresponding to each interval  $(\theta_i, \theta_{i+1})$  we have

(13) 
$$\|\partial_{\theta}\varepsilon\|^2_{L_{1/r}(Q)} \le h^2 \|\partial_{\theta\theta}u\|^2_{L_{1/r}(Q)}$$

Since both  $|| \cdot ||_{L_{1/r}(Q)}$  and  $\Pi$  are continuous, inequality (13) can be shown to hold for all  $u \in U'$  by a density argument. Likewise, for  $\varepsilon(r, \vartheta)$  we have

$$\left|\varepsilon(r,\vartheta)\right|^{2} = \left|\int_{\theta_{i}}^{\vartheta} \partial_{\theta}\varepsilon(r,\zeta) \,\mathrm{d}\zeta\right|^{2} \le h \int_{\theta_{i}}^{\theta_{i+1}} \left|\partial_{\theta}\varepsilon(r,\zeta)\right|^{2} \,\mathrm{d}\zeta.$$

After integrating and using (13), we have

(14) 
$$\|\varepsilon\|_{L^{2}_{r}(Q)}^{2} \leq h^{4} \|\partial_{\theta\theta}u\|_{L^{2}_{1/r}(Q)}^{2}$$

Finally, an estimate must be found for  $\|\partial_r \varepsilon\|_{L^2_r(Q)}$ . Once again, we consider a smooth function u and take into account that  $\partial_r \varepsilon(r, \theta_i) = 0$  for all  $r \in (0, 1)$  and  $i = 1, \ldots, N$ . Hence

$$\partial_r \varepsilon(r, \vartheta) = \int_{\theta_i}^{\vartheta} \partial_{r\theta} \varepsilon(r, \zeta) \,\mathrm{d}\zeta$$

and it follows that

(15) 
$$\left\|\partial_r \varepsilon\right\|_{L^2_r(Q)}^2 \le h^2 \left\|\partial_{r\theta} \varepsilon\right\|_{L^2_{1/r}(Q)}^2.$$

In the same way as (12) is obtained, we apply the mean value theorem to the function  $\partial_r(\Pi u)(r, \vartheta)$  for  $\vartheta \in (\theta_i, \theta_{i+1})$  and obtain

$$\left(\partial_{r\theta}(\Pi u)(r,\vartheta)\right)\left(\theta_{i+1}-\theta_i\right) = \int_{\theta_i}^{\theta_{i+1}} \partial_{r\theta} u(r,\zeta) \,\mathrm{d}\zeta$$

and therefore we can establish that

(16) 
$$\|\partial_{r\theta}\Pi u\|_{L^2_r(Q)} \le \|\partial_{r\theta} u\|_{L^2_r(Q)}.$$

Given (16), we have the following error estimate for all smooth functions u:

(17) 
$$\begin{aligned} \left\|\partial_{r}\varepsilon\right\|_{L^{2}_{r}(Q)}^{2} &\leq 2h^{2}\left(\left\|\partial_{r\theta}u\right\|_{L^{2}_{r}(Q)}^{2}+\left\|\partial_{r\theta}\Pi u\right\|_{L^{2}_{r}(Q)}^{2}\right) \\ &\leq 4h^{2}\left\|\partial_{r\theta}u\right\|_{L^{2}_{r}(Q)}^{2} \end{aligned}$$

which, by a density argument, holds for all  $u \in U'$ . Therefore, putting (13), (14), and (17) together, we obtain the required bound. More precisely, we have

$$\|u - \Pi u\|_{\tilde{H}^{1}(Q)}^{2} \leq h^{2} \left(h^{2} + 4\right) \left(\|\partial_{r\theta} u\|_{L_{r}^{2}(Q)}^{2} + \|\partial_{\theta\theta} u\|_{L_{1/r}(Q)}^{2}\right).$$

# 4. Constructing semi-discrete solutions with SBFEM

In order to solve our problem, the scaled boundary finite element method rewrites the formulation (7) as a system of ordinary differential equations; this is carried out in Section 4.1. This is possible thanks to the representation of semi-discrete functions in  $U_0^s$  as the product of *r*-dependent functions  $u_i(r)$  with  $\theta$ -dependent test functions. The resulting system of differential equations is supplemented with additional conditions arising from regularity requirements and the boundary conditions. This system can be solved by an analytical method under certain conditions, a process which is described in Section 4.2.

4.1. Rewriting the Poisson equation as an ODE. Let us consider  $u_s = \sum_{i=1}^{N} u_i e_i$  and  $v_s = \sum_{i=1}^{N} v_i e_i$ . An immediate consequence of the definition of the space of semi-discrete functions  $U_0^s$  is that the bilinear form  $a: U_0^s \times U_0^s \to \mathbb{R}$  and the linear form  $b: U_0^s \to \mathbb{R}$  may be rewritten as follows:

$$\begin{aligned} a(u_s, v_s) &= \sum_{i,j=1}^N \int_0^1 \int_0^\Theta \left( u'_i(r) v'_j(r) e_i(\theta) e_j(\theta) + \frac{1}{r^2} u_i(r) v_j(r) e'_i(\theta) e'_j(\theta) \right) r \, \mathrm{d}r \, \mathrm{d}\theta \\ &= \sum_{i,j=1}^N \left( \int_0^1 u'_i(r) v'_j(r) r \, \mathrm{d}r \int_0^\Theta e_i(\theta) e_j(\theta) \, \mathrm{d}\theta \right. \\ &+ \int_0^1 u_i(r) v_j(r) \frac{\mathrm{d}r}{r} \int_0^\Theta e'_i(\theta) e'_j(\theta) \, \mathrm{d}\theta \right) \\ &= \sum_{i,j=1}^n \left( \mathcal{A}_{ij} \int_0^1 u'_i(r) v'_j(r) r \, \mathrm{d}r + \mathcal{B}_{ij} \int_0^1 u_i(r) v_j(r) \frac{\mathrm{d}r}{r} \right) \end{aligned}$$

and

$$b(v_s) = \sum_{j=1}^N \int_0^1 \int_0^\Theta f(r,\theta) v_j(r) e_j(\theta) r \, \mathrm{d}r \, \mathrm{d}\theta$$
$$= \sum_{j=1}^N \int_0^1 F_j(r) v_j(r) r \, \mathrm{d}r,$$

where

$$A_{ij} = \int_0^{\Theta} e_i(\theta) e_j(\theta) d\theta$$
$$B_{ij} = \int_0^{\Theta} e'_i(\theta) e'_j(\theta) d\theta$$
$$F_j(r) = \int_0^{\Theta} f(r, \theta) e_j(\theta) d\theta.$$

We now proceed formally with the derivation of the differential equation. To this aim, we will use the following integration by parts formula

$$\int_0^1 u_i'(r)v_j'(r)r\,\mathrm{d}r = -\int_0^1 u_i''(r)v_j(r)r\,\mathrm{d}r - \int_0^1 u_i'(r)v_j(r)\mathrm{d}r + u_i'(1)v_j(1)$$

which is clearly valid for smooth enough  $u_i$  and  $v_i$ .

In order to simplify our notation we introduce a name for the space of the coefficients in  $U^s$  as follows

$$\boldsymbol{U}_{0}^{s} = \left\{ \boldsymbol{u} = (u_{i})_{i=1}^{N} : u_{i} \in H_{r}^{1}(0,1) \text{ for } 1 \leq i \leq N, \sum_{i=1}^{N} u_{i}e_{i} \in U_{0}^{s} \right\}.$$

It follows that if  $u_s \in U_0^s$  is smooth enough, it solves  $a(u_s, v_s) = b(v_s)$  for all  $v_s \in U_0^s$  if and only if

(18) 
$$\sum_{j=1}^{N} \int_{0}^{1} v_{j}(r) \sum_{i=1}^{N} \left[ -r \mathbf{A}_{ij} u_{i}''(r) - \mathbf{A}_{ij} u_{i}'(r) + \frac{1}{r} \mathbf{B}_{ij} u_{i}(r) - F_{j}(r) r \right] \mathrm{d}r = 0$$

holds for all  $\boldsymbol{v} = (v_i)_{i=1}^N \in \boldsymbol{U}_0^s$ . Moreover, as a result of the fundamental lemma of calculus of variations, (18) holds if and only if

(19) 
$$\sum_{j=1}^{N} \left( r \mathbf{A}_{ij} u_j''(r) + \mathbf{A}_{ij} u_j'(r) - \frac{1}{r} \mathbf{B}_{ij} u_j(r) \right) = r F_i(r) \text{ for a.e. } r \in (0,1)$$

is satisfied for all i = 2, ..., N - 1, since  $v_1 = v_N = 0$  by  $\boldsymbol{v} \in \boldsymbol{U}_0^s$ . Because we seek a solution  $\boldsymbol{u} \in \boldsymbol{U}_0^s$ , we must also enforce the boundary conditions  $u_1 = u_N = 0$  and  $u_i(1) = 0$  for i = 2, ..., N - 1. Furthermore, we impose the compatibility condition  $u_i(0) = u_j(0)$  for all i, j = 1, ..., N to avoid a singularity at the center of the disk. In his, we guarantee that the function  $u = \sum_{i=1}^N u_i e_i$  belongs to  $\tilde{H}_0^1(Q)$ . Denote by **A** and **B** the matrices in  $\mathbb{R}^{N \times N}$  with components  $A_{ij}$  and  $B_{ij}$  and by

Denote by **A** and **B** the matrices in  $\mathbb{R}^{N \times N}$  with components  $A_{ij}$  and  $B_{ij}$  and by F(r) the vector function with components  $F_i(r)$ . We modify the columns and rows in **A** and **B** and we set  $F_i(r) = 0$  for i = 1 and N in order to enforce  $u_1 = u_N = 0$ . Then, a semi-discrete solution of the Poisson equation  $u = (u_i)_{i=1}^N$  will satisfy the following conditions:

(20) 
$$\begin{cases} r^{2} \mathbf{A} \boldsymbol{u}'' + r \mathbf{A} \boldsymbol{u}' - \mathbf{B} \boldsymbol{u} = r^{2} \boldsymbol{F} & \text{for } r \in (0, 1), \\ \boldsymbol{u}(1) = 0, \\ u_{i}(0) = u_{j}(0) & \text{for all } i, j = 1, \dots, N. \end{cases}$$

SBFEM provides a methodology for the construction of solutions for (20), see the next section. Whenever such solutions exist and satisfy the integration by parts formula for all  $\boldsymbol{v} \in \boldsymbol{U}_0^s$ , then they correspond with the unique solution in  $U_0^s$ .

4.2. Solving the ODE analytically. Following [9], a solution to (20) can be constructed by finding a family of functions that satisfy the homogeneous ODE together with a particular solution. In order to construct the homogeneous solution, we shall define the matrix  $\mathbf{E} \in \mathbb{R}^{2N \times 2N}$  by

$$\mathbf{E} = \begin{pmatrix} 0 & \mathbf{A}^{-1} \\ \mathbf{B} & 0 \end{pmatrix}.$$

Note that **E** arises when the ODE in (20) is rewritten as a first order differential equation by introducing the additional variable  $\boldsymbol{q}(r) = r \mathbf{A} \boldsymbol{u}'(r)$ . If  $(\lambda, (\boldsymbol{\phi}, \boldsymbol{\psi})^T)$  is an eigenpair of **E**, with  $\boldsymbol{\phi}, \boldsymbol{\psi} \in \mathbb{C}^N$ , we can see that

(21) 
$$\mathbf{B}\boldsymbol{\phi} = \lambda^2 \mathbf{A}\boldsymbol{\phi}.$$

Furthermore, for  $\boldsymbol{u}(r) = r^{\lambda} \boldsymbol{\phi}$  we have that

(22) 
$$r^2 \mathbf{A} \boldsymbol{u}'' + r \mathbf{A} \boldsymbol{u}' - \mathbf{B} \boldsymbol{u} = 0$$

for all r > 0. That is,  $\mathbf{u}(r) = r^{\lambda} \boldsymbol{\phi}$  is a homogeneous solution of the ODE in (20). This idea can be extended in such a way that we get homogeneous solutions as a linear combination of N linearly independent functions that satisfy (22). Indeed, first note that  $\mathbf{E}$  is a Hamiltonian matrix and therefore, for every eigenvalue  $\lambda \in \mathbb{C}$ , we will also have that  $-\lambda$ ,  $\overline{\lambda}$  and  $-\overline{\lambda}$  are eigenvalues of  $\mathbf{E}$ . Consider the following subset of N eigenpairs of  $\mathbf{E}$ :

$$\left\{ (\lambda_i, \begin{pmatrix} \boldsymbol{\phi}_i \\ \boldsymbol{\psi}_i \end{pmatrix}) : \operatorname{Re}(\lambda_i) \ge 0, \quad \boldsymbol{\phi}_i, \boldsymbol{\psi}_i \in \mathbb{C}^N, \quad i = 1, \dots, N \right\}.$$

From equation (21), we see that the pair  $(\lambda_i^2, \phi_i)$  is an eigenpair of the one dimensional Laplace problem discretised with piecewise polynomials. As a result,  $\lambda_i \in \mathbb{R}$  and  $\phi_i \in \mathbb{R}^N$ . Moreover, the vectors  $(\phi_i)$  form a basis of  $\mathbb{R}^N$ . Then, for any *N*-tuple of real numbers  $c_1, \ldots, c_N$ , we have that

$$\boldsymbol{u}_h(r) = \sum_{i=1}^N c_i r^{\lambda_i} \boldsymbol{\phi}_i$$

is a homogeneous solution to the ODE in (20). Given a particular solution  $u_p$  for which

$$r^{2}\mathbf{A}\boldsymbol{u}_{p}^{\prime\prime}+r\mathbf{A}\boldsymbol{u}_{p}^{\prime}-\mathbf{B}\boldsymbol{u}_{p}=r^{2}\boldsymbol{F}\quad\text{on }(0,1),$$

the constants  $c_1, \ldots c_N$  are then set by enforcing the boundary conditions  $\boldsymbol{u}_h(1) + \boldsymbol{u}_p(1) = 0$ . The resulting function  $\boldsymbol{u} = \boldsymbol{u}_h + \boldsymbol{u}_p$  is a solution of (20).

**Remark 10.** In the construction of  $\boldsymbol{u}$  we only consider non-negative eigenvalues because we require  $u_i(r) \in H_r^1(0,1)$  for all i = 1, ..., N. Moreover the condition  $u_i(0) = u_j(0)$  for all i, j = 1, ..., N holds because if  $\lambda_i = 0$  then  $\boldsymbol{\phi}$  is in the kernel of **B**, which consists of vectors whose entries are all equal.

The analytical construction of a particular solution relies on the form of  $\mathbf{F}(r)$ . For example, if  $\mathbf{F}(r) = r^{\alpha} \mathbf{f}$ , where  $\mathbf{f} \in \mathbb{R}^{N}$  and  $\alpha \in \mathbb{R}$ , then it is simple to see that  $\mathbf{u}_{p}(r) = r^{\alpha+2} \phi_{p}$ , with  $\phi_{p} \in \mathbb{R}^{N}$  solving the linear system

$$\left( (\alpha + 2)^2 \mathbf{A} - \mathbf{B} \right) \boldsymbol{\phi}_p = \boldsymbol{f},$$

is a particular solution of (19).

#### 5. Numerical examples

One of the main features of the method presented in the previous sections is that it can achieve high order of convergence also in the presence of singular solutions. We are going to show this behavior with two simple examples. The approximate solutions to (19) are obtained by means of the analytical method described in Section 4.2. In these tests, the basis functions  $e_i$  in the angular direction are piecewise polynomials of order 1 or higher.

5.1. A first numerical test. We take  $\Theta = 3\pi/2$  and consider the function

$$u_e(r,\theta) = r^{\frac{2}{3}} \sin\left(\frac{2}{3}\theta\right).$$

This function satisfies  $\Delta \hat{u}_e = 0$  on  $\Omega$ , where  $\hat{u}_e = \Phi^{-1}(u_e)$ , so it is a solution to the homogeneous problem: find  $\hat{u} \in H^1(\Omega)$  such that

$$-\Delta \hat{u} = 0 \quad \text{in } \Omega$$
$$\hat{u} = \hat{u}_e \quad \text{on } \partial \Omega.$$

Our theory can be easily extended to accommodate non-homogeneous boundary conditions. Moreover, it is well-known that  $\hat{u}_e$  belongs to  $H^1(\Omega)$  but not to  $H^2(\Omega)$ . Indeed, this follows from the fact that the following inequality holds for any 0 < R < 1 and  $0 < \varepsilon < R$ 

$$\begin{aligned} \|\hat{u}_e\|_{2,\Omega}^2 &\ge \int_{\varepsilon}^R \int_0^{\frac{3}{2}\pi} \left(\frac{\partial u_e}{\partial r}\right)^2 r \,\mathrm{d}\theta \,\mathrm{d}r = \left(\frac{4}{81} \int_0^{\frac{3}{2}\pi} \sin^2\left(\frac{2}{3}\theta\right) \,\mathrm{d}\theta\right) \int_{\varepsilon}^R r^{-\frac{5}{3}} \,\mathrm{d}r \\ &= C\left(\varepsilon^{-\frac{2}{3}} - R^{-\frac{2}{3}}\right) \end{aligned}$$

which tends to infinity as  $\varepsilon$  goes to 0. On the other hand, it can be easily seen that  $u_e$  belongs to U' and this makes it possible to use the result of Theorem 9 which implies, in particular, that first order elements achieve second order of convergence in  $L^2$  even in presence of a corner singularity.

We report in Figure 2 the results of our numerical test that confirm our theoretical findings. We include also higher order approximations (up to order six) and a convergence plot in  $H^1$ .

5.2. A second numerical test. For the second numerical test, we consider a slightly more complicated example. Once again, we set  $\Theta = 3\pi/2$  and consider the function

$$v_e(r,\theta) = r^{\frac{2}{3}} \left( \left( 1 - \frac{4\theta}{3\pi} \right) \cos\left(2\theta/3\right) - \frac{4}{3\pi} \log\left(r\right) \sin\left(2\theta/3\right) \right).$$

For this function we once again have that  $\Delta \hat{v}_e = 0$ , where  $\hat{v}_e = \Phi^{-1}(v_e)$ , and  $v_e = r^{2/3}$  at  $\theta = 0$  and  $\theta = \Theta$ . This function therefore is a solution to the problem: find  $\hat{u} \in H^1(\Omega)$  such that

$$\begin{aligned} -\Delta \hat{u} &= 0 \quad \text{in } \Omega \\ \hat{u} &= \hat{v}_{e} \quad \text{on } \partial \Omega. \end{aligned}$$

The results to our numerical tests are presented in Figure 3 for polynomial orders 1 and 2. The resulting convergence rates confirm the theoretical predictions from Section 3.



FIGURE 2. Convergence plots of  $L^2$  and  $H^1$ -errors for the first numerical test with basis functions  $e_i$  of polynomial order 1, 2, 4, and 6.



FIGURE 3. Convergence plots of  $L^2$  and  $H^1$ -errors for the second numerical test with basis functions  $e_i$  of polynomial order 1 and 2.

# References

- Hauke Gravenkamp, Albert A. Saputra, Chongmin Song, and Carolin Birk, Efficient wave propagation simulation on quadtree meshes using SBFEM with reduced modal basis, International Journal for Numerical Methods in Engineering 110 (2016), no. 12, 1119–1141.
- 2. Alois Kufner, Weighted sobolev spaces, Teubner-Texte zur Mathematik, B.G. Teubner, 1985.
- Lei Liu, Junqi Zhang, Chongmin Song, Carolin Birk, Albert A. Saputra, and Wei Gao, Automatic three-dimensional acoustic-structure interaction analysis using the scaled boundary finite element method, Journal of Computational Physics 395 (2019), 432 – 460.
- Ean Tat Ooi, Chongmin Song, and Francis Tin-Loi, A scaled boundary polygon formulation for elasto-plastic analyses, Computer Methods in Applied Mechanics and Engineering 268 (2014), 905 – 937.

- 5. Ean Tat Ooi, Chongmin Song, Francis Tin-Loi, and Zhenjun Yang, *Polygon scaled boundary finite elements for crack propagation modelling*, International Journal for Numerical Methods in Engineering **91** (2012), no. 3, 319–342.
- Chongmin Song and John P. Wolf, The scaled boundary finite-element method—alias consistent infinitesimal finite-element cell method—for elastodynamics, Computer Methods in Applied Mechanics and Engineering 147 (1997), no. 3, 329 – 355.
- Chongmin Song and John P. Wolf, The scaled boundary finite element method—alias consistent infinitesimal finite element cell method—for diffusion, International Journal for Numerical Methods in Engineering 45 (1999), no. 10, 1403–1431.
- Chongmin Song and John P. Wolf, Semi-analytical representation of stress singularities as occurring in cracks in anisotropic multi-materials with the scaled boundary finite-element method, Computers & Structures 80 (2002), no. 2, 183 – 197.
- 9. John Wolf, Scaled boundary finite element method, John Wiley & Sons, 2003.
- 10. William P. Ziemer, Weakly differentiable functions, Springer-Verlag, Berlin, Heidelberg, 1989.

HUMBOLDT-UNIVERSITÄT ZU BERLIN, GERMANY, KING ABDULLAH UNIVERSITY OF SCIENCE AND TECHNOLOGY, SAUDI ARABIA AND UNIVERSITY OF TWENTE, ENSCHEDE, THE NETHERLANDS *Email address*: fb@math.hu-berlin.de

KING ABDULLAH UNIVERSITY OF SCIENCE AND TECHNOLOGY (KAUST), SAUDI ARABIA AND UNIVERSITY OF PAVIA, ITALY

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, UK *Email address:* gonzalezdedi@maths.ox.ac.uk