

ITERATED CONSTRUCTIONS OF COMPLETELY NORMAL POLYNOMIALS

ANIBAL ARAVENA[†]

ABSTRACT. The $R_{\sigma,t}$ -transform introduced by Bassa and Menares can be used to construct families of irreducible polynomials in $\mathbb{F}_q[x]$. This iterative construction is a generalization of Cohen's R -transform. For this transform, Chapman proved that under some conditions, the polynomials in the resulting family are completely normal. In this paper we establish conditions ensuring that the polynomials obtained by using the $R_{\sigma,t}$ -transform are completely normal polynomials and we give a simple proof of Chapman's result.

Keywords: Finite Fields; normal polynomials; R -transform.

1. INTRODUCTION

Let q be a prime power number and n a positive integer. An element $\alpha \in \mathbb{F}_{q^n}$ is said normal in the extension $\mathbb{F}_{q^n}/\mathbb{F}_q$ if the elements $\{\alpha, \alpha^q, \dots, \alpha^{q^{n-1}}\}$ are linearly independent over \mathbb{F}_q . Furthermore, if for each divisor d of n , α is normal in the extension $\mathbb{F}_{q^n}/\mathbb{F}_{q^d}$, then α is called a completely normal element in the extension $\mathbb{F}_{q^n}/\mathbb{F}_q$. Let f be an irreducible polynomial in $\mathbb{F}_q[x]$, we say that f is normal (completely normal) over $\mathbb{F}_q[x]$ if any of its roots is normal (completely normal) in the extension $\mathbb{F}_{q^{\deg f}}/\mathbb{F}_q$.

Starting with a polynomial $g_0(x) \in \mathbb{F}_q[x]$, we focus on an iterative construction introduced in [1] to get a family $\{g_k\}_{k \geq 0}$ of polynomials in $\mathbb{F}_q[x]$ of growing degree. Our aim is to find suitable hypothesis ensuring that $\{g_k\}_{k \geq 0}$ are all completely normal polynomials over $\mathbb{F}_q[x]$. To define it and state the basic results we need some previous notations. Let

$$\mathrm{GL}_2(\mathbb{F}_q) = \left\{ \sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{F}_q) : ad - bc \neq 0 \right\}.$$

For $\alpha \in \overline{\mathbb{F}}_q \cup \{\infty\}$, $\sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{F}_q)$, we define the operation $\sigma \cdot \alpha$ by

$$\sigma \cdot \alpha = \begin{cases} \frac{a\alpha+b}{c\alpha+d} & \text{if } c\alpha + d \neq 0 \\ \infty & \text{if } c\alpha + d = 0 \end{cases}, \quad \sigma \cdot \infty = \begin{cases} \frac{a}{c} & \text{if } c \neq 0 \\ \infty & \text{if } c = 0 \end{cases}.$$

Also, for a degree n polynomial f in $\mathbb{F}_q[x]$, we define the polynomial $P_\sigma(f)$ by

$$P_\sigma(f)(x) = (cx + d)^n f\left(\frac{ax + b}{cx + d}\right).$$

Also, for any positive integer t , let $S_t : \mathbb{F}_q[x] \rightarrow \mathbb{F}_q[x]$ be the map given by $S_t(f)(x) = f(x^t)$. Finally, for a degree n polynomial g , we define the $R_{\sigma,t}$ -transform by

$$g^{R_{\sigma,t}}(x) = P_{\sigma^{-1}} \circ S_t \circ P_\sigma(g)(x).$$

If $g(a/c) \neq 0$, we define the element $\eta(g; \sigma) \in \mathbb{F}_q$ by

$$\eta(g; \sigma) = (\sigma^{-1} \cdot \infty)^n \frac{g(\sigma \cdot 0)}{g(\sigma \cdot \infty)} = \left(-\frac{d}{c}\right)^n \cdot g\left(\frac{b}{d}\right) \cdot g\left(\frac{a}{c}\right)^{-1}$$

with the convention $\eta(g; \sigma) = (-d/a)^n \cdot g(b/d)$ if $c = 0$ and $\eta(g; \sigma) = (-b/c)^n \cdot g(a/c)^{-1}$ if $d = 0$.

[†]Pontificia Universidad Católica de Chile.

The author was partially supported by FONDECYT 1171329.

The following Theorem is about the irreducibility of the iterated sequence $\{g_k\}_{k \geq 0}$ constructed using the $R_{\sigma,t}$ -transform.

Theorem 1.1 ([1], Theorem 1.2). *Consider a finite field \mathbb{F}_q . Let $t \geq 2$ be an integer such that every prime factor of t divides $q-1$. Let $\sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{F}_q)$. Let $g(x) \neq x - a/c$ be a monic irreducible polynomial in $\mathbb{F}_q[x]$ of degree n . If $q \equiv 3 \pmod{4}$ and t is even assume moreover that n is even.*

Assume that, for all prime numbers $\ell|t$, the element $\eta(g; \sigma)$ is not an ℓ -th power in \mathbb{F}_q . Define $g_0 = g$ and let $g_k = g_{k-1}^{R_{\sigma,t}}$ for $k \geq 1$. Then $\{g_k\}_{k \geq 0}$ forms an infinite sequence of irreducible polynomials, with $\deg g_k = t^k \cdot n$.

An important particular case is the R -transform, previously introduced by S.Cohen [2]:

$$g^R(x) = (2x)^{\deg g} \cdot g\left(\frac{1}{2}\left(x + \frac{1}{x}\right)\right).$$

This corresponds to use $t = 2$, $\sigma^* = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ and we have that $\eta(\sigma^*, g) = g(1)g(-1)^{-1}$. Moreover, we have the following result.

Theorem 1.2 ([3], Theorem 1). *Let $q \equiv 1 \pmod{4}$ be a prime power, and let $g_1 = x^2 + ax + 1$ be an irreducible quadratic polynomial. Define g_k recursively by $g_{k+1} = g_k^R$. If α_k is a zero of g_k , then α_k is a completely normal element in the extension $\mathbb{F}_{q^n}/\mathbb{F}_q$.*

Our main Theorem is a generalization of this result for the $R_{\sigma,t}$ -transform

Theorem 1.3. *Let $g(x) = x - A \in \mathbb{F}_q[x]$ with $A \neq -a/c$, $t \geq 2$, $\sigma \in \text{GL}_2(\mathbb{F}_q)$ be as in Theorem 1.1. Let $\{g_k\}_{k \geq 0}$ the sequence constructed by iteration of the $R_{\sigma,t}$ -transform. Suppose $dc \neq 0$, then we have the following results:*

- i) *If $ab = 0$, then $\{g_k\}_{k \geq 0}$ is a sequence of completely normal polynomials over $\mathbb{F}_q[x]$.*
- ii) *If $ab \neq 0$, and the element $\frac{bc}{ad}$ is an ℓ -th power for some prime ℓ dividing t , then $\{g_k\}_{k \geq 0}$ is a sequence of completely normal polynomials over $\mathbb{F}_q[x]$.*
- iii) *If $ab \neq 0$, $d = -c$ and $A \neq 0$, then $\{g_k\}_{k \geq 0}$ is a sequence of completely normal polynomials over $\mathbb{F}_q[x]$.*

In section 4, we show how to recover Chapman's result from Theorem 1.3. In section 5, we prove a Theorem more general than Theorem 1.3 in that the starting polynomial does not need to be linear. We also provide explicit examples.

2. PRELIMINARIES IN ALGEBRA

This section is a brief summary about the basic results that we will use in the proof of Theorem 1.3. For more details and proofs see [4] and [5].

Let $\tau : \overline{\mathbb{F}}_q \rightarrow \overline{\mathbb{F}}_q$ be the Frobenius map defined by $\tau(\alpha) = \alpha^q$. For every $n \in \mathbb{N}$, τ induces an automorphism of \mathbb{F}_{q^n} that fixes \mathbb{F}_q . Let $f(x) = \sum_{i=0}^m a_i x^i$ be a polynomial in $\mathbb{F}_q[x]$, we define $f(\tau) \in \text{End}(\mathbb{F}_{q^n})$ by $f(\tau)(\alpha) = \sum_{i=0}^m a_i \tau^i(\alpha)$.

Assume that $(n, q) = 1$ and let

$$x^n - 1 = p_1 p_2 \dots p_r$$

be the factorization of $x^n - 1$ into irreducible factors over $\mathbb{F}_q[x]$. Samaev [5] gives the following characterisation of normal elements in the extension $\mathbb{F}_{q^n}/\mathbb{F}_q$.

Theorem 2.1 ([5], Lemma 2.1). *Let $W_l = \{\alpha \in \mathbb{F}_{q^n} : p_l(\tau)(\alpha) = 0\}$ for $l = 1, \dots, r$. Then*

$$\mathbb{F}_{q^n} = \bigoplus_{l=1}^r W_l$$

is a direct sum, where each W_l is a vector space over \mathbb{F}_q of dimension $\deg p_l$. Furthermore, an element $\alpha \in \mathbb{F}_{q^n}$ is normal if and only if for $\alpha = \sum_{l=1}^r \alpha_l$ with $\alpha_l \in W_l$, we have that $\alpha_l \neq 0$ for all $l = 1, \dots, r$.

Each V_l is irreducible τ -invariant i.e. $\tau(W_l) \subset W_l$ and it has no proper τ -invariant subspace. Also the decomposition is unique, namely, if there exist $\{V_l\}_{l=1}^s$ irreducible τ -invariant subspaces such that

$$\mathbb{F}_{q^n} = \bigoplus_{l=1}^s V_l,$$

then $s = r$ and, after rearranging the order of V_l 's if necessary, $V_l = W_l$ for $l = 1, \dots, r$.

We define an equivalence relation \sim on $\{0, 1, \dots, n-1\}$ by $l_1 \sim l_2$ iff $l_1 \equiv l_2 q^i \pmod n$ for some i . For $l \in \{0, 1, \dots, n-1\}$, let M_l be its equivalence class. Then

$$|M_l| = \min\{m \geq 1 : l \equiv l q^m \pmod n\}$$

and there exist $S \subset \{0, 1, \dots, n-1\}$ such that $\cup_{l \in S} M_l = \{0, 1, \dots, n-1\}$ and $M_l \cap M_j = \emptyset$ if $l, j \in S$ and $l \neq j$.

Theorem 2.2 ([5], Theorem 2.2). *Let $x^n - A \in \mathbb{F}_q[x]$ be an irreducible polynomial. Let $\theta \in \mathbb{F}_{q^n}$ be a root of $x^n - A$ and let S, M_l be the sets defined previously. For $l \in S$, let V_l be the subspace of \mathbb{F}_{q^n} spanned over \mathbb{F}_q by the elements $\{\theta^m : m \in M_l\}$. Then each V_l is an irreducible τ -invariant subspace and*

$$\mathbb{F}_{q^n} = \bigoplus_{l \in S} V_l.$$

In particular, an element $\alpha \in \mathbb{F}_{q^n}$ is normal if and only if for $\alpha = \sum_{l \in S} \alpha_l$ with $\alpha_l \in V_l$, we have that $\alpha_l \neq 0$ for all $l \in S$.

3. MAIN LEMMA

If f is a degree n polynomial in $\mathbb{F}_q[x]$, the polynomial $P_\sigma(f)$ is given by

$$P_\sigma(f)(x) = (cx + d)^n f\left(\frac{ax + b}{cx + d}\right).$$

An useful result about this transformation is the following Proposition.

Proposition 3.1 ([1], Proposition 3.1 and Lemma 3.2). *Let $f \in \mathbb{F}_q[x]$ be a polynomial of degree n and $\sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{F}_q)$. Then*

- i) $P_\sigma(f)$ is of degree n if and only if $f(\sigma \cdot \infty) \neq 0$.
- ii) If f is irreducible, then $P_\sigma(f)$ is irreducible in $\mathbb{F}_q[x]$
- iii) If f satisfies $f(\sigma \cdot \infty) \neq 0$ and $\alpha_1, \dots, \alpha_n$ are the roots of $P_\sigma(f)$, then the roots of f are given by β_1, \dots, β_n with

$$\beta_i = \sigma \cdot \alpha_i, \quad \text{for } i = 1, \dots, n$$

Using Theorem 2.2 and Proposition 3.1, we can find conditions such that the polynomial $P_\sigma(f)$ will be normal over $\mathbb{F}_q[x]$. The following Lemma is motivated by [[6] Theorem 3.4.1].

Lemma 3.2 (Main Lemma). *Let $\sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{F}_q)$ and $f(x) = x^n - A \neq x - a/c$ be an irreducible polynomial in $\mathbb{F}_q[x]$. Let $g = P_\sigma(f)$ and α a root of g . Then:*

- i) If $a = 0$, then g is normal if and only if $d \neq 0$, $n = p$, with p prime and q a primitive root mod p .
- ii) If $c = 0$, then g is normal if and only if $b \neq 0$, $n = p$, with p prime and q is a primitive root mod p .
- iii) If $ac \neq 0$, then g is normal if and only if $c^{n-1}dA - ba^{n-1} \neq 0$ if and only if the trace of α over \mathbb{F}_q is not zero.

Proof. Since f is irreducible and different from $x - a/c$, then $f(\sigma \cdot \infty) \neq 0$. Let α be a root of g , by Proposition 3.1 iii), then $\theta = \sigma \cdot \alpha$ is a root of $f(x) = x^n - A$ and $\sigma^{-1} \cdot \theta = \alpha$.

Suppose first that $ac = 0$. If $a = 0$, then $c \neq 0$ since $\det \sigma \neq 0$, so $\sigma^{-1} = \frac{1}{-bc} \begin{pmatrix} d & -b \\ -c & 0 \end{pmatrix}$.

Therefore

$$\alpha = \frac{d\theta - b}{-c\theta} = \frac{dA - b\theta^{n-1}}{-cA} = -\frac{d}{c} + \frac{b}{cA}\theta^{n-1} = \alpha_0 + \alpha_{n-1} \quad \alpha_0 \in V_0, \alpha_1 \in V_{n-1},$$

where V_0 and V_{n-1} are the sets defined in Theorem 2.2. It follows that α is normal if and only if $d \neq 0$ and $M_{n-1} = \{1, 2, \dots, n-1\}$. Since

$$|M_{n-1}| = \min\{m \in \mathbb{N} | (n-1)q^{m-1} \equiv n-1 \pmod{n}\} = \min\{m \in \mathbb{N} | q^{m-1} \equiv 1 \pmod{n}\}$$

and $(q, n) = 1$, then $|M_{n-1}|$ divides $\phi(n)$, where ϕ is the Euler's totient function. In particular $|M_{n-1}| \leq \phi(n)$ so $M_{n-1} = \{1, 2, \dots, n-1\}$ iff $|M_{n-1}| = n-1 \leq \phi(n)$ iff $n = p$ prime and q is a primitive root mod n . This proves i).

If $c = 0$, then $a \neq 0$ since $\det \sigma \neq 0$, so $\sigma^{-1} = \frac{1}{ad} \begin{pmatrix} d & -b \\ 0 & a \end{pmatrix}$ and

$$\alpha = \frac{d\theta - b}{a} = -\frac{b}{a} + \frac{d}{a}\theta = \alpha_0 + \alpha_1 \quad \alpha_0 \in V_0, \alpha_1 \in V_1$$

Then using Theorem 2.2 we see that α is normal if and only if $b \neq 0$ and $M_1 = \{1, 2, \dots, n-1\}$ and by a similar argument this is equivalent to $n = p$ with p prime and q a primitive root mod n , so this proves ii).

Now suppose that $ac \neq 0$. Since $f(\sigma \cdot \infty) \neq 0$ i.e. $a^n - c^n A \neq 0$, then

$$\begin{aligned} \alpha &= \frac{d\theta - b}{-c\theta + a} \cdot \left(\frac{\sum_{i=0}^{n-1} a^i (c\theta)^{n-1-i}}{\sum_{i=0}^{n-1} a^i (c\theta)^{n-1-i}} \right) \\ &= \frac{\sum_{i=1}^{n-1} \theta^i [da^{n-i}c^{i-1} - bc^i a^{n-1-i}] + c^{n-1}dA - ba^{n-1}}{a^n - c^n A} \\ &= \frac{\sum_{i=1}^{n-1} \theta^i (c/a)^{i-1} a^{n-2} \det \sigma + c^{n-1}dA - ba^{n-1}}{a^n - c^n A} \\ &= \sum_{l \in S} \alpha_l, \end{aligned}$$

where

$$\alpha_l = \begin{cases} \frac{a^{n-2} \det \sigma}{a^n - c^n A} \sum_{i \in M_l} \theta^i (c/a)^{i-1} & \text{if } l \neq 0 \\ \frac{c^{n-1}dA - ba^{n-1}}{a^n - c^n A} & \text{if } l = 0 \end{cases}.$$

Claim: $\alpha_l \neq 0$ for all $l \neq 0$.

Assume for contradiction that $\alpha_l = 0$ for some l different from zero. Then

$$\frac{a^{n-2} \det \sigma}{a^n - c^n A} \sum_{i \in M_l} \theta^i (c/a)^{i-1} = 0,$$

and since $a, c, \det \sigma \neq 0$, this relation defines a polynomial in $\mathbb{F}_q[x]$ with degree less than n having θ as a root, but $f(x) = x^n - A$ is irreducible and $f(\theta) = 0$, so $\alpha_l \neq 0$.

By using Theorem 2.2, we conclude that α is normal if and only if

$$\alpha_0 = \frac{c^{n-1}dA - ba^{n-1}}{a^n - c^n A} \neq 0,$$

if and only if $c^{n-1}dA - ba^{n-1} \neq 0$.

Otherwise, since g is irreducible by Theorem 3.1, g is the minimal polynomial for α over $\mathbb{F}_q[x]$ and

$$\begin{aligned} g &= P_\sigma(x^n - A) \\ &= (ax + b)^n - A(cx + d)^n \\ &= x^n(a^n - c^n A) + x^{n-1}(nba^{n-1} - nc^{n-1}dA) + \text{terms of degree } < (n-1), \end{aligned}$$

then the trace of α over \mathbb{F}_q is $n \left(\frac{c^{n-1}dA - ba^{n-1}}{a^n - c^n A} \right)$ and since $x^n - A$ is irreducible, necessary $(n, q) = 1$. This relation proves *iii*). ■

To use our Main Lemma in the proof of Theorem 1.3, we need the following Lemma.

Lemma 3.3. *Let $t \geq 2, \sigma \in \text{GL}_2(\mathbb{F}_q)$ and $g_0 \neq x - a/c$ be as in Theorem 1.1. For $k \geq 0$ define $f_k = P_\sigma(g_k)$. Then, f_k is irreducible. Moreover, we have that $f_k(x) = f_0(x^{t^k})$ and $P_{\sigma^{-1}}(f_k) = g_k$.*

Proof. See proof of Theorem 1.1 in [1]. ■

Now, we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. Let $\alpha_k \in \overline{\mathbb{F}_q}$ be any root of g_k and let $s|t^k$ be a divisor. The first step of the proof is to show that α_k is root of a polynomial of the form $P_{\sigma^{-1}}(x^{t^k/s} - A_{k,s})$, with $x^{t^k/s} - A_{k,s} \in \mathbb{F}_{q^s}[x]$ irreducible.

Since $g_0 = x - A \neq x - a/c$, then

$$f_0 = P_\sigma(g_0) = (ax + b) - (cx + d)(A) = (a - Ac)(x - \sigma^{-1} \cdot A) = \lambda(x - \sigma^{-1} \cdot A),$$

where $\lambda \in \mathbb{F}_q^*$ ($g_0 \neq x - a/c$). Therefore by Lemma 3.3, $f_k(x) = \lambda(x^{t^k} - \sigma^{-1} \cdot A)$ is irreducible and

$$g_k = P_{\sigma^{-1}}(f_k) = P_{\sigma^{-1}}(\lambda(x^{t^k} - \sigma^{-1} \cdot A)) = \lambda P_{\sigma^{-1}}(x^{t^k} - \sigma^{-1} \cdot A).$$

Also, we note that $g_k(\sigma \cdot \infty) \neq 0$ for all $k \geq 0$. Indeed, if $k = 0$, then $g_0(\sigma \cdot \infty) \neq 0$ because $g_0 \neq x - a/c$. For $k \geq 1$, g_k is an irreducible polynomial of degree bigger than 1, so it cannot have a root in \mathbb{F}_q . In the same way, we have that $f_k(\sigma^{-1} \cdot \infty) \neq 0$ for all $k \geq 0$. Therefore, applying Proposition 3.1 *iii*) and Lemma 3.3, we have that $\theta_k := \sigma^{-1} \cdot \alpha_k$ is a root of f_k .

Moreover, the polynomial $x^{t^k/s} - (\theta_k)^{t^k/s}$ is irreducible in $\mathbb{F}_{q^s}[x]$ and has θ_k as a root. Indeed, the polynomial $x^s - A \in \mathbb{F}[x]$ has $\theta^{t^k/s}$ as a root, but $f_k(x) = \lambda(x^{t^k} - \sigma^{-1} \cdot A)$ irreducible, then by counting degrees we have that the polynomials $x^{t^k/s} - (\theta_k)^{t^k/s}$ and $x^s - A \in \mathbb{F}[x]$ are irreducible in $\mathbb{F}_{q^s}[x]$ and $\mathbb{F}_q[x]$ respectively. Therefore, using again Proposition 3.1 *iii*), we have that α_k is root of the polynomial $P_{\sigma^{-1}}(x^{t^k/s} - (\theta_k)^{t^k/s})$. Taking $A_{k,s} = (\theta_k)^{t^k/s}$ we prove the first step.

The second step is to show that, under the assumptions of Theorem 1.3, the polynomial $P_{\sigma^{-1}}(x^{t^k/s} - (\theta_k)^{t^k/s})$ is normal over $\mathbb{F}_{q^s}[x]$, so we will have that α_k is normal in the extension $\mathbb{F}_{q^{t^k}} / \mathbb{F}_{q^s}$.

Assume for contradiction that $P_{\sigma^{-1}}(x^{t^k/s} - (\theta_k)^{t^k/s})$ is not normal over $\mathbb{F}_{q^s}[x]$. Since $dc \neq 0$, applying Lemma 3.2 *iii*) with σ^{-1} and $x^{t^k/s} - (\theta_k)^{t^k/s}$, we have that

$$(1) \quad (-c)^{t^k/s-1} a (\theta_k)^{t^k/s} + b d^{t^k/s-1} = 0.$$

To prove *i*), assume first that $b = 0$. Then equation 1 becomes

$$(-c)^{t^k/s-1} a (\theta_k)^{t^k/s} + b d^{t^k/s-1} = (-c)^{t^k/s-1} a (\theta_k)^{t^k/s} = 0,$$

but this is true iff $c = 0$, $a = 0$ or $\theta_k = 0$ (equivalently $\sigma^{-1} \cdot A = 0$). Now $c = 0$ cannot be true because $dc \neq 0$. The relation $a = 0$ cannot be true either since $\det \sigma \neq 0$. If $\sigma^{-1} \cdot A = 0$, then the polynomials $f_k(x) = x^{t^k} - \sigma^{-1}(A) = x^{t^k}$ are not irreducible, but this is a contradiction by Lemma 3.3. Then the polynomial $P_{\sigma^{-1}}(x^{t^k/s} - (\theta_k)^{t^k/s})$ is normal over $\mathbb{F}_{q^s}[x]$.

Since s is an arbitrary divisor of t^k , we have that α_k is completely normal in the extension $\mathbb{F}_{q^{t^k}}/\mathbb{F}_q$ for all $k \geq 0$.

If $a = 0$, then equation 1 becomes $bd^{t^k/s-1} = 0$ and by a similar argument this is a contradiction. So the conclusion is the same as $b = 0$. This proves *i*).

Now, assume that $ab \neq 0$. We rewrite equation 1 as

$$(2) \quad (\theta_k)^{t^k/s} = - \left(-\frac{d}{c} \right)^{t^k/s-1} \frac{b}{a} = \left(-\frac{d}{c} \right)^{t^k/s} \frac{bc}{ad}.$$

Claim: α_k is normal in the extension $\mathbb{F}_{q^{t^k}}/\mathbb{F}_{q^s}$ for each $s|t^k$ divisor different from 1.

This comes from the fact that $x^{t^k} - \sigma^{-1} \cdot A$ is the minimal polynomial of θ_k over $\mathbb{F}_q[x]$, therefore necessarily $(\theta_k)^{t^k/s} \notin \mathbb{F}_q$ for all $s|t^k$ divisor different of 1, but this is a contradiction with the equation 2 because $\sigma \in \text{GL}_2(\mathbb{F}_q)$.

For $s = 1$, equation 2 is equivalent to $\sigma^{-1} \cdot A = \left(-\frac{d}{c} \right)^{t^k} \frac{bc}{ad}$, and using

$$\eta(g_0, \sigma) = \left(-\frac{d}{c} \right) \cdot g_0 \left(\frac{b}{d} \right) \cdot g_0 \left(\frac{a}{c} \right)^{-1} = \left(-\frac{d}{c} \right) \left(\frac{b}{d} - A \right) \left(\frac{a}{c} - A \right)^{-1} = \sigma^{-1} \cdot A,$$

we have that $\eta(g_0, \sigma) = \left(-\frac{d}{c} \right)^{t^k} \frac{bc}{ad}$.

Assume that $\frac{bc}{ad}$ is a ℓ -th power for some ℓ dividing t . Then, $\eta(g, \sigma)$ is a ℓ -th power, but this is a contradiction since g satisfies the conditions of Theorem 1.1. Therefore g_k is normal over $\mathbb{F}_q[x]$ and by the previous Claim, this polynomial is completely normal over $\mathbb{F}_q[x]$. This proves *ii*).

Now, assume that $d = -c$, and $A \neq 0$. Then, $\sigma \cdot (\sigma^{-1} \cdot A) = A \neq 0$, so $\sigma^{-1} \cdot A \neq -b/a$ if and only if $a(\sigma^{-1} \cdot A) + b \neq 0$. Therefore

$$(-c)^{t^k-1} a(\sigma^{-1} \cdot A) + bd^{t^k-1} = d^{t^k-1} [a(\sigma^{-1} \cdot A) + b] \neq 0$$

for all $k \geq 0$. Then g_k is normal over $\mathbb{F}_q[x]$ for all k and by the previous Claim, this polynomial is completely normal over $\mathbb{F}_q[x]$. Thus proving *iii*). ■

4. CHAPMAN'S THEOREM

In this section, we give a proof of Chapman's Theorem using Theorem 1.3.

We recall that the R -transform corresponds to the case $t = 2$ and $\sigma^* = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ in the $R_{\sigma,t}$ -transform.

Proof of Chapman's Theorem 1.2. We will show that there exist a linear polynomial $g_0 \in \mathbb{F}_q[x]$, such that $g_1 = g_0^R$ and it satisfies the conditions of Theorem 1.1.

Since g_1 is an irreducible self reciprocal polynomial, its roots are given by γ and $1/\gamma$, so the element $A = (\gamma + \gamma^{-1})/2 = -a/2$ belongs to \mathbb{F}_q .

Claim: The polynomial $g_0 = x - A \in \mathbb{F}_q[x]$ satisfies the required conditions.

If $g_0 = x - 1$, then $a = -2$ but this is a contradiction because $g_1 = x^2 + ax + 1$ is irreducible. The polynomial g_0^R is monic and degree 2. Moreover, using that $g_0(A) = 0$ and

$$g_0^R = (2x)g_0 \left(\frac{x + 1/x}{2} \right),$$

we have that g_0^R has γ and $1/\gamma$ as its roots, this proves $g_0^R = g_1$. Also

$$\eta(g_0; \sigma) = \sigma^{-1} \cdot A = \frac{-1 - A}{-1 + A} = \frac{-2 + a}{-2 - a} = \left(\frac{1}{2 + a} \right)^2 (-4 + a^2),$$

and since $-4 + a^2$ is minus the discriminant of the irreducible polynomial g_1 and -1 is a square in \mathbb{F}_q ($q \equiv 1 \pmod{4}$), then the element $\eta(g_0; \sigma)$ is a quadratic nonresidue in \mathbb{F}_q , so g_0 satisfies the conditions of Theorem 1.1. Then we conclude using Theorem 1.3 *ii*) ■

5. GENERALIZATION OF THEOREM 1.3

In this section, we show a generalization of Theorem 1.3. This result allows us to construct families of completely normal polynomials $\{g_0\}_{k \geq 0}$ with the starting polynomial g_0 not necessarily linear. We need first the followings results.

Proposition 5.1 ([7], Theorem 1). *Let $A \in \mathbb{F}_q^*$ with multiplicative order e . Then the polynomial $x^t - A$ is irreducible if and only if the integer t satisfies the following conditions:*

- i) $(t, (q-1)/e) = 1$
- ii) For all prime factors ℓ of t , we have that ℓ divides e
- iii) If $4|n$, then $4|q-1$

Unlike the proof of Theorem 1.3, we will use the following Theorem about the irreducibility of the polynomials $\{g_k\}_{k \geq 0}$.

Theorem 5.2 ([1], Theorem 3.3). *Let $g_0 \in \mathbb{F}_q[x]$ be an irreducible polynomial of degree n and let $t \geq 2$ be a positive integer. If $q \equiv 3 \pmod{4}$ and t is even assume moreover that n is even. For $k \geq 1$ define $g_k = g_{k-1}^{R_{\sigma,t}}$. Assume that*

- i) $g_0(\sigma \cdot \infty) \neq 0$.
- ii) g_1 is irreducible.

Then, $\{g_k\}_{k \geq 0}$ is a sequence of irreducible polynomials over $\mathbb{F}_q[x]$ with $\deg g_k = t^k n$.

Theorem 5.3. *Suppose that the polynomial $f_0(x) = x^n - A$ is irreducible over $\mathbb{F}_q[x]$. If t is even, assume moreover that $q \equiv 1 \pmod{4}$. If the polynomial $g_0 = P_{\sigma^{-1}}(f_0)$ satisfies the conditions of Theorem 5.2 with $dc \neq 0$, then we have the following results.*

- i) *If $ab = 0$, then $\{g_k\}_{k \geq 0}$ is a sequence of completely normal polynomials over $\mathbb{F}_q[x]$.*
- ii) *If $ab \neq 0$, and the element $\frac{bc}{ad}$ is an ℓ -th power for some prime ℓ dividing tn , then $\{g_k\}_{k \geq 0}$ is a sequence of completely normal polynomials over $\mathbb{F}_q[x]$.*
- iii) *If $ab \neq 0$, $d = -c$ and $A \neq -b/a$, then $\{g_k\}_{k \geq 0}$ is a sequence of completely normal polynomials over $\mathbb{F}_q[x]$.*

Proof. Let α_k be a root of g_k . Similarly as in proof of the Theorem 1.3, for each s divisor of t^k , α_k is root of the polynomial $P_{\sigma^{-1}}(x^{nt^k/s} - (\theta_k)^{nt^k/s})$, with θ_k root of $f_k(x) = x^{t^k} - A$ and $x^{nt^k/s} - (\theta_k)^{nt^k/s} \in \mathbb{F}_{q^s}[x]$ irreducible.

Since $dc \neq 0$, applying Lemma 3.2 iii), we have that α_k is normal in the extension $\mathbb{F}_{q^{nt^k}} / \mathbb{F}_{q^s}$ if and only if

$$(-c)^{nt^k/s-1} aA + b d^{nt^k/s-1} \neq 0.$$

The proof of i) and iii) is the same as in Theorem 1.3. For ii) we only have to show that g_k is normal over $\mathbb{F}_q[x]$. Since $ab \neq 0$, the condition

$$(-c)^{nt^k-1} aA + b d^{nt^k-1} \neq 0,$$

is equivalent to

$$A \neq \left(-\frac{d}{c}\right)^{nt^k} \frac{bc}{ad}.$$

Let e be the multiplicative order of A , since $f_0(x) = x^n - A$ and $f_1(x) = x^{nt} - A$ are irreducible, by Proposition 5.1 i), we have that $(n, (q-1)/e) = 1$ and $(nt, (q-1)/e) = 1$, therefore A is not an ℓ -th power in \mathbb{F}_q for all prime $\ell|nt$. So, if the element $\frac{bc}{ad}$ is an ℓ -th power in \mathbb{F}_q with $\ell|nt$, then necessarily $A \neq \left(-\frac{d}{c}\right)^{nt^k} \frac{bc}{ad}$, it follows that g_k is normal over $\mathbb{F}_q[x]$. This proves ii). ■

We can construct explicitly families of completely normal polynomials using Theorem 5.3 and the following Lemma.

Lemma 5.4. *Suppose that the polynomial $x^{nt} - A$ with $A \neq -d/c$ is irreducible over $\mathbb{F}_q[x]$. If t is even assume moreover that $q \equiv 1 \pmod{4}$. Then the polynomial $g_0 = P_{\sigma^{-1}}(x^n - A)$ satisfies the conditions of Theorem 5.2.*

Proof. We define $f_0(x) = x^n - A$. Since $f_0 \neq x - a/c$ is irreducible, then $f_0(\sigma^{-1} \cdot \infty) \neq 0$, so applying Proposition 3.3 i) and ii) with $g_0 = P_{\sigma^{-1}}(f_0)$, we have that g_0 is irreducible and $g_0(\sigma \cdot \infty) \neq 0$. Similarly we have that $P_{\sigma^{-1}}(f_1)$ irreducible and by a direct calculation, this polynomial correspond to g_1 . This proves the Lemma. ■

For example, using Proposition 5.1, the polynomial $x^6 - 3$ is irreducible in $\mathbb{F}_7[x]$. Then for $t = 3$ and $\sigma = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \in \text{GL}_2(\mathbb{F}_q)$, we have that the element $\frac{bc}{ad} = 4$ is an square in \mathbb{F}_7 and $\sigma^{-1} = \begin{pmatrix} 2 & 3 \\ 3 & 3 \end{pmatrix}$. So, applying Theorem 5.3 ii), with $n = 2$ and $g_0(x) = P_{\sigma^{-1}}(x^2 - 3)$, we have a family $\{g_k\}_{k \geq 0}$ of completely normal polynomials over $\mathbb{F}_7[x]$ with $\deg g_k = 2 \cdot 3^k$ and the starting polynomial

$$g_0(x) = P_{\sigma^{-1}}(x^2 - 3) = (2x + 3)^2 - 3(3x + 2)^2 = 5x^2 + 4x + 4.$$

For \mathbb{F}_{31} , $\text{ord}(5) = 3$ and $\text{ord}(2) = 5$. So, the polynomial $x^{15} - 10$ is irreducible in $\mathbb{F}_{31}[x]$. Taking $t = 5$ and $\sigma = \begin{pmatrix} 2 & 2 \\ 2 & -2 \end{pmatrix} \in \text{GL}_2(\mathbb{F}_{31})$, we have that $A = 10 \neq -2/2 = -1$ and $\sigma^{-1} = \begin{pmatrix} 8 & 8 \\ 8 & 23 \end{pmatrix}$. Then, using Theorem 5.3 iii) with $g_0 = P_{\sigma^{-1}}(x^3 - 10)$, we get a family $\{g_k\}_{k \geq 0}$ of completely normal polynomials with $\deg g_k = 3 \cdot 5^k$ and starting polynomial

$$g_0(x) = P_{\sigma^{-1}}(x^3 - 10) = (8x + 8)^3 - 10(8x + 23)^3 = 11x^3 + x^2 + 2x + 21$$

ACKNOWLEDGEMENTS

The author thanks Ricardo Menares to motivate him to write this paper and his invaluable comments on making this work more readable.

REFERENCES

- [1] Alp Bassa and Ricardo Menares. The r-transform as a power map and its generalisations to higher degree, 2019 *arxiv.1909.02608v1*.
- [2] Stephen D. Cohen. The explicit construction of irreducible polynomials over finite fields. *Designs, Codes and Cryptography*, 2:169–174, 1992.
- [3] Robin Chapman. Completely normal elements in iterated quadratic extensions of finite fields. *Finite Fields and Their Applications*, 3(1):1 – 10, 1997.
- [4] Ian F Blake, Shuhong Gao, and Ronald C Mullin. Specific irreducible polynomials with linearly independent roots over finite fields. *Linear Algebra and its Applications*, 253(1):227 – 249, 1997.
- [5] I A Semaev. Construction of polynomials irreducible over a finite field with linearly independent roots. *Mathematics of the USSR-Sbornik*, 63(2):507–519, feb 1989.
- [6] Shuhong Gao. *Normal Bases over Finite Fields*. PhD thesis, University of Waterloo, 1993.
- [7] Stephen Cohen. On irreducible polynomials of certain types in finite fields. *Mathematical Proceedings of the Cambridge Philosophical Society*, 66:335 – 344, 09 1969.

ANIBAL ARAVENA. PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE, FACULTAD DE MATEMÁTICAS, VICUÑA MACKENNA 4860, SANTIAGO, CHILE.

E-mail address: akaravena@uc.cl