

Cosmic evolution in novel-Gauss Bonnet Gravity

Gaurav Narain^{a,*} and Hai-Qing Zhang^{a,b†}

^a *Center for Gravitational Physics, Department of Space Science, Beihang University, Beijing 100191, China.*

^b *International Research Institute for Multidisciplinary Science, Beihang University, Beijing 100191, China*

In this short paper we investigate any non-trivial effect the novel Gauss-Bonnet gravity may give rise in the cosmic evolution of the Universe in four spacetime dimensions. We start by considering a generic Friedmann-Lemaître-Robertson-Walker (FLRW) metric respecting homogeneity and isotropicity in arbitrary space-time dimension D . The metric depends on two functions: scale factor and lapse. Plugging this metric in novel Einstein-Gauss-Bonnet (EGB) gravity action, doing an integration by parts and then take the limit of $D \rightarrow 4$ give us a dynamical action in four spacetime dimensions for scale-factor and lapse. The peculiar rescaling of Gauss-Bonnet coupling by factor of $D - 4$ results in a non-trivial contribution in the action of the theory. In this paper we study this action. We investigate the dynamics of scale-factor and behavior of lapse in an empty Universe (no matter). Due to complexity of the problem we study the theory to first order in Gauss-Bonnet coupling and solve system of equation to the first order. We compute the first order correction to the on-shell action of the empty Universe and find that its sign is opposite of the leading order part. We discuss its consequences.

I. INTRODUCTION

General relativity although is a very good theory of gravity which works over a large range of energy, but it seems to be lacking a complete description of gravity. It is expected to get modified at short distances and/or in deep infrared. At short distances such modification are motivated as the quantum theory of GR has problems: theory is non-renormalizable and lacks the ability to predict [1–7]. In deep infrared the observations supporting dark-matter and dark energy [8–11] are key motivations for modification of gravity at long distances [12].

It is generally seen that any kind of modification introduced to Einstein-Hilbert gravity leads to some undesirable problems. For example at high-energies to address issues of renormalizability, a higher-derivative modification although resolves renormalizability [13–15] but introduces a nasty problem of non-unitarity. Some efforts have been made to tackle it [16–19], and the direction has seen a recent uprising where the interest has been rekindled following works in asymptotic safety approach [20–22] and ‘Agravity’ [23]. Issues of unitarity arises in such higher-derivative gravity theories as the equation of motion contains more than two time-derivatives of metric. Lovelock gravity [24–26] are a special class of higher-derivative gravity whose equation of motion remains second order in time, and in fact Einstein gravity can be extended by a whole tower of higher-curvature terms where the equation of motion still remains second-order in time.

In four spacetime dimension the Lovelock gravity also known as Gauss-Bonnet gravity is topological and doesn’t contribute in the dynamical evolution of metric. However, they play a central role in the path-integral quantization of gravity where it is used to classify topolo-

gies. Recently, it has been noticed in [27] that the Gauss-Bonnet gravity term in the action can contribute non-trivially if its coefficient in action has been rescaled by $(D - 4)$, where D is spacetime dimensionality. This motivation for rescaling arises following works [28, 29] where it was observed that Gauss-Bonnet contribution to equation of motion is proportional to $(D - 4)$. Such rescaling introduces non-trivial features coming from Gauss-Bonnet in four spacetime dimensions. This has generated tremendous interest in novel Gauss-Bonnet gravity.

The novel Gauss-Bonnet gravity [27] action is following

$$S = \frac{1}{16\pi G} \int d^D x \sqrt{-g} \left[-2\Lambda + R + \frac{\alpha}{D-4} \left(R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4R_{\mu\nu} R^{\mu\nu} + R^2 \right) \right], \quad (1)$$

where G is the Newton’s gravitational constant, Λ is cosmological constant term and α is the Gauss-Bonnet coupling. The Gauss-Bonnet coefficient is defined with a $(D - 4)$ factor in denominator. G has mass dimension M^{2-D} , Λ has mass dimension M^2 , while α has mass dimension M^{-2} .

By now the action in eq. (1) has been explored in the context of spherical black holes [30–32], star-like solutions [33], radiating solutions [39], collapsing solutions [40], even extensions to more higher-curvature Lovelock gravity theories [40, 41]. There are already a numerous investigation on the thermodynamic behavior [42–44] of these objects quasi-normal modes [34], strong cosmic-censorship [35], bending of light [36], lower-dimensional solutions [37, 38]. Using the recent observations of black-hole shadows constraints on its coupling parameter has also been investigated [45–48].

It is noticed in [49–52] that under an integration by parts the novel Gauss-Bonnet gravity action gets rid of $(D - 4)$ factors, thereby leading to a well-defined $D \rightarrow 4$ limit at the level of action. Their strategy is inspired by Kaluza-Klein compactification where they decompose the metric in two parts $\mathcal{M}_D = \mathcal{M}_4 \otimes \mathcal{M}_{D-4}$:

* gaunarain@gmail.com

† hqzhang@buaa.edu.cn

four-dimensional manifold \mathcal{M}_4 and extra-dimension piece \mathcal{M}_{D-4} . They write their action under this decomposition and smoothly remove the extra-dimension piece by taking limit. One then gets Horndeski type gravity. A similar study was done in [50, 53] using ADM decomposition. There they noticed that for a well-defined limit and a consistent theory in four dimensions one either break (a part of) the diffeomorphism invariance or have an extra degree of freedom, in agreement with the Lovelock theorem [53].

Inspired by the above work we decided to investigate the cosmological settings in the novel-Gauss-Bonnet gravity. We start by considering the most general metric respecting spatial homogeneity and isotropicity in D -spacetime dimensions. This is a generalisation of FLRW metric in D -dimensions consisting of two unknown time-dependent functions: lapse and scale-factor. This is different from the KK decomposition that has been studied in [49, 52, 53]. Plugging the generalised FLRW metric in action of theory and performing an integration by parts results in action where $D \rightarrow 4$ limit can be smoothly taken without encountering divergences. This leaves us with a well-defined action for scale factor and lapse, which gets non-trivial corrections from Gauss-Bonnet gravity part. In this paper we study this and investigate the dynamics of scale-factor and lapse in empty matter free Universe.

The paper is organized as follows: in section II we write the generic FLRW metric and compute its action, in section III we study the action for flat Universe and study the equation of motion perturbatively. We conclude with a discussion in section IV.

II. FLRW METRIC

We start by considering generalization of FLRW metric in arbitrary spacetime dimension whose dimensionality is D . We write the metric in polar co-ordinates $\{t_p, r, \theta, \dots\}$

$$ds^2 = -N_p^2(t_p)dt_p^2 + a^2(t_p) \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega_{D-2}^2 \right], \quad (2)$$

where $N_p(t_p)$ is lapse function, $a(t_p)$ is scale-factor, $k = (0, \pm 1)$ is the curvature, and $d\Omega_{D-2}$ is the metric corresponding to unit sphere in $D-2$ spatial dimensions. This metric is conformally related to flat metric and hence its Weyl-tensor $C_{\mu\nu\rho\sigma} = 0$. The nonzero entries of Riemann tensor [54–57] are

$$\begin{aligned} R_{0i0j} &= - \left(\frac{\ddot{a}}{a} - \frac{\dot{a}\dot{N}_p}{aN_p} \right) g_{ij}, \\ R_{ijkl} &= \left(\frac{k}{a^2} + \frac{\dot{a}^2}{N_p^2 a^2} \right) (g_{ik}g_{jl} - g_{il}g_{jk}), \end{aligned} \quad (3)$$

where g_{ij} is the spatial part of the FLRW metric and $(\dot{})$ denotes derivative with respect to t_p . The non-zero

components of the Ricci-tensor are

$$\begin{aligned} R_{00} &= -(D-1) \left(\frac{\ddot{a}}{a} - \frac{\dot{a}\dot{N}_p}{aN_p} \right), \\ R_{ij} &= \left[\frac{(D-2)(kN_p^2 + \dot{a}^2)}{N_p^2 a^2} + \frac{\ddot{a}N_p - \dot{a}\dot{N}_p}{aN_p^3} \right] g_{ij}, \end{aligned} \quad (4)$$

while the Ricci-scalar is given by

$$R = 2(D-1) \left[\frac{\ddot{a}N_p - \dot{a}\dot{N}_p}{aN_p^3} + \frac{(D-2)(kN_p^2 + \dot{a}^2)}{2N_p^2 a^2} \right]. \quad (5)$$

The good thing about FLRW metric is that it is Weyl-flat which implies that Riemann tensor can be expressed in terms of Ricci-tensor and Ricci scalar as follow

$$\begin{aligned} R_{\mu\nu\rho\sigma} &= \frac{R_{\mu\rho}g_{\nu\sigma} - R_{\mu\sigma}g_{\nu\rho} + R_{\nu\sigma}g_{\mu\rho} - R_{\nu\rho}g_{\mu\sigma}}{D-2} \\ &\quad - \frac{R(g_{\mu\rho}g_{\nu\sigma} - g_{\mu\sigma}g_{\nu\rho})}{(D-1)(D-2)}. \end{aligned} \quad (6)$$

This identity is valid for all conformally flat metrics and allows one to express

$$R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} = \frac{4}{D-2}R_{\mu\nu}R^{\mu\nu} - \frac{2R^2}{(D-1)(D-2)}. \quad (7)$$

By making use of this identity for conformally flat metrics in the Gauss-Bonnet action one can obtain a simplified action of the theory. In such cases we have

$$\begin{aligned} &\int d^D x \sqrt{-g} (R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 4R_{\mu\nu}R^{\mu\nu} + R^2) \\ &= \frac{D-3}{D-2} \int d^D x \sqrt{-g} \left(-R_{\mu\nu}R^{\mu\nu} + \frac{DR^2}{D-1} \right). \end{aligned} \quad (8)$$

On plugging the FRW metric of eq. (2) in the action in eq. (1) one can get an action for $a(t_p)$ and $N_p(t_p)$. On doing integration by parts it is noticed that the resulting terms are independent of factors of $(D-4)$, which cancels off. This resulting action in $D = 4$ is given by

$$\begin{aligned} S &= \frac{\pi k^{-3/2}}{8G} \int dt_p \left[(3k - \Lambda a)N_p a - \frac{3aa'}{N_p} \right. \\ &\quad \left. + \frac{3\alpha}{a} \left\{ \frac{(kN_p^2 + a'^2)^2}{N_p^3} + \frac{4ka'^2}{N_p} + \frac{4a'^4}{N_p} \right\} \right], \end{aligned} \quad (9)$$

where (\prime) denotes derivative with respect to t_p . Here one notices that the Gauss-Bonnet term gives a non-trivial contribution in $D = 4$ which is possible due to particular style of defining the Gauss-Bonnet coupling parameter. With this action one can do further analysis. To make the theory more appealing one can rescale lapse and scale factor in following manner

$$N_p(t_p)dt_p = \frac{N(t)}{a(t)}dt, \quad q(t) = a^2(t). \quad (10)$$

This transformation changes our original metric in eq. (2) into following

$$ds^2 = -\frac{N^2}{q(t)}dt^2 + q(t) \left[\frac{dr^2}{1-kr^2} + r^2 d\Omega_{D-2}^2 \right], \quad (11)$$

and our action in eq. (9) changes to following simple form.

$$S = \frac{\pi k^{-3/2}}{16G} \int dt \left[(6k - 2\Lambda q)N - \frac{3\dot{q}^2}{2N} + \frac{3\alpha}{8N^3 q} (4kN^2 + \dot{q}^2)(4kN^2 + 5\dot{q}^2) \right], \quad (12)$$

where $(\dot{})$ represent derivative with respect to t . It should be noticed that the action doesn't contains any derivative of N , which happens as we have performed integration by parts previously. This is an interesting higher-derivative action which only depends on q , \dot{q} and N . This action further acquires simplicity when $k = 0$.

III. FLAT SPACE $k = 0$

The action acquires a simplified structure in flat space ($k = 0$), this is also motivated physically as our physical Universe is also observed to be spatially flat to a high accuracy. The action in this case is given by,

$$S_{k=0} = \frac{V_3}{16\pi G} \int dt \left[-2\Lambda q N - \frac{3\dot{q}^2}{2N} + \frac{15\alpha\dot{q}^4}{8N^3 q} \right], \quad (13)$$

where V_3 is the volume corresponding to 3-dimensional space. As there is no derivative term corresponding to N appearing in action so variation of action with respect to N will result in a constraint. On the other hand, variation of action with respect to $q(t)$ will give evolution equation for $q(t)$. In the gauge $\dot{N} = 0$ we have $N(t) = N_c$, and the equation of motion for $q(t)$ is given by

$$-2N_c\Lambda + \frac{3\ddot{q}}{N_c} + \frac{45\alpha}{8N_c^3} \left(\frac{\dot{q}^4}{q^2} - \frac{4\dot{q}^2\ddot{q}}{q} \right) = 0. \quad (14)$$

This is a higher-derivative equation. The higher-derivative contribution is novel here which doesn't arise if the Gauss-Bonnet coupling wasn't rescaled by factor of $(D-4)$. In principle one has to solve for $q(t)$ from the above equation for the boundary conditions

$$q(t=0) = b_0, \quad q(t=1) = b_1. \quad (15)$$

Then plug the $q(t)$ -solution back in to the action in eq. (13) where now we are in constant- N gauge, integrate with respect to time and obtain an action for the constant lapse N_c . One then look for saddle points solution for N_c which are obtained by varying this action with respect to N_c . This will be the full saddle point solution of theory.

In practice this is not always possible. In the current case the evolution equation for $q(t)$ is quite complicated

and involve higher-derivates. We therefore try to solve the system perturbatively. We start by expanding $q(t)$ in powers of α .

$$q(t) = q_0(t) + \alpha q_1(t) + \dots, \quad (16)$$

where q_0 is zeroth-order solution while q_1 is the first order solution.

A. zeroth-order

At the zeroth order we have

$$\ddot{q} = \frac{2N_c^2\Lambda}{3}. \quad (17)$$

It is a linear second order ODE which can be solved analytically exactly. The solution obeying the boundary condition stated in eq. (15) is given by

$$q_0(t) = \frac{\Lambda N_c^2}{3}(t^2 - t) + b_0(1 - t) + b_1 t. \quad (18)$$

Plugging it back into action in eq. (13) and integrating with respect to t one gets zeroth-order part of action for N_c . This is given by

$$S_0 = \frac{V_3}{16\pi G} \left[-\frac{3(b_0 - b_1)^2}{2N_c} - (b_0 + b_1)N_c\Lambda + \frac{N_c^3\Lambda^2}{18} \right]. \quad (19)$$

Once we have the action for N_c , one can obtain saddle points of this by varying it with respect to N_c and looking for extrema. Then we see that $\partial S_0 / \partial N_c = 0$ on solving gives N_0 . Its solutions obey the equation

$$\frac{3(b_0 - b_1)^2}{2N_0^2} - (b_0 + b_1)\Lambda + \frac{N_0^2\Lambda^2}{6} = 0. \quad (20)$$

This is a quadratic in N_0^2 and will consist of four solutions which are given by

$$(N_0)_{\pm,\pm} = \pm \sqrt{\frac{3}{\Lambda}} \left(\sqrt{b_1} \pm \sqrt{b_0} \right). \quad (21)$$

At the zeroth order we notice that there are four saddle point solutions. At this level we don't receive any correction from the Gauss-Bonnet term and they agree with the known saddles in the context of Lorentzian quantum cosmology [58, 59]. Corresponding to each $(N_0)_{\pm\pm}$ we have corresponding $(q_0)_{\pm\pm}$. Each one of them lead to a different FLRW metric. Corresponding to each of them we have an on-Shell action, which is given by

$$S_0^{\text{on-shell}} = \mp \frac{V_3}{4\pi G} \sqrt{\frac{\Lambda}{3}} \left(\sqrt{b_1^3} \pm \sqrt{b_0^3} \right). \quad (22)$$

B. First order

At first order in α the equations becomes more involved. The evolution of $q(t)$ at first order is dictated by following equation

$$\ddot{q}_1 = -\frac{15}{8N_c^2} \left(\frac{\dot{q}_0^4}{q_0^2} - \frac{4\dot{q}_0^2\ddot{q}_0}{q_0} \right), \quad (23)$$

where q_0 is the zeroth order solution to $q(t)$ obtained above. The boundary conditions for $q_1(t)$ can be obtained from eq. (15) and those of q_0 . This implies that

$$q_1(t=0) = q_1(t=1) = 0. \quad (24)$$

The ODE for q_1 satisfying the these boundary conditions can be solved and its solution is given by

$$\begin{aligned} q_1(t) = & \frac{5N_c^2\Lambda^2 t(t-1)}{3} - \frac{5U}{4N_c^2} \left[\left(b_0 - b_1 + \frac{N_c^2\Lambda}{3} \right) (t-1) \right. \\ & \times \tan^{-1} \left(\frac{3(b_0 - b_1) + N_c^2\Lambda}{U} \right) + \left(b_0 - b_1 - \frac{N_c^2\Lambda}{3} \right) \\ & \times t \tan^{-1} \left(\frac{3(b_1 - b_0) + N_c^2\Lambda}{U} \right) + \left(b_1 - b_0 \right. \\ & \left. \left. + \frac{N_c^2\Lambda(2t-1)}{3} \right) \tan^{-1} \left(\frac{3(b_1 - b_0) + N_c^2\Lambda(1-2t)}{U} \right) \right] \end{aligned} \quad (25)$$

where

$$U = \sqrt{6(b_0 + b_1)^2 N_c^2 \Lambda - 9(b_0 - b_1)^2 - N_c^4 \Lambda^2}. \quad (26)$$

Once we have obtained the first order correction to $q(t)$, we can plug it back in action in eq. (13) and perform the t -integration. This results in a first order corrected action for N_c .

$$\begin{aligned} S_1 = S_0 + & \frac{V_3\alpha}{16\pi G} \left[\frac{5(b_0 - b_1)^2\Lambda}{N_c} - \frac{5(b_0 + b_1)N_c\Lambda^2}{3} \right. \\ & + \frac{10N_c^3\Lambda^3}{27} + \frac{15U^3}{108N_c^3} \left\{ \tan^{-1} \left(\frac{3(b_0 - b_1) + N_c^2\Lambda}{U} \right) \right. \\ & \left. \left. + \tan^{-1} \left(\frac{3(b_1 - b_0) + N_c^2\Lambda}{U} \right) \right\} + \dots \right]. \end{aligned} \quad (27)$$

This first order corrected action can be varied with respect to N_c again to obtain the first order correction to saddle points. To obtain this we substitute

$$N_c = N_0 + \alpha N_1 + \dots \quad (28)$$

Then N_1 can be obtained from

$$\left. \frac{\partial S_0}{\partial N_c} \right|_{N_c \rightarrow (N_0 + \alpha N_1)} = 0. \quad (29)$$

From this equation we have

$$N_1 = -\frac{5N_0\Lambda}{2}, \quad (30)$$

where N_0 is given by eq. (20). Once the expression of N_0 given in eq. (20) is plugged back in above, we get the full first order corrected lapse which is given by

$$N_c = s_1 \sqrt{\frac{3}{\Lambda}} \left(1 - \frac{5\alpha\Lambda}{2} + \dots \right) (\sqrt{b_1} + s_2 \sqrt{b_0}), \quad (31)$$

where $s_{1,2} = \{\pm, \pm\}$. The first order corrected on-shell action is given by

$$S_1^{\text{on-shell}} = \left(1 - \frac{5\alpha\Lambda}{6} + \dots \right) S_0^{\text{on-shell}}, \quad (32)$$

where $S_0^{\text{on-shell}}$ is the zeroth order on-shell action given in eq. (22).

IV. DISCUSSION AND CONCLUSION

In this short paper we investigate the non-trivial effects that arise in novel-Gauss-Bonnet gravity, where the coefficient in front of the Gauss-Bonnet term in action has been appropriately rescaled by factor of $(D-4)$. We investigate cosmological spacetimes respecting homogeneity and isotropicity. We consider a most general spacetime respecting such symmetry in D -spacetime dimensions. It consists of two functions: scale factor $a(t_p)$ and lapse $N_p(t_p)$. Plugging this in the gravitational action, doing integration by parts and taking the limit $D \rightarrow 4$, we are able to obtain a dynamical action for the scale factor $a(t_p)$. We notice that an integration by parts (and discarding a surface term) allows us to obtain an action for the resulting theory where the $D-4$ terms cancels off completely. This allows us to perform a smooth $D \rightarrow 4$ limit without witnessing divergences. Similar observations were also made in [49, 53] where the authors considered KK decomposition of metric. This interesting observation indicates that probably something deep is happening which requires more careful analysis.

Once the action of theory is obtained one can study various aspects of it. We notice that a redefinition of scale factor and lapse lead us to a simplified theory for the new variables: $q(t)$ and $N(t)$. The resulting action is a function of N , q and \dot{q} only. The action doesn't contain any derivative term for the lapse, indicating that its dynamical nature comes from the dynamics of $q(t)$ indirectly. On varying the action with respect to $q(t)$ we obtain equation of motion. This is a non-linear higher-derivative ODE, and it might be hard to find an analytic closed form solution of same. We solve the system perturbatively, doing perturbation in the Gauss-Bonnet coupling. We compute the first order correction to Einstein-Hilbert solution for q and N , for the case of empty Universe. Using these we also compute the first order correction to the on-shell action. We notice that the sign of the first order correction to on-shell action is negative. This can have serious consequences in euclidean path-integral as such correction where $\alpha > 0$ will lead to exponential growth. This perhaps maybe considered a shortcoming of this theory.

Moreover, this will also indicate that only one sign of coupling is allowed.

Given that we have an action of theory in four space-time dimensions it is then natural to investigate the quantum gravity path-integral and the corrections it achieves in the novel Gauss-Bonnet gravity. Full gravitational path-integral although quite complicated to deal with, still many nice features of it can be understood when degree of freedoms are reduced substantially. We will

present this in our next publication [60].

Acknowledgements

GN will like to thank Nirmalya Kajuri and Avinash Raju for useful discussion. GN is supported by “Zhuoyue” Fellowship (ZYPH2018-03). H. Q. Z. is supported by the National Natural Science Foundation of China (Grants No. 11675140, No. 11705005, and No. 11875095).

-
- [1] G. 't Hooft and M. J. G. Veltman, “One loop divergencies in the theory of gravitation,” *Ann. Inst. H. Poincaré Phys. Theor. A* **20**, 69 (1974).
 - [2] S. Deser, H. S. Tsao and P. van Nieuwenhuizen, “Non-renormalizability of Einstein Yang-Mills Interactions at the One Loop Level,” *Phys. Lett.* **50B**, 491 (1974). doi:10.1016/0370-2693(74)90268-8
 - [3] S. Deser and P. van Nieuwenhuizen, “One Loop Divergences of Quantized Einstein-Maxwell Fields,” *Phys. Rev. D* **10**, 401 (1974). doi:10.1103/PhysRevD.10.401
 - [4] S. Deser and P. van Nieuwenhuizen, “Nonrenormalizability of the Quantized Dirac-Einstein System,” *Phys. Rev. D* **10**, 411 (1974). doi:10.1103/PhysRevD.10.411
 - [5] M. H. Goroff and A. Sagnotti, “Quantum Gravity At Two Loops,” *Phys. Lett.* **160B**, 81 (1985). doi:10.1016/0370-2693(85)91470-4
 - [6] M. H. Goroff and A. Sagnotti, “The Ultraviolet Behavior of Einstein Gravity,” *Nucl. Phys. B* **266**, 709 (1986). doi:10.1016/0550-3213(86)90193-8
 - [7] A. E. M. van de Ven, “Two loop quantum gravity,” *Nucl. Phys. B* **378**, 309 (1992). doi:10.1016/0550-3213(92)90011-Y
 - [8] A. G. Riess *et al.* [Supernova Search Team], “Observational evidence from supernovae for an accelerating universe and a cosmological constant,” *Astron. J.* **116** (1998) 1009 doi:10.1086/300499 [astro-ph/9805201].
 - [9] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], “Measurements of Omega and Lambda from 42 high redshift supernovae,” *Astrophys. J.* **517** (1999) 565 doi:10.1086/307221 [astro-ph/9812133].
 - [10] C. Armendariz-Picon, T. Damour and V. F. Mukhanov, “ k - inflation,” *Phys. Lett. B* **458** (1999) 209 doi:10.1016/S0370-2693(99)00603-6 [hep-th/9904075].
 - [11] C. Armendariz-Picon, V. F. Mukhanov and P. J. Steinhardt, “A Dynamical solution to the problem of a small cosmological constant and late time cosmic acceleration,” *Phys. Rev. Lett.* **85** (2000) 4438 doi:10.1103/PhysRevLett.85.4438 [astro-ph/0004134].
 - [12] M. Maggiore and M. Mancarella, “Nonlocal gravity and dark energy,” *Phys. Rev. D* **90**, no. 2, 023005 (2014) doi:10.1103/PhysRevD.90.023005 [arXiv:1402.0448 [hep-th]].
 - [13] K. S. Stelle, “Renormalization of Higher Derivative Quantum Gravity,” *Phys. Rev. D* **16**, 953 (1977). doi:10.1103/PhysRevD.16.953.
 - [14] A. Salam and J. A. Strathdee, “Remarks on High-energy Stability and Renormalizability of Gravity Theory,” *Phys. Rev. D* **18**, 4480 (1978). doi:10.1103/PhysRevD.18.4480
 - [15] J. Julve and M. Tonin, “Quantum Gravity with Higher Derivative Terms,” *Nuovo Cim. B* **46**, 137 (1978). doi:10.1007/BF02748637
 - [16] G. Narain and R. Anishetty, “Short Distance Freedom of Quantum Gravity,” *Phys. Lett. B* **711**, 128 (2012) doi:10.1016/j.physletb.2012.03.070 [arXiv:1109.3981 [hep-th]].
 - [17] G. Narain and R. Anishetty, “Unitary and Renormalizable Theory of Higher Derivative Gravity,” *J. Phys. Conf. Ser.* **405**, 012024 (2012) doi:10.1088/1742-6596/405/1/012024 [arXiv:1210.0513 [hep-th]].
 - [18] G. Narain, “Signs and Stability in Higher-Derivative Gravity,” *Int. J. Mod. Phys. A* **33** (2018) no.04, 1850031 doi:10.1142/S0217751X18500318 [arXiv:1704.05031 [hep-th]].
 - [19] G. Narain, “Exorcising Ghosts in Induced Gravity,” *Eur. Phys. J. C* **77** (2017) no.10, 683 doi:10.1140/epjc/s10052-017-5249-z [arXiv:1612.04930 [hep-th]].
 - [20] A. Codello and R. Percacci, “Fixed points of higher derivative gravity,” *Phys. Rev. Lett.* **97** (2006) 221301 doi:10.1103/PhysRevLett.97.221301 [hep-th/0607128].
 - [21] M. R. Niedermaier, “Gravitational Fixed Points from Perturbation Theory,” *Phys. Rev. Lett.* **103** (2009) 101303. doi:10.1103/PhysRevLett.103.101303
 - [22] D. Benedetti, P. F. Machado and F. Saueressig, “Asymptotic safety in higher-derivative gravity,” *Mod. Phys. Lett. A* **24** (2009) 2233 doi:10.1142/S0217732309031521 [arXiv:0901.2984 [hep-th]].
 - [23] A. Salvio and A. Strumia, “Agravity,” *JHEP* **1406**, 080 (2014) doi:10.1007/JHEP06(2014)080 [arXiv:1403.4226 [hep-ph]].
 - [24] D. Lovelock, “The Einstein tensor and its generalizations,” *J. Math. Phys.* **12** (1971), 498-501 doi:10.1063/1.1665613
 - [25] D. Lovelock, “The four-dimensionality of space and the einstein tensor,” *J. Math. Phys.* **13** (1972), 874-876 doi:10.1063/1.1666069
 - [26] C. Lanczos, “A Remarkable property of the Riemann-Christoffel tensor in four dimensions,” *Annals Math.* **39** (1938), 842-850 doi:10.2307/1968467
 - [27] D. Glavan and C. Lin, “Einstein-Gauss-Bonnet gravity in 4-dimensional space-time,” *Phys. Rev. Lett.* **124** (2020) no.8, 081301 doi:10.1103/PhysRevLett.124.081301 [arXiv:1905.03601 [gr-qc]].
 - [28] A. Mardones and J. Zanelli, “Lovelock-Cartan theory of gravity,” *Class. Quant. Grav.* **8** (1991), 1545-1558 doi:10.1088/0264-9381/8/8/018
 - [29] T. Torii and H. a. Shinkai, “N+1 formalism in Einstein-Gauss-Bonnet gravity,” *Phys. Rev. D* **78** (2008), 084037 doi:10.1103/PhysRevD.78.084037 [arXiv:0810.1790 [gr-qc]].

- [30] A. Kumar and R. Kumar, “Bardeen black holes in the novel $4D$ Einstein-Gauss-Bonnet gravity,” [arXiv:2003.13104 [gr-qc]].
- [31] P. G. Fernandes, “Charged Black Holes in AdS Spaces in $4D$ Einstein Gauss-Bonnet Gravity,” [arXiv:2003.05491 [gr-qc]].
- [32] R. Kumar and S. G. Ghosh, “Rotating black holes in the novel $4D$ Einstein-Gauss-Bonnet gravity,” [arXiv:2003.08927 [gr-qc]].
- [33] D. D. Doneva and S. S. Yazadjiev, “Relativistic stars in $4D$ Einstein-Gauss-Bonnet gravity,” [arXiv:2003.10284 [gr-qc]].
- [34] A. Aragn, R. Bcar, P. Gonzlez and Y. Vsquez, “Perturbative and nonperturbative quasinormal modes of $4D$ Einstein-Gauss-Bonnet black holes,” [arXiv:2004.05632 [gr-qc]].
- [35] A. K. Mishra, “Quasinormal modes and Strong Cosmic Censorship in the novel $4D$ Einstein-Gauss-Bonnet gravity,” [arXiv:2004.01243 [gr-qc]].
- [36] M. Heydari-Fard, M. Heydari-Fard and H. Sepangi, “Bending of light in novel $4D$ Gauss-Bonnet-de Sitter black holes by Rindler-Ishak method,” [arXiv:2004.02140 [gr-qc]].
- [37] S. Nojiri and S. D. Odintsov, “Novel cosmological and black hole solutions in Einstein and higher-derivative gravity in two dimensions,” *EPL* **130** (2020) no.1, 10004 doi:10.1209/0295-5075/130/10004 [arXiv:2004.01404 [hep-th]].
- [38] R. A. Hennigar, D. Kubiznak, R. B. Mann and C. Pollack, “Lower-dimensional Gauss-Bonnet Gravity and BTZ Black Holes,” [arXiv:2004.12995 [gr-qc]].
- [39] S. G. Ghosh and S. D. Maharaj, “Radiating black holes in the novel $4D$ Einstein-Gauss-Bonnet gravity,” [arXiv:2003.09841 [gr-qc]].
- [40] A. Casalino, A. Colleaux, M. Rinaldi and S. Vicentini, “Regularized Lovelock gravity,” [arXiv:2003.07068 [gr-qc]].
- [41] R. Konoplya and A. Zhidenko, “Black holes in the four-dimensional Einstein-Lovelock gravity,” *Phys. Rev. D* **101** (2020) no.8, 084038 doi:10.1103/PhysRevD.101.084038 [arXiv:2003.07788 [gr-qc]].
- [42] C. Y. Zhang, P. C. Li and M. Guo, “Greybody factor and power spectra of the Hawking radiation in the novel $4D$ Einstein-Gauss-Bonnet de-Sitter gravity,” [arXiv:2003.13068 [hep-th]].
- [43] R. A. Konoplya and A. F. Zinhailo, “Grey-body factors and Hawking radiation of black holes in $4D$ Einstein-Gauss-Bonnet gravity,” [arXiv:2004.02248 [gr-qc]].
- [44] S. W. Wei and Y. X. Liu, “Extended thermodynamics and microstructures of four-dimensional charged Gauss-Bonnet black hole in AdS space,” [arXiv:2003.14275 [gr-qc]].
- [45] S. W. Wei and Y. X. Liu, “Testing the nature of Gauss-Bonnet gravity by four-dimensional rotating black hole shadow,” [arXiv:2003.07769 [gr-qc]].
- [46] R. Roy and S. Chakrabarti, “A study on black hole shadows in asymptotically de Sitter spacetimes,” [arXiv:2003.14107 [gr-qc]].
- [47] M. Guo and P. C. Li, “The innermost stable circular orbit and shadow in the novel $4D$ Einstein-Gauss-Bonnet gravity,” [arXiv:2003.02523 [gr-qc]].
- [48] X. X. Zeng, H. Q. Zhang and H. Zhang, “Shadows and photon spheres with spherical accretions in the four-dimensional Gauss-Bonnet black hole,” [arXiv:2004.12074 [gr-qc]].
- [49] H. Lu and Y. Pang, “Horndeski Gravity as $D \rightarrow 4$ Limit of Gauss-Bonnet,” [arXiv:2003.11552 [gr-qc]].
- [50] T. Kobayashi, “Effective scalar-tensor description of regularized Lovelock gravity in four dimensions,” [arXiv:2003.12771 [gr-qc]].
- [51] S. Mahapatra, “A note on the total action of $4D$ Gauss-Bonnet theory,” [arXiv:2004.09214 [gr-qc]].
- [52] R. A. Hennigar, D. Kubiznak, R. B. Mann and C. Pollack, “On Taking the $D \rightarrow 4$ limit of Gauss-Bonnet Gravity: Theory and Solutions,” [arXiv:2004.09472 [gr-qc]].
- [53] K. Aoki, M. A. Gorji and S. Mukohyama, “A consistent theory of $D \rightarrow 4$ Einstein-Gauss-Bonnet gravity,” [arXiv:2005.03859 [gr-qc]].
- [54] N. Deruelle and L. Farina-Busto, “The Lovelock Gravitational Field Equations in Cosmology,” *Phys. Rev. D* **41** (1990), 3696 doi:10.1103/PhysRevD.41.3696
- [55] F. Tangherlini, “Schwarzschild field in n dimensions and the dimensionality of space problem,” *Nuovo Cim.* **27** (1963), 636-651 doi:10.1007/BF02784569
- [56] F. Tangherlini, “Dimensionality of Space and the Pulsating Universe,” *Nuovo Cim.* **91** (1986), 209-217
- [57] G. Cognola, R. Myrzakulov, L. Sebastiani and S. Zerbini, “Einstein gravity with Gauss-Bonnet entropic corrections,” *Phys. Rev. D* **88** (2013) no.2, 024006 doi:10.1103/PhysRevD.88.024006 [arXiv:1304.1878 [gr-qc]].
- [58] J. Feldbrugge, J. L. Lehnert and N. Turok, “Lorentzian Quantum Cosmology,” *Phys. Rev. D* **95** (2017) no.10, 103508 doi:10.1103/PhysRevD.95.103508 [arXiv:1703.02076 [hep-th]].
- [59] J. Feldbrugge, J. L. Lehnert and N. Turok, “No smooth beginning for spacetime,” *Phys. Rev. Lett.* **119** (2017) no.17, 171301 doi:10.1103/PhysRevLett.119.171301 [arXiv:1705.00192 [hep-th]].
- [60] G. Narain, H. Q. Zhang, in preparation.