NRCPS-HE-27-2020

# *Parton Distribution Functions and Tensorgluons*

arXiv:2005.09700v2 [hep-ph] 2 Jun 2020

*Roland Kirschner*[a,1] *and George Savvidy*[b,2]

[a] *Institut für Theoretische Physik, Universität Leipzig*
*Augustusplatz 10, D-04109 Leipzig, Germany*
[b] *Institute of Nuclear and Particle Physics*
*Demokritos National Research Center, Ag. Paraskevi, Athens, Greece*

[1] *kirschne@itp.uni-leipzig.de*, [2] *savvidy@inp.demokritos.gr*

**Abstract**

We further consider a possibility that inside the proton and, more generally, inside the hadrons, there are additional partons - tensorgluons, which can carry a part of the proton momentum. The tensorgluons have zero electric charge, like gluons, but have a larger spin. Inside the proton a nonzero density of the tensorgluons can be generated by the emission of tensorgluons by gluons. The last mechanism is typical for non-Abelian tensor gauge theories. The process of gluon splitting suggests that part of the proton momentum that was carried by neutral partons is shared between vector and tensorgluons. We derive the regularised evolution equations for the parton distribution functions that take into account these new processes. In particular, this will allow to solve numerically the extended DGLAP equations and to find out the ratio of densities between gluons and tensorgluons.

# 1 *Introduction*

It was predicted that the Bjorken scaling should be broken by logarithms of a transverse momentum $Q^2$ and that these deviations from the scaling law can be computed for the deep inelastic structure functions [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13]. In the leading logarithmic approximation the results can be formulated in the parton language [14] by assigning the well determined $Q^2$ dependence to the parton densities [1, 2, 15, 16, 3].

In this article we shall further consider a possibility that inside the proton and, more generally, inside the hadrons there are additional partons - tensorgluons, which can carry a part of the proton momentum [17, 18, 19, 20, 21]. It was proposed in [11] that a possible emission of tensorgluons inside proton will produce a tensorgluon "cloud" of neutral tensorgluon partons in addition to the quark and gluon "clouds". Tensorgluons have zero electric charge, like gluons, but have a larger spin. Inside the proton a nonzero density of the tensorgluons can be generated by the emission of tensorgluons by gluons [18, 19, 20, 21]. The last mechanism is typical for non-Abelian tensor gauge theories, in which there exists a gluon-tensor-tensor vertex of order $g$. The number of gluons changes not only because a quark may radiate a gluon or because a gluon may split into a quark-antiquark pair or into two gluons [15, 16, 3], but also because a gluon can split into two tensorgluons [18, 19, 20, 21, 22, 23, 24]. The process of gluon splitting into tensorgluons suggests that part of the proton momentum which was carried by neutral partons is shared between vector and tensorgluons.

The proposed model was formulated in terms of a field theory Lagrangian, which describes the interaction of the gluons with their massless partners of higher spin, the tensorgluons [18, 19, 20, 21]. The gauge invariant Lagrangian for the lower-rank tensor gauge fields has the form [18, 19, 20]:

$$\begin{aligned}\mathcal{L} = &-\frac{1}{4}G^a_{\mu\nu}G^a_{\mu\nu} - \frac{1}{4}G^a_{\mu\nu,\lambda}G^a_{\mu\nu,\lambda} - \frac{1}{4}G^a_{\mu\nu}G^a_{\mu\nu,\lambda\lambda} \\ &+ \frac{1}{4}G^a_{\mu\nu,\lambda}G^a_{\mu\lambda,\nu} + \frac{1}{4}G^a_{\mu\nu,\nu}G^a_{\mu\lambda,\lambda} + \frac{1}{2}G^a_{\mu\nu}G^a_{\mu\lambda,\nu\lambda} + ...\end{aligned} \tag{1.1}$$

where the the field strength tensors have the form:

$$\begin{aligned}G^a_{\mu\nu} &= \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + g f^{abc}\ A^b_\mu\ A^c_\nu, \\ G^a_{\mu\nu,\lambda} &= \partial_\mu A^a_{\nu\lambda} - \partial_\nu A^a_{\mu\lambda} + g f^{abc}(\ A^b_\mu\ A^c_{\nu\lambda} + A^b_{\mu\lambda}\ A^c_\nu\ ), \\ G^a_{\mu\nu,\lambda\rho} &= \partial_\mu A^a_{\nu\lambda\rho} - \partial_\nu A^a_{\mu\lambda\rho} + g f^{abc}(\ A^b_\mu\ A^c_{\nu\lambda\rho} + A^b_{\mu\lambda}\ A^c_{\nu\rho} + A^b_{\mu\rho}\ A^c_{\nu\lambda} + A^b_{\mu\lambda\rho}\ A^c_\nu\ ),\end{aligned} \tag{1.2}$$

..... . .....

The first term in (1.1) corresponds to the standard Yang-Mills Lagrangian. The expression for the full Lagrangian which describes the infinite tour of interacting high-rank tensor gauge bosons can be found in [18, 19, 20]. For illustration we shall present the next term of the Lagrangian that describes the interaction of the rank-3 tensor gauge boson

$$\begin{aligned}\mathcal{L} = &-\frac{1}{4}G^a_{\mu\nu,\lambda\rho}G^a_{\mu\nu,\lambda\rho} - \frac{1}{8}G^a_{\mu\nu,\lambda\lambda}G^a_{\mu\nu,\rho\rho} - \frac{1}{2}G^a_{\mu\nu,\lambda}G^a_{\mu\nu,\lambda\rho\rho} - \frac{1}{8}G^a_{\mu\nu}G^a_{\mu\nu,\lambda\lambda\rho\rho} + \\ &+\frac{1}{3}G^a_{\mu\nu,\lambda\rho}G^a_{\mu\lambda,\nu\rho} + \frac{1}{3}G^a_{\mu\nu,\nu\lambda}G^a_{\mu\rho,\rho\lambda} + \frac{1}{3}G^a_{\mu\nu,\nu\lambda}G^a_{\mu\lambda,\rho\rho} + \\ &+\frac{1}{3}G^a_{\mu\nu,\lambda}G^a_{\mu\lambda,\nu\rho\rho} + \frac{2}{3}G^a_{\mu\nu,\lambda}G^a_{\mu\rho,\nu\lambda\rho} + \frac{1}{3}G^a_{\mu\nu,\nu}G^a_{\mu\lambda,\lambda\rho\rho} + \frac{1}{3}G^a_{\mu\nu}G^a_{\mu\lambda,\nu\lambda\rho\rho}.\end{aligned} \tag{1.3}$$

The characteristic property of the model is that all interaction vertices between gluons and tensorgluons have the universal *dimensionless coupling constant* $g$ in four-dimensional space-time (1.1). That is, the cubic interaction vertices have only first order derivatives and the quartic vertices have no derivatives at all. These are familiar properties of the standard Yang-Mills field theory. In order to understand the physical properties of the model it was important to study the tree-level scattering amplitudes. A very powerful spinor helicity technique [25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42] was used to calculate high-order tree-level diagrams with the participation of tensorgluons in [22].

Here we shall further develop the regularisation technique proposed earlier for the splitting amplitudes. The present paper is organised as follows. In section two the basic formulae for scattering amplitude and splitting functions are recalled, definitions and notations are specified, the details of the regularisation scheme are presented. In section three we derive the regularised evolution equations for the parton distribution functions that take into account the creation of tensorgluons. Section four contains concluding remarks and summarises the physical consequences of the tensorgluons hypothesis.

## 2 *Splitting Functions*

It was proposed in [11] that a possible emission of tensorgluons inside proton will produce a tensorgluon "cloud" of neutral tensorgluon partons in addition to the quark and gluon "clouds". Our goal is to specify the regularisation of the generalised DGLAP equations [3, 5, 6, 7, 8, 9, 10, 4] that was introduced in [11].

In the generalized Yang-Mills theory [18, 19, 20, 21] all interaction vertices between gluons and tensorgluons have *dimensionless coupling constant*. Using these vertices one can compute the scattering amplitudes of gluons and tensorgluons. The color-ordered scattering amplitudes involving two tensorgluons of helicities $h = \pm s$, one negative helicity gluon and $(n-3)$ gluons of positive helicity were found in [22]. These scattering amplitudes have been used to extract splitting amplitudes of gluons and tensorgluons [23]. Since the collinear limits of the scattering amplitudes are responsible for parton evolution, one can extract from these expressions the splitting probabilities for tensorgluons [11]:

$$\begin{aligned} P_{TG}(z) &= C_2(G)\left[\frac{z^{2s+1}}{(1-z)_+^{2s-1}} + \frac{(1-z)^{2s+1}}{z^{2s-1}}\right], \\ P_{GT}(z) &= C_2(G)\left[\frac{1}{z(1-z)_+^{2s-1}} + \frac{(1-z)^{2s+1}}{z}\right], \\ P_{TT}(z) &= C_2(G)\left[\frac{z^{2s+1}}{(1-z)_+} + \frac{1}{(1-z)_+ z^{2s-1}}\right]. \end{aligned} \tag{2.4}$$

The invariant operator $C_2$ for the representation R is defined by the equation $t^a t^a = C_2(R)\,1$ and $tr(t^a t^b) = T(R)\delta^{ab}$. These functions satisfy the relations

$$P_{TG}(z) = P_{TG}(1-z), \quad P_{GT}(z) = P_{TT}(1-z), \qquad z < 1. \tag{2.5}$$

The kernel $P_{TG}(z)$ has a meaning of variation per unit time of the probability density of finding a tensorgluon inside the gluon, $P_{GT}(z)$ - of finding gluon inside the tensorgluon and $P_{TT}(z)$ - of finding tensorgluon inside the tensorgluon. One should define the regularisation procedure for the singular factors $(1-z)^{-2s+1}$ and $z^{-2s+1}$ reinterpreting them as the distributions $(1-z)_+^{-2s+1}$ and $z_+^{-2s+1}$. The regularisation has been defined in the following way [11]:

$$\begin{aligned} \int_0^1 dz \frac{f(z)}{(1-z)_+^{2s-1}} &= \int_0^1 dz \frac{f(z) - \sum_{k=0}^{2s-2} \frac{(-1)^k}{k!} f^{(k)}(1)(1-z)^k}{(1-z)^{2s-1}}, \\ \int_0^1 dz \frac{f(z)}{z_+^{2s-1}} &= \int_0^1 dz \frac{f(z) - \sum_{k=0}^{2s-2} \frac{1}{k!} f^{(k)}(0) z^k}{z^{2s-1}}, \\ \int_0^1 dz \frac{f(z)}{z_+(1-z)_+} &= \int_0^1 dz \frac{f(z) - (1-z)f(0) - zf(1)}{z(1-z)}, \end{aligned} \tag{2.6}$$

where $f(z)$ is any test function that is sufficiently regular at the end points and, as one can see, the defined substraction guarantees the convergence of the integrals. Using the

same arguments as in the standard case [3] we should add the delta function terms into the definition of the diagonal kernels so that they will completely determine the behaviour of $P_{qq}(z)$ , $P_{GG}(z)$ and $P_{TT}(z)$ functions.

One should add $\delta(z-1)$ to the diagonal kernels $P_{qq}(z)$, $P_{GG}(z)$ and $P_{TT}(z)$ with the coefficients that have been determined by using the momentum sum rule:

$$
\begin{aligned}
P_{qq}(z) &= C_2(R)\left[\frac{1+z^2}{(1-z)_+} + \frac{3}{2}\delta(z-1)\right], \\
P_{GG}(z) &= 2C_2(G)\left[\frac{z}{(1-z)_+} + \frac{1-z}{z} + z(1-z)\right] + \frac{\sum_s (12s^2-1)C_2(G) - 4n_f T(R)}{6}\,\delta(z-1), \\
P_{TT}(z) &= C_2(G)\left[\frac{z^{2s+1}}{(1-z)_+} + \frac{1}{(1-z)_+ z^{2s-1}} + \sum_{j=1}^{2s+1}\frac{1}{j}\,\delta(z-1)\right].
\end{aligned} \tag{2.7}
$$

Thus we completed the definition of the kernels appearing in the evolution equations (3.10). For completeness we shall present also quark and gluon splitting functions [3]:

$$P_{Gq}(z) = C_2(R)\frac{1+(1-z)^2}{z}, \tag{2.8}$$

$$P_{qG}(z) = T(R)[z^2+(1-z)^2], \tag{2.9}$$

where $C_2(G) = N, C_2(R) = \frac{N^2-1}{2N}, T(R) = \frac{1}{2}$ for the SU(N) groups.

At the end of this section we shall discuss, what type of processes could also be included into the evolution equations (3.10). In (3.10) we ignore contribution of the high-spin fermions $\tilde{q}^i$ of spin $s+1/2$, which are the partners of the standard quarks [18, 19, 20, 21], supposing that they are even heavier than the top quark. That is, all kernels $P_{q\tilde{q}}, P_{G\tilde{q}}, P_{T\tilde{q}}, P_{\tilde{q}\tilde{q}}, P_{\tilde{q}q}, P_{\tilde{q}G}, P_{\tilde{q}T}$ and $P_{Tq}$ with the emission of $\tilde{q}^i$ are taken to be zero. These terms can be included in the case of very high energies, but at modern energies it seems safe to ignore these contributions. In the evolution equation for tensorgluons in (3.10) one could also include the kernels $P_{TT'}$ that describe the emission of tensorgluons by tensorgluons, the $TT'T''$ vertex [18, 19, 20, 21]. That also can be done, and the number of evolution equations in that case will tend to infinity. In this article we shall limit ourselves by considering only emissions that always involve the standard gluons and spin-2 tensors ignoring infinite "stairs" of transitions between tensorgluons.

## 3 *Regularisation of Generalised DGLAP Equations*

The deep inelastic structure functions can be expressed in terms of parton densities [3, 5, 6, 7, 8, 9, 10, 4]. If $q^i(x,Q^2)$ is the density of quarks of type i (summed over colors) inside a nucleon target with fraction x of the proton longitudinal momentum in the infinite momentum frame, then the unpolarised structure functions can be represented in the following form:

$$2F_1(x,Q^2) = F_2(x,Q^2)/x = \sum_i e_i^2[q^i(x,Q^2) + \bar{q}^i(x,Q^2)].$$

The $Q^2$ dependence of the parton densities is described by the integro-differential equations for quark $q^i(x,t)$ and gluon densities $G(x,t)$, where $t = \ln(Q^2/Q_0^2)$ [3, 5, 6, 7, 8, 9, 10, 4]. If there is an additional emission of tensorgluons in the proton, then one should introduce the corresponding density $T(x,t)$ of tensorgluons and the integro-differential equations that describe the $Q^2$ dependence of parton densities in this general case has the following form [11]:

$$\frac{dq^i(x,t)}{dt} = \frac{\alpha(t)}{2\pi}\int_x^1 \frac{dy}{y}[\sum_{j=1}^{2n_f} q^j(y,t)\ P_{q^iq^j}(\frac{x}{y}) + G(y,t)\ P_{q^iG}(\frac{x}{y})], \tag{3.10}$$

$$\frac{dG(x,t)}{dt} = \frac{\alpha(t)}{2\pi}\int_x^1 \frac{dy}{y}[\sum_{j=1}^{2n_f} q^j(y,t)\ P_{Gq^j}(\frac{x}{y}) + G(y,t)\ P_{GG}(\frac{x}{y}) + T(y,t)\ P_{GT}(\frac{x}{y})],$$

$$\frac{dT(x,t)}{dt} = \frac{\alpha(t)}{2\pi}\int_x^1 \frac{dy}{y}[G(y,t)\ P_{TG}(\frac{x}{y}) + T(y,t)\ P_{TT}(\frac{x}{y})].$$

The $\alpha(t)$ is the running coupling ($\alpha = g^2/4\pi$) and has the following form [17]:

$$\frac{\alpha}{\alpha(t)} = 1 + b\ \alpha\ t \quad , \tag{3.11}$$

where

$$b = \frac{\sum_s (-1)^{2s}(12s^2-1)C_2(G) - 4n_f T(R)}{12\pi} \tag{3.12}$$

is the one-loop Callan-Symanzik coefficient. In particular, the presence of the spin-two tensorgluons in the proton will give

$$b = \frac{58C_2(G) - 4n_f T(R)}{12\pi}. \tag{3.13}$$

The density of tensorgluons $T(x,t)$ changes when a gluon splits into two tensorgluons or when a tensorgluon radiates a gluon. This evolution is described by the last equation (3.10).

The tensorgluon kernels (2.4) are singular at the boundary values similar to the case of the standard kernels (2.8), though the singularities are of higher order compared to the standard case. The '+' prescription in

$$P(z) = \begin{pmatrix} P_{qq}(z) & 2n_f P_{qG}(z) & 0 \\ P_{Gq}(z) & P_{GG}(z) & P_{GT}(z) \\ 0 & P_{TG}(z) & P_{TT}(z) \end{pmatrix} \tag{3.14}$$

is defined as

$$[g(x)]_+ = g(x) - \delta(1-x)\int_0^1 g(z)dz, \tag{3.15}$$

and so

$$\int_x^1 f(z)[g(z)]_+ dz = \int_x^1 [f(z) - f(1)]g(z)dz - f(1)\int_x^1 g(z)dz. \tag{3.16}$$

Considering the splitting probabilities for spin two tensorgluons we have to define '+++' prescription as

$$[g(x)]_{+++} = g(x) - \delta(1-x)\int_0^1 g(z)dz - \delta^{'}(1-x)\int_0^1 g(z)(1-z)dz - \\ -\frac{1}{2}\delta^{''}(1-x)\int_0^1 g(z)(1-z)^2 dz, \tag{3.17}$$

and so

$$\int_x^1 f(z)[g(z)]_{+++}dz = \int_x^1 [f(z) - f(1) + f^{'}(1)(1-z) - \frac{1}{2}f^{''}(1)(1-z)^2]g(z)dz - \\ -\int_0^x [f(1) - f^{'}(1)(1-z) + \frac{1}{2}f^{''}(1)(1-z)^2]g(z)dz. \tag{3.18}$$

For the helicity-2 tensorgluons, the $s = 2$ in (2.4), we will have

$$\begin{aligned} P_{TG}(z) &= C_2(G)\left[\frac{z^5}{(1-z)^3} + \frac{(1-z)^5}{z^3}\right], \\ P_{GT}(z) &= C_2(G)\left[\frac{1}{z(1-z)^3} + \frac{(1-z)^5}{z}\right], \\ P_{TT}(z) &= C_2(G)\left[\frac{z^5}{(1-z)} + \frac{1}{(1-z)z^3} + \sum_{j=1}^{5}\frac{1}{j}\,\delta(1-z)\right] \end{aligned} \tag{3.19}$$

and the regularisation of these kernels can be performed using the regularisation prescription (3.18). The regular splitting functions will take the following form:

$$\int_x^1 P_{TG}(z) f(\frac{x}{z})dz = \int_x^1 \left[\frac{z^5}{(1-z)^3}_{+++} + \frac{(1-z)^5}{z^3}\right] f(\frac{x}{z})dz = I_1 + I_2 + I_3, \tag{3.20}$$

$$I_1 = \int_x^1 \frac{dz}{(1-z)^3}\Big(z^5 f(\frac{x}{z}) - f(x) + (5f(x) - xf^{'}(x))(1-z) - \\ - (10f(x) - 4xf^{'}(x) + \frac{1}{2}x^2 f^{''}(x))(1-z)^2\Big),$$

$$I_2 = -\int_0^x \frac{dz}{(1-z)^3}\Big(f(x) - (5f(x) - xf^{'}(x))(1-z) + \\ + (10f(x) - 4xf^{'}(x) + \frac{1}{2}x^2 f^{''}(x))(1-z)^2\Big),$$

$$I_3 = \int_x^1 \frac{(1-z)^5}{z^3} f(\frac{x}{z})dz.$$

For the gluon-tensor splitting function we will get:

$$\int_x^1 P_{GT}(z) f(\frac{x}{z})dz = \int_x^1 \left[\frac{3-3z+z^2}{(1-z)^3}_{+++} + \frac{1+(1-z)^5}{z}\right] f(\frac{x}{z})dz = J_1 + J_2 + J_3,$$

$$J_1 = \int_x^1 \frac{dz}{(1-z)^3}\Big((3-3z+z^2) f(\frac{x}{z}) - f(x) - (f(x) + xf^{'}(x))(1-z) - \\ - (f(x) + 2xf^{'}(x) + \frac{1}{2}x^2 f^{''}(x))(1-z)^2\Big),$$

$$J_2 = -\int_0^x \frac{dz}{(1-z)^3}[f(x) + (f(x) + xf^{'}(x)(1-z) + (f(x) + 2xf^{'}(x) + \frac{1}{2}x^2 f^{''}(x)(1-z)^2],$$

$$J_3 = \int_x^1 \frac{1+(1-z)^5}{z} f(\frac{x}{z})dz. \tag{3.21}$$

Using the regularisation (3.16) for the tensor-tensor splitting function we will get the following expression:

$$\int_x^1 P_{TT}(z) f(\frac{x}{z})dz = \int_x^1 \left[\frac{1+z^5}{(1-z)_+} + \frac{1+z+z^2}{z^3} + \sum_{j=1}^{5}\frac{1}{j}\,\delta(1-z)\right] f(\frac{x}{z})dz = \\ = \int_x^1 \frac{dz}{(1-z)}\left[(1+z^5) f(\frac{x}{z}) - 2f(x)\right] + f(x)\left[2\ln(1-x) + \frac{137}{60}\right] + \\ + \int_x^1 \frac{1+z+z^2}{z^3} f(\frac{x}{z})dz. \tag{3.22}$$

All new splitting functions which have been added to the standard evolution equation are now well defined and can be calculated using the above equations (3.20)-(3.22).

# 4 *Conclusion*

Let us summarise the physical consequences of the tensorgluons hypothesis. Among all parton distributions, the gluon density $G(x,t)$ is one of the least constrained functions since it does not couple directly to the photon in deep-inelastic scattering measurements

of the proton $F_2$ structure function. Therefore it is only indirectly constrained by scaling violations and by the momentum sum rule which resulted in the fact that only half of the proton momentum is carried by charged constituents - the quarks - and that the other part is ascribed to the neutral constituents. As it was suggested in [11], the process of gluon splitting leads to the emission of tensorgluons and therefore a part of the proton momentum that is carried by the neutral constituents here is shared between gluons and tensorgluons. The density of neutral partons in the proton is therefore given by the sum of two functions: $G(x,t) + T(x,t)$, where $T(x,t)$ is the density of the tensorgluons. Thus tensorgluons can carry a part of nucleon momentum together with gluons. Because tensorgluons have a larger spin, they can influence the spin structure of the nucleon. This contribution appears in the next to leading order, compared to the gluons and can provide a substantial screening effect due to the larger spin of tensorgluons. The details can be found in [53]. To disentangle these contributions and to decide which piece of the neutral partons is the contribution of gluons and which one is of the tensorgluons one should measure the helicities of the neutral components, which seems to be a difficult task. The other test of the proposed model will be the consistency of the mild $Q^2$ behaviour of the moments of the structure functions with the experimental data.

In supersymmetric extensions of the Standard Model [47, 48] the gluons and quarks have natural partners - gluinos of spin s=1/2 and squarks of spin s=0. If the gluinos appear as elementary constituents of the hadrons, then the theory predicts the existence of new hadronic states, the R-hadrons [49, 50]. These new hadronic states can be produced in ordinary hadronic collisions, and they decay into ordinary hadrons with the radiation of massless photino - the massless partner of the photon, which takes out a conserved R-parity. Depending on the model the gluinos may be massless or massive, depending on the remaining unbroken symmetries and the representation content of the theory. The existing experimental data give evidence that most probably they have to be very heavy [51, 52].

It seems that phenomenological limitation on the existence of the R-hadrons is much stronger that the limitation on the existence of the tensorgluons inside the ordinary hadrons. This is because gluinos change the statistics of the ordinary hadrons: the proton has to have a partner, the R-proton, which is a boson and is indeed a new hadron.

The existence of tensorgluon partons inside the proton does not predict a new hadronic state, a proton remains a proton. The tensorgluons will alternate the parton distribution

functions of a proton. The question is to which extent the tensorgluons will change the parton distribution functions. The regularisation of the splitting amplitudes developed above (3.20)-(3.22) will allow a numerical solution of the generalised DGLAP evolution equations (3.10) for the parton distribution functions that takes into account the processes of emission of tensorgluons by gluons. The integration can be performed using the algorithms developed in [43, 44, 45, 46] and will allow to find out the ratio of densities between gluons and tensorgluons.

# References


[1] D. J. Gross and F. Wilczek, *Ultraviolet Behavior Of Non-Abelian Gauge Theories,* Phys. Rev. Lett. **30**, 1343 (1973).

[2] H. D. Politzer, *Reliable Perturbative Results For Strong Interactions?,* Phys. Rev. Lett. **30**, 1346 (1973).

[3] G. Altarelli and G. Parisi, *Asymptotic Freedom in Parton Language,* Nucl. Phys. B **126** (1977) 298.

[4] Y. L. Dokshitzer, *Calculation of the Structure Functions for Deep Inelastic Scattering and e+ e- Annihilation by Perturbation Theory in Quantum Chromodynamics.,* Sov. Phys. JETP **46** (1977) 641 [Zh. Eksp. Teor. Fiz. **73** (1977) 1216].

[5] V. N. Gribov and L. N. Lipatov, *Deep inelastic e p scattering in perturbation theory,* Sov. J. Nucl. Phys. **15** (1972) 438 [Yad. Fiz. **15** (1972) 781].

[6] V. N. Gribov and L. N. Lipatov, *e+ e- pair annihilation and deep inelastic e p scattering in perturbation theory,* Sov. J. Nucl. Phys. **15** (1972) 675 [Yad. Fiz. **15** (1972) 1218].

[7] L. N. Lipatov, *The parton model and perturbation theory,* Sov. J. Nucl. Phys. **20** (1975) 94 [Yad. Fiz. **20** (1974) 181].

[8] V. S. Fadin, E. A. Kuraev and L. N. Lipatov, *On the Pomeranchuk Singularity in Asymptotically Free Theories,* Phys. Lett. B **60** (1975) 50.

[9] E. A. Kuraev, L. N. Lipatov and V. S. Fadin, *The Pomeranchuk Singularity in Non-abelian Gauge Theories,* Sov. Phys. JETP **45** (1977) 199 [Zh. Eksp. Teor. Fiz. **72** (1977) 377].

[10] I. I. Balitsky and L. N. Lipatov, *The Pomeranchuk Singularity in Quantum Chromodynamics,* Sov. J. Nucl. Phys. **28** (1978) 822 [Yad. Fiz. **28** (1978) 1597].

[11] G. Savvidy, *Proton structure and tensor gluons,* J. Phys. A **47** (2014) 35, 355401 [arXiv:1310.0856 [hep-th]].

[12] N. Cabibbo and R. Petronzio, *Two Stage Model Of Hadron Structure: Parton Distributions And Their Q**2 Dependence,* Nucl. Phys. B **137** (1978) 395.

[13] L. V. Gribov, E. M. Levin and M. G. Ryskin, *Singlet Structure Function at Small x: Unitarization of Gluon Ladders,* Nucl. Phys. B **188** (1981) 555.

[14] J. D. Bjorken and E. A. Paschos, *Inelastic Electron Proton and gamma Proton Scattering, and the Structure of the Nucleon,* Phys. Rev. **185** (1969) 1975.

[15] D. J. Gross and F. Wilczek, *Asymptotically Free Gauge Theories. 1,* Phys. Rev. D **8** (1973) 3633.

[16] D. J. Gross and F. Wilczek, *Asymptotically Free Gauge Theories. 2.,* Phys. Rev. D **9** (1974) 980.

[17] G. Savvidy, *Asymptotic freedom of non-Abelian tensor gauge fields,* Phys. Lett. B **732** (2014), 150-155 doi:10.1016/j.physletb.2014.03.022

[18] G. Savvidy, *Non-Abelian tensor gauge fields: Generalization of Yang-Mills theory*, Phys. Lett. B **625** (2005) 341

[19] G. Savvidy, *Non-abelian tensor gauge fields. I,* Int. J. Mod. Phys. A **21** (2006) 4931;

[20] G. Savvidy, *Non-abelian tensor gauge fields. II,* Int. J. Mod. Phys. A **21** (2006) 4959;

[21] G. Savvidy, *Extension of the Poincaré Group and Non-Abelian Tensor Gauge Fields,* Int. J. Mod. Phys. A **25** (2010) 5765 [arXiv:1006.3005 [hep-th]].

[22] G. Georgiou and G. Savvidy, *Production of non-Abelian tensor gauge bosons. Tree amplitudes and BCFW recursion relation,* Int. J. Mod. Phys. A **26** (2011) 2537 [arXiv:1007.3756 [hep-th]].

[23] I. Antoniadis and G. Savvidy, *Conformal invariance of tensor boson tree amplitudes,* Mod. Phys. Lett. A **27** (2012) 1250103 [arXiv:1107.4997 [hep-th]].

[24] R. Kirschner and G. Savvidy, *Yangian and SUSY symmetry of high spin parton splitting amplitudes in generalised Yang-Mills theory,* Mod. Phys. Lett. A **32** (2017) no.23, 1750121 doi:10.1142/S0217732317501218 [arXiv:1701.06660 [hep-th]].

[25] F. A. Berends, R. Kleiss, P. De Causmaecker, R. Gastmans and T. T. Wu, *Single Bremsstrahlung Processes In Gauge Theories,* Phys. Lett. B **103** (1981) 124.

[26] R. Kleiss and W. J. Stirling, *Spinor Techniques For Calculating P Anti-P* $\to W^{+-}/Z^0$ *+ Jets,* Nucl. Phys. B **262** (1985) 235.

[27] Z. Xu, D. H. Zhang and L. Chang, *Helicity Amplitudes for Multiple Bremsstrahlung in Massless Nonabelian Gauge Theories,* Nucl. Phys. B **291** (1987) 392.

[28] J. F. Gunion and Z. Kunszt, *Improved Analytic Techniques For Tree Graph Calculations And The G G Q Anti-Q Lepton Anti-Lepton Subprocess,* Phys. Lett. B **161** (1985) 333.

[29] L. J. Dixon, *Calculating scattering amplitudes efficiently,* arXiv:hep-ph/9601359.

[30] S. J. Parke and T. R. Taylor, *An Amplitude for n Gluon Scattering,* Phys. Rev. Lett. **56** (1986) 2459.

[31] F. A. Berends and W. T. Giele, *Recursive Calculations for Processes with n Gluons,* Nucl. Phys. B **306** (1988) 759.

[32] E. Witten, *Perturbative gauge theory as a string theory in twistor space,* Commun. Math. Phys. **252** (2004) 189 [arXiv:hep-th/0312171].

[33] F. Cachazo, P. Svrcek and E. Witten, *Gauge theory amplitudes in twistor space and holomorphic anomaly,* JHEP **0410** (2004) 077 [hep-th/0409245].

[34] F. Cachazo, *Holomorphic anomaly of unitarity cuts and one-loop gauge theory amplitudes*, hep-th/0410077.

[35] R. Britto, F. Cachazo and B. Feng, *New recursion relations for tree amplitudes of gluons,* Nucl. Phys. B **715** (2005) 499 [arXiv:hep-th/0412308].

[36] R. Britto, F. Cachazo, B. Feng and E. Witten, *Direct proof of tree-level recursion relation in Yang-Mills theory,* Phys. Rev. Lett. **94** (2005) 181602 [arXiv:hep-th/0501052].

[37] P. Benincasa and F. Cachazo, *Consistency Conditions on the S-Matrix of Massless Particles,* arXiv:0705.4305 [hep-th].

[38] F. Cachazo, P. Svrcek and E. Witten, *MHV vertices and tree amplitudes in gauge theory,* JHEP **0409** (2004) 006 [arXiv:hep-th/0403047].

[39] G. Georgiou, E. W. N. Glover and V. V. Khoze, *Non-MHV Tree Amplitudes in Gauge Theory,* JHEP **0407** (2004) 048 [arXiv:hep-th/0407027].

[40] N. Arkani-Hamed and J. Kaplan, *On Tree Amplitudes in Gauge Theory and Gravity,* JHEP **0804** (2008) 076 [arXiv:0801.2385 [hep-th]].

[41] F. A. Berends and W. T. Giele, *Multiple Soft Gluon Radiation in Parton Processes,* Nucl. Phys. B **313** (1989) 595.

[42] M. L. Mangano and S. J. Parke, *Quark - Gluon Amplitudes in the Dual Expansion,* Nucl. Phys. B **299** (1988) 673.

[43] M. Botje, *QCDNUM: Fast QCD Evolution and Convolution,* Comput. Phys. Commun. **182** (2011), 490-532 doi:10.1016/j.cpc.2010.10.020 [arXiv:1005.1481 [hep-ph]].

[44] V. Bertone and M. Botje, *A C++ interface to QCDNUM,* [arXiv:1712.08162 [hep-ph]].

[45] M. Miyama and S. Kumano, *Numerical solution of Q**2 evolution equations in a brute force method,* Comput. Phys. Commun. **94** (1996), 185-215 doi:10.1016/0010-4655(96)00013-6 [arXiv:hep-ph/9508246 [hep-ph]].

[46] A. Vogt, *Efficient evolution of unpolarized and polarized parton distributions with QCD-PEGASUS,* Comput. Phys. Commun. **170** (2005), 65-92 doi:10.1016/j.cpc.2005.03.103 [arXiv:hep-ph/0408244 [hep-ph]].

[47] P. Fayet, *Supersymmetry and Weak, Electromagnetic and Strong Interactions,* Phys. Lett. B **64** (1976) 159.

[48] P. Fayet, *Spontaneously Broken Supersymmetric Theories of Weak, Electromagnetic and Strong Interactions,* Phys. Lett. B **69** (1977) 489.

[49] G. R. Farrar and P. Fayet, *Phenomenology of the Production, Decay, and Detection of New Hadronic States Associated with Supersymmetry,* Phys. Lett. B **76** (1978) 575.

[50] P. Fayet, *Massive Gluinos,* Phys. Lett. B **78** (1978) 417.

[51] G. R. Farrar and P. Fayet, *Bounds on R-Hadron Production from Calorimetry Experiments,* Phys. Lett. B **79** (1978) 442.

[52] I. Antoniadis, C. Kounnas and R. Lacaze, *Light Gluinos in Deep Inelastic Scattering,* Nucl. Phys. B **211** (1983) 216.

[53] S. Konitopoulos and G. Savvidy, *Proton structure, its spin and tensor gluons,* EPJ Web Conf. **125** (2016), 04016 doi:10.1051/epjconf/201612504016