

Combining Cluster Sampling and Link-Tracing Sampling to Estimate Totals and Means of Hidden Populations in Presence of Heterogeneous Probabilities of Links

Martín H. Félix Medina

Reporte Técnico No. FCFM-UAS-2020-01

Serie: Investigación

21 de Mayo de 2020



Facultad de Ciencias Físico-Matemáticas

Universidad Autónoma de Sinaloa

Ciudad Universitaria, Culiacán Sinaloa

México

Combining Cluster Sampling and Link-Tracing Sampling to Estimate Totals and Means of Hidden Populations in Presence of Heterogeneous Probabilities of Links

Martín H. Félix Medina*

Facultad de Ciencias Físico-Matemáticas de la
Universidad Autónoma de Sinaloa

Abstract

We propose Horvitz-Thompson-like and Hájek-like estimators of the total and mean of the values of a variable of interest associated with the elements of a hard-to-reach population sampled by the variant of link-tracing sampling proposed by Félix-Medina and Thompson (2004). As examples of this type of population are drug users, homeless people and sex workers. In this sampling variant, a frame of venues or places where the members of the population tend to gather, such as parks and bars, is constructed. The frame is not assumed to cover the whole population. An initial cluster sample of elements is selected from the frame, where the clusters are the venues, and the elements in the initial sample are asked to name their contacts who are also members of the population. The sample size is increased by including in the sample the named elements who are not in the initial sample. The proposed estimators do not use design-based inclusion probabilities, but model-based inclusion probabilities which are derived from a model proposed by Félix-Medina et al. (2015) and are estimated by maximum likelihood estimators. The inclusion probabilities are assumed to be heterogeneous, that is, that they depend on the sampled people. Estimates of the variances of the proposed estimators are obtained by bootstrap and they are used to construct confidence intervals of the totals and means. The performance of the proposed estimators and confidence intervals is evaluated by two numerical studies, one of them based on real data, and the results show that their performance is acceptable.

Key words: *Chain-referral sampling, capture-recapture, Hájek estimator, Horvitz-Thompson estimator, maximum likelihood estimator, snowball sampling*

*mhfelix@uas.edu.mx

Resumen

En este trabajo se proponen estimadores tipo Horvitz-Thompson y tipo Hájek del total y la media de los valores de una variable de interés asociados con los elementos de una población de difícil detección que se muestrea mediante la variante del muestreo por bola de nieve propuesta por Félix-Medina y Thompson (2004). Ejemplos de este tipo de población son drogadictos, indigentes y trabajadoras sexuales. En esta variante de muestreo se construye un marco muestral de sitios donde los miembros de la población tienden a reunirse, tales como parques y bares. No se supone que el marco muestral cubre a toda la población. Del marco muestral se selecciona una muestra inicial por conglomerados de elementos, donde los conglomerados son los sitios, y se les pide a los elementos en la muestra inicial que nombren a sus contactos que también sean miembros de la población. El tamaño muestral se incrementa al incluir en la muestra a los elementos nombrados que no están en la muestra inicial. Los estimadores que se proponen no usan probabilidades de inclusión basadas en el diseño, sino basadas en un modelo propuesto por Félix-Medina et al. (2015) y se estiman mediante estimadores máximo verosímiles. Las probabilidades de inclusión se consideran que son heterogéneas, esto es, que dependen de las personas muestreadas. Estimaciones de las varianzas de los estimadores propuestos se obtienen mediante bootstrap y se usan para construir intervalos de confianza de los totales y medias. Los desempeños de los estimadores e intervalos de confianza propuestos se evalúan mediante dos estudios numéricos, uno de ellos basado en datos reales, y los resultados muestran que sus desempeños son aceptables.

1 Introduction

The problem of selecting samples from hidden or hard-to-detect populations, such as drug users, sex workers and homeless people, that allow reasonably valid statistical inferences is challenging because of the following factors: (i) lack of appropriate sampling frames for those populations; (ii) rareness of those populations; (iii) elusiveness of their members to be sampled; (iv) difficulty in identifying their elements due to a stigmatized or illegal behaviour; (v) difficulty in locating their members, and (v) hardness in persuading their elements to participate in the study, among others. See Tourangeau (2014) for a detailed discussion about these and other issues. Because of these factors, conventional sampling methods are not appropriate for this type of population, and consequently several specially tailored sampling methods that take into account the particular characteristics of those populations have been proposed. Among these methods we can mention multiplicity sampling, venue-based sampling, link-tracing sampling and capture-recapture sampling. See Spreen (1992), Magnani et al. (2005), Kalton (2009), Marpsat and Razafindratsima (2010), UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance (2010), Lee et al. (2014), Spreen and Bogaerts (2015) and Heckathorn and Cameron (2017) for descriptions of these methods. In addition, special estimation methods based on conventional samples, such as the scale up method (Killworth et al. 1998a and 1998b, Bernard et al. 2010, McCormick et al. 2010 and Maltiel et al. 2015) or on a combination of some of the previously mentioned sampling designs, such as the multiplier method (UNAIDS/WHO Working Group on Global HIV/AIDS

and STI Surveillance 2010, Johnston et al. 2013 and Men and Gustafson 2017) and the one step network based method (Dombrowski et al. 2012 and Khan et al. 2018) have been proposed.

It is worth noting that most of the recently published research papers on sampling from hidden populations focus on estimating the population size. (See Cheng et al. 2020 for a review and analysis, from an asymptotic approach, of most of these methods.) Thus, the scale up, multiplier and one step network methods were developed with this goal in mind. The interest in developing methods for estimating the size of a hidden population is mainly because information about this parameter allows the design of appropriate plans to address the problems associated with this type of population. However, information about other population parameters, such as average monthly spending on drugs and average age at which consumption begins in a population of drug addicts, and average weekly number of clients and average weekly income in a population of sex workers, is also important because it increases our knowledge about the population, and in addition, this knowledge could be used to improve the plans for its care that are based only on its population size.

On the other hand, among the sampling designs for hidden populations that allow estimating parameters different from or in addition to the population size, we have venue-based sampling and link-tracing sampling. Venue-based sampling (MacKellar et al., 1996) is a probability sampling method specifically developed to estimate the means of variables of interest, and particularly proportions. The method consists in carrying an ethnographic study to construct a sampling frame of venues where the members of the population tend to gather. Venues are not only sites, such as bars, parks and street locations, but could be combinations of sites, days of the week and time segments. For instance, a venue could be a specific bar from 4:00 p.m. to 12:00 a.m. on Fridays and Saturdays, whereas another venue could be the same bar, over the same time segment, but from Monday to Thursday. Furthermore, some venues could be events such as gay parades. A probability sample of venues is selected, and from each chosen venue a sort of systematic sample of members of the population who are present at the venue is selected. For each sampled element the values of the variables of interest associated with that person are recorded, and in addition, information about his or her attendance to the venues in the sampling frame is obtained so that his or her inclusion probability can be estimated, and consequently, the estimators of the means can be computed. It is evident that the estimates obtained by using this sampling design are valid only for the portion of the population that attends the venues in the frame. Therefore, the extension of the results to the entire population requires the assumption that with respect to the distributions of the variables of interest there are no differences between the members who visit the places in the frame and those who do not.

Link-tracing sampling (LTS) is an umbrella term that encompasses a set of sampling designs in which an initial sample of elements from the target population is selected and every sampled person is asked to name his or her contacts (defined according to a certain criterion) who are also members of the target population. The elements in the initial sample form the wave zero, and the named elements who are not in the initial sample form the wave one. People in wave one might also be asked to name their contacts. The named elements who have not been previously sampled form the wave two. The sampling procedure might continue

in this way until a stopping rule is satisfied. For instance, a specified number of waves or a specified sample size. Several variants of LTS have been proposed. For example, Klov Dahl (1989) developed a variant, known as random walk, in which the size of each wave is equal to one. Heckathorn (1997 and 2002) proposed a variant of LTS, called respondent driven sampling (RDS), in which every person who is already included in the sample is asked to name a small fixed number of his or her contacts randomly selected. RDS was originally proposed to estimate proportions of some subpopulations of the population of interest, such as the proportions of women and people under 18 years of age in a population of drug users. However, subsequently, Volz and Heckathorn (2008) developed estimators of the total and mean of a quantitative response variable and Handcock et al. (2014), as well as Crawford et al. (2018), developed Bayesian models which allow inferences about the population size. It is worth noting that RDS is a sampling design that has been extensively used worldwide to select samples from different hard-to-reach populations and according to Johnston et al. (2016), it has the endorsement of different organizations such as the US Centers for Disease Control and Prevention, UNAIDS, WHO and others. Another variant of LTS is the one proposed by Frank and Snijders (1994). In this variant, the initial sample is assumed to be selected by Bernoulli sampling, that is, every element of the population has the same probability of being included in the initial sample and the inclusions are independent. Furthermore, those authors assumed that the probability that a specific element of the population be named as a contact by a particular person in the initial sample, which in this document is called link probability, is a constant, that is, it does not depend on the named person nor on the person who names. This supposition is known as the homogeneity assumption. Finally, it is worth noting that Thompson and Frank (2000) proposed a stochastic block model to estimate population proportions from samples selected by a pretty general LTS design. They assumed that the population is divided into two subpopulations and that the probability of a link between two elements of the population depends on the subpopulations to which those elements belong. From their assumed model they construct maximum likelihood estimators of the proportions of those subpopulations. Chow and Thompson (2003) used the previous model and derived estimators from a Bayesian approach. Finally, St. Clair and O'Connell (2012) extended the previous Bayesian model to estimate the population mean of a quantitative variable of interest.

In this work we consider the problem of estimating the total and the mean of a variable of interest, such as the weekly drug expense of a drug user, the number of weekly clients of a sex worker, and an indicator (positive=1/negative=0) of a person's drug use, from a sample selected by the LTS variant proposed by Félix-Medina and Thompson (2004). This sampling variant was devised to avoid the assumption of an initial Bernoulli sample required by the sampling variant proposed by Frank and Snijders (1994), which is difficult to satisfy in real-world applications. To achieve this goal, those authors proposed that a sampling frame of venues or places where the members of the population tend to gather be constructed, as in venue-based sampling. The frame is not assumed to cover the whole population, but only a portion of it. Then, a simple random sample without replacement (SRSWOR) of venues is selected and the members of the population who belong to each sampled venue are sampled. Next, from each sampled venue its elements are asked to name their contacts who are also

members of the population, either they belong or not to the portion covered by the frame. It is important to indicate that, as in venue-based sampling, the venues in the frame could be combinations of places and time segments. However, in this LTS variant, an element in the portion covered by the frame is assumed that can be assigned to only one venue by using a specified criterion, for instance, the venue where the person spends most of his or her time. Félix-Medina and Thompson (2004) proposed maximum likelihood estimators (MLEs) of the population size which were derived under the assumption that the probability that a person is linked to a sampled venue, that is, that he or she is a contact of an element in that venue, depends on the venue, but not on the named person. This means that the estimators were derived under the assumption of homogeneous link probabilities. Later, Félix-Medina et al. (2015) derived MLEs of the population size under the assumption that the link probabilities also depend on the named persons, that is, that the probabilities are heterogeneous. In this work, we used the model proposed by these authors to construct model-based Horvitz-Thompson-like estimators (HTLEs) and model-based Hájek-like estimators (HKLEs) of the total and the mean of a variable of interest. It should be noted that Félix-Medina and Monjardin (2010) also considered the problem addressed in this work, but they proposed estimators of the total and the mean derived under the assumption of homogeneous link probabilities. Thus, our work is an extension of theirs.

The structure of this paper is as follows. In Section 2 we introduce the LTS variant proposed by Félix-Medina and Thompson (2004), as well as the notation to be used throughout this paper. In Section 3 we present the models and the MLEs of the population sizes proposed by Félix-Medina et al. (2015). In Section 4 we develop the strategy to construct the proposed model-based HTLEs and HKLEs of the total and the mean. In Section 5 we describe the variant of bootstrap that is proposed to construct the estimators of the variances of the HTLEs and HKLEs of the totals and means, as well as the confidence intervals of these parameters. In Section 6 we present the results of two numerical studies carried out to observe the performance of the proposed estimators and confidence intervals and to compare their performance with that of the proposed by Félix-Medina and Monjardin (2010). Finally, in Section 7 we state some conclusions and suggestions for future research.

2 Sampling design and notation

In this work we consider the variant of LTS proposed by Félix-Medina and Thompson (2004) which we will describe next. Let U be a finite population of an unknown number τ of people. A portion U_1 of U is assumed to be covered by a sampling frame of N venues A_1, \dots, A_N , where the members of the population can be found with high probability. As in ordinary cluster sampling, each person in U_1 is assumed that can be assigned to only one venue in the frame, for instance, the venue where he or she spends most of his or her time. Notice that this does not imply that a person could not be found in different venues. Let m_i denote the number of members of the population that belong, that is, that are assigned to the venue A_i , $i = 1, \dots, N$. From the previous assumption it follows that the number of people in U_1 is $\tau_1 = \sum_{i=1}^N m_i$ and the number of people in the portion $U_2 = U - U_1$ of U that is not covered

by the frame is $\tau_2 = \tau - \tau_1$.

The sampling design is as follows. A SRSWOR S_A of n venues A_1, \dots, A_n is selected from the frame. The m_i members of the population who belong to the sampled venue A_i are identified and their associated y -values of a variable of interest y are recorded, $i = 1, \dots, n$. Let S_0 be the set of people in the initial sample. Notice that the size of S_0 is $m = \sum_1^n m_i$. The people in each sampled venue are asked to name other members of the population. We will say that a person and a venue are linked if any of the people who belong to that venue names him or her. Let $x_{ij}^{(k)} = 1$ if person $j \in U_k - A_i$ is linked to venue $A_i \in S_A$ and $x_{ij}^{(k)} = 0$ if $j \in A_i$ or j is not linked to A_i , $i = 1, \dots, n$; $k = 1, 2$. For each named person, the following information is recorded: the value of the variable of interest y associated with him or her, the sampled venues that are linked to him or her, and the subset of U : $U_1 - S_0$, a specific $A_i \in S_A$ or U_2 , that contains him or her. Let S_1 be the set of people in $U_1 - S_0$ who are linked to at least one venue in S_A , and let S_2 be the set of people in U_2 who are linked to at least one venue in S_A . We will denote by r_k the size of S_k , $k = 1, 2$. Finally, let $S_1^* = S_0 \cup S_1$ and $S_2^* = S_2$ be the sets of the sampled people from U_1 and U_2 , respectively. Notice that the respective sizes of these sets are $m + r_1$ and r_2 .

We will end this section by making the following remark. Spreen and Bogaerts (2015) recently proposed a sampling procedure to estimate the size of a hidden population called B-Graph sampling, which could be considered as a particular case of the LTS variant proposed by Félix-Medina and Thompson (2004). To see this, let us firstly describe the B-Graph sampling procedure. In this method a set of incomplete sampling frames of the population of interest is obtained. For instance, in the case of a population of drug users, two incomplete sampling frames could be the register of drug users who were detained by the police and that of drug users who were attended by health clinics. An initial SRSWOR of elements is selected from the pooled sampling frame, which is the one formed by the union of the several sampling frames. Then, every sampled element is asked to name his or her contacts who are also members of the population, either they are or not in the pooled frame. For every named person, the number of elements in the initial sample who name him or her is obtained. To estimate the size of the portion of the population that is not included in the pooled frame, those authors made the following analogy between B-Graph sampling and multiple capture-recapture sampling: each person in the initial sample of their sampling design is considered as a sampling occasion of the capture-recapture procedure. Thus, the size of the portion of the population that is not covered by the pooled frame is estimated by using a capture-recapture estimator, for instance, any of the proposed by Chao (1987), Chao (1988) or Zelterman (1988). The size of the entire population is estimated by the sum of the number of elements in the pooled frame and the estimate obtained by the capture-recapture estimator. Thus, if in the variant of LTS proposed by Félix-Medina and Thompson (2004), τ_1 were known, and every m_i were equal to 1, so that $N = \tau_1$, then the only problem would be to estimate τ_2 , which is the problem considered by Spreen and Bogaerts (2015). The difference is that those authors used Chao's (1988) estimator, whereas Félix-Medina and Thompson (2004) and Félix-Medina et al. (2015) used MLEs.

3 Maximum likelihood estimators of the population sizes

Félix-Medina et al. (2015) proposed MLEs of the population sizes τ_1 , τ_2 and τ , which were derived from the following assumptions. The values m_1, \dots, m_N are considered as realizations of the random variables M_1, \dots, M_N , which are supposed to be independent and identically distributed Poisson random variables with mean λ_1 . This implies that the joint conditional distribution of the vector of variables $\mathbf{M}_s = (M_1, \dots, M_n, \tau_1 - M)$, where $M = \sum_1^n M_i$, given that $\sum_1^N M_i = \tau_1$, is multinomial with parameter of size τ_1 and vector of probabilities $(1/N, \dots, 1/N, 1 - n/N)$. The values $x_{ij}^{(k)}$ s are assumed to be realizations of the random variables $X_{ij}^{(k)}$ s, which given the sample S_A of venues, are supposed to be independent Bernoulli random variables with means $p_{ij}^{(k)}$ s, where the means or link probabilities $p_{ij}^{(k)}$ s are given by the following Rasch model:

$$p_{ij}^{(k)} = \Pr(X_{ij}^{(k)} = 1 | S_A, \alpha_{(k)i}, \beta_{(k)j}) = \frac{\exp(\alpha_{(k)i} + \beta_{(k)j})}{1 + \exp(\alpha_{(k)i} + \beta_{(k)j})}, j \in U_k - A_i; \quad i = 1, \dots, n. \quad (1)$$

As is indicated in Félix-Medina et al. (2015), this model was considered by Coull and Agresti (1999) in the context of multiple capture-recapture sampling. The parameter $\alpha_{(k)i}$ is a fixed (not random) effect that represents the potential that the venue A_i has of forming links with people in $U_k - A_i$, and $\beta_{(k)j}$ is a random effect that represents the propensity of the person $j \in U_k$ to be linked to a sampled venue. Those authors suppose that $\beta_{(k)j}$ is normally distributed with mean 0 and unknown variance σ_k^2 and that these variables are independent. The parameter σ_k^2 determines the degree of heterogeneity of the $p_{ij}^{(k)}$ s: great values of σ_k^2 imply high degrees of heterogeneity.

Henceforth all probability statements will be conditioned on the sample S_A of venues unless otherwise is specified. Now, let $\mathbf{X}_j^{(k)} = (X_{1j}^{(k)}, \dots, X_{nj}^{(k)})$ be the n -dimensional vector of link indicator variables $X_{ij}^{(k)}$ s associated with the j -th person in $U_k - S_0$, and let $\Omega = \{\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n : x_i = 0 \text{ or } x_i = 1; i = 1, \dots, n\}$. Then the probability that $\mathbf{X}_j^{(k)}$ equals $\mathbf{x} = (x_1, \dots, x_n) \in \Omega$, that is, the probability that the j -th person in $U_k - S_0$ is linked to only the venues $A_i \in S_A$ such that the i -th element x_i of \mathbf{x} equals 1, is

$$\Pr(\mathbf{X}_j^{(k)} = \mathbf{x} | \alpha_k, \beta_{(k)j}) = \prod_{i=1}^n [p_{ij}^{(k)}]^{x_i} [1 - p_{ij}^{(k)}]^{1-x_i} = \prod_{i=1}^n \frac{\exp[x_i(\alpha_{(k)i} + \beta_{(k)j})]}{1 + \exp(\alpha_{(k)i} + \beta_{(k)j})},$$

where $\alpha_k = (\alpha_{(k)1}, \dots, \alpha_{(k)n})$. Therefore, the probability that the vector of link indicator variables associated with a randomly selected person in $U_k - S_0$ equals \mathbf{x} is

$$\pi_{(k)\mathbf{x}}(\alpha_k, \sigma_k) = \int \prod_{i=1}^n \frac{\exp[x_i(\alpha_{(k)i} + \sigma_k z)]}{1 + \exp(\alpha_{(k)i} + \sigma_k z)} \phi(z) dz,$$

where $\phi(\cdot)$ denotes the probability density function of the standard normal distribution $[N(0, 1)]$.

Félix-Medina et al. (2015), following Coull and Agresti (1999), use the Gaussian quadrature method to obtain the following approximation to $\pi_{(k)\mathbf{x}}$:

$$\tilde{\pi}_{(k)\mathbf{x}} = \tilde{\pi}_{(k)\mathbf{x}}(\boldsymbol{\alpha}_k, \sigma_k) = \sum_{t=1}^q \prod_{i=1}^n \frac{\exp[x_i(\alpha_{(k)i} + \sigma_k z_t)]}{1 + \exp(\alpha_{(k)i} + \sigma_k z_t)} \nu_t, \quad (2)$$

where q is a fixed constant and $\{z_t\}$ and $\{\nu_t\}$ are obtained from tables (see Table 25.5 in Abramowitz and Stegun 1964) or statistical software (see R library statmod developed by Giner and Smyth 2016).

Similarly, for person j in $A_i \in S_A$, let $\mathbf{X}_j^{(A_i)} = (X_{1j}^{(A_i)}, \dots, X_{i-1j}^{(A_i)}, X_{i+1j}^{(A_i)}, \dots, X_{nj}^{(A_i)})$ be the $(n-1)$ -dimensional vector of link indicator variables $X_{ij}^{(A_i)}$ s associated with that person, and let $\Omega_{-i} = \{\mathbf{x} = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{R}^{n-1} : x_j = 0 \text{ or } x_j = 1; j = 1, \dots, n; j \neq i\}$. Then, the probability that $\mathbf{X}_j^{(A_i)}$ equals $\mathbf{x} = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \Omega_{-i}$, that is, the probability that the j -th person in A_i is linked to only the venues $A_{i'} \in S_A$, $i' \neq i$, such that the i' -th element $x_{i'}$ of \mathbf{x} equals 1, is

$$\Pr(\mathbf{X}_j^{(A_i)} = \mathbf{x} | \boldsymbol{\alpha}_1, \beta_{(1)j}) = \prod_{i' \neq i}^n [p_{ij'}^{(1)}]^{x_{i'}} [1 - p_{ij'}^{(1)}]^{1-x_{i'}} = \prod_{i' \neq i}^n \frac{\exp[x_{i'}(\alpha_{(1)i'} + \beta_{(1)j})]}{1 + \exp(\alpha_{(1)i'} + \beta_{(1)j})}$$

and the Gaussian quadrature approximation to the probability $\pi_{(A_i)\mathbf{x}}(\boldsymbol{\alpha}_1, \sigma_1)$ that the vector of link indicator variables associated with a randomly person selected from the sampled venue A_i equals the $(n-1)$ -dimensional vector $\mathbf{x} = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$ is

$$\tilde{\pi}_{(A_i)\mathbf{x}} = \tilde{\pi}_{(A_i)\mathbf{x}}(\boldsymbol{\alpha}_1, \sigma_1) = \sum_{t=1}^q \prod_{i' \neq i}^n \frac{\exp[x_{i'}(\alpha_{(1)i'} + \sigma_1 z_t)]}{1 + \exp(\alpha_{(1)i'} + \sigma_1 z_t)} \nu_t. \quad (3)$$

Under the previous assumptions, Félix-Medina et al. (2015) construct the likelihood function of τ_k , $\boldsymbol{\alpha}_k$ and σ_k , $k = 1, 2$, which is proportional to a product of several multinomial distributions. One multinomial distribution is the conditional distribution of the vector of variables $\mathbf{M}_s = (M_1, \dots, M_n, \tau_1 - M)$, given that $\sum_1^N M_i = \tau_1$, and that was indicated at the beginning of this section. Two others are the multinomial distributions of the vector of variables $\left(R_{\mathbf{x}}^{(k)}\right)_{\mathbf{x} \in \Omega}$, $k = 1, 2$, where $R_{\mathbf{x}}^{(1)}$ and $R_{\mathbf{x}}^{(2)}$ denote the number of people in $U_1 - S_0$ and in U_2 , respectively, whose vectors of link indicator variables $X_{ij}^{(k)}$, $k = 1, 2$, are equal to the vector $\mathbf{x} \in \Omega$. These distributions have parameters of size $\tau_1 - m$, in the case of $k = 1$, and τ_2 , in the case of $k = 2$, and vector of probabilities $(\tilde{\pi}_{(k)\mathbf{x}})_{\mathbf{x} \in \Omega}$, $k = 1, 2$. Finally, for each $i = 1, \dots, n$, we have a multinomial distribution of the vector of variables $\left(R_{\mathbf{x}}^{(A_i)}\right)_{\mathbf{x} \in \Omega_{-i}}$, where $R_{\mathbf{x}}^{(A_i)}$ denotes the number of people in A_i whose vectors of link indicator variables $X_{ij}^{(A_i)}$ are equal to the vector $\mathbf{x} \in \Omega_{-i}$. This distribution has parameters of size m_i and vector of probabilities $(\tilde{\pi}_{(A_i)\mathbf{x}})_{\mathbf{x} \in \Omega_{-i}}$. Those authors proposed maximum likelihoods estimators of τ_k , $\boldsymbol{\alpha}_k$ and σ_k , $k = 1, 2$, whose values are obtained by numerically maximizing the likelihood function. They called these estimators unconditional maximum likelihood estimators

(UMLE) and denoted them as $\hat{\tau}_k^{(U)}$, $\hat{\alpha}_k^{(U)}$ and $\hat{\sigma}_k^{(U)}$, $k = 1, 2$. Although these estimators do not have closed forms, the authors provided the following asymptotic approximations for $\hat{\tau}_1^{(U)}$ and $\hat{\tau}_2^{(U)}$:

$$\hat{\tau}_1^{(U)} = \frac{M + R_1}{1 - (1 - n/N)\tilde{\pi}_{(1)0}(\hat{\alpha}_1^{(U)}, \hat{\sigma}_1^{(U)})} \quad \text{and} \quad \hat{\tau}_2^{(U)} = \frac{R_2}{1 - \tilde{\pi}_{(2)0}(\hat{\alpha}_2^{(U)}, \hat{\sigma}_2^{(U)})}, \quad (4)$$

where R_1 and R_2 denote the numbers of distinct people in $U_1 - S_0$ and U_2 , respectively, that are linked to at least one venue in S_A . Notice that these are not close forms because $\hat{\alpha}_k^{(U)}$ and $\hat{\sigma}_k^{(U)}$ depend on $\hat{\tau}_k^{(U)}$. Once $\hat{\tau}_1^{(U)}$ and $\hat{\tau}_2^{(U)}$ are obtained, the UMLE of τ is $\hat{\tau}^{(U)} = \hat{\tau}_1^{(U)} + \hat{\tau}_2^{(U)}$.

Also, Félix-Medina et al. (2015), following Coull and Agresti (1999), used Sanathanan's (1972) approach to derive conditional MLEs $\hat{\alpha}_k^{(C)}$ and $\hat{\sigma}_k^{(C)}$ of α_k and σ_k , $k = 1, 2$, given R_k . The values of these estimators are obtained by maximizing numerically the corresponding conditional likelihood functions. Then, they showed that the conditional MLEs $\hat{\tau}_1^{(C)}$ and $\hat{\tau}_2^{(C)}$ of τ_1 and τ_2 are given by (4), but replacing $\hat{\alpha}_k^{(U)}$ and $\hat{\sigma}_k^{(U)}$ by $\hat{\alpha}_k^{(C)}$ and $\hat{\sigma}_k^{(C)}$. Notice that in this case the expressions (4) are closed forms. The conditional MLE of τ is $\hat{\tau}^{(C)} = \hat{\tau}_1^{(C)} + \hat{\tau}_2^{(C)}$.

4 Estimators of the total and mean

In this section we will focus on the problem of estimating the total and the mean of the values of a variable of interest y . Let $y_j^{(k)}$ be the value of y associated with the j -th element of U_k , $j = 1, \dots, \tau_k$, $k = 1, 2$. In this work we will suppose that the y -values are fixed numbers and not random variables. Notice that this assumption is the one made in traditional sampling. Then $Y_k = \sum_{j \in U_k} y_j^{(k)}$ and $\bar{Y}_k = Y_k/\tau_k$ represent the total and the mean of the portion U_k , $k = 1, 2$, of the population. Similarly, $Y = Y_1 + Y_2$ and $\bar{Y} = Y/\tau$ represent the total and the mean of the whole population U .

We cannot compute the design-based inclusion probabilities of the sampled elements because we do not know the venues in the frame that are linked to each sampled person, we however can compute conditional model-based inclusion probabilities given the venues $A_i \in S_A$. These probabilities are given by

$$\pi_{(1)j}(\alpha_1, \sigma_1, \beta_{(1)j}) = 1 - (1 - n/N) \prod_{i=1}^n (1 - p_{ij}^{(1)}) \quad \text{if } j \in U_1, \quad \text{and} \quad (5)$$

$$\pi_{(2)j}(\alpha_2, \sigma_2, \beta_{(2)j}) = 1 - \prod_{i=1}^n (1 - p_{ij}^{(2)}) \quad \text{if } j \in U_2. \quad (6)$$

The probabilities $\pi_{(k)j}(\alpha_k, \sigma_k, \beta_{(k)j})$ s are not known because depend on unknown parameters. However, we could estimate them by estimating those parameters and replacing in (5) and (6) the parameters by their estimates. Both UMLEs and CMLEs of α_k and σ_k have already been derived by Félix-Medina et al. (2015). We will next derive a predictor of the random effect $\beta_{(k)j}$.

Thus, if the element $j \in U_k - S_0$, $k = 1, 2$, then the conditional joint probability density function of the vector $\mathbf{X}_j^{(k)}$ of link indicator variables associated with that element and the random effect $\beta_{(k)j}$ is

$$f(\mathbf{x}_j^{(k)}, \beta_{(k)j} | \boldsymbol{\alpha}_k, \sigma_k) = \Pr(\mathbf{X}_j^{(k)} = \mathbf{x}_j^{(k)} | \beta_{(k)j}, \boldsymbol{\alpha}_k) f(\beta_{(k)j} | \sigma_k) \\ \propto \prod_{i=1}^n [p_{ij}^{(k)}]^{x_{ij}^{(k)}} [1 - p_{ij}^{(k)}]^{1-x_{ij}^{(k)}} \exp[-(\beta_{(k)j})^2 / 2\sigma_k^2],$$

whereas if the element $j \in A_{i'} \in S_A$, $i' = 1, \dots, n$, then

$$f(\mathbf{x}_j^{(A_{i'})}, \beta_{(1)j} | \boldsymbol{\alpha}_1, \sigma_1) \propto \prod_{i \neq i'}^n [p_{ij}^{(1)}]^{x_{ij}^{(A_{i'})}} [1 - p_{ij}^{(1)}]^{1-x_{ij}^{(A_{i'})}} \exp[-(\beta_{(1)j})^2 / 2\sigma_1^2].$$

We will propose as a predictor of $\beta_{(k)j}$ the conditional expected value of $\beta_{(k)j}$ given $\mathbf{X}_j^{(k)} = \mathbf{x}_j^{(k)}$, evaluated either at the UMLEs $\hat{\boldsymbol{\alpha}}_k^{(U)}$ and $\hat{\sigma}_k^{(U)}$ or at the CMLEs $\hat{\boldsymbol{\alpha}}_k^{(C)}$ and $\hat{\sigma}_k^{(C)}$, that is

$$\hat{\beta}_{(k)j}^{(a)} = E(\beta_{(k)j} | \mathbf{x}_j^{(k)}, \hat{\boldsymbol{\alpha}}_k^{(a)}, \hat{\sigma}_k^{(a)}) = \frac{\int \beta_{(k)j} f(\mathbf{x}_j^{(k)}, \beta_{(k)j} | \hat{\boldsymbol{\alpha}}_k^{(a)}, \hat{\sigma}_k^{(a)}) d\beta_{(k)j}}{\int f(\mathbf{x}_j^{(k)}, \beta_{(k)j} | \hat{\boldsymbol{\alpha}}_k^{(a)}, \hat{\sigma}_k^{(a)}) d\beta_{(k)j}}, \quad a = U, C.$$

We will approximate $\hat{\beta}_{(k)j}^{(a)}$ by using the Gaussian quadrature method, that is by

$$\tilde{\beta}_{(k)j}^{(a)} = \frac{\hat{\sigma}_k^{(a)} \sum_{t=1}^q z_t \prod_{i=1}^n \left\{ \exp[x_i(\hat{\alpha}_{(k)i}^{(a)} + \hat{\sigma}_k^{(a)} z_t)] / [1 + \exp(\hat{\alpha}_{(k)i}^{(a)} + \hat{\sigma}_k^{(a)} z_t)] \right\} \nu_t}{\sum_{t=1}^q \prod_{i=1}^n \left\{ \exp[x_i(\hat{\alpha}_{(k)i}^{(a)} + \hat{\sigma}_k^{(a)} z_t)] / [1 + \exp(\hat{\alpha}_{(k)i}^{(a)} + \hat{\sigma}_k^{(a)} z_t)] \right\} \nu_t} \\ = \frac{\hat{\sigma}_k^{(a)} \sum_{t=1}^q z_t \left\{ \exp(\hat{\sigma}_k^{(a)} z_t \sum_{i=1}^n x_i) / \prod_{i=1}^n [1 + \exp(\hat{\alpha}_{(k)i}^{(a)} + \hat{\sigma}_k^{(a)} z_t)] \right\} \nu_t}{\sum_{t=1}^q \left\{ \exp(\hat{\sigma}_k^{(a)} z_t \sum_{i=1}^n x_i) / \prod_{i=1}^n [1 + \exp(\hat{\alpha}_{(k)i}^{(a)} + \hat{\sigma}_k^{(a)} z_t)] \right\} \nu_t}, \quad a = U, C, \quad (7)$$

if $j \in U_k - S_0$, $k = 1, 2$, and by

$$\tilde{\beta}_{(1)j}^{(a)} = \frac{\hat{\sigma}_1^{(a)} \sum_{t=1}^q z_t \left\{ \exp(\hat{\sigma}_1^{(a)} z_t \sum_{i \neq i'}^n x_i) / \prod_{i \neq i'}^n [1 + \exp(\hat{\alpha}_{(1)i}^{(a)} + \hat{\sigma}_1^{(a)} z_t)] \right\} \nu_t}{\sum_{t=1}^q \left\{ \exp(\hat{\sigma}_1^{(a)} z_t \sum_{i \neq i'}^n x_i) / \prod_{i \neq i'}^n [1 + \exp(\hat{\alpha}_{(1)i}^{(a)} + \hat{\sigma}_1^{(a)} z_t)] \right\} \nu_t}, \quad a = U, C,$$

if $j \in A_{i'} \in S_A$, $i' = 1, \dots, n$.

The previous expressions imply that $\tilde{\beta}_{(k)j}^{(a)}$ depends on the x_i s through their sum, that is, on the number of venues that are linked to the element j , but not on the particular venues to which that element is linked. Thus, if two persons j and j' in $U_k - S_0$ are linked to the same number of venues in S_A , the predictors $\tilde{\beta}_{(k)j}^{(a)}$ and $\tilde{\beta}_{(k)j'}^{(a)}$ are equal one another. The same happens for two persons in $A_i \in S_A$.

Thus, model-based Horvitz-Thompson-like estimators (HTLEs) of the totals Y_k , $k = 1, 2$, and Y are

$$\hat{Y}_{HT.k}^{(a)} = \sum_{j \in S_k^*} y_j^{(k)} / \hat{\pi}_{(k)j}^{(a)} (\hat{\alpha}_k^{(a)}, \hat{\sigma}_k^{(a)}, \tilde{\beta}_{(k)j}^{(a)}), \quad k = 1, 2, \text{ and } \hat{Y}_{HT}^{(a)} = \hat{Y}_{HT.1}^{(a)} + \hat{Y}_{HT.2}^{(a)}, \quad a = U, C. \quad (8)$$

Similarly, model-based HTLEs of the means \bar{Y}_k and \bar{Y} are

$$\hat{\bar{Y}}_{HT.k}^{(a)} = \hat{Y}_{HT.k}^{(a)} / \hat{\tau}_k^{(a)}, \quad k = 1, 2, \quad \text{and} \quad \hat{\bar{Y}}_{HT}^{(a)} = \hat{Y}_{HT}^{(a)} / \hat{\tau}^{(a)}, \quad a = U, C.$$

Notice that if we set $y_j^{(k)} = 1$, for $j = 1, \dots, \tau_k$ and $k = 1, 2$, then $Y_k = \tau_k$, $k = 1, 2$, and $Y = \tau$. Therefore, HTLEs of τ_k and τ are $\hat{\tau}_{HT.k}^{(a)} = \hat{Y}_{HT.k}^{(a)}$, $k = 1, 2$, and $\hat{\tau}_{HT}^{(a)} = \hat{Y}_{HT}^{(a)}$, $a = U, C$, where $\hat{Y}_{HT.k}^{(a)}$ and $\hat{Y}_{HT}^{(a)}$ are given by (8) with $y_j^{(k)} = 1$.

We could also define Hájek-like estimators (HKLEs) of the population totals and means. Thus, HKLEs of the means \bar{Y}_k and \bar{Y} are

$$\hat{\bar{Y}}_{HK.k}^{(a)} = \hat{Y}_{HT.k}^{(a)} / \hat{\tau}_{HT.k}^{(a)}, \quad k = 1, 2, \quad \text{and} \quad \hat{\bar{Y}}_{HK}^{(a)} = \hat{Y}_{HT}^{(a)} / \hat{\tau}_{HT}^{(a)}, \quad a = U, C,$$

and HKLEs of the totals Y_k and Y are

$$\hat{Y}_{HK.k}^{(a)} = \hat{\bar{Y}}_{HK.k}^{(a)} \hat{\tau}_k^{(a)}, \quad k = 1, 2, \quad \text{and} \quad \hat{Y}_{HK}^{(a)} = \hat{\bar{Y}}_{HK}^{(a)} \hat{\tau}^{(a)}, \quad a = U, C.$$

5 Bootstrap variance estimators and confidence intervals

We propose the use of bootstrap to construct estimators of the variances of the proposed estimators of the totals and means, as well as confidence intervals (CIs) for those population parameters. The proposed bootstrap variant is obtained by combining the bootstrap version for finite populations proposed by Booth et al. (1994) and the parametric bootstrap variant (see Davison and Hinkley, 1997, Ch. 2). This version of bootstrap is an extension of the one used by Félix-Medina et al. (2015) to construct CIs based on their proposed MLEs of the population sizes.

Hereinafter, we will denote by $\lfloor x \rfloor$, the greatest integer less than or equal to $x \in \mathbb{R}$. The steps of the proposed bootstrap procedure are the following. (i) Construct a population vector \mathbf{m}_{Boot} of N values of m_i s by means of the following procedure. If N/n is an integer, repeat N/n times the observed sample of n cluster sizes $\mathbf{m}_s = \{m_1, \dots, m_n\}$. If N/n is not an integer, that is, if $N = an + b$, where a and b , $b < n$, are positive integers, then repeat a times \mathbf{m}_s and add to this set a SRSWOR of b values of m_i s selected from \mathbf{m}_s . If the sum of the elements of the vector \mathbf{m}_{Boot} is greater than the value $\hat{\tau}_1^{(a)}$, $a = U, C$, (depending of the type of estimator that is being considered), delete one element at a time from \mathbf{m}_{Boot} starting from the N -th element until the sum is less than or equal to $\hat{\tau}_1^{(a)}$. Let N_{Boot} be the final number of elements in \mathbf{m}_{Boot} . (ii) For each $k = 1, 2$, construct a population vector $\hat{\alpha}_{(k)Boot}^{(a)}$ of length N_{Boot} whose elements are the estimates $\hat{\alpha}_{(k)i}^{(a)}$ s of the $\alpha_{(k)i}$ s associated with the clusters whose

sizes m_i s are in \mathbf{m}_{Boot} . (iii) For each $k = 1, 2$, construct a population vector $\hat{\beta}_{(k)Boot}^{(a)}$ of length $\hat{\tau}_k^{(a)}$ whose first $m + r_1$ elements in the case of $k = 1$, or whose first r_2 elements in the case of $k = 2$, are the estimates $\hat{\beta}_{(k)j}^{(a)}$ s of the $\beta_{(k)j}$ s associated with the people in S_k^* , and each one of the remaining elements is the estimate $\hat{\beta}_{(k)0}^{(a)}$ of $\beta_{(k)0}$ obtained using (7) with $x_i = 0$, $i = 1, \dots, n$. (iv) For each $k = 1, 2$, construct a population vector $\hat{y}_{(k)Boot}^{(a)}$ of length $\hat{\tau}_k^{(a)}$ whose first $m + r_1$ elements in the case of $k = 1$, or whose first r_2 elements in the case of $k = 2$, are the y -values associated with the elements in S_k^* , and the remaining elements are estimates of the y -values associated with the people in $U_k - S_k^*$ and obtained using the following procedure. If the variable of interest y is continuous, then fit a simple linear regression model to the data $(\hat{\pi}_{(k)j}^{(a)}(\hat{\alpha}_k^{(a)}, \hat{\sigma}_k^{(a)}, \tilde{\beta}_{(k)j}^{(a)}), y_j^{(k)})$, $j \in S_k^*$. Next, predict the y -value associated with the j -th element in $U_k - S_k^*$ by using a value sampled from the normal distribution with mean equals to the quantity obtained by evaluating the fitted model at the estimate $\hat{\pi}_{(k)0}^{(a)}(\hat{\alpha}_k^{(a)}, \hat{\sigma}_k^{(a)}, \tilde{\beta}_{(k)0}^{(a)})$ of the inclusion probability of an element in $U_k - S_k^*$ and variance equals to the estimate of the variance of the error terms of the regression model. If the design matrix is numerically singular, then predict the y -value associated with $j \in U_k - S_k^*$ by a value sampled from the normal distribution with mean and variance given by the sample mean and sample variance, respectively, of the y -values associated with the elements in S_k^* . On the other hand, if the variable of interest y is binary, then fit a simple logistic regression model to the data $(\hat{\pi}_{(k)j}^{(a)}(\hat{\alpha}_k^{(a)}, \hat{\sigma}_k^{(a)}, \tilde{\beta}_{(k)j}^{(a)}), y_j^{(k)})$, $j \in S_k^*$. Next, predict the y -value associated with the j -th element in $U_k - S_k^*$ by using a value sampled from the Bernoulli distribution with success probability equals to the quantity obtained by evaluating the fitted model at $\hat{\pi}_{(k)0}^{(a)}(\hat{\alpha}_k^{(a)}, \hat{\sigma}_k^{(a)}, \tilde{\beta}_{(k)0}^{(a)})$. If the design matrix is numerically singular, then predict the y -value associated with $j \in U_k - S_k^*$ by a value sampled from the Bernoulli distribution with success probability equals to the sample mean of the y -values of the elements in S_k^* . (v) Select a SRSWOR of n values m_i from \mathbf{m}_{Boot} . Let $S_A^{Boot} = \{i_1, \dots, i_n\}$ be the set of indices of the m_i s in the sample. In addition, let $A_i^{Boot} = (\sum_{t=1}^{i-1} m_t, \sum_{t=1}^i m_t] \cap \mathbb{Z}$ be the set of indices j associated with the elements in the cluster whose index is $i \in S_A^{Boot}$, where m_t is the t -th element of \mathbf{m}_{Boot} and \mathbb{Z} is the set of the integer numbers. Finally, let $S_0^{Boot} = \cup_{i \in S_A^{Boot}} A_i^{Boot}$ be the set of indices j associated with the elements in the clusters whose indices are in S_A^{Boot} . (vi) For each $k = 1, 2$, $i \in S_A^{Boot}$ and $j \in \{1, \dots, \lfloor \hat{\tau}_1^{(a)} \rfloor\} - A_i^{Boot}$ in the case of $k = 1$, or $j \in \{1, \dots, \lfloor \hat{\tau}_2^{(a)} \rfloor\}$ in the case of $k = 2$, generate a value $x_{ij}^{(k)}$ by sampling from the Bernoulli distribution with success probability equals to the value obtained by evaluating (1) at the i -th element of the vector $\hat{\alpha}_{(k)Boot}^{(a)}$ and the j -th element of the vector $\hat{\beta}_{(k)Boot}^{(a)}$. (vii) Compute the estimates of the sizes τ_1 , τ_2 and τ ; those of the totals Y_1 , Y_2 and Y , and those of the means \bar{Y}_1 , \bar{Y}_2 and \bar{Y} using the same procedure as that used to compute the original estimates. (viii) Repeat the steps (v)-(vii) a large enough number B of times.

To construct the CIs for the population totals and means we could use any of the different bootstrap alternatives that have been proposed. For instance, if we did not want to assume any probability distribution for an estimator, we could use the basic or the percentile method. (See Davison and Hinkley, 1997, Ch. 5, for descriptions of these methods.) Although this

type of alternative has good properties of robustness, it requires a large number B of bootstrap samples, say $B = 1000$, and this might be a serious problem if the estimator require much time to be computed. On the other hand, if we were willing to assume a distribution probability for an estimator, we could use the B bootstrap estimates that were computed using that estimator to estimate its variance and construct the CI using the assumed distribution and the estimated variance. In this case, the number B of required bootstrap samples is not so large, say $50 \leq B \leq 200$ is generally enough. We will follow this approach using some ideas taken from Félix-Medina et al. (2015).

Thus, as in that paper, we will estimate the variance of an estimator $\hat{\theta}$ of the population parameter θ , by using Huber's proposal 2 to jointly estimate the parameters of location and scale from the bootstrap sample of B values $\hat{\theta}_b$. (See Staudte and Sheather, 1990, Sec. 4.5, for a description of this method.) In particular, the estimate of the parameter of scale is an estimate of the standard deviation $\sqrt{\hat{V}(\hat{\theta})}$ of $\hat{\theta}$. The idea behind the use of this estimator is that it yields an estimate of the standard deviation that is robust to very large values $\hat{\theta}_b$ which are likely to occur.

To construct the CIs we will use the following approach. (i) If the parameter is τ_k , $k = 1, 2$, or τ , then, as in Félix-Medina et al. (2015), we will assume that $\hat{\tau}_k^{(a)} - \nu_k$ is lognormally distributed, where $\hat{\tau}_k^{(a)}$, $a = U, C$, is an estimator of τ_k and ν_k is the number of sampled elements from U_k . Thus, a CI for τ_k is $\left(\nu_k + (\hat{\tau}_k^{(a)} - \nu_k)/c_k, \nu_k + (\hat{\tau}_k^{(a)} - \nu_k) \times c_k \right)$, where $c_k = \exp \left\{ z_{\alpha/2} \sqrt{\ln[1 + \hat{V}(\hat{\tau}_k^{(a)})/(\hat{\tau}_k^{(a)} - \nu_k)^2]} \right\}$, $z_{\alpha/2}$ is the upper $\alpha/2$ point of the standard normal distribution and $\hat{V}(\hat{\tau}_k^{(a)})$ is an estimate of the variance of $\hat{\tau}_k^{(a)}$. (See Williams et al., 2002, Sec. 14.2, for a description of this type of CI.) A CI for τ is built analogously. The values of ν_1 , ν_2 and ν that are used in the CIs for τ_1 , τ_2 and τ are $m + r_1$, r_2 and $m + r_1 + r_2$, respectively. (ii) If the parameter is \bar{Y}_k , $k = 1, 2$, or \bar{Y} , and it is a proportion, that is, the y -value associated with an element is equal to one if the element has a characteristic of interest and is equal to zero otherwise, then we will assume that the number of sampled elements with the characteristic of interest has a binomial distribution and a CI for \bar{Y} will be constructed using the proposal of Korn and Graubard (1998), which is an adaptation of the Clopper-Pearson CI for a proportion in the case of complex samples. Thus, a CI for \bar{Y}_k is $\left(\nu_{(k)1}^{(a)} F_{\nu_{(k)1}^{(a)}, \nu_{(k)2}^{(a)}}(\alpha/2) [\nu_{(k)2}^{(a)} + \nu_{(k)1}^{(a)} F_{\nu_{(k)1}^{(a)}, \nu_{(k)2}^{(a)}}(\alpha/2)], \nu_{(k)3}^{(a)} F_{\nu_{(k)3}^{(a)}, \nu_{(k)4}^{(a)}}(1 - \alpha/2) [\nu_{(k)4}^{(a)} + \nu_{(k)3}^{(a)} F_{\nu_{(k)3}^{(a)}, \nu_{(k)4}^{(a)}}(1 - \alpha/2)] \right)$, where $\nu_{(k)1}^{(a)} = 2y_k^{(a)}$, $\nu_{(k)2}^{(a)} = 2(n_k^{(a)} - y_k^{(a)} + 1)$, $\nu_{(k)3}^{(a)} = 2(y_k^{(a)} + 1)$, $\nu_{(k)4}^{(a)} = 2(n_k^{(a)} - y_k^{(a)})$, $y_k^{(a)} = n_k^{(a)} \hat{Y}_k^{(a)}$, $n_k^{(a)} = \hat{Y}_k^{(a)}(1 - \hat{Y}_k^{(a)})/\hat{V}(\hat{Y}_k^{(a)})$, $\hat{Y}_k^{(a)}$ is an estimator of \bar{Y}_k , $\hat{V}(\hat{Y}_k^{(a)})$ is an estimate of the variance of $\hat{Y}_k^{(a)}$ and $F_{d_1, d_2}(\beta)$ is the β quantile of the F distribution with d_1 and d_2 degrees of freedom. A CI for \bar{Y} is built analogously. (iii) If the parameter is \bar{Y}_k , $k = 1, 2$, or \bar{Y} , and it is the mean of the y -values a continuous variable of interest or if it is Y_k , $k = 1, 2$, or Y , that is, it is the total of the y -values of a continuous or a binary variable of interest, we will assume that the estimator $\hat{Y}_k^{(a)}$ (or $\hat{Y}_k^{(a)}$) is normally

distributed. Thus a CI for \bar{Y}_k is $\left(\hat{Y}_k^{(a)} - z_{\alpha/2} \sqrt{\hat{V}(\hat{Y}_k^{(a)})}, \hat{Y}_k^{(a)} + z_{\alpha/2} \sqrt{\hat{V}(\hat{Y}_k^{(a)})} \right)$, where $z_{\alpha/2}$ is the upper $\alpha/2$ point of the standard normal distribution and $\hat{V}(\hat{Y}_k^{(a)})$ is an estimate of the variance of $\hat{Y}_k^{(a)}$. CIs for Y_k , \bar{Y} and Y are built analogously.

6 Monte Carlo studies

In order to observe the performance of the proposed estimators and CIs and to compare their performance with the ones proposed by Félix-Medina and Monjardin (2010), which were derived under the assumption of homogeneity of the link probabilities, we carried out two numerical studies. In the first study we used data from the National Longitudinal Study of Adolescent Health collected during the 1994-1995 school year to construct a population, whereas in the second one we used artificial data to construct two populations with specific characteristics. Both studies were carried out using the R software environment for statistical computing (R Core Team, 2018).

6.1 Populations constructed using data from the National Longitudinal Study of Adolescent Health

In this Monte Carlo Study we used data from the National Longitudinal Study of Adolescent Health (Add Health) to construct a population. The Add Health is a longitudinal study of a representative sample of more than 90000 adolescents who in the years 1994-95 were in grades 7-12 in the United States. The participants were followed through adolescence and the transition to adulthood with the goal of helping to explain the causes of adolescent health and health behavior. The sample of students was selected by a stratified probability proportional to size cluster sampling design, where the clusters were the high schools and the strata were defined in terms of region, urbanicity, school type, etc. For each of the 84 selected high schools, one of its feeder middle school was selected with probability proportional to the number of contributed students to the high school. Each student in the representative sample was asked to named up to 5 male and 5 female friends within his or her high school or in the feeder school, and in addition, to complete an in-school questionnaire. Thus, the collected information can be modeled as a directed network, where the nodes are the sampled students and their referred friends, and a directed arc from node i to node j is considered to exist if student i names student j as a friend. See Harris (2013) for a description of this study.

A subset of the data obtained in the Add Health study is contained in Linton Freeman's web page: <http://moreno.ss.uci.edu/data.html#ahealth>. In our numerical study we used data from this subset corresponding to the high school and its feeder in Community 50 to construct a population U of $\tau = 2497$ elements divided into subpopulations U_1 and U_2 of sizes $\tau_1 = 1800$ and $\tau_2 = 697$, respectively. The elements assigned to U_1 were those at positions labeled with odd numbers in the data file and that named at least one friend plus a simple random sample of elements at positions labeled with even numbers and that named at least one friend. These elements were grouped into $N = 150$ clusters of sizes m_i , $i = 1, \dots, N$, obtained

by sampling from a negative binomial distribution with mean and variance equal to 12 and 24, respectively. The elements assigned to U_2 were the remaining elements in the data file that were named as a friend by at least one element in U_1 . Once the subpopulation U_k was constructed, the $N \times \tau_k$ matrix \mathbf{X}_k of values $x_{ij}^{(k)}$ s of the link indicator variables $X_{ij}^{(k)}$ s was constructed, $k = 1, 2$. We considered as response variables the following: “Number of friends” (named by each element) and “Gender” (1=male, 0=female). The totals and means of the variable “Number of friends” were $(Y_1, Y_2, Y) = (10101, 2729, 12830)$ and $(\bar{Y}_1, \bar{Y}_2, \bar{Y}) = (5.612, 3.915, 5.138)$, and those of the variable “Gender” were $(Y_1, Y_2, Y) = (838, 361, 1200)$ and $(\bar{Y}_1, \bar{Y}_2, \bar{Y}) = (0.466, 0.518, 0.481)$. For an initial sample size $n = 20$, which was the size used in this study, the values of the Pearson correlation coefficients between the values of the variable “Number of friends” and those of the inclusion probabilities associated with subpopulations U_1 and U_2 were $\rho(y^{(1)}, \pi_{(1)}) \approx 0.36$ and $\rho(y^{(2)}, \pi_{(2)}) \approx 0.29$, respectively, whereas the corresponding values for the variable “Gender” were $\rho(y^{(1)}, \pi_{(1)}) \approx -0.04$ and $\rho(y^{(2)}, \pi_{(2)}) \approx -0.07$.

The Monte Carlo study was carried out by repeatedly selecting r samples from the population U using the sampling design described in Section 2. Thus, a SRSWOR of $n = 20$ values m_i was selected from the population of $N = 150$ values. For each selected value m_i , the values $x_{ij}^{(k)}$ s of the link indicator variables $X_{ij}^{(k)}$, $j = 1, \dots, \tau_k$, were obtained from the matrix \mathbf{X}_k , $k = 1, 2$. Furthermore, for each element $j \in U_k$ that was sampled, its associated values $y_j^{(k)}$ s of both response variables were recorded. From each selected sample, the following estimators of the population sizes τ_1 , τ_2 and τ were computed: the UMLEs and CMLEs $\hat{\tau}_1^{(a)}$, $\hat{\tau}_2^{(a)}$ and $\hat{\tau}^{(a)}$, $a = U, C$, proposed by Félix-Medina et al. (2015); both types of HTLEs $\hat{\tau}_{HT,1}^{(a)}$, $\hat{\tau}_{HT,2}^{(a)}$ and $\hat{\tau}_{HT}^{(a)}$, proposed in this work: the ones based on the UMLEs ($a = U$) and those based on the CMLEs ($a = C$) of the inclusion probabilities; the MLEs $\hat{\tau}_{ML,1}^{(H)}$, $\hat{\tau}_{ML,2}^{(H)}$ and $\hat{\tau}_{ML}^{(H)}$ proposed by Félix-Medina and Thompson (2004) and derived under the assumption of homogeneity of the link probabilities, as well as the Bayesian-assisted estimators $\hat{\tau}_{BA,1}^{(H)}$, $\hat{\tau}_{BA,2}^{(H)}$ and $\hat{\tau}_{BA}^{(H)}$ proposed by Félix-Medina and Monjardin (2006) and derived also under the homogeneity assumption. We also computed the following estimators of the population totals Y_1 , Y_2 and Y , and means \bar{Y}_1 , \bar{Y}_2 and \bar{Y} : the two types of HTLEs $\hat{Y}_{HT,1}^{(a)}$, $\hat{Y}_{HT,2}^{(a)}$ and $\hat{Y}_{HT}^{(a)}$ and $\hat{Y}_{HT,1}^{(a)}$, $\hat{Y}_{HT,2}^{(a)}$ and $\hat{Y}_{HT}^{(a)}$, and the two types of HKLEs $\hat{Y}_{HK,1}^{(a)}$, $\hat{Y}_{HK,2}^{(a)}$ and $\hat{Y}_{HK}^{(a)}$ and $\hat{Y}_{HK,1}^{(a)}$, $\hat{Y}_{HK,2}^{(a)}$ and $\hat{Y}_{HK}^{(a)}$ proposed in this work. One type was based on the UMLEs ($a = U$) and the other on the CMLEs ($a = C$) of the inclusion probabilities. Finally, the following two types of HTLEs of the population totals and means proposed by Félix-Medina and Monjardin (2010) and derived under the assumption of homogeneity of the link probabilities were computed: $\hat{Y}_{a,1}^{(H)}$, $\hat{Y}_{a,2}^{(H)}$ and $\hat{Y}_a^{(H)}$ and $\hat{Y}_{a,1}^{(H)}$, $\hat{Y}_{a,2}^{(H)}$ and $\hat{Y}_a^{(H)}$. Here, one type was based on the MLEs ($a = ML$) and the other on the Bayesian-assisted estimators ($a = BA$) of the inclusion probabilities.

The performance of an estimator $\hat{\theta}$ of a parameter θ was evaluated by its relative bias (r-bias), the square root of its relative mean square error ($\sqrt{\text{r-mse}}$), the median of its relative estimation error (mdre), and the median of its absolute relative estimation error (mdare) defined as $\text{r-bias} = \sum_1^r (\hat{\theta}_i - \theta) / (r\theta)$, $\sqrt{\text{r-mse}} = \sqrt{\sum_1^r (\hat{\theta}_i - \theta)^2 / (r\theta^2)}$, $\text{mdre} = \text{median}\{(\hat{\theta}_i - \theta) / \theta\}$

and $\text{mdare} = \text{median}\{ |(\hat{\theta}_i - \theta)/\theta| \}$, respectively, where $\hat{\theta}_i$, is the value of $\hat{\theta}$ obtained in the i -th sample, $i = 1, \dots, r$. In the case of the point estimators of the population sizes, totals and means their performance was evaluated using $r = 5000$ samples.

We also computed estimators of the variances of these point estimators. In the case of the estimators derived under the assumption of heterogeneous link probabilities we used the bootstrap variance estimators described in Section 5 based on $B = 50$ bootstrap samples. In the case of the estimators based on the assumption of homogeneous link probabilities we used the variance estimators described in the papers in which the estimators were presented. The performance of a variance estimator $\hat{V}(\hat{\theta})$ of the variance $V(\hat{\theta})$ of $\hat{\theta}$ was also evaluated by its r-bias, $\sqrt{r\text{-mse}}$, mdre and mdre, where $V(\hat{\theta})$ was computed by the sample variance of the $\hat{\theta}_i$, $i = 1, \dots, r$. Because of the time required to computed the bootstrap variance estimators we used $r = 500$ samples, whereas in the case of the estimators of the variances of the point estimators derived under the assumption of homogeneous link probabilities we used $r = 5000$ samples.

From each point estimator and its associated variance estimator a 95% CI was computed for the corresponding parameter. In the case of the estimators based on the assumption of heterogeneous link probabilities the CIs were computed as was described in Section 5, whereas in the case of those based on the homogeneity assumption the CIs were Wald type CIs. The performance of a CI was evaluated by its coverage probability (cp) defined as the proportion of samples in which the parameter is inside the interval, and by both its mean relative length (mrl) and median relative length (mdrl) defined as the sample mean and median of the lengths of the r intervals divided by the value of the parameter, respectively. In the case of the CI based on point estimators derived under the assumption of heterogeneous link probabilities we used $r = 500$ samples, whereas in the case of the estimators derived under the assumption of homogeneous link probabilities we used $r = 5000$ samples.

It is worth noting that, in this and in the following study, in the case of the estimators that were derived under the assumption of heterogeneous link probabilities, we only present the outcomes corresponding to the estimators based on the UMLEs of these probabilities because their performance was very similar to that of the estimators based on the CMLEs of the link probabilities. In the case of the estimators derived under the assumption of homogeneous link probabilities, we only present the outcomes corresponding to the estimators based on the MLEs of these probabilities because their performance was very similar to that of the estimators based on the Bayesian assisted estimators of the link probabilities. In addition, in the descriptions of the results of numerical studies we will use the convention that the performance of a point estimator will be considered as acceptable if both its r-bias (or mdre) and its $\sqrt{r\text{-mse}}$ (or mdare) are around or are lesser than 0.1. Similarly, that the performance of a 95% CI is acceptable if its cp is around or greater than 0.9 and its mrl (or mdrl) is around or is lesser than 0.4 ($= 4 \times 0.1$).

The results of the study on the point estimators are shown in Table 1 and in Figure 1. We can see that regardless of the type of response variable, the performance of each estimator that was derived under the heterogeneity assumption was pretty acceptable. The values of its mdre and mdare (summary statistics that are not affected by large values of the estimators) were generally small. The exceptions were the estimators $\hat{Y}_{HK.2}^{(U)}$ and $\hat{\bar{Y}}_{HK.2}^{(U)}$, which in the case of

Table 1: Simulation results obtained for the point and standard deviation estimators in a population constructed using data from the National Longitudinal Study of Adolescent Health.

Resp. variable	Point estimators								Standard deviation estimators							
	Num. of friends				Gender				Num. of friends				Gender			
Estimator	r b i a s	\sqrt{r} m s e	m d r e	m d a r e	r b i a s	\sqrt{r} m s e	m d r e	m d a r e	r b i a s	\sqrt{r} m s e	m d r e	m d a r e	r b i a s	\sqrt{r} m s e	m d r e	m d a r e
UMLEs $\hat{\tau}_1^{(U)}$	-.01	.06	-.01	.04	-.01	.06	-.01	.04	-.15	.23	-.17	.19	-.15	.23	-.17	.19
of $\hat{\tau}_2^{(U)}$.06	.25	.00	.13	.06	.25	.00	.13	.08	1.3	-.29	.43	.08	1.3	-.29	.43
sizes $\hat{\tau}^{(U)}$.01	.08	.00	.05	.01	.08	.00	.05	.05	.99	-.24	.33	.05	.99	-.24	.33
HTLEs $\hat{\tau}_{HT,1}^{(U)}$	-.04	.07	-.04	.05	-.04	.07	-.04	.05	-.15	.23	-.16	.18	-.15	.23	-.16	.18
of $\hat{\tau}_{HT,2}^{(U)}$	-.05	.15	-.07	.10	-.05	.15	-.07	.10	.13	.81	-.10	.31	.13	.81	-.10	.31
sizes $\hat{\tau}_{HT}^{(U)}$	-.05	.08	-.05	.05	-.05	.08	-.05	.05	-.04	.47	-.17	.25	-.04	.47	-.17	.25
HTLEs $\hat{Y}_{HT,1}^{(U)}$.00	.06	.01	.04	-.07	.09	-.07	.07	-.14	.22	-.16	.17	-.15	.22	-.16	.17
of $\hat{Y}_{HT,2}^{(U)}$.07	.17	.05	.09	-.06	.16	-.08	.11	.12	.75	-.10	.31	.19	.81	-.04	.29
totals $\hat{Y}_{HT}^{(U)}$.02	.06	.02	.04	-.07	.09	-.07	.07	-.07	.36	-.16	.21	-.02	.45	-.13	.23
HTLEs $\hat{\hat{Y}}_{HT,1}^{(U)}$.01	.02	.01	.02	-.06	.07	-.06	.06	-.15	.19	-.15	.16	-.03	.13	-.03	.09
of $\hat{\hat{Y}}_{HT,2}^{(U)}$.03	.10	.04	.08	-.10	.13	-.09	.09	-.14	.42	-.27	.32	.03	.42	-.10	.24
means $\hat{\hat{Y}}_{HT}^{(U)}$.01	.08	.02	.03	-.07	.06	-.07	.07	-.09	.70	-.33	.40	.10	.67	-.09	.20
HKLEs $\hat{Y}_{HK,1}^{(U)}$.04	.08	.04	.05	-.03	.08	-.03	.05	-.15	.23	-.17	.18	-.16	.23	-.18	.19
of $\hat{Y}_{HK,2}^{(U)}$.19	.33	.13	.15	.04	.24	-.01	.13	.05	1.2	-.30	.43	.12	1.3	-.26	.40
totals $\hat{Y}_{HK}^{(U)}$.08	.11	.07	.07	-.01	.09	-.02	.06	.03	.89	-.22	.30	.06	.96	-.21	.31
HKLEs $\hat{\hat{Y}}_{HK,1}^{(U)}$.05	.05	.05	.05	-.02	.04	-.02	.03	-.12	.16	-.13	.13	-.06	.14	-.06	.09
of $\hat{\hat{Y}}_{HK,2}^{(U)}$.13	.13	.13	.13	-.02	.04	-.02	.03	.01	.16	.00	.10	.31	.37	.30	.30
means $\hat{\hat{Y}}_{HK}^{(U)}$.07	.07	.07	.07	-.02	.03	-.02	.02	-.09	.29	-.15	.20	.01	.16	-.00	.10
Homo. $\hat{\tau}_1^{(H)}$	-.14	.15	-.14	.14	-.14	.15	-.14	.14	-.42	.43	-.42	.42	-.42	.43	-.42	.42
MLEs $\hat{\tau}_2^{(H)}$	-.24	.25	-.24	.24	-.24	.25	-.24	.24	-.24	.29	-.26	.26	-.24	.29	-.26	.26
sizes $\hat{\tau}^{(H)}$	-.17	.18	-.17	.17	-.17	.18	-.17	.17	-.45	.45	-.45	.45	-.45	.45	-.45	.45
Homo. $\hat{Y}_{HT,1}^{(H)}$	-.09	.11	-.09	.09	-.16	.17	-.16	.16	-.20	.21	-.19	.19	-.22	.23	-.22	.22
HTLEs $\hat{Y}_{HT,2}^{(H)}$	-.13	.16	-.14	.14	-.25	.26	-.25	.25	-.17	.23	-.19	.20	-.19	.25	-.21	.22
totals $\hat{Y}_{HT}^{(H)}$	-.10	.11	-.10	.10	-.19	.20	-.19	.19	-.28	.28	-.27	.27	-.31	.31	-.31	.31
Homo. $\hat{\hat{Y}}_{HT,1}^{(H)}$.06	.06	.06	.06	-.02	.04	-.02	.03	-.18	.19	-.18	.18	-.13	.14	-.13	.13
HTLEs $\hat{\hat{Y}}_{HT,2}^{(H)}$.14	.14	.14	.14	-.02	.04	-.02	.03	-.10	.12	-.11	.11	.12	.14	.12	.12
means $\hat{\hat{Y}}_{HT}^{(H)}$.08	.02	.08	.08	-.02	.03	-.02	.03	-.25	.26	-.26	.26	-.13	.14	-.13	.13

Notes: Results for point estimators are based on 5000 samples; those for sd estimators derived under the heterogeneity assumption are based on 500 samples and those derived under the homogeneity assumption are based on 5000 samples. Average sampling fractions are $f_1 = 0.46$ and $f_2 = 0.40$. No convergence problems were observed, except in one sample in which the point estimators of the parameters of U_2 could not be computed.




Figure 1: Boxplots for the values of the estimators of the population sizes, totals and means in the population constructed using data from the National Longitudinal Study of Adolescent Health.

the response variable “Number of friends” presented some problems of bias. The UMLEs of the population sizes were practically unbiased (in terms of the values of their mdre), whereas the HTLEs slightly underestimated the population sizes. The performance of the HTLEs of the population totals and means was better in the case of the response variable “Number of friends” than in the case of the response variable “Gender”. In the first case, the estimators presented small positive values of r-bias (or mdre), whereas in the second case they presented relatively small negative values of r-bias; however, the magnitudes of the values of their r-bias and $\sqrt{r\text{-mse}}$ were smaller in the first case than in the second one. The opposite situation happened with the HKLEs which presented better performance in the case of the variable “Gender” than in the case of the variable “Number of friends”. Notice from Figure 1 that the distributions of the estimators of the parameters corresponding to U_1 , that is, τ_1 , Y_1 and \bar{Y}_1 are approximately symmetrical; those of the parameters τ_2 and Y_2 are skewed to the right, as well as those of $\hat{\tau}^{(U)}$ and $\hat{Y}_{HK}^{(U)}$; those of the HTLEs of τ and Y are also approximately symmetrical as well as those of the HKLEs of the means, whereas those of $\hat{Y}_{HT.2}^{(U)}$ and $\hat{Y}_{HT}^{(U)}$ are skewed to the left. Notice also that each of these distributions are heavy-tailed as shown by the presence of many outlying points on the box plots.

The performance of the estimators of the population sizes and totals derived under the assumption of homogeneous link probabilities presented problems of underestimation which

produced large values of $\sqrt{r\text{-mse}}$ regardless of the type of response variable. However, the estimators of the population means performed acceptably, with the exception of the estimator $\hat{Y}_{HT.2}^{(H)}$ which presented a relatively large bias. Notice that these estimators of the population means performed pretty well in the case of the binary response variable: very small values of the r-bias and $\sqrt{r\text{-mse}}$. Finally, notice that the distributions of these estimators are approximately symmetrical and heavy-tailed.

The results of the study about the standard deviation estimations are also shown in Table 1. We can see that every one of the estimators presented problems of underestimation (in terms of their mdre). The exceptions, perhaps, were the estimators of the standard deviations of the HTLEs $\hat{Y}_{HT.1}^{(U)}$, $\hat{Y}_{HT.2}^{(U)}$ and $\hat{Y}_{HT}^{(U)}$, which in the case of the response variable “Gender” presented acceptable values of mdre.

The results on the 95% CIs are shown in Table 2. We can see that the CIs of the population sizes based on the UMLEs $\hat{\tau}_1^{(U)}$, $\hat{\tau}_2^{(U)}$ and $\hat{\tau}^{(U)}$ presented acceptable performance, although the mdrl of the CI of τ_2 was relatively large. The CIs of the population sizes based on the HTLEs $\hat{\tau}_{HT.1}^{(U)}$, $\hat{\tau}_{HT.2}^{(U)}$ and $\hat{\tau}_{HT}^{(U)}$ presented relatively low values of the cp (between 0.82 and 0.9), but acceptable values of their mdrl. The performance of the CIs of the population totals based on the HTLEs $\hat{Y}_{HT.1}^{(U)}$, $\hat{Y}_{HT.2}^{(U)}$ and $\hat{Y}_{HT}^{(U)}$ had good performance in the case of the response variable “Number of friends”: both their cp and mdrl presented acceptable values; however, in the case of the response variable “Gender” their performance was not good because their cp were somehow low: between 0.73 and 0.85. The performance of the CIs of the population means was neither good: although their mdrl were very short, in the case of the response variable “Number of friends” their cp were low (between 0.77 and 0.81), and in the case of the variable “Gender” they were very low (0.53 in the case of the CIs of \bar{Y}_1 and \bar{Y} , and 0.93 in the case of the CI of \bar{Y}_2). Nevertheless the small cp of the CIs of the population proportions, these CIs still provide valuable information about the proportions. For instance, in the case of the CI for \bar{Y}_1 , the value of the r-bias of $\hat{Y}_{HT.1}^{(U)}$ was -0.06 , and since $\bar{Y}_1 = 0.466$, it follows that the average value of $\hat{Y}_{HT.1}^{(U)}$ was $0.94 \times \bar{Y}_1 = 0.438$. In addition, since the mrl of the CI was 0.12, it follows that the average values of the lower and upper limits of the CI were 0.410 and 0.466, respectively. Consequently, from this point of view, the performance of the CI was acceptable despite its small cp. The performance of CIs of the population totals based on the HKLEs was moderately acceptable because of their relatively low cp and the relatively large mrl of the CI of Y_2 . In the case of the CIs of the population means based on this type of estimator, their performance was acceptable in the case of the variable “Gender”, but their cp were very small in the case of the variable “Number of friends”. However, these CIs still provide valuable information because the “average” CIs of $\bar{Y}_1 = 5.612$, $\bar{Y}_2 = 3.915$ and $\bar{Y} = 5.138$ were (5.75, 6.03), (3.93, 4.55) and (5.35, 5.65), respectively. Finally, in the case of the CIs constructed under the assumption of homogeneous link probabilities their cp were very low; however, their mdrl were very short, and since the values of the r-bias of the estimators of the means were small, we could also conclude that the performance of these CIs of the means was still acceptable. Notice also that in the case of the response variable “Gender”, the cp of the CIs of the means were not so low (between 0.77 and 0.96).

From the results of this study, we have that inferences based on the UMLEs of the pop-

Table 2: Simulation results obtained for the confidence intervals in a population constructed using data from the National Longitudinal Study of Adolescent Health.

Response variable		CIs based on UMLEs of sizes			CIs based on HTLEs of sizes			CIs based on HTLEs of totals			CIs based on HTLEs of means		
		$\hat{\tau}_1^{(U)}$	$\hat{\tau}_2^{(U)}$	$\hat{\tau}^{(U)}$	$\hat{\tau}_{HT.1}^{(U)}$	$\hat{\tau}_{HT.2}^{(U)}$	$\hat{\tau}_{HT}^{(U)}$	$\hat{Y}_{HT.1}^{(U)}$	$\hat{Y}_{HT.2}^{(U)}$	$\hat{Y}_{HT}^{(U)}$	$\hat{Y}_{HT.1}^{(U)}$	$\hat{Y}_{HT.2}^{(U)}$	$\hat{Y}_{HT}^{(U)}$
Number of friends	cp	.89	.95	.93	.82	.90	.82	.90	.97	.94	.81	.77	.77
	mrl	.22	1.1	.36	.20	.67	.24	.20	.68	.23	.06	.32	.15
	mdrl	.21	.68	.26	.19	.52	.21	.20	.55	.21	.06	.27	.11
Gender	cp	.89	.95	.93	.82	.90	.82	.73	.85	.74	.53	.93	.53
	mrl	.22	1.1	.36	.20	.67	.24	.22	.66	.26	.12	.33	.15
	mdrl	.21	.68	.26	.19	.52	.21	.22	.54	.23	.12	.29	.13
		CIs based on HKLEs of totals			CIs based on HKLEs of means								
		$\hat{Y}_{HK.1}^{(U)}$	$\hat{Y}_{HK.2}^{(U)}$	$\hat{Y}_{HK}^{(U)}$	$\hat{Y}_{HK.1}^{(U)}$	$\hat{Y}_{HK.2}^{(U)}$	$\hat{Y}_{HK}^{(U)}$						
Number of friends	cp	.84	.98	.88	.02	.16	.09						
	mrl	.22	1.1	.33	.05	.16	.07						
	mdrl	.22	.71	.25	.05	.16	.06						
Gender	cp	.84	.92	.88	.85	.99	.86						
	mrl	.24	1.0	.37	.11	.19	.10						
	mdrl	.23	.68	.28	.11	.19	.10						
		CIs based on homogeneous MLEs of sizes			CIs based on homogeneous HTLEs of totals			CIs based on homogeneous HTLEs of means					
		$\hat{\tau}_1^{(H)}$	$\hat{\tau}_2^{(H)}$	$\hat{\tau}^{(H)}$	$\hat{Y}_{HT.1}^{(H)}$	$\hat{Y}_{HT.2}^{(H)}$	$\hat{Y}_{HT}^{(H)}$	$\hat{Y}_{HT.1}^{(H)}$	$\hat{Y}_{HT.2}^{(H)}$	$\hat{Y}_{HT.1}^{(H)}$			
Number of friends	cp	.04	.10	.01	.42	.49	.28	.01	.05	.00			
	mrl	.11	.22	.10	.16	.28	.14	.04	.14	.05			
	mdrl	.11	.22	.10	.16	.27	.14	.04	.14	.05			
Gender	cp	.04	.10	.01	.10	.11	.02	.81	.96	.77			
	mrl	.11	.22	.10	.17	.25	.14	.10	.16	.09			
	mdrl	.11	.22	.10	.17	.25	.14	.10	.16	.08			

Notes: Results for confidence intervals derived under the heterogeneity assumption are based on 500 samples and those derived under the homogeneity assumption are based on 5000 samples. Sampling fractions are $f_1 = 0.46$ and $f_2 = 0.40$. No convergence problems were observed.

ulation totals are pretty acceptable, those based on the HTLEs of the population sizes are acceptable, but not as good as those based on the UMLEs. In the case of the response variable “Number of friends”, inferences about the population totals and means based on the HTLEs are acceptable, although the 95% CIs of the means tended to have relatively low values of the cp, and they are better than those based on the HKLEs. However, in the case of the variable “Gender” the opposite happened. This result is consequence of the higher correlation between the inclusion probabilities and the variable “Number of friends” than between the inclusion probabilities and the variable “Gender”. (See Thompson, 2002, Ch. 6.)

6.2 Populations constructed using artificial data

We constructed two populations whose characteristics are described in Table 3. The difference between the two populations is that in Population I the link probabilities were generated by using expression (1), that is, under the assumed model, whereas in Population II they were generated by the following latent-class model used by Pledger (2000) in the context of capture-recapture studies: $p_{ij} = \exp[\mu^{(k)} + \alpha_i^{(k)} + \beta_j^{(k)} + (\alpha\beta)_{ij}^{(k)}] / \{1 + \exp[\mu^{(k)} + \alpha_i^{(k)} + \beta_j^{(k)} + (\alpha\beta)_{ij}^{(k)}]\}$, $i = 1, \dots, n$; $j = 1, 2$, and $k = 1, 2$. In this model, the people in U_k are divided into two classes according to their propensities to be linked to the sample clusters. The probability that a person in U_k is in class j is $p_j^{(k)}$ and it is the same for each person in U_k . The values of the parameters that appear in each of the two expressions of the link probabilities were set so that when the size of the initial sample of clusters is $n = 15$, in both populations the sampling fractions were $f_1 = 0.5$ in U_1 and $f_2 = 0.4$ in U_2 . Notice from Table 1 that associated with each element of each population there are two values of two response variables. One variable is a continuous variable whose value associated with the j -th element of U_k was obtained by sampling from a non-central chi-square distribution with two degrees of freedom and non-centrality parameter $\psi_j^{(k)}$. The other variable is a binary variable whose value associated with that element was obtained from a Bernoulli distribution with mean $\phi_j^{(k)}$. The values of the parameters that appear in the expressions of $\psi_j^{(k)}$ and $\phi_j^{(k)}$ were set so that the values of the population means of the continuous variable in both populations were $\bar{Y}_1 \approx 50$ and $\bar{Y}_2 \approx 40$, whereas the corresponding values of the binary variable were $\bar{Y}_1 \approx 0.3$ and $\bar{Y}_2 \approx 0.2$. Furthermore, they were set so that for $n = 15$, in Population I the values of the Pearson correlation coefficients between the values of the continuous response variable and those of the inclusion probabilities were $\rho(y^{(1)}, \pi_{(1)}) \approx 0.8$ and $\rho(y^{(2)}, \pi_{(2)}) \approx 0.7$, whereas the corresponding values for the binary response variable were $\rho(y^{(1)}, \pi_{(1)}) \approx 0.3$ and $\rho(y_j^{(2)}, \pi_{(2)}) \approx 0.27$. In the case of Population II and continuous response variable those values were $\rho(y^{(1)}, \pi_{(1)}) \approx 0.15$ and $\rho(y^{(2)}, \pi_{(2)}) \approx 0.1$, whereas the corresponding values for the binary variable were $\rho(y^{(1)}, \pi_{(1)}) \approx 0$ and $\rho(y^{(2)}, \pi_{(2)}) \approx 0.1$.

The Monte Carlo study was carried out as in the previous study, except that the size of the initial sample of clusters was $n = 15$ and that for each selected value m_i , the value $x_{ij}^{(k)}$ of the link indicator variable $X_{ij}^{(k)}$ was generated from the Bernoulli distribution with mean $p_{ij}^{(k)}$ (see its expression in Table 3).

Table 3: Parameters of simulated populations.

Population I	Population II
$N = 150$ $M_i \sim \text{zero trunc. neg. binom. distribution}$ $E(M_i) = 8, \quad V(M_i) = 24$ $\tau_1 = 1208, \tau_2 = 400, \tau = 1608$ $p_{ij}^{(k)} = \frac{\exp(\alpha_i^{(k)} + \beta_j^{(k)})}{1 + \exp(\alpha_i^{(k)} + \beta_j^{(k)})}$ $i = 1, \dots, N, j = 1, \dots, \tau_k, k = 1, 2$ $\alpha_i^{(k)} = \frac{c_k}{0.001 + M_i^{1/4}}, i = 1, \dots, N, k = 1, 2$ $c_1 = -5.45, \quad c_2 = -5.85$ $\beta_j^{(k)} \sim N(0, 1), j = 1, \dots, \tau_k$	$N = 150$ $M_i \sim \text{zero trunc. neg. binom. distribution}$ $E(M_i) = 8, \quad V(M_i) = 24$ $\tau_1 = 1208, \tau_2 = 400, \tau = 1608$ $p_{ij}^{(k)} = \frac{\exp(\mu^{(k)} + \alpha_i^{(k)} + \beta_j^{(k)} + (\alpha\beta)_{ij}^{(k)})}{1 + \exp(\mu^{(k)} + \alpha_i^{(k)} + \beta_j^{(k)} + (\alpha\beta)_{ij}^{(k)})}$ $i = 1, \dots, N, j = 1, 2, k = 1, 2$ $\mu^{(1)} = 0.25, \quad \mu^{(2)} = 0.05$ $\alpha_i^{(k)} = \frac{c_k}{0.001 + M_i^{1/4}}, i = 1, \dots, N, k = 1, 2$ $c_1 = -12, \quad c_2 = -12$ $\beta_1^{(k)} = 1.5, \beta_2^{(k)} = 0, k = 1, 2$ $(\alpha\beta)_{i1}^{(k)} \sim N(0, 1.25^2), (\alpha\beta)_{i2}^{(k)} = 0, k = 1, 2$ $i = 1, \dots, N$ $p_1^{(k)} = 0.3, p_2^{(k)} = 0.7, k = 1, 2$
<p>Continuous response variable:</p> $Y_j^{(k)} \sim \chi_2^2(\psi_j^{(k)}), j = 1, \dots, \tau_k, k = 1, 2$ $\psi_j^{(k)} = 5 + \frac{d_k \exp(\beta_j^{(k)})}{1 + \exp(\beta_j^{(k)})}, j = 1, \dots, \tau_k, k = 1, 2$ $d_1 = 87, \quad d_2 = 65$ $\rho(y^{(1)}, \pi_{(1)}) = 0.79, \quad \rho(y^{(2)}, \pi_{(2)}) = 0.72$ $Y_1 = 60390.34, Y_2 = 15945.89, Y = 76336.22$ $\bar{Y}_1 = 49.99, \bar{Y}_2 = 39.87, \bar{Y} = 47.47$	<p>Continuous response variable:</p> $Y_j^{(k)} \sim \chi_2^2(\psi_j^{(k)}), j = 1, \dots, \tau_k, k = 1, 2$ $\psi_j^{(k)} = 5 + \frac{d_k \exp(\mu_k + \beta_j^{(k)})}{1 + \exp(\mu_k + \beta_j^{(k)})}, j = 1, \dots, \tau_k, k = 1, 2$ $d_1 = 65.05, \quad d_2 = 50.05$ $\rho(y^{(1)}, \pi_{(1)}) = 0.16, \quad \rho(y^{(2)}, \pi_{(2)}) = 0.11$ $Y_1 = 60289.03, Y_2 = 16112.78, Y = 76401.81$ $\bar{Y}_1 = 49.91, \bar{Y}_2 = 40.28, \bar{Y} = 47.51$
<p>Binary response variable:</p> $Y_j^{(k)} \sim \text{Bernoulli}(\phi_j^{(k)}), j = 1, \dots, \tau_k, k = 1, 2$ $\phi_j^{(k)} = \frac{g_k \exp(\beta_j^{(k)})}{1 + \exp(\beta_j^{(k)})}, j = 1, \dots, \tau_k, k = 1, 2$ $g_1 = 0.6, \quad g_2 = 0.39$ $\rho(y^{(1)}, \pi_{(1)}) = 0.29, \quad \rho(y^{(2)}, \pi_{(2)}) = 0.27$ $Y_1 = 366, Y_2 = 82, Y = 448$ $\bar{Y}_1 = 0.303, \bar{Y}_2 = 0.205, \bar{Y} = 0.279$	<p>Binary response variable:</p> $Y_j^{(k)} \sim \text{Bernoulli}(\phi_j^{(k)}), j = 1, \dots, \tau_k, k = 1, 2$ $\phi_j^{(k)} = \frac{g_k \exp(\mu_k + \beta_j^{(k)})}{1 + \exp(\mu_k + \beta_j^{(k)})}, j = 1, \dots, \tau_k, k = 1, 2$ $g_1 = 0.46, \quad g_2 = 0.33$ $\rho(y^{(1)}, \pi_{(1)}) = -0.01, \quad \rho(y^{(2)}, \pi_{(2)}) = 0.09$ $Y_1 = 365, Y_2 = 79, Y = 444$ $\bar{Y}_1 = 0.302, \bar{Y}_2 = 0.198, \bar{Y} = 0.276$

Table 4: Relative biases, square roots of relative mean square errors and medians of relative errors and absolute relative errors of the estimators of the population sizes, totals and means.

Population	I								II							
Sampling rates	$f_1 = 0.5$				$f_2 = 0.4$				$f_1 = 0.5$				$f_2 = 0.4$			
Resp. variable	Continuous				Binary				Continuous				Binary			
Estimator	r b i a s	\sqrt{r} m s e	m d r e	m d a r e	r b i a s	\sqrt{r} m s e	m d r e	m d a r e	r b i a s	\sqrt{r} m s e	m d r e	m d a r e	r b i a s	\sqrt{r} m s e	m d r e	m d a r e
UMLEs $\hat{\tau}_1^{(U)}$	-0.00	.08	-.01	.05	-0.00	.08	-.01	.05	-0.09	.13	-.08	.09	-0.09	.13	-.08	.09
of $\hat{\tau}_2^{(U)}$.06	.37	-.01	.16	.06	.37	-.01	.16	-.16	1.9	-.27	.28	-.16	1.9	-.27	.28
sizes $\hat{\tau}^{(U)}$.01	.11	.01	.06	.01	.11	.01	.06	-.10	.48	-.12	.12	-.10	.48	-.12	.12
HTLEs $\hat{\tau}_{HT.1}^{(U)}$	-0.11	.13	-.11	.11	-0.11	.13	-.11	.11	-.15	.18	-.15	.15	-.15	.18	-.15	.15
of $\hat{\tau}_{HT.2}^{(U)}$	-.19	.24	-.21	.21	-.19	.24	-.21	.21	-.28	.35	-.32	.32	-.28	.35	-.32	.32
sizes $\hat{\tau}_{HT}^{(U)}$	-.13	.14	-.13	.13	-.13	.14	-.13	.13	-.18	.20	-.18	.18	-.18	.20	-.18	.18
HTLEs $\hat{Y}_{HT.1}^{(U)}$	-0.00	.06	-.00	.04	.01	.07	.01	.05	-.14	.17	-.14	.14	-.16	.19	-.16	.16
of $\hat{Y}_{HT.2}^{(U)}$	-.06	.17	-.08	.12	.03	.19	.01	.11	-.27	.34	-.30	.31	-.21	.32	-.25	.26
totals $\hat{Y}_{HT}^{(U)}$	-.01	.06	-.01	.04	.02	.07	.02	.05	-.17	.18	-.17	.17	-.16	.19	-.17	.17
HTLEs $\hat{\hat{Y}}_{HT.1}^{(U)}$.00	.03	.00	.02	.02	.05	.02	.03	-.06	.07	-.07	.07	-.08	.09	-.08	.08
of $\hat{\hat{Y}}_{HT.2}^{(U)}$	-.08	.14	-.07	.08	.01	.16	.01	.10	-.06	.10	-.05	.05	.02	.14	.03	.09
means $\hat{\hat{Y}}_{HT}^{(U)}$	-.02	.08	-.01	.03	.01	.09	.01	.04	-.06	.11	-.06	.06	-.06	.15	-.06	.06
HKLEs $\hat{Y}_{HK.1}^{(U)}$.11	.14	.11	.11	.13	.16	.13	.13	-.07	.12	-.07	.08	-.09	.14	-.09	.10
of $\hat{Y}_{HK.2}^{(U)}$.24	.47	.15	.18	.35	.57	.25	.25	-.15	1.9	-.26	.27	-.06	2.7	-.20	.22
totals $\hat{Y}_{HK}^{(U)}$.15	.18	.14	.14	.18	.21	.17	.17	-.08	.45	-.09	.10	-.08	.49	-.09	.10
HKLEs $\hat{\hat{Y}}_{HK.1}^{(U)}$.12	.12	.12	.12	.14	.14	.14	.14	.02	.02	.02	.02	-.01	.04	-.01	.03
of $\hat{\hat{Y}}_{HK.2}^{(U)}$.17	.17	.17	.17	.27	.30	.27	.27	.02	.03	.02	.02	.11	.16	.11	.11
means $\hat{\hat{Y}}_{HK}^{(U)}$.13	.13	.13	.13	.17	.17	.17	.17	.02	.03	.02	.02	.03	.05	.03	.03
Homo. $\hat{\tau}_1^{(H)}$	-.31	.31	-.31	.31	-.31	.31	-.31	.31	-.25	.27	-.26	.26	-.25	.27	-.26	.26
MLEs $\hat{\tau}_2^{(H)}$	-.40	.41	-.40	.40	-.40	.41	-.40	.40	-.35	.36	-.37	.37	-.35	.36	-.37	.37
sizes $\hat{\tau}^{(H)}$	-.33	.33	-.33	.33	-.33	.33	-.33	.33	-.27	.29	-.28	.28	-.27	.29	-.28	.28
Homo. $\hat{Y}_{HT.1}^{(H)}$	-.19	.19	-.19	.19	-.17	.17	-.17	.17	-.24	.25	-.24	.24	-.26	.28	-.26	.26
HTLEs $\hat{Y}_{HT.2}^{(H)}$	-.29	.29	-.29	.29	-.21	.23	-.21	.21	-.33	.35	-.35	.35	-.27	.31	-.29	.30
totals $\hat{Y}_{HT}^{(H)}$	-.21	.21	-.21	.21	-.18	.18	-.18	.18	-.26	.27	-.26	.26	-.26	.27	-.26	.26
Homo. $\hat{\hat{Y}}_{HT.1}^{(H)}$.17	.17	.17	.17	.20	.21	.20	.20	.02	.02	.02	.02	-.01	.04	-.01	.03
HTLEs $\hat{\hat{Y}}_{HT.2}^{(H)}$.20	.20	.20	.20	.33	.35	.32	.32	.02	.03	.02	.02	.12	.16	.12	.12
means $\hat{\hat{Y}}_{HT}^{(H)}$.18	.12	.18	.18	.24	.14	.23	.23	.03	.03	.03	.03	.02	.08	.02	.03

Notes: Results are based on 5000 samples. In Population I the percentages of samples in which the estimators derived under the heterogeneity assumption of the population parameters associated with U_1 , U_2 and U were not computed because of numerical convergence problems were 0%, 0.02% and 0.02%, respectively, whereas in Population II the corresponding percentages were 0.36%, 8.7% and 9.0%. In the computation of the estimators derived under the homogeneity assumption no convergence problems were presented.




Figure 2: Boxplots for the values of the estimators of the population sizes, totals and means in Populations I and II.

The results of the study on the point estimators are shown in Table 4 and in Figure 2. We can see that in the case of Population I, the performance of the UMLEs of the population sizes was acceptable: they did not show problems of bias and only the estimator $\hat{\tau}_2^{(U)}$ presented problems of instability. The distribution of the estimator $\hat{\tau}_1^{(U)}$ was symmetrical, whereas those of the estimators $\hat{\tau}_2^{(U)}$ and $\hat{\tau}^{(U)}$ were skewed to the right with long tails. For that reason the values of their $\sqrt{r\text{-mse}}$ were greater than those of their mdare. On the other hand, the performance of the HTLEs $\hat{\tau}_{HT.1}^{(U)}$, $\hat{\tau}_{HT.2}^{(U)}$ and $\hat{\tau}_{HT}^{(U)}$ was not good because they presented relatively large negative biases. The performance of the HTLEs $\hat{Y}_{HT.1}^{(U)}$, $\hat{Y}_{HT.2}^{(U)}$ and $\hat{Y}_{HT}^{(U)}$ of the population totals and that of the HTLEs $\hat{\hat{Y}}_{HT.1}^{(U)}$, $\hat{\hat{Y}}_{HT.2}^{(U)}$ and $\hat{\hat{Y}}_{HT}^{(U)}$ of the population means was acceptable in both cases continuous and binary response variables: they did not present problems of bias nor problems of instability. It is worth noting that in the case of the continuous response variable the distributions of $\hat{Y}_{HT.1}^{(U)}$ and $\hat{Y}_{HT}^{(U)}$ were relatively symmetrical; those of $\hat{Y}_{HT.2}^{(U)}$ and $\hat{\hat{Y}}_{HT.1}^{(U)}$ were somewhat skewed to the right, and those of $\hat{\hat{Y}}_{HT.2}^{(U)}$ and $\hat{\hat{Y}}_{HT}^{(U)}$ were skewed to the left, whereas in the case of the binary response variable the distribution of every one of the estimators was symmetrical, except those of $\hat{Y}_{HT.2}^{(U)}$ and $\hat{\hat{Y}}_{HT}^{(U)}$ which were skewed to the right and to the left, respectively. The performance of the HKLEs of the population totals and means was not good because they presented relatively large positive biases. Finally, the performance of the estimators derived under the assumption of homogeneous link probabilities was not good: the estimators of the population sizes and population totals presented serious problems of underestimation, whereas those of the population means serious problems of overestimation.

In the case of the Population II, the performance of the UMLEs $\hat{\tau}_1^{(U)}$ and $\hat{\tau}^{(U)}$ was still acceptable, but less good than that observed in Population I. The performance of $\hat{\tau}_2^{(U)}$ was not good: it presented a large negative bias and a large variability. Notice that the distribution of $\hat{\tau}_1^{(U)}$ was symmetrical and those of $\hat{\tau}_2^{(U)}$ and $\hat{\tau}^{(U)}$ were skewed to the right with long tails. The performance of the HTLEs $\hat{\tau}_{HT.1}^{(U)}$, $\hat{\tau}_{HT.2}^{(U)}$ and $\hat{\tau}_{HT}^{(U)}$ was not good: their biases were large and caused large values of their $\sqrt{r\text{-mse}}$; in fact, the values of both r-bias and $\sqrt{r\text{-mse}}$ were greater than those observed in Population I. The shape of their distributions were similar to those of the corresponding estimators $\hat{\tau}_1^{(U)}$, $\hat{\tau}_2^{(U)}$ and $\hat{\tau}^{(U)}$. The performance of the HTLEs $\hat{Y}_{HT.1}^{(U)}$, $\hat{Y}_{HT.2}^{(U)}$ and $\hat{Y}_{HT}^{(U)}$ of the population totals was not good in both cases continuous and binary response variables. It is worth noting that their distributions were similar to those of the HTLEs $\hat{\tau}_{HT.1}^{(U)}$, $\hat{\tau}_{HT.2}^{(U)}$ and $\hat{\tau}_{HT}^{(U)}$. The performance of the HTLEs $\hat{\hat{Y}}_{HT.1}^{(U)}$, $\hat{\hat{Y}}_{HT.2}^{(U)}$ and $\hat{\hat{Y}}_{HT}^{(U)}$ of the population means was acceptable: their biases and variances were not large. In the case of the continuous response variable the distribution of $\hat{\hat{Y}}_{HT.1}^{(U)}$ was skewed to the right, whereas those of $\hat{\hat{Y}}_{HT.2}^{(U)}$ and $\hat{\hat{Y}}_{HT}^{(U)}$ were skewed to the left. In the case of the binary response variable, the distribution of $\hat{\hat{Y}}_{HT.1}^{(U)}$ was skewed to the right, that of $\hat{\hat{Y}}_{HT.2}^{(U)}$ was more or less symmetrical, and that of $\hat{\hat{Y}}_{HT}^{(U)}$ was skewed to the left. The performance of the HKLEs of the population totals and means was pretty acceptable in both cases continuous and binary response variables, except that of the estimator of Y_2 which presented problems of subestimation. The distributions of $\hat{Y}_{HK.1}^{(U)}$, $\hat{\hat{Y}}_{HK.1}^{(U)}$ and $\hat{\hat{Y}}_{HK.2}^{(U)}$ were symmetrical, those of $\hat{Y}_{HK.2}^{(U)}$ and

Table 5: Relative biases, square roots of relative mean square errors and medians of relative errors and absolute relative errors of the deviation standard estimators of the estimators of the population sizes, totals and means.

Population	I								II							
Sampling rates	$f_1 = 0.5$				$f_2 = 0.4$				$f_1 = 0.5$				$f_2 = 0.4$			
Response variable	Continuous				Binary				Continuous				Binary			
Deviation standard estimator	r b i a s	$\sqrt{\frac{r}{m}}$ s e	m d r e	m d a r e	r b i a s	$\sqrt{\frac{r}{m}}$ s e	m d r e	m d a r e	r b i a s	$\sqrt{\frac{r}{m}}$ s e	m d r e	m d a r e	r b i a s	$\sqrt{\frac{r}{m}}$ s e	m d r e	m d a r e
UMLEs $\widehat{sd}_B(\hat{\tau}_1^{(U)})$.15	.29	.12	.16	.15	.29	.12	.16	-.19	.27	-.20	.21	-.19	.27	-.20	.21
of $\widehat{sd}_B(\hat{\tau}_2^{(U)})$	2.0	6.7	.22	.58	2.0	6.7	.22	.58	.26	10.5	-.59	.64	.26	10.5	-.59	.64
sizes $\widehat{sd}_B(\hat{\tau}^{(U)})$	1.8	5.6	.35	.39	1.8	5.6	.35	.39	.23	7.3	-.30	.33	.23	7.3	-.30	.33
HTLEs $\widehat{sd}_B(\hat{\tau}_{HT,1}^{(U)})$.15	.29	.11	.15	.15	.29	.11	.15	-.30	.35	-.31	.31	-.30	.35	-.31	.31
of $\widehat{sd}_B(\hat{\tau}_{HT,2}^{(U)})$.85	1.6	.42	.48	.85	1.6	.42	.48	-.11	1.9	-.48	.55	-.11	1.9	-.48	.55
sizes $\widehat{sd}_B(\hat{\tau}_{HT}^{(U)})$.55	.90	.36	.36	.55	.90	.36	.36	-.18	.94	-.32	.35	-.18	.94	-.32	.35
HTLEs $\widehat{sd}_B(\hat{Y}_{HT,1}^{(U)})$.20	.32	.16	.19	.12	.26	.10	.15	-.29	.35	-.30	.30	-.23	.30	-.24	.25
of $\widehat{sd}_B(\hat{Y}_{HT,2}^{(U)})$.78	1.5	.33	.45	.63	1.3	.26	.41	-.09	1.9	-.47	.54	.02	1.6	-.35	.44
totals $\widehat{sd}_B(\hat{Y}_{HT}^{(U)})$.57	.89	.40	.40	.46	.72	.34	.34	-.18	.82	-.31	.34	-.13	.59	-.22	.26
HTLEs $\widehat{sd}_B(\hat{Y}_{HT,1}^{(U)})$.01	.18	-.01	.11	.16	.23	.15	.16	-.50	.54	-.51	.51	-.11	.27	-.13	.17
of $\widehat{sd}_B(\hat{Y}_{HT,2}^{(U)})$.35	.65	.22	.35	.24	.44	.17	.21	-.20	.67	-.38	.44	.06	.50	-.06	.24
means $\widehat{sd}_B(\hat{Y}_{HT}^{(U)})$.84	1.8	.20	.43	.65	1.3	.19	.28	-.35	.73	-.50	.52	-.02	.60	-.16	.22
HKLEs $\widehat{sd}_B(\hat{Y}_{HK,1}^{(U)})$.20	.32	.16	.19	.12	.26	.09	.15	-.18	.27	-.19	.21	-.16	.26	-.16	.19
of $\widehat{sd}_B(\hat{Y}_{HK,2}^{(U)})$	1.9	6.3	.14	.56	1.6	5.9	.09	.52	.26	10.3	-.59	.63	.32	9.6	-.50	.57
totals $\widehat{sd}_B(\hat{Y}_{HK}^{(U)})$	1.9	5.6	.34	.37	1.7	5.2	.33	.37	.22	6.7	-.27	.30	.18	5.2	-.21	.26
HKLEs $\widehat{sd}_B(\hat{Y}_{HK,1}^{(U)})$	-.12	.18	-.13	.14	.11	.18	.10	.12	-.10	.20	-.12	.15	.05	.23	.04	.15
of $\widehat{sd}_B(\hat{Y}_{HK,2}^{(U)})$.09	.23	.06	.15	.11	.25	.09	.15	.07	.35	.00	.20	.03	.33	-.03	.20
means $\widehat{sd}_B(\hat{Y}_{HK}^{(U)})$.06	.30	-.01	.14	.16	.29	.13	.14	-.13	.47	-.24	.29	.08	.32	.06	.16
Homo. $\widehat{sd}(\hat{\tau}_1^{(H)})$	-.42	.42	-.42	.42	-.42	.42	-.42	.42	-.74	.75	-.75	.75	-.74	.75	-.75	.75
MLEs $\widehat{sd}(\hat{\tau}_2^{(H)})$	-.20	.26	-.22	.23	-.20	.26	-.22	.23	-.48	.66	-.59	.60	-.48	.66	-.59	.60
sizes $\widehat{sd}(\hat{\tau}^{(H)})$	-.41	.42	-.42	.42	-.41	.42	-.42	.42	-.69	.71	-.71	.71	-.69	.71	-.71	.71
Homo. $\widehat{sd}(\hat{Y}_{HT,1}^{(H)})$	-.24	.25	-.24	.24	-.18	.19	-.18	.18	-.69	.71	-.72	.72	-.62	.64	-.65	.65
HTLEs $\widehat{sd}(\hat{Y}_{HT,2}^{(H)})$	-.11	.21	-.14	.17	-.08	.19	-.10	.14	-.47	.65	-.58	.59	-.31	.54	-.42	.45
totals $\widehat{sd}(\hat{Y}_{HT}^{(H)})$	-.25	.27	-.26	.26	-.17	.19	-.18	.18	-.66	.68	-.69	.69	-.58	.60	-.61	.61
Homo. $\widehat{sd}(\hat{Y}_{HT,1}^{(H)})$	-.38	.39	-.39	.39	-.16	.18	-.16	.16	-.21	.29	-.23	.24	-.09	.25	-.11	.18
HTLEs $\widehat{sd}(\hat{Y}_{HT,2}^{(H)})$	-.19	.22	-.20	.20	-.11	.15	-.12	.13	-.12	.29	-.16	.22	-.12	.29	-.15	.22
means $\widehat{sd}(\hat{Y}_{HT}^{(H)})$	-.37	.38	-.38	.38	-.16	.18	-.17	.17	-.26	.36	-.30	.31	-.13	.25	-.15	.19

Notes: Bootstrap standard deviation estimators \widehat{sd}_B were computed using 50 bootstrap samples and their results are based on 500 replicated samples. In Population II the percentages of replicated samples in which the estimators \widehat{sd}_B were not computed because of convergence problems were 0.8%, 8.2% and 8.8% in U_1 , U_2 and U , respectively, whereas in Population I the respective percentages were all 0%. Results on sd estimators derived under the homogeneity assumption are based on 5000 replicated samples and no convergence problems were presented.

Table 6: Coverage probabilities and means and medians of relative lengths of the 95% confidence intervals of the population sizes, totals and means.

Population		I						II					
Sampling rates		$f_1 = 0.5$			$f_2 = 0.4$			$f_1 = 0.5$			$f_2 = 0.4$		
Response variable		Continuous			Binary			Continuous			Binary		
95% CI		cp	mrl	mdrl	cp	mrl	mdrl	cp	mrl	mdrl	cp	mrl	mdrl
UMLEs of sizes	$CI(\hat{\tau}_1^{(U)})$.95	.37	.36	.95	.37	.36	.76	.29	.29	.76	.29	.29
	$CI(\hat{\tau}_2^{(U)})$.97	5.6	1.8	.97	5.6	1.8	.51	1.7	.44	.51	1.7	.44
	$CI(\hat{\tau}^{(U)})$.98	1.4	.52	.98	1.4	.52	.63	.54	.26	.63	.54	.26
HTLEs of sizes	$CI(\hat{\tau}_{HT.1}^{(U)})$.78	.31	.30	.78	.31	.30	.44	.24	.24	.44	.24	.24
	$CI(\hat{\tau}_{HT.2}^{(U)})$.85	1.2	.85	.85	1.2	.85	.36	.66	.34	.36	.66	.34
	$CI(\hat{\tau}_{HT}^{(U)})$.78	.39	.33	.78	.39	.33	.28	.27	.21	.28	.27	.21
HTLEs of totals	$CI(\hat{Y}_{HT.1}^{(U)})$.96	.29	.28	.96	.32	.32	.42	.24	.23	.53	.29	.28
	$CI(\hat{Y}_{HT.2}^{(U)})$.90	1.1	.81	.98	1.4	1.0	.28	.56	.33	.63	.82	.49
	$CI(\hat{Y}_{HT}^{(U)})$.96	.35	.31	.98	.38	.35	.27	.25	.21	.47	.30	.26
HTLEs of means	$CI(\hat{\hat{Y}}_{HT.1}^{(U)})$.95	.10	.09	.96	.19	.19	.19	.08	.07	.55	.18	.18
	$CI(\hat{\hat{Y}}_{HT.2}^{(U)})$.98	.57	.52	.96	.79	.75	.91	.23	.18	.95	.58	.51
	$CI(\hat{\hat{Y}}_{HT}^{(U)})$.98	.33	.21	.96	.41	.29	.38	.11	.08	.84	.22	.19
HKLEs of totals	$CI(\hat{Y}_{HK.1}^{(U)})$.80	.34	.33	.66	.38	.37	.74	.28	.28	.78	.33	.33
	$CI(\hat{Y}_{HK.2}^{(U)})$.98	4.1	1.6	.99	6.4	2.0	.45	1.3	.43	.75	1.9	.60
	$CI(\hat{Y}_{HK}^{(U)})$.94	1.1	.50	.79	1.3	.54	.64	.44	.26	.75	.53	.31
HKLEs of means	$CI(\hat{\hat{Y}}_{HK.1}^{(U)})$.00	.07	.07	.17	.19	.19	.48	.03	.03	.96	.17	.17
	$CI(\hat{\hat{Y}}_{HK.2}^{(U)})$.00	.13	.12	.48	.55	.54	.77	.07	.07	.84	.47	.44
	$CI(\hat{\hat{Y}}_{HK}^{(U)})$.00	.08	.08	.08	.21	.20	.30	.04	.04	.92	.18	.17
Homo.	$CI(\hat{\tau}_1^{(H)})$.00	.08	.08	.00	.08	.08	.05	.10	.10	.05	.10	.10
MLEs	$CI(\hat{\tau}_2^{(H)})$.00	.17	.16	.00	.17	.16	.09	.24	.19	.09	.24	.19
sizes	$CI(\hat{\tau}^{(H)})$.00	.07	.07	.00	.07	.07	.02	.10	.09	.02	.10	.09
Homo.	$CI(\hat{Y}_{HT.1}^{(H)})$.00	.10	.10	.01	.14	.14	.09	.12	.11	.10	.15	.14
HTLEs	$CI(\hat{Y}_{HT.2}^{(H)})$.00	.21	.20	.32	.32	.32	.10	.25	.20	.26	.38	.32
totals	$CI(\hat{Y}_{HT}^{(H)})$.00	.09	.09	.00	.13	.13	.03	.11	.10	.06	.15	.14
Homo.	$CI(\hat{\hat{Y}}_{HT.1}^{(H)})$.00	.04	.04	.00	.13	.13	.17	.03	.03	.91	.14	.13
HTLEs	$CI(\hat{\hat{Y}}_{HT.2}^{(H)})$.00	.08	.08	.11	.39	.39	.66	.06	.06	.75	.40	.38
means	$CI(\hat{\hat{Y}}_{HT}^{(H)})$.00	.04	.04	.00	.13	.13	.13	.03	.03	.86	.14	.13

Notes: Results on confidence intervals derived under the assumption of heterogeneous link probabilities are based on 500 samples. In Population II the percentages of replicated samples in which the CIs were not computed because of convergence problems were 0.8%, 8.2% and 8.8% in U_1 , U_2 and U , respectively, whereas in Population I the respective percentages were all 0%. Results on CIs derived under the homogeneity assumption are based on 5000 replicated samples and no convergence problems were presented.

$\hat{Y}_{HK}^{(U)}$ were skewed to the right, whereas that of $\hat{Y}_{HK}^{(U)}$ was slightly skewed to the left in the case of the continuous response variable and symmetrical and the case of the binary variable. As in the previous study, the better performance of the HKLEs than that of the HTLEs in the case of Population II is result of the weaker correlations between the response variables and the inclusion probabilities. The performance of the estimators of the population sizes and population totals derived under the assumption of homogeneous link probabilities was bad: they presented serious problems of underestimation. However, the performance of the estimators $\hat{Y}_{HT,1}^{(H)}$, $\hat{Y}_{HT,2}^{(H)}$ and $\hat{Y}_{HT}^{(H)}$ was pretty acceptable in both cases continuous and binary response variables: they presented small biases and small variances, and in addition, their distributions were more or less symmetrical. We consider that the small bias of each one of these estimators could be explained because the bias of each estimator of the population total was practically the same as the bias of the corresponding estimator of the population size, and consequently, their biases were canceled out when the quotient was computed.

In Table 5 we present the results of the study on the standard deviation estimators. As can be seen, in general the results are not good, but there are some exceptions. In Population I, every one of the bootstrap estimators of the standard deviations of the point estimators that were derived under the heterogeneity assumption presented a large positive bias, except the estimators of the standard deviations of some estimators of parameters of the population U_1 , that is, τ_1 , Y_1 and \bar{Y}_1 , and those of the estimators $\hat{Y}_{HK,1}^{(U)}$, $\hat{Y}_{HK,2}^{(U)}$ and $\hat{Y}_{HK}^{(U)}$ which showed moderate values of bias and variability. The standard deviation estimators corresponding to the point estimators derived under the homogeneity assumption showed a large negative bias. In the case of Population II, and regardless of the type of response variable, continuous or binary, each one of the standard deviation estimators, either that derived under the homogeneity assumption as well as that not derived under that assumption, presented a large negative bias and a large variance. The exceptions were again the estimators $\hat{Y}_{HK,1}^{(U)}$, $\hat{Y}_{HK,2}^{(U)}$ and $\hat{Y}_{HK}^{(U)}$, which in the case of the binary response variable presented small values of bias and moderate values of variability.

The results of the Monte Carlo study on the confidence intervals are shown in Table 6. In the case of Population I, and regardless of the type of response variable, continuous or binary, every one of the CIs that was derived under the heterogeneity assumption presented good coverage probabilities, except the CIs obtained from the HTLEs $\hat{\tau}_{HT,1}^{(U)}$, $\hat{\tau}_{HT,2}^{(U)}$ and $\hat{\tau}_{HT}^{(U)}$, and from the HKLEs $\hat{Y}_{HK,1}^{(U)}$ and $\hat{Y}_{HK}^{(U)}$, which presented relatively low values of the cp, and those obtained from the HKLEs of the population means which showed very low values of the cp. Notice that in the case of the continuous response variable the CIs obtained from these estimators presented null values of the coverage probabilities. On the other hand, with respect to the relative lengths of the CIs, we have that the CIs of the parameters associated with U_2 , that is, τ_2 , Y_2 and \bar{Y}_2 , showed very large lengths, and consequently their performance was not good. The exception was the CI obtained from the HKLE $\hat{Y}_{HK,2}^{(U)}$, which showed a small value of mdlr. The relative lengths of the CIs of τ_1 , Y_1 and \bar{Y}_1 , as well as those of τ , Y and \bar{Y} were acceptable, and consequently their performance was also acceptable. The CIs derived under the homogeneity assumption presented null values of their cp, which were consequence of the large biases of the corresponding point estimators. Therefore, despite the

adequate values of their relative lengths, their performance was bad. In the case of Population II, the cp of the CIs that were constructed under the assumption of heterogeneous link probabilities were in general small. Only those associated with the CIs for \bar{Y}_2 (for both types of response variables), and particularly those corresponding to the CIs of the means based on the HKLEs were acceptable, say greater than 0.9. The small values of the cp were consequence of the biases of the corresponding point estimators. The relative lengths of these CIs were acceptable. Thus, based on the values of the cp and relative lengths, only the CIs of the population means of the binary variable obtained from the HKLEs had good performance, as well as those of mean \bar{Y}_2 of both variables (continuous and binary) obtained from the HTLE $\hat{Y}_{HT.2}$. The CIs of τ_1 based on $\hat{\tau}_1^{(U)}$ and those of the means of the binary variable based on the HTLEs showed regular performance. It is worth noting that even though the CIs of the means based on the HTLEs showed low values of the cp, the average CIs of the means $\bar{Y}_1 = 49.99$, $\bar{Y}_2 = 39.87$ and $\bar{Y} = 47.47$ of the continuous variable were (42.40, 45.90), (34.29, 41.47) and (42.72, 46.52), respectively, whereas the average CIs of the means $\bar{Y}_1 = 49.91$, $\bar{Y}_2 = 40.28$ and $\bar{Y} = 47.51$ of the binary variable were (41.43, 50.41), (31.22, 51.76) and (40.15, 49.17), respectively; therefore, these CIs still provide valuable information. Finally, the CIs constructed under the assumption of homogeneous link probabilities also presented small values of their cp, and consequently their performance was not good. The only exception were the CIs of the population means of the binary response variable where their cp showed moderate values, that is, between 0.75 and 0.91. Thus, based on the values of their cp and mdrl these CIs had regular performance. However, the average CIs of the means of the continuous variable were (50.24, 51.74), (39.47, 41.86) and (48.18, 49.61), whereas those of the means of the binary variable were (46.17, 52.66), (37.46, 52.77) and (45.37, 51.55); consequently, the information provided by these CIs provide is useful.

From the results of this study we have that inferences based on the UMLEs of the population totals are pretty acceptable, although the coverage probabilities of the CIs are affected when the assumed model of the link probabilities is not satisfied. Inferences based on the HTLEs of the population sizes are not good. HTLEs of the totals and means perform well when the correlation between the values of the response variable and the inclusion probabilities is not too small, say greater than or equal to 0.3. For values of the correlation coefficient less than 0.3, the HTLEs of the means are the only ones that perform acceptably, although the values of the coverage probabilities of the CIs are affected. HKLEs of the totals and means tend to perform acceptably when the values of the correlation coefficient between the response variable and the inclusion probabilities are small. In particular, HKLEs of the means perform pretty well in this situation.

7 Conclusions and suggestions for future research

In this work we have considered the link-tracing sampling variant proposed by Félix-Medina and Thompson (2004) and have proposed Horvitz-Thompson-like and Hájek-like estimators of population totals and means. This work extends that of Félix-Medina and Monjadin (2010) by assuming heterogeneous, rather than homogeneous, link probabilities which are modeled

by a Rasch model used by Félix-Medina et al. (2015). The variances of the proposed estimators are estimated by a variant of bootstrap which extends the variant used by the previously cited authors by estimating the variances of estimators of totals and means, in addition to the variances of estimators of population sizes. This variant of bootstrap allows the estimation of variances when the response variable is either continuous, discrete or binary. Large sample confidence intervals for the population parameters are constructed by assuming normal distributions of the estimators of the parameters, except in the case of the estimators of the population sizes, where log-normal distributions are assumed. In addition, confidence intervals for proportions are constructed using Korn and Graubard's (1998) proposal.

We evaluated the performance of the proposed estimators by means of two Monte Carlo studies. In the first study a finite population was constructed using data from the Add Health study. The result of this numerical study are promising; thus, if this population were more or less representative of the populations that could be found in applications of this methodology, then reliable inferences would be expected to be obtained. However, the results of the second study in which two populations were constructed using simulated data show that erroneous inferences might be obtained if some model assumptions were not satisfied. In particular, we found that if any of the assumptions is satisfied, then reliable inferences about population sizes, totals and means are obtained. Furthermore, we found that the assumption of the Poisson distribution of the sizes M_i s of the venues A_i s does not need to be satisfied to obtain reliable inferences. Nevertheless, we also found out that severe deviations from the Rasch model of the link probabilities lead to erroneous inferences, and that inferences about population sizes and totals are affected in greater extent than inferences about population means. In fact, inferences about the population means seem to be robust to deviations from the assumed models. In addition, we came upon that in any situation, the performance of the proposed bootstrap variance estimators is at most just good.

In the light of these results, we consider that the following issues are worthy of future research: (i) To develop a model for the link probabilities $p_{ij}^{(k)}$ s that be robust to deviations from the assumed model. For instance, to model $p_{ij}^{(k)}$ as a quadratic function of $\alpha_i^{(k)}$ and $\beta_j^{(k)}$, or to assume that the $\beta_j^{(k)}$ s are T-Student distributed instead of normally distributed, or to change the Rasch model by the latent classes model proposed by Pledger (2000) and which was used in the second Monte Carlo study to generate the values $x_{ij}^{(k)}$ s in Population II. (ii) To improve the proposed bootstrap variance estimators. For example, to predict the values of the response variable associated with the nonsampled elements by using a quadratic or a nonparametric regression model instead of a simple linear regression model. (iii) To enhance the proposed CIs of the population sizes, totals and means. For instance, using the bootstrap percentile method which does not require assuming a probability distribution for the estimator of the parameter of interest.

Acknowledgements

This research was supported by Grant PROFAPI-2011/057 of Universidad Autónoma de Sinaloa.

References

- Abramowitz, M., and I. A. Stegun. 1964. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. ninth Dover printing, tenth GPO printing. New York: Dover.
- Bernard, H. R., T. Hallett, A. Iovita, E. C. Johnsen, R. Lyster, C. McCarty, M. Mahy, M. J. Salganik, T. Saliuk, O. Scutelniciuc, G. A. Shelley, P. Sirinirund, S. Weir, and D. F. Stroup. 2010. "Counting hard-to-count populations: The network scale-up method for public health." *Sexually Transmitted Infections* 86 (Suppl. 2): ii11–ii15.
- Booth, J. G., R. W. Butler, and P. Hall. 1994. "Bootstrap methods for finite populations." *Journal of the American Statistical Association* 89:1282–1289.
- Chao, A. 1987. "Estimating the population size for capture-recapture data with unequal catchability." *Biometrics* 43:783–791.
- . 1988. "Estimating animal abundance with capture frequency data." *Journal of Wildlife Management* 52:295–300.
- Cheng, S., Eck, D. J., and F. W. Crawford. 2020. "Estimating the size of a hidden finite set: Large-sample behavior of estimators." *Statistics Surveys* 14:1–31.
- Chow, M., and S. K. Thompson. 2003. "Estimation with link-tracing sampling designs - A Bayesian approach." *Survey Methodology* 29:197–205.
- Coull, B. A., and A. Agresti. 1999. "The use of mixed logit models to reflect heterogeneity in capture-recapture studies." *Biometrics* 55:294–301.
- Crawford, F. W., Wu J., and R. Heimer. 2018. "Hidden population size estimation from respondent-driven sampling: A network approach." *Journal of the American Statistical Association* 113:755–766.
- Davison, A.C., and D.V. Hinkley. 1997. *Bootstrap Methods and their Applications*. New York: Cambridge University Press.
- Dombrowski, K., B. Khan, T. Wendel, K. McLean, E. Misshula, and R. Curtis. 2012. "Estimating the Size of the Methamphetamine-Using Population in New York City Using Network Sampling Techniques." *Advances in Applied Sociology* 2:245–252.
- Félix-Medina, M. H., and P. E. Monjardin. 2006. "Combining link-tracing sampling and cluster sampling to estimate the size of hidden populations: A Bayesian assisted approach." *Survey Methodology* 32:187–195.
- . 2010. "Combining link-tracing sampling and cluster sampling to estimate totals and means of hidden human populations." *Journal of Official Statistics* 26:603–631.
- Félix-Medina, M. H., P. E. Monjardin, and A. N. Aceves-Castro. 2015. "Combining link-tracing sampling and cluster sampling to estimate the size of a hidden population in presence of heterogeneous link-probabilities." *Survey Methodology*, pp. 349–376.

- Félix-Medina, M. H., and S. K. Thompson. 2004. "Combining cluster sampling and link-tracing sampling to estimate the size of hidden populations." *Journal of Official Statistics* 20:19–38.
- Frank, O., and T. Snijders. 1994. "Estimating the size of hidden populations using snowball sampling." *Journal of Official Statistics* 10:53–67.
- Giner, G., and G. K. Smyth. 2016. "statmod: probability calculations for the inverse Gaussian distribution." *R Journal* 8:339–351.
- Handcock, M. S., K. J. Gile, and C. M. Mar. 2014. "Estimating hidden population size using respondent-driven sampling data." *Electronic Journal of Statistics* 8:1491–1521.
- Harris, K. M. 2013. "The add health study: Design and accomplishments." Available at: www.cpc.unc.edu/projects/addhealth/data/guides/DesignPaperWIIIV.pdf.
- Heckathorn, D. D. 1997. "Respondent driven sampling: a new approach to the study of hidden samples." *Social Problems* 44:174–199.
- . 2002. "Respondent-driven sampling II: deriving valid population estimates from chain-referral samples of hidden populations." *Social Problems* 49:11–34.
- Heckathorn, D. D., and C. J. Cameron. 2017. "Snowball and multiplicity to respondent-driven sampling." *Annual Review of Sociology* 43:101–119.
- Johnston, L. G., A. J. Hakim, S. Dittrich, J. Burnett, E. Kim, and R. G. White. 2016. "A systematic review of published respondent-driven sampling surveys collecting behavioral and biologic data." *AIDS and Behavior* 20:1754–1776.
- Johnston, L. G., D. Prybylski, H. F. Raymond, A. Mirzazadeh, C. Manopaiboon, and W. McFarland. 2013. "Incorporating the service multiplier method in respondent-driven sampling surveys to estimate the size of hidden and hard-to-reach populations: Case studies from around the world." *Sexually Transmitted Diseases* 40:304–310.
- Kalton, G. 2009. "Methods for oversampling rare populations in social surveys." *Survey Methodology* 35:125–141.
- Khan, B., H.-W. Lee, I. Fellows, and K. Dombrowski. 2018. "One-step estimation of networked population size: Respondent-driven capture-recapture with anonymity." *PLoS ONE* 13 (4): 1–39. e0195959. <https://doi.org/10.1371/journal.pone.0195959>.
- Killworth, P., E. Johnsen, C. McCarty, G. Shelley, and H. Bernard. 1998a. "A social network approach to estimating seroprevalence in the United States." *Social Networks* 20:23–50.
- Killworth, P., C. McCarty, H. Bernard, G. Shelley, and E. Johnsen. 1998b. "Estimation of seroprevalence, rape and homelessness in the United States using a social network approach." *Evaluation Review* 22:289–308.
- Klov Dahl, A. S. 1989. "Urban Social Networks: Some Methodological Problems and Possibilities." In *The Small World*, edited by M. Kochen, 176–210. Norwood, NJ: Ablex.

- Korn, E. L., and B. I. Graubard. 1998. "Confidence intervals for proportions with small expected number of positive counts estimated from survey data." *Survey Methodology* 24:193–201.
- Lee, S., J. Wagner, R. Valliant, and S. Heeringa. 2014. "Recent developments of sampling hard-to-survey populations: An assessment." In *Hard-to-Survey Populations*, edited by R. Tourangeau, B. Edwards, T. Johnson, K. Wolter, and N. Bates (Eds.), 424–444. Cambridge: Cambridge University Press.
- MacKellar, D., L. Valleroy, J. Karon, G. Lemp, and R. Janssen. 1996. "The Young Men's Survey: Methods for estimating HIV seroprevalence and risk factors among young men who have sex with men." *Public Health Reports* 111(Suppl. 1):138–44.
- Magnani, R., K. Sabin, T. Saidel, and D. Heckathorn. 2005. "Review of sampling hard-to-reach populations for HIV surveillance." *AIDS* 19:S67–S72.
- Maltiel, R., A. E. Raftery, T. H. McCormick, and A. J. Baraff. 2015. "Estimating population size using the network scale up method." *The Annals of Applied Statistics* 9:1247–1277.
- Marpsat, M., and N. Razafindratsima. 2010. "Survey methods for hard-to-reach populations: Introduction to the special issue." *Methodological Innovations Online* 5:3–16.
- McCormick, T. H., M. J. Salganik, and T. Zheng. 2010. "How many people do you know? Efficiently estimating personal network size." *Journal of the American Statistical Association* 105:59–70.
- Meng, V. Y., and P. Gustafson. 2017. "Inferring population size: extending the multiplier method to incorporate multiple traits with a likelihood-based approach." *Stat* 6:4–13.
- Pledger, S. 2000. "Unified maximum likelihood estimates for closed capture-recapture models using mixtures." *Biometrics* 56:434–442.
- R Core Team. 2018. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Sanathanan, L. 1972. "Estimating the size of a multinomial population." *Annals of Mathematical Statistics* 43:142–152.
- Spreen, M. 1992. "Rare populations, hidden populations and link-tracing designs: What and why?" *Bulletin de Méthodologie Sociologique* 36:34–58.
- Spreen, M., and S. Bogaerts. 2015. "B-Graph Sampling to Estimate the Size of a Hidden Population." *Journal of Official Statistics* 31:723–736.
- Staudte, R. G., and S. J. Sheather. 1990. *Robust Estimation and Testing*. New York: Wiley.
- St. Clair, K., and D. O'Connell. 2012. "A Bayesian model for estimating population means using a link-tracing sampling design." *Biometrics* 68:165–173.
- Thompson, S. K. 2002. *Sampling, Second edition*. New Jersey: Wiley.
- Thompson, S.K., and O. Frank. 2000. "Model-based estimation with link-tracing sampling designs." *Survey Methodology* 26:87–98.

- Tourangeau, R. 2014. "Defining hard-to-survey populations." In *Hard-to-Survey Populations*, edited by R. Tourangeau, B. Edwards, T. Johnson, K. Wolter, and N. Bates (Eds.), 3–20. Cambridge: Cambridge University Press.
- UNAIDS, World Health Organization Working Group on Global HIV/AIDS, and STI Surveillance. 2010. *Guidelines on estimating the size of populations most at risk to HIV*. UNAIDS – Joint United Nations Programme on HIV/AIDS.
- Volz, E., and D. Heckathorn. 2008. "Probability based estimation theory for respondent driven sampling." *Journal of Official Statistics* 24:79–97.
- Williams, B. K., J. D. Nichols, and M. J. Conroy. 2002. *Analysis and Management of Animal Populations*. San Diego, California: Academic Press.
- Zelterman, D. 1988. "Robust estimation in truncated discrete distributions with application to capture-recapture experiments." *Journal of Statistical Planning and Inference* 18:225–237.