

RATE OF CONVERGENCE OF THE ATTRACTORS FOR REACTION-DIFFUSION EQUATIONS WITH NONLINEAR BOUNDARY CONDITIONS

FLANK D. M. BEZERRA AND LEONARDO PIRES

ABSTRACT. In this paper we study the rate of convergence of the asymptotic dynamics of reaction-diffusion equations with nonlinear Robin boundary conditions. We show how the rate of convergence of the global attractors can be affected by the variation of the potentials, boundary conditions and vectors fields.

Mathematical Subject Classification 2010: 34D45, 35B40, 41A25, 37L05, 37B35.

Keywords: global attractor; Morse-Smale semigroup; rate of convergence; reaction-diffusion equations; Robin boundary condition

CONTENTS

1. Introduction	1
2. Functional setting	4
3. Convergence of the resolvent operators	7
3.1. Resolvent operators	7
3.2. Eigenvalues, eigenfunctions and equilibria	9
4. Convergence of linear and nonlinear semigroups	11
5. Convergence of local unstable manifolds	13
6. Convergence of global attractors	16
7. Scalar reaction diffusion problem	17
8. Closing Remarks and Acknowledgments	20
Appendix A. General theory of rate of convergence of attractors	20
Appendix B. Shadowing theory and rate of convergence of attractors	23
References	25

1. INTRODUCTION

The continuity of the attractors for small perturbations of semilinear parabolic equations is a fundamental property to understand the qualitative properties present in the model that the equation represents. Under small changes in the initial data and parameters, a precise model of differential equations must have the permanence and continuity of bounded solutions

defined for all time, in addition some type of proximity or convergence of its solutions it must be proved. These topics were covered in several works where conditions were presented to guarantee the existence and continuity of attractors for semigroups generated by solutions of parabolic equations in a infinite dimensional Banach phase space, see for example [8, 11] and [12]. The problem is divided in two step; firstly, to show the continuity of the global attractors; secondly, once continuity is obtained, the question that arises is whether we can estimate it, that is, can we measure how fast the dynamics approach each other under small perturbations?

The answer known as rate of convergence of attractors is a problem which has attracted the attention of many researchers over the last years; see for instance [6] and the recent works [1, 3, 4, 5, 7, 9] and [10]. These works do not consider nonlinear boundary conditions, to our best knowledge, so a relevant question is how boundary conditions influence the estimates for the continuity of atrartores?

In this paper we are concerned with the rate of convergence of the asymptotic dynamics of reaction-diffusion equations with nonlinear Robin boundary conditions. Our main goal is to show how the rate of convergence of the attractors can be affected by the variation of the potentials, boundary conditions and vector fields. We will show explicitly how these terms added in the equation contribute to the speed of convergence of attractors. In particular, if the problem is one-dimensional, then we will significantly improve the rate of convergence obtained in [1], using the fact that the scalar equation generate a Morse-Smale semigroup in the energy space.

We consider the family of parabolic equations

$$(1.1) \quad \begin{cases} \partial_t u^\varepsilon - \operatorname{div}(p_\varepsilon(x) \nabla u^\varepsilon) + (\lambda + V_\varepsilon(x))u^\varepsilon = f^\varepsilon(u^\varepsilon) & \text{in } \Omega \times (0, \infty), \\ \frac{\partial u^\varepsilon}{\partial \vec{n}_\varepsilon} + (\lambda + b_\varepsilon(x))u^\varepsilon = g^\varepsilon(u^\varepsilon) & \text{on } \Gamma \times [0, \infty), \end{cases}$$

where $0 \leq \varepsilon \leq \varepsilon_0 \leq 1$, $\Omega \subset \mathbb{R}^N$ is a bounded smooth domain and Γ is the boundary of Ω . Here, $\frac{\partial u}{\partial \vec{n}_\varepsilon} = p_\varepsilon(x) \langle \nabla u, \vec{n}_\varepsilon \rangle$ denotes the conormal derivative and we chooce $\lambda \in \mathbb{R}$ sufficiently large such that $\operatorname{essinf}_{x \in \Omega} \{\lambda + V_\varepsilon\} \geq m_0$ and $\operatorname{essinf}_{x \in \Gamma} \{\lambda + b_\varepsilon\} \geq m_0$, for some $m_0 > 0$. The source f^ε acts on Ω and the potentials $V_\varepsilon(x)$ and $b_\varepsilon(x)$ are given function on Ω and Γ , respectively. The diffusion coefficients p_ε are smooth strictly positive function on Ω , such that, for $\varepsilon \in [0, \varepsilon_0]$,

$$(1.2) \quad 0 < m_0 \leq p_\varepsilon(x) \quad \text{and} \quad \|p_\varepsilon - p_0\|_{L^\infty(\Omega)} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0^+.$$

The potentials and the coefficients of terms of boundary verify

$$(1.3) \quad \|V_\varepsilon - V_0\|_{L^\infty(\Omega)} \leq \eta(\varepsilon) \quad \text{and} \quad \|b_\varepsilon - b_0\|_{L^\infty(\Gamma)} \leq \tau(\varepsilon),$$

where $\eta, \tau : [0, \varepsilon_0] \rightarrow [0, \infty)$ with $\eta(\varepsilon) \rightarrow 0^+, \tau(\varepsilon) \rightarrow 0^+$ as $\varepsilon \rightarrow 0^+$. It is possible to consider V_ε and b_ε in more general L^p spaces as we can see in [16] but in order to simplify the arguments we will consider convergences only in L^∞ .

If we denote $f^\varepsilon : H^1(\Omega) \rightarrow L^2(\Omega)$ and $g^\varepsilon : H^1(\Gamma) \rightarrow H^{-\frac{1}{2}}(\Gamma)$ the Nemitskii operators associated with the functions f^ε and g^ε , respectively, we assume the following convergence

$$(1.4) \quad \|f^\varepsilon(u) - f^0(u)\|_{L^2(\Omega)} \leq \kappa(\varepsilon) \quad \text{and} \quad \|g^\varepsilon(v) - g^0(v)\|_{H^{-\frac{1}{2}}(\Gamma)} \leq \xi(\varepsilon),$$

where $u \in H^1(\Omega)$, $v \in H^1(\Gamma)$ and $\kappa, \xi : [0, \varepsilon_0] \rightarrow [0, \infty)$ with $\kappa(\varepsilon) \rightarrow 0^+, \xi(\varepsilon) \rightarrow 0^+$ as $\varepsilon \rightarrow 0^+$.

Our goal is to study the behavior of solutions of (1.1) as $\varepsilon \rightarrow 0^+$. Heuristically, we have

$$\partial_t u^\varepsilon - \operatorname{div}(p_\varepsilon(x) \nabla u^\varepsilon) + (\lambda + V_\varepsilon(x))u^\varepsilon = f^\varepsilon(u^\varepsilon) \text{ in } \Omega \times (0, \infty),$$

and from the convergence properties (1.3) and (1.4) we have in the limit

$$\partial_t u^0 - \operatorname{div}(p_0(x) \nabla u^0) + (\lambda + V_0(x))u^0 = f^0(u^0) \text{ in } \Omega \times (0, \infty).$$

Analogously, we have

$$\frac{\partial u^\varepsilon}{\partial \vec{n}_\varepsilon} + (\lambda + b_\varepsilon(x))u^\varepsilon = p_\varepsilon(x) \langle \nabla u^\varepsilon, \vec{n}_\varepsilon \rangle + (\lambda + b_\varepsilon(x))u^\varepsilon = g^\varepsilon(u^\varepsilon) \text{ on } \Gamma \times [0, \infty),$$

and from the convergence properties (1.2), (1.3) and (1.4) we have in the limit

$$\frac{\partial u^0}{\partial \vec{n}_0} + (\lambda + b_0(x))u^0 = p_0(x) \langle \nabla u^0, \vec{n}_0 \rangle + (\lambda + b_0(x))u^0 = g^0(u^0) \text{ on } \Gamma \times [0, \infty).$$

In the next sections we will show that the asymptotic behavior of the equation (1.1) for $\varepsilon \neq 0$ is associated with the equation (1.1) for $\varepsilon = 0$. In fact we will prove that the convergence of eigenvalue, eigenfunction, equilibrium points, spectral projections, local invariant unstable manifolds and global attractors as $\varepsilon \rightarrow 0^+$ can be estimate by

$$(1.5) \quad [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)]^l, \quad N \geq 2, \quad 0 < l < 1$$

and for the scalar situation $N = 1$ we can estimate by

$$(1.6) \quad |\log(\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon))| [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)].$$

Notice that (1.6) goes to zero faster than (1.5) as $\varepsilon \rightarrow 0^+$ and these estimates is composed by convergence of the potentials, boundary terms, and vector fields.

The terms l and \log in estimates above in fact represent a loss in the rates of convergence (1.5) and (1.6). A result without these terms is still an open problem which can be called an optimal rate problem. We already know that in the situation of a family of PDEs whose asymptotic behavior is dictated by an ODE it is possible to obtain the optimal rate, see [9]. In addition, constructing explicit estimates that give rise to the rate of convergence of attractors

can be hard-working as we can see in [10], where an elliptic problem involving a divergent operator with localized large diffusion was studied.

In order to prove the rate of convergence (1.5) and (1.6) for equation (1.1) we organize this paper as follows. In Section 2 we introduce the appropriate functional setting to deal with problem (1.1) and we state the results on the well posedness, regularity and existence of global attractors. In Section 3 we show the convergence with rate of the resolvent operators, eigenvalues, eigenfunction, spectral projection and equilibrium points. The Section 4 is reserved to obtain the convergence of linear and nonlinear semigroups and to obtain one part of rate of convergence. In Section 5 we study the local unstable manifold around a equilibrium point through local linearization. In Section 6 we present our main result on the rate of convergence of attractors. The Section 7 we deal with the special one dimensional case. We conclude with two appendices: Appendix A is related to the general theory of rate of convergence of attractors, and Appendix B deals with Shadowing Theory and its applications in the convergence of attractors.

2. FUNCTIONAL SETTING

In this section, we introduce the functional framework to study (1.1) and, we present a fundamental result on equivalence between the norms of the fractional powers space and the Sobolev space $H^1(\Omega)$.

The natural form to study (1.1) is as evolution equation in the Sobolev space $H^s(\Omega)$, $s > 0$ whose dual space we denote by $H^{-s}(\Omega)$. The duality between these spaces will be denoted by $\langle \cdot, \cdot \rangle_{-s, s}$. In particular the scalar product in $L^2(\Omega)$ we will be denoted by $\langle \cdot, \cdot \rangle$. For the boundary terms we consider the trace space $H^s(\Gamma)$ and the the trace operator $\gamma_{tr} : H^s(\Omega) \rightarrow H^{s-\frac{1}{2}}(\Gamma)$, for $s > \frac{1}{2}$. Since $H^{\frac{1}{2}}(\Gamma) \hookrightarrow L^2(\Gamma) \hookrightarrow H^{-\frac{1}{2}}(\Gamma) \hookrightarrow H^{-1}(\Gamma)$ we define, for $f \in H^1(\Omega)$

$$\langle \gamma_{tr}(f), \varphi \rangle_{-1, 1} = \int_{\Gamma} f \varphi \, d\sigma, \quad \forall \varphi \in H^1(\Omega).$$

We also consider the normal derivative operator, relative to the normal divergent operator $-\operatorname{div}(p_{\varepsilon}(x)\nabla u)$, defined as follows: if $u \in \{z \in H^1(\Omega) : -\operatorname{div}(p_{\varepsilon}(x)\nabla z) \in L^2(\Omega)\}$ then $\frac{\partial u}{\partial n_{\varepsilon}} \in H^{-\frac{1}{2}}(\Gamma)$ and it is defined as

$$(2.1) \quad \left\langle \frac{\partial u}{\partial n_{\varepsilon}}, \gamma_{tr}(v) \right\rangle_{-\frac{1}{2}, \frac{1}{2}} = \int_{\Gamma} \operatorname{div}(p_{\varepsilon}(x)\nabla u) v \, dx = - \int_{\Gamma} p_{\varepsilon}(x) \nabla u \nabla v \, dx,$$

for every $v \in H^1(\Omega)$.

We define the unbounded linear operator $A_{\varepsilon} : D(A_{\varepsilon}) \subset L^2(\Omega) \rightarrow L^2(\Omega)$ by

$$A_{\varepsilon} = -\operatorname{div}(p_{\varepsilon}(x)\nabla u) + (\lambda + V_{\varepsilon})u$$

with domain

$$D(A_\varepsilon) = \left\{ u \in H^2(\Omega) : \frac{\partial u}{\partial \vec{n}_\varepsilon} + (\lambda + b_\varepsilon(x))u = 0 \text{ on } \Gamma \right\}.$$

The operator A_ε is selfadjoint and has compact resolvent in $L^2(\Omega)$. Moreover, if $\mu > \lambda_\varepsilon^1$, the first eigenvalue of A_ε , then the operator $\mu I + A_\varepsilon$ is positive. If we denote by X_ε^β its fractional power space, we have $X_\varepsilon^1 = D(A_\varepsilon)$, $X_\varepsilon^{\frac{1}{2}} = H^1(\Omega)$, $X_\varepsilon^0 = L^2(\Omega)$, $X_\varepsilon^{-\frac{1}{2}} = H^{-1}(\Omega)$, see [13]. By interpolation $X_\varepsilon^\beta \hookrightarrow H^{2\beta}(\Omega)$, for $0 \leq \beta \leq \frac{1}{2}$ and by duality $H^{-2\beta}(\Omega) \hookrightarrow X_\varepsilon^{-\beta}$ for $0 < \beta \leq \frac{1}{2}$, where both embedding constants are uniform in ε .

Notice that the choice of λ in (1.1) is not restrictive since we can translate the operator A_ε for a parameter μ so that $\mu + A_\varepsilon$ is positive and then the fractional power space is defined. Thus we fix λ just to write the inner product in a clear form and then we study the variational formulation of the elliptic divergent operator. In fact the operator A_ε is the realization in $L^2(\Omega)$ of the canonical isomorphism between $H^1(\Omega)$ and its dual $H^{-1}(\Omega)$, defined by means of the bilinear form

$$(2.2) \quad \langle A_\varepsilon u, v \rangle_{-1,1} = \int_\Omega p_\varepsilon(x) \nabla u \nabla v \, dx + \int_\Omega (\lambda + V_\varepsilon) uv \, dx + \int_\Gamma (\lambda + b_\varepsilon(x)) uv \, d\sigma,$$

for every $u, v \in H^1(\Omega)$. In this way, by (2.1), we can rewrite (2.2) as

$$\langle A_\varepsilon u, v \rangle_{-1,1} = \langle -\operatorname{div}(p_\varepsilon(x) \nabla u) + (\lambda + V_\varepsilon)u, v \rangle_\Omega + \left\langle \frac{\partial u}{\partial \vec{n}_\varepsilon}, \gamma_{tr}(v) \right\rangle_{-\frac{1}{2}, \frac{1}{2}},$$

for $u \in D(A_\varepsilon)$ and $v \in H^1(\Omega)$.

Lemma 2.1. The inner product in $X_\varepsilon^{\frac{1}{2}}$ given by

$$\langle u, v \rangle_{X_\varepsilon^{\frac{1}{2}}} = \int_\Omega p_\varepsilon(x) \nabla u \nabla v \, dx + \int_\Omega (\lambda + V_\varepsilon) uv \, dx + \int_\Gamma (\lambda + b_\varepsilon(x)) uv \, d\sigma = \langle A_\varepsilon u, v \rangle_{-1,1}$$

gives a norm in $H^1(\Omega)$, equivalent to the usual one.

Proof. We have $\operatorname{ess\,inf}_{x \in \Omega} \{\lambda + V_\varepsilon\}, \operatorname{ess\,inf}_{x \in \Gamma} \{\lambda + b_\varepsilon\} \geq m_0$ and by (1.2) it follows that

$$m_0 \|u\|_{H^1(\Omega)} \leq \|u\|_{X_\varepsilon^{\frac{1}{2}}} \leq M_0 \|u\|_{H^1(\Omega)},$$

where $\sup_{0 \leq \varepsilon \leq \varepsilon_0} \{\|p_\varepsilon\|_{L^\infty(\Omega)}, \|\lambda + V_\varepsilon\|_{L^\infty(\Omega)}, \|\lambda + b_\varepsilon\|_{L^\infty(\Gamma)}\} \leq M_0$. \square

Now, if u is a solution of (1.1), multiplying (1.1) by a test function φ and integrating by parts over Ω we have

$$\int_\Omega \partial_t u^\varepsilon \varphi \, dx + \int_\Omega p_\varepsilon(x) \nabla u^\varepsilon \nabla \varphi \, dx + \int_\Omega (\lambda + V_\varepsilon) u^\varepsilon \varphi \, dx = \int_\Omega f^\varepsilon(u^\varepsilon) \varphi \, dx + \int_\Gamma g^\varepsilon(\gamma_{tr}(u^\varepsilon)) \gamma_{tr}(\varphi) \, d\sigma.$$

But

$$\langle g^\varepsilon(\gamma_{tr}(u^\varepsilon)), \gamma_{tr}(\varphi) \rangle_\Gamma = \int_\Gamma g^\varepsilon(\gamma_{tr}(u^\varepsilon)) \gamma_{tr}(\varphi) \, d\sigma = \langle \gamma_{tr}(g^\varepsilon(u^\varepsilon)), \gamma_{tr}(\varphi) \rangle_{-1,1},$$

then

$$\langle \partial_t u^\varepsilon, \varphi \rangle_{-1,1} + \langle A_\varepsilon u^\varepsilon, \varphi \rangle_{-1,1} = \langle f^\varepsilon(u^\varepsilon), \varphi \rangle_{-1,1} + \langle g^\varepsilon(\gamma_{tr}(u^\varepsilon)), \gamma_{tr}(\varphi) \rangle_{-1,1}.$$

Hence, to obtain appropriated semilinear formulation for (1.1) we consider nonlinear maps, in the form $h^\varepsilon(u) = f_\Omega^\varepsilon(u) + g_\Gamma^\varepsilon(u)$ with

$$(2.3) \quad \langle h^\varepsilon(u), \varphi \rangle = \langle f_\Omega^\varepsilon(u), \varphi \rangle_\Omega + \langle g_\Gamma^\varepsilon(u), \varphi \rangle_\Gamma,$$

where h^ε acts from $H^1(\Omega)$ to $H^{-\beta}(\Omega)$, for $-\beta < 0$ in the interval $[-\frac{1}{2}, -\frac{1}{4}]$. In fact, the geometric theory for abstract parabolic problems under nonlinear boundary conditions require that the Nemitskii functional acts on the fractional power spaces $X_\varepsilon^\alpha \rightarrow X_\varepsilon^{-\beta}$ where $-\beta < 0 < \alpha$ and $\alpha + \beta \leq 1$, see [13]. We are interested in the L^2 -theory so, we take $\alpha = \frac{1}{2}$ which implies $-\frac{1}{2} \leq -\beta$. On the other hand, if $-\beta + 1 \geq \frac{3}{4}$, the space $X_\varepsilon^{-\beta+1}$ incorporates the boundary condition, thus we must have $-\beta + 1 < \frac{3}{4}$, that is, $-\beta < -\frac{1}{4}$.

From now on, we choose $-\frac{1}{2} \leq -\beta < -\frac{1}{4}$ and we rewrite (1.1) as

$$(2.4) \quad \begin{cases} \frac{du^\varepsilon}{dt} + A_\varepsilon u^\varepsilon = h^\varepsilon(u^\varepsilon), \\ u^\varepsilon(0) = u_0^\varepsilon \in H^1(\Omega), \quad \varepsilon \in [0, \varepsilon_0], \end{cases}$$

where the nonlinearity $h^\varepsilon = f_\Omega^\varepsilon + g_\Gamma^\varepsilon : H^1(\Omega) \rightarrow H^{-\beta}(\Omega)$ is given by (2.3).

We just have to formulate the problem in the phase space $H^1(\Omega)$, now we assume that $f^\varepsilon, g^\varepsilon$ satisfy the standard growth, sign and dissipative conditions stated in [2] in order to ensure that (2.4), for each $\varepsilon \in [0, \varepsilon_0]$, is globally well posed and its solutions are classical and continuously differentiable with respect to the initial data. Therefore we are able to consider in $H^1(\Omega)$ the family of nonlinear semigroups $\{T_\varepsilon(t); t \geq 0\}_{\varepsilon \in [0, \varepsilon_0]}$ defined by $T_\varepsilon(t) = u^\varepsilon(t, u_0^\varepsilon)$, $t \geq 0$, where $u^\varepsilon(t, u_0^\varepsilon)$ is the solution of (2.4) through $u_0^\varepsilon \in H^1(\Omega)$ and

$$(2.5) \quad T_\varepsilon(t)u_0^\varepsilon = e^{-A_\varepsilon t}u_0^\varepsilon + \int_0^t e^{-A_\varepsilon(t-s)}h^\varepsilon(T_\varepsilon(s)u_0^\varepsilon)ds, \quad t \geq 0,$$

where $e^{-A_\varepsilon(\cdot)}$ is the linear semigroup generated by operator $-A_\varepsilon$. The existence of attractor and uniform bounds for semigroups $\{T_\varepsilon(t); t \geq 0\}$ associated with (2.4) are also established in [2]. In fact we have the following result.

Theorem 2.2. The nonlinear semigroup $\{T_\varepsilon(t); t \geq 0\}$ associated with (2.4) has global attractor \mathcal{A}_ε in $H^1(\Omega)$. Furthermore

$$\sup_{\varepsilon \in [0, \varepsilon_0]} \sup_{w \in \mathcal{A}_\varepsilon} \|w\|_{H^1(\Omega)} < \infty \quad \text{and} \quad \sup_{\varepsilon \in [0, \varepsilon_0]} \sup_{w \in \mathcal{A}_\varepsilon} \|w\|_{L^\infty(\Omega)} < \infty,$$

for some $\varepsilon_0 > 0$.

Once the uniform bound in $L^\infty(\Omega)$ for the attractors has been obtained, we may perform a cutoff to the nonlinearities $f^\varepsilon, g^\varepsilon$ so that the new nonlinearity is globally Lipschitz and globally bounded with bounded derivatives up to second order, it coincides with the original one in a L^∞ -neighborhood of all the attractors and is strictly dissipative outside this neighborhood. This

guarantees that the system with the new nonlinearities have attractor which coincide exactly with the original ones. In addition it follows from (1.4) and (2.3) that

$$(2.6) \quad \|h^\varepsilon(u) - h^0(v)\|_{H^{-\beta}(\Omega)} \leq C\|u - v\|_{H^1(\Omega)} + \kappa(\varepsilon) + \xi(\varepsilon), \text{ for all } u, v \in H^1(\Omega).$$

3. CONVERGENCE OF THE RESOLVENT OPERATORS

In this section we study the convergence of the resolvent operators and its consequences.

3.1. Resolvent operators. We study the elliptic problem $A_\varepsilon u^\varepsilon = v$ to obtain the first part of the rate of convergence. We show the convergence of the resolvent operators A_ε^{-1} to A_0^{-1} and we establish that the rate of this convergence is of order $\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)$.

Lemma 3.1. For $v = v_\Omega + v_\Gamma : H^1(\Omega) \rightarrow H^{-\beta}(\Omega)$ with $\|v\|_{\mathcal{L}(H^1(\Omega), H^{-\beta}(\Omega))} \leq 1$ and $\varepsilon \in [0, \varepsilon_0]$, let u^ε be the solution of elliptic problem $A_\varepsilon u^\varepsilon = v$. Then there is a constant $C > 0$ independent of ε such that

$$(3.1) \quad \|u^\varepsilon - u^0\|_{H^1(\Omega)} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)].$$

Proof. Since u^ε is a weak solution of the problem $A_\varepsilon u^\varepsilon = v$, we have

$$(3.2) \quad \int_\Omega p_\varepsilon \nabla u^\varepsilon \nabla \varphi \, dx + \int_\Omega (\lambda + V_\varepsilon) u^\varepsilon \varphi \, dx + \int_\Gamma (\lambda + b_\varepsilon) u^\varepsilon \varphi \, d\sigma = \int_\Omega v \varphi \, dx + \int_\Gamma v \varphi \, d\sigma,$$

for all test function φ and $\varepsilon \in [0, \varepsilon_0]$.

If we take $\varphi = u^\varepsilon - u^0$ in (3.2), we obtain

$$\begin{aligned} \int_\Omega p_\varepsilon \nabla u^\varepsilon (\nabla u^\varepsilon - \nabla u^0) \, dx + \int_\Omega (\lambda + V_\varepsilon) u^\varepsilon (u^\varepsilon - u^0) \, dx + \int_\Gamma (\lambda + b_\varepsilon) u^\varepsilon (u^\varepsilon - u^0) \, d\sigma \\ = \int_\Omega v (u^\varepsilon - u^0) \, dx + \int_\Gamma v (u^\varepsilon - u^0) \, d\sigma, \end{aligned}$$

and

$$\begin{aligned} \int_\Omega p_0 \nabla u^0 (\nabla u^\varepsilon - \nabla u^0) \, dx + \int_\Omega (\lambda + V_0) u^0 (u^\varepsilon - u^0) \, dx + \int_\Gamma (\lambda + b_0) u^0 (u^\varepsilon - u^0) \, d\sigma \\ = \int_\Omega v (u^\varepsilon - u^0) \, dx + \int_\Gamma v (u^\varepsilon - u^0) \, d\sigma. \end{aligned}$$

Now, subtracting the above expressions, we have

$$\begin{aligned} \int_\Omega p_\varepsilon |\nabla u^\varepsilon - \nabla u^0|^2 \, dx + \int_\Omega (\lambda + V_\varepsilon) |u^\varepsilon - u^0|^2 \, dx + \int_\Gamma (\lambda + b_\varepsilon) |u^\varepsilon - u^0|^2 \, d\sigma \\ \leq \int_\Omega |p_\varepsilon - p_0| \nabla u^0 (\nabla u^\varepsilon - \nabla u^0) \, dx + \int_\Omega |V_\varepsilon - V_0| u^0 (u^\varepsilon - u^0) \, dx + \int_\Gamma |b_\varepsilon - b_0| u^0 (u^\varepsilon - u^0) \, d\sigma, \end{aligned}$$

which implies by Hölder's inequality

$$\begin{aligned}
\|u^\varepsilon - u^0\|_{H^1(\Omega)}^2 &\leq \int_{\Omega} |p_\varepsilon - p_0| \nabla u^0 (\nabla u^\varepsilon - \nabla u^0) dx + \int_{\Omega} |V_\varepsilon - V_0| u^0 (u^\varepsilon - u^0) dx \\
&\quad + \int_{\Gamma} |b_\varepsilon - b_0| u^0 (u^\varepsilon - u^0) d\sigma \\
&\leq C \|\nabla u^\varepsilon - \nabla u^0\|_{L^2(\Omega)} \|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + C \|V_\varepsilon - V_0\|_{L^2(\Omega)} \|u^\varepsilon - u^0\|_{L^2(\Omega)} \\
&\quad + C \|b_\varepsilon - b_0\|_{L^\infty(\Gamma)} \|u^\varepsilon - u^0\|_{L^2(\Omega)},
\end{aligned}$$

where C depends on u^0 and embedding constants, but independent of ε . Finally, from (1.2) and (1.3), we obtain (3.1). \square

As an immediate consequence of the previous Lemma 3.1, we can prove

Theorem 3.2. There exists constant $C > 0$ independent of ε such that

$$\|A_\varepsilon^{-1} - A_0^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)].$$

Furthermore, there exists $\phi \in (\frac{\pi}{2}, \pi)$ such that for all

$$\mu \in \Sigma_{\nu, \phi} = \{\mu \in \mathbb{C} : |\arg(\mu + \nu)| \leq \phi\} \setminus \{\mu \in \mathbb{C} : |\mu + \nu| \leq r\},$$

for some $r > 0$,

$$(3.3) \quad \|(\mu + A_\varepsilon)^{-1} - (\mu + A_0)^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)],$$

where $C = C(\mu) > 0$ is a constant independent of ε .

Proof. The first part is an immediate consequence of Lemma 3.1. Indeed, let $A_\varepsilon u^\varepsilon = v$, where $v = v_\Omega + v_\Gamma : H^1(\Omega) \rightarrow H^{-\beta}(\Omega)$ with $\|v\|_{\mathcal{L}(H^1(\Omega), H^{-\beta}(\Omega))} \leq 1$; that is, $u^\varepsilon = A_\varepsilon^{-1}v$. Hence the result follows by (3.1).

Now, let $\rho(A_\varepsilon)$ be the resolvent set of the operator A_ε , $\varepsilon \in [0, \varepsilon_0]$. If $\mu \in \rho(-A_\varepsilon) \cap \rho(-A_0)$, we choose $\varphi \in (\frac{\pi}{2}, \pi)$ suitable in order to get the sectorial estimates

$$\|(\mu + A_\varepsilon)^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega))} \leq \frac{M_\varphi}{|\mu|}, \quad \varepsilon \in [0, \varepsilon_0],$$

where M_φ independent of ε , see [1] for more details.

Therefore

$$\|A_\varepsilon(\mu + A_\varepsilon)^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega), L^2(\Omega))} \leq 1 + M_\varphi, \quad \varepsilon \in [0, \varepsilon_0].$$

But, for $v : H^1(\Omega) \rightarrow H^{-\beta}(\Omega)$ we can write

$$(3.4) \quad A_\varepsilon^{\frac{1}{2}}[(\mu + A_\varepsilon)^{-1} - (\mu + A_0)^{-1}]v = A_\varepsilon(\mu + A_\varepsilon)^{-1}A_\varepsilon^{\frac{1}{2}}(A_\varepsilon^{-1} - A_0^{-1})A_0(\mu + A_0)^{-1}v$$

and thus

$$\begin{aligned}
& \|(\mu + A_\varepsilon)^{-1} - (\mu + A_0)^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \\
& \leq \|A_\varepsilon(\mu + A_\varepsilon)^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega), L^2(\Omega))} \|A_\varepsilon^{\frac{1}{2}}(A_\varepsilon^{-1} - A_0^{-1})\|_{\mathcal{L}(H^{-\beta}(\Omega), L^2(\Omega))} \|A_0(\mu + A_0)^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega), L^2(\Omega))} \\
& \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)],
\end{aligned}$$

for some constant $C = C(\varphi, \mu) > 0$ independent of ε . \square

3.2. Eigenvalues, eigenfunctions and equilibria. Thanks to Theorem 3.2 we will establish the rate of convergence of eigenvalues, eigenfunctions and spectral projections associated to operators A_ε .

The spectrum $\sigma(-A_\varepsilon)$ of $-A_\varepsilon$, ordered and counting multiplicity is given by

$$(3.5) \quad \dots - \lambda_m^\varepsilon < -\lambda_{m-1}^\varepsilon < \dots < -\lambda_0^\varepsilon$$

with $\{\varphi_i^\varepsilon\}_{i=0}^\infty$ the eigenfunctions related, for any $\varepsilon \in [0, \varepsilon_0]$. We consider the spectral projection onto the space generated by the first m eigenvalues, i.e., if ω is an appropriated closed curve in the resolvent set $\rho(-A_0)$ of $-A_0$ around $\{-\lambda_0^0, \dots, -\lambda_{m-1}^0\}$, then we define the spectral projection

$$Q_\varepsilon = \frac{1}{2\pi i} \int_\omega (\mu + A_\varepsilon)^{-1} d\mu, \quad \varepsilon \in [0, \varepsilon_0].$$

Theorem 3.3. The family of projections Q_ε converges to Q_0 in the uniform operator topology of $\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))$ as $\varepsilon \rightarrow 0^+$ and

$$\|Q_\varepsilon - Q_0\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)],$$

where $C > 0$ independent of ε .

Proof. Note that

$$Q_\varepsilon - Q_0 = \frac{1}{2\pi i} \int_\omega [(\mu + A_\varepsilon)^{-1} - (\mu + A_0)^{-1}] d\mu$$

and using the Theorem 3.2, we obtain

$$\|Q_\varepsilon(\mu_0) - Q_0(\mu_0)\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)].$$

\square

Next we present the convergence of the eigenvalues.

Corollary 3.4. If $\lambda_0 \in \sigma(A_0)$ and if there is $\lambda_\varepsilon \in \sigma(A_\varepsilon)$ such that $\lambda_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} \lambda_0$, then

$$|\lambda_\varepsilon - \lambda_0| \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)].$$

Proof. If $-\lambda_0$ is an isolated eigenvalue for $-A_0$, we may define Q_ε as above and if there is a λ_ε which is an eigenvalue of A_ε such that $\lambda_\varepsilon \xrightarrow{\varepsilon \rightarrow 0^+} \lambda_0$, then the composition $Q_\varepsilon Q_0$ is an isomorphism between $\text{Im}(Q_0)$ and $\text{Im}(Q_\varepsilon)$ (using extension to the dual space). Therefore, for each ε suitably small, there exists $u_\varepsilon \in \text{Im}(Q_0)$, $\|u_\varepsilon\|_{H^1(\Omega)} = 1$ such that $Q_\varepsilon u_\varepsilon$ is an eigenvector of A_ε associated to λ_ε and

$$|\lambda_\varepsilon - \lambda_0| \leq \|\lambda_\varepsilon Q_0 u_\varepsilon - \lambda_\varepsilon Q_\varepsilon u_\varepsilon\|_{H^1(\Omega)} + \|\lambda_\varepsilon Q_\varepsilon u_\varepsilon - \lambda_0 Q_0 u_\varepsilon\|_{H^1(\Omega)}.$$

The result follows by Theorem (3.3). \square

The equilibrium solutions of (2.4) are those which are independent of time, i.e., for $\varepsilon \in [0, \varepsilon_0]$, they are the solutions of the elliptic problem $A_\varepsilon u^\varepsilon - h^\varepsilon(u^\varepsilon) = 0$. We denote by \mathcal{E}_ε the set of the equilibrium solutions of (2.4) and we say that $u_*^\varepsilon \in \mathcal{E}_\varepsilon$ is hyperbolic if

$$\sigma(A_\varepsilon - (h^\varepsilon)'(u_*^\varepsilon)) \cap \{\mu \in \mathbb{C}; \text{Re}(\mu) = 0\} = \emptyset.$$

Since hyperbolicity of the equilibrium is a quite common property for reaction diffusion equations, we assume \mathcal{E}_0 is composed of finitely many hyperbolic equilibrium points.

The next results prove that the family $\{\mathcal{E}_\varepsilon\}_{\varepsilon \in [0, \varepsilon_0]}$ is continuous at $\varepsilon = 0$, thus for ε sufficiently small, \mathcal{E}_ε is composed by a finite number of hyperbolic equilibrium points.

Theorem 3.5. Let $u_*^0 \in \mathcal{E}_0$. Then for each $\varepsilon > 0$ sufficiently small, there is $\delta > 0$ such that the equation $A_\varepsilon u - h^\varepsilon(u) = 0$ has only solution $u_*^\varepsilon \in \{u \in H^1(\Omega); \|u - u_*^0\|_{H^1(\Omega)} \leq \delta\}$. Moreover

$$(3.6) \quad \|u_*^\varepsilon - u_*^0\|_{H^1(\Omega)} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)].$$

Proof. The proof is standard, see [8, Chapter 14] except for the estimate (3.6). The functions u_*^ε and u_*^0 given by

$$u_*^0 = (A_0 + R_0)^{-1}[h^0(u_*^0) + R_0 u_*^0] \quad \text{and} \quad u_*^\varepsilon = (A_\varepsilon + R_\varepsilon)^{-1}[h^\varepsilon(u_*^\varepsilon) + R_\varepsilon u_*^\varepsilon],$$

where $R_0 = -(h^0)'(u_*^0)$ and $R_\varepsilon = -(h^\varepsilon)'(u_*^\varepsilon)$. Thus

$$\begin{aligned} \|u_*^\varepsilon - u_*^0\|_{H^1(\Omega)} &\leq \|(A_\varepsilon + R_\varepsilon)^{-1}[h^\varepsilon(u_*^\varepsilon) - h^0(u_*^0) + R_\varepsilon(u_*^\varepsilon - u_*^0)]\|_{H^1(\Omega)} \\ &\quad + \|[(A_\varepsilon + R_\varepsilon)^{-1} - (A_0 + R_0)^{-1}][h^0(u_*^0) + R_\varepsilon u_*^0]\|_{H^1(\Omega)} \\ &\quad + \|(A_\varepsilon + R_0)^{-1}[(R_\varepsilon - R_0)u_*^0]\|_{H^1(\Omega)}. \end{aligned}$$

Using an identity similar to (3.4) and Theorem 3.2 we obtain

$$\|[(A_\varepsilon + R_\varepsilon)^{-1} - (A_0 + R_0)^{-1}][h^0(u_*^0) + R_\varepsilon u_*^0]\|_{H^1(\Omega)} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)].$$

Now we denote $z^\varepsilon = h^\varepsilon(u_*^\varepsilon) - h^0(u_*^0) + R_\varepsilon(u_*^\varepsilon - u_*^0)$. Since h^ε is continuously differentiable, for all $\delta > 0$, there is $\varepsilon > 0$ sufficiently small such that

$$\|(A_\varepsilon + R_\varepsilon)^{-1} z^\varepsilon\|_{H^1(\Omega)} \leq \delta \|(A_\varepsilon + R_\varepsilon)^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} [\|u_*^\varepsilon - u_*^0\|_{H^1(\Omega)} + \kappa(\varepsilon) + \xi(\varepsilon)].$$

$$\|(A_\varepsilon + R_0)^{-1}[(R_\varepsilon - R_0)u_*^0]\|_{H^1(\Omega)} \leq \delta \|(A_\varepsilon + R_0)^{-1}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} [\|u_*^\varepsilon - u_*^0\|_{H^1(\Omega)} + \kappa(\varepsilon) + \xi(\varepsilon)].$$

We choose $\delta > 0$ sufficiently small such that

$$\|u_*^\varepsilon - u_*^0\|_{H^1(\Omega)} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)] + \frac{1}{2}\|u_*^\varepsilon - u_*^0\|_{H^1(\Omega)}.$$

□

As a consequence of the Theorem 3.5 we have the following result.

Corollary 3.6. The family $\{\mathcal{E}_\varepsilon\}_{\varepsilon \in (0, \varepsilon_0]}$ is continuous at $\varepsilon = 0$; that is, if $\mathcal{E}_0 = \{u_*^{0,1}, \dots, u_*^{0,k}\}$ then for $\varepsilon > 0$ sufficiently small, $\mathcal{E}_\varepsilon = \{u_*^{\varepsilon,1}, \dots, u_*^{\varepsilon,k}\}$ and

$$\|u_*^{\varepsilon,i} - u_*^{0,i}\|_{H^1(\Omega)} \leq C[\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)], \quad i = 1, \dots, k.$$

4. CONVERGENCE OF LINEAR AND NONLINEAR SEMIGROUPS

In this section we obtain the rate of convergence of the linear and nonlinear semigroups. Here, we see the arising of the exponent $0 < l < 1$ in the rate (1.5). The main difficulty is the presence of singularities when we use the variation of the constants formula due the immersions between the fractional power spaces and the phase space L^2 . Our solution to this problem is to make an interpolation using appropriate exponents in order to improve the exponential decay of the linear semigroup.

Theorem 4.1. Let $\theta \in (0, \frac{1}{2}]$, then there exist $\alpha > 0$ and $C > 0$ independent of ε such that

$$(4.1) \quad \|e^{-A_\varepsilon t} - e^{-A_0 t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C e^{-\alpha t} [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)]^{2\theta} t^{-\frac{1+\beta}{2}-\theta},$$

where $t > 0$ and $\varepsilon \in (0, \varepsilon_0]$.

Proof. For each $\varepsilon \in [0, \varepsilon_0]$, the operators A_ε are self-adjoint and by Theorem 3.2, A_ε^{-1} converges to A_0^{-1} as $\varepsilon \rightarrow 0^+$. Thus we can define a appropriate closed rectifiable simple curve with trace in the resolvent set of A_0 such that, for $\alpha < \lambda_1^0$, the first eigenvalue of A_0 , and we can choose $0 < \varepsilon \leq \varepsilon_0$ sufficiently small and constant $C = C(\alpha) > 0$ such that

$$(4.2) \quad \|e^{-A_\varepsilon t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C e^{-\alpha t} t^{-\frac{1+\beta}{2}},$$

for $t > 0$ and $\varepsilon \in [0, \varepsilon_0]$, see [9] for more details on how to get the above estimates. Therefore we obtain

$$(4.3) \quad \begin{aligned} \|e^{-A_\varepsilon t} - e^{-A_0 t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} &\leq \|e^{-A_\varepsilon t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} + \|e^{-A_0 t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \\ &\leq C e^{-\alpha t} t^{-\frac{1+\beta}{2}}. \end{aligned}$$

On the other hand, the linear semigroup is given by

$$e^{-A_\varepsilon t} = \frac{1}{2\pi i} \int_{\omega} e^{\mu t} (\mu + A_\varepsilon)^{-1} d\mu,$$

where $\varepsilon \in [0, \varepsilon_0]$ and ω is the boundary of sector $\Sigma_{\nu, \phi}$ as in Theorem 3.2, oriented in such a way that the imaginary part of μ increases as μ runs in ω . Hence

$$\|e^{-A_\varepsilon t} - e^{-A_0 t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq \frac{1}{2\pi i} \int_{\omega} e^{\mu t} |(\mu + A_\varepsilon)^{-1} - (\mu + A_0)^{-1}| |d\mu|.$$

By (3.3) we obtain

$$(4.4) \quad \|e^{-A_\varepsilon t} - e^{-A_0 t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C e^{-\alpha t} [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)] t^{-1}.$$

Finally, interpolating (4.3) and (4.4) with $1 - 2\theta$ and 2θ , respectively, we obtain (4.1). \square

On the convergence of nonlinear semigroups we have:

Theorem 4.2. Let $u^\varepsilon, u^0 \in H^1(\Omega)$ and $\theta \in (0, \frac{1}{2}]$, then there exist constants $C > 0$ and $L > 0$ such that

$$(4.5) \quad \begin{aligned} & \|T_\varepsilon(t)u^\varepsilon - T_0(t)u^0\|_{H^1(\Omega)} \\ & \leq C e^{Lt} t^{-\frac{1+\beta}{2}-\theta} [\|u^\varepsilon - u^0\|_{H^1(\Omega)} + [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)]^{2\theta} + \kappa(\varepsilon) + \xi(\varepsilon)], \end{aligned}$$

for all $t \geq 0$ and $\varepsilon \in (0, \varepsilon_0]$.

Proof. Since the nonlinear semigroup is given by (2.5), we have

$$\begin{aligned} \|T_\varepsilon(t)u^\varepsilon - T_0(t)u^0\|_{H^1(\Omega)} & \leq \|e^{-A_\varepsilon t}(u^\varepsilon - u^0)\|_{H^1(\Omega)} + \|(e^{-A_\varepsilon t} - e^{-A_0 t})u^0\|_{H^1(\Omega)} \\ & \quad + \int_0^t \|e^{-A_\varepsilon(t-s)}(h^\varepsilon(T_\varepsilon(s)u^\varepsilon) - h^0(T_0(s)u^0))\|_{H^1(\Omega)} ds \\ & \quad + \int_0^t \|(e^{-A_\varepsilon(t-s)} - e^{-A_0(t-s)})h^0(T_0(s)u^0)\|_{H^1(\Omega)} ds. \end{aligned}$$

Consequently from (2.6) and (4.1) we have

$$\begin{aligned} \|T_\varepsilon(t)u^\varepsilon - T_0(t)u^0\|_{H^1(\Omega)} & \leq C[\|u^\varepsilon - u^0\|_{H^1(\Omega)} + (\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon))^{2\theta}] t^{-\frac{1}{2}-\theta} e^{-\alpha t} \\ & \quad + C \int_0^t (t-s)^{-\frac{1+\beta}{2}} e^{-\alpha(t-s)} \|T_\varepsilon(s)u^\varepsilon - T_0(s)u^0\|_{H^1(\Omega)} ds + C[\kappa(\varepsilon) + \xi(\varepsilon)], \end{aligned}$$

and using the singular Gronwall's inequality (see [8, Chapter 6]) we obtain (4.5). \square

Remark 4.3. The rate of convergence of eigenvalues, eigenfunctions and equilibrium points is better than rate of convergence of the nonlinear semigroup. This fact is related to the estimates that we obtained for the linear semigroup in the fractional power space and, we will see that the rate of convergence of attractors of problems (2.4) has a loss with respect to the rate of resolvent operators. A class of problems where the rate of convergence of the resolvent operators is the same rate of convergence of the attractors is presented in [9].

5. CONVERGENCE OF LOCAL UNSTABLE MANIFOLDS

In this section we use the general theory of invariant manifold to estimate the convergence of local unstable manifolds around an equilibrium point. We conclude the exponential attraction of the global attractors \mathcal{A}_ε in the phase space $H^1(\Omega)$ and we show how the nonlinear boundary conditions affects the estimates obtained in [1].

Let $u_*^\varepsilon \in \mathcal{E}_\varepsilon$ be an equilibrium for (2.4) and consider its linearization

$$(5.1) \quad \frac{dw^\varepsilon}{dt} + \overline{A}_\varepsilon w^\varepsilon = h^\varepsilon(w^\varepsilon + u_*^\varepsilon) - h^\varepsilon(u_*^\varepsilon) - (h^\varepsilon)'(u_*^\varepsilon)w^\varepsilon,$$

where $w^\varepsilon = u^\varepsilon - u_*^\varepsilon$ and $\overline{A}_\varepsilon = A_\varepsilon - (h^\varepsilon)'(u_*^\varepsilon)$. Let $\bar{\gamma}$ be a smooth, closed, simple, rectifiable curve in the resolvent set of $-\overline{A}_\varepsilon$, oriented counterclockwise evolving the first m positive eigenvalues. Thus we can choose an $\varepsilon > 0$ sufficiently small such that $\{\bar{\gamma}\} \subset \rho(-\overline{A}_\varepsilon)$ and then we define the bounded linear operator

$$\overline{Q}_\varepsilon^+ = \frac{1}{2\pi i} \int_{\bar{\gamma}} (\mu + \overline{A}_\varepsilon)^{-1} d\mu.$$

The operator \overline{A}_ε is self-adjoint and there is a $\alpha > 0$ and $M \geq 1$ such that, for all $0 \leq \varepsilon \leq \varepsilon_0$,

$$\begin{aligned} \|e^{-\overline{A}_\varepsilon t} \overline{Q}_\varepsilon^+\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} &\leq M e^{\alpha t}, \quad t \leq 0 \quad \text{and} \\ \|e^{-\overline{A}_\varepsilon t} (I - \overline{Q}_\varepsilon^+)\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} &\leq M t^{-\frac{1+\beta}{2}} e^{-\alpha t}, \quad t > 0. \end{aligned}$$

Moreover we can prove all convergence results of Section 3 for \overline{A}_ε rather than A_ε , for example, we have

$$\|e^{-\overline{A}_\varepsilon t} - e^{-\overline{A}_0 t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C e^{-\alpha t} [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)]^{2\theta} t^{-\frac{1+\beta}{2}-\theta}, \quad t > 0.$$

Using the decomposition $H^1(\Omega) = \overline{Q}_\varepsilon^+(H^1(\Omega)) \oplus (I - \overline{Q}_\varepsilon^+)(H^1(\Omega))$, the solution w^ε of (5.1) can be decomposed as $w^\varepsilon = v^\varepsilon + z^\varepsilon$, with $v^\varepsilon = \overline{Q}_\varepsilon^+ w^\varepsilon$ and $z^\varepsilon = (I - \overline{Q}_\varepsilon^+) w^\varepsilon$. If we define the operators $B_\varepsilon := \overline{A}_\varepsilon \overline{Q}_\varepsilon^+$ and $\tilde{A}_\varepsilon := \overline{A}_\varepsilon (I - \overline{Q}_\varepsilon^+)$, then we can rewrite (5.1) as

$$(5.2) \quad \begin{cases} \frac{dv^\varepsilon}{dt} + B_\varepsilon v^\varepsilon = H_\varepsilon(v^\varepsilon, z^\varepsilon), & t > 0, \\ \frac{dz^\varepsilon}{dt} + \tilde{A}_\varepsilon z^\varepsilon = G_\varepsilon(v^\varepsilon, z^\varepsilon), & t > 0, \end{cases}$$

where

$$H_\varepsilon(v^\varepsilon, z^\varepsilon) := \overline{Q}_\varepsilon^+ [h^\varepsilon(v^\varepsilon + z^\varepsilon + u_*^\varepsilon) - h^\varepsilon(u_*^\varepsilon) - (h^\varepsilon)'(u_*^\varepsilon)(v^\varepsilon + z^\varepsilon)]$$

and

$$G_\varepsilon(v^\varepsilon, z^\varepsilon) := (I - \overline{Q}_\varepsilon^+) [h^\varepsilon(v^\varepsilon + z^\varepsilon + u_*^\varepsilon) - h^\varepsilon(u_*^\varepsilon) - (h^\varepsilon)'(u_*^\varepsilon)(v^\varepsilon + z^\varepsilon)].$$

The maps H_ε and G_ε are continuously differentiable with $H_\varepsilon(0, 0) = 0 = G_\varepsilon(0, 0) \in H^{-\beta}(\Omega)$ and $H'_\varepsilon(0, 0) = 0 = G'_\varepsilon(0, 0) \in \mathcal{L}(H^1(\Omega), H^{-\beta}(\Omega))$. Hence, given $\rho_o > 0$, there are $0 < \bar{\varepsilon} = \bar{\varepsilon}_{\rho_o} \leq \varepsilon_0$ and $\delta = \delta_{\rho_o} > 0$ such that if $\|v\|_{\overline{Q}_\varepsilon^+ H^1(\Omega)} + \|z\|_{H^1(\Omega)} < \delta$ and $\varepsilon \leq \varepsilon_0$, then

$$(5.3) \quad \|H_\varepsilon(v, z)\|_{\overline{Q}_\varepsilon^+ (H^1(\Omega))} \leq \rho_o \quad \text{and} \quad \|G_\varepsilon(v, z)\|_{H^{-\beta}(\Omega)} \leq \rho_o;$$

$$(5.4) \quad \|H_\varepsilon(v, z) - H_\varepsilon(\bar{v}, \bar{z})\|_{\overline{Q}_\varepsilon^+(H^1(\Omega))} \leq \rho_o(\|v - \bar{v}\|_{\overline{Q}_\varepsilon^+(H^1(\Omega))} + \|z - \bar{z}\|_{H^1(\Omega)});$$

$$(5.5) \quad \|G_\varepsilon(v, z) - G_\varepsilon(\bar{v}, \bar{z})\|_{H^{-\beta}(\Omega)} \leq \rho_o(\|v - \bar{v}\|_{\overline{Q}_\varepsilon^+(H^1(\Omega))} + \|z - \bar{z}\|_{H^1(\Omega)}).$$

Theorem 5.1. Given $D > 0$ and $\Delta > 0$, $\vartheta \in (0, 1)$ and $\bar{\rho} > 0$ such that

$$(5.6) \quad \begin{aligned} \rho_o M \alpha^{-\frac{1}{2}} \Gamma\left(\frac{1}{2}\right) &\leq D \quad \rho_o M \Gamma\left(\frac{1}{2}\right) \frac{M(1+\Delta)}{(2\alpha - \rho M(1+\Delta))^{\frac{1}{2}}} \leq \Delta \\ \rho_o M \alpha^{-\frac{1}{2}} \Gamma\left(\frac{1}{2}\right) &\left[1 + \frac{\rho_o M(1+\Delta)\alpha^{-\frac{1}{2}}}{(2\alpha - \rho_o M(1+\Delta))^{\frac{1}{2}}}\right] \leq \vartheta \end{aligned}$$

are satisfied for all $\rho_o \in (0, \bar{\rho})$. Assume that H_ε and G_ε satisfies (5.3)-(5.5), with $0 < \rho_o \leq \bar{\rho}$ for all $(v, z) \in \overline{Q}_\varepsilon^+ H^1(\Omega) \times (I - \overline{Q}_\varepsilon^+) H^1(\Omega)$. Then, there exists $s_\varepsilon^* : \overline{Q}_\varepsilon^+ H^1(\Omega) \rightarrow (I - \overline{Q}_\varepsilon^+) H^1(\Omega)$ such that the local unstable manifold of u_ε^* is given as the graph of the map s_ε^* ,

$$W^u(u_\varepsilon^*) = \{(v, z) \in H^1(\Omega); z = s_\varepsilon^*(v), v \in \overline{Q}_\varepsilon^+ H^1(\Omega)\}.$$

The map s_ε^* satisfy

$$\|s_\varepsilon^*\| := \sup_{v \in \overline{Q}_\varepsilon^+(H^1(\Omega))} \|s_\varepsilon^*(v)\|_{H^1(\Omega)} \leq D, \quad \|s_\varepsilon^*(v) - s_\varepsilon^*(\tilde{v})\|_{H^1(\Omega)} \leq \Delta \|v - \tilde{v}\|_{\overline{Q}_\varepsilon^+(H^1(\Omega))},$$

and there is $C > 0$ independent of ε and $0 < \theta < 1$ such that

$$(5.7) \quad \|s_\varepsilon^* - s_0^*\| \leq C[(\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon))^\theta + \kappa(\varepsilon) + \xi(\varepsilon)]$$

Furthermore, given $0 < \gamma < \alpha$, there is $0 < \rho_1 \leq \bar{\rho}$ and $C > 0$, independent of ε , such that, for any solution $[t_0, \infty) \ni t \mapsto (v^\varepsilon(t), z^\varepsilon(t)) \in H^1(\Omega)$ of (5.2),

$$(5.8) \quad \|z^\varepsilon(t) - s_\varepsilon^*(v^\varepsilon(t))\|_{H^1(\Omega)} \leq C e^{-\gamma(t-t_0)} \|z^\varepsilon(t_0) - s_\varepsilon^*(v^\varepsilon(t_0))\|_{H^1(\Omega)}, \text{ for all } t \geq t_0.$$

Proof. The proof is standard and was made in several papers, see e.g. [1, 8, 13]. In [1] it was shown how to estimate the convergence (5.7). Once we have established the spaces and estimates, we outline what was done in these works.

Consider the set

$$\Sigma_\varepsilon = \left\{ s : \overline{Q}_\varepsilon^+(H^1(\Omega)) \rightarrow (I - \overline{Q}_\varepsilon^+)(H^1(\Omega)) : \|s\| \leq D, \|s(v) - s(\tilde{v})\|_{H^1(\Omega)} \leq \Delta \|v - \tilde{v}\|_{\overline{Q}_\varepsilon^+ H^1(\Omega)} \right\}.$$

It is not difficult to see that $(\Sigma_\varepsilon, \|\cdot\|)$ is a complete metric space.

Given $s_\varepsilon \in \Sigma_\varepsilon$ and $\Theta \in \overline{Q}_\varepsilon(H^1(\Omega))$, denote by $v^\varepsilon(t) = \psi(t, \tau, \Theta, s_\varepsilon)$ the solution of

$$\begin{cases} \frac{dv^\varepsilon}{dt}(t) + B_\varepsilon v^\varepsilon(t) = H_\varepsilon(v^\varepsilon(t), s_\varepsilon^*(v^\varepsilon(t))), & t < \bar{\tau} \\ v^\varepsilon(\bar{\tau}) = \Theta. \end{cases}$$

Define the map $\Psi_\varepsilon : \Sigma_\varepsilon \rightarrow \Sigma_\varepsilon$ by

$$\Psi_\varepsilon(s_\varepsilon)\Theta = \int_{-\infty}^{\bar{\tau}} e^{-\tilde{A}_\varepsilon(\bar{\tau}-s)} G_\varepsilon(v^\varepsilon(s), s_\varepsilon(v^\varepsilon(s))) ds.$$

Note that, from (5.3) and (5.6), we have $\|\Psi(s_\varepsilon)(\Theta)\|_{H^1(\Omega)} \leq D$.

Moreover, if $\Theta, \tilde{\Theta} \in \overline{Q}_\varepsilon(H^1(\Omega))$, $s_\varepsilon, \tilde{s}_\varepsilon \in \Sigma_\varepsilon$, $v^\varepsilon(t) = \psi(t, \bar{\tau}, \Theta, s_\varepsilon)$ and $\tilde{v}^\varepsilon(t) = \psi(t, \bar{\tau}, \Theta, \tilde{s}_\varepsilon)$, it is easy to see that

$$\phi(t) \leq M\|\Theta - \tilde{\Theta}\|_{H^1(\Omega)} + M\rho_o(1 + \Delta) \int_t^{\bar{\tau}} \phi(s)ds + M\rho_o\alpha^{-1}\|s_\varepsilon - \tilde{s}_\varepsilon\|$$

where $\phi(t) = e^{-\alpha(t-\bar{\tau})}\|v^\varepsilon(t) - \tilde{v}^\varepsilon(t)\|_{H^1(\Omega)}$ and using Gronwall's inequality

$$\|v^\varepsilon(t) - \tilde{v}^\varepsilon(t)\|_{H^1(\Omega)} \leq M \left(\|\Theta - \tilde{\Theta}\|_{H^1(\Omega)} + \rho_o\alpha^{-1}\|s_\varepsilon - \tilde{s}_\varepsilon\| \right) e^{(\alpha - M\rho_o(1+\Delta))(t-\bar{\tau})}.$$

From this we obtain that

$$\begin{aligned} \|\Psi(s_\varepsilon)(\Theta) - \Psi(\tilde{s}_\varepsilon)(\tilde{\Theta})\|_{H^1(\Omega)} &\leq \rho_o M \alpha^{-\frac{1}{2}} \Gamma\left(\frac{1}{2}\right) \left[1 + \frac{\rho_o M (1 + \Delta) \alpha^{-\frac{1}{2}}}{(2\alpha - \rho_o M (1 + \Delta))^{\frac{1}{2}}} \right] \|s_\varepsilon - \tilde{s}_\varepsilon\| + \\ &+ \frac{\rho_o M^2 (1 + \Delta)}{(2\alpha - \rho_o M (1 + \Delta))^{\frac{1}{2}}} \Gamma\left(\frac{1}{2}\right) \|\Theta - \tilde{\Theta}\|_{\overline{Q}_\varepsilon^+ H^1(\Omega)}. \end{aligned}$$

and

$$\|\Psi(s_\varepsilon)(\Theta) - \Psi(\tilde{s}_\varepsilon)(\tilde{\Theta})\|_{H^1(\Omega)} \leq \Delta \|\Theta - \tilde{\Theta}\|_{\overline{Q}_\varepsilon^+ H^1(\Omega)} + \vartheta \|s_\varepsilon - \tilde{s}_\varepsilon\|.$$

Hence, $\Psi_\varepsilon : \Sigma_\varepsilon \rightarrow \Sigma_\varepsilon$ is a contraction. Therefore, there a fixed point $s_\varepsilon^* = \Psi(s_\varepsilon^*)$ in Σ_ε .

Now, we prove that $\{(v^\varepsilon, s_\varepsilon^*(v^\varepsilon)); v \in \overline{Q}_\varepsilon^+ H^1(\Omega)\}$ is invariant for (5.2). Let $(v_0^\varepsilon, z_0^\varepsilon) \in W^u(u_\varepsilon^*)$, $z_0^\varepsilon = s_\varepsilon^*(v_0^\varepsilon)$. Denote by $v_\varepsilon^*(t)$ the solution of the initial value problems

$$(5.9) \quad \begin{cases} \frac{dv^\varepsilon}{dt} + B_\varepsilon v^\varepsilon = H_\varepsilon(v^\varepsilon, s_\varepsilon^*(v^\varepsilon)) \\ v^\varepsilon(0) = v_0^\varepsilon. \end{cases}$$

This defines a curve $(v_\varepsilon^*(t), s_\varepsilon^*(v_\varepsilon^*(t))) \in W^u(u_\varepsilon^*)$, $t \in \mathbb{R}$. Also, the only solution of

$$z_t^\varepsilon + \tilde{A}_\varepsilon z^\varepsilon = G_\varepsilon(v_\varepsilon^*(t), s_\varepsilon^*(v_\varepsilon^*(t)))$$

which remains bounded as $t \rightarrow -\infty$ must be

$$z_\varepsilon^*(t) = \int_{-\infty}^t e^{\tilde{A}_\varepsilon(t-s)} G_\varepsilon(v_\varepsilon^*(s), s_\varepsilon^*(v_\varepsilon^*(s))) ds = s_\varepsilon^*(v_\varepsilon^*(t)).$$

This proves the invariance of the graph of s_ε^* . To prove that the graph of s_ε^* is the unstable manifold assume the exponential attraction of the graph of s_ε^* uniformly in ε ; i.e., if $u^\varepsilon(t) = z^\varepsilon(t) + v^\varepsilon(t)$ is a solution of (5.2) with $v^\varepsilon(t) = \overline{Q}_\varepsilon u^\varepsilon(t)$. If, given $\gamma < \alpha$, there exists $\rho_1 > 0$ such that (5.8) holds for any $0 < \rho_o \leq \rho_1$, it is easy to see that, when $z^\varepsilon(t)$ remains bounded as $t \rightarrow -\infty$, it follows that (making $t_0 \rightarrow -\infty$ in (5.8)) that $z^\varepsilon(t) = s_\varepsilon^*(v^\varepsilon(t))$ for all $t \in \mathbb{R}$.

The proof of (5.8) can be carried out as [10], using the singular Gronwall's inequality.

Finally, we obtain the rate of convergence (5.7). For $\Theta \in \overline{Q}_\varepsilon^+(H_0^1(\Omega))$, we have

$$\begin{aligned} \|s_*^\varepsilon(\Theta) - s_*^0(\Theta)\|_{H^1(\Omega)} &\leq \int_{-\infty}^{\bar{\tau}} \|e^{-\tilde{A}_\varepsilon(\tau-r)} G_\varepsilon(v^\varepsilon, s_*^\varepsilon(v^\varepsilon)) - e^{-\tilde{A}_\varepsilon(\tau-r)} G_\varepsilon(v^0, s_*^0(v^0))\|_{H^1(\Omega)} dr \\ &\quad + \int_{-\infty}^{\bar{\tau}} \|e^{-\tilde{A}_\varepsilon(\tau-r)} G_\varepsilon(v^0, s_*^0(v^0)) - e^{-\tilde{A}_0(\tau-r)} G_0(v^0, s_*^0(v^0))\|_{H^1(\Omega)} dr \\ &\quad + \int_{-\infty}^{\bar{\tau}} \|e^{-\tilde{A}_\varepsilon(\tau-r)} G_0(v^0, s_*^0(v^0)) - e^{-\tilde{A}_0(\tau-r)} G_0(v^0, s_*^0(v^0))\|_{H^1(\Omega)} dr. \end{aligned}$$

Since we can estimate $G_\varepsilon - G_0$ by $Q_\varepsilon^+ - Q_0^+$ these integrals are estimated as in the Theorem 4.2, and using the singular Gronwall's inequality. Repeating the argument for estimate $v^\varepsilon - v^0$ we obtain (5.7). \square

Remark 5.2. The projection $\overline{Q}_\varepsilon^+$ has finite rank hence the unstable manifold can be considered as a finite dimensional object, in fact $W^u(u_*^\varepsilon) \approx \mathbb{R}^m$ and the semigroup restrict to this manifold is conjugate to that generated by a vector field in \mathbb{R}^m that is, it is given by ordinary equation (5.9). Notice that the rate of convergence of the nonlinear boundary conditions is added in the rate of convergence of unstable manifolds.

6. CONVERGENCE OF GLOBAL ATTRACTORS

In this section we prove the main result of this work. We used all previous results to put the problem (1.1) in the conditions of Corollary A.3 in the Appendix A. Here, we consider the dimension $N \geq 2$. The scalar case will be addressed in the next section.

Let X be a Banach space. We denote by $\text{dist}_H(A, B)$ the Hausdorff semidistance between $A, B \subset X$, defined as

$$\text{dist}_H(A, B) = \sup_{a \in A} \inf_{b \in B} \|a - b\|_X,$$

and we define the symmetric Hausdorff metric by

$$d_H(A, B) = \max\{\text{dist}_H(A, B), \text{dist}_H(B, A)\}.$$

Definition 6.1. We say that a family $\{\mathcal{A}_\varepsilon\}_{\varepsilon \in \Lambda}$ of subsets of X is continuous at ε_0 if

$$d_H(\mathcal{A}_\varepsilon, \mathcal{A}_{\varepsilon_0}) \xrightarrow{\varepsilon \rightarrow \varepsilon_0} 0.$$

Theorem 6.2. The family of attractors $\{\mathcal{A}_\varepsilon\}_{\varepsilon \in [0, \varepsilon_0]}$ of (2.4) for $N \geq 2$ is continuous at $\varepsilon = 0$. Moreover we have the following rate of convergence

$$(6.1) \quad d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)]^l, \quad \text{with } 0 < l < 1.$$

Proof. It follows from Theorem 4.2 that for $u, v \in H^1(\Omega)$,

$$\text{dist}_H(T_\varepsilon(t)u, T_0(t)v) \leq Ce^{Lt}(\|u - v\|_{H^1(\Omega)} + \delta(\varepsilon)), \quad t \geq 1,$$

where $\delta(\varepsilon) = [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)]^{2\theta} + \kappa(\varepsilon) + \xi(\varepsilon)$, with $C, L > 0$ and $0 < \theta \leq \frac{1}{2}$.

By Theorem 2.2 the set $D = \cup_{\varepsilon \in [0, \varepsilon_0]} \mathcal{A}_\varepsilon$ is uniformly bounded and by (5.8) the attractor \mathcal{A}_ε attracts exponentially, i.e.

$$\sup_{\varepsilon \in [0, \varepsilon_0]} \text{dist}_H(T_\varepsilon(t)D, \mathcal{A}_\varepsilon) \leq \Theta(t), \quad t \geq 1$$

where $\Theta(t) = Ce^{-\gamma(t-t_0)}$.

Now applying the Corollary A.3 in the Appendix A we get

$$d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)]^{2\theta} + \kappa(\varepsilon) + \xi(\varepsilon)^{\frac{\gamma}{\gamma+L}},$$

which implies (6.1) taking $\theta = \frac{1}{2}$ and $l = \frac{\gamma}{\gamma+L}$. \square

7. SCALAR REACTION DIFFUSION PROBLEM

In this section we study the problem (2.4) for the case $N = 1$. In this scalar situation we will have a family of Morse Smale problems with gap condition in the eigenvalues. This property allows us to obtain a finite dimensional invariant manifold and then we reduce the flows in the attractors to ordinary differential equations in the Euclidian Space \mathbb{R}^M , where $M = m$ is defined in (3.5). Once this is done we will use the results of the Appendix B to obtain a better rate of convergence, than the one of the previous section, for the convergence of the attractors of (2.4).

Consider the scalar parabolic equations

$$(7.1) \quad \begin{cases} \partial_t u^\varepsilon - \partial_x(p_\varepsilon(x)\partial_x u^\varepsilon) + (\lambda + V_\varepsilon(x))u^\varepsilon = f^\varepsilon(u^\varepsilon), & 0 < x < 1, \quad t > 0, \\ \partial_x u^\varepsilon + (\lambda + b_\varepsilon(x))u^\varepsilon = g^\varepsilon(u^\varepsilon), & x \in \{0, 1\}, \end{cases}$$

where $\varepsilon \in [0, \varepsilon_0]$, and $p_\varepsilon, V_\varepsilon, b_\varepsilon, f^\varepsilon$ and g^ε satisfies (1.2) (1.3) and (1.4).

We can rewrite (7.1) abstractly as in (2.4), where the nonlinearity is given by $h^\varepsilon = f_\Omega^\varepsilon + g_\Gamma^\varepsilon : H^1(\Omega) \rightarrow H^{-\beta}(\Omega)$, for $-\beta \in [-\frac{1}{2}, -\frac{1}{4}]$ and $\Omega = (0, 1)$. Thus, with standard conditions on the nonlinearities, (2.4) is well posed in $H^1(\Omega)$ and the solutions through u_0^ε satisfies (2.5). Moreover, the nonlinear semigroup has global attractor \mathcal{A}_ε in $H^1(\Omega)$, such that $\overline{\cup_{\varepsilon \in [0, \varepsilon_0]} \mathcal{A}_\varepsilon}$ is compact.

Moreover the scalar equation (7.1) is a Morse-Smale problem (see [14]) in the sense that the unstable and stable local manifolds (of different equilibrium points) has transversal intersection which implies geometrical structural stability of diagram phase space. Consequently, the nonlinear semigroups are Morse-Smale semigroup for $\varepsilon \in [0, \varepsilon_0]$.

Theorem 7.1. For each $w_0 \in \overline{\cup_{\varepsilon \in [0, \varepsilon_0]} \mathcal{A}_\varepsilon}$, there is constant $C > 0$ independent of ε such that

$$\|T_\varepsilon(1)w_0 - T_0(1)w_0\|_{H^1(\Omega)} \leq C\delta(\varepsilon)|\log(\delta(\varepsilon))|,$$

where $\delta(\varepsilon) = \|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)$.

Proof. As in the Theorem 4.1 we have for, $\varepsilon \in [0, \varepsilon_0]$,

$$\|e^{-A_\varepsilon t} - e^{-A_0 t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C e^{-\alpha t} t^{-\frac{1+\beta}{2}}$$

and

$$\|e^{-A_\varepsilon t} - e^{-A_0 t}\|_{\mathcal{L}(H^{-\beta}(\Omega), H^1(\Omega))} \leq C e^{-\alpha t} [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)] t^{-1}.$$

Notice that the terms $t^{-\frac{1+\beta}{2}}$ and t^{-1} in the the estimates above originates a singularity in the formula of the variation of the constants. This is the main difficulty in estimating the nonlinear semigroups. In the Theorem 4.2 we performed an interpolation of these terms together with the rate of convergence of resolvent operators which resulted in the considerable loss in the rate of convergence of attractors (6.1). In the situation where the limiting problems is Morse-Smale, the authors in [4] had the same problem, however they used the following estimate (placed in our context). If we denote $l_\varepsilon(t) = \min\{t^{-\frac{1+\beta}{2}}, \|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)\} t^{-1}$, then

$$\int_{-\infty}^{\bar{\tau}} l_\varepsilon(\bar{\tau} - r) e^{-\alpha(\bar{\tau} - r)} dr \leq C [\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)] \log(\|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon)).$$

Since the nonlinear semigroup is given by (2.5), then for $0 < t \leq 1$, we have

$$\begin{aligned} \|T_\varepsilon(t)w_0 - T_0(t)w_0\|_{H^1(\Omega)} &\leq \|(e^{-A_\varepsilon t} - e^{-A_0 t})w_0\|_{H^1(\Omega)} \\ &\quad + \int_0^t \|e^{-A_\varepsilon(t-s)} f(T_\varepsilon(s)w_0) - e^{-A_0(t-s)} f(T_0(s)w_0)\|_{H^1(\Omega)} ds, \end{aligned}$$

Now, as in Theorem 4.2, we obtain

$$\|T_\varepsilon(t)w_0 - T_0(t)w_0\|_{H^1(\Omega)} \leq C\delta(\varepsilon)|\log(\delta(\varepsilon))| + C\delta(\varepsilon)|\log(\delta(\varepsilon))|e^{Kt},$$

where $K > 0$. Now the result follows taking $t = 1$. \square

We saw in the Section 3 that or each $\varepsilon \in [0, \varepsilon_0]$, the spectrum $\sigma(-A_\varepsilon)$ of $-A_\varepsilon$, ordered and counting multiplicity is given by $\dots - \lambda_m^\varepsilon < -\lambda_{m-1}^\varepsilon < \dots < -\lambda_0^\varepsilon$. Moreover is true the following gap condition

$$|\lambda_m^\varepsilon - \lambda_{m-1}^\varepsilon| \rightarrow \infty \quad \text{as} \quad m \rightarrow \infty.$$

This property enable us to find a finite dimension invariant manifold as well as in [3] and [17].

Theorem 7.2. For sufficiently large m and ε small there is an invariant manifold \mathcal{M}_ε for the problem (2.4) given by

$$\mathcal{M}_\varepsilon = \{u^\varepsilon \in H^1(\Omega); u^\varepsilon = Q_\varepsilon u^\varepsilon + s_*^\varepsilon(Q_\varepsilon u^\varepsilon)\}, \quad \varepsilon \in [0, \varepsilon_0],$$

where, Q_ε is the spectral projection and $s_*^\varepsilon : Y_\varepsilon \rightarrow Z_\varepsilon$ is a Lipschitz continuous map satisfying

$$\|s_*^\varepsilon - s_*^0\| = \sup_{v \in Y_\varepsilon} \|s_*^\varepsilon(v) - s_*^0(v)\|_{H^1(\Omega)} \leq C\delta(\varepsilon)|\log(\delta(\varepsilon))|,$$

where $\delta(\varepsilon) = \|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)$, $C > 0$ is constant independent of ε and $Y_\varepsilon = Q_\varepsilon H^1(\Omega)$ and $Z_\varepsilon = (I - Q_\varepsilon)H^1(\Omega)$. The invariant manifold \mathcal{M}_ε is exponentially attracting and the attractor \mathcal{A}_ε of the problem (2.4) lies in \mathcal{M}_ε . The flow on \mathcal{A}_ε is given by

$$u^\varepsilon(t) = v^\varepsilon(t) + s_*^\varepsilon(v^\varepsilon(t)), \quad t \in \mathbb{R},$$

where $v^\varepsilon(t)$ satisfy

$$(7.2) \quad \frac{dv^\varepsilon}{dt} + A_\varepsilon^+ v^\varepsilon = Q_\varepsilon h^\varepsilon(v^\varepsilon + s_*^\varepsilon(v^\varepsilon(t))).$$

Proof. The proof follows the same steps of the Theorem 5.1, see [3] and [17] for more details. \square

Thus the flow in \mathcal{A}_ε is given by ODE (7.2). Since Q_ε has finite rank we can consider $v^\varepsilon \in \mathbb{R}^M$ and $H_\varepsilon(v^\varepsilon) = Q_\varepsilon h^\varepsilon(v^\varepsilon + s_*^\varepsilon(v^\varepsilon(t)))$ a continuously differentiable map in \mathbb{R}^M . For each $\varepsilon \in [0, \varepsilon_0]$, we denote $\tilde{T}_\varepsilon = \tilde{T}_\varepsilon(1)$, where $\tilde{T}_\varepsilon(\cdot)$ is the semigroup generated by solution $v^\varepsilon(\cdot)$ of (7.2) in \mathbb{R}^M . We have the following convergences

$$(7.3) \quad \|\tilde{T}_\varepsilon - \tilde{T}_0\|_{C^1(\mathbb{R}^M, \mathbb{R}^M)} \xrightarrow{\varepsilon \rightarrow 0^+} 0 \quad \text{and} \quad \|\tilde{T}_\varepsilon - \tilde{T}_0\|_{L^\infty(\mathbb{R}^M, \mathbb{R}^M)} \leq C\delta(\varepsilon)|\log(\delta(\varepsilon))|,$$

where the last estimate can be proved as well as in Theorem 7.1.

Therefore we have a Morse-Smale semigroup in \mathbb{R}^M and using techniques of shadowing in the Appendix B, we are ready to prove the main result of this section.

Theorem 7.3. Let \mathcal{A}_ε , $\varepsilon \in [0, \varepsilon_0]$, be the global attractor for the problem (2.4). Then there is constant $C > 0$ independent of ε such that

$$d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq C\delta(\varepsilon)|\log(\delta(\varepsilon))|,$$

where $\delta(\varepsilon) = \|p_\varepsilon - p_0\|_{L^\infty(\Omega)} + \eta(\varepsilon) + \tau(\varepsilon) + \kappa(\varepsilon) + \xi(\varepsilon)$.

Proof. For each $\varepsilon \in [0, \varepsilon_0]$ we denote $T_\varepsilon = T_\varepsilon(1)$. Given $u^\varepsilon \in \mathcal{A}_\varepsilon$, by invariance there is $w^\varepsilon \in \mathcal{A}_\varepsilon$ such that $u^\varepsilon = T_\varepsilon w^\varepsilon$ so we can write $w^\varepsilon = Q_\varepsilon w^\varepsilon + s_*^\varepsilon(Q_\varepsilon w^\varepsilon)$, where $Q_\varepsilon w^\varepsilon \in \bar{\mathcal{A}}_\varepsilon$ with $\bar{\mathcal{A}}_\varepsilon = Q_\varepsilon \mathcal{A}_\varepsilon$ the projected attractor in \mathbb{R}^M . Thus

$$\begin{aligned} \|u^\varepsilon - u^0\|_{H^1(\Omega)} &= \|T_\varepsilon w^\varepsilon - T_0 w^0\|_{H^1(\Omega)} \leq \|T_\varepsilon w^\varepsilon - T_\varepsilon w^0\|_{H^1(\Omega)} + \|T_\varepsilon w^0 - T_0 w^0\|_{H^1(\Omega)} \\ &\leq C\delta(\varepsilon)|\log(\delta(\varepsilon))|. \end{aligned}$$

But

$$\begin{aligned}
\|w^\varepsilon - w^0\| &= \|Q_\varepsilon w^\varepsilon - Q_0 w^0\|_{\mathbb{R}^M} + \|s_*^\varepsilon(Q_\varepsilon w^\varepsilon) - s_*^0(Q_0 w^0)\|_{H^1(\Omega)} \\
&\leq \|Q_\varepsilon w^\varepsilon - Q_0 w^0\|_{\mathbb{R}^M} + \|s_*^\varepsilon(Q_\varepsilon w^\varepsilon) - s_*^\varepsilon(Q_0 w^0)\|_{H^1(\Omega)} + \|s_*^\varepsilon(Q_0 w^0) - s_*^0(Q_0 w^0)\|_{H^1(\Omega)} \\
&\leq C\|Q_\varepsilon w^\varepsilon - Q_0 w^0\|_{\mathbb{R}^M} + C\delta(\varepsilon)|\log(\delta(\varepsilon))|,
\end{aligned}$$

which implies

$$d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq d_H(\bar{\mathcal{A}}_\varepsilon, \bar{\mathcal{A}}_0) + C\delta(\varepsilon)|\log(\delta(\varepsilon))|.$$

The result follows by (7.3) and Theorem B.6. \square

8. CLOSING REMARKS AND ACKNOWLEDGMENTS

We conclude by saying that the rate of convergence of nonlinear boundary conditions affects the rate of attractors in a pleasant way, that is, their speeds are added to the final speed of the entire dynamics and does not represent a loss. In fact, the exponent l obtained in the Theorem 6.2 is the same as that obtained in the paper [1]. This fact is surprising because the variational formulation for problems with nonlinear boundary conditions involves considering the dual space $H^{-\beta}(\Omega)$ and consequently the immersion results between fractional power spaces must be estimated, this can be seen in the linear and nonlinear semigroup estimates where the exponent $-\beta$ of these space is explicitly visible. Our result directs a path to the problem of the optimal rate, in fact the works [9] and [10] seemed to show us that phase space is an obstruction to the optimal rate but as we have just seen the dual spaces have a positive role in this question. The problem of optimal rate is still open: would the rate of resolvent operator be the same as that of attractors?

The authors would like to thank the professor José M. Arrieta for pointing out some misprints and making valuable suggestions. The authors are grateful to the anonymous referee for carefully reading and valuable comments.

APPENDIX A. GENERAL THEORY OF RATE OF CONVERGENCE OF ATTRACTORS

Let $\{T_\varepsilon(\cdot)\}_{\varepsilon \in \Lambda}$ be a family of semigroups on a Banach space X , where Λ is a topology space. We will assume that the family $\{T_\varepsilon(\cdot)\}_{\varepsilon \in \Lambda}$ converges in some appropriate sense to $T_{\varepsilon_0}(\cdot)$ as $\varepsilon \rightarrow \varepsilon_0$ in Λ .

Definition A.1. If $T_\varepsilon(\cdot)$ is a family of semigroups with attractors \mathcal{A}_ε , then we say that $\{\mathcal{A}_\varepsilon\}_{\varepsilon \in \Lambda}$ is equi-attracting if

$$\sup_{\varepsilon \in \Lambda} \text{dist}_H(T_\varepsilon(t)B, \mathcal{A}_\varepsilon) \xrightarrow{t \rightarrow \infty} 0,$$

for each bounded subset B of X .

We now see that equi-attraction implies the continuity of attractors, we can also obtain a rate of convergence of attractors with respect to the underlying parameters.

Theorem A.2. Let $T_\varepsilon(\cdot)$ be a family of semigroups with attractors \mathcal{A}_ε and set $D = \bigcup_{\varepsilon \in \Lambda} \mathcal{A}_\varepsilon$. If there is a strictly decreasing function $\Theta : [t_0, \infty) \rightarrow (0, \infty)$ such that

$$\sup_{\varepsilon \in \Lambda} \text{dist}_H(T_\varepsilon(t)D, \mathcal{A}_\varepsilon) \leq \Theta(t), \quad t \geq t_0$$

and, there are $C, L > 0$ and a function $\delta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow \varepsilon_0$ such that

$$\text{dist}_H(T_\varepsilon(t)x, T_0(t)y) \leq Ce^{Lt}(\|x - y\|_X + \delta(\varepsilon)), \quad t \geq t_0.$$

Then

$$d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq \min_{\nu \in \Theta([0, \infty))} 2\{Ce^{L\Theta^{-1}(\nu)}\delta(\varepsilon) + \nu\}.$$

Proof. For $t \geq t_0$ we have

$$\begin{aligned} \text{dist}_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) &\leq \text{dist}_H(T_\varepsilon(t)\mathcal{A}_\varepsilon, T_0(t)\mathcal{A}_\varepsilon) + \text{dist}_H(T_0(t)\mathcal{A}_\varepsilon, \mathcal{A}_0) \\ &\leq \sup_{x \in \mathcal{A}_\varepsilon} \text{dist}_H(T_\varepsilon(t)x, T_0(t)x) + \text{dist}_H(T_0(t)\mathcal{A}_\varepsilon, \mathcal{A}_0) \\ &\leq Ce^{Lt}\delta(\varepsilon) + \Theta(t). \end{aligned}$$

Given $\nu \in \Theta([0, \infty))$ and $t = \Theta^{-1}(\nu)$, we get

$$\text{dist}_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq Ce^{L\Theta^{-1}(\nu)}\delta(\varepsilon) + \nu.$$

□

In particular, if the equi-attraction is of exponential order, then we have optimal rate of convergence.

Corollary A.3. Assume that the conditions of Theorem A.2 are satisfied with $\Theta(t) = ce^{-\gamma t}$ for some $c \geq 1$, $\gamma > 0$ and for all $t \in [t_0, \infty)$. Then there is constant $\bar{c} > 0$ independent of ε , such that

$$d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq \bar{c}\delta(\varepsilon)^{\frac{\gamma}{\gamma+L}}.$$

Proof. Take $\Theta^{-1}(\nu) = \log(\frac{c}{\nu})^{\frac{1}{\gamma}}$, we obtain

$$(A.1) \quad d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq 2 \min_{\nu \in (0, c]} \left[\delta(\varepsilon) \left(\frac{c}{\nu} \right)^{\frac{L}{\gamma}} + \nu \right].$$

The minimum of the right-hand side of (A.1) occurs when $\nu = c \left(\frac{L}{\gamma} \right)^{\frac{\gamma}{\gamma+L}} \delta(\varepsilon)^{\frac{\gamma}{\gamma+L}}$. Since the left-hand side of (A.1) is independent of ν , it follows that

$$d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq 2c \left[\left(\frac{L}{\gamma} \right)^{\frac{-L}{\gamma+L}} + \left(\frac{L}{\gamma} \right)^{\frac{\gamma}{\gamma+L}} \right] \delta(\varepsilon)^{\frac{\gamma}{\gamma+L}}.$$

□

Example A.4. Consider the following problem

$$\begin{cases} \varepsilon \frac{d^2 x}{dt^2} + \frac{dx}{dt} = \mu x + f(x), & x \in \mathbb{R}^N \\ x(0) = x_0 \in \mathbb{R}^N \\ \frac{dx}{dt}(0) = v_0 \in \mathbb{R}^N. \end{cases}$$

Assume that $\varepsilon \in [0, 1]$, $\mu \geq 1$, $f : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a C^1 -function which is globally Lipschitz, globally bounded and with symmetric Jacobian matrix at every point. If we write the above equation in the form of a system with variables x and $v = \varepsilon \frac{dx}{dt}$ we have

$$(A.2) \quad \begin{cases} \frac{d}{dt} \begin{bmatrix} x \\ v \end{bmatrix} = \begin{bmatrix} 0 & \frac{1}{\varepsilon} I \\ -\mu I & -\frac{1}{\varepsilon} I \end{bmatrix} \begin{bmatrix} x \\ v \end{bmatrix} + \begin{bmatrix} 0 \\ f(x) \end{bmatrix}, & \begin{bmatrix} x \\ v \end{bmatrix} \in \mathbb{R}^N \times \mathbb{R}^N, \\ \begin{bmatrix} x \\ v \end{bmatrix}(0) = \begin{bmatrix} x_0 \\ v_0 \end{bmatrix} \in \mathbb{R}^N \times \mathbb{R}^N. \end{cases}$$

The solutions of (A.2) are globally defined and the solution operator family $T_\varepsilon(\cdot)$ defines a semigroup in $Z = \mathbb{R}^N \times \mathbb{R}^N$ which has attractor \mathcal{A}_ε . The problem (A.2) can be rewritten as

$$\begin{cases} \frac{d}{dt} \begin{bmatrix} I & I \\ -\varepsilon \mu I & 0 \end{bmatrix} \begin{bmatrix} x \\ v \end{bmatrix} = - \begin{bmatrix} x \\ v \end{bmatrix} + \begin{bmatrix} I & I \\ -\varepsilon \mu I & 0 \end{bmatrix} \begin{bmatrix} 0 \\ f(x) \end{bmatrix}, & \begin{bmatrix} x \\ v \end{bmatrix} \in \mathbb{R}^N \times \mathbb{R}^N, \\ \begin{bmatrix} x \\ v \end{bmatrix}(0) = \begin{bmatrix} x_0 \\ \varepsilon v_0 \end{bmatrix} \in \mathbb{R}^N \times \mathbb{R}^N. \end{cases}$$

Since ε goes to zero, one would expect that the dynamical properties of (A.2) are given by

$$\begin{cases} \frac{d}{dt} \begin{bmatrix} I & I \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ v \end{bmatrix} = - \begin{bmatrix} x \\ v \end{bmatrix} + \begin{bmatrix} I & I \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ f(x) \end{bmatrix}, & \begin{bmatrix} x \\ v \end{bmatrix} \in \mathbb{R}^N \times \mathbb{R}^N, \\ \begin{bmatrix} x \\ v \end{bmatrix}(0) = \begin{bmatrix} x_0 \\ \varepsilon v_0 \end{bmatrix} \in \mathbb{R}^N \times \mathbb{R}^N \end{cases}$$

which corresponds to $v = 0$ and

$$(A.3) \quad \begin{cases} \frac{dx}{dt} = -\mu x + f(x) \\ x(0) = x_0 \in \mathbb{R}^N. \end{cases}$$

Notice that the solutions for (A.3) are globally defined and the solution operator family $R_0(\cdot)$ defines a semigroup in \mathbb{R}^N . To compare the dynamics of these two problems we should find a way to see the dynamics of (A.3) in Z . That is done simply defining

$$T_0(t) \begin{bmatrix} x_0 \\ v_0 \end{bmatrix} = \begin{bmatrix} R_0(t)x_0 \\ 0 \end{bmatrix}, \quad t > 0, \quad T_0(0) = I,$$

and noting that $T_0(\cdot)$ is a semigroup (singular at zero) with attractor \mathcal{A}_0 . We can prove (see [7]) that there are constants $\varepsilon_0 > 0$, $\gamma > 0$, $t_0 > 0$ and $C > 0$ independent of ε such that

$$\text{dist}_H \left(T_\varepsilon(t) \bigcup_{\varepsilon \in [0, \varepsilon_0]} \mathcal{A}_\varepsilon, \mathcal{A}_\varepsilon \right) \leq C e^{-\gamma t}, \quad t \geq t_0$$

and for any $\bar{x}, \bar{y} \in Z$,

$$\|T_\varepsilon(t)\bar{x} - T_0(t)\bar{y}\|_Z \leq C e^{Lt}, \|\bar{x} - \bar{y}\|_Z + \varepsilon^\alpha, \quad t \geq t_0, \quad \alpha < 1.$$

Then by Corollary A.3

$$d_H(\mathcal{A}_\varepsilon, \mathcal{A}_0) \leq C\varepsilon^{\frac{\alpha\gamma}{\gamma+L}}.$$

APPENDIX B. SHADOWING THEORY AND RATE OF CONVERGENCE OF ATTRACTORS

In this section we make a brief overview of the some results on Shadowing Theory presented in [4] and [17]. This theory enable us estimate the convergence of attractors by continuity of nonlinear semigroups. Since the semigroups are given by variation of constants formula we can use the finite dimension to obtain estimates for the linear semigroup and then estimate the continuity of nonlinear semigroups.

We notice that the arguments work in finite dimension, therefore applying the techniques used here in more general problems where the dynamics can not be described by an ODE it seems not capable of working successfully.

Let $T : \mathbb{R}^N \rightarrow \mathbb{R}^N$ be a continuous function. Recall that the discrete dynamical system generated by T is defined by $T^0 = I_{\mathbb{R}^N}$ and, for $k \in \mathbb{N}$, $T^k = T \circ \dots \circ T$ is the k th iterate of T . The notions of Morse-Smale systems, hyperbolic fixed points, stable and unstable manifolds for a function T are similar to the continuous case; see e.g. [12].

Definition B.1. A trajectory (or global solution) of the discrete dynamical system generated by T is a sequence $\{x_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}^N$, such that, $x_{n+1} = T(x_n)$, for all $n \in \mathbb{Z}$.

Definition B.2. We say that a sequence $\{x_n\}_{n \in \mathbb{Z}}$ is a δ -pseudo-trajectory of T if

$$\|Tx_n - x_{n+1}\|_{\mathbb{R}^N} \leq \delta, \text{ for all } n \in \mathbb{Z}.$$

Definition B.3. We say that a point $x \in \mathbb{R}^N$ ε -shadows a δ -pseudo-trajectory $\{x_k\}$ on $U \subset \mathbb{R}^N$ if the following inequality holds

$$\|T^k x - x_k\|_{\mathbb{R}^N} \leq \varepsilon, \text{ for all } k \in \mathbb{Z},$$

Definition B.4. The map T has the Lipschitz Shadowing Property (LpSP) on $U \subset \mathbb{R}^N$, if there are constants $L, \delta_0 > 0$ such that for any $0 < \delta \leq \delta_0$, any δ -pseudo-trajectory of T in U is $(L\delta)$ -shadowed by a trajectory of T in \mathbb{R}^N , i.e., for any sequence $\{x_k\}_k \subset U \subset \mathbb{R}^N$ with

$$\|Tx_k - x_{k+1}\|_{\mathbb{R}^N} \leq \delta \leq \delta_0, \quad k \in \mathbb{Z},$$

there is a point $x \in X$ such that the following inequality holds

$$\|T^k x - x_k\|_{\mathbb{R}^N} \leq L\delta, \quad k \in \mathbb{Z}.$$

Let $T : \mathbb{R}^M \rightarrow \mathbb{R}^M$ be a Morse Smale function which has attractor \mathcal{A} . Since \mathcal{A} is compact and has all dynamic of the system, we can restrict our attention on a neighborhood $\mathcal{N}(\mathcal{A})$ of \mathcal{A} , thus we consider the space $C^1(\mathcal{N}(\mathcal{A}), \mathbb{R}^M)$ with the C^1 -topology.

The next result can be found in [3]. It describes an application of Shadowing Theory to rate of convergence of attractors.

Proposition B.5. Let $T_1, T_2 : X \rightarrow X$ be maps which has global attractors $\mathcal{A}_1, \mathcal{A}_2$. Assume that $\mathcal{A}_1, \mathcal{A}_2 \subset \mathcal{U} \subset X$, that T_1, T_2 have both the LpSP on \mathcal{U} , with parameters L, δ_0 and $\|T_1 - T_2\|_{\mathcal{L}^\infty(\mathcal{U}, X)} \leq \delta$. Then we have

$$\text{dist}_H(\mathcal{A}_1, \mathcal{A}_2) \leq \|T_1 - T_2\|_{\mathcal{L}^\infty(\mathcal{U}, X)}.$$

Proof. Since T_1 and T_2 has the LpSP on \mathcal{U} . Take a trajectory $\{y_n\}_n$ of T_2 in \mathcal{A}_2 , then $\{y_n\}_n$ is a δ -pesudo-trajectory of T_1 with $\delta = \|T_1 - T_2\|_{\mathcal{L}^\infty(\mathcal{U}, X)} \leq \delta_0$. By LpSP there is a trajectory $\{x_n\}_n \subset X$ of T_1 such that $\|x_n - y_n\|_X \leq L\delta$ for al $n \in \mathbb{Z}$, hence $\{x_n\}_n \subset \mathcal{A}_1$. Since $\{y_n\}_n$ is arbitrary the result follows. \square

As a consequence of the Proposition B.5, we have the following result.

Theorem B.6. Let $T : \mathbb{R}^M \rightarrow \mathbb{R}^M$ be a Morse-Smale function with attractor \mathcal{A} . Then there exist constant $L > 0$, a neighborhood $\mathcal{N}(\mathcal{A})$ of \mathcal{A} and a neighborhood $\mathcal{N}(T)$ of T in the $C^1(\mathcal{N}(\mathcal{A}), \mathbb{R}^M)$ topology such that, for any $T_1, T_2 \in \mathcal{N}(T)$ with attractors $\mathcal{A}_1, \mathcal{A}_2$, respectively, we have

$$\text{dist}_H(\mathcal{A}_1, \mathcal{A}_2) \leq L\|T_1 - T_2\|_{L^\infty(\mathcal{N}(\mathcal{A}), \mathbb{R}^M)}.$$

Proof. In [15] was proved that a structurally stable dynamical system on a compact manifold has the LpSP and it is know that a Morse-Smale system is structurally stable. Moreover, In [3] was proved that a discrete Morse-Smale semigroup T has the LpSP in a neighborhood $\mathcal{N}(\mathcal{A})$ of its attractor \mathcal{A} . Hence, the result follows by the Proposition B.5. \square

Example B.7. Consider the set $Q = \{(x, y) \in \mathbb{R} \times \mathbb{R}^{N-1} : 0 \leq x \leq 1, |y| < 1\}$. The thin domain is defined by $Q_\varepsilon = \{(x, \varepsilon y) \in \mathbb{R} \times \mathbb{R}^{N-1} : (x, y) \in Q\}$, $\varepsilon \in (0, 1]$. This domain is obtained by shrinking the set Q by a factor ε in the $N - 1$ direction given by the variable y . Note that Q_ε collapses to a straight segment $(0, 1)$ as ε goes to zero.

We consider the boundary value problem associated with a reaction-diffusion equation

$$(B.1) \quad \begin{cases} \partial_t u - \Delta u + \alpha u = f(u) & \text{in } Q_\varepsilon \times [0, \infty) \\ \frac{\partial u}{\partial \nu_\varepsilon} = 0 & \text{on } \partial Q_\varepsilon \times (0, \infty), \end{cases}$$

where $\alpha > 0$, ν_ε the unit outward normal to ∂Q_ε and $f : \mathbb{R} \rightarrow \mathbb{R}$ a C^2 -function satisfying




FIGURE 1. Thin Domain

- (i) $|f'(s)| \leq C(1 + |s|^{\rho-1})$, $s \in \mathbb{R}$, for some $\rho > 1$,
- (ii) $f(s)s \leq 0$, $|s| \geq M$, for some $M > 0$.

The limit problem of (B.1) as $\varepsilon \rightarrow 0$ is given by

$$(B.2) \quad \begin{cases} \partial_t u - \partial_x^2 u + \alpha u = f(u) & \text{in } (0, 1), \\ \partial_x u(0) = \partial_x u(1) = 0. \end{cases}$$

The equation (B.1) is well posed in $H^1(Q_\varepsilon)$ and the equation (B.2) is well posed in $H^1(0, 1)$, i.e., if we define the operators $A_\varepsilon = -\Delta u + \alpha u$ and $A_0 u = -\partial_x^2 u + \alpha u$, the global solutions of the Cauchy problem of type (2.4) with $f_\Omega^\varepsilon = f$ and $g_\Gamma^\varepsilon \equiv 0$ generates a nonlinear semigroup $T_\varepsilon(\cdot)$, and this semigroups has global attractor \mathcal{A}_ε , for all $\varepsilon \in [0, 1]$. Moreover $T_0(\cdot)$ is a Morse-Smale semigroup.

In order to understand the attractor \mathcal{A}_0 in $H^1(Q_\varepsilon)$, we define $E_\varepsilon : H^1(0, 1) \rightarrow H^1(Q_\varepsilon)$ by $(E_\varepsilon u)(x, y) = u(x)$, for all $u \in H^1(0, 1)$. Assuming that the equilibrium points of (B.2) are hyperbolic, was proved in [4] the following results

- (i) $\|A_\varepsilon^{-1} - E_\varepsilon A_0^{-1}\|_{\mathcal{L}(L^2(0,1), H^1(0,1))} \leq C\varepsilon$;
- (ii) $\|T_\varepsilon(1) - E_\varepsilon T_0(1)\|_{\mathcal{L}(H^1(0,1))} \leq C\varepsilon |\log(\varepsilon)|$;
- (iii) There is a finite dimensional invariant manifold \mathcal{M}_ε given by graph of Lipschitz functions s_ε such that $\mathcal{A}_\varepsilon \subset \mathcal{M}_\varepsilon$ and the flow can be reduced to finite dimension, i.e., we can consider an ODE generating a semigroup $\overline{T}_\varepsilon(\cdot)$ in \mathbb{R}^M with an attractor $\overline{\mathcal{A}}_\varepsilon$, where $M = \dim(\mathcal{M}_\varepsilon)$. Moreover, we have the following estimates
 - (a) $\|\overline{T}_\varepsilon(1) - \overline{T}_0(1)\|_{\mathcal{L}(\mathbb{R}^M)} \leq C\varepsilon |\log(\varepsilon)|$;
 - (b) $\sup_{v \in \mathbb{R}^M} \|s_\varepsilon(v) - s_0(v)\|_{H^1(0,1)} \leq C\varepsilon |\log(\varepsilon)|$.

Therefore the convergence of attractors of B.1 and B.2 can be estimated by

$$d_H(\mathcal{A}_\varepsilon, E_\varepsilon \mathcal{A}_0) \leq C\varepsilon^{\frac{N+1}{2}} |\log(\varepsilon)|,$$

where $C > 0$ is a constant independent of ε .

REFERENCES

- [1] J. M. Arrieta, F. D. M. Bezerra, and A. N. Carvalho. Rate of convergence of attractors for some singular perturbed parabolic problems. *Topological Methods in Nonlinear Analysis*, 41:229–253, 2013.

- [2] J. M. Arrieta, A. N. Carvalho, and A. Rodríguez-Bernal. Attractors for parabolic problems with nonlinear boundary condition. uniform bounds. *Commun. in partial differential equations*, 25:1–37, 2000.
- [3] J. M. Arrieta and E. Santamaria. Estimates on the distance of inertial manifolds. *Discrete and continuous dynamical systems*, 34(10):3921–3944, 2014.
- [4] J. M. Arrieta and E. Santamaria. Distance of attractors of reaction-diffusion equations in thin domains. *Journal of Differential Equations*, 63:4222–4266, 2017.
- [5] J. M. Arrieta and E. Santamaria. $c^{1,\theta}$ —estimates on the distance of inertial manifolds. *Collectanea Mathematica*, 69:315–336, 218.
- [6] A. Babin and M. Vishik. *Attractors of Evolution Equations*. Elsevier, 1992.
- [7] A. Carvalho, J. Cholewa, and T. Dlotko. Equi-exponential attraction and rate of convergence of attractors for singularly perturbed evolution equations. *Proceedings of the Royal Society of Edinburgh*, 144(1):13–51, 2014.
- [8] A. N. Carvalho, J. Langa, and J. Robinson. *Attractors for infinite-dimensional non-autonomous dynamical systems*. Springer, 2010.
- [9] A. N. Carvalho and L. Pires. Rate of convergence of attractors for singularly perturbed semilinear problems. *Journal of Mathematical Analysis and Applications*, 452:258–296, 2017.
- [10] A. N. Carvalho and L. Pires. Parabolic equations with localized large diffusion: rate of convergence of attractors. *Topological Methods in Nonlinear Analysis*, 53(1):1–23, 2019.
- [11] J. Hale and G. Raugel. Lower semi-continuity of attractors of gradient systems and applications. *Ann Mat Pura Appl*, 154:281–326, 1989.
- [12] J. K. Hale. *Asymptotic behavior of dissipative systems*, volume 25. Mathematical surveys and monographs, 1988.
- [13] D. Henry. *Geometric theory of semilinear parabolic equations*. Number 840 in Springer-Verlag. Lecture Notes in Mathematics, 1980.
- [14] D. Henry. Some infinite-dimensional morse-smale systems defined by parabolic partial differential equations. *Journal of differential equations*, 59:165–205, 1985.
- [15] S. Y. Pilyugin. *Shadowing in Dynamical Systems*, volume 1706 of *Lectures Notes in Mathematics*. Springer, 1999.
- [16] A. Rodríguez-Bernal. Localized spatial homogenizations and large diffusion. *SIAM Journal of Mathematical Analysis*, 29(6):1361–1380, 1998.
- [17] E. Santamaria. *Distance of attractors of evolutionary equations*. PhD thesis, Universidad Complutense de Madrid, 2013.

(F. D. M. Bezerra) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58051-900 JOÃO PESSOA PB, BRAZIL

E-mail address: flank@mat.ufpb.br

(Leonardo Pires) DEPARTAMENTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE ESTADUAL DE PONTA GROSSA, 84030-900 PONTA GROSSA PR, BRAZIL

E-mail address: lpires@uepg.br