CATEGORIES OVER QUANTUM AFFINE ALGEBRAS AND MONOIDAL CATEGORIFICATION

MASAKI KASHIWARA, MYUNGHO KIM, SE-JIN OH, AND EUIYONG PARK

ABSTRACT. Let $U'_q(\mathfrak{g})$ be a quantum affine algebra of untwisted affine ADE type, and $\mathscr{C}^0_{\mathfrak{g}}$ the Hernandez-Leclerc category of finite-dimensional $U'_q(\mathfrak{g})$ -modules. For a suitable infinite sequence $\widehat{w}_0 = \cdots s_{i-1} s_{i_0} s_{i_1} \cdots$ of simple reflections, we introduce subcategories $\mathscr{C}^{[a,b]}_{\mathfrak{g}}$ of $\mathscr{C}^0_{\mathfrak{g}}$ for all $a \leq b \in \mathbb{Z} \sqcup \{\pm\infty\}$. Associated with a certain chain \mathfrak{C} of intervals in [a,b], we construct a real simple commuting family $M(\mathfrak{C})$ in $\mathscr{C}^{[a,b]}_{\mathfrak{g}}$, which consists of Kirillov-Reshetikhin modules. The category $\mathscr{C}^{[a,b]}_{\mathfrak{g}}$ provides a monoidal categorification of the cluster algebra $K(\mathscr{C}^{[a,b]}_{\mathfrak{g}})$, whose set of initial cluster variables is $[M(\mathfrak{C})]$. In particular, this result gives an affirmative answer to the monoidal categorification conjecture on $\mathscr{C}^-_{\mathfrak{g}}$ by Hernandez-Leclerc since it is $\mathscr{C}^{[-\infty,0]}_{\mathfrak{g}}$, and is also applicable to $\mathscr{C}^0_{\mathfrak{g}}$ since it is $\mathscr{C}^{[-\infty,\infty]}_{\mathfrak{g}}$.

1. INTRODUCTION

Let $U'_q(\mathfrak{g})$ be a quantum affine algebra. The category $\mathscr{C}_{\mathfrak{g}}$ of finite-dimensional integrable modules over $U'_q(\mathfrak{g})$ has been intensively studied due to its rich structure. For instances, every object M in $\mathscr{C}_{\mathfrak{g}}$ has its left M^* and right dual *M , and the *q*-characters of Kirillov-Reshetikhin modules in $\mathscr{C}_{\mathfrak{g}}$ provide a solution of the *T*-system, a system of differential equations appearing in solvable lattice models ([2, 5, 10, 19]).

On the other hand, the cluster algebras were introduced by Fomin and Zelivinsky in [3] to investigate upper global bases and total positivity in an aspect of combinatorics.

Interestingly, it is proved in [6, 7, 9] that the Grothendieck rings $K(\mathcal{C})$ of monoidal subcategories $\mathcal{C} = \mathscr{C}_N$ $(N \in \mathbb{Z}_{\geq 1}), \mathscr{C}_Q, \mathscr{C}_{\mathfrak{g}}^-$ of $\mathscr{C}_{\mathfrak{g}}$ have cluster algebra structures \mathscr{A} , and conjectured that every cluster monomial corresponds to the isomorphism class of

Date: May 22, 2020.

²⁰¹⁰ Mathematics Subject Classification. 17B37, 81R50, 18D10.

Key words and phrases. Monoidal categorification, Quantum affine algebra, Cluster algebra, Kirillov-Reshetikhin module, T-system.

The research of M. Kashiwara was supported by Grant-in-Aid for Scientific Research (B) 15H03608, Japan Society for the Promotion of Science.

The research of M. Kim was supported by the National Research Foundation of Korea(NRF) Grant funded by the Korea government(MSIP) (NRF-2017R1C1B2007824).

The research of S.-j. Oh was supported by the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2019R1A2C4069647).

a real simple module in \mathcal{C} ; that is, \mathcal{C} is expected to be a monoidal categorification of \mathscr{A} . The conjectures for \mathscr{C}_N ($N \in \mathbb{Z}_{\geq 1}$) of untwisted affine ADE types are proved in [6, 8, 20] and [23]. Also, the conjecture for the subcategory $\mathscr{C}_Q \subset \mathscr{C}_g$, determined by a Q-data $\mathcal{Q} = (Q, \phi_Q)$ ([4, 17]), is proved in [12] via the quantum affine Weyl-Schur duality functor \mathcal{F}_Q ([11, 13, 18, 21]) from the category \mathcal{C}_{QH} of finite-dimensional graded modules over the symmetric quiver Hecke algebra to \mathscr{C}_Q . More precisely, the category \mathcal{C}_{QH} provides a monoidal categorification of the quantum cluster algebra $A_q(\mathfrak{n})$, the quantum unipotent coordinate algebra of finite simply-laced type ([1]). Since \mathcal{F}_Q is an exact monoidal functor preserving simplicity, we can prove the conjecture for \mathscr{C}_Q in an indirect way. However, this method could not be applicable to other \mathcal{C} directly.

Recently, in [14], the authors of the present paper (KKOP) developed Z-valued invariants $\Lambda, \Lambda^{\infty}, \tilde{\Lambda}, \mathfrak{d}$ for pairs of modules in $\mathscr{C}_{\mathfrak{g}}$, which is extracted from distinguished $U'_q(\mathfrak{g})$ -module homomorphisms, called *R-matrices*. Furthermore, KKOP provided a criterion for a monoidal subcategory $\mathscr{C} \subset \mathscr{C}_{\mathfrak{g}}$ to become a monoidal categorification of a cluster algebra by using those invariants. This paper can be understood as a continuation of [14], since we will apply the above criterion to various subcategories \mathscr{C} of $\mathscr{C}_{\mathfrak{g}}$, including $\mathscr{C}_{\mathfrak{g}}^0, \mathscr{C}_{\mathfrak{g}}^-$ and \mathscr{C}_N . We also give their initial monoidal seeds in a uniform manner.

Let \mathfrak{g}_0 be a finite-dimensional simple Lie algebra of ADE type with a Cartan matrix $\mathsf{A} = (a_{ij})_{i,j\in I_0}$, W the Weyl group generated by simple reflections s_i $(i \in I_0)$, \mathfrak{g} the untwisted affine Kac-Moody algebra associated with \mathfrak{g}_0 , and $U'_q(\mathfrak{g})$ the quantum affine algebra associated with \mathfrak{g} . In [6], Hernandez-Leclerc defined the full subcategory $\mathscr{C}^0_{\mathfrak{g}}$ of $\mathscr{C}_{\mathfrak{g}}$. Since every simple modules in $\mathscr{C}_{\mathfrak{g}}$ is a tensor product of suitable parameter shifts of simple modules in $\mathscr{C}_{\mathfrak{g}}^0$, it is enough to consider subcategories of $\mathscr{C}^0_{\mathfrak{g}}$.

By extending a reduced expression $s_{i_1}s_{i_2}\cdots s_{i_\ell}$ of the longest element w_0 of the Weyl group W, we obtain an infinite sequence

(1.1)
$$\widehat{w}_0 = \cdots s_{i_{-2}} s_{i_{-1}} s_{i_0} s_{i_1} s_{i_2} \cdots$$

of simple reflections satisfying properties (a) and (b) in Section 2, and then we define fundamental modules $V[k]^{\widehat{w}_0}$ $(k \in \mathbb{Z})$. For each *interval* $[a, b] = \{k \in \mathbb{Z} \mid a \leq k \leq b\}$ with $a \leq b \in \mathbb{Z} \sqcup \{\pm \infty\}$, we define the subcategory $\mathscr{C}_{\mathfrak{g}}^{[a,b]}$ of $\mathscr{C}_{\mathfrak{g}}^{0}$ which is the smallest full monoidal subcategory containing $V[k]^{\widehat{w}_0}$ for all $k \in [a, b]$. Then $\mathscr{C}_{\mathfrak{g}}^{0}$ is nothing but $\mathscr{C}_{\mathfrak{g}}^{[-\infty,+\infty]}$ and the subcategory $\mathscr{C}_{\mathfrak{g}}^{-}$ introduced by Hernandez-Leclerc ([9]) can be identified with $\mathscr{C}_{\mathfrak{g}}^{[-\infty,0]}$ (Remark 2.3).

We say that an interval [a, b] is an *i*-box if $i_a = i_b$. For each *i*-box [a, b], we define a simple module M[a, b], which can be understood as a quantum affine analogue of the determinantial module (see Remark 3.2). In Theorem 3.1, we show that M[a, b] is a Kirillov-Reshetikhin module and give a sufficient condition for the simplicity of the tensor product $M[a, b] \otimes M[a', b']$ for *i*-boxes [a, b] and [a', b']. Then we define the notion of an admissible chain $\mathfrak{C} = \{[a_k, b_k] \subset [a, b] \mid 1 \leq k \leq b - a + 1, i_{a_k} = i_{b_k}\}$ of *i*-boxes for an interval [a, b] satisfying certain properties (Definition 3.3). For each admissible chain \mathfrak{C} , the family of Kirillov-Reshetikhin modules $M(\mathfrak{C}) = \{M[a_k, b_k]\}_{1 \leq k \leq b-a+1}$ in $\mathscr{C}_{\mathfrak{g}}^{[a,b]}$ forms a commuting family of real simple modules (Theorem 3.4).

The next step is to show that $K(\mathscr{C}_{\mathfrak{g}}^{[a,b]})$ has a cluster algebra structure, $\mathscr{C}_{\mathfrak{g}}^{[a,b]}$ provides a monoidal categorification of $K(\mathscr{C}_{\mathfrak{g}}^{[a,b]})$, and any admissible chain \mathfrak{C} gives a monoidal seed $M(\mathfrak{C})$ (Theorem 4.3). Based on the criterion in [14], we shall prove this by showing the assertion for a special chain \mathfrak{C} , and then by extending it to a general \mathfrak{C} . Namely, we proceed by proving

- (i) the existence of a Λ -admissible monoidal seed \mathscr{S} of $K(\mathscr{C}_{\mathfrak{g}}^{[a,b]})$ whose set of initial cluster variable modules is $M(\mathfrak{C})$ for some admissible chain \mathfrak{C} ,
- (ii) the existence of sequences of mutations among the $M(\mathfrak{C})$'s only employing T-systems,

which implies that any admissible chain \mathfrak{C} gives a Λ -admissible monoidal seed for all $\mathscr{C}_{\mathfrak{g}}^{[a,b]}$. In particular, we prove that \mathscr{C} is a monoidal categorification of the cluster algebra $K(\mathscr{C})$ for $\mathscr{C} = \mathscr{C}_{\mathfrak{g}}^{0}$ and $\mathscr{C} = \mathscr{C}_{\mathfrak{g}}^{-}$. Note that we need in step (i) above the existence of the cluster algebra structure on $K(\mathscr{C}_{\mathfrak{g}}^{-})$ proved in [9].

This paper is an announcement whose details will appear elsewhere.

2. Subcategories

We take the algebraic closure \mathbf{k} of $\mathbb{C}(q)$ inside $\bigcup_{m>0} \mathbb{C}((q^{1/m}))$ as the base field for $U'_q(\mathfrak{g})$. Recall that $\mathscr{C}_{\mathfrak{g}}$ is the category of finite-dimensional integrable modules over $U'_q(\mathfrak{g})$. There is a family $\{V(\varpi_i)_c \mid i \in I_0, c \in \mathbf{k}^{\times}\}$ in $\mathscr{C}_{\mathfrak{g}}$ of simple modules, called the fundamental modules.

For simple modules M and N in $\mathscr{C}_{\mathfrak{g}}$, we say that M and N strongly commute if $M \otimes N$ is simple, and M is real if $M^{\otimes k}$ is simple for all $k \in \mathbb{Z}_{\geq 1}$.

Let us denote by Ψ the quiver whose set of vertices is

$$I_0 := \{ (i,k) \in I_0 \times \mathbb{Z} \mid k \equiv d(1,i) \mod 2 \}.$$

and the arrows of Ψ consist of two types:

(2.1)
$$\begin{cases} (A) & (i,t) \to (j,s) \text{ with } d(i,j) = 1 \text{ and } s - t = 1, \\ (B) & (i,s+2) \to (i,s). \end{cases}$$

Here d(i, j) denotes the distance between the vertices i and j in the Dynkin diagram of \mathfrak{g}_0 and $1 \in I_0$ is an arbitrary chosen element.

We say that an infinite sequence

$$\widehat{w}_0 = \cdots s_{i-1} s_{i_0} s_{i_1} \cdots$$

of simple reflections in the Braid group $B(\mathfrak{g}_0)$ ([16]) of type \mathfrak{g}_0 is *admissible* if (a) there exists a sequence $\{t_k\}_{k\in\mathbb{Z}}$ of integers such that

- (1) $(i_k, t_k) \in I_0$, (2) $t_{k^+} = t_k + 2$, and
- (3) $t_k > t_{k'}$ if k > k' and $d(i_k, i_{k'}) = 1$.
- (b) $s_{i_k} \cdots s_{i_{k+\ell-1}} = w_0$ for all $k \in \mathbb{Z}$, where ℓ denotes the length of longest element $w_0 \in W$.

Here, for $k \in \mathbb{Z}$ and $j \in I_0$, we set

$$k^{+} := \min\{p \mid k < p, \ i_{k} = i_{p}\},\$$

$$k^{-} := \max\{p \mid p < k, \ i_{k} = i_{p}\},\$$

$$k(j)^{+} := \min\{p \mid k \leq p, \ i_{p} = j\},\$$

$$k(j)^{-} := \max\{p \mid p \leq k, \ i_{p} = j\}.$$

Remark 2.1. (i) We have $i_{k+\ell} = i_k^*$, where * denotes the involution on I_0 induced by w_0 .

- (ii) \widehat{w}_0 completely determines $\{(i_k, t_k)\}_{k \in \mathbb{Z}}$ up to an even translation.
- (iii) For every $k \in \mathbb{Z}$, the reduced expression $s_{i_k} \cdots s_{i_{k+\ell-1}}$ in (b) is *adapted* to some Dynkin quiver Q of type \mathfrak{g}_0 . Conversely, for any Dynkin quiver Q of type \mathfrak{g}_0 , there exists a sequence \widehat{w}_0 satisfying (a) and (b) such that $s_{i_1} \cdots s_{i_\ell}$ is adapted to Q.

For each $k \in \mathbb{Z}$, we define the fundamental module

$$V[k]^{w_0} := V(\varpi_{i_k})_{(-q)^{t_k}}.$$

Then we have

$$V[k^{\pm}]^{\widehat{w}_0} \simeq V[k]^{\widehat{w}_0}_{(-q)^{\pm 2}}$$
 and $V[k+\ell]^{\widehat{w}_0} = \mathscr{D}(V[k]^{\widehat{w}_0}),$

where \mathscr{D} is the right dual functor.

Definition 2.2. For each interval [a, b], we denote by $\mathscr{C}_{\mathfrak{g}}^{[a,b]}$ the smallest full subcategory of $\mathscr{C}_{\mathfrak{g}}$ satisfying the following conditions:

- (i) it is stable under taking subquotients, extensions, tensor products and
- (ii) it contains $V[k]^{\widehat{w}_0}$ for all $a \leq k \leq b$ and the trivial module 1.

Remark 2.3. Many of known subcategories \mathcal{C} of $\mathscr{C}_{\mathfrak{g}}$ can be identified with $\mathscr{C}_{\mathfrak{g}}^{[a,b]}$ by taking suitable [a,b]:

- (1) $\mathscr{C}_{\mathfrak{g}}^{[-\infty,\infty]}$ coincides with the subcategory $\mathscr{C}_{\mathfrak{g}}^{0}$.
- (2) The subcategory $\mathscr{C}_{\mathcal{Q}}$ associated to a Q-data \mathcal{Q} coincides with $\mathscr{C}_{\mathfrak{g}}^{[a,b]}$ for some interval [a,b] with $|[a,b]| := b a + 1 = \ell$.

4

CATEGORIES OVER QUANTUM AFFINE ALGEBRAS AND MONOIDAL CATEGORIFICATION 5

(3) By taking $s_{i_1} \cdots s_{i_\ell}$ in (b) as adapted to the Dynkin quiver Q with $\{1, 2\} \ni \phi_Q(k) \equiv d(1, i_k) \pmod{2}$ and $t_k = \phi(i_k)$ for $1 \leq k \leq |I_0|$, \mathscr{C}_N can be identified with $\mathscr{C}_{\mathfrak{g}}^{[a,0]}$ where $a = 1 - (N \times |I_0|)$, and $\mathscr{C}_{\mathfrak{g}}^-$ can be identified with $\mathscr{C}_{\mathfrak{g}}^{[-\infty,0]}$. Those subcategories $\mathscr{C}_N, \mathscr{C}_{\mathfrak{g}}^-$ of $\mathscr{C}_{\mathfrak{g}}^0$ are introduced in [6, 9].

3. Real simple commuting family associated to an admissible chain of i-boxes

Let us fix an admissible sequence \widehat{w}_0 and $\{t_k\}_{k\in\mathbb{Z}}$. We write V[k] for $V[k]^{\widehat{w}_0}$. We say that an interval c = [a, b] is an *i*-box if $i_a = i_b$. For each *i*-box [a, b], the module M[a, b] in $\mathscr{C}^0_{\mathfrak{a}}$ is defined as follows:

$$M[a,b] := \operatorname{hd}(V[b] \otimes V[b^{-}] \otimes \cdots \otimes V[a^{+}] \otimes V[a]),$$

where hd(M) for $M \in \mathscr{C}_{\mathfrak{g}}$ denotes the head of M. In particular, V[a] = M[a, a].

Theorem 3.1. (i) M[a, b] is a Kirillov-Reshetikhin module with a dominant extremal weight $s \varpi_{i_a}$ where $s = |\{k \mid a \leq k \leq b, i_k = i_a\}|$.

(ii) For *i*-boxes [a, b] and [c, d], M[a, b] and M[c, d] commutes if either

 $a^- < c \leqslant d < b^+ \quad or \quad c^- < a \leqslant b < d^+.$

(iii) For any *i*-box [a, b], there exists an exact sequence in terms of M[a, b]'s as follows:

$$(3.1) \quad 0 \longrightarrow \bigotimes_{d(i_a,j)=1} M[a(j)^+, b(j)^-] \to M[a^+, b] \otimes M[a, b^-] \to M[a, b] \otimes M[a^+, b^-] \to 0,$$

We call it a T-system.

Remark 3.2. For any reduced expression $\underline{w}_0 = s_{j_1} \cdots s_{j_\ell}$ of w_0 (not necessarily adapted) and [a, b] with $j_a = j_b$ and $b - a + 1 \leq \ell$, there exists a real simple module $D[a, b]^{\underline{w}_0}$ in \mathcal{C}_{QH} of type \mathfrak{g}_0 , called the *determinantial module*, and there exists an exact sequence (called the T-system)

$$0 \longrightarrow \bigotimes_{d(i_a,j)=1} \mathcal{D}[a(j)^+, b(j)^-] \to \mathcal{D}[a^+, b] \otimes \mathcal{D}[a, b^-] \to \mathcal{D}[a, b] \otimes \mathcal{D}[a^+, b^-] \longrightarrow 0$$

in C_{QH} , which is analogous to (3.1). More precisely, when \underline{w}_0 is adapted to some Dynkin quiver Q of type \mathfrak{g}_0 , quantum affine Weyl-Schur duality functor \mathcal{F}_Q associated with some Q-data $Q = (Q, \phi_Q)$ transforms the above exact sequence in C_{QH} to the T-system (3.1). Thus M[a, b] can be understood as a quantum affine analogue of the determinantial module. (See [7, Proposition 4.1] and [12] for more detail.)

For any interval c := [a, b], we introduce *i*-boxes

$$[a,b] := [a,b(i_a)^-], \quad \{a,b] := [a(i_b)^+,b] \text{ and}$$

 $\mathcal{L}(c) := [a-1,b], \quad \mathcal{R}(c) := \{a,b+1].$

Definition 3.3. A chain \mathfrak{C} of *i*-boxes

$$(c_k = [a_k, b_k])_{1 \le k \le l} \quad (l \in \mathbb{Z}_{\ge 1} \sqcup \{\infty\})$$

is called admissible if the interval $\widetilde{c}_k = [\widetilde{a}_k, \widetilde{b}_k] := \bigcup_{1 \leq j \leq k} [a_j, b_j]$ satisfies $|\widetilde{c}_k| = k$ and one of the following two statements.

- (1) $c_k = \mathcal{L}(\widetilde{c}_{k-1}),$
- (2) $c_k = \mathcal{R}(\widetilde{c}_{k-1}).$

(Please do not confuse l and ℓ .) The sequence of intervals $\{\widetilde{c}_k\}_{1 \leq k \leq l}$ is called the envelope of the chain \mathfrak{C} and \widetilde{c}_l is called the range of \mathfrak{C} .

Thus, for an admissible chain \mathfrak{C} of *i*-boxes, we can associate a pair (a, \mathfrak{T}) consisting of an integer *a* and a sequence $\mathfrak{T} = (T_1, T_2, \ldots, T_{l-1})$ such that (i) $T_i \in \{\mathcal{L}, \mathcal{R}\}$ ($1 \leq i \leq l-1$), (ii) $a = a_1 = b_1$,

(iii) $[\widetilde{a}_k, \widetilde{b}_k] = \begin{cases} [\widetilde{a}_{k-1} - 1, \widetilde{b}_{k-1}] & \text{if } T_{k-1} = \mathcal{L}, \\ [\widetilde{a}_{k-1}, \widetilde{b}_{k-1} + 1] & \text{if } T_{k-1} = \mathcal{R}. \end{cases}$

Hence we have $c_k = [a_k, b_k] = T_{k-1}[\tilde{a}_{k-1}, \tilde{b}_{k-1}]$ $(k \ge 2)$, and the interval \tilde{c}_k is obtained from \tilde{c}_{k-1} by adding an element from the left or from the right according that $T_{k-1} = \mathcal{L}$ or $T_{k-1} = \mathcal{R}$.

For an admissible chain $\mathfrak{C} = (c_k)_{1 \leq k \leq l}$ with the associated pair (a, \mathfrak{T}) and for $1 \leq s < l$, we say that an *i*-box c_s is *movable* if s = 1 or $T_{s-1} \neq T_s$ $(s \geq 2)$. For a movable c_s in \mathfrak{C} , we define a new admissible chain $B_s(\mathfrak{C})$ whose associated pair (a', \mathfrak{T}') is given

(i)
$$\begin{cases} a' = a \pm 1 & \text{if } s = 1 \text{ and } T_1 = \mathcal{R} \text{ (resp. } \mathcal{L}), \\ a' = a & \text{if } s > 1, \end{cases}$$

(ii) $T'_k = T_k \text{ for } k \notin \{s - 1, s\}, \text{ and}$
(iii) $T'_k \neq T_k \text{ for } k \in \{s - 1, s\}.$

That is, $B_s(\mathfrak{C})$ is the admissible chain obtained from \mathfrak{C} by moving \tilde{c}_s by 1 to the right or to the left inside \tilde{c}_{s+1} .

Theorem 3.4. Let $\mathfrak{C} = (c_k)_{1 \leq k \leq l}$ be an admissible chain and set

$$M(\mathfrak{C}) := \{ M[a_k, b_k] \mid 1 \leq k \leq l \}.$$

Then we have the followings:

(a) $M(\mathfrak{C})$ is a commuting family of real simple modules.

- (b) If M[c,d] commutes with all $M[a_k,b_k]$ and $[c,d] \subset [\widetilde{a}_l,\widetilde{b}_l]$, then $[c,d] \in \mathfrak{C}$.
- (c) For another admissible chain $\mathfrak{C}' = (c'_k)_{1 \leq k \leq l}$ with the same range, there exists a finite sequence $(t_1, t_2, \ldots, t_r) \in \{1, 2, \ldots, l\}^r$ such that

$$B_{t_r}(\cdots(B_{t_2}(B_{t_1}(\mathfrak{C}))\cdots)=\mathfrak{C}'.$$

CATEGORIES OVER QUANTUM AFFINE ALGEBRAS AND MONOIDAL CATEGORIFICATION 7

4. MONOIDAL CATEGORIFICATION

Let $K = K_{\text{ex}} \sqcup K_{\text{fr}}$ be a countable index set. Let \mathscr{C} be a full subcategory of $\mathscr{C}^0_{\mathfrak{g}}$ stable under taking subquotients, extensions and tensor products.

A monoidal seed in \mathscr{C} is a pair $\mathscr{S} = (\{M_i\}_{i \in K}, B)$ consisting of a commuting family $\{M_i\}_{i \in K}$ of real simple objects in \mathscr{C} and a \mathbb{Z} -valued $K \times K_{ex}$ -matrix $\widetilde{B} = (b_{ij})_{(i,j) \in K \times K_{ex}}$ such that

- (i) for each $j \in K_{ex}$, there exist finitely many $i \in K$ such that $b_{ij} \neq 0$,
- (ii) the principal part $B := (b_{ij})_{i,j \in K_{ex}}$ is skew-symmetric. For $i \in K$, we call M_i the *i*-th cluster variable module of \mathscr{S} .

For a monoidal seed $\mathscr{S} = (\{M_i\}_{i \in K}, \widetilde{B})$, let $\Lambda^{\mathscr{S}} = (\Lambda^{\mathscr{S}}_{ij})_{i,j \in K}$ be the skew-symmetric matrix given by $\Lambda^{\mathscr{S}}_{ij} = \Lambda(M_i, M_j)$ (see [15]).

A monoidal seed $\mathscr{S} = (\{M_i\}_{i \in K}, \widetilde{B})$ is called Λ -admissible if (i) $(\Lambda^{\mathscr{S}}\widetilde{B})_{jk} = -2\delta_{jk}$ for $(j,k) \in K \times K_{ex}$, and (ii) for each $k \in K_{ex}$, there exist a simple object M'_i of \mathcal{C} commuting

(ii) for each $k \in K_{ex}$, there exist a simple object M'_k of \mathcal{C} commuting with M_i for any $i \neq k$ and an exact sequence in \mathscr{C}

(4.1)
$$0 \longrightarrow \bigotimes_{b_{ik}>0} M_i^{\otimes b_{ik}} \to M_k \otimes M_k' \to \bigotimes_{b_{ik}<0} M_i^{\otimes (-b_{ik})} \longrightarrow 0.$$

Under the following two assumptions

- (a) there exists a Λ -admissible monoidal seed $\mathscr{S} = (\{M_i\}_{i \in K}, \widetilde{B})$ in \mathscr{C} ,
- (b) $K(\mathscr{C})$ is isomorphic to the cluster algebra $\mathscr{A}([\mathscr{S}])$,

KKOP ([14, Theorem 6.10]) proved that \mathscr{C} provides a monoidal categorification of $\mathscr{A}([\mathscr{S}])$. Here $[\mathscr{S}] := ([\{M_i\}]_{i \in K}, \widetilde{B})$ is a seed in $K(\mathscr{C})$, and $\mathscr{A}([\mathscr{S}])$ denotes the cluster algebra with the initial seed $[\mathscr{S}]$.

Set $\widehat{I_0}^- := \widehat{I_0} \cap (I_0 \times \mathbb{Z}_{\leq 0})$ and let Ψ^- be the full subquiver of Ψ whose set of vertices is $\widehat{I_0}^-$. In [9], Hernandez-Leclerc proved that $\mathscr{A}^- := K(\mathscr{C}_{\mathfrak{g}}^-)$ has a cluster algebra structure whose initial cluster variable modules $\{\mathcal{M}_{(i,t)}\}_{(i,t)\in\widehat{I_0}^-}$ consist of certain KR-modules. For a suitable choice of \widehat{w}_0 (Remark 2.3), we have $\mathscr{C}_{\mathfrak{g}}^- = \mathscr{C}_{\mathfrak{g}}^{[-\infty,0]}$ and $\{\mathcal{M}_{(i,t)}\}_{(i,t)\in\widehat{I_0}^-}$ can be described as $M(\mathfrak{C}^-)$ for the following admissible chain \mathfrak{C}^- of *i*-boxes:

$$\mathfrak{C}^{-} = (0, \mathfrak{T} = (\mathcal{L}, \mathcal{L}, \mathcal{L}, \ldots)).$$

More precisely, for $(i, t) = (i_a, t_a)$ $(a \leq 0)$, we have

$$\mathcal{M}_{(i,t)} = M[a,0].$$

The following theorem gives an affirmative answer for the conjecture on $\mathscr{C}_{\mathfrak{g}}^-$:

Theorem 4.1. The monoidal seed

$$\mathscr{S}^- := (M(\mathfrak{C}^-), \widetilde{B}^-)$$
 is Λ -admissible,

where \widetilde{B}^- is the matrix associated with Ψ^- . Hence $\mathscr{C}_{\mathfrak{g}}^-$ provides a monoidal categorification of \mathscr{A}^- .

Now we shall generalize the above theorem to an arbitrary $\mathscr{C}^{[a,b]}_{\mathfrak{a}}$.

Proposition 4.2. Let $\mathfrak{C} = (c_k)_{1 \leq k \leq l}$ be an admissible chain of *i*-boxes with the range [a, b] and the envelope $\{\widetilde{c}_k\}_{1 \leq k \leq l}$. Assume that $\mathscr{C} := \mathscr{C}_{\mathfrak{g}}^{[a,b]}$ provides a monoidal categorification of $K(\mathscr{C})$ with a Λ -admissible monoidal seed $(M(\mathfrak{C}), \widetilde{B})$. Let c_s be a movable *i*-box of \mathfrak{C} and set $\mathfrak{C}' = B_s(\mathfrak{C})$.

- (i) If $\widetilde{c}_{s+1} \neq c_{s+1}$, then $M(\mathfrak{C}')$ is equal to $M(\mathfrak{C})$ up to a permutation.
- (ii) If c̃_{s+1} = c_{s+1}, then M(𝔅') is the monoidal mutation of M(𝔅) at s. Moreover the corresponding exact sequence (4.1) is given by the T-system (3.1).

The above proposition and Theorem 3.4 show that all $M(\mathfrak{C})$ are mutation equivalent. Now we state our main theorem:

Theorem 4.3. For any admissible chain $\mathfrak{C} = (c_k)_{1 \leq k \leq l}$ for $l \in \mathbb{Z}_{\geq 1} \sqcup \{\infty\}$ with the range

 $\widetilde{c}_l = [a, b] \quad for \ a \leqslant b \in \mathbb{Z} \sqcup \{\pm \infty\},$

there exists a Λ -admissible monoidal seed \mathscr{S} of $\mathscr{C}_{\mathfrak{g}}^{[a,b]}$ such that

- (i) its set of cluster variable modules is $M(\mathfrak{C})$,
- (ii) its set of frozen variable modules is $\{M[a(i)^+, b(i)^-] \mid i \in I_0, -\infty < a(i)^+ \leq b(i)^- < +\infty\},\$

and

(iii) $K(\mathscr{C}^{[a,b]}_{\mathfrak{g}})$ has the cluster algebra structure with the initial seed $[\mathscr{S}]$, and $\mathscr{C}^{[a,b]}_{\mathfrak{g}}$ provides a monoidal categorification of $\mathscr{A}([\mathscr{S}]) \simeq K(\mathscr{C}^{[a,b]}_{\mathfrak{g}})$.

By Remark 2.3, we have the following corollary:

Corollary 4.4. The Grothendieck ring $K(\mathscr{C}^0_{\mathfrak{g}})$ has a cluster algebra strucure, and $\mathscr{C}^0_{\mathfrak{g}}$ provides a monoidal categorification of $K(\mathscr{C}^0_{\mathfrak{g}})$.

Remark 4.5. We can generalize the above results to an arbitrary quantum affine algebra $U'_q(\mathfrak{g})$ by applying a similar framework with the results in [13, 18, 21, 22].

References

- A. Berenstein and A. Zelevinsky, Quantum cluster algebras, Adv. Math. 195 (2005), no. 2, 405– 455.
- [2] E. Frenkel and N. Yu. Reshetikhin, The q-characters of representations of quantum affine algebras and deformations of W-algebras, Recent developments in quantum affine algebras and related topics, Contemp. Math. 248 (1999), 163–205.

CATEGORIES OVER QUANTUM AFFINE ALGEBRAS AND MONOIDAL CATEGORIFICATION 9

- [3] S. Fomin and A. Zelevinsky, Cluster algebras I. Foundations, J. Amer. Math. Soc. 15 (2002), no. 2, 497–529.
- [4] R. Fujita and S-j. Oh, Combinatorics of twisted Auslander-Reiten quivers and Representations of quantum affine algebras, in preparation.
- [5] D. Hernandez, The Kirillov-Reshetikhin conjecture and solutions of T-systems, J. Reine Angew. Math. 596 (2006), 63–87.
- [6] D. Hernandez and B. Leclerc, Cluster algebras and quantum affine algebras, Duke Math. J. 154 (2010), no. 2, 265–341.
- [7] _____, Quantum Grothendieck rings and derived Hall algebras, J. Reine Angew. Math. 701 (2015), 77–126.
- [8] _____, Monoidal categorifications of cluster algebras of type A and D, in Symmetries, Integrable Systems and Representations, Springer Proc. Math. Stat. 40 (2013), 175–193.
- [9] _____, A cluster algebra approach to q-characters of Kirillov-Reshetikhin modules, J. Eur. Math. Soc. 18 (2016), no. 5, 1113–1159.
- [10] A. Kuniba, T. Nakanishi and J. Suzuki, Functional relations in solvable lattice models. I. Functional relations and representation theory, Internat. J. Modern Phys. A 9, (1994) no. 30, 5215– 5266.
- [11] S.-J. Kang, M. Kashiwara and M. Kim, Symmetric quiver Hecke algebras and R-matrices of quantum affine algebras II, Duke Math. J. 164 no.8 (2015), 1549–1602.
- [12] S.-J. Kang, M. Kashiwara, M. Kim and S-j. Oh, Monoidal categorification of cluster algebras, J. Amer. Math. Soc. 31 (2018), no. 2, 349–426.
- [13] _____, Symmetric quiver Hecke algebras and R-matrices of quantum affine algebras IV, Selecta Math. (N.S.), 22 (2016) 1987–2015.
- [14] M. Kashiwara, M. Kim, S.-j. Oh and E. Park, Monoidal categorification and quantum affine algebras, Compos. Math. 156 (2020), no. 5, 1039-1077
- [15] _____, Block decomposition for quantum affine algebras by the associated simply-laced root system, arXiv:2003.03265v1.
- [16] _____, Braid group action on the module category of quantum affine algebras, arXiv:2004.04939v1.
- [17] _____, *PBW* theoretic approach to the module category of quantum affine algebras, arXiv:2005.04838v2.
- [18] M. Kashiwara and S-j. Oh, Categorical relations between Langlands dual quantum affine algebras: Doubly laced types, J. Algebraic Combin., 49 (2019), 401–435.
- [19] H. Nakajima, Quiver varieties and finite-dimensional representations of quantum affine algebras, J. Amer. Math. Soc., 14 (2001), no. 1, 145–238.
- [20] _____, Quiver varieties and cluster algebras, Kyoto J. Math.51(2011), no. 1, 71–126.
- [21] S.-j. Oh and T. Scrimshaw, Categorical relations between Langlands dual quantum affine algebras: Exceptional cases, Comm. Math. Phys., 368 (1) (2019), 295–367.
- [22] S.-j. Oh and U. Suh, Twisted and folded Auslander-Reiten quiver and applications to the representation theory of quantum affine algebras, J. Algebra, 535 (2019), 53–132.
- [23] F. Qin, Triangular bases in quantum cluster algebras and monoidal categorification conjectures. Duke Math. J., 166 (12) (2017), 2337–2442.

(M. Kashiwara) Kyoto University Institute for Advanced Study, Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan & Korea Institute for Advanced Study, Seoul 02455, Korea

 $E\text{-}mail \ address: \texttt{masaki@kurims.kyoto-u.ac.jp}$

(M. Kim) DEPARTMENT OF MATHEMATICS, KYUNG HEE UNIVERSITY, SEOUL 02447, KOREA *E-mail address*: mkim@khu.ac.kr

(S.-j. Oh) DEPARTMENT OF MATHEMATICS, EWHA WOMANS UNIVERSITY, SEOUL 03760, KOREA *E-mail address*: sejin0920gmail.com

(E. Park) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SEOUL, SEOUL 02504, KOREA *E-mail address*: epark@uos.ac.kr

10