

Quasinormal modes and their anomalous behavior for black holes in $f(R)$ gravity

Almendra Aragón,^{1,*} P. A. González,^{1,†} Eleftherios Papantonopoulos,^{2,‡} and Yerko Vásquez^{3,§}

¹*Facultad de Ingeniería y Ciencias, Universidad Diego Portales,
Avenida Ejército Libertador 441, Casilla 298-V, Santiago, Chile.*

²*Physics Division, National Technical University of Athens, 15780 Zografou Campus, Athens, Greece.*

³*Departamento de Física y Astronomía, Facultad de Ciencias, Universidad de La Serena,
Avenida Cisternas 1200, La Serena, Chile.*

(Dated: March 12, 2024)

We study the propagation of scalar fields in the background of an asymptotically de-Sitter black hole solution in $f(R)$ gravity. The aim of this work is to analyze in modified theories of gravity the existence of an anomalous decay rate of the quasinormal modes (QNMs) of a massive scalar field which was recently reported in Schwarzschild black holes backgrounds, in which the longest-lived modes are the ones with higher angular number, for a scalar field mass smaller than a critical value, while that beyond this value the behavior is inverted. We study the QNMs for various overtone numbers and they depend on a parameter β which appears in the metric and characterizes the $f(R)$ gravity. For small β , i.e small deviations from the Schwarzschild-dS black hole the anomalous behaviour in the QNMs is present for the photon sphere modes, and the critical value of the mass of the scalar field depends on the parameter β while for large β , i.e large deviations the anomalous behaviour and the critical mass does not appear. Also, the critical mass of the scalar field increases when the overtone number increases until the $f(R)$ gravity parameter β approaches the near extremal limit at which the critical mass of the scalar field does not depend anymore on the overtone number. The imaginary part of the quasinormal frequencies is always negative leading to a stable propagation of the scalar fields in this background.

PACS numbers: 04.40.-b, 95.30.Sf, 98.62.Sb

Contents

I. Introduction	2
II. $f(R)$ Modified Gravity	3
III. Scalar perturbations	4
A. The photon sphere family	6
1. Photon sphere modes	6
2. Anomalous decay rate	7
3. Analysis using the WKB method	9
B. The de Sitter family	13
C. The dominance family	14
IV. Conclusions	15
Acknowledgments	16
References	16

*Electronic address: almendra.aragon@mail.udp.cl

†Electronic address: pablo.gonzalez@udp.cl

‡Electronic address: lpapa@central.ntua.gr

§Electronic address: yvasquez@userena.cl

I. INTRODUCTION

Modified theories of gravity in which the Einstein-Hilbert action is replaced with a generic form of $f(R)$ gravity have been introduced in an attempt to describe the early and late cosmological evolution. Another motivation to study such theories is the understanding of the existence of dark energy and dark matter consistent with the recent observations, without introducing new material ingredients that have not yet been detected by experiments [1–5]. Modifying the action not only affects the dynamics of the universe, it can also alter the dynamics at the galactic or solar system scales. Therefore, modified theories of gravity with curvature corrections, provide a deeper understanding of General Relativity (GR).

Assuming that the gravitational Lagrangian is not only a linear function of R , variable modified theories of gravity were considered describing the cosmic evolution at early times, in which the gravitational Lagrangian contains some of the four possible second-order curvature invariants. Also models in which higher order curvature invariants as functions of the Ricci scalar were introduced in the gravitational Lagrangian resulting to various $f(R)$ gravity models [6]–[15]. Although such theories exclude contributions from any curvature invariants other than R , they could also avoid the Ostrogradski instability [16] which proves to be problematic for general higher derivative theories [17].

One of the first modifications of the Einstein Lagrangian density was proposed in [18]. A more natural modification of the Einstein Lagrangian is to add terms R^n , like the Starobinsky model $f(R) = R + \alpha R^2$ [19]. For $n < 0$ such corrections become important in the late universe and can lead to self-accelerating vacuum solutions [20–23]. However, these models suffer from instabilities [24, 25] and there are strong constraints from the solar system [26]. A wide range of phenomena can be explained by considering different $f(R)$ functions. Some discussions have been performed, such as on gravitational wave detection [27, 28], early-time inflation [29], cosmological phases [30–32], the singularity problem [33], the stability of the solutions [34–36], and other different branches have been studied [37].

Spherically symmetric static solutions have been studied in $f(R)$ theories of gravity. In particular, it was shown that for a large class of models the Schwarzschild-de Sitter (SdS) metric is an exact solution of the field equations. However, in addition to the SdS metric, $f(R)$ theories typically also have new different solutions [38]. Also, static spherically symmetric perfect fluid solutions have been studied by matching the outside SdS-metric to the metric inside the mass distribution that leads to additional constraints that limit the allowed fluid configurations [39], while in [40] black hole solution were found with and without electric charge, and exact spherically symmetric solutions were discussed in [41–46]. Also, exact charged black hole solutions in $f(R)$ gravity theories with dynamic curvature in D -dimensions were discussed in [47].

A way to probe the behaviour and the stability of black holes is the study of their quasinormal modes (QNMs) and quasinormal frequencies (QNFs) [48–52]. The QNMs depend on the black hole parameters and probe field parameters, and on the fundamental constants of the system and they are independent of the initial conditions of the perturbations. The QNMs are described by complex frequencies, $\omega = \omega_R + i\omega_I$, in which the real part ω_R determines the oscillation timescale of the modes, while the complex part ω_I determines their exponential decaying timescale (for a review on QNM modes see [50, 53]).

The QNMs and QNFs have been calculated using various numerical and analytical techniques [54–59]. In the case of gravitational perturbations it was found that for the Schwarzschild and Kerr black hole background the longest-lived modes are always the ones with lower angular number ℓ . In the case of a massive probe scalar field it was found [60–63], at least for the overtone $n = 0$, that if the mass of the scalar field is small, then the longest-lived QNMs are those with a high angular number ℓ , whereas if the mass of the scalar field is large the longest-lived modes are those with a low angular number ℓ . This inverted behaviour appears when the mass of the scalar field exceeds a critical value, which corresponds to the value of the scalar field mass where the behavior of the decay rate of the QNMs is inverted and can be obtained from the condition $Im(\omega)_\ell = Im(\omega)_{\ell+1}$ in the *eikonal* limit, that is when $\ell \rightarrow \infty$. This anomalous decay rate for small mass scale of the scalar field was recently discussed in [64].

The study of the anomalous behaviour of QNMs was extended to other asymptotic geometries, such as, Schwarzschild-de Sitter and Schwarzschild-AdS black holes in [65]. It was found that the same behaviour is present in the Schwarzschild-de Sitter background, i.e., the absolute values of the imaginary part of the QNFs decay when the angular harmonic numbers increase if the mass of the scalar field is smaller than the critical mass, and they grow when the angular harmonic numbers increase, if the mass of the scalar field is larger than the critical mass. Also it was found that the increase of the value of the cosmological constant results in the increase of the value of the critical mass of the scalar field. It was also found that the anomalous behaviour is present in Schwarzschild-AdS black holes backgrounds; however, there is not a critical mass where the behavior in the decay rate is inverted. Also anomalous decay of QNMs in accelerating black holes was recently reported in [66], as well as, for Reissner-Nordström black holes [67]. Furthermore, the anomalous behaviour of QNMs was observed for fermionic field perturbations in Schwarzschild-de Sitter black holes [68].

Most of the $f(R)$ models have a very interesting property. If a conformal transformation is carried out from the original Jordan frame to the Einstein frame then, the conformal metric appears to be coupled minimally to gravity and

a new scalar field appears which is coupled minimally to the conformal metric and also a scalar potential is generated. The resulted theory can be considered as a scalar-tensor theory with a geometric (gravitational) scalar field. Then it was shown in [69, 70], that this geometric scalar field cannot dress a $f(R)$ black hole with hair, therefore, the non-hair theorem is respected in these models. Matter was introduced in [71] and a R^2 gravity model was discussed having a conformally coupled scalar field with associated Higgs-like potential and a static spherically symmetric black hole was found. However, the conformally coupled scalar field could not give to the black hole hair, because it is regulated by the same parameters that appear in the metric. A scalar field minimally coupled to gravity with a self-interacting potential in a $f(R)$ theory was discussed in [72]. An exact hairy black hole solution was found similar to the BTZ black hole. Also the process of scalarization of $f(R)$ gravity theories was investigated in [73].

As we already discussed the knowledge of the QNMs and QNFs can give us vital information of the behaviour, properties and the stability of black holes. In this work we will consider a specific $f(R)$ theory which accepts a black hole solution [74]. In this theory the Ricci scalar receives non-linear correction terms which they introduce new parameters in the metric functions of the black hole solution, expressing the change of the curvature altering the strength of the gravitational force. In this theory we will consider a massless and a massive test scalar field scattered off the background black hole and we will calculate the QNMs and QNFs. The motivation of this work is to calculate the scalar field perturbations and study the behaviour of QNMs and QNFs and see how they are effected by the presence of non-linear curvature terms which appear explicitly in the metric functions.

The manuscript is organized as follows: In Sec. II we give a brief review of the model considered for $f(R)$ gravity and its black hole solution. In Sec. III, we study the scalar field stability and we calculate the QNFs of scalar perturbations numerically by using the pseudospectral Chebyshev method for each family of modes. In Sec. III A 3, we perform an analysis using the WKB method to get some analytical insight. Finally, our conclusions are in Sec. IV.

II. $f(R)$ MODIFIED GRAVITY

We consider a generic action in $f(R)$ gravity depending on the Ricci scalar R given by

$$S = \frac{1}{2k} \int d^4x \sqrt{-g} f(R) + S_m, \quad (1)$$

where S_m is the matter content of the theory. In [74] a specific ansatz for the function $f(R)$ was considered

$$f(R) = R + \Lambda + \frac{R + \Lambda}{R/R_0 + 2/\alpha} \ln \frac{R + \Lambda}{R_c}, \quad (2)$$

where Λ corresponds to the cosmological constant, R_c is a constant of integration and $R_0 = 6\alpha^2/d^2$, with α and d being free parameters. Then considering a spherically symmetric metric

$$ds^2 = B(r)dt^2 - A(r)dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (3)$$

where $A(r) = B(r)^{-1}$, a vacuum spherically symmetric solution was found in [74] with $B(r) = 1 - \frac{2M}{r} + \beta r - \frac{\Lambda r^2}{3}$, where $\beta = \alpha/d$ is a real constant. For the range $R \gg \Lambda$ and $R/R_0 \gg 2/\alpha$ the action reduces to

$$f(R) = R + R_0 \ln \left(\frac{R}{R_c} \right), \quad (4)$$

and satisfies the solar system range of $r \ll d$. On the other hand, at large scales, for $\alpha \ll 1$ and $R \sim R_0 \sim \Lambda$, the action tends to $f(R) \sim R + \Lambda$, and agrees with cosmological observations. For metric (3) the curvature scalar is given by

$$R = \frac{6\beta}{r} - 4\Lambda, \quad (5)$$

and the action tends to a constant value asymptotically

$$f(R) \approx \left(-3\Lambda - \frac{3R_0\alpha\Lambda \ln(-3\Lambda/R_c)}{2R_0 - 4\alpha\Lambda} \right) + \mathcal{O}(r^{-1}). \quad (6)$$

It is worth mentioning that the solution in both regimes was obtained as perturbation around the Einstein-Hilbert action and it was shown that within this approximation the solutions are consistent with the modified gravity equations.

Also, the parameters of the theory were constrained by using the pioneer data, the flat rotation curve of galaxies and the cosmic microwave background radiation and Supernova Type Ia gold sample data giving $\beta \approx 10^{-26} m^{-1}$, $\beta d \approx 10^{-6}$ and $\beta d \ll 10^{-3}$, respectively [74]. However, the spacecraft's heat loss is a more plausible explanation for the anomaly than the cosmological one [75].

Our analysis consider a range of values of β , such that the metric represents a black hole solution with an event horizon and a cosmological horizon as it shown in Fig. 1. In this figure for a fixed value of the cosmological constant, we show the transition among black holes, near extremal ($r_H \approx r_\Lambda$) and naked singularity, when the β parameter goes from positive to negative values. Notice that, for $\beta = 0$, the spacetime is described by Schwarzschild-dS black hole. So, for β positive (negative) the cosmological horizon radius is larger (less) than Schwarzschild-de Sitter black hole, while for β positive (negative) the event horizon radius is less (larger) than Schwarzschild-dS black hole, when the spacetime describes black hole solutions.




FIG. 1: The behaviour of $B(r)$ as a function of r for different values of the parameters β with $M = 1$ and $\Lambda = 0.04$.

III. SCALAR PERTURBATIONS

In order to obtain the QNMs of scalar perturbations in the background of the metric (3) we consider the Klein-Gordon equation

$$\frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu \varphi) = -m^2 \varphi, \quad (7)$$

with suitable boundary conditions for a black hole geometry. In the above expression m is the mass of the scalar field φ . Now, by means of the following ansatz

$$\varphi = e^{-i\omega t} R(r) Y(\Omega), \quad (8)$$

the Klein-Gordon equation reduces to

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 B(r) \frac{dR}{dr} \right) + \left(\frac{\omega^2}{B(r)} - \frac{\ell(\ell+1)}{r^2} - m^2 \right) R(r) = 0, \quad (9)$$

where $\ell = 0, 1, 2, \dots$ corresponds to the eigenvalue of the Laplacian on the two-sphere. Now, defining $R(r) = \frac{F(r)}{r}$ and by using the tortoise coordinate r^* given by $dr^* = \frac{dr}{B(r)}$, the Klein-Gordon equation can be written as

$$\frac{d^2 F(r^*)}{dr^{*2}} - V_{eff}(r) F(r^*) = -\omega^2 F(r^*), \quad (10)$$

that corresponds to a one-dimensional Schrödinger-like equation with an effective potential $V_{eff}(r)$, parametrized as $V_{eff}(r^*)$ and given by

$$V_{eff}(r) = \frac{B(r)}{r^2} (\ell(\ell+1) + m^2 r^2 + B'(r)r), \quad (11)$$

and substituting the metric function $B(r)$ in the effective potential in Eq. (11) we obtain

$$V_{eff}(r) = \frac{1}{3} (3\beta^2 - \ell(\ell+1)\Lambda - 2\Lambda + 3m^2) + \frac{3\beta\ell(\ell+1) + 3\beta - 6m^2M + 2\Lambda M}{3r} + \frac{1}{9}r^2 (2\Lambda^2 - 3\Lambda m^2) + r(\beta m^2 - \beta\Lambda) - \frac{4M^2}{r^4} + \frac{-2\ell(\ell+1)M + 2M}{r^3} + \frac{\ell(\ell+1)}{r^2}. \quad (12)$$

Now, by considering $m = m_c$, as the value of the mass that cancels the divergence of order r^2 , i.e., $m_c = \sqrt{2\Lambda/3}$, and $\ell = 0$, the effective potential is

$$V_{eff}(r) = \beta^2 - \frac{4M^2}{r^4} + \frac{2M}{r^3} - \frac{2\Lambda M - 3\beta}{3r} - \frac{\beta\Lambda r}{3}. \quad (13)$$

So, we can note that for β negative (positive) the potential diverges positive (negative) at infinity, and in principle the existence of a critical mass could not be present in this case, due to the arguments given in [65].

Now, in order to solve numerically the differential equation (9) by using the pseudospectral Chebyshev method [76], it is convenient to perform the change of variable $y = (r - r_H)/(r_\Lambda - r_H)$. Thus, the values of the radial coordinate are limited to the range $[0, 1]$, and the radial equation (9) can be written as

$$B(y)R''(y) + \left(\frac{2(r_\Lambda - r_H)B(y)}{r_H + (r_\Lambda - r_H)y} + B'(y) \right) R'(y) + (r_\Lambda - r_H)^2 \left(\frac{\omega^2}{B(y)} - \frac{\ell(\ell+1)}{(r_H + (r_\Lambda - r_H)y)^2} - m^2 \right) R(y) = 0. \quad (14)$$

Also, in order to propose an ansatz for $R(y)$ we consider its behavior in the vicinity of the horizon ($y \rightarrow 0$)

$$R(y) = C_1 e^{-\frac{i\omega(r_\Lambda - r_H)}{B'(0)} \ln y} + C_2 e^{\frac{i\omega(r_\Lambda - r_H)}{B'(0)} \ln y}, \quad (15)$$

and at the cosmological horizon ($y = 1$)

$$R(y) = D_1 e^{-\frac{i\omega(r_\Lambda - r_H)}{B'(1)} \ln(1-y)} + D_2 e^{\frac{i\omega(r_\Lambda - r_H)}{B'(1)} \ln(1-y)}. \quad (16)$$

So, an ansatz that satisfy the requirements of only ingoing waves at the event horizon and at the cosmological horizon is

$$R(y) = e^{-\frac{i\omega(r_\Lambda - r_H)}{B'(0)} \ln y} e^{\frac{i\omega(r_\Lambda - r_H)}{B'(1)} \ln(1-y)} F(y). \quad (17)$$

Then, by inserting the above ansatz for $R(y)$ in Eq. (14), it is possible to obtain an equation for the function $F(y)$. The solution for the function $F(y)$ is assumed to be a finite linear combination of the Chebyshev polynomials, then it is inserted in the differential equation for $F(y)$. Also, the interval $[0, 1]$ is discretized at the Chebyshev collocation points. Then, the differential equation is evaluated at each collocation point. So, a system of algebraic equations is obtained, and it corresponds to a generalized eigenvalue problem, which is solved numerically to obtain the QNFs. It is worth mentioning that, for $\beta = 0$, the spacetime is described by the Schwarzschild-de Sitter black hole. The complex family of QNMs for this geometry was determined in [77] by using the WKB and Pöschl-Teller method. Also, it was shown that the frequencies all have a negative imaginary part, which means that the propagation of scalar field is stable in this background, and the presence of the cosmological constant leads to decrease of the real oscillation frequency and to a slower decay. High overtones were studied in [78]. Also, a family of purely imaginary modes was reported in [79] and an analysis of the photon sphere (PS) modes and de Sitter (dS) modes was recently performed in Ref. [65].

A. The photon sphere family

1. Photon sphere modes

The photon sphere QNMs, are represented by damped oscillations whose decay rate (ω_I) is directly connected to the instability timescale of null geodesics at the photon sphere. The photon sphere is a spherical trapping region of space where gravity is strong enough that photons are forced to travel in unstable circular orbits around a black hole. This region has a strong pull in the control of decay of perturbations and the QNMs with large frequencies. For instance, the decay timescale is related to the instability timescale of null geodesics near the photon sphere. For asymptotically dS black holes we find a family that can be traced back to the photon sphere and refer to them as PS modes. The dominant modes of this family are approached in the eikonal limit, where $\ell \rightarrow \infty$, for the mass of the scalar field lower than a critical value, and can be very well approximated with the WKB method. For this family of modes, higher angular momentum represents smaller decay rates, such that in the limit of interest, the most representative modes are those for which $\ell \rightarrow \infty$ [80, 81].

Now, in order to observe the behavior of photon sphere modes, we plot the behavior of the fundamental QNFs $n_{PS} = 0$ ¹ of massless scalar fields as a function of $M\beta$ with $\ell = 0$, see Fig. 2. We can observe that for small values of $M\beta$, the decay rate decreases when the cosmological constant increases. However, for large $M\beta$, the behavior is opposite, i.e, the decay rate increases. On the other hand, the frequency of the oscillation increases when $M\beta$ increases and decreases when the cosmological constant increases. In Fig. 3 we consider $\ell = 1$, note that the behavior of the QNFs is equivalent to the observed for $\ell = 0$ and small $M\beta$; however, there is not an inverted behavior, for the range of $M\beta$ values considered. Thus, when the cosmological constant increases the decay rate decreases, for small $M\beta$ and when $M\beta$ increases the decay rate converges to the same value and it does not depend on the value of the cosmological constant. On the other hand, the frequency of the oscillation increases when $M\beta$ increases and decreases when the cosmological constant increases, the same behavior was observed for $\ell = 0$.




FIG. 2: The behavior of $Im(\omega)M$ (left panel) and $Re(\omega)M$ (right panel) for the fundamental QNF as a function of $M\beta$ for massless scalar field with $\ell = 0$ and different values of the cosmological constant.

¹ n_{PS} corresponds to the overtone number of photon sphere modes




FIG. 3: The behaviour of $Im(\omega)M$ (left panel) and $Re(\omega)M$ (right panel) for the fundamental QNF as a function of $M\beta$ for massless scalar field with $\ell = 1$ and different values of the cosmological constant.

2. Anomalous decay rate

Now, we will study if in $f(R)$ modified gravity, in the presence of a cosmological constant, the photon sphere modes present an anomalous behavior, i.e the longest-lived modes are the ones with higher angular number, and also the presence of a critical mass, where beyond this value the behavior mentioned is inverted. So, we plot the behavior of $-Im(\omega)M$ as a function of the scalar field mass for different values of the parameters $M\beta$, ℓ and $n_{PS} = 0$ (left panels), and $n_{PS} = 1$ with $\ell > n_{PS}$ (right panels)², for a fixed values of the black hole mass and the cosmological constant, see Fig. 4. Also, we consider larger values of the parameter $M\beta$ with $n_{PS} = 0$, see Fig. 5. We can observe that the anomalous behavior along with the critical mass can be present for a certain range of values of the parameter $M\beta$. See, Fig. 4, where the anomalous behavior occurs, and the critical mass appears, for $M\beta = -0.06, -0.01$, and 0.01 ; and see Fig. 5, where the anomalous behavior does occur for $M\beta = 0.10$ and it does not occur and the critical mass does not appear, for $M\beta = 0.15$, and 0.20 . Note that, for $M\beta = 0.20$, the longest-lived modes always are the ones with smaller angular number, but for $M\beta = 0.15$, we cannot say the same.

As mentioned, when $M\beta$ increases the metric function could represents naked singularity, extremal black hole, near extremal black hole and black hole solutions. So, one of the effects of $M\beta$ is tuning the different cases, which have an important effect in the anomalous behavior, that is, for the near extremal case, see Fig. 4 top panel, the value of the critical mass is the same for the overtone numbers $n_{PS} = 0$ and $n_{PS} = 1$, which does not occur for the other case where the value of the critical mass increases when the overtone number increases. Also, we can observe that when the $M\beta$ parameter increases the critical mass is shifted by the effect of the $M\beta$ parameter, decreasing. So, there is a critical value of $M\beta$ where the critical mass is zero thereby there is not an anomalous behavior in the QNMs for larger values of $M\beta$. To explain this effect, we could say that the deviation of Schwarzschild-dS black hole, where the anomalous behavior and the critical mass have been observed, is given by the parameter $M\beta$ which also appears in the metric function. So for small values of this parameter, i.e small deviations Schwarzschild-dS black hole, the anomalous behavior occurs and the critical mass appears. However, for larger values of the parameter $M\beta$ i.e large deviations from Schwarzschild-dS black hole, the QNMs have a different behavior, where either the anomalous behavior and the critical mass are not present, this cut off point **could** numerically occurs for $M\beta \approx 0.10$, see Fig. 2, where the QNFs present an inverted behavior with respect to $M\beta$ for massless scalar field.

² We have left outside the case for $\ell = 0$, because the imaginary part of the QNFs exhibits a behavior other than $\ell > 0$. In general, this different behavior occurs for $\ell < n_{PS}$.




FIG. 4: The behavior of $-Im(\omega)M$ as a function of the scalar field mass for different values of the parameter $\ell = 0, 1, 2, 30$ and $n_{PS} = 0$ (left panels), and $\ell = 1, 2, 30$ and $n_{PS} = 1$ (right panels) with $M^2\Lambda = 0.04$, and from top to bottom $M\beta = -0.06, -0.01$, and 0.01 , respectively.




FIG. 5: The behavior of $-Im(\omega)M$ as a function of the scalar field mass mM for different values of the parameter $\ell = 0, 1, 2, 30$ and $n_{PS} = 0$ with $M^2\Lambda = 0.04$, and $M\beta = 0.1$ (top panel), $M\beta = 0.15$ (center panel) and $M\beta = 0.20$ (bottom panel).

3. Analysis using the WKB method

In this section, we use the method based on Wentzel-Kramers-Brillouin (WKB) in order to get some analytical insight of the behavior of the QNFs in the eikonal limit $\ell \rightarrow \infty$ [82–87]. It is worth mentioning that this method can be used for effective potentials which have the form of potential barriers that approach to a constant value at the horizon and spatial infinity [52]. The method consider the behavior of the effective potential near its maximum value r_{max}^* . So, by using the Taylor series expansion, the potential around its maximum is given by

$$V(r^*) = V(r_{max}^*) + \sum_{i=2}^{i=\infty} \frac{V^{(i)}}{i!} (r^* - r_{max}^*)^i. \quad (18)$$

where

$$V^{(i)} = \frac{d^i}{dr^{*i}} V(r^*)|_{r^*=r_{max}^*}, \quad (19)$$

corresponds to the i -th derivative of the potential with respect to r^* evaluated at the maximum of the potential r_{max}^* . Using the WKB approximation up to 6th order the QNFs are given by the following expression [88]

$$\omega^2 = V(r_{max}^*) - 2iU, \quad (20)$$

where

$$\begin{aligned} U = & N\sqrt{-V^{(2)}/2} + \frac{i}{64} \left(-\frac{1}{9} \frac{V^{(3)2}}{V^{(2)2}} (7 + 60N^2) + \frac{V^{(4)}}{V^{(2)}} (1 + 4N^2) \right) \\ & + \frac{N}{2^{3/2}288} \left(\frac{5}{24} \frac{V^{(3)4}}{(-V^{(2)})^{9/2}} (77 + 188N^2) + \frac{3}{4} \frac{V^{(3)2}V^{(4)}}{(-V^{(2)})^{7/2}} (51 + 100N^2) \right. \\ & \left. + \frac{1}{8} \frac{V^{(4)2}}{(-V^{(2)})^{5/2}} (67 + 68N^2) + \frac{V^{(3)}V^{(5)}}{(-V^{(2)})^{5/2}} (19 + 28N^2) + \frac{V^{(6)}}{(-V^{(2)})^{3/2}} (5 + 4N^2) \right), \end{aligned}$$

and $N = n_{PS} + 1/2$, with $n_{PS} = 0, 1, 2, \dots$, is the overtone number. Now, by defining $L^2 = \ell(\ell + 1)$, we find that for large values of ℓ , the maximum of the potential is approximately at

$$r_{max} \approx r_0 + \frac{1}{L^2} r_1 + \mathcal{O}(L^{-4}),$$

where

$$\begin{aligned} r_0 \approx & 3M - \frac{9M^2}{2}\beta + \frac{27M^3}{2}\beta^2 - \frac{405M^4}{8}\beta^3 + \mathcal{O}(\beta^4), \\ r_1 \approx & -\frac{1}{3}M(-1 + 9M^2(3m^2 - 2\Lambda))(-1 + 9\Lambda M^2) + \frac{1}{2}M^2(-1 + 9M^2(3m^2 - 2\Lambda))(-1 + 63M^2\Lambda)\beta + \\ & \frac{9}{4}M^3(2 + 9M^2\Lambda(13 + 216M^2\Lambda) - 27m^2(M^2 + 108M^4\Lambda))\beta^2 + \frac{9}{8}M^4(-35 + 9M^2(-2\Lambda(98 + 1485\Lambda M^2) + \\ & m^2(84 + 4455\Lambda M^2)))\beta^3 + \mathcal{O}(\beta^4), \end{aligned}$$

and

$$V(r_{max}^*) \approx V_0 L^2 + V_1 + \mathcal{O}(L^{-2}),$$

where

$$\begin{aligned} V_0 \approx & \left(\frac{1}{27M^2} - \frac{\Lambda}{3} \right) + \frac{\beta}{3M} + \frac{\beta^2}{4} - \frac{M\beta^3}{4} + \mathcal{O}(\beta^4), \\ V_1 \approx & -\frac{(2 + 9M^2(3m^2 - 2\Lambda))(-1 + 9\Lambda M^2)}{81M^2} + \left(\frac{10}{27M} + M \left(2m^2 - \frac{8\Lambda}{3} \right) + 3M^3(3m^2 - 2\Lambda)\Lambda \right) \beta + \\ & \left(\frac{23}{18} + \frac{1}{4}M^2(-3m^2(4 + 45\Lambda M^2) + 2\Lambda(8 + 45\Lambda M^2)) \right) \beta^2 + \frac{1}{4}M(-2 + 3M^2(3m^2(4 + 63\Lambda M^2) - 2\Lambda(8 + \\ & 63\Lambda M^2)))\beta^3 + \mathcal{O}(\beta^4). \end{aligned} \quad (21)$$

The second derivative of the potential evaluated at r_{max}^* yields the expression

$$V^{(2)}(r_{max}^*) \approx V_0^{(2)} L^2 + V_1^{(2)} + \mathcal{O}(L^{-2}), \quad (22)$$

where

$$\begin{aligned} V_0^{(2)} &\approx -\frac{2(1-9\Lambda M^2)^2}{729M^4} - \frac{2((-7+9\Lambda M^2)(-1+9\Lambda M^2))}{243M^3} \beta + \left(-\frac{32}{81M^2} + \frac{13\Lambda}{9} + \Lambda^2 M^2\right) \beta^2 \\ &\quad - \frac{(8+3\Lambda M^2(2+9\Lambda M^2))}{9M} \beta^3 + \mathcal{O}(\beta^4), \\ V_1^{(2)} &\approx \frac{2(1-9\Lambda M^2)^2(-8+9M^2(m^2(6-135\Lambda M^2)+2\Lambda(-1+45\Lambda M^2)))}{6561M^4} + \\ &\quad \frac{2(-1+9\Lambda M^2)(64+27M^2(4\Lambda-54\Lambda^2 M^2(3+11\Lambda M^2)+m^2(-16+9\Lambda M^2(23+99\Lambda M^2))))}{2187M^3} \beta + \\ &\quad \frac{(-116+9M^2(-3m^2(-25+9\Lambda M^2(13+18\Lambda M^2(8+63\Lambda M^2))))+\Lambda(5+27\Lambda M^2(14+3\Lambda M^2(41+252\Lambda M^2))))}{243M^2} \beta^2 \\ &\quad + \left(-\frac{344}{243M} + 4M(m^2-\Lambda) + 5M^3(9m^2-10\Lambda) + 6M^5(126m^2-107\Lambda)\Lambda^2 + 2106M^7(3m^2-2\Lambda)\Lambda^3\right) \beta^3 + \mathcal{O}(\beta^4), \end{aligned}$$

and the higher derivatives of the potential evaluated at r_{max}^* yields the expressions

$$\begin{aligned} V^{(3)}(r_{max}^*) &\approx \left(-\frac{4(-1+9\Lambda M^2)^3}{6561M^5} + \frac{4(1-9\Lambda M^2)^2}{243M^4} \beta + \frac{(-13+9\Lambda M^2)(-1+9\Lambda M^2)}{81M^3} \beta^2 + \frac{53-9\Lambda M^2(16+9\Lambda M^2)}{81M^2} \beta^3 \right. \\ &\quad \left. + \mathcal{O}(\beta^4)\right) L^2 + \mathcal{O}(L^0), \\ V^{(4)}(r_{max}^*) &\approx \left(-\frac{16(-1+9\Lambda M^2)^3}{19683M^6} - \frac{16(1-9\Lambda M^2)^2(-11+18\Lambda M^2)}{6561M^5} \beta + \frac{4(-1+9\Lambda M^2)(-7+27\Lambda M^2)}{81M^4} \beta^2 + \right. \\ &\quad \left. \frac{4(131+45\Lambda M^2(-20+9\Lambda M^2))}{243M^3} \beta^3 + \mathcal{O}(\beta^4)\right) L^2 + \mathcal{O}(L^0), \\ V^{(5)}(r_{max}^*) &\approx \left(-\frac{40(1-9\Lambda M^2)^4}{59049M^7} - \frac{40(13-9\Lambda M^2)(1-9\Lambda M^2)^3}{19683M^6} \beta + \frac{20(1-9\Lambda M^2)^2(-137+9\Lambda M^2(28+9\Lambda M^2))}{6561M^5} \beta^2 \right. \\ &\quad \left. + \frac{20(1-9\Lambda M^2)(-371+9\Lambda M^2(149+9\Lambda M^2(5+9\Lambda M^2)))}{2187M^4} \beta^3 + \mathcal{O}(\beta^4)\right) L^2 + \mathcal{O}(L^0), \\ V^{(6)}(r_{max}^*) &\approx \left(-\frac{16(1-9\Lambda M^2)^4(4+15\Lambda M^2)}{177147M^8} + \right. \\ &\quad \frac{64(-1+9\Lambda M^2)^3(5+6\Lambda M^2)}{19683M^7} \beta - \frac{4(1-9\Lambda M^2)^2(1517+9\Lambda M^2(-302+81\Lambda M^2))}{19683M^6} \beta^2 \\ &\quad \left. + \frac{4(-1+9\Lambda M^2)(15655+9\Lambda M^2(-9161+9\Lambda M^2(841+81\Lambda M^2)))}{19683M^5} \beta^3 + \mathcal{O}(\beta^4)\right) L^2 + \mathcal{O}(L^0). \end{aligned}$$

Then, using the expressions above with Eq. (20), we obtain the following analytical expression for the QNFs for large values of ℓ and small values of $M\beta$

$$\omega \approx \omega_{-1} L + \omega_0 + \omega_1 L^{-1} + \omega_2 L^{-2} + \mathcal{O}(L^{-3}), \quad (23)$$

where

$$\begin{aligned}
\omega_{-1} &\approx \frac{\sqrt{1-9\Lambda M^2}}{3\sqrt{3}M} + \frac{\sqrt{3}}{2\sqrt{1-9\Lambda M^2}}\beta - \frac{3\sqrt{3}M(2+9\Lambda M^2)}{8(1-9\Lambda M^2)^{3/2}}\beta^2 \\
&\quad + \frac{3\sqrt{3}M^2\sqrt{1-9\Lambda M^2}(-16+9\Lambda M^2(-13+18\Lambda M^2))}{16(-1+9\Lambda M^2)^3}\beta^3 + \mathcal{O}(\beta^4), \\
\omega_0 &\approx -i\frac{\sqrt{1-9\Lambda M^2}}{6\sqrt{3}M} - i\frac{4-9\Lambda M^2}{4\sqrt{3}\sqrt{1-9\Lambda M^2}}\beta + i\frac{3\sqrt{3}M(1+9\Lambda M^2(1+9\Lambda M^2))}{16(1-9\Lambda M^2)^{3/2}}\beta^2 \\
&\quad + i\frac{3\sqrt{3}M^2(-8+9\Lambda M^2(-13+9\Lambda M^2(-13+63\Lambda M^2)))}{32(1-9\Lambda M^2)^{5/2}}\beta^3 + \mathcal{O}(\beta^4), \\
\omega_1 &\approx -\frac{\sqrt{1-9\Lambda M^2}(-34+9M^2(-108m^2+61\Lambda))}{648M\sqrt{3}} + \frac{314-45\Lambda M^2(59+180\Lambda M^2)+972m^2(M^2+18\Lambda M^4)}{432\sqrt{3}\sqrt{1-9\Lambda M^2}}\beta \\
&\quad - \frac{-710M-5832M^7(135m^2-46\Lambda)\Lambda^2+108M^3(72m^2+5\Lambda)+243\Lambda M^5(396m^2+13\Lambda)}{576\sqrt{3}(1-9\Lambda M^2)^{3/2}}\beta^2 + \\
&\quad M^2\left(-1262+9M^2(\Lambda(-466+27\Lambda M^2(335+6\Lambda M^2(301-2736\Lambda M^2))))+324m^2(10+3\Lambda M^2(31+ \right. \\
&\quad \left. 18\Lambda M^2(-31+126\Lambda M^2))))\right)\beta^3 / \left(384\sqrt{3}(1-9\Lambda M^2)^{5/2}\right) + \mathcal{O}(\beta^4), \\
\omega_2 &\approx -i\frac{(137+45M^2(-648m^2+401\Lambda))(1-9\Lambda M^2)^{3/2}}{23328M\sqrt{3}} \\
&\quad + i\frac{\sqrt{1-9\Lambda M^2}(590+17496m^2M^2(2+27\Lambda M^2)-9\Lambda M^2(3593+30807\Lambda M^2))}{5184\sqrt{3}}\beta \\
&\quad + i\frac{M(7949+9M^2(648m^2(-23+9\Lambda M^2(-11+711\Lambda M^2))+\Lambda(8714-9\Lambda M^2(7858+252747\Lambda M^2))))}{6912\sqrt{3}\sqrt{1-9\Lambda M^2}}\beta^2 \\
&\quad + iM^2\left(-23674+81M^2(\Lambda(-2219+3\Lambda M^2(70514+627708\Lambda M^2-8621073\Lambda^2M^4)) \right. \\
&\quad \left. +72m^2(154+27\Lambda M^2(-65+27\Lambda M^2(-119+977\Lambda M^2))))\right)\beta^3 / \left(13824\sqrt{3}(1-9\Lambda M^2)^{3/2}\right) + \mathcal{O}(\beta^4).
\end{aligned}$$

The term proportional to $1/L^2$ is zero at the value of the critical mass m_c , which is given by

$$\begin{aligned}
m_c M &\approx \sqrt{\frac{137+18045M^2\Lambda}{29160}} - \frac{(1886-4185M^2\Lambda)M\beta}{60\sqrt{10}\sqrt{137+18045M^2\Lambda}} - \frac{3(4634227+134256255M^2\Lambda+1091552625M^4\Lambda^2)M^2\beta^2}{400\sqrt{10}(137+18045M^2\Lambda)^{3/2}} \\
&\quad - \frac{9(5019824246+76442598769M^2\Lambda-4248331450335M^4\Lambda^2-85433253134175M^6\Lambda^3)M^3\beta^3}{800\sqrt{10}(137+18045M^2\Lambda)^{5/2}} + \dots \quad (24)
\end{aligned}$$

and it is valid for small values of β and $n_{PS} = 0$. In all the above expressions we have performed a Taylor's expansion around $\beta = 0$ in order to obtain expressions that will be easy to handle analytically. For $\beta = 0$ we recover the result of critical mass for Schwarzschild-dS black holes [65], and for $\Lambda M^2 = 0.04$ and $n_{PS} = 0$ we obtain from (24) the value $m_c M = 0.186, 0.175, 0.168$, and 0.130 for $\beta = -0.06, -0.01, 0.01$, and 0.10 , respectively, which agrees with the numerical results shown in Fig. 4, thereby the value of the critical mass decreases when the beta parameter increases and it increases when ΛM^2 increases, see Fig. 6. Also, note that for small values of ΛM^2 there is a range of values for $M\beta$ where $m_c M$ becomes negative. Also, the WKB method proposes a critical scalar field mass for $M\beta = 0.20$, contrary to the observed via the pseudospectral Chebyshev method. However, the analysis performed with the WKB method is valid only for small values of β , because we have performed a Taylor's expansion around $\beta = 0$ to obtain the formula (24). Now, in order to check the correctness and accuracy of the pseudospectral Chebyshev method with respect to the analytical expression for the QNFs given by Eq. (23), we show in Table I, the values obtained with both methods. Also, we show the relative error, which is defined by

$$\epsilon_{Re(\omega)} = \frac{|Re(\omega_1)| - |Re(\omega_0)|}{|Re(\omega_0)|} \cdot 100\%, \quad (25)$$

$$\epsilon_{Im(\omega)} = \frac{|Im(\omega_1)| - |Im(\omega_0)|}{|Im(\omega_0)|} \cdot 100\%, \quad (26)$$

where ω_1 corresponds to the result from Eq. (23) and ω_0 denotes the result with the pseudospectral Chebyshev method. We can observe that the error does not exceed 0.0008 % in the imaginary part, and 0.0010 % in the real




FIG. 6: The behavior of the $m_c M$ as a function of $M\beta$ for different values of ΛM^2 with $n_{PS} = 0$.

part. Also, as it was observed, the frequencies all have a negative imaginary part, which means that the propagation of massive scalar fields is stable in this background.

TABLE I: Fundamental quasinormal frequencies ($n_{PS} = 0$) for massive scalar fields with $\ell = 10, 20$, and 30 in the background of black hole solution of $f(R)$ gravity with $M^2\Lambda = 0.04$, and $M\beta = 0.01$.

$mM = 0.02$				
ℓ	Pseudospectral method	WKB	$\epsilon_{Re(\omega)}$	$\epsilon_{Im(\omega)}$
10	$1.72651001 - 0.08356503i$	$1.72652389 - 0.08356574i$	0.0008	0.0008
20	$3.37194090 - 0.08348702i$	$3.37197238 - 0.08348760i$	0.0009	0.0007
30	$5.01712051 - 0.08347182i$	$5.01716774 - 0.08347239i$	0.0009	0.0007
$mM = 0.15$				
ℓ	Pseudospectral method	WKB	$\epsilon_{Re(\omega)}$	$\epsilon_{Im(\omega)}$
10	$1.72802056 - 0.08348145i$	$1.72803867 - 0.08348207i$	0.0010	0.0007
20	$3.37271554 - 0.08346511i$	$3.37274759 - 0.08346568i$	0.0010	0.0007
30	$5.01764129 - 0.08346193i$	$5.01768870 - 0.08346249i$	0.0009	0.0007
$mM = 0.30$				
ℓ	Pseudospectral method	WKB	$\epsilon_{Re(\omega)}$	$\epsilon_{Im(\omega)}$
10	$1.73263370 - 0.08322691i$	$1.73266525 - 0.08322649i$	0.0018	0.0005
20	$3.37508142 - 0.08339825i$	$3.37511532 - 0.08339874i$	0.0010	0.0006
30	$5.01923188 - 0.08343171i$	$5.01927986 - 0.083432269i$	0.0010	0.0007

As we mentioned, the Taylor's series have been performed around $\beta = 0$, so the WKB analysis is true for small values of β . In table II, we show the error in the quasinormal frequency between the WKB and pseudospectral method, for different values of mM , and $M\beta$. Note that for $M\beta = 0.20$ the error is 18 % in the imaginary part, and 31 % in the real part, approximately, that mean a loss of robustness and diminished validity in the results. Also, we can observe that the error increases slightly, when mM increases.

TABLE II: Fundamental quasinormal frequencies ($n_{PS} = 0$) for massive scalar fields with $\ell = 30$ in the background of black hole solution of $f(R)$ gravity with $M^2\Lambda = 0.04$, and different values of $M\beta$.

$mM = 0.02$				
$M\beta$	Pseudospectral method	WKB	$\epsilon_{Re(\omega)}$	$\epsilon_{Im(\omega)}$
-0.06	2.08314131 - 0.03055262i	2.25222758 - 0.03267338i	8.11689	6.94134
-0.01	4.35540688 - 0.07032092i	4.35546389 - 0.07032160i	0.00130895	0.000966995
0.01	5.01712051 - 0.08347182i	5.01716774 - 0.08347239i	0.000941377	0.000682865
0.10	7.42982610 - 0.13698770i	7.70221565 - 0.14017416i	3.66616	2.32609
0.15	8.55684679 - 0.16467397i	9.67805702 - 0.17771512i	13.1031	7.91938
0.20	9.59615293 - 0.19154767i	12.58587755 - 0.22617900i	31.1554	18.0797
$mM = 0.15$				
$M\beta$	Pseudospectral method	WKB	$\epsilon_{Re(\omega)}$	$\epsilon_{Im(\omega)}$
-0.06	2.08341635 - 0.03055125i	2.25253125 - 0.03267230i	8.11719	6.9426
-0.01	4.35588702 - 0.07031330i	4.35594414 - 0.07031399i	0.00131133	0.000981322
0.01	5.01764129 - 0.08346193i	5.01768870 - 0.08346249i	0.000944866	0.000670965
0.10	7.43044497 - 0.13696970i	7.70288435 - 0.14015649i	3.66653	2.32664
0.15	8.55749294 - 0.16465262i	9.67890740 - 0.17769503i	13.1045	7.92117
0.20	9.59681747 - 0.19152351i	12.58708647 - 0.22615793i	31.159	18.0836
$mM = 0.30$				
$M\beta$	Pseudospectral method	WKB	$\epsilon_{Re(\omega)}$	$\epsilon_{Im(\omega)}$
-0.06	2.08425622 - 0.03054706i	2.25345872 - 0.03266898i	8.11812	6.9464
-0.01	4.35735346 - 0.07029005i	4.35741099 - 0.07029073i	0.0013203	0.00096742
0.01	5.01923188 - 0.08343171i	5.01927986 - 0.08343226i	0.000955923	0.000659222
0.10	7.43233530 - 0.13691473i	7.70492677 - 0.14010254i	3.66764	2.32832
0.15	8.55946661 - 0.16458739i	9.68150472 - 0.17763367i	13.1087	7.92666
0.20	9.59884736 - 0.19144971i	12.59077885 - 0.22609358i	31.1697	18.0955

B. The de Sitter family

The de Sitter family of modes, are related to the accelerated expansion of the Universe, which is related to the surface gravity of the cosmological horizon of pure dS space. They correspond to purely imaginary modes which can be very well approximated by the pure dS QNMs. The pure dS modes are determined by two branches [89, 90], and the most important for our purposes corresponds to the lowest lying solution, given by

$$\omega_{\text{pure dS}} = -i\sqrt{\frac{\Lambda}{3}}\left(2n + \ell + \frac{3}{2} \pm \sqrt{\frac{9}{4} - 3\frac{m^2}{\Lambda}}\right), \quad (27)$$

where n is the overtone number.

In order to visualize the behavior of the QNFs for the dS modes, in Fig. 7 we plot $-Im(\omega)M$ for $\ell = 1$ and different values of the cosmological constant, note that there is a slowly decay rate when $M\beta$ increases, and when the cosmological constant decreases. Also, as before, we will study the behavior of the dS QNFs as a function of the




FIG. 7: The behavior of the fundamental dS mode ($n_{dS} = 0$) $-Im(\omega)M$ as a function of $M\beta$ obtained by using the pseudospectral Chebyshev method for massless scalar field with $\ell = 1$ and different values of the cosmological constant.

scalar field mass. So, we obtain some QNFs for different values of the parameters ℓ and $n_{dS} = 0$ ³, for a fixed values of the black hole mass and the cosmological constant, see table III for $M\beta = 0.01$. As we mentioned, this family is dominant for small masses and it can acquire a real part becoming complex, which depend on the scalar field mass and the angular number. Also, it is possible to observe in the table III that the decay rate always increases when the angular number increases, thereby the longest-lived modes are the ones with lower angular number and the anomalous behavior is not present in this family.

TABLE III: Fundamental dS modes ($n_{dS} = 0$) for massive scalar fields with $\ell = 0, 1, 2$ in the background of black hole solution of $f(R)$ gravity with $M^2\Lambda = 0.04$, and $M\beta = 0.01$.

ℓ	$mM = 0.02$	$mM = 0.05$	$mM = 0.10$
0	$-0.00096728i$	$-0.00623134i$	$-0.02841651i$
1	$-0.11303750i$	$-0.11914334i$	$-0.14367078i$
2	$-0.22668753i$	$-0.23290179i$	$0.39967626 - 0.25614428i$
ℓ	$mM = 0.15$	$mM = 0.20$	$mM = 0.25$
0	$-0.09297689i$	$0.09799882 - 0.17514088i$	$0.14338038 - 0.17214279i$
1	$-0.20354608i$	$0.23373131 - 0.25290461i$	$0.22754315 - 0.23202223i$
2	$0.40221808 - 0.25380624i$	$0.40581565 - 0.25048547i$	$0.41051976 - 0.24611626i$
ℓ	$mM = 0.30$	$mM = 0.35$	$mM = 0.40$
0	$0.17714386 - 0.16988129i$	$0.20756224 - 0.16809914i$	$0.23661390 - 0.16676332i$
1	$0.25091348 - 0.20524307i$	$0.27737304 - 0.19237303i$	$0.30313974 - 0.18446717i$
2	$0.41643654 - 0.24058189i$	$0.42382467 - 0.23371594i$	$0.43324930 - 0.22546868i$

C. The dominance family

In order to observe the dominance between the photon and de Sitter family we plot in Fig. 8 the imaginary part of the QNFs as a function of the scalar field mass, for different overtone numbers and $\ell = 0$, where the black points correspond to the purely imaginary QNFs, and the gray color points correspond to the complex QNFs for small values of the parameter $M\beta = 0.01$. The behavior is similar to the observed in Ref. [65]. We can recognize the two families for zero mass of the scalar field, a family of complex QNFs, and a purely imaginary family. As we have seen, the purely imaginary modes belong to the family of de Sitter modes, while the complex ones corresponds to the photon sphere modes.

In order to interpret this figure, first observe the black points, and $n_{dS} = 0$. This family is dominant for small masses and near $mM = 0.17$ it can acquire a real part becoming complex due to the continuity between the black and gray points, and also could be connect with the dS modes with $-Im(\omega)M \approx 1$. Also, for low values of the mass, it is possible to observe that for $n_{dS} = 1$ and $n_{dS} = 2$, can combine acquiring a real part becoming complex for $mM \approx 0.05$; the same occurs for the overtone numbers $n_{dS} = 3$ and $n_{dS} = 4$, and for $mM \approx 0.025$ becoming complex. Besides, for higher overtones numbers, there are dS modes purely imaginary for all the range of mass considered. On the other hand, for the photon sphere modes, gray points, we can see that for $n_{PS} = 0$ and small mass this branch is subdominant; however, for $mM \approx 0.155$ it begins to dominate.

Now, for a bigger value of the parameter $M\beta = 0.20$, see Fig. 9. As before, the black points correspond to the purely imaginary QNFs, and the gray points correspond to the complex QNFs. Note that for low values of the scalar field mass dS modes are dominant; however for larger mass the photon sphere modes are dominant. Also, it is possible to observe that dS modes with $n_{dS} = 0$ could to connect with dS modes with $n_{dS} = 1$ ($-Im(\omega)M \approx 1.10$). Besides, for higher overtones numbers, there is a purely imaginary family of dS modes for all the range of mass considered. Also, for the photon sphere modes, colored points, we can see that for $n_{PS} = 0$ and small mass this family is subdominant; however, for $mM \approx 0.13$ it begins to dominate.

³ n_{dS} corresponds to the overtone number of de Sitter modes




FIG. 8: The behavior of the imaginary part of the quasinormal frequencies $-Im(\omega)M$ as a function of the scalar field mass mM for different overtone numbers with $\ell = 0$, $M\beta = 0.01$, and $M^2\Lambda = 0.04$. Black points for purely imaginary QNFs and gray points for complex QNFs.




FIG. 9: The behavior of the imaginary part of the quasinormal frequencies $-Im(\omega)M$ as a function of the scalar field mass mM for different overtone numbers with $\ell = 0$, $M\beta = 0.20$, and $M^2\Lambda = 0.04$. Black points for purely imaginary QNFs and gray points for complex QNFs.

IV. CONCLUSIONS

In this work, we considered an asymptotically de-Sitter black hole solution of a specific $f(R)$ modified gravity as background. In order to see if strong curvature effects, that arise from non-linear terms in R in the action, can modify the behaviour of the QNMs. Thus, we studied the propagation of a probe scalar field and we analyzed the presence of anomalous behaviour in the quasinormal modes spectrum. First, we showed that the QNMs spectrum consist of two families: the photon sphere modes and the de Sitter modes, and their behaviour depends on the parameter β which characterizes the considered black hole solution of the $f(R)$ gravity.

For the photon sphere modes we found that for massless scalar field and small values of the parameter $M\beta$, the decay rate decreases when the cosmological constant increases, the behaviour is similar to the observed for Schwarzschild-dS black holes [65]. On the other hand, for large values of the parameter $M\beta$ the behaviour is inverted, i.e. the decay

rate increases when the cosmological constant increases.

Also, we found that the photon sphere modes show an anomalous behaviour that depends on the value of the parameter β and the mass of the scalar field. As in the case of the Schwarzschild-dS black holes, which corresponds to $\beta = 0$, there is a critical value of the mass of the scalar field, which depends on the cosmological constant, beyond which the QNMs have an anomalous behaviour. However, in the $f(R)$ gravity this critical value depends, not only on cosmological constant but also on the new parameter β , which expresses the presence of non-linear terms in the curvature. We found that the critical value of the scalar mass decreases when the parameter $M\beta$ increases, and there is a critical value of $M\beta$ where the critical mass is zero indicating that there is not an anomalous behaviour of the QNMs. Also, we saw that the critical value of the mass of the scalar field increases when the overtone number increases; however when the parameter $M\beta$ approaches a near extremal black hole, i.e. when the cosmological horizon approximates to the event horizon, the critical value of the mass of the scalar field does not depend of the overtone number.

On the other hand, for the dS modes we observed that for massless scalar field they are purely imaginary, and for massive scalar field they can acquire an oscillatory part becoming complex for some value of the mass and higher. Also, we do not found an anomalous behaviour for this family of modes.

Also, we found that the imaginary part of the quasinormal frequencies is always negative leading to a stable propagation of the scalar fields in this background for both families of modes. The PS family or the dS family may dominates the evolution of the perturbation depending on the values of the parameters. We showed that for small values of $M^2\Lambda$ the dS modes dominate for small values of the scalar field mass, while that the photon sphere modes dominate for larger values of the scalar field mass. Also, the decay rate of the PS modes increase with $M\beta$, while that the decay rate of the dS modes decrease, so it can occurs that for low values of $M\beta$ the PS modes dominate and for high values of $M\beta$ the dS modes dominate, depending on the value of $M^2\Lambda$.

An interesting extension of this work is to calculate the QNMs and QNFs of the $f(R)$ gravity theory if we allow the matter fields to backreact with the metric. Then we can study the properties of the resulting hairy black holes [72, 73] and possibly see the interplay effects of the geometric scalar present in the $f(R)$ theory and the matter scalar field minimally coupled to the modified gravity theory.

Acknowledgments

P. A. G. acknowledges the hospitality of the Universidad de La Serena where part of this work was undertaken. Y.V. acknowledge support by the Dirección de Investigación y Desarrollo de la Universidad de La Serena, Grant No. PR18142.

-
- [1] S. Nojiri and S. D. Odintsov, “Introduction to modified gravity and gravitational alternative for dark energy,” Conf C **0602061**, 06 (2006) [*Int. J. Geom. Meth. Mod. Phys.* **4**, 115 (2007)] [hep-th/0601213].
 - [2] E. J. Copeland, M. Sami and S. Tsujikawa, “Dynamics of dark energy,” *Int. J. Mod. Phys. D* **15**, 1753 (2006) [hep-th/0603057].
 - [3] S. Nojiri and S. D. Odintsov, “Unified cosmic history in modified gravity: from $F(R)$ theory to Lorentz non-invariant models,” *Phys. Rept.* **505**, 59 (2011) [arXiv:1011.0544 [gr-qc]].
 - [4] T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, “Modified Gravity and Cosmology,” *Phys. Rept.* **513**, 1 (2012) [arXiv:1106.2476 [astro-ph.CO]].
 - [5] S. Capozziello and M. Francaviglia, “Extended Theories of Gravity and their Cosmological and Astrophysical Applications,” *Gen. Rel. Grav.* **40**, 357-420 (2008) [arXiv:0706.1146 [astro-ph]].
 - [6] A. De Felice and S. Tsujikawa, “ $f(R)$ theories,” *Living Rev. Rel.* **13**, 3 (2010), [arXiv:1002.4928 [gr-qc]].
 - [7] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov, L. Sebastiani and S. Zerbini, “A Class of viable modified $f(R)$ gravities describing inflation and the onset of accelerated expansion,” *Phys. Rev. D* **77** (2008) 046009 [arXiv:0712.4017 [hep-th]].
 - [8] L. Pogosian and A. Silvestri, “The pattern of growth in viable $f(R)$ cosmologies,” *Phys. Rev. D* **77** (2008) 023503 Erratum: [*Phys. Rev. D* **81** (2010) 049901] [arXiv:0709.0296 [astro-ph]].
 - [9] P. Zhang, “Testing $f(R)$ gravity against the large scale structure of the universe.,” *Phys. Rev. D* **73** (2006) 123504 [astro-ph/0511218].
 - [10] B. Li and J. D. Barrow, “The Cosmology of $f(R)$ gravity in metric variational approach,” *Phys. Rev. D* **75** (2007) 084010 [gr-qc/0701111].
 - [11] Y. S. Song, H. Peiris and W. Hu, “Cosmological Constraints on $f(R)$ Acceleration Models,” *Phys. Rev. D* **76** (2007) 063517 [arXiv:0706.2399 [astro-ph]].
 - [12] S. Nojiri and S. D. Odintsov, “Modified $f(R)$ gravity unifying R^{**m} inflation with Lambda CDM epoch,” *Phys. Rev. D* **77** (2008) 026007 [arXiv:0710.1738 [hep-th]].

- [13] S. Nojiri and S. D. Odintsov, “Unifying inflation with LambdaCDM epoch in modified $f(R)$ gravity consistent with Solar System tests,” *Phys. Lett. B* **657** (2007) 238 [arXiv:0707.1941 [hep-th]].
- [14] S. Capozziello, C. A. Mantica and L. G. Molinari, “Cosmological perfect-fluids in $f(R)$ gravity,” *Int. J. Geom. Meth. Mod. Phys.* **16** (2018) no.01, 1950008 [arXiv:1810.03204 [gr-qc]].
- [15] J. Vainio and I. Vilja, “ $f(R)$ gravity constraints from gravitational waves,” *Gen. Rel. Grav.* **49** (2017) no.8, 99 [arXiv:1603.09551 [astro-ph.CO]].
- [16] M. Ostrogradsky, “Memoires sur les equations differentielles, relatives au probleme des isoperimetres,” *Mem. Acad. St. Petersburg* **6** (1850) no.4, 385.
- [17] R. P. Woodard, “Avoiding dark energy with $1/r$ modifications of gravity,” *Lect. Notes Phys.* **720** (2007) 403 [astro-ph/0601672].
- [18] H. A. Buchdahl, *Mon. Not. Roy. Astron. Soc.* **150**, 1 (1970).
- [19] A. A. Starobinsky, “A New Type of Isotropic Cosmological Models Without Singularity,” *Phys. Lett. B* **91** (1980) 99 [*Phys. Lett.* **91B** (1980) 99] [*Adv. Ser. Astrophys. Cosmol.* **3** (1987) 130].
- [20] S. M. Carroll, M. Hoffman and M. Trodden, “Can the dark energy equation - of - state parameter w be less than -1 ?,” *Phys. Rev. D* **68** (2003) 023509 [astro-ph/0301273].
- [21] S. M. Carroll, V. Duvvuri, M. Trodden and M. S. Turner, “Is cosmic speed - up due to new gravitational physics?,” *Phys. Rev. D* **70** (2004) 043528 [astro-ph/0306438].
- [22] S. Capozziello, “Curvature quintessence,” *Int. J. Mod. Phys. D* **11** (2002) 483 [gr-qc/0201033].
- [23] S. Capozziello, V. F. Cardone, S. Carloni and A. Troisi, “Curvature quintessence matched with observational data,” *Int. J. Mod. Phys. D* **12** (2003) 1969 [astro-ph/0307018].
- [24] M. E. Soussa and R. P. Woodard, “The force of gravity from a Lagrangian containing inverse powers of the ricci scalar,” *Gen. Rel. Grav.* **36** (2004) 855 [astro-ph/0308114].
- [25] V. Faraoni and S. Nadeau, “The Stability of modified gravity models,” *Phys. Rev. D* **72** (2005) 124005 [gr-qc/0511094].
- [26] T. Chiba, “ $1/R$ gravity and scalar - tensor gravity,” *Phys. Lett. B* **575** (2003) 1 [astro-ph/0307338].
- [27] C. Corda, “Primordial production of massive relic gravitational waves from a weak modification of General Relativity,” *Astropart. Phys.* **30**, 209 (2008) [arXiv:0812.0483 [gr-qc]].
- [28] C. Corda, “Massive relic gravitational waves from $f(R)$ theories of gravity: Production and potential detection,” *Eur. Phys. J. C* **65**, 257 (2010) [arXiv:1007.4077 [gr-qc]].
- [29] K. Bamba and S. D. Odintsov, “Inflation and late-time cosmic acceleration in non-minimal Maxwell- $F(R)$ gravity and the generation of large-scale magnetic fields,” *JCAP* **0804**, 024 (2008) [arXiv:0801.0954 [astro-ph]].
- [30] S. Nojiri and S. D. Odintsov, “Modified $f(R)$ gravity consistent with realistic cosmology: From matter dominated epoch to dark energy universe,” *Phys. Rev. D* **74**, 086005 (2006) [hep-th/0608008].
- [31] S. Nojiri and S. D. Odintsov, “Modified gravity and its reconstruction from the universe expansion history,” *J. Phys. Conf. Ser.* **66** (2007) 012005 [hep-th/0611071].
- [32] S. Nojiri and S. D. Odintsov, “The Future evolution and finite-time singularities in $F(R)$ -gravity unifying the inflation and cosmic acceleration,” *Phys. Rev. D* **78**, 046006 (2008) [arXiv:0804.3519 [hep-th]].
- [33] T. Kobayashi and K. i. Maeda, “Can higher curvature corrections cure the singularity problem in $f(R)$ gravity?,” *Phys. Rev. D* **79**, 024009 (2009) [arXiv:0810.5664 [astro-ph]].
- [34] V. Faraoni, “Modified gravity and the stability of de Sitter space,” *Phys. Rev. D* **72**, 061501 (2005) [gr-qc/0509008].
- [35] S. Capozziello, S. Nojiri, S. D. Odintsov and A. Troisi, “Cosmological viability of $f(R)$ -gravity as an ideal fluid and its compatibility with a matter dominated phase,” *Phys. Lett. B* **639**, 135 (2006) [astro-ph/0604431].
- [36] L. Amendola, R. Gannouji, D. Polarski and S. Tsujikawa, “Conditions for the cosmological viability of $f(R)$ dark energy models,” *Phys. Rev. D* **75**, 083504 (2007) [gr-qc/0612180].
- [37] M. Akbar and R. G. Cai, “Thermodynamic Behavior of Field Equations for $f(R)$ Gravity,” *Phys. Lett. B* **648**, 243 (2007) [gr-qc/0612089].
- [38] T. Multamaki and I. Vilja, “Spherically symmetric solutions of modified field equations in $f(R)$ theories of gravity,” *Phys. Rev. D* **74** (2006) 064022 [astro-ph/0606373].
- [39] T. Multamaki and I. Vilja, “Static spherically symmetric perfect fluid solutions in $f(R)$ theories of gravity,” *Phys. Rev. D* **76** (2007) 064021 [astro-ph/0612775].
- [40] A. de la Cruz-Dombriz, A. Dobado and A. L. Maroto, “Black Holes in $f(R)$ theories,” *Phys. Rev. D* **80** (2009) 124011 Erratum: [*Phys. Rev. D* **83** (2011) 029903] [arXiv:0907.3872 [gr-qc]].
- [41] S. H. Hendi, B. Eslam Panah and S. M. Mousavi, “Some exact solutions of $F(R)$ gravity with charged (a)dS black hole interpretation,” *Gen. Rel. Grav.* **44**, 835 (2012) [arXiv:1102.0089 [hep-th]].
- [42] L. Sebastiani and S. Zerbini, “Static Spherically Symmetric Solutions in $F(R)$ Gravity,” *Eur. Phys. J. C* **71** (2011) 1591 [arXiv:1012.5230 [gr-qc]].
- [43] S. H. Hendi and D. Momeni, “Black hole solutions in $F(R)$ gravity with conformal anomaly,” *Eur. Phys. J. C* **71**, 1823 (2011) [arXiv:1201.0061 [gr-qc]].
- [44] S. Asgari and R. Saffari, “Vacuum Solution of a Linear Red-Shift Based Correction in $f(R)$ Gravity,” *Gen. Rel. Grav.* **44**, 737 (2012) [arXiv:1104.5108 [gr-qc]].
- [45] S. G. Ghosh, S. D. Maharaj and U. Papnoi, “Radiating Kerr-Newman black hole in $f(R)$ gravity,” *Eur. Phys. J. C* **73**, no. 6, 2473 (2013) [arXiv:1208.3028 [gr-qc]].
- [46] S. H. Hendi, B. Eslam Panah and R. Saffari, “Exact solutions of three-dimensional black holes: Einstein gravity versus $F(R)$ gravity,” *Int. J. Mod. Phys. D* **23**, no. 11, 1450088 (2014) [arXiv:1408.5570 [hep-th]].
- [47] Z. Y. Tang, B. Wang and E. Papantonopoulos, “Exact charged black hole solutions in D -dimensions in $f(R)$ gravity,”

- [arXiv:1911.06988 [gr-qc]].
- [48] T. Regge and J. A. Wheeler, “*Stability of a Schwarzschild singularity*,” *Phys. Rev.* **108**, 1063 (1957).
 - [49] F. J. Zerilli, “*Gravitational field of a particle falling in a schwarzschild geometry analyzed in tensor harmonics*,” *Phys. Rev. D* **2**, 2141 (1970).
 - [50] K. D. Kokkotas and B. G. Schmidt, “*Quasinormal modes of stars and black holes*,” *Living Rev. Rel.* **2**, 2 (1999) [gr-qc/9909058].
 - [51] H. P. Nollert, “*TOPICAL REVIEW: Quasinormal modes: the characteristic ‘sound’ of black holes and neutron stars*,” *Class. Quant. Grav.* **16**, R159 (1999).
 - [52] R. A. Konoplya and A. Zhidenko, “*Quasinormal modes of black holes: From astrophysics to string theory*,” *Rev. Mod. Phys.* **83**, 793 (2011) [arXiv:1102.4014 [gr-qc]].
 - [53] E. Berti, V. Cardoso and A. O. Starinets, “*Quasinormal modes of black holes and black branes*,” *Class. Quant. Grav.* **26**, 163001 (2009) [arXiv:0905.2975 [gr-qc]].
 - [54] S. I. Finazzo, R. Rougemont, M. Zaniboni, R. Critelli and J. Noronha, “*Critical behavior of non-hydrodynamic quasinormal modes in a strongly coupled plasma*,” *JHEP* **1701**, 137 (2017) [arXiv:1610.01519 [hep-th]].
 - [55] P. A. González, E. Papantonopoulos, J. Saavedra and Y. Vásquez, “*Superradiant Instability of Near Extremal and Extremal Four-Dimensional Charged Hairy Black Hole in anti-de Sitter Spacetime*,” *Phys. Rev. D* **95**, no. 6, 064046 (2017) [arXiv:1702.00439 [gr-qc]].
 - [56] E. Abdalla, B. Cuadros-Melgar, R. Fontana, J. de Oliveira, E. Papantonopoulos and A. Pavan, “*Instability of a Reissner-Nordström-AdS black hole under perturbations of a scalar field coupled to the Einstein tensor*,” *Phys. Rev. D* **99**, no.10, 104065 (2019) [arXiv:1903.10850 [gr-qc]].
 - [57] P. A. Gonzalez, Y. Vasquez and R. N. Villalobos, “*Perturbative and nonperturbative fermionic quasinormal modes of Einstein-Gauss-Bonnet-AdS black holes*,” *Phys. Rev. D* **98**, no. 6, 064030 (2018) [arXiv:1807.11827 [gr-qc]].
 - [58] R. Bécár, P. A. González, E. Papantonopoulos and Y. Vásquez, “*Quasinormal modes of three-dimensional rotating Hořava AdS black hole and the approach to thermal equilibrium*,” *Eur. Phys. J. C* **80**, no. 7, 600 (2020) doi:10.1140/epjc/s10052-020-8169-2 [arXiv:1906.06654 [gr-qc]].
 - [59] A. Aragón, R. Bécár, P. A. González and Y. Vásquez, “*Perturbative and nonperturbative quasinormal modes of 4D Einstein-Gauss-Bonnet black holes*,” *Eur. Phys. J. C* **80** (2020) no.8, 773 [arXiv:2004.05632 [gr-qc]].
 - [60] R. A. Konoplya and A. V. Zhidenko, “*Decay of massive scalar field in a Schwarzschild background*,” *Phys. Lett. B* **609**, 377 (2005) [gr-qc/0411059].
 - [61] R. A. Konoplya and A. Zhidenko, “*Stability and quasinormal modes of the massive scalar field around Kerr black holes*,” *Phys. Rev. D* **73**, 124040 (2006) [gr-qc/0605013].
 - [62] S. R. Dolan, “*Instability of the massive Klein-Gordon field on the Kerr spacetime*,” *Phys. Rev. D* **76**, 084001 (2007) [arXiv:0705.2880 [gr-qc]].
 - [63] O. J. Tattersall and P. G. Ferreira, “*Quasinormal modes of black holes in Horndeski gravity*,” *Phys. Rev. D* **97**, no. 10, 104047 (2018) [arXiv:1804.08950 [gr-qc]].
 - [64] M. Lagos, P. G. Ferreira and O. J. Tattersall, “*Anomalous decay rate of quasinormal modes*,” *Phys. Rev. D* **101**, no. 8, 084018 (2020) [arXiv:2002.01897 [gr-qc]].
 - [65] A. Aragón, P. A. González, E. Papantonopoulos and Y. Vásquez, “*Anomalous decay rate of quasinormal modes in Schwarzschild-dS and Schwarzschild-AdS black holes*,” *JHEP* **2008** (2020) 120 [arXiv:2004.09386 [gr-qc]].
 - [66] K. Destounis, R. D. B. Fontana and F. C. Mena, “*Counterexamples to strong cosmic censorship in asymptotically flat black hole spacetimes*,” [arXiv:2006.01152 [gr-qc]].
 - [67] R. D. B. Fontana, P. A. González, E. Papantonopoulos and Y. Vásquez, “*Anomalous decay rate of quasinormal modes in Reissner-Nordström black holes*,” *Phys. Rev. D* **103** (2021) no.6, 064005 [arXiv:2011.10620 [gr-qc]].
 - [68] A. Aragón, R. Bécár, P. A. González and Y. Vásquez, “*Massive Dirac quasinormal modes in Schwarzschild-de Sitter black holes: Anomalous decay rate and fine structure*,” *Phys. Rev. D* **103** (2021) no.6, 064006 [arXiv:2009.09436 [gr-qc]].
 - [69] P. Cañate, L. G. Jaime and M. Salgado, “*Spherically symmetric black holes in $f(R)$ gravity: Is geometric scalar hair supported ?*,” *Class. Quant. Grav.* **33**, no.15, 155005 (2016) [arXiv:1509.01664 [gr-qc]].
 - [70] P. Cañate, “*A no-hair theorem for black holes in $f(R)$ gravity*,” *Class. Quant. Grav.* **35**, no.2, 025018 (2018)
 - [71] J. Sultana and D. Kazanas, “*A no-hair theorem for spherically symmetric black holes in R^2 gravity*,” *Gen. Rel. Grav.* **50**, no.11, 137 (2018) [arXiv:1810.02915 [gr-qc]].
 - [72] T. Karakasis, E. Papantonopoulos, Z. Y. Tang and B. Wang, “*Black holes of $(2+1)$ -dimensional $f(R)$ gravity coupled to a scalar field*,” [arXiv:2101.06410 [gr-qc]].
 - [73] Z. Y. Tang, B. Wang, T. Karakasis and E. Papantonopoulos, “*Curvature Scalarization of Black Holes in $f(R)$ Gravity*,” [arXiv:2008.13318 [gr-qc]].
 - [74] R. Saffari and S. Rahvar, “ *$f(R)$ Gravity: From the Pioneer Anomaly to the Cosmic Acceleration*,” *Phys. Rev. D* **77**, 104028 (2008) [arXiv:0708.1482 [astro-ph]].
 - [75] S. G. Turyshev, V. T. Toth, G. Kinsella, S. C. Lee, S. M. Lok and J. Ellis, “*Support for the thermal origin of the Pioneer anomaly*,” *Phys. Rev. Lett.* **108** (2012), 241101 [arXiv:1204.2507 [gr-qc]].
 - [76] J. P. Boyd, Chebyshev and Fourier Spectral Methods. Dover Books on Mathematics. Dover Publications, Mineola, NY, second ed., 2001.
 - [77] A. Zhidenko, “*Quasinormal modes of Schwarzschild de Sitter black holes*,” *Class. Quant. Grav.* **21**, 273 (2004)
 - [78] R. A. Konoplya and A. Zhidenko, “*High overtones of Schwarzschild-de Sitter quasinormal spectrum*,” *JHEP* **0406**, 037 (2004)
 - [79] A. Jansen, “*Overdamped modes in Schwarzschild-de Sitter and a Mathematica package for the numerical computation of*

- quasinormal modes*,” *Eur. Phys. J. Plus* **132**, no. 12, 546 (2017)
- [80] V. Cardoso, J. L. Costa, K. Destounis, P. Hintz and A. Jansen, “*Quasinormal modes and Strong Cosmic Censorship*,” *Phys. Rev. Lett.* **120**, no.3, 031103 (2018) [arXiv:1711.10502 [gr-qc]].
 - [81] K. Destounis, “*Charged Fermions and Strong Cosmic Censorship*,” *Phys. Lett. B* **795**, 211-219 (2019) [arXiv:1811.10629 [gr-qc]].
 - [82] B. Mashhoon, “*Quasi-normal modes of a black hole*,” Third Marcel Grossmann Meeting on General Relativity 1983.
 - [83] B. F. Schutz and C. M. Will, “*Black Hole Normal Modes: A Semianalytic Approach*,” *Astrophys. J. Lett.* **291**, L33 (1985).
 - [84] S. Iyer and C. M. Will, “*Black Hole Normal Modes: A WKB Approach. 1. Foundations and Application of a Higher Order WKB Analysis of Potential Barrier Scattering*,” *Phys. Rev. D* **35**, 3621 (1987).
 - [85] R. A. Konoplya, “*Quasinormal behavior of the d-dimensional Schwarzschild black hole and higher order WKB approach*,” *Phys. Rev. D* **68**, 024018 (2003) [gr-qc/0303052].
 - [86] J. Matyjasek and M. Opala, “*Quasinormal modes of black holes. The improved semianalytic approach*,” *Phys. Rev. D* **96**, no. 2, 024011 (2017) [arXiv:1704.00361 [gr-qc]].
 - [87] R. A. Konoplya, A. Zhidenko and A. F. Zinhailo, “*Higher order WKB formula for quasinormal modes and grey-body factors: recipes for quick and accurate calculations*,” *Class. Quant. Grav.* **36**, 155002 (2019) [arXiv:1904.10333 [gr-qc]].
 - [88] Y. Hatsuda, “*Quasinormal modes of black holes and Borel summation*,” *Phys. Rev. D* **101**, no. 2, 024008 (2020) [arXiv:1906.07232 [gr-qc]].
 - [89] D. P. Du, B. Wang and R. K. Su, “*Quasinormal modes in pure de Sitter space-times*,” *Phys. Rev. D* **70**, 064024 (2004) [hep-th/0404047].
 - [90] A. Lopez-Ortega, “*Quasinormal modes of D-dimensional de Sitter spacetime*,” *Gen. Rel. Grav.* **38**, 1565-1591 (2006) [arXiv:gr-qc/0605027 [gr-qc]].