A COMPARISON PRINCIPLE FOR HIGHER ORDER NONLINEAR HYPOELLIPTIC HEAT OPERATORS ON GRADED LIE GROUPS

MICHAEL RUZHANSKY AND NURGISSA YESSIRKEGENOV

ABSTRACT. In this paper we present a comparison principle for higher order nonlinear hypoelliptic heat operators on graded Lie groups. Moreover, using the comparison principle we obtain blow-up type results and global in *t*-boundedness of solutions of nonlinear equations for the heat *p*-sub-Laplacian on stratified Lie groups. In particular, this paper generalises and extends previous results obtained by the first author and Suragan in [RS18].

1. INTRODUCTION

A connected simply connected Lie group \mathbb{G} is called a graded Lie group if its Lie algebra admits a gradation. The graded Lie groups form the subclass of homogeneous nilpotent Lie groups admitting homogeneous hypoelliptic left-invariant differential operators ([Mil80], [tER97], see also a discussion in [FR16, Section 4.1]). These operators are called Rockland operators from the Rockland conjecture, solved by Helffer and Nourrigat [HN79]. So, we understand by a Rockland operator any leftinvariant homogeneous hypoelliptic differential operator on \mathbb{G} . Thus, the considered setting includes the higher order operators on \mathbb{R}^n as well as higher order hypoelliptic invariant differential operators on the Heisenberg group, on general stratified Lie groups, and on general graded Lie groups.

Let us also recall that the standard Lebesgue measure is the Haar measure for \mathbb{G} . Let $\Omega \subset \mathbb{G}$ be a bounded set with smooth boundary. We denote the Sobolev space by $S^{a,p}(\Omega) = S^{a,p}_{\mathcal{R}}(\Omega)$, for a > 0 and $p \in (1, \infty) \cup \{\infty_o\}$, defined by the norm

$$||u||_{S^{a,p}(\Omega)} := \left(\int_{\Omega} \left(\left| \mathcal{R}^{\frac{a}{\nu}} u(x) \right|^p + |u(x)|^p \right) dx \right)^{\frac{1}{p}}, \tag{1.1}$$

where ν is the homogeneous order of the Rockland operator \mathcal{R} . We have allowed ourselves to write $\|\cdot\|_{L^{\infty}(\mathbb{G})} = \|\cdot\|_{L^{\infty_{o}}(\mathbb{G})}$ for the supremum norm, in the notation of [FR16, Chapter 4]. Let us also define the functional class $S_{0}^{a,p}(\Omega)$ to be the completion of $C_{0}^{\infty}(\Omega)$ in the norm (1.1). For a general discussion of Sobolev spaces on graded Lie groups we refer to [FR16, Chapter 4] and [FR17].

²⁰¹⁰ Mathematics Subject Classification. 35G20, 22E30.

Key words and phrases. Nonlinear hypoelliptic heat equation, Rockland operator, p-sub-Laplacian, comparison principle, graded Lie group, stratified Lie group, global solution, blow-up.

MR was supported by the EPSRC Grant EP/R003025/1, by the Leverhulme Research Grant RPG-2017-151, and by the FWO Odysseus 1 grant G.0H94.18N: Analysis and Partial Differential Equations. NY was supported by the FWO Odysseus 1 grant G.0H94.18N: Analysis and Partial Differential Equations and by the LMS Early Career Fellowship grant ECF-1819-12.

In this paper we study the higher order nonlinear hypoelliptic heat equation for u = u(t, x),

$$u_{t} - \sum_{j=1}^{n_{2}} \mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} \left(\left| \mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} u \right|^{p_{j}-2} \mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} u \right) u = \alpha \sum_{i=1}^{n_{1}} |u|^{q_{i}-1} u + \beta \sum_{j=1}^{n_{2}} |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u|^{r_{j}} + \gamma \sum_{k=1}^{n_{3}} |u|^{s_{k}-1} u$$

$$(1.2)$$

for $x \in \Omega$ and t > 0, with the initial-boundary conditions

$$u = 0, \quad x \in \partial\Omega, \quad t > 0, \tag{1.3}$$

$$u(0,x) = u_0(x), \quad x \in \Omega, \tag{1.4}$$

where $a_1, a_2 \geq 0$, and $\alpha, \beta, \gamma \in \mathbb{R}$, and $n_1, n_2, n_3 \in \mathbb{N}$, $p_j > 1$ and

$$q_i \left\{ \begin{array}{ll} \geq 1, & \text{if } \alpha > 0, \\ > 0, & \text{if } \alpha < 0, \end{array} \right. s_k \left\{ \begin{array}{ll} \geq 1, & \text{if } \gamma > 0, \\ > 0, & \text{if } \gamma < 0, \end{array} \right. r_j \left\{ \begin{array}{ll} > 1, & \text{if } \beta > 0, \\ > 0, & \text{if } \beta < 0. \end{array} \right.$$
(1.5)

Here, ν_1 and ν_2 are the homogeneous orders of the Rockland operators \mathcal{R}_1 and \mathcal{R}_2 , respectively. We also assume that the initial data satisfies

$$u_0 \in S_0^{a,\infty}(\Omega), \quad u_0 \ge 0,$$

where $a = \max\{a_1, a_2\}.$

Definition 1.1. Set $Q_T = (0,T) \times \Omega$, $S_T = (0,T) \times \partial\Omega$, $\partial Q_T = S_T \cup \{\{0\} \times \overline{\Omega}\}$, $p = \max\{p_j\}$ and $m = \max\{p_j, q_i, r_j, s_k\}$. A nonnegative function u(t, x) is called a weak super- (sub-) solution of (1.2)-(1.4) on Q_T if it satisfies

$$u \in C([0,T) \times \overline{\Omega}) \cap L^m((0,T); S_0^{a,m}(\Omega)), \ \partial_t u \in L^2((0,T); L^2(\Omega)),$$
$$u(0,x) \ge (\le) u_0, \ u|_{\partial\Omega} \ge (\le) 0,$$

$$\begin{aligned} \iint_{Q_T} \partial_t u \phi + \sum_{j=1}^{n_2} |\mathcal{R}_1^{\frac{a_1}{\nu_1}} u|^{p_j - 2} \mathcal{R}_1^{\frac{a_1}{\nu_1}} u \cdot \mathcal{R}_1^{\frac{a_1}{\nu_1}} \phi dx dt \\ \geq (\leq) \iint_{Q_T} \left(\alpha \sum_{i=1}^{n_1} |u|^{q_i - 1} u + \beta \sum_{j=1}^{n_2} |\mathcal{R}_2^{\frac{a_2}{\nu_2}} u|^{r_j} + \gamma \sum_{k=1}^{n_3} |u|^{s_k - 1} u \right) \phi dx dt, \end{aligned}$$

for all $\phi \in C(\overline{Q_T}) \cap L^p((0,T); S_0^{a_1,p}(\Omega))$ such that $\phi \ge 0, \phi|_{S_T} = 0$. Then *u* is called a weak solution if it is a super-solution and a sub-solution. Here and after, we use T_{\max} to denote the maximal existence time.

Our goal in this paper is to give a simple proof of a comparison principle for the initial boundary value problem for higher order nonlinear hypoelliptic heat operators on graded Lie groups using pure algebraic relations, inspired by the works [Att12] and [ZL13].

The structure of this paper is as follows. Section 2 establishes a comparison principle for the problem (1.2)-(1.4). Then, in Section 3, using the comparison principle, we investigate the blow-up or the boundedness of solution of (1.2)-(1.4) depending on the signs of α , β , γ , and relations between parameters p_j , q_i , r_j , s_k , and on u_0 .

2. A Comparison principle on graded Lie groups

In this section we state a comparison principle for the problem (1.2)-(1.4).

Theorem 2.1. Assume that $u, v \in L_{loc}^{\infty}$ $((0,T); S^{a,\infty}(\Omega))$ are sub- and super-solutions of (1.2)-(1.4), respectively. Assume also that at least one of the parameters α , β and γ be positive or $\alpha = \beta = \gamma = 0$. Let $r_j \geq \frac{p_j}{2}$ if $\beta > 0$. Then we have $u \leq v$ on Q_T .

Remark 2.2. In the special case $n_1 = n_2 = n_3 = 1$, $\beta = 0$ and $\alpha \gamma \leq 0$, Theorem 2.1 was obtained in [RS18].

The proof of the comparison principle mostly based on the following algebraic lemma (see e.g. [Att12, Lemma 2.1]).

Lemma 2.3. Let $\sigma > 1$. For all $\vec{a}, \vec{b} \in \mathbb{R}^N$, we have

$$\left\langle |\vec{a}|^{\sigma-2}\vec{a} - |\vec{b}|^{\sigma-2}\vec{b}, \vec{a} - \vec{b} \right\rangle \ge \frac{4}{\sigma^2} ||\vec{a}|^{\frac{\sigma-2}{2}}\vec{a} - |\vec{b}|^{\frac{\sigma-2}{2}}\vec{b}\Big|^2.$$

Proof of Theorem 2.1. First, let us consider the case α, β and $\gamma > 0$. Denote $\phi := \max\{u-v, 0\}$, hence $\phi(0, x) = 0$ and $\phi(t, x)|_{x \in \partial\Omega} = 0$. By the definitions of sub- and super-solutions, using ϕ as the test function, for any $\tau \in (0, T)$, we have

$$\frac{1}{2} \int_{\Omega} \phi^{2}(\tau, x) dx = \int_{0}^{\tau} \int_{\Omega} \frac{1}{2} \partial_{t} \left(\phi^{2}(t, x) \right) dx dt = \int_{0}^{\tau} \int_{\Omega} \partial_{t} \phi \phi dx dt \\
\leq -\sum_{j=1}^{n_{2}} \int_{0}^{\tau} \int_{\{\phi(t, \cdot)>0\}} \left(|\mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} u|^{p_{j}-2} \mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} u - |\mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} v|^{p_{j}-2} \mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} v \right) \cdot \mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} \phi dx dt \\
+ \beta \sum_{j=1}^{n_{2}} \int_{0}^{\tau} \int_{\{\phi(t, \cdot)>0\}} \left(|\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u|^{r_{j}} - |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v|^{r_{j}}| \right) \phi dx dt \\
+ \alpha \sum_{i=1}^{n_{1}} \int_{0}^{\tau} \int_{\{\phi(t, \cdot)>0\}} \left(|u|^{q_{i}-1}u - |v|^{q_{i}-1}v) \phi dx dt \right) \\
+ \gamma \sum_{k=1}^{n_{3}} \int_{0}^{\tau} \int_{\{\phi(t, \cdot)>0\}} \left(|u|^{s_{k}-1}u - |v|^{s_{k}-1}v) \phi dx dt \right) \\
I_{4}$$
(2.1)

By Lemma 2.3, for I_1 we have

$$I_{1} \geq \sum_{j=1}^{n_{2}} \frac{4}{p_{j}^{2}} \int_{0}^{\tau} \int_{\{\phi(t,\cdot)>0\}} \left| |\mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} u|^{\frac{p_{j}-2}{2}} \mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} u - |\mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} v|^{\frac{p_{j}-2}{2}} \mathcal{R}_{1}^{\frac{a_{1}}{\nu_{1}}} v \right|^{2} dx dt.$$
(2.2)

Let us now estimate the term I_2 . We put $h(s) = s^{\frac{2r_j}{p_j}}$ for $s \ge 0$. Given that $r_j \ge \frac{p_j}{2}$, we have $h'(s) = \frac{2r_j}{p_j} s^{\frac{2r_j - p_j}{p_j}}$. Then, by the mean value theorem we have $\left| \left| \mathcal{R}_2^{\frac{a_2}{\nu_2}} u \right|^{r_j} - \left| \mathcal{R}_2^{\frac{a_2}{\nu_2}} v \right|^{r_j} \right|^2 \le Ch'(\theta)^2 \left| \left| \mathcal{R}_2^{\frac{a_2}{\nu_2}} u \right|^{p_j/2} - \left| \mathcal{R}_2^{\frac{a_2}{\nu_2}} v \right|^{p_j/2} \right|^2$, for some $0 \leq \theta \leq \max\left(|\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}}u|^{p_{j}/2}, |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}}v|^{p_{j}/2}\right)$.

A direct computation yields that

$$\left| |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u|^{p_{j}/2} - |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v|^{p_{j}/2} \right|^{2} \leqslant \left| |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u|^{(p_{j}-2)/2} \mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u - |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v|^{(p_{j}-2)/2} \mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v \right|^{2}.$$

Taking into account $u, v \in L^{\infty}((0,T); S^{a,\infty}(\Omega))$, it follows that

$$\left| |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u|^{r_{j}} - |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v|^{r_{j}} \right|^{2} \leqslant C \left| |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u|^{(p_{j}-2)/2} \mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u - |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v|^{(p_{j}-2)/2} \mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v \right|^{2}, \quad (2.3)$$

where C is a positive constant depending on r_j, p_j and $\max\{|\mathcal{R}_2^{\frac{a_2}{\nu_2}}u|^{p_j/2}, |\mathcal{R}_2^{\frac{a_2}{\nu_2}}v|^{p_j/2}\}$. On the other hand, by Young's inequality we have

$$I_{2} \leqslant \sum_{j=1}^{n_{2}} \epsilon_{j} \int_{0}^{\tau} \int_{\{\phi(t,\cdot)>0\}} \left| |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u|^{r_{j}} - |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v|^{r_{j}} \right|^{2} dx dt + \sum_{j=1}^{n_{2}} C(\epsilon_{j}) \int_{0}^{\tau} \int_{\{\phi(t,\cdot)>0\}} \phi^{2} dx dt. \quad (2.4)$$

A combination of (2.3) and (2.4) leads to

$$I_{2} \leqslant C \sum_{j=1}^{n_{2}} \epsilon_{j} \int_{0}^{\tau} \int_{\{\phi(t,\cdot)>0\}} \left| |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u|^{(p_{j}-2)/2} \mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} u - |\mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v|^{(p_{j}-2)/2} \mathcal{R}_{2}^{\frac{a_{2}}{\nu_{2}}} v \right|^{2} dx dt + \sum_{j=1}^{n_{2}} C(\epsilon_{j}) \int_{0}^{\tau} \int_{\{\phi(t,\cdot)>0\}} \phi^{2} dx dt. \quad (2.5)$$

For I_3 , by the mean value theorem we obtain

$$I_3 \le \sum_{i=1}^{n_1} q_i \|u\|_{L^{\infty}}^{q_i-1} \int_0^\tau \int_{\{\phi(t,\cdot)>0\}} \phi^2 dx dt.$$
(2.6)

Similarly, for I_4 we have

$$I_4 \le \sum_{k=1}^{n_3} s_k \|u\|_{L^{\infty}}^{s_k-1} \int_0^\tau \int_{\{\phi(t,\cdot)>0\}} \phi^2 dx dt.$$
(2.7)

Choosing $0 < \epsilon_j < 4/(\beta C p_j^2)$ and combining the estimates (2.1), (2.2), (2.5), (2.6) and (2.7), we obtain for any $\tau \in (0, T)$ that

$$\int_{\Omega} \phi^2(\tau) dx$$

$$\leq C\left(\alpha, \beta, \epsilon, q_i, s_k, r_j, p_j, \|u\|_{L^{\infty}}, \max\{|\mathcal{R}_2^{\frac{a_2}{\nu_2}}u|^{p_j/2}, |\mathcal{R}_2^{\frac{a_2}{\nu_2}}v|^{p_j/2}\}\right) \int_0^{\tau} \int_{\Omega} \phi^2 dx dt. \quad (2.8)$$

Then by Gronwall's lemma we conclude that $\phi \equiv 0$ almost everywhere.

The case $\alpha = \beta = \gamma = 0$ is trivial.

Now, we discuss the case, when not all, but at least one of the parameters α , β , γ is positive. Note that I_3 is positive, since for $q_i > 0$ we have

$$\begin{cases} |u|^{q_i-1}u - |v|^{q_i-1}v = u^{q_i} - v^{q_i} > 0, & \text{for } u > v > 0\\ |u|^{q_i-1}u - |v|^{q_i-1}v = u^{q_i} + |v|^{q_i} > 0, & \text{for } u > 0 > v\\ |u|^{q_i-1}u - |v|^{q_i-1}v = -|u|^{q_i} + |v|^{q_i} > 0, & \text{for } 0 > u > v. \end{cases}$$

Similarly, one can verify that I_4 is positive for $s_k > 0$. Therefore, in the case when $\alpha < 0$ or $\beta < 0$ (or $\gamma < 0$) by dropping I_3 or I_2 (or I_4), respectively, we can always get (2.8).

3. Some applications to nonlinear equations for the heat p-sub-Laplacian

In this section, we give some applications of Theorem 2.1 to nonlinear equations for the heat *p*-sub-Laplacian on stratified Lie groups. These groups are an important class of graded Lie groups, investigated thoroughly by Folland [Fol75]. There are many different, equivalent ways to define a stratified Lie group (see, for example, [BLU07, FS82] or [FR16, RS19] for the Lie group and Lie algebra points of view, respectively). A Lie group $\mathbb{G} = (\mathbb{R}^N, \circ)$ is called a stratified Lie group if it satisfies the following two conditions:

• for every $\lambda > 0$ the dilation $\delta_{\lambda} : \mathbb{R}^N \to \mathbb{R}^N$ defined by

$$\delta_{\lambda}(x) \equiv \delta_{\lambda}\left(x^{'}, \dots, x^{(r)}\right) := \left(\lambda x^{'}, \dots, \lambda^{r} x^{(r)}\right)$$

is an automorphism of the group \mathbb{G} , where $x' \equiv x^{(1)} \in \mathbb{R}^{N_1}$ and $x^{(k)} \in \mathbb{R}^{N_k}$ for $k = 2, \ldots, r$ with $N_1 + \cdots + N_r = N$ and $\mathbb{R}^N = \mathbb{R}^{N_1} \times \cdots \times \mathbb{R}^{N_r}$.

• let X_1, \ldots, X_{N_1} be the left invariant vector fields on \mathbb{G} such that $X_j(0) = \frac{\partial}{\partial x_j}|_0$ for $j = 1, \ldots, N_1$. Then, for every $x \in \mathbb{R}^N$ the Hörmander condition

 $\operatorname{rank}(\operatorname{Lie}\{X_1,\ldots,X_{N_1}\})=N$

holds, that is, X_1, \ldots, X_{N_1} with their iterated commutators span the whole Lie algebra of the group \mathbb{G} .

Let us also recall that the left invariant vector field X_j has an explicit form given by (see, e.g. [FR16, Section 3.1.5])

$$X_{j} = \frac{\partial}{\partial x'_{j}} + \sum_{l=2}^{r} \sum_{m=1}^{N_{l}} a_{j,m}^{(l)} \left(x', \dots, x^{(l-1)} \right) \frac{\partial}{\partial x_{m}^{(l)}}.$$
 (3.1)

Throughout this section, we will also use the following notations

$$\nabla_H := (X_1, \ldots, X_{N_1})$$

for the horizontal gradient,

$$\mathcal{L}_p f := \nabla_H \left(|\nabla_H f|^{p-2} \nabla_H f \right), \quad 1$$

for the p-sub-Laplacian, and

$$|x'| = \sqrt{x_1'^2 + \dots + x_{N_1}'^2}$$

for the Euclidean norm on \mathbb{R}^{N_1} .

Using (3.1) one can observe that (see, e.g. [RS17])

$$|\nabla_H | x'|^b | = b |x'|^{b-1}, \qquad (3.3)$$

and

$$\nabla_H \left(\frac{x'}{|x'|^b} \right) = \frac{N_1 - b}{|x'|^b} \tag{3.4}$$

for all $b \in \mathbb{R}, x' \in \mathbb{R}^{N_1}$ and $|x'| \neq 0$.

Let us first consider the following initial boundary value problem for the *p*-sub-Laplacian, 1 ,

$$\begin{cases}
 u_t - \mathcal{L}_p u = \alpha |u|^{q-1} u + \beta |\nabla_H u|^r, & x \in \Omega, \quad t > 0, \\
 u(t, x) = 0, & x \in \partial\Omega, \quad t > 0, \\
 u(0, x) = u_0(x), & x \in \Omega,
\end{cases}$$
(3.5)

where $u_0(x) \ge 0$, $u_0(x) \ne 0$, $u_0 \in S_0^{1,\infty}(\Omega)$, and the parameters α , β , q and r will be determined later. By Definition 1.1 let us recall that T_{max} is the maximal existence time of a weak solution of (3.5).

Theorem 3.1. Let $\Omega \subset \mathbb{G}$ be a bounded open set in a stratified Lie group with N_1 being the dimension of the first stratum. Assume that α , β , p, q and r in (3.5) satisfy one of the following conditions:

(i) $\alpha < 0, \beta > 0, and p, r > 1$ with $p \le r + 1$ and $p/2 \le r < q$; (ii) $\alpha > 0, \beta < 0, and p > 1, q \ge 1$ with p < r + 1 and $q \le r$.

Then a weak solution of (3.5) is globally in t-bounded, that is, there exists a constant M depending only on p, q, r, α , β , N_1 , Ω and u_0 such that for every T > 0 we have $0 \le u \le M$ on (0, T).

Proof of Theorem 3.1. Part (i). For convenience, we assume that $\beta = -\alpha = 1$. Set $R' := \max_{x=(x',x'')\in\Omega} |x'|$. Then, since Ω is bounded, we get $R' < \infty$. For any $x = (x',x'') \in \Omega$, let $x_0 = (x'_0, x''_0) \in \mathbb{G} \setminus \Omega$ and $\varepsilon \in (0,1)$ be such that $\varepsilon \leq |x'_0 - x'| < R' + 1$. We also introduce the following notations

$$V_1(t,x) := K_1 e^{\sigma_1 R}, \quad R = |x' - x'_0|, \quad x = (x', x'') \in \Omega,$$

and

$$\mathcal{M}_p w := w_t - \mathcal{L}_p w + w^q - |\nabla_H w|^r.$$

Let us now find suitable positive K_1 and σ_1 such that $V_1(t, x)$ is a super-solution of (3.5). By using the identities (3.3) and (3.4), we observe that

$$|\nabla_H V_1|^r = K_1^r e^{\sigma_1 R r} \sigma_1^r$$

and

$$\begin{aligned} \mathcal{L}_{p}V_{1} &= \nabla_{H} \left(\left| \nabla_{H}K_{1}e^{\sigma_{1}R} \right|^{p-2} \nabla_{H}K_{1}e^{\sigma_{1}R} \right) \\ &= \nabla_{H} \left(K_{1}^{p-2}\sigma_{1}^{p-2}e^{\sigma_{1}|x'-x_{0}'|(p-2)}K_{1}\sigma_{1}e^{\sigma_{1}|x'-x_{0}'|} \frac{x'-x_{0}'}{|x'-x_{0}'|} \right) \\ &= \nabla_{H} \left(K_{1}^{p-1}\sigma_{1}^{p-1}e^{\sigma_{1}|x'-x_{0}'|(p-1)} \frac{x'-x_{0}'}{|x'-x_{0}'|} \right) \\ &= K_{1}^{p-1}\sigma_{1}^{p-1}\sigma_{1}(p-1)e^{\sigma_{1}|x'-x_{0}'|(p-1)} + K_{1}^{p-1}\sigma_{1}^{p-1}e^{\sigma_{1}|x'-x_{0}'|(p-1)} \frac{N_{1}-1}{|x'-x_{0}'|} \\ &= (p-1)\sigma_{1}^{p}K_{1}^{p-1}e^{(p-1)\sigma_{1}R} + \frac{N_{1}-1}{R}\sigma_{1}^{p-1}K_{1}^{p-1}e^{\sigma_{1}R(p-1)}. \end{aligned}$$

Thus, we have

$$\mathcal{M}_{p}V_{1} = -(p-1)\sigma_{1}^{p}K_{1}^{p-1}e^{(p-1)\sigma_{1}R} - \frac{N_{1}-1}{R}\sigma_{1}^{p-1}K_{1}^{p-1}e^{(p-1)\sigma_{1}R} + K_{1}^{q}e^{q\sigma_{1}R} - K_{1}^{r}e^{r\sigma_{1}R}\sigma_{1}^{r}.$$

Now, we need to find σ_{1} and K_{1} such that $\mathcal{M}_{p}V_{1} \ge 0$, that is,

$$(p-1)\sigma_1^p K_1^{p-1} e^{(p-1)\sigma_1 R} + \frac{N_1 - 1}{R} \sigma_1^{p-1} K_1^{p-1} e^{(p-1)\sigma_1 R} + K_1^r e^{r\sigma_1 R} \sigma_1^r \leqslant K_1^q e^{q\sigma_1 R}.$$

Multiplying both sides of the inequality by $K_1^{-p+1}e^{-(p-1)\sigma_1 R}$, we derive that

$$(p-1)\sigma_1^p + \frac{N_1 - 1}{R}\sigma_1^{p-1} + K_1^{r-p+1}e^{(r-p+1)\sigma_1R}\sigma_1^r \leqslant K_1^{q+1-p}e^{(q+1-p)\sigma_1R}.$$
 (3.6)

Taking into account $\varepsilon \leq R < R' + 1$, we see that in order to prove (3.6) it is sufficient to show

$$(p-1)\sigma_1^p + \frac{N_1 - 1}{\varepsilon}\sigma_1^{p-1} + K_1^{r-p+1}e^{(r-p+1)\sigma_1(R'+1)}\sigma_1^r \leqslant K_1^{q+1-p}$$

Thus, to have $\mathcal{M}_p V_1 \ge 0$ we can choose

$$\sigma_1 = \frac{1}{(r-p+1)(R'+1)},$$

$$K_1 = \max\left\{ (2e\sigma_1^r)^{1/(q-r)}, \left(2\left((p-1)\sigma_1^p + \frac{N_1-1}{\varepsilon}\sigma_1^{p-1} \right) \right)^{1/(q+1-p)} \right\}$$

when r+1 > p, and

$$\sigma_1 = 1, \quad K_1 = \max\left\{2^{1/(q-r)}, \left(2\left(p-1+\frac{N_1-1}{\varepsilon}\right)\right)^{1/(q+1-p)}\right\}$$

when r + 1 = p. We also need that $K_1 \ge ||u_0||_{L^{\infty}(\Omega)}$ such that $V_1(0, x) = K_1 e^{\sigma_1 R} \ge u_0(x)$. Obviously, we also have $V_1(t, x) \ge 0 = u(t, x)$ on $\partial\Omega$. Therefore, $V_1(t, x)$ is a super-solution of (3.5). Then, Theorem 2.1 concludes that

$$0 \le u(t,x) \le K_1 e^{\sigma_1(R'+1)} < \infty, \quad R' = \max_{x = (x',x'') \in \Omega} |x'|.$$
(3.7)

Note that the right-hand side of (3.7) is independent of t, hence u(t, x) is globally in t-bounded.

Part (ii). In this case, we may assume that $\alpha = -\beta = 1$. We recall from Part (i) that $R' = \max_{x=(x',x'')\in\Omega} |x'| < \infty$ and $\varepsilon \leq |x'_0 - x'| < R' + 1$ for any $x = (x',x'') \in \Omega$, where $x_0 = (x'_0, x''_0) \in \mathbb{G} \setminus \Omega$ and $\varepsilon \in (0, 1)$.

First, let us consider the case r > q. Here, we will use the following notations

$$V_2(t,x) := \frac{K_2}{\sigma_2} R^{\sigma_2}, \quad \sigma_2 = \frac{p}{p-1}, \quad R = |x' - x'_0|, \quad x \in \Omega,$$

and

$$\mathcal{N}_p w := w_t - \mathcal{L}_p w - w^q + |\nabla_H w|^r.$$

Now, we need to find a suitable positive K_2 such that $V_2(t, x)$ is a super-solution of (3.5). By using the identities (3.3) and (3.4), we observe that

$$\mathcal{L}_{p}V_{2} = \nabla_{H} \left(\left| \nabla_{H} \left(\frac{K_{2}}{\sigma_{2}} R^{\sigma_{2}} \right) \right|^{p-2} \nabla_{H} \left(\frac{K_{2}}{\sigma_{2}} R^{\sigma_{2}} \right) \right)$$
$$= \left(\frac{K_{2}}{\sigma_{2}} \right)^{p-1} \nabla_{H} \left(\sigma_{2}^{p-2} R^{(\sigma_{2}-1)(p-2)} \sigma_{2} R^{\sigma_{2}-1} \frac{x'-x'_{0}}{|x'-x'_{0}|} \right)$$
$$= K_{2}^{p-1} \nabla_{H} \left(R^{(\sigma_{2}-1)(p-1)} \frac{x'-x'_{0}}{|x'-x'_{0}|} \right)$$
$$= N_{1} K_{2}^{p-1}.$$

Then we have

$$\mathcal{N}_p V_2 = -N_1 K_2^{p-1} + K_2^r R^{\frac{r}{p-1}} - \left(\frac{K_2}{\sigma_2}\right)^q R^{\frac{qp}{p-1}}$$

From this, we have

$$\mathcal{N}_p V_2 \ge 0 \iff K_2^r R^{\frac{r}{p-1}} \ge N_1 K_2^{p-1} + \left(\frac{K_2}{\sigma_2}\right)^q R^{\frac{qp}{p-1}}.$$

Thus, it is sufficient to choose K_2 such that

$$K_2^r R^{\frac{r}{p-1}} \ge 2N_1 K_2^{p-1},\tag{3.8}$$

$$K_2^r R^{\frac{r}{p-1}} \ge 2\left(\frac{K_2}{\sigma_2}\right)^q R^{\frac{qp}{p-1}}.$$
 (3.9)

,

Note that the inequality (3.8) is satisfied if we take

$$K_2 \ge \left(\frac{2N_1}{\varepsilon^{\frac{r}{p-1}}}\right)^{\frac{1}{r-p+1}}$$

provided that r > p - 1. We divide inequality (3.9) by $K_2^q R^{\frac{r}{p-1}}$ to derive

$$K_2^{r-q} \ge \frac{2}{\sigma_2^q} R^{\frac{qp-r}{p-1}}.$$

For $qp \ge r$, we can set

$$K_2 \ge \left(\frac{2}{\sigma_2^q}\right)^{\frac{1}{r-q}} (R'+1)^{\frac{qp-r}{(p-1)(r-q)}},$$

while for qp < r, we can set

$$K_2 \ge \left(\frac{2}{\sigma_2^q}\right)^{\frac{1}{r-q}} \varepsilon^{\frac{qp-r}{(p-1)(r-q)}}.$$

We also need that $K_2 \ge \frac{\sigma_2 \|u_0\|_{L^{\infty}}}{\varepsilon^{\sigma_2}}$ to have $V_2(0, x) \ge u_0$. Thus, taking K_2 as follows $\int \sigma_2 \|u_0\|_{L^{\infty}} \left(2N_1\right)^{\frac{1}{r-p+1}} \left(2\right)^{\frac{1}{r-q}} \left(2V_1\right)^{\frac{q_p-r}{r-q}} \left(2\right)^{\frac{1}{r-q}} \left(2\right)^{\frac{1}{r-q}} \left(2V_1\right)^{\frac{1}{r-q}} \left($

$$K_2 \ge \max\left\{\frac{\sigma_2 \|u_0\|_{L^{\infty}}}{\varepsilon^{\sigma_2}}, \left(\frac{2N_1}{\varepsilon^{\frac{r}{p-1}}}\right)^{r-p+1}, \left(\frac{2}{\sigma_2^q}\right)^{r-q} (R'+1)^{\frac{qp-r}{(p-1)(r-q)}}, \left(\frac{2}{\sigma_2^q}\right)^{r-q} \varepsilon^{\frac{qp-r}{(p-1)(r-q)}}\right\}$$
we obtain $M |V| \ge 0$ and $V(0, \pi) \ge q$. It is clear that $V(t, \pi) \ge 0 = u(t, \pi)$ on $\partial\Omega$

we obtain $\mathcal{N}_p V_2 \geq 0$ and $V_2(0, x) \geq u_0$. It is clear that $V_2(t, x) \geq 0 = u(t, x)$ on $\partial \Omega$. Therefore, $V_2(t, x)$ is a super-solution of (3.5). Then, Theorem 2.1 concludes that

$$0 \le u(t, x) \le \frac{K_2(R'+1)^{\frac{p}{p-1}}}{\sigma_2} < \infty.$$

In the case when r = q, we can take

$$\sigma_3 \ge \max\left\{1, 2^{1/r}(R'+1)\right\},\$$

and

$$K_{3} \ge \max\left\{\varepsilon^{-\sigma_{3}} \|u_{0}\|_{L^{\infty}}, \left(\frac{2((p-1)(\sigma_{3}-1)+N_{1}-1)}{\varepsilon^{(r-p+1)(\sigma_{3}-1)+1}}\right)^{\frac{1}{r-p+1}}\right\},$$

such that the function $V_3(t,x) = K_3 R^{\sigma_3}$ is a super-solution of (3.5). By the same procedure, one can obtain the uniform boundedness of u(t,x).

Theorem 3.2. Let $\alpha > 0$, $\beta < 0$, p > 1 and r > 0. If $q > \max\{p - 1, r, 1\}$, then the solution of the problem (3.5) blows up in finite time for some large $u_0 > 0$.

Proof of Theorem 3.2. For convenience, let us assume that $\alpha = -\beta = 1$. Set

$$v(t, |x'|) := \frac{1}{(1 - \delta t)^{k_1}} F\left(\frac{|x'|}{(1 - \delta t)^{k_2}}\right), \quad t_0 \le t < \frac{1}{\delta}, \tag{3.10}$$

where

$$F(y) := 1 + \frac{A}{\sigma} - \frac{y^{\sigma}}{\sigma A^{\sigma-1}}, \quad y \ge 0, \quad \sigma = \frac{p}{p-1},$$

and

$$k_1 = \frac{1}{q-1}, \quad 0 < k_2 < \min\left\{\frac{q-p+1}{p(q-1)}, \frac{q-r}{r(q-1)}\right\}, \quad A > \frac{k_1}{k_2}, \quad \delta < \frac{1}{k_1\left(1+\frac{A}{\sigma}\right)}.$$
(3.11)

Then, it can be noted that v(t, |x'|) is positive and smooth when $t \in [t_0, \frac{1}{\delta})$ and $|x'| < R_1(1 - \delta t)^{k_2}$, where $R_1 := (A^{\sigma-1}(A + \sigma))^{1/\sigma}$.

We want to show that v(t, |x'|) is a sub-solution of (3.5). For $y = \frac{|x'|}{(1-\delta t)^{k_2}}$, by a direct calculation we have

$$\mathcal{N}_{p}v = v_{t} - \mathcal{L}_{p}v - v^{q} + |\nabla_{H}v|^{r}$$

$$= \frac{\delta \left(k_{1}F + k_{2}yF'\right)}{(1 - \delta t)^{k_{1}+1}} - \frac{\left(|F'|^{p-2}F'\right)' + \frac{N_{1}-1}{y}|F'|^{p-2}F'}{(1 - \delta t)^{(p-2)(k_{1}+k_{2})+(k_{1}+2k_{2})}}$$

$$- \frac{F^{q}}{(1 - \delta t)^{qk_{1}}} + \frac{|F'|^{r}}{(1 - \delta t)^{r(k_{1}+k_{2})}}.$$

Note that $k_1q = k_1 + 1 > (p-2)(k_1 + k_2) + k_1 + 2k_2$ and $k_1 + 1 > (k_1 + k_2)r$ by (3.11). Observe that

$$\left(\left|F'\right|^{p-2}F'\right)' + \frac{N_1 - 1}{y}\left|F'\right|^{p-2}F' = -\frac{N_1}{A}, \quad 0 < y < R_1.$$

Let us now show that $\mathcal{N}_p v \leq 0$ for all $t \in [t_0, \frac{1}{\delta})$ and $0 \leq y \leq R_1$. In the case $0 \leq y \leq A$, from the representation of F(y) we note that

$$1 \le F(y) \le 1 + \frac{A}{\sigma}$$
 and $-1 \le F'(y) \le 0.$

Then, we can take $t_0 = t_0(p, q, r, \delta, N_1, A)$ close to $\frac{1}{\delta}$ such that

$$\mathcal{N}_{p}v \leq \frac{1}{(1-\delta t)^{k_{1}+1}} \left(\delta k_{1} \left(1 + \frac{A}{\sigma} \right) - 1 + \frac{N_{1}}{A} \left(1 - \delta t_{0} \right)^{1-2k_{2}-(p-2)(k_{1}+k_{2})} + \left(1 - \delta t_{0} \right)^{k_{1}+1-r(k_{1}+k_{2})} \right) \leq 0. \quad (3.12)$$

In the case $A \leq y \leq R_1$, we have

$$0 \le F(y) \le 1$$
 and $-\left(\frac{R_1}{A}\right)^{\sigma-1} \le F'(y) \le -1$

Similarly as above, one verifies that

$$\mathcal{N}_{p}v \leq \frac{1}{(1-\delta t)^{k_{1}+1}} \left(\delta(k_{1}-k_{2}A) + \frac{N_{1}}{A} (1-\delta t_{0})^{1-2k_{2}-(p-2)(k_{1}+k_{2})} + \left(\frac{R_{1}}{A}\right)^{r(\sigma-1)} (1-\delta t_{0})^{k_{1}+1-r(k_{1}+k_{2})} \right) \leq 0. \quad (3.13)$$

From (3.12) and (3.13), we conclude that $\mathcal{N}_p v \leq 0$ for all $t \in [t_0, \frac{1}{\delta})$ and $|x'| < R_1(1-\delta t)^{k_2}$.

Next, we estimate u_0 . By the group translation, without loss of generality we may assume that Ω contains the unit element of the group \mathbb{G} . Then, we can take suitable t_0 such that $R_1(1 - \delta t_0)^{k_2} < \max_{x=(x',x'')\in\Omega} |x'|$ and $u_0 \ge v(t_0, \cdot)$ in $\Omega \cap \{x = (x', x'') :$ $|x'| < R_1(1 - \delta t_0)^{k_2}\}$ for some large $u_0 > 0$. Then, taking into account $v \le 0$ when $|x'| \ge R_1(1 - \delta t)^{k_2}$, we obtain that $u_0 \ge v(t_0, \cdot)$ in $\overline{\Omega}$. Obviously, we also have $v \le 0$ when $(t, x) \in (t_0, \frac{1}{\delta}) \times \partial \Omega$. Thus, the comparison principle (Theorem 2.1) implies that

$$u(t,x) \ge v(t+t_0,x), \quad t \in \left[t_0,\frac{1}{\delta}\right), \quad |x'| < R_1(1-\delta t)^{k_2}.$$

On the other hand, by the definition of v we have $\lim_{t \to 1/\delta} v(t,0) \to \infty$. Consequently, u must blow up at a finite time $T \leq \frac{1}{\delta} - t_0 < \infty$.

Theorem 3.3. Assume that $\alpha < 0$, $\beta > 0$, p, r > 1 and q > 0 in (3.5) satisfy one of the following conditions:

- $r > \max\{p, q\};$
- r = q > p, and $\beta \gg |\alpha|$.

There exists M > 0 such that if $\int_{\Omega} u_0^{\frac{2r-p}{r-p}} dx > M$, then $T_{max} < \infty$.

Proof of Theorem 3.3. Assume for a contradiction that $T_{\text{max}} = \infty$. By C_1 and C_2 we denote positive constants which may vary from line to line. Set $\kappa = r/(r-p)$ and

$$y(t) = \frac{1}{\kappa+1} \int_{\Omega} u^{\kappa+1} dx. \text{ Then, using } \kappa - 1 = \frac{p}{r-p} = \frac{p\kappa}{r}, \text{ we have}$$
$$y'(t) = \beta \int_{\Omega} u^{\kappa} |\nabla_{H}u|^{r} dx - \kappa \int_{\Omega} u^{\kappa-1} |\nabla_{H}u|^{p} dx - |\alpha| \int_{\Omega} u^{q+\kappa} dx$$
$$= \beta \int_{\Omega} u^{\kappa} |\nabla_{H}u|^{r} dx - \kappa \int_{\Omega} (u^{\kappa} |\nabla_{H}u|^{r})^{p/r} dx - |\alpha| \int_{\Omega} u^{q+\kappa} dx.$$

For r > q, using Hölder's and Young's inequalities we get

$$\int_{\Omega} \left(u^{\kappa} |\nabla_{H} u|^{r} \right)^{p/r} dx \leq \left(\int_{\Omega} u^{\kappa} |\nabla_{H} u|^{r} dx \right)^{p/r} |\Omega|^{(r-p)/r}$$
$$\leq \epsilon \frac{p}{r} \int_{\Omega} u^{\kappa} |\nabla_{H} u|^{r} dx + C(\epsilon) \frac{r-p}{r} |\Omega|^{r}$$

and

$$\int_{\Omega} u^{q+\kappa} dx = \int_{\Omega} \left(u^{r+\kappa} \right)^{\frac{q+\kappa}{r+\kappa}} dx \le \left(\int_{\Omega} u^{r+\kappa} dx \right)^{\frac{q+\kappa}{r+\kappa}} |\Omega|^{\frac{r-q}{r+\kappa}} \\ \le \varepsilon \frac{q+\kappa}{r+\kappa} \int_{\Omega} u^{r+\kappa} dx + C(\varepsilon) \frac{r-q}{r+\kappa} |\Omega|.$$

Then, by Poincaré's (see, e.g. [RS17, Formula 1.10]) and reverse Hölder's inequalities we obtain

$$y'(t) \geq \frac{\beta r - \epsilon p}{r} \int_{\Omega} u^{\kappa} |\nabla_{H}u|^{r} dx - |\alpha| \varepsilon \frac{q + \kappa}{r + \kappa} \int_{\Omega} u^{r + \kappa} dx - C$$

$$= \frac{\beta r - \epsilon p}{r} \left(\frac{r}{r + \kappa}\right)^{r} \int_{\Omega} |\nabla_{H}u^{\frac{r + \kappa}{r}}|^{r} dx - |\alpha| \varepsilon \frac{q + \kappa}{r + \kappa} \int_{\Omega} u^{r + \kappa} dx - C$$

$$\geq \left(\frac{\beta r - \epsilon p}{r} \left(\frac{r}{r + \kappa}\right)^{r} C' - |\alpha| \varepsilon \frac{q + \kappa}{r + \kappa}\right) \int_{\Omega} u^{r + \kappa} dx - C$$

$$= C_{1} \int_{\Omega} u^{r + \kappa} dx - C$$

$$\geq C_{1} \left(\int_{\Omega} u^{\kappa + 1} dx\right)^{\frac{r + \kappa}{\kappa + 1}} |\Omega|^{\frac{1 - r}{\kappa + 1}} - C_{2}$$

$$\geq C_{1} \left(\int_{\Omega} u^{\kappa + 1} dx\right)^{\frac{r + \kappa}{\kappa + 1}} - C_{2}.$$

Thus, we have obtained

$$y'(t) \ge C_1 y^{\frac{r+\kappa}{\kappa+1}}(t) - C_2,$$

where $C_1 = C_1(p, r, q, \alpha, \beta, \epsilon, \varepsilon, \Omega, N_1)$ and $C_2 = C_2(p, r, q, \alpha, \beta, \epsilon, \varepsilon, \Omega, N_1) > 0$ with suitable ϵ and ε , and N_1 is the dimension of the first stratum of the group \mathbb{G} . Set

$$M > \left(\frac{2C_2}{C_1}\right)^{\frac{\kappa+1}{r+\kappa}},$$

then if y(0) > M, we have

$$y'(t) \ge \frac{C_1 y^{\frac{r+\kappa}{\kappa+1}}(t)}{2}.$$
 (3.14)

A contradiction then follows by integrating (3.14), hence $T_{\text{max}} < \infty$.

In the case r = q, the proof above is still valid for $\beta \gg |\alpha|$.

As another application of the comparison principle, we now investigate the following initial boundary value problem for the *p*-sub-Laplacian, 1 ,

$$\begin{cases} u_t - \mathcal{L}_p u = \alpha \sum_{i=1}^{n_1} |u|^{q_i - 1} u + \gamma \sum_{i=1}^{n_1} |u|^{s_i - 1} u, & x \in \Omega, \quad t > 0, \\ u(t, x) = 0, & x \in \partial\Omega, \quad t > 0, \\ u(0, x) = u_0(x), & x \in \Omega, \end{cases}$$
(3.15)

where $u_0(x) \ge 0$, $u_0(x) \ne 0$, $u_0 \in S_0^{1,\infty}(\Omega)$, and the parameters α , γ , q_i and s_i will be determined later.

Theorem 3.4. Let $\Omega \subset \mathbb{G}$ be a bounded open set in a stratified Lie group with N_1 being the dimension of the first stratum. Let $\tilde{s} = \min\{s_i\}$ and $\tilde{q} = \min\{q_i\}$. Assume that α , γ , q_i and s_i in (3.15) satisfy one of the following conditions:

(i) $\alpha > 0$, $\gamma < 0$, and $q_i \ge 1$ with $1 and <math>s_i < q_i$;

(ii) $\alpha < 0, \gamma > 0$, and $s_i \ge 1$ with $1 and <math>s_i > q_i$.

Then a weak solution of (3.15) is globally in t-bounded, that is, there exists a constant M depending only on p, q_i , s_i , α , γ , N_1 , Ω and u_0 such that for every T > 0 we have $0 \le u \le M$ on (0, T).

Remark 3.5. We refer to [RS18, Section 3] for a similar investigation when $\alpha = -\gamma = 1, n_1 = 1$.

Proof of Theorem 3.4. We only prove Part (i), since Part (ii) is actually the same, but only α and q_i are swapped by β and s_i , respectively. For convenience, we assume that $\alpha = -\gamma = 1$. We recall that $R' = \max_{x=(x',x'')\in\Omega} |x'| < \infty$ and $\varepsilon \leq |x'_0 - x'| < R' + 1$ for any $x = (x', x'') \in \Omega$, where $x_0 = (x'_0, x''_0) \in \mathbb{G} \setminus \Omega$ and $\varepsilon \in (0, 1)$. We also employ the following notations

$$V_4(t,x) := \frac{K_4}{\sigma_4} R^{\sigma_4}, \quad \sigma_4 = \frac{p}{p-1}, \quad R = |x' - x'_0|, \quad x = (x', x'') \in \Omega,$$

and

$$\mathcal{K}_p w := w_t - \mathcal{L}_p w - \sum_{i=1}^{n_1} w^{q_i} + \sum_{i=1}^{n_1} w^{s_i}.$$

Now, we look for a suitable positive K_4 such that $V_4(t, x)$ is a super-solution of (3.15). Then we have

$$\mathcal{K}_p V_4 = -N_1 K_4^{p-1} - \sum_{i=1}^{n_1} \left(\frac{K_4}{\sigma_4}\right)^{q_i} R^{\frac{q_i p}{p-1}} + \sum_{i=1}^{n_1} \left(\frac{K_4}{\sigma_4}\right)^{s_i} R^{\frac{s_i p}{p-1}}.$$

From this, we note that

$$\mathcal{K}_{p}V_{4} \ge 0 \iff \sum_{i=1}^{n_{1}} \left(\frac{K_{4}}{\sigma_{4}}\right)^{s_{i}} R^{\frac{s_{i}p}{p-1}} \ge N_{1}K_{4}^{p-1} + \sum_{i=1}^{n_{1}} \left(\frac{K_{4}}{\sigma_{4}}\right)^{q_{i}} R^{\frac{q_{i}p}{p-1}}$$

So, it is sufficient to choose K_4 such that

$$\sum_{i=1}^{n_1} \left(\frac{K_4}{\sigma_4}\right)^{s_i} R^{\frac{s_i p}{p-1}} \ge 2N_1 K_4^{p-1}, \tag{3.16}$$

$$\left(\frac{K_4}{\sigma_4}\right)^{s_i} R^{\frac{s_i p}{p-1}} \ge 2 \left(\frac{K_4}{\sigma_4}\right)^{q_i} R^{\frac{q_i p}{p-1}}.$$
(3.17)

The inequality (3.16) is satisfied if we take

$$K_4 \ge (2N_1)^{\frac{1}{\overline{s}-p+1}} \left(\sum_{i=1}^{n_1} \frac{\varepsilon^{\frac{s_i p}{p-1}}}{\sigma_4^{s_i}}\right)^{-\frac{1}{\overline{s}-p+1}},$$

provided that $\tilde{s} = \min\{s_i\} > p-1$. Dividing the inequality (3.17) by $K_4^{q_i} \frac{R^{\frac{s_i p}{p-1}}}{\sigma_4^{s_i}}$ we get

$$K_4^{s_i - q_i} \ge 2\sigma_4^{s_i - q_i} R^{\frac{p(q_i - s_i)}{p-1}}$$

that is,

$$K_4 \ge 2\sigma_4 \varepsilon^{-\frac{p}{p-1}}$$

We also need that $K_4 \geq \frac{\sigma_4 ||u_0||_{L^{\infty}}}{\varepsilon^{\sigma_4}}$ to ensure $V_4(0, x) \geq u_0$. Thus, choosing K_4 as follows

$$K_4 \ge \max\left\{\frac{\sigma_4 \|u_0\|_{L^{\infty}}}{\varepsilon^{\sigma_4}}, (2N_1)^{\frac{1}{\overline{s}-p+1}} \left(\sum_{i=1}^{n_1} \frac{\varepsilon^{\frac{s_i p}{p-1}}}{\sigma_4^{s_i}}\right)^{-\frac{1}{\overline{s}-p+1}}, 2\sigma_4 \varepsilon^{-\frac{p}{p-1}}\right\},$$

we obtain $\mathcal{K}_p V_4 \geq 0$ and $V_4(0, x) \geq u_0$. Clearly, we also have $V_4(t, x) \geq 0 = u(t, x)$ on $\partial \Omega$. Therefore, we can conclude that $V_4(t, x)$ is a super-solution of (3.15). Then, the comparison principle yields that

$$0 \le u(t,x) \le \frac{K_4(R'+1)^{\frac{p}{p-1}}}{\sigma_4} < \infty, \quad R' = \max_{x=(x',x'')\in\Omega} |x'|.$$
(3.18)

Since the right-hand side of (3.18) is independent of t, we can conclude that u(t, x) is globally in t-bounded.

By the same procedure as in the proof of Theorem 3.2, one can obtain the following result for the problem (3.15) when $n_1 = 1$:

Theorem 3.6. Let $\alpha > 0$, $\gamma < 0$, p > 1 and s > 0. If $q > \max\{s, p - 1, 1\}$, then the solution of the problem (3.15) blows up in finite time for some large $u_0 > 0$.

Proof of Theorem 3.6. As in the proof of Theorem 3.2, one can show that the same function v from (3.10) is a sub-solution of the problem (3.15). Then, the comparison principle (Theorem 2.1) concludes the proof.

References

- [Att12] A. Attouchi. Well-posedness and gradient blow-up estimate near the boundary for a Hamilton-Jacobi equation with degenerate diffusion. J. Differential Equations, 253:2474–2492, 2012.
- [BLU07] A. Bonfiglioli, E. Lanconelli and F. Uguzzoni. *Stratified Lie groups and potential theory for their sub-Laplacians*, Springer, Berlin/Heidelberg, 2007.
- [ZL13] Z. Zhang and Y. Li. Blowup and existence of global solutions to nonlinear parabolic equations with degenerate diffusion. *EJDE*, 2013(264):1–17, 2013.
- [FR16] V. Fischer and M. Ruzhansky. Quantization on nilpotent Lie groups, volume 314 of Progress in Mathematics. Birkhäuser, 2016. (open access book)
- [FR17] V. Fischer and M. Ruzhansky. Sobolev spaces on graded groups. Ann. Inst. Fourier (Grenoble), 67:1671–1723, 2017.

M. RUZHANSKY AND N. YESSIRKEGENOV

- [Fol75] G. B. Folland. Subelliptic estimates and function spaces on nilpotent Lie groups. Ark. Mat., 13:161–207, 1975.
- [FS82] G. B. Folland and E. M. Stein. Hardy spaces on homogeneous groups, volume 28 of Mathematical Notes. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1982.
- [HN79] B. Helffer and J. Nourrigat. Caracterisation des opérateurs hypoelliptiques homogènes invariants à gauche sur un groupe de Lie nilpotent gradué. Comm. Partial Differential Equations, 4(8):899–958, 1979.
- [Mil80] K. G. Miller. Parametrices for hypoelliptic operators on step two nilpotent Lie groups. Comm. Partial Differential Equations, 5(11):1153–1184, 1980.
- [RS17] M. Ruzhansky and D. Suragan. On horizontal Hardy, Rellich, Caffarelli-Kohn-Nirenberg and p-sub-Laplacian inequalities on stratified groups. J. Differential Equations, 262:1799– 1821, 2017.
- [RS18] M. Ruzhansky and D. Suragan. A comparison principle for nonlinear heat Rockland operators on graded groups. *Bull. London Math. Soc.*, 50:753–758, 2018.
- [RS19] M. Ruzhansky and D. Suragan. Hardy inequalities on homogeneous groups, volume 537 of Progress in Mathematics. Birkhäuser, 2019. (open access book)
- [tER97] A. F. M. ter Elst and D. W. Robinson. Spectral estimates for positive Rockland operators. In Algebraic groups and Lie groups, volume 9 of Austral. Math. Soc. Lect. Ser., pages 195–213. Cambridge Univ. Press, Cambridge, 1997.

MICHAEL RUZHANSKY:

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS GHENT UNIVERSITY, BELGIUM

AND

School of Mathematical Sciences

QUEEN MARY UNIVERSITY OF LONDON, UNITED KINGDOM

 $E\text{-}mail\ address\ michael.ruzhansky@ugent.be$

NURGISSA YESSIRKEGENOV:

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS GHENT UNIVERSITY, BELGIUM

AND DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, UNITED KINGDOM AND INSTITUTE OF MATHEMATICS AND MATHEMATICAL MODELLING, KAZAKHSTAN *E-mail address* nurgissa.yessirkegenov@gmail.com