

# LOCAL RIGIDITY OF TEICHMÜLLER SPACE WITH THE THURSTON METRIC

HUIPING PAN

**ABSTRACT.** We show that every  $\mathbb{R}$ -linear surjective isometry between the cotangent spaces to the Teichmüller space equipped with the Thurston norm is induced by some isometry between the underlying hyperbolic surfaces. This is an analogue of Royden's theorem concerning the Teichmüller norm.

**Keywords:** Teichmüller space, Thurston metric, Royden theorem, curve complex, rigidity.

AMS MSC2020: 30F60, 32G15

## 1. INTRODUCTION

**1.1. Background.** Let  $S_{g,n}$  be an oriented surface of genus  $g$  with  $n$  punctures whose Euler characteristic is negative:  $\chi(S_{g,n}) = 2 - 2g - n < 0$ . Let  $\mathcal{T}(S_{g,n})$  and  $\mathcal{MCG}^\pm(S_{g,n})$  be respectively the Teichmüller space and the extended mapping class group of  $S_{g,n}$ .

In his famous paper, Royden [Roy71] proved the *infinitesimal rigidity* of the Teichmüller metric which states that for closed surfaces of genus at least two, every complex linear surjective isometry between the space of integrable holomorphic quadratic differentials on Riemann surfaces are induced by some biholomorphic map between the underlying Riemann surfaces. Based on this, he showed that every (global) isometry of the Teichmüller space with the *Teichmüller metric* is an element of the extended mapping class group, and that every local isometry is the restriction of a global isometry. Since then the isometry rigidity problem of the Teichmüller metric has been extensively studied (see [Roy71], [EK74a], [EK74b], [EG96], [Lak97], [AP98], [Iva01], [Mar03], [EM03], [FW10], [MM13]). This result was extended by Earle-Kra in [EK74a] to the case of punctured Riemann surfaces, and in [EK74b] to the case of isometries between different Teichmüller spaces. Later, it was further extended to bordered Riemann surfaces by Earle-Gardiner in [EG96] and to Riemann surfaces with infinitely many punctures by Lakic in [Lak97], where in both cases the Teichmüller spaces are of infinite dimensional. The problem was completely settled for general surfaces which are not of exceptional finite type by Markovic in [Mar03]. The proofs in these papers are based on a detailed analysis of the space of integral holomorphic quadratic differentials which serves as the cotangent space to the Teichmüller space.

In [Iva01], Ivanov provided a different proof of the *global isometry rigidity* from the point of view of Mostow rigidity. A key role in the proof is played by the *curve complex*, which is an analogue of the Tits building. By analysing the asymptotic behaviour of pairs of Teichmüller geodesic rays, Ivanov showed that every isometry of the Teichmüller metric induces an automorphism of the curve complex. Then he applied the *automorphism rigidity of curve complexes* to show that every isometry is an element of the extended mapping class group.

Ivanov's idea has been extended to other metrics on the Teichmüller spaces. In [MW02], Masur-Wolf proved the *global isometry rigidity* of the Teichmüller space  $\mathcal{T}(S_{g,n})$  with the *Weil-Petersson metric*, provided that  $(g, n) \notin \{(0, 3), (0, 4), (1, 1), (1, 2)\}$ . Later, it was extended by Brock-Margalit ([BM07]) to all hyperbolic surfaces of finite area via the *automorphism rigidity of pants complex* ([Mar04]).

In [Wal14], Walsh proved the *global isometry rigidity* of the Teichmüller space  $\mathcal{T}(S_{g,n})$  with the *Thurston metric*, provided that  $(g, n) \notin \{(0, 4), (1, 1), (1, 2)\}$ . Walsh's proof is based on the *automorphism rigidity of the curve complex* as well as the *horofunction compactification* of  $\mathcal{T}(S_{g,n})$ . Very recently, Dumas-Lenzhen-Rafi-Tao ([DLRT20]) proved the *infinitesimal rigidity*, the *local isometry rigidity*, and the *global isometry rigidity* of  $\mathcal{T}(S_{1,1})$  with the Thurston metric. The idea of their proof is to recognize lengths and intersection numbers of curves on  $X$  from the features of the unit sphere in the tangent space to  $\mathcal{T}(S_{1,1})$ , and then apply the Fenchel-Nilsen coordinates to identify the underlying hyperbolic surfaces.

In [FW10], Farb and Weinberger proved a general rigidity for all complete, finite covolume and  $\mathcal{MCG}(S_{g,n})$  invariant Finsler metrics on  $\mathcal{T}(S_{g,n})$  from a totally different perspective, which states that the group of isometries contains  $\mathcal{MCG}(S_{g,n})$  as a subgroup of finite index.

The goal of this paper is to consider the *infinitesimal rigidity* of the *Thurston metric*.

**1.2. The Thurston metric.** The Thurston metric on  $\mathcal{T}(S_{g,n})$  is defined as following:<sup>1</sup>

$$(1) \quad d_{\text{Th}}(X, Y) = \log \sup_{\alpha \in \mathcal{C}^0(S_{g,n})} \frac{\ell_\alpha(Y)}{\ell_\alpha(X)},$$

where  $\mathcal{C}^0(S_{g,n})$  is the set of (isotopy classes of) simple closed curves on  $S_{g,n}$  and  $\ell_\alpha(X)$  is the hyperbolic length of the geodesic representative of  $\alpha$  on  $X$ . Thurston showed in [Thu98] that  $d_{\text{Th}}$  is a Finsler metric, i.e. it is induced by a (nonsymmetric) norm on the tangent bundle:

$$(2) \quad \|v\|_{\text{Th}} := \sup_{\lambda \in \mathcal{PML}(S)} (d_X \log \ell_\lambda)[v], \quad \forall v \in T_X \mathcal{T}(S_{g,n}),$$

where  $\mathcal{PML}(S_{g,n}) = \mathcal{ML}(S_{g,n})/\mathbb{R}_{>0}$  is the space of projective measured laminations and  $\mathcal{ML}(S_{g,n})$  is the space of measured laminations on  $S_{g,n}$ . We call the above asymmetric norm the *Thurston infinitesimal norm*.

**Theorem 1.1** ([DLRT20], Theorem 6.1). *Let  $S_{g,n}$  be an orientable surface of finite type with negative Euler characteristic. Then the Thurston norm function  $T\mathcal{T}(S_{g,n}) \rightarrow \mathbb{R}$  is locally Lipschitz.*

The unit sphere in the cotangent bundle has a very nice description as following.

**Theorem 1.2** ([Thu98], Theorem 5.1). *For any hyperbolic surface  $X$  of finite type, the map*

$$\begin{aligned} \mathcal{PD}_X : \mathcal{PML}(S_{g,n}) &\longrightarrow T_X^* \mathcal{T}(S_{g,n}) \\ [\mu] &\longmapsto d_X \log \ell_\mu \end{aligned}$$

*embeds  $\mathcal{PML}(S_{g,n})$  as the boundary of a convex neighbourhood of the origin. This convex neighbourhood is dual to the unit ball  $\{v \in T_X \mathcal{T}(S_{g,n}) : \|v\|_{\text{Th}} \leq 1\}$ .*

---

<sup>1</sup>Technically, Thurston defined  $d_{\text{Th}}(X, Y)$  as the logarithm of the minimum of Lipschitz constants among Lipschitz homeomorphisms from  $X$  to  $Y$  which are homotopic to the identity, and proved (1) in [Thu98, Theorem 8.5]. Here we use the latter as the definition of the Thurston metric for the sake of convenience.

The following rigidity questions were raised by Papadopoulos, Ther  t, Dumas, and Rafi in [PT07a, Problem IV] and [Su16, Problem 2.1, Problem 2.6].

- Question 1.** (1) Does the Thurston infinitesimal norm determine the underlying hyperbolic surfaces?  
 (2) Is each local isometry of Teichm  ller space with the Thurston metric induced by an element of the extended mapping class group?

**Remark 1.3.** For  $\mathcal{T}(S_{1,1})$ , Question 1 was answered affirmatively by Dumas-Lenzhen-Rafi-Tao in [DLRT20].

**1.3. Main results.** We answer Question 1 affirmatively.

**Theorem 1.4.** *Let  $S_{g,n}$  be an orientable surface of finite type with negative Euler characteristic. Then every surjective  $\mathbb{R}$ -linear isometry*

$$\Phi^* : (T_X^* \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}}) \rightarrow (T_Y^* \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}})$$

*is induced by some isometry  $\phi : X \rightarrow Y$  such that  $\Phi^*(d_X \ell_\mu) = d_Y \ell_{\phi\mu}$  for every  $\mu \in \mathcal{ML}(S_{g,n})$ . In particular,  $Y$  belongs to the  $\mathcal{MCG}^\pm(S_{g,n})$  orbit of  $X$ .*

We can also use the Thurston infinitesimal norm to distinguish the topology of the underlying hyperbolic surfaces.

**Theorem 1.5.** *Let  $S_{g,n}$  and  $S_{g',n'}$  be two orientable surfaces of finite type with negative Euler characteristic. If there exists a surjective  $\mathbb{R}$ -linear isometry*

$$\Phi : (T_X \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}}) \rightarrow (T_Y \mathcal{T}(S_{g',n'}), \|\bullet\|_{\text{Th}})$$

*for some  $X \in \mathcal{T}(S_{g,n})$  and  $Y \in \mathcal{T}(S_{g',n'})$ , then  $(g, n) = (g', n')$ .*

Based on the infinitesimal rigidities above, we obtain both the local rigidity and the global rigidity of the Thurston metric, as well as an analogue of the Patterson's theorem.

**Theorem 1.6** (Local rigidity). *Let  $S_{g,n}$  be an orientable surface of finite type with negative Euler characteristic. Let  $U$  be a connected open set in  $\mathcal{T}(S_{g,n})$ . Then any isometric embedding  $(U, d_{\text{Th}}) \rightarrow (\mathcal{T}(S_{g,n}), d_{\text{Th}})$  is the restriction to  $U$  of some element of  $\mathcal{MCG}^\pm(S_{g,n})$ .*

**Theorem 1.7** (Global rigidity). *Let  $S_{g,n}$  be an orientable surface of finite type with negative Euler characteristic. Then every isometry of  $\mathcal{T}(S_{g,n})$  is an element of  $\mathcal{MCG}^\pm(S_{g,n})$ . In particular,*

$$\text{Isom}(\mathcal{T}(S_{g,n}), d_{\text{Th}}) = \begin{cases} \mathcal{MCG}^\pm(S_{g,n})/\mathbb{Z}_2, & \text{if } (g, n) \in \{(2, 0), (1, 1), (1, 2)\}, \\ \mathcal{MCG}^\pm(S_{g,n})/\mathbb{Z}_2 \oplus \mathbb{Z}_2, & \text{if } (g, n) = (0, 4), \\ \mathcal{MCG}^\pm(S_{g,n}), & \text{otherwise.} \end{cases}$$

**Theorem 1.8** (Topological rigidity). *Let  $S_{g,n}$  and  $S_{g',n'}$  be two orientable surfaces of finite type with negative Euler characteristics. Then  $(\mathcal{T}(S_{g,n}), d_{\text{Th}})$  and  $(\mathcal{T}(S_{g',n'}), d_{\text{Th}})$  are isometric if and only if  $(g, n) = (g', n')$ .*

**Remark 1.9.** • Theorem 1.4 and Theorem 1.6 were proved By Dumas-Lenzhen-Rafi-Tao [DLRT20] if  $(g, n) = (1, 1)$ .

- Theorem 1.7 was proved Dumas-Lenzhen-Rafi-Tao [DLRT20] if  $(g, n) = (1, 1)$ , and by Walsh [Wal14] if  $(g, n) \notin \{(0, 4), (1, 1), (1, 2)\}$ .
- Theorem 1.8 was first proved by Walsh [Wal14] if  $\{(g, n), (g', n')\}$  is not one of the following pairs:  $\{(0, 4), (1, 1)\}, \{(0, 5), (1, 2)\}, \{(0, 6), (2, 0)\}$ .
- For surfaces of low complexity, Theorem 1.8 is different from the case of Teichm  ller metric. For the Teichm  ller metric, Patterson [Pat72] showed that

$\mathcal{T}(S_{g,n})$  and  $\mathcal{T}(S_{g',n'})$  are isometric if and only if  $(g, n) = (g', n')$  or  $\{(g, n), (g'n')\}$  is one of the following pairs:

$$\{(0, 4), (1, 1)\}, \{(0, 5), (1, 2)\}, \{(0, 6), (2, 0)\}.$$

- Recently, Theorem 1.4-1.8 are proved independently by Huang-Ohshika-Papadopoulos [HOP21] using a differently method.

**Remark 1.10.** For more research about the Thurston metric, we refer to [Thu98], [Liu00], [PT07a], [CR07], [The07], [PT07b], [LRT12], [PT12], [LPST13], [Wal14], [PS14], [The14], [LRT15], [PS15], [PS16], [PY17], [AD19], [HP19], [DLRT20], and [HOP21].

**1.4. Idea of proof.** Our proof is based on the linear structures on  $\mathcal{ML}(S_{g,n})$  induced from the cotangent spaces of the Teichmüller space via Theorem 1.2. We show that the induced linear structures depends dramatically on the underlying hyperbolic surfaces (see Theorem 3.1). By analysing the maximal flats (facets) in the unit sphere of the tangent space, we show that every surjective  $\mathbb{R}$ -linear isometry between tangent spaces induces an isomorphism of the curve complexes for non-exceptional surfaces (see Proposition 4.3). The linearity rigidity of linear structures on  $\mathcal{ML}(S_{g,n})$  and the automorphism rigidity of the curve complex then enable us to prove Theorem 1.4 and Theorem 1.5 in this case. For exceptional surfaces, we adapt the idea from [DLRT20]. The proofs of Theorem 1.6, Theorem 1.7 and Theorem 1.8 are standard assuming Theorem 1.1 and Theorem 1.4.

**Acknowledgements.** I am grateful to Kasra Rafi for sharing with me the question about the local rigidity of Thurston metric. I thank Weixu Su for his comments. Finally, I would like to thank the referee for his/her comments and suggestions. This work is supported by NSFC 11901241.

## 2. PRELIMINARIES

**2.1. Teichmüller space and measured laminations.** Continuing with the same notation as in the introduction, let  $S_{g,n}$  be an oriented surface of genus  $g$  with  $n$  punctures whose Euler characteristic is negative:  $\chi(S_{g,n}) = 2 - 2g - n < 0$ . A *marked hyperbolic surface* is a pair  $(X, f)$ , where  $X$  is a hyperbolic surface and  $f : S_{g,n} \rightarrow X$  is an orientation-preserving homeomorphism. Two marked hyperbolic surfaces  $(X, f)$  and  $(X', f')$  are said to be *equivalent* if there exists an isometry  $X \rightarrow X'$  which is isotopic to  $f' \circ f^{-1}$ . The *Teichmüller space*  $\mathcal{T}(S_{g,n})$  is the set of equivalence classes of marked hyperbolic surfaces. Sometimes, we simply denote an equivalence class by  $X$ . With the topology induced by the Thurston metric (defined in (1)),  $\mathcal{T}(S_{g,n})$  is homeomorphic to  $\mathbb{R}^{6g-6+2n}$ .

A *measured lamination* is a lamination on  $S_{g,n}$  equipped with a transverse invariant measure. Every measured lamination  $\mu$  induces a functional over  $\mathcal{C}^0(S_{g,n})$ , the set of isotopy classes of simple closed curves on  $S_{g,n}$ , by associating to each  $\alpha \in \mathcal{C}^0(S_{g,n})$  the *intersection number*  $i(\mu, \alpha)$ :

$$i(\mu, \alpha) := \inf_{\alpha'} \int_{\alpha'} d\mu$$

where  $\alpha'$  ranges over all simple closed curves  $\alpha'$  isotopic to  $\alpha$ . Two measured laminations are said to be *equivalent* if they induce the same functional on  $\mathcal{C}^0(S_{g,n})$ . Let  $\mathcal{ML}(S_{g,n})$  be the space of equivalence classes of measured laminations equipped with the weak-\* topology induced from the space of functionals over  $\mathcal{C}^0(S_{g,n})$ . Let  $\mathcal{PML}(S_{g,n}) := \mathcal{ML}(S_{g,n})/\mathbb{R}_{>0}$  be the space of projective classes of measured laminations. Thurston showed that  $\mathcal{ML}(S_{g,n})$  is homeomorphic to  $\mathbb{R}^{6g-6+2n}$ , and that the set of weighted simple closed curves is dense in  $\mathcal{ML}(S_{g,n})$  (see [FLP12] and [CB88] for more details).

**2.2. Mapping class group action.** Let  $\text{Homeo}^\pm(S_{g,n})$  (resp.  $\text{Homeo}^+(S_{g,n})$ ) be the group of homeomorphisms (resp. orientation-preserving homeomorphisms) of  $S_{g,n}$ . Let  $\text{Homeo}_0(S_{g,n})$  be the subgroup of homeomorphisms isotopic to the identity. Then the mapping class group  $\mathcal{MCG}^+(S_{g,n})$  of  $S_{g,n}$  is defined to be:

$$\mathcal{MCG}^+(S_{g,n}) = \text{Homeo}^+(S_{g,n}) / \text{Homeo}_0(S_{g,n})$$

and the extended mapping class group  $\mathcal{MCG}^\pm(S_{g,n})$  is defined to be:

$$\mathcal{MCG}^\pm(S_{g,n}) = \text{Homeo}^\pm(S_{g,n}) / \text{Homeo}_0(S_{g,n}).$$

The extended mapping class group acts on  $\mathcal{T}(S_{g,n})$  as follows. Each orientation-preserving mapping class  $\phi \in \mathcal{MCG}^+S_{g,n}$  acts on  $\mathcal{T}(S_{g,n})$  by pre-composition:

$$\phi \cdot (X, f) := (X, f \circ \phi^{-1}).$$

We now consider the action of orientation-reversing homeomorphisms. For every hyperbolic surface  $X$ , let  $\overline{X}$  be  $X$  with the opposite orientation. Let  $j_X : X \rightarrow \overline{X}$  be the identity map. In particular,  $j_X$  is an orientation-reversing isometry. Now the image of  $(X, f)$  under  $\psi$  is defined to be:

$$\psi \cdot (X, f) := (\overline{X}, j_X \circ f \circ \psi^{-1}).$$

By definition, we see that  $\mathcal{MCG}^\pm(S_{g,n})$  acts on  $(\mathcal{T}(S_{g,n}), d_{\text{Th}})$  by isometries.

**2.3. Stretch lines.** A geodesic lamination on  $X \in \mathcal{T}(S_{g,n})$  is a lamination whose leaves are hyperbolic geodesics. A geodesic lamination is said to be *maximal* if each component of  $X - \lambda$  is an ideal hyperbolic triangle.

**Theorem 2.1** ([Thu98], Corollary 4.2). *For any complete hyperbolic structure  $X$  of finite area on a surface  $S$ , for any maximal geodesic lamination  $\lambda$  not all of whose leaves go to a cusp at both ends there is a new hyperbolic structure*

$$\text{stretch}(X, \lambda, t)$$

*depending analytically on  $t > 0$  such that*

(a) *the identity map is Lipschitz with Lipschitz constant*

$$L(X, \text{stretch}(X, \lambda, t)) = \exp(t)$$

(b) *the identity map exactly expands arc length of  $\lambda$  by the constant factor  $\exp(t)$ , while for each point on  $X - \lambda$  the Lipschitz constant is strictly less than  $\exp(t)$ .*

A direct consequence is the following infinitesimal version.

**Corollary 2.2.** *Let  $v \in T_X \mathcal{T}(S_{g,n})$  be the vector tangent to the stretch line  $\text{stretch}(X, \lambda, t)$ . Then*

$$(d_X \log \ell_\mu)(v) \leq 1, \forall \mu \in \mathcal{ML}(S_{g,n})$$

and

$$(d_X \log \ell_\mu)(v) = 1 \iff \text{supp} \mu \subset \lambda$$

where  $\text{supp} \mu$  represents the supporting lamination of  $\mu$ .

Using stretch lines, Thurston ([Thu98, Theorem 8.5]) showed that for every pair of points in  $\mathcal{T}(S_{g,n})$ , there exists a geodesic with respect to  $d_{\text{Th}}$ , which is a concatenation of stretch lines. In general, such geodesics are not unique. An interesting question is to describe the set of the union of all geodesics between two points, see [DLRT20] for the case of punctured torus.

**2.4. Curve complex.** For  $(g, n) \neq (0, 4), (1, 1)$ , the *curve complex*  $\mathcal{C}(S_{g,n})$  is a complex whose vertices are isotopy classes of simple closed curves on  $S_{g,n}$  and whose  $k$ -simplices are collections of  $k + 1$  isotopy classes of simple closed curves on  $S_{g,n}$  which can be realized disjointly. For our purpose, we need the following connectedness and rigidity of  $\mathcal{C}(S_{g,n})$ .

**Theorem 2.3** ([Har81], see also [Put08]).  $\mathcal{C}(S_{g,n})$  is connected.

**Theorem 2.4** ([Iva97], [Kor99], [Luo00]). (a) Suppose that  $\{(g, n), (g', n')\}$  is not one of the following three pairs:  $\{(0, 4), (1, 1)\}$ ,  $\{(0, 5), (1, 2)\}$ ,  $\{(0, 6), (2, 0)\}$ . then  $\mathcal{C}(S_{g,n})$  and  $\mathcal{C}(S_{g',n'})$  are isomorphic if and only if  $(g, n) = (g', n')$ .

(b) Suppose that  $S_{g,n}$  is not a sphere with four or fewer punctures, nor a torus with two or fewer punctures. Then every automorphism of  $\mathcal{C}(S_{g,n})$  is an element of  $\mathcal{MCG}^\pm(S_{g,n})$ , and

$$\text{Aut}(\mathcal{C}(S_{g,n})) := \begin{cases} \mathcal{MCG}^\pm(S_{g,n})/\mathbb{Z}_2, & \text{if } (g, n) = (2, 0) \\ \mathcal{MCG}^\pm(S_{g,n}), & \text{otherwise} \end{cases}$$

Moreover, any automorphism of  $\mathcal{C}(S_{1,2})$  preserving the set of vertices represented by separating curves is induced by an extended mapping class.

We also need the curve complexes of surfaces with low complexity. The curve complex  $\mathcal{C}(S_{0,4})$  (resp.  $\mathcal{C}(S_{1,1})$ ) is a graph whose vertices are isotopy classes of simple closed curves on  $S_{0,4}$  (resp.  $S_{1,1}$ ) and whose edges are pairs of isotopy classes of simple closed curves with geometric intersection number one (resp. two).

For simple closed curves  $\alpha$  and  $\beta$ , we denote by  $D_\alpha^n \beta$  the  $n$ -th Dehn twist of  $\beta$  along  $\alpha$ . We say that an isomorphism  $\iota : \mathcal{C}(S_{g,n}) \rightarrow \mathcal{C}(S_{g',n'})$  is *compatible with Dehn twists* if  $\iota(D_\alpha \beta) = D_{\iota(\alpha)} \iota(\beta)$  holds for any simple closed curves  $\alpha$  and  $\beta$ .

**Lemma 2.5** ([Luo00], Lemma 2.1). Suppose that  $\{(g, n), (g', n')\}$  is one of the following three pairs:  $\{(0, 4), (1, 1)\}$ ,  $\{(0, 5), (1, 2)\}$ ,  $\{(0, 6), (2, 0)\}$ . then  $\mathcal{C}(S_{g,n}) \simeq \mathcal{C}(S_{g',n'})$ . Moreover any isomorphism  $\mathcal{C}(S_{g,n}) \rightarrow \mathcal{C}(S_{g',n'})$  is compatible with the Dehn twists.

*Sketch of the proof.* Let  $r : S_{2,0} \rightarrow S_{2,0}$  be a hyperelliptic involution. Then  $r$  fixes every simple closed curve. The quotient map  $\pi : S_{2,0} \rightarrow S_{2,0}/\sim$  induces an isomorphism  $\pi^* : \mathcal{C}(S_{2,0}) \rightarrow \mathcal{C}(S_{0,6})$ . Any other isomorphism is a precomposition of  $\pi^*$  with some automorphism of  $\mathcal{C}(S_{2,0})$ . The compatibility with Dehn twists then follows from the automorphism rigidity of  $\mathcal{C}(S_{2,0})$  in Theorem 2.4. By considering subgraphs of  $\mathcal{C}(S_{2,0})$  and  $\mathcal{C}(S_{0,6})$ , one gets isomorphisms  $\mathcal{C}(S_{1,2}) \rightarrow \mathcal{C}(S_{0,5})$  and  $\mathcal{C}(S_{1,1}) \rightarrow \mathcal{C}(S_{0,4})$  as well as the compatibility.  $\square$

**Remark 2.6.** Let  $\iota$  be either an automorphism of  $(\mathcal{C}(S_{1,2}))$  or an isomorphism from  $(\mathcal{C}(S_{1,2}))$  to  $(\mathcal{C}(S_{0,5}))$  which sends some non-separating curve  $\alpha$  to a separating curve  $\iota(\alpha)$ . Considering the subgraphs induced by the curves disjoint from  $\alpha$  and  $\iota(\alpha)$  respectively, we see that  $\iota$  induces an isomorphism  $\iota^* : \mathcal{C}(S_{0,4}) \rightarrow \mathcal{C}(S_{1,1})$ .

**2.5. Exceptional surface  $S_{1,1}$ .** We recall some results from [DLRT20] in the case of punctured torus  $S_{1,1}$ . Notice that for both  $S_{0,4}$  and  $S_{1,1}$ , the Teichmüller spaces are of dimension two. Accordingly, the unit tangent spheres are of dimension one. Let  $\alpha$  be a simple closed curve. Let

$$F_X(\alpha) := \{v \in T_X^1 \mathcal{T}(S_{g,n}) : (d_X \log \ell_\alpha)(v) = 1\}$$

be the facet in the unit tangent sphere  $T_X^1 \mathcal{T}(S_{g,n})$  corresponding to  $\alpha$ . Let  $|F_X(\alpha)|$  be the length of  $F_X(\alpha)$  with respect to the Thurston norm  $\|\cdot\|_{\text{Th}}$ .



**Proposition 2.7** ([DLRT20], Proposition 6.7). *Let  $X \in \mathcal{T}(S_{1,1})$ . There exists a constant  $C_1 = C_1(X)$  depending on  $X$ , such that for every simple closed curve  $\alpha$ ,*

$$\frac{1}{C_1} \ell_\alpha(X)^2 e^{-\ell_\alpha(X)} \leq |F_X(\alpha)| \leq C_1 \ell_\alpha(X)^2 e^{-\ell_\alpha(X)}.$$

This implies

**Theorem 2.8** ([DLRT20], Theorem 6.8). *Let  $X \in \mathcal{T}(S_{1,1})$ . Let  $\alpha$  and  $\beta$  be two simple closed curves with  $i(\alpha, \beta) = 1$ . Let  $\beta_n = D_\alpha^n \beta$ . Then*

$$\lim_{n \rightarrow \infty} \frac{|\log |F_X(\beta_n)||}{|n|} = \ell_\alpha(X).$$

For each simple closed curve  $\alpha$  on  $S_{0,4}$  or  $S_{1,1}$ , there are two canonical maximal laminations  $\alpha^+$  and  $\alpha^-$ <sup>2</sup>. Let  $v_X(\alpha^+)$  and  $v_X(\alpha^-)$  be respectively the unit tangent vectors to the stretch paths  $\text{stretch}(X, \alpha^+, t)$  and  $\text{stretch}(X, \alpha^-, t)$ . Then  $v_X(\alpha^+)$  and  $v_X(\alpha^-)$  are the endpoints of  $F_X(\alpha)$ . Hence

$$|F_X(\alpha)| = \|v_X(\alpha^+) - v_X(\alpha^-)\|_{\text{Th}}.$$

Fix a simple closed curve  $\alpha$ . For  $\gamma \neq \alpha$ , let  $(v_X(\gamma^-), v_X(\alpha^+))$  (resp.  $(v_X(\gamma^+), v_X(\alpha^-))$ ) be the subinterval of  $T_X^1 \mathcal{T}(S_{1,1}) \setminus F_X(\alpha)$  whose endpoints are  $v_X(\gamma^-)$  and  $v_X(\alpha^+)$  (resp.  $v_X(\gamma^+)$  and  $v_X(\alpha^-)$ ).

Combining Theorem 2.8 and the train track approximate, Dumas-Lenzhen-Rafi-Tao also proved the following, which is contained in the proof of [DLRT20, Theorem 1.4] and which we summarize as a proposition.

**Proposition 2.9** ([DLRT20]). *Let  $X \in \mathcal{T}(S_{1,1})$ . There exists a constant  $L_1$  depending on  $X$  such that for any simple closed curve  $\alpha$  with  $\ell_\alpha(X) > L_1$ , there exist a simple closed curve  $\beta$  with  $i(\alpha, \beta) = 1$  and a positive integer  $N$ , such that*

- for each  $n > N$ ,  $F_X(\beta_n)$  is the longest facet in  $(v_X(\beta_{n-1}^-), v_X(\alpha^+))$ , and
- for each  $n < -N$ ,  $F_X(\beta_n)$  is the longest facet in  $(v_X(\beta_{n+1}^+), v_X(\alpha^-))$ ,

where  $\beta_n = D_\alpha^n \beta$ .

**2.6. Exceptional surface  $S_{0,4}$ .** Using the same idea as in [DLRT20], one can prove analogues for  $S_{0,4}$ . Namely

**Proposition 2.10.** *Let  $X \in \mathcal{T}(S_{0,4})$ . There exists a constant  $C_2 = C_2(X)$  depending on  $X$ , such that for every simple closed curve  $\alpha$ ,*

$$\frac{1}{C_2} \ell_\alpha(X)^2 e^{-\ell_\alpha(X)/2} \leq |F_X(\alpha)| \leq C_2 \ell_\alpha(X)^2 e^{-\ell_\alpha(X)/2}.$$

**Theorem 2.11.** *Let  $X \in \mathcal{T}(S_{0,4})$ . Let  $\alpha$  and  $\beta$  be two simple closed curves with  $i(\alpha, \beta) = 2$ . Let  $\beta_n = D_\alpha^n \beta$ . Then*

$$\lim_{n \rightarrow \infty} \frac{|\log |F_X(\beta_n)||}{|n|} = \ell_\alpha(X).$$

---

<sup>2</sup>For  $S_{1,1}$ ,  $\alpha^+ \setminus \alpha$  (resp.  $\alpha^- \setminus \alpha$ ) consists of three bi-infinite simple arcs  $\{\delta, \eta_1, \eta_2\}$  such that  $\delta$  spirals to the left (resp. right) at both ends around  $\alpha$  while  $\eta_i$  spirals to left (resp. right) around  $\alpha$  at one end and approaches to the puncture at the other end. For  $S_{0,4}$ ,  $\alpha^+ \setminus \alpha$  (resp.  $\alpha^- \setminus \alpha$ ) consists of six bi-infinite simple arcs  $\{\delta_1, \delta_2, \eta_1, \eta_2, \eta_3, \eta_4\}$  such that  $\delta_i$  spirals to the left (resp. right) at both ends around  $\alpha$  while  $\eta_i$  spirals to left (resp. right) around  $\alpha$  at one end and approaches to a puncture at the other end.

**Proposition 2.12.** *Let  $X \in \mathcal{T}(S_{0,4})$ . There exists a constant  $L_2$  depending on  $X$  such that for any simple closed curve  $\alpha$  with  $\ell_\alpha(X) > L_2$ , there exist a simple closed curve  $\beta$  with  $i(\alpha, \beta) = 2$  and a positive integer  $N$ , such that*

- for each  $n > N$ ,  $F_X(\beta_n)$  is the longest facet in  $(v_X(\beta_{n-1}^-), v_X(\alpha^+))$ , and
- for each  $n < -N$ ,  $F_X(\beta_n)$  is the longest facet in  $(v_X(\beta_{n+1}^+), v_X(\alpha^-))$ ,

where  $\beta_n = D_\alpha^n \beta$ .

**Remark 2.13.** Theorem 2.11 is a direct consequence of Proposition 2.10. The proof of Proposition 2.12 is exactly the same as that of Proposition 2.9 except that one replace Proposition 2.7 and Theorem 2.8 by Proposition 2.10 and Theorem 2.11. Proposition 2.10 is a slightly different from Proposition 2.9 in the sense that  $|F_X(\alpha)|$  is comparable to  $\ell_\alpha(X)^2 e^{-\ell_\alpha(X)/2}$  in Proposition 2.10 while it is comparable to  $\ell_\alpha(X)^2 e^{-\ell_\alpha(X)}$  in Proposition 2.7. The difference is due to the calculation of Fenchel-Nielsen coordinates. Namely, in this case the equation in the fourth line in [DLRT20, Page 50] is replaced by the following:

$$\Delta(t) = \tau_\alpha(X_t^+) - \tau_\alpha(X_t^-) = 4e^t \log \frac{e^{\ell_\alpha(X)} + 1}{e^{\ell_\alpha(X)/2} - 1} - 4 \log \frac{e^{e^t \ell_\alpha(X)} + 1}{e^{e^t \ell_\alpha(X)/2} - 1}.$$

The calculation of this equation is almost the same as the one demonstrated in the proof of [DLRT20, Theorem 5.1]. We omit the details.

### 3. LINEAR STRUCTURES FOR THE SPACE OF MEASURED LAMINATIONS

By identifying  $\mathcal{PML}(S_{g,n})$  with  $\{\mu \in \mathcal{ML}(S_{g,n}) : \ell_\mu(X) = 1\}$ , we obtain a homeomorphism

$$\begin{aligned} \mathcal{D}_X : \mathcal{ML}(S_{g,n}) &\longrightarrow T_X^* \mathcal{T}(S_{g,n}) \\ \mu &\longmapsto d_X \ell_\mu. \end{aligned}$$

In this way, we associate a linear structure to  $\mathcal{ML}(S_{g,n})$  by pulling back the linear structure on the cotangent space  $T_X^* \mathcal{T}(S_{g,n})$ .

Notice that for each  $\mu \in \mathcal{ML}(S_{g,n})$ , we have

$$\|d_X \log \ell_\mu\|_{\text{Th}, X} = 1.$$

Therefore,

$$\|d_X \ell_\mu\|_{\text{Th}, X} = \ell_\mu(X), \quad \forall \mu \in \mathcal{ML}(S_{g,n}), \quad \forall X \in \mathcal{T}(S_{g,n}).$$

For two marked hyperbolic surfaces  $X, Y \in \mathcal{T}(S_{g,n})$ , let us consider a homeomorphism  $\Gamma_{X,Y} : T_X^* \mathcal{T}(S_{g,n}) \longrightarrow T_Y^* \mathcal{T}(S_{g,n})$  which is the unique continuous extension of

$$(3) \quad \begin{aligned} \Gamma_{X,Y} : T_X^* \mathcal{T}(S_{g,n}) - \{0\} &\longrightarrow T_Y^* \mathcal{T}(S_{g,n}) - \{0\} \\ d_X \ell_\mu &\longmapsto \frac{\ell_\mu(X)}{\ell_\mu(Y)} d_Y \ell_\mu. \end{aligned}$$

Therefore,

$$\|\Gamma_{X,Y}(d_X \ell_\mu)\|_{\text{Th}, Y} = \|d_X \ell_\mu\|_{\text{Th}, X} = \ell_\mu(X).$$

In particular,  $\Gamma_{X,Y}(0) = 0$ .

In general,  $\Gamma_{X,Y}$  is not linear. In fact, we have the following rigidity.

**Theorem 3.1** (Linearity rigidity). *Suppose that  $S_{g,n}$  is not a sphere with four or fewer punctures, nor a torus with one or fewer punctures. Let  $X, Y \in \mathcal{T}(S_{g,n})$ . Then  $\Gamma_{X,Y}$  is linear if and only if  $X = Y$ .*



*Proof.* If  $X = Y$ , then  $\Gamma_{X,Y}$  is the identity map, which is linear. We now consider the converse. In the following, we assume that  $\Gamma_{X,Y}$  is linear.

Let  $\alpha$  and  $\beta$  be two disjoint, non-isotopic simple closed curves. Then  $\alpha + \beta$  is also a measured lamination. Therefore,

$$\begin{aligned}\Gamma_{X,Y}(d_X \ell_\alpha + d_X \ell_\beta) &= \Gamma_{X,Y}(d_X(\ell_{\alpha+\beta})) \\ &= \frac{\ell_{\alpha+\beta}(X)}{\ell_{\alpha+\beta}(Y)} d_Y \ell_{\alpha+\beta} \quad (\text{by (3)}) \\ &= \frac{\ell_\alpha(X) + \ell_\beta(X)}{\ell_\alpha(Y) + \ell_\beta(Y)} (d_Y \ell_\alpha + d_Y \ell_\beta).\end{aligned}$$

On the other hand, it follows from the linearity of  $\Gamma_{X,Y}$  that

$$\begin{aligned}\Gamma_{X,Y}(d_X \ell_\alpha + d_X \ell_\beta) &= \Gamma_{X,Y}(d_X \ell_\alpha) + \Gamma_{X,Y}(d_X \ell_\beta) \\ &= \frac{\ell_\alpha(X)}{\ell_\alpha(Y)} d_Y \ell_\alpha + \frac{\ell_\beta(X)}{\ell_\beta(Y)} d_Y \ell_\beta.\end{aligned}$$

By Theorem 1.2, we see that  $d_Y \ell_\alpha$  and  $d_Y \ell_\beta$  are linearly independent in  $T^* \mathcal{T}(S_{g,n})$ . Comparing the two equations above, we have

$$\frac{\ell_\alpha(X)}{\ell_\alpha(Y)} = \frac{\ell_\alpha(X) + \ell_\beta(X)}{\ell_\alpha(Y) + \ell_\beta(Y)} = \frac{\ell_\beta(X)}{\ell_\beta(Y)}.$$

Now, Let  $\alpha, \delta$  be an arbitrary pair of simple closed curves. By the connectedness of the curve complex (Theorem 2.3), there exists a sequence of simple closed curves  $\alpha_0 = \alpha, \alpha_1, \dots, \alpha_k = \delta$ , such that  $\alpha_i$  and  $\alpha_{i+1}$  are disjoint for each  $i = 0, 1, \dots, k-1$ . It then follows from the discussion above that

$$\frac{\ell_{\alpha_0}(X)}{\ell_{\alpha_0}(Y)} = \frac{\ell_{\alpha_1}(X)}{\ell_{\alpha_1}(Y)} = \dots = \frac{\ell_{\alpha_k}(X)}{\ell_{\alpha_k}(Y)}.$$

In particular,

$$\frac{\ell_\alpha(X)}{\ell_\alpha(Y)} = \frac{\ell_\delta(X)}{\ell_\delta(Y)}.$$

By the arbitrariness of  $\alpha$  and  $\delta$ , we see that there exists a constant  $K$ , such that

$$\frac{\ell_\gamma(X)}{\ell_\gamma(Y)} \equiv K, \quad \forall \gamma \in \mathcal{C}^0(S_{g,n}).$$

This implies that  $K = 1$ . (Otherwise,  $d_{\text{Th}}(Y, X) < 0$  if  $K < 1$ , or  $d_{\text{Th}}(X, Y) < 0$  if  $K > 1$ .) Therefore,  $Y = X$ . □

#### 4. FLATS IN THE UNIT TANGENT SPHERES

There are flat places on the unit sphere  $T_X^1 \mathcal{T}(S_{g,n})$  of  $T_X \mathcal{T}(S_{g,n})$ . A *facet* is a maximal flat portion of  $T_X^1 \mathcal{T}(S_{g,n})$  which has maximum possible dimension so that it has interior.

**Theorem 4.1** ([Thu98], Theorem 10.1). *There is a bijection between the set of facets on the unit sphere  $T_X^1 \mathcal{T}(S_{g,n})$  and the set of simple closed curves. In other words, every facet is contained in a plane  $d_X \log \ell_\alpha = 1$  for some simple closed curve  $\alpha$ .*

For each simple closed curve  $\alpha$ , let

$$F_X(\alpha) := \{v \in T_X^1 \mathcal{T}(S_{g,n}) : (d_X \log \ell_\alpha)(v) = 1\}$$

be the corresponding facet obtained in Theorem 4.1.

**Lemma 4.2.** *Let  $\alpha, \beta \in \mathcal{C}^0(S_{g,n})$  be two simple closed curves. Then*

$$i(\alpha, \beta) = 0 \iff \partial F_X(\alpha) \cap \partial F_X(\beta) \neq \emptyset.$$

*Proof.* (i) Assume that  $i(\alpha, \beta) = 0$ . Let  $\lambda$  be a maximal geodesic lamination which contains both  $\alpha$  and  $\beta$ . Let  $v$  be the vector tangent to the stretch path  $\text{stretch}(X, \lambda, t)$ . Let  $X_t := \text{stretch}(X, \lambda, t)$ . Then

$$\frac{\ell_\beta(X_t)}{\ell_\beta(X)} = \frac{\ell_\alpha(X_t)}{\ell_\alpha(X)} = e^t,$$

which implies that  $(d_X \log \ell_\alpha)(v) = (d_X \log \ell_\beta)(v) = 1$ . In particular,

$$v \in \partial F_X(\alpha) \cap \partial F_X(\beta) \neq \emptyset.$$

(ii) Assume that  $\partial F_X(\alpha) \cap \partial F_X(\beta) \neq \emptyset$ . Let  $v \in \partial F_X(\alpha) \cap \partial F_X(\beta)$ . Then

$$(4) \quad \|v\|_{\text{Th}} = \sup_{\mu \in \mathcal{ML}(S)} (d_X \log \ell_\mu)(v) = (d_X \log \ell_\alpha)(v) = (d_X \log \ell_\beta)(v) = 1.$$

For each  $0 \leq s \leq 1$ ,

$$\begin{aligned} & \|s \cdot d_X \log \ell_\alpha + (1-s) \cdot d_X \log \ell_\beta\|_{\text{Th}} \\ &= \sup_{v' \in T_X^1 \mathcal{T}(S_{g,n})} (s \cdot d_X \log \ell_\alpha + (1-s) \cdot d_X \log \ell_\beta)(v') \\ &\geq (s \cdot d_X \log \ell_\alpha + (1-s) \cdot d_X \log \ell_\beta)(v) \\ &= 1. \quad (\text{by (4)}) \end{aligned}$$

On the other hand,

$$\|s \cdot d_X \log \ell_\alpha + (1-s) \cdot d_X \log \ell_\beta\|_{\text{Th}} \leq s \|d_X \log \ell_\alpha\|_{\text{Th}} + (1-s) \|d_X \log \ell_\beta\|_{\text{Th}} = 1.$$

Consequently,

$$\|s \cdot d_X \log \ell_\alpha + (1-s) \cdot d_X \log \ell_\beta\|_{\text{Th}} = 1, \quad \forall 0 \leq s \leq 1.$$

In particular, the segment

$$\{s \cdot d_X \log \ell_\alpha + (1-s) \cdot d_X \log \ell_\beta : 0 \leq s \leq 1\} \subset T_X^* \mathcal{T}(S_{g,n})$$

is contained in the unit sphere of the cotangent space  $T_X^* \mathcal{T}(S_{g,n})$ . It then follows from Theorem 1.2 that for each  $0 \leq s \leq 1$ , there exists a unique  $\mu_s \in \mathcal{PML}(S)$  such that

$$(5) \quad s \cdot d_X \log \ell_\alpha + (1-s) \cdot d_X \log \ell_\beta = d_X \log \ell_{\mu_s}.$$

Suppose to the contrary that  $i(\alpha, \beta) > 0$ . Let  $\widehat{\mu}_s$  be a maximal geodesic lamination obtained from  $\mu_s$  by adding isolated leaves. In other words,  $\mu_s$  is the maximal measured geodesic lamination contained in  $\widehat{\mu}_s$ . Let  $v_s \in T_X \mathcal{T}(S_{g,n})$  be the vector tangent to the stretch path  $\text{stretch}(X, \widehat{\mu}_s, t)$  at  $X$ . Then by Corollary 2.2,

$$(6) \quad (d_X \log \ell_\mu)(v_s) \leq 1, \quad \forall \mu \in \mathcal{ML}(S_{g,n})$$

and

$$(7) \quad (d_X \log \ell_\mu)(v_s) = 1 \iff \mu \subset \mu_s.$$

On the other hand, since  $i(\alpha, \beta) > 0$ , it follows either  $\alpha$  or  $\beta$  is not contained in  $\widehat{\mu}_s$ . Consequently,

$$(d_X \log \ell_\alpha)(v_s) < 1$$

or

$$(d_X \log \ell_\beta)(v_s) < 1.$$

Combined with (6), this implies that

$$(s \cdot d_X \log \ell_\alpha + (1-s) \cdot d_X \log \ell_\beta)(v_s) < 1 = (d_X \log \ell_{\mu_s})(v_s), \quad \forall 0 < s < 1.$$

This contradicts to (5), which proves the lemma.  $\square$

**Proposition 4.3.** *Suppose that  $S_{g,n}$  and  $S_{g',n'}$  are not spheres with four or fewer punctures, nor tori with one or fewer punctures. Then every surjective  $\mathbb{R}$ -linear isometry*

$$\Phi : (T_X \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}}) \rightarrow (T_Y \mathcal{T}(S_{g',n'}), \|\bullet\|_{\text{Th}})$$

*induces an isomorphism  $\widehat{\Phi} : \mathcal{C}(S_{g,n}) \rightarrow \mathcal{C}(S_{g',n'})$  such that for every  $\alpha \in \mathcal{C}^0(S_{g,n})$ ,*

- (a)  $\Phi(F_X(\alpha)) = F_Y(\widehat{\Phi}(\alpha))$ , and
- (b)  $\widehat{\Phi}(\alpha)$  is separating if and only if  $\alpha$  is separating.

*Proof.* By assumption,  $\Phi$  induces a bijection between the set of facets of  $T_X^1 \mathcal{T}(S_{g,n})$  and the set of faces of  $T_Y^1 \mathcal{T}(S_{g',n'})$ . By Theorem 4.1,  $\Phi$  also induces a bijection  $\widehat{\Phi}$  between  $\mathcal{C}^0(S_{g,n})$  and  $\mathcal{C}^0(S_{g',n'})$ . Moreover, it follows from Lemma 4.2 that  $\widehat{\Phi}$  preserves the disjointness of simple closed curves. This implies that  $\widehat{\Phi} : \mathcal{C}(S_{g,n}) \rightarrow \mathcal{C}(S_{g',n'})$  is a simplicial isomorphism. This proves the statement (a).

Let us now consider (b). If neither  $(g,n)$  nor  $(g',n')$  equals  $(1,2)$ , the statement follows from Theorem 2.4 and Lemma 2.5. Now assume that one of  $(g,n)$  and  $(g',n')$  equals  $(1,2)$ , say  $(g,n)$ , then by Lemma 2.5  $(g',n') = (1,2)$  or  $(0,5)$ .

Notice that the map  $\Phi$  induces an  $\mathbb{R}$ -linear isometry

$$\Gamma : (T_X^* \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}}) \rightarrow (T_Y^* \mathcal{T}(S_{g',n'}), \|\bullet\|_{\text{Th}})$$

so that

$$\Gamma \left( \frac{d_X \ell_\alpha}{\ell_\alpha(X)} \right) = \frac{d_Y \ell_{\widehat{\Phi}(\alpha)}}{\ell_{\widehat{\Phi}(\alpha)}(Y)}$$

for every simple closed  $\alpha$ . Since the set of weighted simple closed curves is dense in  $\mathcal{ML}(S_{g,n})$ , it follows that the identity above also holds for any measured lamination. Applying an argument similar to the proof of Theorem 3.1, we see that there exists a positive constant  $K$  such that  $\forall \alpha \in \mathcal{C}^0(S_{g,n})$

$$(8) \quad \frac{\ell_\alpha(X)}{\ell_{\widehat{\Phi}(\alpha)}(Y)} \equiv K.$$

By Theorem 3.1 and Lemma 2.5, we see that  $\widehat{\Phi}$  is compatible with Dehn twists. Namely,

$$(9) \quad \widehat{\Phi}(D_\alpha^n \beta) = D_{\widehat{\Phi}(\alpha)}^n \widehat{\Phi}(\beta)$$

holds for any integer  $n$  and any simple closed curves  $\alpha$  and  $\beta$ , where  $D_\alpha^n \beta$  is the  $n$ -th Dehn twist of  $\beta$  along  $\alpha$ . Considering the length functions, we have

$$(10) \quad \lim_{n \rightarrow \infty} \frac{\ell_{D_{\widehat{\Phi}(\alpha)}^n \widehat{\Phi}(\beta)}(Y)}{|n| \ell_{\widehat{\Phi}(\alpha)}(Y) i(\widehat{\Phi}(\alpha), \widehat{\Phi}(\beta))} = 1$$

and

$$(11) \quad \lim_{n \rightarrow \infty} \frac{\ell_{D_\alpha^n \beta}(X)}{|n| \ell_\alpha(X) i(\alpha, \beta)} = 1,$$

where  $i(\cdot, \cdot)$  represents the geometric intersection number.

Suppose to the contrary that  $\widehat{\Phi}$  sends some non-separating curve  $\alpha$  to a separating curve  $\widehat{\Phi}(\alpha)$ . Let  $\gamma$  be a separating curve on  $S_{1,2}$  with  $i(\alpha, \gamma) = 0$ . Consider the component of  $S_{1,2} - \gamma$  which contains  $\alpha$ , and the component of  $S_{g',n'} - \widehat{\Phi}(\gamma)$  which contains  $\widehat{\Phi}(\alpha)$ . By Remark 2.6, we see that  $\widehat{\Phi}$  induces an isomorphism  $\mathcal{C}(S_{1,1}) \rightarrow$

$\mathcal{C}(S_{0,4})$  which is still denoted by  $\widehat{\Phi}$  for simplicity. Let  $\beta$  be simple closed curve on  $S_{g',n'} - \widehat{\Phi}(\gamma)$  with  $i(\alpha, \beta) = 1$ . Then  $\widehat{\Phi}(\beta)$  is a simple closed curve on  $S_{g',n'} - \widehat{\Phi}(\gamma)$  with  $i(\widehat{\Phi}(\alpha), \widehat{\Phi}(\beta)) = 2$ . Combined with Equations (9), (10) and (11), this implies that

$$\lim_{n \rightarrow \infty} \frac{\ell_{D_\alpha^n \beta}(X)}{\ell_{\widehat{\Phi}(D_\alpha^n \beta)}(Y)} = \frac{1}{2} \frac{\ell_\alpha(X)}{\ell_{\widehat{\Phi}(\alpha)}(Y)},$$

which contradicts Equation (8). This completes the proof.  $\square$

## 5. PROOF OF THEOREMS

In this final section, we shall prove Theorem 1.4, 1.5, 1.6, 1.7, 1.8, which are stated in the introduction.

*Proof of Theorem 1.4.* We divide the proof into two cases depending on whether  $(g, n)$  belongs to  $\{(0, 4), (1, 1)\}$  or not.

**Case 1:**  $(g, n) \notin \{(0, 4), (1, 1)\}$ .

By duality,  $\Phi^*$  induces an  $\mathbb{R}$ -linear surjective isometry

$$\Phi : (T_Y \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}}) \rightarrow (T_X \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}}).$$

By Proposition 4.3 and Theorem 2.4, we see that there exists a mapping class  $\phi \in \mathcal{MCG}^\pm(S)$  such that for every simple closed curve  $\alpha$

$$\Phi(F_X(\alpha)) = F_Y(\phi\alpha),$$

where

$$F_X(\alpha) = \{v \in T_X^1 \mathcal{T}(S_{g,n}) : (d_X \log \ell_\alpha)(v) = 1\}$$

and

$$F_Y(\phi\alpha) = \{u \in T_Y^1 \mathcal{T}(S_{g,n}) : (d_Y \log \ell_{\phi\alpha})(u) = 1\}.$$

Therefore,

$$(12) \quad \Phi^*(d_X \log \ell_\alpha) = d_Y \log \ell_{\phi\alpha}.$$

for every simple closed curve  $\alpha$ .

To complete the proof, it remains to show that  $\phi : X \rightarrow Y$  is (isotopic to) an isometry.

Notice that  $\phi$  induces an  $\mathbb{R}$ -linear surjective isometry:

$$\begin{aligned} \widehat{\phi} : T_Y^* \mathcal{T}(S_{g,n}) &\longrightarrow T_{\phi^{-1}Y}^* \mathcal{T}(S_{g,n}) \\ d_Y \ell_\mu &\longmapsto d_{\phi^{-1}Y} \ell_{\phi^{-1}\mu}. \end{aligned}$$

Consequently,

$$\widehat{\phi} \circ \Phi^* : T_X^* \mathcal{T}(S_{g,n}) \longrightarrow T_{\phi^{-1}Y}^* \mathcal{T}(S_{g,n})$$

is also an  $\mathbb{R}$ -linear surjective isometry. Moreover, by (12), we see that for every simple closed curve  $\alpha$ ,

$$\widehat{\phi} \circ \Phi^*(d_X \log \ell_\alpha) = d_{\phi^{-1}Y} \log \ell_\alpha.$$

Then

$$\begin{aligned} \widehat{\phi} \circ \Phi^*(d_X \ell_\alpha) &= \ell_\alpha(X) \cdot (\widehat{\phi} \circ \Phi^*)(d_X \log \ell_\alpha) \\ &= \ell_\alpha(X) \cdot d_{\phi^{-1}Y} \log \ell_\alpha \\ &= \frac{\ell_\alpha(X)}{\ell_\alpha(\phi^{-1}Y)} d_{\phi^{-1}Y} \ell_\alpha \end{aligned}$$

for every simple closed curve  $\alpha$ . Since the set of weighted simple closed curves is dense in  $\mathcal{ML}(S_{g,n})$ , it follows that  $\widehat{\phi} \circ \Phi^*$  coincides with  $\Gamma_{X,\phi^{-1}Y}$  on a dense subset of  $T_X^* \mathcal{T}(S_{g,n})$  (see (3) for the definition of  $\Gamma_{X,\phi^{-1}Y}$ ). By continuity, this implies that

$$\widehat{\phi} \circ \Phi^* = \Gamma_{X,\phi^{-1}Y}.$$

In particular,  $\Gamma_{X,g^{-1}Y}$  is linear. It then follows from Theorem 3.1 that  $X = \phi^{-1}Y$ .

**Case 2:**  $(g, n) \in \{(0, 4), (1, 1)\}$ .

The case that  $(g, n) = (1, 1)$  is treated in [DLRT20]. The proof for the case  $(g, n) = (0, 4)$  is essentially the same as that of the case  $(g, n) = (1, 1)$ . For completeness, we include a proof for  $(g, n) = (0, 4)$ .

By Theorem 4.1, the isometry

$$\Phi : (T_X \mathcal{T}(S_{0,4}), \|\bullet\|_{\text{Th}}) \rightarrow (T_Y \mathcal{T}(S_{0,4}), \|\bullet\|_{\text{Th}})$$

induces a bijection  $\widehat{\Phi} : \mathcal{C}^0(S_{0,4}) \rightarrow \mathcal{C}^0(S_{0,4})$  between the sets of simple closed curves such that

$$\Phi(F_X(\alpha)) = F_Y(\widehat{\Phi}(\alpha)).$$

Let  $\alpha$  a simple closed curve such that both  $\ell_\alpha(X) > L_2$  where  $L_2$  is the constant Proposition 2.12. By Proposition 2.12, there exist  $N > 0$ ,  $n_0 \in \mathbb{Z}$ , and a simple closed curve  $\beta$  with  $i(\beta, \alpha) = 2$ , such that for all  $n > N$ ,

$$(13) \quad \widehat{\Phi}(D_\alpha^n(\beta)) = D_{\widehat{\Phi}(\alpha)}^{n+n_0}(\widehat{\Phi}(\beta)).$$

Since  $\Phi$  is an isometry, it follows that  $|F_X(D_\alpha^n(\beta))| = |F_Y(D_{\widehat{\Phi}(\alpha)}^{n+n_0}(\gamma))|$ . Combined with Theorem 2.11, this implies that

$$\ell_\alpha(X) = \ell_{\widehat{\Phi}(\alpha)}(Y).$$

Moreover, there exists  $N_2 > 0$  such that for all  $n > N_2$ , we have

$$\ell_{D_\alpha^n(\beta)}(X) > L_2, \quad \ell_{D_{\widehat{\Phi}(\alpha)}^{n+n_0}\gamma}(Y) > L_2.$$

Applying the argument above to  $D_\alpha^n\beta$  and  $\widehat{\Phi}(D_\alpha^n\beta) = D_{\widehat{\Phi}(\alpha)}^{n+n_0}(\gamma)$ , we see that for all  $n > N_2$ ,

$$(14) \quad \ell_{D_\alpha^n(\beta)}(X) = \ell_{D_{\widehat{\Phi}(\alpha)}^{n+n_0}\gamma}(Y).$$

Let  $m > N_2$  be such that  $d\ell_\alpha$  and  $d\ell_{D_\alpha^m\beta}$  are linearly independent in  $T_X^* \mathcal{T}(S_{0,4})$ . Since  $i(\alpha, D_\alpha^m\beta) = i(\widehat{\Phi}(\alpha), D_{\widehat{\Phi}(\alpha)}^{m+n_0}\gamma) = 2$ , it follows that there exists a mapping class  $f \in \mathcal{MCG}^\pm(S_{g,n})$  such that  $Y = fX$  and

$$f(\alpha) = \widehat{\Phi}(\alpha), \quad f(D_\alpha^m\beta) = \widehat{\Phi}(D_\alpha^m\beta) = D_{\widehat{\Phi}(\alpha)}^{m+n_0}\gamma.$$

Notice that  $f$  induces an  $\mathbb{R}$ -linear isometry

$$\begin{aligned} f^* : (T_X^* \mathcal{T}(S_{0,4}), \|\bullet\|_{\text{Th}}) &\longrightarrow (T_{fX}^* \mathcal{T}(S_{0,4}), \|\bullet\|_{\text{Th}}) \\ d\ell_\mu &\longmapsto d\ell_{f\mu}. \end{aligned}$$

Consequently, the composition map

$$(f^*)^{-1} \circ \Phi^* : (T_X^* \mathcal{T}(S_{0,4}), \|\bullet\|_{\text{Th}}) \longrightarrow (T_X^* \mathcal{T}(S_{0,4}), \|\bullet\|_{\text{Th}})$$

is an  $\mathbb{R}$ -linear isometry with

$$(f^*)^{-1} \circ \Phi^*(d\ell_\alpha) = d\ell_\alpha, \quad (f^*)^{-1} \circ \Phi^*(d\ell_{D_\alpha^m\beta}) = d\ell_{D_\alpha^m\beta}.$$

Since  $d\ell_\alpha$  and  $d\ell_{D_\alpha^n\beta}$  are linearly independent, this implies that  $(f^*)^{-1} \circ \Phi^*$  is the identity map. Therefore  $\Phi^* = f^*$ . This completes the proof.  $\square$

*Proof of Theorem 1.5.* Considering the dimension  $\dim_{\mathbb{R}}(\mathcal{T}(S_{g,n})) = 6g - 6 + 2n$ , we see that  $(g, n) \in \{(0, 4), (1, 1)\}$  if and only if  $(g', n') \in \{(0, 4), (1, 1)\}$ . We divide the proof into two cases depending on whether  $(g, n)$  and  $(g', n')$  belong to  $\{(0, 4), (1, 1)\}$  or not.

**Case 1:**  $(g, n) \notin \{(0, 4), (1, 1)\}$  and  $(g', n') \notin \{(0, 4), (1, 1)\}$ . In this case, Proposition 4.3 implies that  $\Phi^*$  induces an isomorphism  $\widehat{\Phi} : \mathcal{C}(S_{g,n}) \rightarrow \mathcal{C}(S_{g',n'})$  which preserves curve type. It then follows from Theorem 2.4 that either

- $S_{g,n}$  and  $S_{g',n'}$  are homeomorphic, or
- $\{S_{g,n}, S_{g',n'}\}$  is one of  $\{S_{0,5}, S_{1,2}\}, \{S_{0,6}, S_{2,0}\}$ .

On the other hand,  $S_{1,1}$  (resp.  $S_{2,0}$ ) contains non-separating simple closed curves, while all simple closed curves on  $S_{0,5}$  (resp.  $S_{0,6}$ ) are separating. This implies that  $\{S_{g,n}, S_{g',n'}\}$  is neither  $\{S_{0,5}, S_{1,2}\}$  nor  $\{S_{0,6}, S_{2,0}\}$ . Therefore,  $S_{g,n}$  and  $S_{g',n'}$  are homeomorphic.

**Case 2:**  $(g, n) \in \{(0, 4), (1, 1)\}$  and  $(g', n') \in \{(0, 4), (1, 1)\}$ . Suppose to the contrary that  $S_{g,n}$  and  $S_{g',n'}$  are not homeomorphic. Then  $\{S_{g,n}, S_{g',n'}\} = \{(0, 4), (1, 1)\}$ . Without loss of generality we may assume that  $S_{g,n} = S_{0,4}$  and  $S_{g',n'} = S_{1,1}$ . By Theorem 4.1, the isometry

$$\Phi : (T_X \mathcal{T}(S_{0,4}), \|\bullet\|_{\text{Th}}) \rightarrow (T_Y \mathcal{T}(S_{1,1}), \|\bullet\|_{\text{Th}})$$

induces a bijection  $\widehat{\Phi} : \mathcal{C}^0(S_{0,4}) \rightarrow \mathcal{C}^0(S_{1,1})$  between the sets of simple closed curves such that

$$\Phi(F_X(\alpha)) = F_Y(\widehat{\Phi}(\alpha)).$$

Let  $\alpha$  a simple closed curve such that both  $\ell_\alpha(X) > L_2$  and  $\ell_{\widehat{\alpha}}(Y) > L_1$ , where  $L_1$  and  $L_2$  are the constants from Proposition 2.9 and Proposition 2.12. By Proposition 2.9 and Proposition 2.12, there exist  $N > 0$ ,  $n_0 \in \mathbb{Z}$ , simple closed curves  $\beta$  with  $i(\beta, \alpha) = 2$  and  $\gamma$  with  $i(\gamma, \widehat{\Phi}(\alpha)) = 1$ , such that for all  $n > N$ ,

$$\widehat{\Phi}(D_\alpha^n \beta) = D_{\widehat{\Phi}(\alpha)}^{n+n_0} \gamma.$$

Since  $\Phi$  is an isometry, it follows that  $|F_X(D_\alpha^n \beta)| = |F_Y(D_{\widehat{\Phi}(\alpha)}^{n+n_0} \gamma)|$ . Combined with Theorem 2.8 and Theorem 2.11, this implies that

$$(15) \quad \ell_\alpha(X) = \ell_{\widehat{\Phi}(\alpha)}(Y).$$

Moreover, there exists  $N_2 > 0$  such that for all  $n > N_2$ , we have

$$\ell_{D_\alpha^n \beta}(X) > L_2, \quad \ell_{D_{\widehat{\Phi}(\alpha)}^{n+n_0} \gamma}(Y) > L_1.$$

Applying the argument above to  $D_\alpha^n \beta$  and  $\widehat{\Phi}(D_\alpha^n \beta) = D_{\widehat{\Phi}(\alpha)}^{n+n_0} \gamma$ , we see that for all  $n > N_2$ ,

$$(16) \quad \ell_{D_\alpha^n \beta}(X) = \ell_{D_{\widehat{\Phi}(\alpha)}^{n+n_0} \gamma}(Y).$$

On the other hand, we have

$$\lim_{n \rightarrow +\infty} \frac{\ell_{D_\alpha^n \beta}(X)}{n} = 2\ell_\alpha(X)$$

and

$$\lim_{n \rightarrow +\infty} \frac{\ell_{D_{\widehat{\Phi}(\alpha)}^{n+n_0} \gamma}(Y)}{n} = \ell_{\widehat{\Phi}(\alpha)}(Y).$$



These contradict Equations (15) and (16). Therefore,  $\{S_{g,n}, S_{g',n'}\} \neq \{S_{0,4}, S_{1,1}\}$ . This completes the proof.  $\square$

*Proof of Theorem 1.6.* The proof we present here follows essentially that of [DLRT20, Theorem 1.5]. Let  $F : (U, d_{\text{Th}}) \rightarrow (\mathcal{T}(S_{g,n}), d_{\text{Th}})$  be an isometric embedding. By [DLRT20, Theorem 6.1] the Thurston norm is locally Lipschitz (locally  $C_{loc}^{0,1}$ ). It then follows from [MT17, Theorem A, Theorem B] that  $F$  is  $C_{loc}^{1,1}$  and its differential is norm-preserving. Therefore, for each  $X \in U$ ,

$$d_X F : T_X \mathcal{T}(S_{g,n}) \longrightarrow T_{F(X)} \mathcal{T}(S_{g,n})$$

is an  $\mathbb{R}$ -linear surjective isometry for the Thurston norm. By duality,  $d_X F$  induces an  $\mathbb{R}$ -linear surjective isometry

$$d_X^* F : (T_{F(X)}^* \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}}) \longrightarrow (T_X^* \mathcal{T}(S_{g,n}), \|\bullet\|_{\text{Th}}).$$

By Theorem 1.4, there exists  $\phi_F(X) \in \mathcal{MCG}^\pm(S_{g,n})$  such that

$$(17) \quad F(X) = \phi_F(X)(X).$$

Let  $j : \mathcal{MCG}^\pm(S_{g,n}) \rightarrow \text{Isom}(\mathcal{T}(S_{g,n}), d_{\text{Th}})$  be the natural homomorphism. Let  $\widehat{U} \subset U$  be the subset of points with trivial stablizer in  $j(\mathcal{MCG}^\pm(S_{g,n}))$ . Then  $\widehat{U}$  is an open and dense subset of  $U$ . Let  $X_0 \in \widehat{U}$ . By the proper discontinuity action of  $\mathcal{MCG}^\pm(S_{g,n})$  on  $\mathcal{T}(S_{g,n})$ , we see that there exist a small neighbourhood  $U_0 \subset U$  of  $X_0$  and a small neighbourhood  $V_0 \subset F(U)$  of  $F(X_0)$ , such that

$$\{\phi \in j(\mathcal{MCG}^\pm(S_{g,n})) : \phi(U_0) \cap V_0 \neq \emptyset\} = \{j(\phi_F(X_0))\}.$$

It then follows from the continuity of  $F$  and (17) that

$$j(\phi_F(X)) = j(\phi_F(X_0))$$

for all  $X$  sufficiently close to  $X_0$ . In other words, the map  $j \circ \phi_F : \widehat{U} \rightarrow \text{Isom}(\mathcal{T}(S_{g,n}), d_{\text{Th}})$  is locally constant. Hence,  $j \circ \phi_F$  is constant on each connected component of  $\widehat{U}$ .

For each  $\psi \in j(\mathcal{MCG}^\pm(S_{g,n}))$ , let  $\text{Fix}_\psi \subset \mathcal{T}(S_{g,n})$  be the subset of fixed points. Notice that  $\text{Fix}_\psi$  is a real submanifold.<sup>3</sup> Let  $U_1$  be the union of those  $\text{Fix}_\psi$  which are of real codimension one. Let  $U_{\geq 2}$  be the union of those  $\text{Fix}_\psi$  which are of real codimension at least two. Combining with the proper continuity action of  $\mathcal{MCG}^\pm(S_{g,n})$  on  $\mathcal{T}(S_{g,n})$ , we see that the complement  $U \setminus U_{\geq 2}$  is connected in  $U$ .

Let  $U' = U_1 \setminus U_{\geq 2}$ . Let  $X'$  be an arbitrary point in  $U'$ . Taking  $W \subset U$  to be a small neighbourhood of  $X'$ , we may assume that  $W \setminus U'$  is contained in  $\widehat{U}$  and has exactly two components, which we denote by  $W_+$  and  $W_-$ . By the discussion above, we see that  $j \circ \phi_F$  is constant on  $W_-$  and on  $W_+$ . Let  $\psi_-$  and  $\psi_+$  be the corresponding values. The continuity of  $F$  implies that  $\psi_+ \circ \psi_-^{-1} \in j(\mathcal{MCG}^\pm(S_{g,n}))$  fixes  $W \cap U'$  pointwise, and is therefore the identity or a reflection. In the latter case,  $F$  would map both sides of  $W \cap U'$  to the same side of  $F(W \cap U')$ . This contradicts the fact that  $F$  is  $C^1$  at  $X'$ . Therefore,  $\psi_+ \circ \psi_-^{-1}$  is the identity. This implies that  $j \circ \phi_F$  is locally constant on  $\widehat{U} \cup U'$ . Notice that  $\widehat{U} \cup U' = U \setminus U_{\geq 2}$ , which is both connected and dense by the previous two paragraphs. Therefore,  $j \circ \phi_F$  extends to a constant function on  $U$  with value  $j \circ \phi_F(X_0)$ .  $\square$

<sup>3</sup>This can be seen using the Teichmüller metric. Since  $\mathcal{T}(S_{g,n})$  is uniquely geodesic in the Teichmüller metric [GL00, Section 7.4] and  $\mathcal{MCG}^\pm(S_{g,n})$  acts as isometries with respect to the Teichmüller metric, it follows that  $\text{Fix}_\psi$  is a totally geodesic submanifold of  $\mathcal{T}(S_{g,n})$ .

*Proof of Theorem 1.7.* Let  $j : \mathcal{MCG}^\pm(S_{g,n}) \rightarrow \text{Isom}(\mathcal{T}(S_{g,n}), d_{\text{Th}})$  be the natural homomorphism. By Theorem 1.6, we see that  $j$  is surjective. The theorem then follows from the following fact ([Bir75, Vir72])

$$\text{Kernel}(j) \simeq \begin{cases} \mathbb{Z}_2, & \text{if } (g, n) \in \{(2, 0), (1, 1), (1, 2)\}, \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2, & \text{if } (g, n) = (0, 4), \\ \{\text{id}\}, & \text{otherwise.} \end{cases}$$

□

*Proof of Theorem 1.8.* This follows from Theorem 1.5 and [DLRT20, Theorem 6.1]. □

## REFERENCES

- [AD19] Daniele Alessandrini and Valentina Disarlo. Generalized stretch lines for surfaces with boundary. *Int Math Res Not IMRN*: 18919–18991, 2022.
- [AP98] Marco Abate and Giorgio Patrizio. Isometries of the Teichmüller metric. *Annali della Scuola Normale Superiore di Pisa. Classe di Scienze. Serie IV*, 26(3):437–452, 1998.
- [Bir75] Joan S. Birman, Braids, links, and mapping class groups, *Ann. Math. Stud.*, vol. 82, Princeton Univ. Press, Princeton, NJ, 1975.
- [BM07] Jeffrey Brock and Dan Margalit. Weil-Petersson isometries via the pants complex. *Proceedings of the American Mathematical Society*, 135(3):795–803, 2007.
- [CB88] Andrew Casson and Steven A Bleiler. *Automorphisms of surfaces after Nielsen and Thurston*. Number 9. Cambridge University Press, 1988.
- [CR07] Young-Eun Choi and Kasra Rafi. Comparison between Teichmüller and Lipschitz metrics. *Journal of the London Mathematical Society. Second Series*, 76(3):739–756, 2007.
- [DLRT20] David Dumas, Anna Lenzhen, Kasra Rafi, and Jing Tao. Coarse and fine geometry of the thurston metric. *Forum of Mathematics, Sigma* (2020), Vol. 8, e28, 58 pages.
- [EG96] Clifford J. Earle and Frederick P. Gardiner. Geometric isomorphisms between infinite-dimensional Teichmüller spaces. *Transactions of the American Mathematical Society*, 348(3):1163–1190, 1996.
- [EK74a] Clifford J. Earle and Irwin Kra. On holomorphic mappings between Teichmüller spaces. In *Contributions to analysis (a collection of papers dedicated to Lipman Bers)*, pages 107–124. 1974.
- [EK74b] Clifford J. Earle and Irwin Kra. On isometries between Teichmüller spaces. *Duke Mathematical Journal*, 41:583–591, 1974.
- [EM03] Clifford J. Earle and V. Markovic. Isometries between the spaces of  $L^1$  holomorphic quadratic differentials on Riemann surfaces of finite type. *Duke Mathematical Journal*, 120(2):433–440, 2003.
- [FLP12] Albert Fathi, François Laudenbach, and Valentin Poénaru. *Thurston’s work on surfaces, Translated from the 1979 French original by Djun M. Kim and Dan Margalit*. Princeton University Press, 2012.
- [FW10] Benson Farb and Shmuel Weinberger. The intrinsic asymmetry and inhomogeneity of Teichmüller space. *Duke Mathematical Journal*, 155(1):91–103, 2010.
- [GL00] Gardiner, Frederick P. and Lakic, Nikola . Quasiconformal Teichmüller theory. *Mathematical Surveys and Monographs*, 76. American Mathematical Society, Providence, RI, 2000. xx+372 pp.
- [Har81] W. J. Harvey. Boundary structure of the modular group. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, volume 97 of *Ann. of Math. Stud.*, pages 245–251. Princeton Univ. Press, Princeton, N.J., 1981.
- [HP19] Yi Huang and Athanase Papadopoulos. Optimal lipschitz maps on one-holed tori and the thurston metric theory of teichmueller space. *arXiv:1909.03639*, 2019.
- [HOP21] Yi Huang, Ken’Ichi Ohshika and Athanase Papadopoulos. The infinitesimal and global Thurston geometry Of Teichmuüller Space. *arXiv:2111.13381*, 2021.
- [Iva97] Nikolai V. Ivanov. Automorphism of complexes of curves and of Teichmüller spaces. *International Mathematics Research Notices, IMRN*, (14):651–666, 1997.
- [Iva01] Nikolai V. Ivanov. Isometries of Teichmüller spaces from the point of view of Mostow rigidity. In *Topology, ergodic theory, real algebraic geometry*, volume 202 of *Amer. Math. Soc. Transl. Ser. 2*, pages 131–149. Amer. Math. Soc., Providence, RI, 2001.
- [Ker83] Steven P. Kerckhoff. The Nielsen realization problem. *Annals of Mathematics. Second Series*, 117(2):235–265, 1983.

- [Kor99] Mustafa Korkmaz. Automorphisms of complexes of curves on punctured spheres and on punctured tori. *Topology and its Applications*, 95(2):85–111, 1999.
- [Lak97] Nikola Lakic. An isometry theorem for quadratic differentials on Riemann surfaces of finite genus. *Transactions of the American Mathematical Society*, 349(7):2951–2967, 1997.
- [Liu00] Li-xin Liu. Asymmetry of Thurston’s pseudometric. *Acta Scientiarum Naturalium Universitatis Sunyatseni. Zhongshan Daxue Xuebao. Ziran Kexue Ban*, 39(6):6–9, 2000.
- [LPST13] L. Liu, A. Papadopoulos, W. Su, and G. Théret. On the classification of mapping class actions on Thurston’s asymmetric metric. *Mathematical Proceedings of the Cambridge Philosophical Society*, 155(3):499–515, 2013.
- [LRT12] Anna Lenzen, Kasra Rafi, and Jing Tao. Bounded combinatorics and the Lipschitz metric on Teichmüller space. *Geometriae Dedicata*, 159:353–371, 2012.
- [LRT15] Anna Lenzen, Kasra Rafi, and Jing Tao. The shadow of a Thurston geodesic to the curve graph. *Journal of Topology*, 8(4):1085–1118, 2015.
- [Luo00] Feng Luo. Automorphisms of Thurston’s space of measured laminations. In *In the tradition of Ahlfors and Bers (Stony Brook, NY, 1998)*, volume 256 of *Contemp. Math.*, pages 221–225. Amer. Math. Soc., Providence, RI, 2000.
- [Mar03] Vladimir Markovic. Biholomorphic maps between Teichmüller spaces. *Duke Mathematical Journal*, 120(2):405–431, 2003.
- [Mar04] Dan Margalit. Automorphisms of the pants complex. *Duke Mathematical Journal*, 121(3):457–479, 2004.
- [MM13] Ryosuke Mineyama and Hideki Miyachi. A characterization of biholomorphic automorphisms of Teichmüller space. *Mathematical Proceedings of the Cambridge Philosophical Society*, 154(1):71–83, 2013.
- [MT17] Vladimir S. Matveev and Marc Troyanov. The Myers-Steenrod theorem for Finsler manifolds of low regularity. *Proceedings of the American Mathematical Society*, 145(6):2699–2712, 2017.
- [MW02] Howard Masur and Michael Wolf. The Weil-Petersson isometry group. *Geometriae Dedicata*, 93:177–190, 2002.
- [PS14] Athanase Papadopoulos and Weixu Su. Thurston’s metric on Teichmüller space and isomorphisms between Fuchsian groups. In *Analysis and geometry of discrete groups and hyperbolic spaces*, RIMS Kôkyûroku Bessatsu, B48, pages 95–109. Res. Inst. Math. Sci. (RIMS), Kyoto, 2014.
- [PS15] Athanase Papadopoulos and Weixu Su. On the Finsler structure of Teichmüller’s metric and Thurston’s metric. *Expositiones Mathematicae*, 33(1):30–47, 2015.
- [PS16] Athanase Papadopoulos and Weixu Su. Thurston’s metric on Teichmüller space and the translation distances of mapping classes. *Annales Academiæ Scientiarum Fennicæ. Mathematica*, 41(2):867–879, 2016.
- [PT07a] Athanase Papadopoulos and Guillaume Théret. On Teichmüller’s metric and Thurston’s asymmetric metric on Teichmüller space. In *Handbook of Teichmüller theory. Vol. I*, volume 11 of *IRMA Lect. Math. Theor. Phys.*, pages 111–204. Eur. Math. Soc., Zürich, 2007.
- [PT07b] Athanase Papadopoulos and Guillaume Théret. On the topology defined by Thurston’s asymmetric metric. *Mathematical Proceedings of the Cambridge Philosophical Society*, 142(3):487–496, 2007.
- [PT12] Athanase Papadopoulos and Guillaume Théret. Some Lipschitz maps between hyperbolic surfaces with applications to Teichmüller theory. *Geometriae Dedicata*, 161:63–83, 2012.
- [Put08] Andrew Putman. A note on the connectivity of certain complexes associated to surfaces. *L’Enseignement Mathématique. Revue Internationale. 2e Série*, 54(3-4):287–301, 2008.
- [PY17] Athanase Papadopoulos and Sumio Yamada. Deforming hyperbolic hexagons with applications to the arc and the Thurston metrics on Teichmüller spaces. *Monatshefte für Mathematik*, 182(4):913–939, 2017.
- [Pat72] David D. Patterson. The Teichmüller spaces are distinct. *Proceedings of the American Mathematical Society*, 35(1), 179–182, 1972.
- [Roy71] H. L. Royden. Automorphisms and isometries of Teichmüller space. In *Advances in the Theory of Riemann Surfaces (Proc. Conf., Stony Brook, N.Y., 1969)*, pages 369–383. Ann. of Math. Studies, No. 66. Princeton Univ. Press, Princeton, N.J., 1971.
- [Su16] W. Su. Problems on the Thurston metric. In *Handbook of Teichmüller theory. Vol. V*, 55–72, IRMA Lect. Math. Theor. Phys., 26, Eur. Math. Soc., Zürich, 2016.
- [The07] Guillaume Theret. On the negative convergence of Thurston’s stretch lines towards the boundary of Teichmüller space. *Annales Academiæ Scientiarum Fennicæ. Mathematica*, 32(2):381–408, 2007.
- [The14] Guillaume Theret. Convexity of length functions and Thurston’s shear coordinates, preprint, arXiv:1408.5771.

- [Thu86] William P. Thurston. Earthquakes in two-dimensional hyperbolic geometry. In *Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984)*, volume 112 of *London Math. Soc. Lecture Note Ser.*, pages 91–112. Cambridge Univ. Press, Cambridge, 1986.
- [Thu98] William P Thurston. Minimal stretch maps between hyperbolic surfaces. *arXiv preprint math/9801039*, 1998.
- [Vir72] Oleg Viro, Links, two-fold branched coverings and braids, *Soviet Math. Sbornik* 87 (2) 216–228, 1972.
- [Wal14] Cormac Walsh. The horoboundary and isometry group of Thurston’s Lipschitz metric. In *Handbook of Teichmüller theory. Vol. IV*, volume 19 of *IRMA Lect. Math. Theor. Phys.*, pages 327–353. Eur. Math. Soc., Zürich, 2014.
- [Wol82] Scott Wolpert. The Fenchel-Nielsen deformation. *Annals of Mathematics. Second Series*, 115(3):501–528, 1982.
- [Wol83] Scott Wolpert. On the symplectic geometry of deformations of a hyperbolic surface. *Annals of Mathematics. Second Series*, 117(2):207–234, 1983.

SCHOOL OF MATHEMATICS, SOUTH CHINA UNIVERSITY OF TECHNOLOGY, WUSHAN RD 381,  
TIANHE, GUANGZHOU, CHINA, 510641

*Email address:* panhp@scut.edu.cn