

# Achieving effective renormalization scale and scheme independence via the Principle of Observable Effective Matching (POEM)

Farrukh A. Chishtie\*

*Department of Applied Mathematics, The University of Western Ontario, London, ON N6A 3K7, Canada*

(Dated: September 18, 2020)

In this work, we explicate a new approach for eliminating renormalization scale and scheme (RSS) dependence in physical observables. We develop this approach by matching RSS dependent observables to a theory which is independent of both these forms of dependencies. We term the fundamental basis behind this approach as the Principle of Observable Effective Matching (POEM), which entails matching of a scale and scheme dependent observable with the fully physical and dynamical scale (PS) dependent theory at the tree and at higher loop orders at which RSS independence is guaranteed. This is aimed towards achieving a so-called “effective” RSS independent expressions, as the resulting dynamical dependence is derived from a particular order in RSS dependent perturbation theory and is sensitive to the matching scale. With this matching, we obtain an “Effective Physical Observable (EPO)”, a finite-order RSS independent version of the RSS dependent observable. We illustrate our approach with a study of the cross section  $R_{e^+e^-}$  for  $e^+e^- \rightarrow$  hadrons, which is demonstrated to achieve scale and scheme independence utilizing the 3- and 4-loop order  $\overline{MS}$  scheme expression in QCD perturbation theory via matching at both one-loop and two-loop orders for obtaining the EPO. At the tree-loop matching, we find physical cutoff  $\Lambda_{eff}$  of the physical observable which is RSS independent. With two-loop matching, we obtain an EPO prediction of  $\frac{3}{11}R_{e^+e^-}^{eff} = 1.056137_{-0.0007}^{+0.0007}$  at  $Q = 31.6 GeV$ , which is in excellent agreement with the experimental value of  $\frac{3}{11}R_{e^+e^-}^{exp} = 1.0527_{-0.005}^{+0.005}$ . Given its new conceptual basis, ease of use and performance, we contend that POEM be explored in its application for obtaining EPOs for other physical observables across domains of high energy theory and phenomenology, as well as other areas of fundamental and applied physics, such as cosmology, statistical and condensed matter physics.

Keywords: Perturbation Theory, Resummation Methods, Perturbative QCD

## I. INTRODUCTION

In perturbative Quantum Field Theory (QFT), infinities are encountered in theoretical expressions of physical observables which require renormalization techniques. Rendering finiteness to such quantities, such as physical cross sections and decay rates introduces renormalization scale and scheme (RSS) dependencies. In contrast to such dependencies, it is surmised and observed that at increasing order in the expansion of a small expansion parameter, along with convergence such dependencies would be reduced and the penultimate expression to be have full dependence only on dynamical physical scales. The issue of RSS dependencies are treated via the Principle of Minimal Sensitivity [1,2]. Other approaches to reducing and/or eliminating scheme dependence includes the approaches of Effective Charge [3], Brodsky-Lepage-Mackenzie (BLM)[4], Renormalization Group (RG) summation [5], RG summation with RS invariants [6-9], Complete Renormalization Group Improvement (CORGI) [10,11], Principle of Maximal Conformality (PMC) [12] and sequential extended BLM (seBLM) [13]. In this work, we detail a new and alternate approach towards achieving RSS in-

dependence, inspired by Effective Field Theory (EFT) techniques. As such, it is based on what we term as the “Principle of Observable Effective Matching” (POEM) and our work is applicable for achieving RSS independence only up to a fixed order in perturbation theory, hence we use the term “effective” for the observable having only Physical Scale (PS) dependence which is also the scale at which matching occurs. We term the derived perturbative expression as an “Effective Physical Observable (EPO)”, which implies these caveats. The paper is outlined as follows: we introduce POEM next and then work out the RSS independent Quantum Chromodynamics (QCD) cross section  $R_{e^+e^-}$  via POEM at the tree, one- and two-loop matching with a RSS independent scheme in the following sections.

## II. THE PRINCIPLE OF OBSERVABLE EFFECTIVE MATCHING (POEM)

Physical observables computed to all orders in perturbative QFT within various RSS approaches are expected to result in scale and scheme independence if results at all orders are computed, and hence must be the same in this limit. This is the starting point for POEM which can be expressed as:

\* Correspondence email address:fachisht@uwo.ca

$$\begin{aligned}
& \lim_{n \rightarrow \infty} O_n^{(i)}(a_n^{(i)}(\mu^{(i)}), m_n^{(i)}(\mu^{(i)}), \mu^{(i)}, Q, M) \quad (1) \\
&= \lim_{l \rightarrow \infty} O_l^{(j)}(a_l^{(j)}(\mu^{(j)}), m_l^{(j)}(\mu^{(j)}), \mu^{(j)}, Q, M) \\
&= O_{full}(a(Q), m(M), Q, M)
\end{aligned}$$

Here  $O$  denotes perturbative contributions to a physical observable such as in a decay rate or a cross-section computed in two schemes denoted by the superscript  $i$  and  $j$  at some loop-order  $n$  and  $l$  respectively, while  $a$  and  $m$  denote coupling and mass respective tied to respective renormalization scales  $\mu^i$  and  $\mu^j$ .  $Q$  and  $m$  denote dynamical PS. Eq. 1 can be further generalized by including multiple schemes, couplings and masses as well as factorization scales and momentum fractions. This equation nevertheless implies that if  $O$  is computed at a certain large limit of perturbation theory, this results in a RSS independent result with dependence only on  $Q$  and  $M$ . While it is rather challenging to find expressions to all orders in perturbation theory, we consider such to be a RSS independent theory or full theory whereby all implicit and explicit renormalization scale within any scheme cancels, and the equalization across in eq. 1 implies an overall RSS independence leading to an independent observable termed as  $O_{full}$  dependent only on scale independent coupling and mass, as well as physical scales,  $Q$  and  $M$ .

While this equation holds for a theory computed at all orders, for practical purposes, and as criteria to derive an effective version of such an all-orders observable, RSS independence holds true at the tree level when no quantum corrections exist and this is also valid at the some  $r$ -loop level at some PS (typically this is at one and two loop level). With these two properties in mind, we propose two conditions which can be applied to achieve an effective RSS independence at an order-by-order basis for truncated expressions of a physical observable. These are:

$$\begin{aligned}
O_n^{(k)}(a_{eff}(\Lambda_{eff}), m_{eff}(\Lambda_{eff}), \Lambda_{eff}, Q, M) \quad (2) \\
= O_{eff}^{(tree)}
\end{aligned}$$

$$\begin{aligned}
O_n^{(k)}(a_n^{(k)}(Q^*), m_n^{(k)}(Q^*), Q^*, Q, M) = \quad (3) \\
O_{eff}^{(r)}(a_{eff}(Q^*), m_{eff}(Q^*), Q^*, Q, M)
\end{aligned}$$

Here,  $O_{eff}$  is conceptualized in a finite-order perturbative physical representation termed as the ‘‘Effective Dynamical Observable’’ (EPO) which is truncated at  $r$ -loop order in order to be consistent with RSS independence, and is dependent only on a PS. Since it is known that RSS independence is guaranteed trivially at

the tree and at higher  $r$ -loop levels at some PS, hence this forms the basis of these equations. We can therefore achieve RSS independence by requiring in eq. 2, a dynamical scheme independent cutoff scale at renormalization  $\mu = \Lambda_{eff}$  for the physical observable in the RSS dependent expression on the left hand side. As it is dependent on dynamical scales, this RSS independent cutoff scale is indeed distinct from typical renormalization coupling and mass cutoffs, which are scheme dependent (see [14] for analysis on RS dependence of the strong coupling constant and its cutoff  $\Lambda_{QCD}$ ). Equation 3 denotes another RSS independence requirement whereby matching is done at a PS  $Q^*$  at  $r$ -loop, via an effective RSS independent dynamical coupling and mass depending on PS, which are denoted as  $a_{eff}$  and  $m_{eff}$  respectively in both equations; though in the RSS dependent expression these are computed at a truncated order,  $n$ , thereby denoting the effective nature of these RSS independence conditions. As such, eq. 2 and 3 allow for a practical realization of eq. 1 at finite order of perturbation theory for our proposed implementation of the POEM approach, and imply RSS independence in the EPO to hold both at the matching scale  $Q^*$ , and for dynamical degrees of freedom,  $Q$  and  $m$ , which can be explicitly stated as:

$$\frac{\partial O_{eff}^{(tree)}}{\partial \mu} = \frac{\partial O_{eff}^{(tree)}}{\partial c_i} = \frac{\partial O_{eff}^{(r)}}{\partial \mu} = \frac{\partial O_{eff}^{(r)}}{\partial c_i} = 0 \quad (4)$$

where  $\mu$  is the renormalization scale, while  $c_i$  are renormalization scheme dependent RG coefficients.

Here, in equations 2 and 3, the  $k$  subscript denotes a particular scheme for a truncated observable  $O$  at a loop order  $n$  (for example the  $\overline{MS}$  scheme) which is matched at the tree and  $r$ -loop level with an effective version of the fully PS dependent version  $O_{eff}$ , whereby  $r$  is to be chosen at the highest loop-order at which scheme independence holds. The rationale behind this requirement is that this allows matching at a perturbative order at which this scheme independence requirement holds fully at the highest order of perturbation theory, hence results derived are then more accurate than at lower orders where scheme independence may still hold. Typically, for most observables,  $r$  is either at one-loop or two-loops of perturbation theory. In the case study below, we study a QCD observable which is scheme independent at 2-loops, however, we do compute the results at both  $r = 1$  and  $r = 2$  loop orders and illustrate as to why matching at the latter order yields slightly different and better results.

Since the POEM is based on EFT and focused directly on the observable itself, it must also be noted that in this approach, the PS or matching scale  $Q^*$  is not arbitrary, but is rather based on the physical observable and the relevant physical process under study. The

value of the coupling and masses must be measured at this scale, and the range of applicability cannot exceed beyond the relevant physical degrees of freedom or go beyond perturbative cutoff of the theory. Moreover, the resummation that happens due to POEM is, in analogy, “integrates out” the unphysical RSS dependencies from the observable to an independent EPO. As such, with POEM we perform an effective “dynamical” renormalization of the RSS dependent physical observable via matching (and resulting resummation) which renders it completely dynamical, with the EPO tied to the physical relevance of the scale of matching with applicability to the underlying theory while preserving physical degrees of freedom. Typically, EFT techniques are applied at the level of operators instead of observables, and deal with physical degrees of freedom, and dealing with RSS dependence as auxiliary variables. Therefore, broadly speaking, our approach via the POEM is to bridge conventional renormalization techniques with EFT techniques, with a first focus on observables in this work. We demonstrate these statements concretely with the case of the cross section, for  $R_{e^+e^-}$ ,  $e^+e^- \rightarrow$  hadrons, where the matching scale  $Q^*$  at the  $Z$ -pole mass is chosen with five-active quarks, not only because the strong coupling constant is measured at this PS in the  $\overline{MS}$  scheme, but also that the center of mass energies,  $Q$ , under consideration are well above the b-quark threshold mass and are within the limits of applicability of the EFT. Hence, the RSS independent EPO derived will be limited in terms of its range of applicability given the reasons for the choice of  $Q^*$ , and with QCD as an underlying EFT. As another example, if a physical observable is related to lower energies is taken much below this threshold, then the matching scale is to be at point at which the strong coupling is measured at a lower scale, say the tau mass, and for 3 to 4 active quark flavors.

We also remark that our approach is distinct from the Effective Charge (EC) approach [3], as in this method there is matching done at one-loop and at a scale in which renormalization logarithms are set to zero, which only allows renormalization scheme independence, while renormalization scale dependence remains in resulting expressions. Via POEM, and for deriving a resultant EPO, the matching at the tree and at  $r$ -loop order at a PS, we overcome these limitations, thereby achieving RSS independence simultaneously. In the case of the QCD cross section ratio,  $R_{e^+e^-}$  studied here, we have an observable which is scheme-independent up to two-loops, and we find that matching at this higher loop yields better results as compared to 1-loop matching, hence, we demonstrate the advantages of matching at a higher loop order with RSS independent results unlike the EC approach which are scale dependent, and are fixed at one-loop order for all observables. Moreover, POEM based results are RSS independent, hence, there are no “commensurate scale relations” [15] as the

observable is dynamical with respect to physical scales, and hence no relative scales exist for the EPO. After achieving RSS invariance, there are no issue of renormalons [16, 17] encountered as well.

### III. ATTAINING EFFECTIVE RSS INDEPENDENCE: $R_{e^+e^-}$ MATCHING AT TREE AND ONE-LOOP ORDER

The form of the cross section  $R_{e^+e^-}$  is given by  $3(\sum_i q_i^2)(1 + R)$  where  $R$  at  $n$ -loop order has a perturbative contribution of order  $a^{n+1}$  in

$$R = R_{\text{pert}} = \sum_{n=0}^{\infty} r_n a^{n+1} = \sum_{n=0}^{\infty} \sum_{m=0}^n T_{n,m} L^m a^{n+1} \quad (5)$$

with  $L \equiv b \ln\left(\frac{\mu}{Q}\right)$ ,  $Q^2$  being the center of mass momentum squared.

The explicit dependence of  $R$  on the renormalization scale parameter  $\mu$  is compensated for by implicit dependence of the “running coupling”  $a(\mu^2)$  on  $\mu$ ,

$$\mu^2 \frac{\partial a}{\partial \mu^2} = \beta(a) = -ba^2(1 + ca + c_2 a^2 + \dots) \quad (6)$$

where  $a \equiv \alpha_s(\mu^2)/\pi$  while  $\alpha_s$  is the QCD strong coupling constant.

For five active flavors of quark, we have,

$$\frac{3}{11} R_{e^+e^-} = 1 + R_{\text{pert}}. \quad (7)$$

The choice of the number of flavors is based on the center of mass energies  $Q$  (for which the EPO will be derived) which namely for those above the b-quark mass.

In the  $\overline{MS}$  renormalization scheme [18] we have,

$$b = 23/12, \quad c = 29/23, \quad c_2 = 9769/6624, \quad (8)$$

$$c_3 = 9.835917120$$

where the values of  $b$  and  $c$  are the same in any mass-independent renormalization scheme while the values of  $c_2$  and  $c_3$  in eqs. (8) are particular to the  $\overline{MS}$  scheme. Furthermore, we find in ref. [19,20] that in the  $\overline{MS}$  scheme

$$T_{0,0} = 1, \quad T_{1,0} = 1.4097, \quad T_{1,1} = 2$$

$$T_{2,0} = -12.76709, \quad T_{2,1} = 8.160539, \quad T_{2,2} = 4$$

$$T_{3,0} = -80.0075, \quad T_{3,1} = -66.54317, \quad T_{3,2} = 29.525095$$

$$T_{3,3} = 8 \quad (9)$$

At the  $Z$ -pole mass ( $M_Z = 91.1876 \text{ GeV}$ ), we have [21] in the  $\overline{MS}$  scheme, (with  $a(\mu)$  defined by eq. (6)),

$$\pi a_{\overline{MS}}(M_Z) = 0.1179 \pm 0.001. \quad (10)$$

Utilizing eq. 2 and 3, we therefore find the following POEM based relations:

$$\begin{aligned} & \frac{3}{11} R_{e^+e^-}^{\overline{MS}}(a_{eff}(\Lambda_{eff}), \Lambda_{eff}) \\ &= 1 + R_{\text{pert}}(a_{eff}(\Lambda_{eff}), \Lambda_{eff}) = \frac{3}{11} R_{e^+e^-}^{eff} = 1 \quad (11) \end{aligned}$$

$$\begin{aligned} & \frac{3}{11} R_{e^+e^-}^{\overline{MS}}(Q^*, Q) = 1 + R_{\text{pert}}(Q^*, Q) \\ &= \frac{3}{11} R_{e^+e^-}^{eff}(Q^*, Q) = 1 + a_{eff}^{1L}(Q^*, Q) \quad (12) \end{aligned}$$

From equation 11, at 3-loops in  $\overline{MS}$ , we find  $a_{eff}^{3L}(\Lambda_{eff}) = 0.3404702$ , while for 4-loops, we obtain  $a_{eff}^{4L}(\Lambda_{eff}) = 0.2094525$ . Since at the tree level, no quantum corrections are present, hence we have, through dimensional transformation,  $a_{eff}(\Lambda_{eff}) = \Lambda_{eff}$  at this level of matching. Hence, for the EPO,  $R_{e^+e^-}^{eff}$ , we obtain scheme independent cutoffs,  $\Lambda_{eff}^{3L} = 340.47 \text{ MeV}$  and  $\Lambda_{eff}^{4L} = 209.45 \text{ MeV}$  at 3- and 4-loops respectively.

With equation 12, which is matching done at the one-loop level, we choose the matching point  $Q^* = M_Z = 91.1876 \text{ GeV}$ , for which the central value of the strong coupling  $a^{MS} = \frac{0.1179}{\pi}$  from eq. 9, which we substitute into eq. 12 to derive  $a_{eff}^{1L}(M_Z, Q)$  as follows:

$$\begin{aligned} a_{eff}^{1L}(M_Z, Q) &= 0.03868 + 0.005973 \ln(M_Z/Q) \\ &+ 0.0009918 (\ln(M_Z/Q))^2 \\ &+ 0.0001117 (\ln(M_Z/Q))^3 \quad (13) \end{aligned}$$

The choice of  $Q^* = M_Z$  and the number of quarks,  $n_f = 5$ , is predicated on deriving the EPO for center of mass energies  $Q$  above the b-quark mass (and also below the t-quark threshold mass).

Since for the EPO we absorb higher order-loop contributions from a RSS dependent scheme at one-loop order for RSS independence, hence,  $R_{e^+e^-}^{eff}(Q)$  is expressed as,

$$\frac{3}{11} R_{e^+e^-}^{eff}(Q) = 1 + a_{eff}^{1L}(Q) \quad (14)$$

where

$$a_{eff}^{1L}(Q) = \frac{a_{eff}^{1L}(Q^*, Q)}{1 - ba_{eff}^{1L}(Q^*, Q) \ln\left(\frac{Q^*}{Q}\right)} \quad (15)$$

is a one-loop beta-function solution in the dynamical EPO with an initial value set at  $a_{eff}^{1L}(\mu_0 = Q^*) = a_{eff}^{1L}(Q^*, Q)$ , which in our case is given by eq. 13.

Since we are at the one-loop matching, we find the one-loop solution to  $a_{eff}(Q)$  which leads to the prediction of  $\frac{3}{11} R_{e^+e^-}^{eff} = 1.056961_{-0.0007}^{+0.0007}$  from eq. 12, at  $Q = 31.6 \text{ GeV}$ , which is in excellent agreement with the experimental value of  $\frac{3}{11} R_{e^+e^-}^{exp} = 1.0527_{-0.005}^{+0.005}$  [22]. We plot the  $Q$  dependence of  $R^{eff}$  at 1-loop matched with 3- and 4-Loop  $\overline{MS}$  in Figure 1, where the both results are similar. At the same time perturbative QCD (pQCD) predictions are found for renormalization scales,  $\mu = M_Z/2$ ,  $\mu = M_Z$  and  $\mu = 2M_Z$  respectively, which are all underestimates as compared to POEM based EPO results, and also indicate high uncertainty across this range due to RSS dependence.




Figure 1. The  $Q$  dependence of Renormalization Scale and Scheme (RSS) independent  $R^{eff}$  at 1-loop matched with 3- and 4-Loop  $\overline{MS}$  and compared to 4-Loop Perturbative QCD in  $\overline{MS}$  scheme for  $R_{e^+e^-}$  referenced at renormalization scales,  $\mu = M_Z/2$ ,  $\mu = M_Z$  and  $\mu = 2M_Z$

#### IV. TWO-LOOP MATCHING

$R_{e^+e^-}$  is scheme-independent at 2-loops, hence this would be applying POEM at the maximum loop order in this case. Utilizing eq. 4 at two-loop level, we find the following scale and scheme independent POEM based relation:

$$\begin{aligned} & \frac{3}{11} R_{e^+e^-}^{\overline{MS}}(Q^*, Q) = 1 + R_{\text{pert}}(Q^*, Q) \\ &= \frac{3}{11} R_{e^+e^-}^{eff}(Q^*, Q) = 1 + a_{eff}^{2L}(Q^*, Q) + T_{1,0} a_{eff}^{2L}(Q^*, Q)^2 \quad (16) \end{aligned}$$

As before, we choose  $Q^* = M_Z$ , and find  $a_{eff}(M_Z, Q)$

as follows:

$$a_{eff}^{2L}(M_Z, Q) = -0.35469 \pm 7.0934 [3.04528 \times 10^{-3} + 1.57512 \times 10^{-6} \ln^3 \left( \frac{M_Z}{Q} \right) + 1.39819 \times 10^{-5} \ln^2 \left( \frac{M_Z}{Q} \right) + 8.41960 \times 10^{-5} \ln \left( \frac{M_Z}{Q} \right)]^{1/2} \quad (17)$$

Since in the EPO, namely,  $R_{e^+e^-}^{eff}$  we absorb higher order-loop contributions from a RSS dependent scheme at  $r = 2$  (two-loop orders) for RSS independence, hence,  $R_{e^+e^-}^{eff}(Q)$  is expressed as

$$\frac{3}{11} R_{e^+e^-}^{eff}(Q) = 1 + a_{eff}^{2L}(Q) + T_{1,0} a_{eff}^{2L}(Q)^2 \quad (18)$$

where

$$a_{eff}^{2L}(Q) = \frac{1}{cW \left[ -\exp \left( \frac{f}{bc} \right) + 1 \right]} \quad (19)$$

where

$$f = \frac{ba_{eff}^{2L}(Q, Q^*) \left[ b \ln \left( \frac{P^2}{Q^2} \right) + c \ln(h) - c \right] - 1}{ca_{eff}^{2L}(Q, Q^*)} \quad (20)$$

and

$$h = \frac{b[ca_{eff}^{2L}(Q, Q^*) + 1]}{a_{eff}^{2L}(Q, Q^*)}. \quad (21)$$

Eqs. 19-21 represent an exact closed form solution of the two-loop beta-function for the physical EPO, which is expressed as a Lambert-W function. We utilize the initial value set at  $a_{eff}(\mu_0 = Q^*) = a_{eff}(Q^*, Q)$ , which in our case is given by the positive root of eq. 17. At the two-loop EPO matching, we find that the prediction is nearly the same as the one-loop EPO, which is  $\frac{3}{11} R_{e^+e^-}^{eff} = 1.056137_{-0.0007}^{+0.0007}$  from eq. 18 at  $Q = 31.6 GeV$ , which in excellent agreement with the experimental value of  $\frac{3}{11} R_{e^+e^-}^{exp} = 1.0527_{-0.005}^{+0.005}$  [22] which is better than our previously derived 1-loop matching EPO result.

We plot the  $Q$  dependence of  $R^{eff}$  at 2-loop matching with 3- and 4-Loop  $\overline{MS}$  in Figure 2 and find the same trends as 1-loop matching, with pQCD predictions lower than EPO results, though these are slightly lower than the 1-loop EPO results. We also show the behavior of RSS independent coupling  $a^{eff}$  at 1- and 2-Loop orders with respect to  $Q$ , in Figure 3, which depict asymptotic freedom as is also seen in the EPO behavior in Figures 1 and 2. We note that the  $a^{eff}$  is

lower for 2-loops as compared to 1-loop matching as depicted in Figure 3, which results in a better prediction of  $\frac{3}{11} R_{e^+e^-}^{eff}$ . In Figure 4, we show that at 2-loops  $r = 2$  which is the highest loop order at which  $\frac{3}{11} R_{e^+e^-}$  is scheme independent, these EPO predictions are slightly lower (and subsequently better) than those found at 1-loop. Based on these findings, we therefore surmise matching at the highest available order at which the observable is scheme-independent to yield best results.

In comparison with our findings, Akrami and Mirjalili [25] present perturbative QCD, RG summation and RS invariants and CORGI approaches estimates of  $R$  at  $Q = 31.6 GeV$  to be  $1.04617_{-0.0006}^{+0.0006}$ ,  $1.04711_{-0.0005}^{+0.0003}$  and  $1.04615_{-0.0008}^{+0.0015}$  at 4-Loops respectively, which are all underestimates as compared to the result derived by POEM. This pattern holds for other experimental measured values at  $Q = 42.5 GeV$  and  $Q = 52.5 GeV$ , which are  $1.0554 \pm 0.2$  [23] and  $1.0745 \pm 0.11$  [24] respectively. With POEM we find  $1.049351_{-0.0005}^{+0.0005}$  and  $1.045737_{-0.0005}^{+0.0005}$  using the 2-Loop EPO matching. These predictions values are higher in value and more accurate than those reported for perturbative QCD, RG summation and RS invariants and CORGI approaches in [25]. However, our predictions are not close to the experimental values as they are based only on photon interactions, and a higher accuracy is expected when electroweak contributions arising from electron-positron annihilation to  $Z$  boson are taken into account. We will address this in a future work, which will also additionally address other processes [25].




Figure 2. The  $Q$  dependence of Renormalization Scale and Scheme (RSS) independent  $R^{eff}$  at 2-loop matched with 3- and 4-Loop  $\overline{MS}$  and compared to 4-Loop perturbative QCD in  $\overline{MS}$  scheme for  $R_{e^+e^-}$  referenced at renormalization scales,  $\mu = M_Z/2$ ,  $\mu = M_Z$  and  $\mu = 2M_Z$  (dashed lines)




Figure 3. The  $Q$  dependence of Renormalization Scale and Scheme (RSS) independent coupling  $a_e^{eff}$  at 1- and 2-Loop orders.




Figure 4. The  $Q$  dependence of Renormalization Scale and Scheme (RSS) independent coupling  $\frac{3}{11}R_{e^+e^-}^{eff}$  derived via matching EPO at 1- and 2-Loop orders.

## V. CONCLUSIONS

In this work, we have introduced a new approach for achieving RSS independence via a principle termed as POEM, which is based on equivalence of physical observables across both scale and scheme dependencies in their perturbative content under certain limits. Inspired by EFT techniques, which involve matching at a physical scale, this approach provides results termed as EPO, which contain only physical scales at fixed order of perturbation theory. In this work, we demonstrate that POEM provides excellent results for the QCD cross section ratio  $R$  and agreement with experiment is better than other comparable estimates based on perturbative QCD, RG summation and RS invariants and CORGI approaches [25]. POEM is distinct from present approaches in its conceptualization, but we do intend to find potential connections and improvements with other approaches including  $\text{PMC}_\infty$  [26], which is based on the PMC [12] in an upcoming work [27]. In this work, we have only incorporated photon interactions, and hence better accuracy for higher values of  $Q$  are expected once contributions from electron-positron annihilation to the Z-boson are also taken into account. We aim to study and explicate POEM's potential in these areas in upcoming studies whereby linkages between conventional renormalization and EFT techniques are further explicated, and as a follow up will also address other electroweak processes using POEM including Standard Model Higgs decays and cross-sections [28]. Based on a new conceptualization and achieving better results for the cross section  $R$ , we contend that POEM is applicable widely to achieve RSS Independence in physical observables across the high energy physics domain including both the Standard Model (SM), Beyond the SM physics, along with areas such as cosmology, statistical and condensed matter physics where RSS dependencies are regularly encountered.

## VI. ACKNOWLEDGMENTS

I thank Gerry McKeon and Thomas G. Steele for helpful discussions and comments on the initial drafts of this work.

- 
- [1] P. M. Stevenson, *Phys. Rev.* **D23**, 2916 (1981); *Phys. Rev.* **D 33**, 3130 (1986).  
 [2] D. G. C. McKeon, *Can. J. Phys.* **61**, 564 (1983); **59**, 1327 (1981).  
 [3] G. Grunberg, *Phys. Lett.* **B 95**, 70 (1980); *Phys. Lett.* **B 110**, 501 (1982).

- [4] S. J. Brodsky, G. P. Lepage and P. B. Mackenzie, *Phys. Rev.* **D 28** (1983) 228. *Phys. Rev.* **D 29**, 2315 (1984); *Phys. Rev.* **D 40**, 680 (1989).  
 [5] M. R. Ahmady, F. A. Chishtie, V. Elias, A. H. Fariborz, N. Fattahi, D. G. C. McKeon, T. N. Sherry and T. G. Steele, *Phys. Rev.* **D66**, 014010 (2002).  
 [6] D. G. C. McKeon, *Phys. Rev.* **D92**, 045031 (2015).

- [7] F. A. Chishtie, D. G. C. McKeon and T.N. Sherry, *Phys. Rev.* **D94**, 054031 (2016).
- [8] F. A. Chishtie and D. G. C. McKeon, *Can. J. Phys.* **96**, 233 (2017).
- [9] F. A. Chishtie and D. G. C. McKeon, *Phys. Rev.* **D95**, 116013 (2017).
- [10] C. J. Maxwell, *Nucl. Phys. B Proc. Suppl.* **86**, 74 (2000). P. Brooks and C. J. Maxwell, *Nucl. Phys.* **B 780**, 76 (2007).
- [11] C. J. Maxwell and A. Mirjalili, *Nucl. Phys.* **B 577**, 209 (2000).
- [12] S. J. Brodsky and X. G. Wu, *Phys. Rev.* **D 85**, 034038 (2012) S. J. Brodsky and L. Di Giustino, *Phys. Rev.* **D 86**, 085026 (2012) S. J. Brodsky and X. G. Wu, *Phys. Rev. Lett.* **109**, 042002 (2012) M. Mojaza, S. J. Brodsky and X. G. Wu, *Phys. Rev. Lett.* **110**, 192001 (2013) S. J. Brodsky, M. Mojaza and X. G. Wu, *Phys. Rev.* **D 89**, 014027 (2014) Y. Ma, S. Q. Wang, H. B. Fu, H. H. Ma, S. J. Brodsky and M. Mojaza, *Rep. Phys.* **78**, 126201 (2015).
- [13] A. L. Kataev and S. V. Mikhailov, *Phys. Rev.* **D 91**, 014007 (2015).
- [14] W. Celmaster and R. J. Gonsalves, *Phys. Rev.* **D 20**, 1420 (1979).
- [15] S. J. Brodsky and H. J. Lu, *Phys. Rev.* **D 51**, 3652 (1995) S. J. Brodsky, G. T. Gabadadze, A. L. Kataev and H. J. Lu, *Phys. Lett.* **B 372**, 133 (1996)
- [16] G. 't Hooft, *The Whys of Subnuclear Physics*, Erice 1977, ed. A. Zichichi, Plenum Press N.Y. (1979).
- [17] M. Beneke, *Phys. Rep.* **37**, 1 (1999).
- [18] T. van Ritbergen, J. A. M. Vermaseren and S.A. Larin, *Phys. Lett.* **B400**, 379 (1997).
- [19] K. G. Chetyrkin, J. H. Kuhn, and A. Kwiatkowski, *Phys. Rep.* **277**, 189 (1996).
- [20] P. A. Baikov, K. G. Chetyrkin, J. H. Kuhn, and J. Rittinger, *J. High Energy Phys.* **07** (2012) 017.
- [21] M. Tanabashi et al. (Particle Data Group), *Phys. Rev.* **D 98**, 030001 (2018) and 2019 update.
- [22] P. Achard et al. (L3 collaboration) *Eur. Phys.J.* **C 47**, 1 (2006).
- [23] B. Adeva et al. (Mark-J Collaboration) *Phys. Rev.* **D 34** 681 (1986).
- [24] K. Abe and et al. (VENUS Collaboration) *Phys. Lett.* **B 234**, 382 (1990).
- [25] M. Akrami and A. Mirjalili, *Phys. Rev.* **D101**, 034007 (2020).
- [26] L. Di Giustino, S. J. Brodsky, S. Q. Wang and X. G. Wu, arXiv:2002.01789 [hep-ph].
- [27] Farrukh A. Chishtie, in preparation.
- [28] Farrukh A. Chishtie, in preparation.