

Dynamics of planar vector fields near a non-smooth equilibrium*

Tao Li, Xingwu Chen[†]

Department of Mathematics, Sichuan University,
Chengdu, Sichuan 610064, P. R. China

Abstract

In this paper we contribute to qualitative and geometric analysis of planar piecewise smooth vector fields, which consist of two smooth vector fields separated by the straight line $y = 0$ and sharing the origin as a non-degenerate equilibrium. In the sense of Σ -equivalence, we provide a sufficient condition for linearization and give phase portraits and normal forms for these linearizable vector fields. This condition is hard to be weakened because there exist vector fields which are not linearizable when this condition is not satisfied. Regarding perturbations, a necessary and sufficient condition for local Σ -structural stability is established when the origin is still an equilibrium of both smooth vector fields under perturbations. In the opposition to this case, we prove that for any piecewise smooth vector field studied in this paper there is a limit cycle bifurcating from the origin, and there are some piecewise smooth vector fields such that for any positive integer m there is a perturbation having exactly m limit cycles bifurcating from the origin. Here m maybe infinity.

2010 MSC: 34A36, 34C41, 37G05, 37G15.

Keywords: limit cycle bifurcation, linearization, non-smooth equilibrium, normal form, structural stability.

1 Introduction and statement of the main results

Let $\mathcal{U} \subset \mathbb{R}^2$ be a bounded open set containing the origin O , \mathfrak{X} be the set of all C^1 vector fields defined on \mathcal{U} and endowed with the C^1 -topology. We consider the piecewise smooth vector field

$$Z(x, y) = \begin{cases} X(x, y) = (X_1(x, y), X_2(x, y)) & \text{if } (x, y) \in \Sigma^+, \\ Y(x, y) = (Y_1(x, y), Y_2(x, y)) & \text{if } (x, y) \in \Sigma^-, \end{cases} \quad (1.1)$$

where $X, Y \in \mathfrak{X}$ and

$$\Sigma^+ = \{(x, y) \in \mathcal{U} : y > 0\} \quad \Sigma^- = \{(x, y) \in \mathcal{U} : y < 0\}.$$

Define Ω as the set of all $Z(x, y)$ satisfying (1.1) and endowed with the product topology. In past two decades, many researchers shift their interest to the study of piecewise smooth vector fields, because such vector fields are ubiquitous in mechanical engineering [4, 8], feedback control systems [1, 14], biological systems [27, 28], electrical circuits [1], etc.

Notice that the piecewise smooth vector field (1.1) is not defined on $\Sigma = \{(x, y) \in \mathcal{U} : y = 0\}$, called *discontinuity line* or *switching line*. Denote the vector field on Σ by Z_Σ , which is usually defined by the

*Supported by NSFC #11871355 and CSC #201906240094.

[†]Author to whom any correspondence should be addressed. Email address: xingwu.chen@hotmail.com (X. Chen).

so-called Filippov convention [13], see Section 2 for a review. Here Z_Σ is naturally defined as X or Y if $X(x, y) \equiv Y(x, y)$ for all $(x, y) \in \Sigma$. The vector field (1.1), together with Z_Σ , are called a *Filippov vector field*. In whole paper, speaking of the vector field $Z \in \Omega$, it always means that $Z = Z_\Sigma$ on Σ . A point at which $Z \in \Omega$ vanishes is said to be an *equilibrium* or *singular point*. Hence, an equilibrium of Z is an equilibrium of either X in Σ^+ or Y in Σ^- or Z_Σ in Σ . Throughout this paper, we call it a *smooth equilibrium* for the first two cases and a *non-smooth equilibrium* for the last case.

Regarding the local dynamics of $Z = (X, Y) \in \Omega$ near a smooth equilibrium, the investigation can be reduced to the local dynamics of the smooth vector field X or Y near this equilibrium and, with the efforts of many researchers, a large number of mature theories and methods have been established (see e.g., [17, 23, 31]). Therefore, we focus on the local dynamics for non-smooth equilibria, which is more difficult than the smooth case because most theories and methods for smooth vector fields are no longer valid for non-smooth ones. Although that, in recent twenty years some excellent results about limit cycle bifurcation, normal form and structural stability were given in textbooks [1, 13] and journal papers [5, 6, 11, 15, 16, 18, 21, 24, 32]. Let $\Omega_0 \subset \Omega$ be the set of all piecewise smooth vector fields satisfying

$$X(0, 0) = Y(0, 0) = (0, 0), \quad \det A^+ \det A^- \neq 0 \quad (1.2)$$

and

$$X_{2x}(0, 0)Y_{2x}(0, 0) > 0, \quad (1.3)$$

where A^+ (resp. A^-) is the Jacobian matrix of X (resp. Y) at O and X_{2x}, Y_{2x} denote the derivatives of X_2, Y_2 with respect to x , respectively. (1.2) means that equilibrium O is non-degenerate for both X and Y , (1.3) means that there exists a hollow neighborhood of O , in which there are no sliding points (see Section 2).

In this paper we study the local dynamics of vector field $Z = (X, Y) \in \Omega_0$ near O , which is a non-smooth equilibrium of Z , i.e., $Z(0, 0) = Z_\Sigma(0, 0) = (0, 0)$. Our first goal is to study the *local Σ -equivalence* between $Z = (X, Y) \in \Omega_0$ and its linear part

$$Z_L(x, y) = \begin{cases} X_L(x, y) = A^+(x, y)^\top & \text{if } (x, y) \in \Sigma^+, \\ Y_L(x, y) = A^-(x, y)^\top & \text{if } (x, y) \in \Sigma^- \end{cases} \quad (1.4)$$

near O . Roughly speaking, the local Σ -equivalence is just the local topological equivalence preserving the switching line Σ . A precise definition of local Σ -equivalence is stated in Section 2. A nonlocal definition of Σ -equivalence, e.g., not in a neighborhood of equilibrium but in the whole domain of definition, was given in [16, Definition 2.20] and [1, Definition 2.30]. One of motivations for this goal comes from the work [10]. In [10, Theorem 2.2], 19 different types of normal forms for $Z \in \Omega$ with (1.2) were obtained by using a continuous piecewise linear change of variables. We notice that in these normal forms the linear parts are normalized but the nonlinear parts are not normalized. So, it is unknown that whether these nonlinear parts can be eliminated after normalization. Another motivation is from smooth vector fields. A smooth vector field is locally topologically equivalent to its linear part near an equilibrium if all eigenvalues of the Jacobian matrix at this equilibrium have nonzero real part (see, e.g., [19] and [31, Theorem 4.7]). Hence, it is a natural question to find conditions such that $Z \in \Omega_0$ is locally Σ -equivalent near O to its linear part Z_L given in (1.4).




Figure 1: Local phase portraits of $Z \in \Omega_1$ near the origin.

Let λ_1^\pm and λ_2^\pm be the eigenvalues of A^\pm , and

$$\Omega_1 = \{Z \in \Omega_0 : \lambda_1^+ \neq \lambda_2^+, \lambda_1^- \neq \lambda_2^-, \ell \neq 0\}, \quad (1.5)$$

where

$$\ell = \begin{cases} \frac{\operatorname{Re}\lambda_1^+}{|\operatorname{Im}\lambda_1^+|} + \frac{\operatorname{Re}\lambda_1^-}{|\operatorname{Im}\lambda_1^-|} & \text{if } \operatorname{Im}\lambda_1^+ \operatorname{Im}\lambda_1^- \neq 0, \\ 1 & \text{if } \operatorname{Im}\lambda_1^+ \operatorname{Im}\lambda_1^- = 0, \end{cases} \quad (1.6)$$

Re and Im denote the real and imaginary part of eigenvalues respectively. We have the first theorem as follows.

Theorem 1.1. *Every $Z \in \Omega_1$ is locally Σ -equivalent to its corresponding piecewise linear vector field Z_L of form (1.4) near the origin. Moreover, the local phase portrait of Z near the origin is one of the 11 phase portraits presented in Figure 1 in the sense of Σ -equivalence.*

Theorem 1.1 is proved in Section 3, where we present a normal form for each one of these 11 kinds of phase portraits shown in Figure 1. We remark that the first part of Theorem 1.1 can be regarded as a generalisation of [31, Theorem 4.7] from smooth vector fields to piecewise smooth vector fields. We clarify some differences between the requirements for eigenvalues in these two theorems as follows. In [31, Theorem 4.7] it is required that all eigenvalues of the Jacobian matrix at a smooth equilibrium have nonzero real part in order that the smooth vector field is topologically equivalent to its linear part near this equilibrium. However, in Theorem 1.1 we require that all eigenvalues of the Jacobian matrixes A^+ and A^- at O , namely the non-smooth equilibrium, satisfy

$$\lambda_1^+ \lambda_2^+ \neq 0, \quad \lambda_1^+ \neq \lambda_2^+, \quad \lambda_1^- \neq \lambda_2^-, \quad \ell \neq 0$$

by the definition of Ω_1 given in (1.5). Comparing the requirements of [31, Theorem 4.7] with our Theorem 1.1, we see that [31, Theorem 4.7] does not allow pure imaginary eigenvalues but Theorem 1.1 allows. On the other hand, by [11, Theorem B] or [18, Theorem 1.2] the condition $\ell \neq 0$ in Theorem 1.1 excludes the case that O is a non-smooth center of the linear part. It is not hard to give an example showing the non-equivalence when O is a non-smooth center of the linear part. Another difference is that [31, Theorem 4.7] allows the Jacobian matrix to have the same eigenvalue, but Theorem 1.1 does not allow this for both Jacobian matrices A^+ and A^- . We give an example to show the non-equivalence when the Jacobian matrix A^+ or A^- has the same eigenvalue in Section 3.

Our second goal is to study the structural stability of $Z \in \Omega_0$ in the sense of Σ -equivalence, i.e., Σ -structural stability as defined in [16, p.1978]. Usually, $Z \in \Omega_0$ is not Σ -structurally stable when the perturbation is inside Ω because O can be destroyed under such a perturbation and the so-called boundary equilibrium bifurcation occurs [24]. Thus the only interest is to consider the Σ -structural stability of $Z \in \Omega_0$ with respect to Ω_0 , i.e., the perturbation is inside Ω_0 . In particular, we focus on the local Σ -structural stability of $Z \in \Omega_0$ near O . Roughly speaking, $Z \in \Omega_0$ is said to be *locally Σ -structurally stable with respect to Ω_0 near O* if any vector field that lies in a sufficiently small neighborhood of Z contained in Ω_0 is locally Σ -equivalent to Z near O .

Theorem 1.2. *$Z \in \Omega_0$ is locally Σ -structurally stable with respect to Ω_0 near the origin if and only if $Z \in \Omega_1$, where Ω_0 is defined above (1.2) and Ω_1 is defined in (1.5).*

Theorem 1.2 is proved in Section 4.

The third goal of this paper is devoted to the study of limit cycle bifurcations, more precisely, identify the existence and number of crossing limit cycles bifurcating from the non-smooth equilibrium O of a piecewise smooth vector field $Z = (X, Y) \in \Omega_0$. Here a limit cycle is said to be a *crossing limit cycle* if it intersects the switching line Σ only at crossing points (see Section 2). Many works about limit cycle bifurcations are done for the case that O is of focus-focus type, i.e., an equilibrium of focus type for both X and Y . See, e.g., [9–12, 22, 25, 32] for the perturbations in Ω_0 and [18, 30] for the perturbations in Ω . Such bifurcation is analogous to the Hopf bifurcation of smooth vector fields. Then a natural question is whether limit cycles can bifurcate from O for other cases, for instance O is of focus-saddle type, focus-node type, etc. Since bifurcations usually depend on the type of local phase portraits of the unperturbed systems and there exist many kinds of possibilities as obtained in Theorem 1.1, in this paper we do not establish the bifurcation diagrams one by one but give some universal results on the limit cycle bifurcations for all unperturbed vector fields in Ω_0 .

Theorem 1.3. *For Ω_0 defined above (1.2) and its subset Ω_1 defined in (1.5), the following statements hold.*

- (1) *For any $Z \in \Omega_0$ and any small neighborhood $N \subset \Omega$ of Z , there exists a vector field in N having a crossing limit cycle bifurcating from the non-smooth equilibrium O of Z .*
- (2) *There exists a $Z_0 \in \Omega_1$ (resp. $\Omega_0 \setminus \Omega_1$) such that, for any $m \in \mathbb{N}^+ \cup \{\infty\}$ and any small neighborhood $N \subset \Omega$ of Z_0 , there exists a vector field in N having exactly m hyperbolic crossing limit cycles bifurcating from the non-smooth equilibrium O of Z_0 .*

Theorem 1.3 is proved in Section 5. Note that even though our main motivation is to consider the case of piecewise smooth vector fields, the set Ω_0 also includes the smooth vector fields with $X \equiv Y$

having O as a non-degenerate equilibrium. Thus it follows from the statement (1) of Theorem 1.3 that limit cycles can bifurcate from a rough focus, saddle or node of smooth vector fields under non-smooth perturbations. This is impossible under smooth perturbations.

This paper is organized as follows. In Section 2 we shortly recall basic notions and results on piecewise smooth vector fields. In Section 3 we give the proof of Theorem 1.1, and an example showing that the vector field in Ω_0 might not be locally Σ -equivalent to its linear part near the origin if the Jacobian matrix A^+ or A^- has the same eigenvalue. The proofs of Theorems 1.2 and 1.3 are given in Sections 4 and 5, respectively.

2 Preliminaries

For the sake of completeness, in this section we shortly review some basic notions and results on piecewise smooth vector fields, especially Filippov vector fields. Section 2.1 contains the definitions of vector field Z_Σ on Σ and all kinds of singularities. Moreover, the local Σ -equivalence is also clarified in Section 2.1. In Section 2.2 we state the pseudo-Hopf bifurcation for a special class of piecewise smooth vector fields in order to prove our results conveniently.

2.1 Notions on piecewise smooth vector fields

Consider the piecewise smooth vector field $Z \in \Omega$ given in (1.1). First we clarify the definition of vector field Z_Σ on Σ by the Filippov convention [13]. To do this, Σ is divided into the *crossing set*

$$\Sigma^c = \{(x, y) \in \Sigma : X_2(x, y) \cdot Y_2(x, y) > 0\},$$

and the *sliding set*

$$\Sigma^s = \{(x, y) \in \Sigma : X_2(x, y) \cdot Y_2(x, y) \leq 0\},$$

as in [13, 24]. The points in Σ^c and Σ^s are called *crossing points* and *sliding points* respectively. For $(x, y) \in \Sigma^c$, X and Y are both transversal to Σ and their normal components have the same sign, so that the orbit passing through (x, y) crosses Σ at (x, y) and it is a continuous, but non-smooth curve. This means that we can define Z_Σ at (x, y) as any one of X and Y . For concreteness, in this paper we specify

$$Z_\Sigma(x, y) = \begin{cases} Y(x, y) & \text{if } (x, y) \in \Sigma^c, \ X_2(x, y) < 0, \\ X(x, y) & \text{if } (x, y) \in \Sigma^c, \ X_2(x, y) > 0. \end{cases}$$

For $(x, y) \in \Sigma^s$, either the normal components of X and Y to Σ have the opposite sign or at least one of them vanishes. In this case Z_Σ is defined such that it is tangent to Σ^s . Particularly, if $Y_2(x, y) \neq X_2(x, y)$,

$$Z_\Sigma(x, y) = \left(\frac{Y_2(x, y)X_1(x, y) - X_2(x, y)Y_1(x, y)}{Y_2(x, y) - X_2(x, y)}, 0 \right)$$

by [13, 24], while if $Y_2(x, y) = X_2(x, y) = 0$, namely (x, y) is a *singular sliding point* (see [24]), we always assume $Z_\Sigma(x, y) = (0, 0)$ in this paper. Sometimes, Z_Σ restricted on Σ^s , denoted by Z^s , is called the *sliding*

vector field of Z and the corresponding equilibria are said to be *pseudoequilibria*. Having the definition of Z_Σ , the flow of Z can be obtained by concatenating the flows of X , Y and Z_Σ as stated in [24].

In the switching line Σ , the boundary $\partial\Sigma^s$ of Σ^s plays an important role in the dynamical analysis of piecewise smooth vector fields. Let $q \in \partial\Sigma^s$. If $X_2(q) = 0, X(q) \neq 0$ (resp. $Y_2(q) = 0, Y(q) \neq 0$), then q is called a *tangency point* of X (resp. Y), see [24]. In addition, a tangency point q of X is called a *fold point* if $X_1(q)X_{2x}(q) \neq 0$ and it is said to be *visible* (resp. *invisible*) when $X_1(q)X_{2x}(q) > 0$ (resp. $X_1(q)X_{2x}(q) < 0$). The above notions can be similarly defined for Y . If q is a fold point of both X and Y , we call it a *fold-fold point* of Z , which can be divided into visible-visible, invisible-invisible and visible-invisible types. If $X(q) = 0$ (resp. $Y(q) = 0$), q is called a *boundary equilibrium* of X (resp. Y). Clearly, a boundary equilibrium must be a pseudoequilibrium.

Regarding piecewise smooth vector fields, there are two types of equivalences, i.e., topological equivalence and Σ -equivalence. We adopt the latter in this paper as it was indicated in Section 1, see [16, Definition 2.20] and [1, Definition 2.30] for the definition of Σ -equivalence. Since we deal with the local dynamics of $Z \in \Omega_0$ near the origin, namely the non-smooth equilibrium, we can localize the definition of the Σ -equivalence as follows.

Definition 2.1. Consider two piecewise smooth vector fields Z_1 and Z_2 in Ω_0 . We say that Z_1 and Z_2 are locally Σ -equivalent near the origin if

- (1) Z_1 and Z_2 are locally topologically equivalent near the origin, i.e., there exist two neighborhoods U and V of the origin, and a homeomorphism $H : U \rightarrow V$ such that H maps the orbits of Z_1 in U onto the orbits of Z_2 in V , preserving the direction of time; and
- (2) the homeomorphism H sends $\Sigma \cap U$ to $\Sigma \cap V$.

As a result, the definition of local Σ -equivalence gives rise to the definition of *local Σ -structural stability* of $Z \in \Omega_0$ with respect to Ω_0 near the origin, that is, $Z \in \Omega_0$ is said to be locally Σ -structurally stable with respect to Ω_0 near the origin, if any vector field that lies in a sufficiently small neighborhood of Z contained in Ω_0 is locally Σ -equivalent to Z near the origin.

2.2 Pseudo-Hopf bifurcation

It is well known that the Hopf bifurcation of smooth vector fields is a main tool to produce limit cycles, where limit cycles bifurcate from a weak focus as the stability of this focus changes. In piecewise smooth vector fields there exists a similar phenomenon, called *pseudo-Hopf bifurcation* (see, e.g., [7, 12, 16, 18, 26]), where limit cycles are created from a *pseudo-focus* as the stability of a sliding segment changes, see Figure 2. Here a point in the switching line is said to be a stable (resp. unstable) pseudo-focus if all orbits near this point turn around and tend to it as the time increases (resp. decreases) as defined in [11]. In order to prove the results of this paper conveniently, we adopt the version given in [12, Proposition 2.3] by considering the special one-parametric piecewise smooth vector field

$$Z_\delta(x, y) = \begin{cases} X(x, y) & \text{if } y > 0, \\ Y(x, y) + (0, \delta)^\top & \text{if } y < 0, \end{cases} \quad (2.1)$$

where $X = (X_1, X_2)$ and $Y = (Y_1, Y_2)$ are C^1 vector fields defined on \mathbb{R}^2 , $\delta \in \mathbb{R}$ is a parameter.




Figure 2: The pseudo-Hopf bifurcation of (2.1) satisfying $X_1(0, 0) < 0 < Y_1(0, 0)$ and the origin is stable.

Proposition 2.1. *For $\delta = 0$ we assume that the origin is a stable (resp. unstable) pseudo-focus formed by an invisible-invisible fold-fold point of the piecewise smooth vector field Z_δ and $X_1(0, 0) < 0 < Y_1(0, 0)$. Then the vector field Z_δ exhibits a pseudo-Hopf bifurcation at $\delta = 0$ for $|\delta|$ sufficiently small, more precisely, there exists some $\delta_0 > 0$ such that Z_δ has a stable (resp. unstable) crossing limit cycle bifurcating from the origin for $-\delta_0 < \delta < 0$ (resp. $\delta_0 > \delta > 0$) and has no crossing limit cycles for $\delta_0 > \delta > 0$ (resp. $-\delta_0 < \delta < 0$).*

The proof of Proposition 2.1 follows directly from the generalized Poincaré-Bendixson Theorem for piecewise smooth vector fields, see [2].

3 Proof of Theorem 1.1

This section is devoted to proving Theorem 1.1. Let $Z = (X, Y) \in \Omega_0$. We start by studying the local sliding dynamics of Z near the origin O .

Lemma 3.1. *For $Z = (X, Y) \in \Omega_0$ there exists a neighborhood $\mathcal{U}_0 \subset \mathcal{U}$ of O such that $\Sigma \cap \mathcal{U}_0$ is separated into two crossing sets by O . In addition, if $X_{2x}(0, 0) > 0$ and $Y_{2x}(0, 0) > 0$, the direction of X and Y on the right (resp. left) crossing set is upward (resp. downward), while if $X_{2x}(0, 0) < 0$ and $Y_{2x}(0, 0) < 0$, the direction of X and Y on the right (resp. left) crossing set is downward (resp. upward).*

Proof. Writing $X_2(x, 0)$ and $Y_2(x, 0)$ around $x = 0$ as

$$X_2(x, 0) = X_{2x}(0, 0)x + \mathcal{O}(x^2), \quad Y_2(x, 0) = Y_{2x}(0, 0)x + \mathcal{O}(x^2), \quad (3.1)$$

we get $X_2(x, 0)Y_2(x, 0) = X_{2x}(0, 0)Y_{2x}(0, 0)x^2 + \mathcal{O}(x^3)$. By the definition of Ω_0 , we get $X_{2x}(0, 0)Y_{2x}(0, 0) > 0$ and then there exists a neighborhood $\mathcal{U}_0 \subset \mathcal{U}$ of O such that $X_2(x, 0)Y_2(x, 0) = 0$ for $(x, 0) = O$ and $X_2(x, 0)Y_2(x, 0) > 0$ for $(x, 0) \in (\mathcal{U}_0 \cap \Sigma) \setminus \{O\}$. It follows from the definition of crossing set that $\{(x, 0) \in \mathcal{U}_0 \cap \Sigma : x < 0\}$ and $\{(x, 0) \in \mathcal{U}_0 \cap \Sigma : x > 0\}$ are two crossing sets separated by O , i.e., the first part of Lemma 3.1 is proved. The second part is obtained directly from (3.1). \square

Our main idea for proving Theorem 1.1 is to provide a normal form for $Z \in \Omega_1 \subset \Omega_0$ such that both Z and the corresponding piecewise linear vector field Z_L are locally Σ -equivalent to this normal form near the origin. Then Z is locally Σ -equivalent to Z_L near the origin, and the local phase portrait of Z is the phase portrait of this normal form in the sense of Σ -equivalence. This concludes the proof of



Figure 3: Local phase portraits of $Z \in \Omega_{ff}$ satisfying (3.2) near O .

Theorem 1.1. Therefore, in what follows we will study the normal forms of $Z \in \Omega_1$ using the method introduced in [5, 6, 16]. Such a method has been successfully applied to obtain the normal forms of piecewise smooth vector fields in Ω near a codimension-zero (resp. codimension-one) singularity in [16] (resp. [5, 6]), and near a Σ -center in [3, 29].

To this end we classify Ω_1 into the following six subsets:

$$\begin{aligned}
\Omega_{ff} &= \{Z \in \Omega_1 : \lambda_1^\pm, \lambda_2^\pm \in \mathbb{C} \setminus \mathbb{R}\}, \\
\Omega_{fn} &= \{Z \in \Omega_1 : \text{either } \lambda_1^+, \lambda_2^+ \in \mathbb{C} \setminus \mathbb{R}, \lambda_1^-, \lambda_2^- \in \mathbb{R}, \lambda_1^- \lambda_2^- > 0 \text{ or } \lambda_1^-, \lambda_2^- \in \mathbb{C} \setminus \mathbb{R}, \lambda_1^+, \lambda_2^+ \in \mathbb{R}, \lambda_1^+ \lambda_2^+ > 0\}, \\
\Omega_{fs} &= \{Z \in \Omega_1 : \text{either } \lambda_1^+, \lambda_2^+ \in \mathbb{C} \setminus \mathbb{R}, \lambda_1^-, \lambda_2^- \in \mathbb{R}, \lambda_1^- \lambda_2^- < 0 \text{ or } \lambda_1^-, \lambda_2^- \in \mathbb{C} \setminus \mathbb{R}, \lambda_1^+, \lambda_2^+ \in \mathbb{R}, \lambda_1^+ \lambda_2^+ < 0\}, \\
\Omega_{nn} &= \{Z \in \Omega_1 : \lambda_1^\pm, \lambda_2^\pm \in \mathbb{R}, \lambda_1^+ \lambda_2^+ > 0, \lambda_1^- \lambda_2^- > 0\}, \\
\Omega_{ns} &= \{Z \in \Omega_1 : \lambda_1^\pm, \lambda_2^\pm \in \mathbb{R}, \text{either } \lambda_1^+ \lambda_2^+ > 0, \lambda_1^- \lambda_2^- < 0 \text{ or } \lambda_1^+ \lambda_2^+ < 0, \lambda_1^- \lambda_2^- > 0\}, \\
\Omega_{ss} &= \{Z \in \Omega_1 : \lambda_1^\pm, \lambda_2^\pm \in \mathbb{R}, \lambda_1^+ \lambda_2^+ < 0, \lambda_1^- \lambda_2^- < 0\}.
\end{aligned}$$

Clearly,

$$\Omega_1 = \Omega_{ff} \cup \Omega_{fn} \cup \Omega_{fs} \cup \Omega_{nn} \cup \Omega_{ns} \cup \Omega_{ss}.$$

Now we study the normal forms for $Z = (X, Y) \in \Omega_{ff}, \Omega_{fn}, \Omega_{fs}, \Omega_{nn}, \Omega_{ns}$ and Ω_{ss} , respectively.

Lemma 3.2. *If $Z = (X, Y) \in \Omega_{ff}$, then Z is locally Σ -equivalent to $Z_{ff} = (X_{ff}, Y_{ff}) \in \Omega_{ff}$ near the origin, where*

$$X_{ff}(x, y) = (\alpha x - y, x + \alpha y), \quad Y_{ff}(x, y) = (\alpha x - y, x + \alpha y),$$

$\alpha = \text{sign} \ell$ and $\ell \neq 0$ is defined in (1.6).

Proof. Because $\Omega_{ff} \subset \Omega_1 \subset \Omega_0$, $Z \in \Omega_{ff}$ satisfies (1.3) by the definition of Ω_0 . Using the change $(x, y) \rightarrow (-x, y)$, we only need to consider the case

$$X_{2x}(0, 0) > 0, \quad Y_{2x}(0, 0) > 0. \quad (3.2)$$

Hence, $\Sigma \cap \mathcal{U}_0$ is separated into two crossing sets by O , and the direction of X and Y on the right (resp. left) crossing set is upward (resp. downward) as it is seen in Lemma 3.1. Recalling [11, Theorem B] and [18, Theorem 1.2], we obtain that O is a stable pseudo-focus if $\ell < 0$ and an unstable pseudo-focus if $\ell > 0$ for $Z \in \Omega_{ff}$ satisfying (3.2), see Figure 3. For $Z_{ff} \in \Omega_{ff}$ it is a linear vector field, and O is a stable focus as shown in (FF-1) of Figure 1 if $\alpha = -1$ and an unstable focus as shown in (FF-2) of Figure 1 if $\alpha = 1$.

Next we prove this lemma for the case $\ell < 0$ and $\alpha = -1$. The case $\ell > 0$ and $\alpha = 1$ can be treated similarly. Consider two sufficiently small neighborhoods $U \subset \mathcal{U}_0$ and $V \subset \mathcal{U}_0$ of O as shown in Figure 4,




Figure 4: The homeomorphism H between $Z \in \Omega_{ff}$ with $\ell < 0$ and Z_{ff} with $\alpha = -1$.

where \mathcal{U}_0 is given in Lemma 3.1, U is surrounded by the closed line segment $\overline{CA} \subset \Sigma$ and the orbital arc of Z from A to C after passing through B , V is surrounded by the closed line segment $\overline{C_1A_1} \subset \Sigma$ and the orbital arc of Z_{ff} from A_1 to C_1 after passing through B_1 . Here overline denotes the closure. We need to construct a homeomorphism H from U to V implying the Σ -equivalence between Z with $\ell < 0$ and Z_{ff} with $\alpha = -1$.

For $Z \in \Omega_{ff}$ satisfying (3.2), O is an anticlockwise rotary equilibrium of focus type of X and Y . Thus, given $P \in \overline{OA}$, there exist a first time $t_1 = t_1(P) \geq 0$ such that $\Phi^+(t_1, P) \in \overline{OB}$, and a first time $t_2 = t_2(\Phi^+(t_1, P)) \geq 0$ such that $\Phi^-(t_2, \Phi^+(t_1, P)) \in \overline{OC}$, where Φ^+ and Φ^- denote the flows of X and Y respectively. This means that we can define a Poincaré map $\mathcal{P} : \overline{OA} \rightarrow \overline{OC}$ by

$$\mathcal{P}(P) = \Phi^-(t_2, \Phi^+(t_1, P)). \quad (3.3)$$

In particular, $\mathcal{P}(O) = O$ and $\mathcal{P}(A) = C$, since A and C lie in the same orbit. Let $(x_P, 0)$ and $(\mathcal{P}_1(x_P), \mathcal{P}_2(x_P))$ be the coordinates of P and $\mathcal{P}(P)$ respectively. Then $\mathcal{P}_2(x_P) = 0$ and $\mathcal{P}_1(x_P)$ is given by

$$\mathcal{P}_1(x_P) = e^{\ell\pi} x_P + O(x_P^2)$$

from [18, Theorem 1.1, Theorem 1.2].

Similarly, denoting the flows of X_{ff} and Y_{ff} by Ψ^+ and Ψ^- respectively, we can define a Poincaré map $\mathcal{Q} : \overline{OA_1} \rightarrow \overline{OC_1}$ by

$$\mathcal{Q}(P) = \Psi^-(s_2, \Psi^+(s_1, P)), \quad (3.4)$$

which satisfies $\mathcal{Q}(O) = O$ and $\mathcal{Q}(A_1) = C_1$, where $s_1 = s_1(P) \geq 0$ is the first time such that $\Psi^+(s_1, P) \in \overline{OB_1}$, and $s_2 = s_2(\Psi^+(s_1, P)) \geq 0$ is the first time such that $\Psi^-(s_2, \Psi^+(s_1, P)) \in \overline{OC_1}$. Let $(\mathcal{Q}_1(x_P), \mathcal{Q}_2(x_P))$ be the coordinates of $\mathcal{Q}(P)$. Then $\mathcal{Q}_2(x_P) = 0$ and a straightway calculation yields

$$\mathcal{Q}_1(x_P) = e^{-2\pi} x_P.$$

Since we are considering the case of $\ell < 0$, according to the linearization and conjugacy theory of smooth map [20], U and V can be chosen to ensure that there exists a homeomorphism $h : [0, x_A] \rightarrow [0, x_{A_1}]$ satisfying

$$h(0) = 0, \quad h(x_A) = x_{A_1}, \quad h(\mathcal{P}_1(x_P)) = \mathcal{Q}_1(h(x_P)), \quad (3.5)$$

where x_A and x_{A_1} are the first coordinates of A and A_1 respectively. Consequently, we define a homeomorphism $H_0 : \overline{OA} \rightarrow \overline{OA_1}$ by

$$H_0(P) = H_0(x_P, 0) = (h(x_P), 0) \quad \text{for } P \in \overline{OA}. \quad (3.6)$$

Clearly, it follows from (3.5) that $H_0(O) = O$, $H_0(A) = A_1$ and $H_0(C) = C_1$.

Given $P \in \overline{OB}$, there exists a first time $t_3 = t_3(P) \leq 0$ such that $\Phi^+(t_3, P) \in \overline{OA}$, since O is an anticlockwise rotary equilibrium of focus type of X . Then $H_0(\Phi^+(t_3, P)) \in \overline{OA_1}$ and there exists a first time $s_3 = s_3(H_0(\Phi^+(t_3, P))) \geq 0$ such that $\Psi^+(s_3, H_0(\Phi^+(t_3, P))) \in \overline{OB_1}$ because O is an anticlockwise rotary focus of X_{ff} . By the arc length parametrization we can identify the orbital arc of X from $\Phi^+(t_3, P)$ to P with the one of X_{ff} from $H_0(\Phi^+(t_3, P))$ to $\Psi^+(s_3, H_0(\Phi^+(t_3, P)))$. Therefore, in this way we can define a homeomorphism $H^+ : \overline{\Sigma^+ \cap U} \rightarrow \overline{\Sigma^+ \cap V}$ that maps \overline{BA} onto $\overline{B_1A_1}$, maps the orbits of X in $\overline{\Sigma^+ \cap U}$ onto the orbits of X_{ff} in $\overline{\Sigma^+ \cap V}$ and satisfies

$$H^+|_{\overline{OA}} = H_0. \quad (3.7)$$

Given $P \in \overline{OC}$, there exists a first time $t_4 = t_4(P) \leq 0$ such that $\Phi^-(t_4, P) \in \overline{OB}$. Then $H^+(\Phi^-(t_4, P)) \in \overline{OB_1}$ from the definition of H^+ , and there exists a first time $s_4 = s_4(H^+(\Phi^-(t_4, P))) \geq 0$ such that $\Psi^-(s_4, H^+(\Phi^-(t_4, P))) \in \overline{OC_1}$. Similarly we can identify the orbital arc of Y from $\Phi^-(t_4, P)$ to P with the one of Y_{ff} from $H^+(\Phi^-(t_4, P))$ to $\Psi^-(s_4, H^+(\Phi^-(t_4, P)))$, and thus define a homeomorphism $H^- : \overline{\Sigma^- \cap U} \rightarrow \overline{\Sigma^- \cap V}$ that maps \overline{BC} onto $\overline{B_1C_1}$, maps the orbits of Y in $\overline{\Sigma^- \cap U}$ onto the orbits of Y_{ff} in $\overline{\Sigma^- \cap V}$ and satisfies

$$H^-|_{\overline{OB}} = H^+|_{\overline{OB}}. \quad (3.8)$$

Moreover, for any $P \in \overline{OC}$ we have

$$\begin{aligned} H^-(P) &= \Psi^-(s_4, H^+(\Phi^-(t_4, P))) = \Psi^-(s_4, \Psi^+(s_3, H_0(\Phi^+(t_3, \Phi^-(t_4, P))))) \\ &= Q(H_0(\Phi^+(t_3, \Phi^-(t_4, P)))) = H_0(\mathcal{P}(\Phi^+(t_3, \Phi^-(t_4, P)))) \\ &= H_0(P) \end{aligned}$$

by (3.3), (3.4), (3.5), (3.6) and the constructions of H^\pm . This implies that

$$H^-|_{\overline{OC}} = H_0|_{\overline{OC}}. \quad (3.9)$$

Let

$$H(P) = \begin{cases} H^+(P) & \text{for } P \in (\Sigma^+ \cup \Sigma) \cap U, \\ H^-(P) & \text{for } P \in (\Sigma^- \cup \Sigma) \cap U. \end{cases} \quad (3.10)$$

Then H is a homeomorphism from U to V because H^\pm are homeomorphisms in their domains and $H^+|_{\overline{BC}} = H^-|_{\overline{BC}}$ by (3.7), (3.8) and (3.9). Furthermore, the construction of H ensures that H maps the orbits of $Z \in \Omega_{ff}$ with $\ell < 0$ in U onto the orbits of Z_{ff} with $\alpha = -1$ in V , preserving the direction of time and the switching line Σ . We eventually conclude that $Z \in \Omega_{ff}$ with $\ell < 0$ and Z_{ff} with $\alpha = -1$ are locally Σ -equivalent near O . \square

Lemma 3.3. *If $Z = (X, Y) \in \Omega_{fn}$, then Z is locally Σ -equivalent to $Z_{fn} = (X_{fn}, Y_{fn}) \in \Omega_{fn}$ near the origin, where*

$$X_{fn}(x, y) = (-y, x), \quad Y_{fn}(x, y) = (2\beta x + y, x + 2\beta y)$$

and

$$\beta = \begin{cases} \text{sign}(\lambda_1^- + \lambda_2^-) & \text{when } \lambda_1^-, \lambda_2^- \in \mathbb{R}, \\ \text{sign}(\lambda_1^+ + \lambda_2^+) & \text{when } \lambda_1^+, \lambda_2^+ \in \mathbb{R}. \end{cases}$$



Figure 5: Local phase portraits of $Z \in \Omega_{fn}$ satisfying (3.2) and $\lambda_1^+, \lambda_2^+ \in \mathbb{C} \setminus \mathbb{R}, \lambda_1^-, \lambda_2^- \in \mathbb{R}, \lambda_1^- \lambda_2^- > 0$ near O .




Figure 6: The homeomorphism H between $Z \in \Omega_{fn}$ with $\lambda_1^- + \lambda_2^- > 0$ and Z_{fn} with $\beta = 1$.

Proof. By $(x, y) \rightarrow (x, -y)$ and $(x, y) \rightarrow (-x, y)$ we only need to consider $Z \in \Omega_{fn}$ satisfying (3.2) and

$$\lambda_1^+, \lambda_2^+ \in \mathbb{C} \setminus \mathbb{R}, \quad \lambda_1^-, \lambda_2^- \in \mathbb{R}, \quad \lambda_1^- \lambda_2^- > 0.$$

In this case, O is an equilibrium of focus type of X and a node of Y by [31, Theorems 4.2, 4.3, 5.1]. Thus, recalling the dynamics on Σ given in Lemma 3.1, we get two different types of the local phase portraits of Z near O depending on the sign of $\lambda_1^- + \lambda_2^-$, namely the stability of O when it is regarded as an equilibrium of Y , see Figure 5. In Figure 5(a), the strong unstable manifold m_s^u lies in the left side of the weak unstable manifold m_w^u , while in Figure 5(b), the strong stable manifold m_s^s lies in the right side of the weak stable manifold m_w^s . Here we use the assumption of $\lambda_1^- \neq \lambda_2^-$ for all vector fields in Ω_1 . Regarding the vector field Z_{fn} , we easily verify that its phase portrait is the one either as shown in (FN-1) of Figure 1 if $\beta = 1$, or as shown in (FN-2) of Figure 1 if $\beta = -1$.

We only consider $\lambda_1^- + \lambda_2^- > 0$ and $\beta = 1$ because the case of $\lambda_1^- + \lambda_2^- < 0$ and $\beta = -1$ is similar. Consider two sufficiently small neighborhoods $U \subset \mathcal{U}_0$ and $V \subset \mathcal{U}_0$ of O as shown in Figure 6, where \mathcal{U}_0 is given in Lemma 3.1, U is surrounded by orbital arc \widehat{AB} of X from A to B , and arc \widehat{BA} on which Y is transverse to it, V is surrounded by orbital arc $\widehat{A_2B_2}$ of X_{fn} from A_2 to B_2 , and arc $\widehat{B_2A_2}$ on which the vector field Y_{fn} is transverse to it. We need to construct a homeomorphism H from U to V providing the Σ -equivalence between $Z \in \Omega_{fn}$ with $\lambda_1^- + \lambda_2^- > 0$ and Z_{fn} with $\beta = 1$.

By the arc length parametrization there exists a homeomorphism $H_0 : \overline{OA} \rightarrow \overline{OA_2}$ such that $H_0(O) = O$ and $H_0(A) = A_2$. Since O is an anticlockwise rotary equilibrium of focus type of X , the forward orbit of X starting from $P \in \overline{OA}$ evolves in $\overline{\Sigma^+ \cap U}$ until it reaches \overline{OB} at a point Q . Then $H_0(P) \in \overline{OA_2}$. Since O is an anticlockwise rotary center of X_{fn} , the forward orbit of X_{fn} starting from $H_0(P)$ evolves in $\overline{\Sigma^+ \cap V}$ until it reaches $\overline{OB_2}$ at a point Q_2 . By the arc length parametrization we can identify the orbital arc of X from P to Q with the one of X_{fn} from $H_0(P)$ to Q_2 . In this way we can define a homeomorphism $H_f : \overline{\Sigma^+ \cap U} \rightarrow \overline{\Sigma^+ \cap V}$ that maps \overline{BA} onto $\overline{B_2A_2}$, maps the orbits of X in $\overline{\Sigma^+ \cap U}$ onto the orbits of X_{fn}

in $\overline{\Sigma^+ \cap V}$ and satisfies

$$H_f|_{\overline{OA}} = H_0. \quad (3.11)$$

Consider the region R_{BOC} surrounded by \overline{OB} , \widehat{BC} and the strong unstable manifold \widehat{OC} , and the corresponding region $R_{B_2OC_2}$ surrounded by $\overline{OB_2}$, $\widehat{B_2C_2}$ and the strong unstable manifold $\widehat{OC_2}$. Given $P \in \overline{OB}$, there exists a unique point $Q \in \widehat{BC}$ such that the backward orbit of Y starting from Q evolves in $\overline{R_{BOC}}$ until it reaches or tends to \overline{OB} at P , since \widehat{OC} is the strong unstable manifold of the node O for Y and we are assuming that the vector field Y on \widehat{BA} is transverse to \widehat{BA} . Analogously, there exists a unique point $Q_2 \in \widehat{B_2C_2}$ such that the backward orbit of Y_{fn} starting from Q_2 evolves in $\overline{R_{B_2OC_2}}$ until it reaches or tends to $\overline{OB_2}$ at $H_f(P)$. Therefore, by the arc length parametrization again we can identify the orbital arc of Y from P to Q with the one of Y_{fn} from $H_f(P)$ to Q_2 , and then define a homeomorphism $H_n^1 : \overline{R_{BOC}} \rightarrow \overline{R_{B_2OC_2}}$ that maps the orbits of Y in $\overline{R_{BOC}}$ onto the orbits of Y_{fn} in $\overline{R_{B_2OC_2}}$ and satisfies

$$H_n^1|_{\overline{OB}} = H_f|_{\overline{OB}}. \quad (3.12)$$

Consider the region R_{COA} surrounded by \widehat{OC} , \widehat{CA} and \overline{OA} , and the corresponding region $R_{C_2OA_2}$ surrounded by $\widehat{OC_2}$, $\widehat{C_2A_2}$ and $\overline{OA_2}$. Regarding arcs \widehat{CA} and $\widehat{C_2A_2}$, we obtain a homeomorphism $H_n^0 : \widehat{CA} \rightarrow \widehat{C_2A_2}$ such that $H_n^0(C) = C_2$ and $H_n^0(A) = A_2$ by the arc length parametrization. Since the choice of U ensures that the vector field Y on $(\widehat{CA} \cup \overline{OA}) \setminus O$ is transverse to $(\widehat{CA} \cup \overline{OA}) \setminus O$, the backward orbit of Y starting from $P \in (\widehat{CA} \cup \overline{OA}) \setminus O$ evolves in $\overline{R_{COA}}$ and finally tends to O . Let $P_2 = H_0(P)$ if $P \in \overline{OA}$ and $P_2 = H_n^0(P)$ if $P \in \widehat{CA}$. Then the backward orbit of Y_{fn} starting from P_2 evolves in $\overline{R_{C_2OA_2}}$ and tends to O . Identify the orbital arc of Y from P to O with the orbital arc of Y_{fn} from P_2 to O . In this way we can define a homeomorphism $H_n^2 : \overline{R_{COA}} \rightarrow \overline{R_{C_2OA_2}}$ that maps the orbits of Y in $\overline{R_{COA}}$ onto the orbits of Y_{fn} in $\overline{R_{C_2OA_2}}$ and satisfies

$$H_n^2|_{\widehat{OC}} = H_n^1|_{\widehat{OC}}, \quad H_n^2|_{\overline{OA}} = H_0, \quad H_n^2|_{\widehat{CA}} = H_n^0. \quad (3.13)$$

Joining the homeomorphisms H_n^1 and H_n^2 , by (3.11), (3.12) and (3.13) we obtain that

$$H_n(P) = \begin{cases} H_n^1(P) & \text{for } P \in \overline{R_{BOC}}, \\ H_n^2(P) & \text{for } P \in \overline{R_{COA}}, \end{cases}$$

is a homeomorphism from $\overline{\Sigma^- \cap U}$ to $\overline{\Sigma^- \cap V}$ that maps the orbits of Y in $\overline{\Sigma^- \cap U}$ onto the orbits of Y_{fn} in $\overline{\Sigma^- \cap V}$ and satisfies $H_n|_{\overline{BA}} = H_f|_{\overline{BA}}$. Consequently, the homeomorphisms H_n and H_f form a homeomorphism $H : U \rightarrow V$ that maps the orbits of $Z \in \Omega_{fn}$ with $\lambda_1^- + \lambda_2^- > 0$ in U onto the orbits of Z_{fs} with $\beta = 1$ in V , preserving the direction of time and the switching line Σ . This concludes the proof of Lemma 3.3. \square

Lemma 3.4. *If $Z = (X, Y) \in \Omega_{fs}$, then Z is locally Σ -equivalent to $Z_{fs} = (X_{fs}, Y_{fs}) \in \Omega_{fs}$ near the origin, where*

$$X_{fs}(x, y) = (-y, x), \quad Y_{fs}(x, y) = (y, x).$$




Figure 7: Local phase portrait of $Z \in \Omega_{fs}$ satisfying (3.2) and $\lambda_1^+, \lambda_2^+ \in \mathbb{C} \setminus \mathbb{R}$, $\lambda_1^-, \lambda_2^- \in \mathbb{R}$, $\lambda_1^- \lambda_2^- < 0$ near O .




Figure 8: The homeomorphism H between $Z \in \Omega_{fs}$ and Z_{fs} .

Proof. Using the changes $(x, y) \rightarrow (x, -y)$ and $(x, y) \rightarrow (-x, y)$, we only need to consider $Z \in \Omega_{fs}$ satisfying (3.2) and

$$\lambda_1^+, \lambda_2^+ \in \mathbb{C} \setminus \mathbb{R}, \quad \lambda_1^-, \lambda_2^- \in \mathbb{R}, \quad \lambda_1^- \lambda_2^- < 0.$$

In this case, O is an equilibrium of focus type of X and a saddle of Y by [31, Theorems 4.2, 4.4, 5.1]. Reviewing the dynamics on Σ given in Lemma 3.1, we depict the local phase portrait of Z near O as shown in Figure 7. The phase portrait of the vector field Z_{fs} is as shown in (FS) of Figure 1.

Consider two sufficiently small neighborhoods $U \subset \mathcal{U}_0$ and $V \subset \mathcal{U}_0$ of O as shown in Figure 8, where \widehat{AB} and $\widehat{A_3B_3}$ are the corresponding orbital arcs, \widehat{BA} (resp. $\widehat{B_3A_3}$) is the arc where the vector field Y (resp. Y_{fs}) is transverse to it. As done in the proof of Lemma 3.3, we can define a homeomorphism $H_f : \overline{\Sigma^+ \cap U} \rightarrow \overline{\Sigma^+ \cap V}$ that maps \overline{BA} onto $\overline{B_3A_3}$, and maps the orbits of X in $\overline{\Sigma^+ \cap U}$ onto the orbits of X_{fs} in $\overline{\Sigma^+ \cap V}$.

In order to complete this proof, next we construct a homeomorphism $H_s : \overline{\Sigma^- \cap U} \rightarrow \overline{\Sigma^- \cap V}$ that maps the orbits of Y in $\overline{\Sigma^- \cap U}$ onto the orbits of X_{fs} in $\overline{\Sigma^- \cap V}$ and satisfies $H_s|_{\overline{BA}} = H_f|_{\overline{BA}}$. Let

$$\widehat{OD} = \{(x, y) \in \overline{\Sigma^- \cap U} : Y_2(x, y) = 0\}, \quad \overline{OD_3} = \{(x, y) \in \overline{\Sigma^- \cap V} : x = 0\},$$

where Y_2 is the ordinate of Y . Then there exists a homeomorphism $H_s^0 : \widehat{OD} \rightarrow \overline{OD_3}$ such that $H_s^0(O) = O$ and $H_s^0(D) = D_3$ by the arc length parametrization. Consider the region R_{BOD} surrounded by \overline{OB} , \widehat{BD} and \widehat{OD} , and the region $R_{B_3OD_3}$ surrounded by $\overline{OB_3}$, $\widehat{B_3D_3}$ and $\overline{OD_3}$. Given $P \in \overline{OB} \cup \widehat{OD}$, there exists a unique point $Q \in \widehat{BD}$ such that the backward orbit of Y starting from Q evolves in $\overline{R_{BOD}}$ until it either reaches $(\overline{OB} \cup \widehat{OD}) \setminus O$ when $P \neq O$ or tends to O when $P = O$, since we require that the vector field Y on \widehat{BD} is transverse to \widehat{BD} . Let $P_3 = H_f(P)$ if $P \in \overline{OB}$ and $P_3 = H_s^0(P)$ if $P \in \widehat{OD}$. We obtain a unique point $Q_3 \in \widehat{B_3D_3}$ such that the backward orbit of Y_{fs} starting from Q_3 evolves in $\overline{R_{B_3OD_3}}$ until it reaches or tends to P_3 . The arc length parametrization allows to identify the orbital arc of Y from Q to P and the one of Y_{fs} from Q_3 to P_3 . In this way we can define a homeomorphism $H_s^1 : \overline{R_{BOD}} \rightarrow \overline{R_{B_3OD_3}}$ that maps

the orbits of Y in $\overline{R_{BOD}}$ onto the orbits of Y_{fs} in $\overline{R_{B_3OD_3}}$ and satisfies

$$H_s^1|_{\overline{OB}} = H_f|_{\overline{OB}}, \quad H_s^1|_{\overline{OD}} = H_s^0. \quad (3.14)$$

A similar argument to the last paragraph yields a homeomorphism $H_s^2 : \overline{R_{DOA}} \rightarrow \overline{R_{D_3OA_3}}$ that maps the orbits of Y in $\overline{R_{DOA}}$ onto the orbits of Y_{fs} in $\overline{R_{D_3OA_3}}$ and satisfies

$$H_s^2|_{\overline{OA}} = H_f|_{\overline{OA}}, \quad H_s^2|_{\overline{OD}} = H_s^0. \quad (3.15)$$

Thus, joining the homeomorphisms H_s^1 and H_s^2 we construct H_s as

$$H_s(P) = \begin{cases} H_s^1(P) & \text{for } P \in \overline{R_{BOD}}, \\ H_s^2(P) & \text{for } P \in \overline{R_{DOA}}. \end{cases}$$

From (3.14) and (3.15), it follows that H_s is a homeomorphism from $\overline{\Sigma^- \cap U}$ to $\overline{\Sigma^- \cap V}$ maps the orbits of Y in $\overline{\Sigma^- \cap U}$ onto the orbits of X_{fs} in $\overline{\Sigma^- \cap V}$ and satisfies $H_s|_{\overline{BA}} = H_f|_{\overline{BA}}$.

Consequently, the homeomorphisms H_s and H_f directly form a homeomorphism $H : U \rightarrow V$ that maps the orbits of $Z \in \Omega_{fs}$ in U onto the orbits of Z_{fs} in V , preserving the direction of time and the switching line Σ . This proves Lemma 3.4. \square

Lemma 3.5. *If $Z = (X, Y) \in \Omega_{nn}$, then Z is locally Σ -equivalent to $Z_{nn} = (X_{nn}, Y_{nn}) \in \Omega_{nn}$ near the origin, where*

$$X_{nn}(x, y) = (2\gamma x + y, x + 2\gamma y), \quad Y_{nn}(x, y) = (2\eta x + y, x + 2\eta y),$$

and

$$\begin{cases} \gamma = \eta = \text{sign}(\lambda_1^+ + \lambda_2^+) & \text{when } (\lambda_1^+ + \lambda_2^+)(\lambda_1^- + \lambda_2^-) > 0, \\ \gamma = -\eta = 1 & \text{when } (\lambda_1^+ + \lambda_2^+)(\lambda_1^- + \lambda_2^-) < 0. \end{cases}$$

Proof. For $Z \in \Omega_{nn}$ we know that O is a node of both X and Y with two different eigenvalues by [31, Theorem 4.3]. Moreover, using the change $(x, y) \rightarrow (-x, y)$ it is enough to consider $Z \in \Omega_{nn}$ satisfying (3.2). In this case, according to the dynamics on Σ given in Lemma 3.1, we get four local phase portraits of Z near O as shown in Figure 9, depending on the sign of $\lambda_1^+ + \lambda_2^+$, namely the stability of O as an equilibrium of X and Y . However, we notice that the phase portrait (d) of Figure 9 can be transformed into (b) of Figure 9 by the change $(x, y) \rightarrow (-x, -y)$, so that there are essentially three different types of the local phase portraits of Z near O . Besides, a simple analysis implies that the phase portrait of Z_{nn} is (NN-1) (resp. (NN-2) and (NN-3)) of Figure 1 if $\gamma = \eta = 1$ (resp. $\gamma = -\eta = 1$ and $\gamma = \eta = -1$).

The homeomorphism between $Z \in \Omega_{nn}$ and Z_{nn} can be constructed by a similar method to the proofs of foregoing lemmas. In fact, consider the case of $\lambda_1^+ + \lambda_2^+ > 0, \lambda_1^- + \lambda_2^- > 0$ and $\gamma = \eta = 1$ as an example. We can choose two sufficiently small neighborhoods $U \subset \mathcal{U}_0$ and $V \subset \mathcal{U}_0$ of O such that Z is transverse to the boundary of U and Z_{nn} is transverse to the boundary of V . Then there is always a homeomorphism $H : \Sigma \cap U \rightarrow \Sigma \cap V$ satisfying $H(O) = O, H(\Sigma_l \cap U) = \Sigma_l \cap V$ and $H(\Sigma_r \cap U) = \Sigma_r \cap V$, where $\Sigma_l = \{(x, 0) \in \mathcal{U} : x < 0\}$ and $\Sigma_r = \{(x, 0) \in \mathcal{U} : x > 0\}$. Like the construction of H_n in the proof of Lemma 3.3, we are able to extend H for $\Sigma^+ \cap U$ and $\Sigma^- \cap U$ respectively, and finally obtain a homeomorphism from U to V that provides the Σ -equivalence between $Z \in \Omega_{nn}$ with $\lambda_1^+ + \lambda_2^+ > 0, \lambda_1^- + \lambda_2^- > 0$ and Z_{nn} with $\gamma = \eta = 1$. That is, Lemma 3.5 holds. \square




Figure 9: Local phase portraits of $Z \in \Omega_{mn}$ satisfying (3.2) near O .



Figure 10: Local phase portraits of $Z \in \Omega_{ns}$ satisfying (3.2), $\lambda_1^+ \lambda_2^+ > 0$ and $\lambda_1^- \lambda_2^- < 0$ near O .

Lemma 3.6. *If $Z = (X, Y) \in \Omega_{ns}$, then Z is locally Σ -equivalent to $Z_{ns} = (X_{ns}, Y_{ns}) \in \Omega_{ns}$ near the origin, where*

$$X_{ns}(x, y) = (2\xi x + y, x + 2\xi y), \quad Y_{ns}(x, y) = (y, x),$$

and

$$\xi = \begin{cases} \text{sign}(\lambda_1^- + \lambda_2^-) & \text{when } \lambda_1^- \lambda_2^- > 0, \\ \text{sign}(\lambda_1^+ + \lambda_2^+) & \text{when } \lambda_1^+ \lambda_2^+ > 0. \end{cases}$$

Proof. Using the changes $(x, y) \rightarrow (x, -y)$ and $(x, y) \rightarrow (-x, y)$, we only need to consider $Z \in \Omega_{ns}$ satisfying (3.2), $\lambda_1^+ \lambda_2^+ > 0$ and $\lambda_1^- \lambda_2^- < 0$. In this case, O is a node of X and a saddle of Y by [31, Theorems 4.3, 4.4]. Combining with the dynamics on Σ given in Lemma 3.1, we get two different types of the local phase portraits of Z near O as shown in Figure 10, depending on the sign of $\lambda_1^+ + \lambda_2^+$. Regarding Z_{ns} , its phase portrait is (NS-1) (resp. (NS-2)) of Figure 1 if $\xi = 1$ (resp. $\xi = -1$).

Consider two sufficiently small neighborhoods $U \subset \mathcal{U}_0$ and $V \subset \mathcal{U}_0$ of O such that Z is transverse to the boundary of U and Z_{ns} is transverse to the boundary of V . For each one of the above two cases, we can define a homeomorphism H with $H(O) = O$ to identify $\Sigma \cap U$ with $\Sigma \cap V$ by the arc length parametrization. Then H can be extended for $\Sigma^+ \cap U$ (resp. $\Sigma^- \cap U$) as the construction of H_n (resp. H_s)




Figure 11: Local phase portrait of $Z \in \Omega_{ss}$ satisfying (3.2) near O .

in the proof of Lemma 3.3 (resp. Lemma 3.4). That is, H is a homeomorphism from U to V that provides Σ -equivalence, and then Lemma 3.6 holds. \square

Lemma 3.7. *If $Z = (X, Y) \in \Omega_{ss}$, then Z is locally Σ -equivalent to $Z_{ss} = (X_{ss}, Y_{ss}) \in \Omega_{ss}$ near the origin, where*

$$X_{ss}(x, y) = (y, x), \quad Y_{ss}(x, y) = (y, x).$$

Proof. For $Z \in \Omega_{ss}$ we know that O is a saddle of both X and Y by [31, Theorem 4.4]. Using the change $(x, y) \rightarrow (-x, y)$ we only need to consider $Z \in \Omega_{ss}$ satisfying (3.2). Together with the dynamics on Σ given in Lemma 3.1, this implies that the local phase portrait of Z near O is as shown in Figure 11. Moreover, the phase portrait of Z_{ss} is (SS) of Figure 1.

Consider two sufficiently small neighborhoods $U \subset \mathcal{U}_0$ and $V \subset \mathcal{U}_0$ of O such that Z is transverse to the boundary of U and Z_{ss} is transverse to the boundary of V . We can define a homeomorphism H with $H(O) = O$ to identify $\Sigma \cap U$ with $\Sigma \cap V$ by the arc length parametrization. Repeating the construction of H_s in the proof of Lemma 3.4, we extend H for $\Sigma^+ \cap U$ and $\Sigma^- \cap U$ respectively, and finally obtain a homeomorphism from U to V that provides Σ -equivalence between $Z \in \Omega_{ss}$ and Z_{ss} . This proves Lemma 3.7. \square

Now we are in a suitable position to prove Theorem 1.1.

Proof of Theorem 1.1. For $Z \in \Omega_{ff}$ (resp. $\Omega_{fn}, \Omega_{fs}, \Omega_{nn}, \Omega_{ns}, \Omega_{ss}$), the corresponding piecewise linear vector field Z_L given in (1.4) is also in Ω_{ff} (resp. $\Omega_{fn}, \Omega_{fs}, \Omega_{nn}, \Omega_{ns}, \Omega_{ss}$). Thus, by Lemmas 3.2-3.7 both Z and Z_L are locally Σ -equivalent to Z_{ff} (resp. $Z_{fn}, Z_{fs}, Z_{nn}, Z_{ns}, Z_{ss}$) near O , which implies that Z is locally Σ -equivalent to Z_L near O if $Z \in \Omega_{ff}$ (resp. $\Omega_{fn}, \Omega_{fs}, \Omega_{nn}, \Omega_{ns}, \Omega_{ss}$). Since $\Omega_1 = \Omega_{ff} \cup \Omega_{fn} \cup \Omega_{fs} \cup \Omega_{nn} \cup \Omega_{ns} \cup \Omega_{ss}$, Z is locally Σ -equivalent to Z_L near O for every $Z \in \Omega_1$. Collecting all non-equivalent phase portraits of $Z_{ff}, Z_{fn}, Z_{fs}, Z_{nn}, Z_{ns}$ and Z_{ss} obtained in Lemmas 3.2-3.7, we get 11 local phase portraits of $Z \in \Omega_1$ near O as shown in Figure 1. \square

From Lemmas 3.2, 3.5 and 3.7 we find that some $Z \in \Omega_1$ are locally Σ -equivalent to smooth linear vector fields near the origin.

As indicated in Section 1, Theorem 1.1 does not allow the same eigenvalue for the Jacobian matrices A^+ and A^- respectively in order that the vector field in Ω_0 is locally Σ -equivalent to its linear part near the



Figure 12: Local phase portraits of Z^* and Z_L^* near O , where (a) is for Z^* and (b) is for Z_L^* .

origin. The next proposition provides an example showing that the vector field in Ω_0 might not be locally Σ -equivalent to its linear part near the origin if the Jacobian matrix A^+ or A^- has the same eigenvalue.

Proposition 3.1. *Consider the piecewise smooth vector field $Z^* = (X^*, Y^*)$ with*

$$X^*(x, y) = (y, x), \quad Y^*(x, y) = \left(x + \frac{1}{2}\Gamma(x, y), x + y + \frac{1}{2}\Gamma(x, y) \right),$$

where

$$\Gamma(x, y) = \begin{cases} (x^2 + y^2)^{1/2} \left(-\frac{1}{2} \ln(x^2 + y^2) \right)^{-3/2} & \text{if } x^2 + y^2 < 1, \\ 0 & \text{if } x^2 + y^2 = 0. \end{cases}$$

Then $Z^* \in \Omega_0$ and it is not locally Σ -equivalent to its linear part $Z_L^* = (X_L^*, Y_L^*)$ near the origin, where $X_L^*(x, y) = (y, x)$ and $Y_L^*(x, y) = (x, x + y)$.

Proof. We start by proving $Z^* = (X^*, Y^*) \in \Omega_0$. In fact, a straightway calculation implies $\Gamma(0, 0) = 0$, $\Gamma_x(0, 0) = \Gamma_y(0, 0) = 0$ and $\Gamma(x, y)$ is continuously differential near O . Thus Y^* is a C^1 vector field having O as a non-degenerate equilibrium, i.e., $Y^*(0, 0) = (0, 0)$ and the determinant of the Jacobian matrix of Y^* at O is nonzero. Clearly, the vector field X^* is also C^1 and O is a linear saddle of it. Accordingly, condition (1.2) holds for Z^* . On the other hand, we have $X_{2x}^*(0, 0) = Y_{2x}^*(0, 0) = 1$, so that (1.3) also holds for Z^* , where X_2^* and Y_2^* are the ordinates of X^* and Y^* respectively. In conclusion, we get $Z^* \in \Omega_0$ from the definition of Ω_0 given above (1.2), and the linear part of Z^* is Z_L^* from $\Gamma_x(0, 0) = \Gamma_y(0, 0) = 0$.

Next we determine the local phase portraits of Z^* and Z_L^* near O in order to prove that Z^* is not locally Σ -equivalent to Z_L^* . Regarding Z^* , O is a saddle of X^* with the unstable manifold $\{(x, y) \in \mathbb{R}^2 : y = x, y > 0\}$ and the stable manifold $\{(x, y) \in \mathbb{R}^2 : y = -x, y > 0\}$. From [31, Example 4.3] we have known that all orbits of Y^* near O starting from the negative x -axis enter into Σ^- and then reach the positive x -axis after a finite time. Thus the local phase portrait of Z^* near O is as shown in Figure 12(a). Regarding Z_L^* , O is an unstable non-diagonalizable node of Y_L^* with the characteristic direction $x = 0$. Due to $X_L^* = X^*$, we conclude that the phase portrait of Z_L^* is as shown in Figure 12(b).

Consider the orbits of Z^* and Z_L^* starting from the negative x -axis. From Figure 12 we observe that these orbits of Z^* intersect the positive x -axis, but the ones of Z_L^* do not. Since any Σ -equivalence sends the orbits of Z^* to the orbits of Z_L^* , preserving the switching line Σ , it also preserves the intersections between the orbits and Σ . Consequently, Z^* cannot be locally Σ -equivalent to Z_L^* near O . \square

4 Proof of Theorem 1.2

Since Ω_{ff} is an open set of $\Omega_1 \subset \Omega_0$, any small perturbation of $Z \in \Omega_{ff}$ inside Ω_0 belongs to Ω_{ff} . In particular, the value of sign function α defined in Lemma 3.2 is the same for $Z \in \Omega_{ff}$ and its perturbation. Thus by Lemma 3.2 both $Z \in \Omega_{ff}$ and any perturbation of it inside Ω_0 are locally Σ -equivalent to the same normal form Z_{ff} near O . This means that $Z \in \Omega_{ff}$ is locally Σ -structurally stable with respect to Ω_0 near O . Similar argument can be applied to Z belonging to Ω_{fn} , Ω_{fs} , Ω_{nn} , Ω_{ns} and Ω_{ss} respectively. Finally, due to $\Omega_1 = \Omega_{ff} \cup \Omega_{fn} \cup \Omega_{fs} \cup \Omega_{nn} \cup \Omega_{ns} \cup \Omega_{ss}$, we conclude that $Z \in \Omega_1$ is locally Σ -structurally stable with respect to Ω_0 near O , that is, the sufficiency holds.

To obtain the necessity, we can equivalently prove that $Z \in \Omega_0$ is not locally Σ -structurally stable with respect to Ω_0 near O if $Z \in \Omega_0 \setminus \Omega_1$. To do this, we classify $\Omega_0 \setminus \Omega_1$ into two subsets:

$$\Omega_2 = \{Z \in \Omega_0 : \text{Im} \lambda_1^+ \text{Im} \lambda_1^- \neq 0, \ell = 0\}, \quad \Omega_3 = \{Z \in \Omega_0 : (\lambda_1^+ - \lambda_2^+)(\lambda_1^- - \lambda_2^-) = 0\}. \quad (4.1)$$

Clearly, $\Omega_0 \setminus \Omega_1 = \Omega_2 \cup \Omega_3$. If $Z = (X, Y) \in \Omega_2$, then O is an equilibrium of focus type for both X and Y . In this case, O is a non-smooth center or pseudo-focus of focus-focus type of Z with the first Lyapunov constant $\ell = 0$ as it was clarified in [11, Theorem B]. Since ℓ only depends on the linear part of Z , we easily obtain a perturbed vector field with $\ell > 0$ and a perturbed one with $\ell < 0$ by perturbing the linear part of Z in Ω_0 . This means that, for any sufficiently small neighborhood of Z in Ω_0 , there always exist two vector fields where O is a pseudo-focus with the different stability. Even limit cycles can bifurcate from O , e.g., [32]. Then these two perturbed vector fields are not locally Σ -equivalent near O , so that $Z \in \Omega_2$ is not locally Σ -structurally stable with respect to Ω_0 near O .

If $Z = (X, Y) \in \Omega_3$, then at least one of $\lambda_1^+ = \lambda_2^+$ and $\lambda_1^- = \lambda_2^-$ holds. Without loss of generality we assume that $\lambda_1^+ = \lambda_2^+$. Writing X near O as

$$X = A^+(x, y)^\top + \Upsilon^+(x, y),$$

where $\Upsilon^+(x, y)$ is the higher order terms and

$$A^+ = \begin{pmatrix} a_{11}^+ & a_{12}^+ \\ a_{21}^+ & a_{22}^+ \end{pmatrix},$$

we get

$$\lambda_1^+ = \lambda_2^+ = \frac{1}{2}(a_{11}^+ + a_{22}^+), \quad (a_{11}^+ - a_{22}^+)^2 + 4a_{12}^+a_{21}^+ = 0, \quad a_{21}^+ \neq 0. \quad (4.2)$$

Here $a_{21}^+ \neq 0$ is due to the fact that $Z \in \Omega_0$ satisfies (1.3). Consider the vector field $Z_\varepsilon = (X_\varepsilon, Y)$ with

$$X_\varepsilon = A_\varepsilon^+(x, y)^\top + \Upsilon^+(x, y)$$

and

$$A_\varepsilon^+ = \begin{pmatrix} a_{11}^+ & \frac{a_{12}^+a_{21}^+ + \varepsilon/4}{a_{21}^+ + \varepsilon} \\ a_{21}^+ + \varepsilon & a_{22}^+ \end{pmatrix}.$$

Then for any sufficiently small neighborhood of Z in Ω_0 , there exists $\varepsilon_0 > 0$ such that Z_ε lies in the neighborhood for all $-\varepsilon_0 < \varepsilon < \varepsilon_0$. Denote the eigenvalues of A_ε^+ by $\lambda_{\varepsilon,1}^+$ and $\lambda_{\varepsilon,2}^+$. It follows from (4.2) that

$$\lambda_{\varepsilon,1}^+ = \lambda_1^+ + \frac{\sqrt{-\varepsilon}}{2}i, \quad \lambda_{\varepsilon,2}^+ = \lambda_1^+ - \frac{\sqrt{-\varepsilon}}{2}i$$

for $-\varepsilon_0 < \varepsilon < 0$, while for $\varepsilon_0 > \varepsilon > 0$,

$$\lambda_{\varepsilon,1}^+ = \lambda_1^+ + \frac{\sqrt{\varepsilon}}{2}, \quad \lambda_{\varepsilon,2}^+ = \lambda_1^+ - \frac{\sqrt{\varepsilon}}{2}.$$

In the case of $-\varepsilon_0 < \varepsilon < 0$, O is a focus of X_ε by [31, Theorem 4.2], so that all orbits of Z_ε near O starting from the positive x -axis enter into Σ^+ and then reach the negative x -axis as t increases (resp. decreases) if $a_{21}^+ > 0$ (resp. < 0). In the case of $\varepsilon_0 > \varepsilon > 0$, O is a diagonalizable node of X_ε by [31, Theorem 4.3], which has two characteristic directions with the nonzero slope due to $a_{21}^+ \neq 0$. Thus all orbits of Z_ε near O starting from the positive x -axis cannot reach the negative x -axis from Σ^+ as t increases (resp. decreases) if $a_{21}^+ > 0$ (resp. < 0). As indicated in the proof of Proposition 3.1, any Σ -equivalence sends the orbits of Z_ε with $-\varepsilon_0 < \varepsilon < 0$ to the orbits of Z_ε with $\varepsilon_0 > \varepsilon > 0$, preserving the switching line Σ and the intersections of Σ and the orbits. Consequently, Z_ε with $-\varepsilon_0 < \varepsilon < 0$ cannot be locally Σ -equivalent to Z_ε with $\varepsilon_0 > \varepsilon > 0$ near O . This means that, for any sufficiently small neighborhood of $Z \in \Omega_3$ in Ω_0 , there are always two vector fields that are not locally Σ -equivalent near O . So $Z \in \Omega_3$ is not locally Σ -structurally stable with respect to Ω_0 near O . Recalling the last paragraph, we conclude the necessity. This ends the proof of Theorem 1.2.

5 Proof of Theorem 1.3

Before proving Theorem 1.3, we study the limit cycle bifurcations by perturbing the following piecewise linear vector field

$$Z_0(x, y) = \begin{cases} X_0(x, y) = (ay, x) & \text{if } y > 0, \\ Y_0(x, y) = (by, x) & \text{if } y < 0, \end{cases} \quad (5.1)$$

where $a, b \in \mathbb{R}$ satisfies $ab \neq 0$.

Proposition 5.1. *Consider the piecewise linear vector field Z_0 with $ab \neq 0$ given in (5.1). Then $Z_0 \in \Omega_1$ if either $a > 0$ or $b > 0$, and $Z_0 \in \Omega_2$ if $a < 0$ and $b < 0$, where $\Omega_1 \subset \Omega_0$ and $\Omega_2 \subset \Omega_0 \setminus \Omega_1$ are defined in (1.5) and (4.1) separately. Moreover, Z_0 has no limit cycles.*

Proof. The first part of Proposition 5.1 follows directly from the definitions of Ω_1 and Ω_2 . Since O is saddle or center of X_0 and Y_0 , it is impossible for Z_0 to have limit cycles totally contained in the half plane $y \geq 0$ or $y \leq 0$. On the other hand, when O is a center of both X_0 and Y_0 , it is a global non-smooth center for Z_0 , so that Z_0 has no limit cycles occupying the half planes $y > 0$ and $y < 0$. Clearly, there also exist no limit cycles occupying the half planes $y > 0$ and $y < 0$ when O is a saddle of X_0 or Y_0 . \square

Next we state two bifurcation results by perturbing the piecewise linear vector field Z_0 given in (5.1).

Proposition 5.2. *Consider the piecewise linear vector field $Z_0 \in \Omega_0$ in (5.1), and the piecewise polynomial vector field*

$$Z_\epsilon^f(x, y) = \begin{cases} X_\epsilon^f(x, y) = ((a - \epsilon)y - \epsilon, x) & \text{if } y > 0, \\ Y_\epsilon^f(x, y) = \left((b - \epsilon)y + \epsilon, x + \epsilon \frac{\partial f(x, \epsilon)}{\partial x} \right) & \text{if } y < 0, \end{cases}$$

where $\epsilon \geq 0$ and

$$f(x, \epsilon) = x \prod_{i=1}^m \left(x^2 - \left(\frac{i\epsilon}{m} \right)^2 \right). \quad (5.2)$$

Then $Z_\epsilon^f = Z_0$ for $\epsilon = 0$. Besides, for any given a and b satisfying $0 < |a| \leq 1/2$ and $0 < |b| \leq 1/2$, there exists $0 < \epsilon_0 < \min\{|a|, |b|\}$ such that for $0 < \epsilon < \epsilon_0$, Z_ϵ^f has exactly m hyperbolic crossing limit cycles Γ_i ($i = 1, 2, \dots, m$) bifurcating from the non-smooth equilibrium O of Z_0 , where Γ_i obeys the algebraic curve

$$\Gamma_i^+ : \frac{1}{2}x^2 - \frac{a-\epsilon}{2}y^2 + \epsilon y = \frac{1}{2} \left(\frac{i\epsilon}{m} \right)^2$$

in the half plane $y \geq 0$ and the algebraic curve

$$\Gamma_i^- : \frac{1}{2}x^2 + \epsilon f(x, \epsilon) - \frac{b-\epsilon}{2}y^2 - \epsilon y = \frac{1}{2} \left(\frac{i\epsilon}{m} \right)^2$$

in the half plane $y \leq 0$. Moreover, Γ_i is stable if $m - i$ is even and unstable if $m - i$ is odd.

Proposition 5.3. Consider the piecewise linear vector field $Z_0 \in \Omega_0$ in (5.1), and the piecewise C^∞ vector field

$$Z_\epsilon^g(x, y) = \begin{cases} X_\epsilon^g(x, y) = ((a - \epsilon)y - \epsilon, x) & \text{if } y > 0, \\ Y_\epsilon^g(x, y) = \left((b - \epsilon)y + \epsilon, x + \epsilon \frac{\partial g(x, \epsilon)}{\partial x} \right) & \text{if } y < 0, \end{cases}$$

where $\epsilon \geq 0$ and $g(x, \epsilon)$ is a C^∞ function given by

$$g(x, \epsilon) = \begin{cases} 0 & \text{if } x \leq 0, \\ e^{-1/x} \sin\left(\frac{\pi\epsilon}{x}\right) & \text{if } x > 0. \end{cases}$$

Then $Z_\epsilon^g = Z_0$ for $\epsilon = 0$. Besides, for any given a and b satisfying $0 < |a| \leq 1/2$ and $0 < |b| \leq 1/2$, there exists $0 < \epsilon_0 < \min\{|a|, |b|\}$ such that for $0 < \epsilon < \epsilon_0$, Z_ϵ^g has infinitely many hyperbolic crossing limit cycles Θ_i ($i \in \mathbb{N}^+$) bifurcating from the non-smooth equilibrium O of Z_0 , where Θ_i obeys the algebraic curve

$$\Theta_i^+ : \frac{1}{2}x^2 - \frac{a-\epsilon}{2}y^2 + \epsilon y = \frac{1}{2} \left(\frac{\epsilon}{i} \right)^2$$

in the half plane $y \geq 0$ and the algebraic curve

$$\Theta_i^- : \frac{1}{2}x^2 + \epsilon g(x, \epsilon) - \frac{b-\epsilon}{2}y^2 - \epsilon y = \frac{1}{2} \left(\frac{\epsilon}{i} \right)^2$$

in the half plane $y \leq 0$. Moreover, Θ_i is stable if i is odd and unstable if i is even.

Propositions 5.2 and 5.3 will be proved later on. If $a = b > 0$ (resp. < 0), then $Z_0 = X_0 = Y_0$ is a linear vector field having O as a saddle (resp. center). Thus our results reveal that any finitely or infinitely many limit cycles can bifurcate from some linear saddle and center under non-smooth perturbations. Besides, observe that Z_ϵ^f and Z_ϵ^g are both piecewise smooth Hamiltonian systems. This means that it is possible for piecewise smooth Hamiltonian systems to have limit cycles, but this cannot occur in smooth Hamiltonian systems as well known.

Now we are in a position to provide the proof of Theorem 1.3.

Proof of Theorem 1.3. For $Z = (X, Y) \in \Omega_0$ we consider the three-parametric perturbed vector field $Z^\epsilon = (X^\epsilon, Y^\epsilon)$ with

$$\begin{aligned} X^\epsilon(x, y) &= (X_1(x, y) - X_{2x}(0, 0)\epsilon_1 + X_{2x}(0, 0)\epsilon_1 x, X_2(x, y)), \\ Y^\epsilon(x, y) &= (Y_1(x, y) + Y_{2x}(0, 0)\epsilon_1 + Y_{2x}(0, 0)\epsilon_1 x + \epsilon_2 Y_2(x, y), Y_2(x, y) + \epsilon_3), \end{aligned}$$

where $\epsilon = (\epsilon_1, \epsilon_2, \epsilon_3) \in \mathbb{R}^3$ is a parameter vector. Clearly, $Z^\epsilon = Z$ for $\epsilon = (0, 0, 0)$, and $Z^\epsilon \in \Omega$. We claim that for any small neighborhood of $\epsilon = (0, 0, 0)$ there always exists ϵ_0 in the neighborhood such that Z^{ϵ_0} has a crossing limit cycle bifurcating from the non-smooth equilibrium O of Z . In fact, fixing $\epsilon_2 = \epsilon_3 = 0$ we have

$$\begin{aligned} X_1^\epsilon(0, 0) &= -X_{2x}(0, 0)\epsilon_1, & X_2^\epsilon(0, 0) &= 0, & X_{2x}^\epsilon(0, 0) &= X_{2x}(0, 0), \\ Y_1^\epsilon(0, 0) &= Y_{2x}(0, 0)\epsilon_1, & Y_2^\epsilon(0, 0) &= 0, & Y_{2x}^\epsilon(0, 0) &= Y_{2x}(0, 0), \end{aligned} \quad (5.3)$$

where $(X_1^\epsilon, X_2^\epsilon)$ and $(Y_1^\epsilon, Y_2^\epsilon)$ are the coordinates of X^ϵ and Y^ϵ respectively. So O is an invisible-invisible fold-fold point of Z^ϵ for $\epsilon_1 > 0$ and $\epsilon_2 = \epsilon_3 = 0$. Besides, all orbits of Z^ϵ near O turn around O because $X_{2x}^\epsilon(0, 0)Y_{2x}^\epsilon(0, 0) = X_{2x}(0, 0)Y_{2x}(0, 0) > 0$. Here $X_{2x}(0, 0)Y_{2x}(0, 0) > 0$ is due that $Z \in \Omega_0$ satisfies (1.3). Thus O is either a non-smooth center or a pseudo-focus of Z^ϵ for $\epsilon_1 > 0$ and $\epsilon_2 = \epsilon_3 = 0$. By the time reversal, without loss of generality next we only work with the case where all orbits of Z^ϵ near O rotate counterclockwise, namely $X_{2x}(0, 0) > 0$ and $Y_{2x}(0, 0) > 0$.

If O is a stable (resp. unstable) pseudo-focus of Z^ϵ for $\epsilon_1 > 0$ and $\epsilon_2 = \epsilon_3 = 0$, a direct application of Proposition 2.1 yields that for given $\epsilon_1 > 0$ and $\epsilon_2 = 0$ there exists $\hat{\epsilon}_3 = \hat{\epsilon}_3(\epsilon_1) > 0$ such that Z^ϵ with $\epsilon_1 > 0$, $\epsilon_2 = 0$ and $-\hat{\epsilon}_3 < \epsilon_3 < 0$ (resp. $0 < \epsilon_3 < \hat{\epsilon}_3$) admits a stable (resp. unstable) crossing limit cycle bifurcating from O . Thus, for any small neighborhood of $\epsilon = (0, 0, 0)$ we can choose some $\epsilon_0 = (\epsilon_{10}, \epsilon_{20}, \epsilon_{30})$ satisfying $\epsilon_{10} > 0$, $\epsilon_{20} = 0$ and $0 < |\epsilon_{30}| < \hat{\epsilon}_3(\epsilon_{10})$ such that Z^{ϵ_0} has a crossing limit cycle bifurcating from O , that is, the claim holds in the case that O is a pseudo-focus.

If O is a non-smooth center of Z^ϵ for $\epsilon_1 > 0$ and $\epsilon_2 = \epsilon_3 = 0$, we can obtain an upper Poincaré map P_U near O which maps a point $(x_0, 0)$ with $x_0 > 0$ to a point $(x_1, 0)$ with $x_1 < 0$, and a lower Poincaré map P_L near O which maps $(x_1, 0)$ to $(x_0, 0)$. When $\epsilon_3 = 0$ and ϵ_2 is perturbed to be $\epsilon_2 \neq 0$, it is easily verify that (5.3) still holds, i.e., O is still an invisible-invisible fold-fold point. In this case, we also can define an upper Poincaré map \tilde{P}_U near O which maps a point $(x_0, 0)$ with $x_0 > 0$ to a point $(x_1, 0)$ with $x_1 < 0$, and a lower Poincaré map \tilde{P}_L near O which maps $(x_1, 0)$ to a point $(x_2, 0)$ with $x_2 > 0$. Clearly, $P_U = \tilde{P}_U$ because X^ϵ is independent of ϵ_2 . Moreover, we can prove that $x_2 > x_0$ if $\epsilon_2 > 0$. In fact, considering the vector field Y^ϵ we define the following two equations

$$\frac{dy}{dx} = \varphi_1(x, y) := \frac{Y_2(x, y)}{Y_1(x, y) + Y_{2x}(0, 0)\epsilon_1 + Y_{2x}(0, 0)\epsilon_1 x} \quad (5.4)$$

for $\epsilon_2 = \epsilon_3 = 0$, and

$$\frac{dy}{dx} = \varphi_2(x, y) := \frac{Y_2(x, y)}{Y_1(x, y) + Y_{2x}(0, 0)\epsilon_1 + Y_{2x}(0, 0)\epsilon_1 x + \epsilon_2 Y_2(x, y)} \quad (5.5)$$

for $\epsilon_2 \neq 0$ and $\epsilon_3 = 0$. Since $Y_1^\epsilon(0, 0) = Y_{2x}(0, 0)\epsilon_1 > 0$, the denominators of $\varphi_1(x, y)$ and $\varphi_2(x, y)$ are positive in a sufficiently small neighborhood of O . Thus $\varphi_1(x, y) \geq \varphi_2(x, y)$ for $\epsilon_2 > 0$, and the equality holds only for $(x, y) = (0, 0)$. Applying the theory of differential inequality to equations (5.4) and (5.5), we obtain the solution of equation (5.4) with the initial value $(x_1, 0)$ always lies above the solution of

equation (5.5) with the initial value $(x_1, 0)$ in the half plane $y \leq 0$. So $x_2 > x_0$ if $\epsilon_2 > 0$, and then O is an unstable pseudo-focus of Z^ϵ for $\epsilon_1 > 0$, $\epsilon_2 > 0$ and $\epsilon_3 = 0$. Repeating the analysis in the last paragraph and using Proposition 2.1, for any small neighborhood of $\epsilon = (0, 0, 0)$ we can choose some $\epsilon_0 = (\epsilon_{10}, \epsilon_{20}, \epsilon_{30})$ satisfying $\epsilon_{10} > 0$, $\epsilon_{20} > 0$ and $0 < \epsilon_{30} < \hat{\epsilon}_3(\epsilon_{10}, \epsilon_{20})$ such that Z^{ϵ_0} has a crossing limit cycle bifurcating from O , that is, the claim also holds in the case that O is a non-smooth center. This, together with the last paragraph, concludes statement (1) because $Z^\epsilon \rightarrow Z$ as $\epsilon \rightarrow 0$.

Let Z_0 be the piecewise linear vector field given in (5.1). Then $Z_0 \in \Omega_1$ if either $a > 0$ or $b > 0$, and $Z_0 \in \Omega_0 \setminus \Omega_1$ if $a < 0$ and $b < 0$ as indicated in Proposition 5.1. Thus statement (2) is a direct conclusion of Propositions 5.2 and 5.3 because $Z_\epsilon^f \rightarrow Z_0$ and $Z_\epsilon^g \rightarrow Z_0$ as $\epsilon \rightarrow 0$. \square

As well known, it is a challenge objective to establish the bifurcation diagram for some bifurcations, particularly for the higher codimension bifurcations, since a higher codimension bifurcation usually consists of too many lower codimension ones. Speaking of bifurcation diagrams, we can obtain an important information from the proof of Theorem 1.3, that is, the bifurcation diagram of any vector field in Ω_0 must contain a bifurcation boundary where the codimension one pseudo-Hopf bifurcation occurs. A complete bifurcation diagram of the vector fields in Ω_0 will be left as a future work. Actually, this is an extremely complex work, since there exist many possible local phase portraits for the unperturbed vector fields as seen in Theorem 1.1, and such a bifurcation has the higher codimension.

Finally, we give the proofs of Propositions 5.2 and 5.3.

Proof of Proposition 5.2. Clearly, $Z_\epsilon^f = Z_0$ for $\epsilon = 0$. The rest of this proof is completed by the following four steps.

Step 1. The upper Poincaré map P_U . Because of $a \neq 0$, we can choose $\epsilon_1 > 0$ such that $\text{sign} a = \text{sign}(a - \epsilon)$ for $0 < \epsilon < \epsilon_1$. In this case, X_ϵ^f has a unique equilibrium $E_X := (0, \epsilon/(a - \epsilon))$, which is a linear center if $a - \epsilon < 0$ and a linear saddle if $a - \epsilon > 0$.

When E_X is linear center, i.e., $a - \epsilon < 0$, it lies in the lower half plane $y < 0$ because of $0 < \epsilon < \epsilon_1$, and then it is not a real equilibrium for Z_ϵ . From the center dynamics and the direction of the vector field X_ϵ^f on the x -axis, it follows that the orbit of X_ϵ^f with $0 < \epsilon < \epsilon_1$ starting from $(x_0, 0)$ with $x_0 > 0$ enters into $y > 0$, and reaches again the x -axis at a point $(x_1, 0)$ with $x_1 < 0$ as t increases.

When E_X is a linear saddle, i.e., $a - \epsilon > 0$, it lies in the upper half plane $y > 0$ because of $0 < \epsilon < \epsilon_1$, and the stable and unstable manifolds of it lie in

$$\left\{ (x, y) \in \mathbb{R}^2 \mid x \neq 0, y = -\frac{x}{\sqrt{a - \epsilon}} + \frac{\epsilon}{a - \epsilon} \right\}, \quad \left\{ (x, y) \in \mathbb{R}^2 \mid x \neq 0, y = \frac{x}{\sqrt{a - \epsilon}} + \frac{\epsilon}{a - \epsilon} \right\},$$

respectively. Thus the stable manifold intersects the x -axis at $(\epsilon/\sqrt{a - \epsilon}, 0)$, and the unstable manifold intersects the x -axis at $(-\epsilon/\sqrt{a - \epsilon}, 0)$. Together with the direction of the vector field X_ϵ^f on $\{(x, 0) \in \mathbb{R}^2 : -\epsilon/\sqrt{a - \epsilon} < x < \epsilon/\sqrt{a - \epsilon}\}$, we get that the orbit of X_ϵ^f with $0 < \epsilon < \epsilon_1$ starting from $(x_0, 0)$ with $0 < x_0 < \epsilon/\sqrt{a - \epsilon}$ enters into $y > 0$ and reaches again the x -axis at a point $(x_1, 0)$ with $-\epsilon/\sqrt{a - \epsilon} < x_1 < 0$ as t increases.

According to the last two paragraphs, we can construct an upper Poincaré map P_U as $x_1 = P_U(x_0, \epsilon)$, which is defined for $0 < x_0 < \varpi_u(\epsilon)$ and $0 < \epsilon < \epsilon_1$, where

$$\varpi_u(\epsilon) = \begin{cases} +\infty & \text{when } E_X \text{ is a center, i.e., } a - \epsilon < 0, \\ \epsilon / \sqrt{a - \epsilon} & \text{when } E_X \text{ is a saddle, i.e., } a - \epsilon > 0. \end{cases} \quad (5.6)$$

Furthermore, calculating the first integral H_X^f of X_ϵ^f we get

$$H_X^f(x, y) = \frac{1}{2}x^2 - \frac{a - \epsilon}{2}y^2 + \epsilon y,$$

so that $P_U(x_0, \epsilon)$ satisfies $H_X^f(x_0, 0) = H_X^f(P_U(x_0), 0)$, i.e.,

$$P_U(x_0, \epsilon) = -x_0 \quad \text{for } 0 < x_0 < \varpi_u(\epsilon) \text{ and } 0 < \epsilon < \epsilon_1. \quad (5.7)$$

Step 2. The lower Poincaré map P_L . Since $b \neq 0$, there exists $\epsilon_2 > 0$ such that $\text{sign} b = \text{sign}(b - \epsilon)$ for $0 < \epsilon < \epsilon_2$. Throughout this step, ϵ_2 can be reduced if necessary. Consider the function

$$F(x, \epsilon) = x + \epsilon \frac{\partial f(x, \epsilon)}{\partial x}.$$

Due to $F(0, 0) = 0$ and $F_x(0, 0) = 1$, by the Implicit Function Theorem there exists a function $x(\epsilon)$ defined for $0 < \epsilon < \epsilon_2$ such that $x(0) = 0$ and $F(x(\epsilon), \epsilon) = 0$. In addition, $x(\epsilon)$ is given by

$$x(\epsilon) = (-1)^{m+1} \epsilon \prod_{i=1}^m \left(\frac{i\epsilon}{m} \right)^2 + O(\epsilon^{2m+2}) = (-1)^{m+1} \frac{(m!)^2}{m^{2m}} \epsilon^{2m+1} + O(\epsilon^{2m+2}). \quad (5.8)$$

By the definition of invisible fold point, $(x(\epsilon), 0)$ is an invisible fold point of Y_ϵ^f for $0 < \epsilon < \epsilon_2$. Combining the direction of Y_ϵ^f on the x -axis, we know that the orbit of Y_ϵ^f near $(x(\epsilon), 0)$ starting from a point $(x_1, 0)$ with $x_1 < x(\epsilon)$ evolves in $y < 0$ until it reaches the x -axis at a point $(x_2, 0)$ with $x_2 > x(\epsilon)$ again. In this case, we can define a lower Poincaré map P_L as $x_2 = P_L(x_1, \epsilon)$ for $x_1 < x(\epsilon)$ closed to $x(\epsilon)$ and $0 < \epsilon < \epsilon_2$. Since the first integral of Y_ϵ^f is

$$H_Y^f(x, y) = \frac{1}{2}x^2 + \epsilon f(x, \epsilon) - \frac{b - \epsilon}{2}y^2 - \epsilon y,$$

$P_L(x_1, \epsilon)$ satisfies

$$\frac{1}{2}x_1^2 + \epsilon f(x_1, \epsilon) = \frac{1}{2}P_L(x_1, \epsilon)^2 + \epsilon f(P_L(x_1, \epsilon), \epsilon). \quad (5.9)$$

Next we precisely determine the definition domain of P_L . Notice that $E_Y := (x(\epsilon), -\epsilon/(b - \epsilon))$ is an equilibrium of Y_ϵ^f for $0 < \epsilon < \epsilon_2$. Calculating the eigenvalues of the Jacobian matrix of Y_ϵ^f at E_Y , we have that E_Y is of focus type if $b - \epsilon < 0$ from [31, Theorem 5.1], and a saddle if $b - \epsilon > 0$ from [31, Theorem 4.4].

When E_Y is of focus type, i.e., $b - \epsilon < 0$, it lies in the upper half plane $y > 0$ because of $\epsilon > 0$. Moreover, O is a linear center of Y_ϵ^f for $\epsilon = 0$ due to $\text{sign} b = \text{sign}(b - \epsilon)$ for $0 < \epsilon < \epsilon_2$. Thus $\epsilon_2 > 0$ can be reduced such that $P_L(x_1, \epsilon)$ is defined for $-1 < x_1 < x(\epsilon)$ and $0 < \epsilon < \epsilon_2$.

When E_Y is saddle, i.e., $b - \epsilon > 0$, it lies in the lower half plane $y < 0$ because of $\epsilon > 0$. E_Y has one stable (resp. unstable) manifold intersecting the x -axis. Let $(x_s, 0)$ (resp. $(x_u, 0)$) be the intersection between the stable (resp. unstable) manifold and the x -axis. Then $H_Y^f(x_u, 0) = H_Y^f(x_s, 0) = H_Y^f(E_Y)$, i.e.,

$$\begin{aligned} \frac{1}{2}(x_u)^2 + \epsilon f(x_u, \epsilon) &= \frac{1}{2}(x_s)^2 + \epsilon f(x_s, \epsilon) = \frac{1}{2}x(\epsilon)^2 + \epsilon f(x(\epsilon), \epsilon) - \frac{b - \epsilon}{2} \left(\frac{-\epsilon}{b - \epsilon} \right)^2 - \epsilon \left(\frac{-\epsilon}{b - \epsilon} \right) \\ &= \frac{\epsilon^2}{2(b - \epsilon)} + O(\epsilon^3), \end{aligned} \quad (5.10)$$

where the last equality is due to (5.8). Solving (5.10) we get

$$x_u = \frac{\epsilon}{\sqrt{b - \epsilon}} + O(\epsilon^2), \quad x_s = -\frac{\epsilon}{\sqrt{b - \epsilon}} + O(\epsilon^2)$$

for $0 < \epsilon < \epsilon_2$ by $x_u > x_s$. Consequently, $P_L(x_1, \epsilon)$ is defined for $x_s < x_1 < x(\epsilon)$ and $0 < \epsilon < \epsilon_2$. Moreover, $x(\epsilon) < P_L(x_1) < x_u$.

In conclusion, we take the definition domain of $P_L(x_1, \epsilon)$ as $\varpi_l(\epsilon) < x_1 < x(\epsilon)$, where

$$\varpi_l(\epsilon) = \begin{cases} -1 & \text{when } E_Y \text{ is of focus type, i.e., } b - \epsilon < 0, \\ x_s = -\frac{\epsilon}{\sqrt{b - \epsilon}} + O(\epsilon^2) & \text{when } E_Y \text{ is a saddle, i.e., } b - \epsilon > 0. \end{cases} \quad (5.11)$$

Step 3. The full Poincaré map P . Take $\epsilon_0 = \min\{\epsilon_1, \epsilon_2\}$ and $\varpi(\epsilon) = |x(\epsilon)|$. In what follows $\epsilon_0 > 0$ can be reduced if necessary. Let

$$I(\epsilon) = (\varpi(\epsilon), \min\{-\varpi_l(\epsilon), \varpi_u(\epsilon)\}).$$

By (5.8) and the definitions of $\varpi_l(\epsilon)$ and $\varpi_u(\epsilon)$ the interval $I(\epsilon)$ is non-empty for $0 < \epsilon < \epsilon_0$. According to the last two steps, we construct P as the composition $P(x_0, \epsilon) = P_L(P_U(x_0, \epsilon), \epsilon)$ for $x_0 \in I(\epsilon)$ and $0 < \epsilon < \epsilon_0$. Hence, a fixed point of $P(x_0, \epsilon)$ in the interval $I(\epsilon)$ corresponds to a crossing periodic orbit of Z_ϵ^f . Furthermore, from (5.2), (5.7) and (5.9) the map $P(x_0, \epsilon)$ satisfies

$$\frac{1}{2}x_0^2 + \epsilon f(-x_0, \epsilon) = \frac{1}{2}P(x_0, \epsilon)^2 + \epsilon f(P(x_0, \epsilon), \epsilon),$$

i.e.,

$$\frac{1}{2}x_0^2 - \epsilon x_0 \prod_{i=1}^m \left(x_0^2 - \left(\frac{i\epsilon}{m} \right)^2 \right) = \frac{1}{2}P(x_0, \epsilon)^2 + \epsilon P(x_0, \epsilon) \prod_{i=1}^m \left(P(x_0, \epsilon)^2 - \left(\frac{i\epsilon}{m} \right)^2 \right). \quad (5.12)$$

Step 4. Crossing limit cycles. Now we study the crossing limit cycles of Z_ϵ^f using the Poincaré map P . Since $0 < |a| \leq 1/2$ and $0 < |b| \leq 1/2$ as assumed in Proposition 5.2, we have $\min\{-\varpi_l(\epsilon), \varpi_u(\epsilon)\} > \sqrt{2}\epsilon + O(\epsilon^2)$ for $0 < \epsilon < \epsilon_0$, so that $i\epsilon/m < \min\{-\varpi_l(\epsilon), \varpi_u(\epsilon)\}$ for all $i = 1, 2, \dots, m$ and $0 < \epsilon < \epsilon_0$. On the other hand, it follows from (5.8) that $i\epsilon/m > |x(\mu)|$, i.e., $i\epsilon/m > \varpi(\epsilon)$, for all $i = 1, 2, \dots, m$ and $0 < \epsilon < \epsilon_0$. So $i\epsilon/m \in I(\epsilon)$ for all $i = 1, 2, \dots, m$ and $0 < \epsilon < \epsilon_0$. Associate with (5.12), x_0 is a fixed point of P in $I(\epsilon)$ if and only if $x_0 = i\epsilon/m$, which implies that Z_ϵ^f has exactly m isolated and nested crossing periodic orbits, namely crossing limit cycles. Moreover, these crossing limit cycles intersect the positive x -axis at $(i\epsilon/m, 0)$, $i = 1, 2, \dots, m$. Using the first integrals H_X^f and H_Y^f , we get that the m limit cycles obey the algebraic curves Γ_i^+ and Γ_i^- defined in Proposition 5.2, $i = 1, 2, \dots, m$.

Finally, in order to determine the hyperbolicity and stability of Γ_i , $i = 1, 2, \dots, m$, taking the derivative with respect to x_0 for (5.12), we have

$$\begin{aligned} \frac{dP}{dx_0} \left(\frac{i\epsilon}{m} \right) &= \frac{\frac{i\epsilon}{m} - 2\epsilon \left(\frac{i\epsilon}{m} \right)^2 \prod_{k=1, k \neq i}^m \left(\left(\frac{i\epsilon}{m} \right)^2 - \left(\frac{k\epsilon}{m} \right)^2 \right)}{\frac{i\epsilon}{m} + 2\epsilon \left(\frac{i\epsilon}{m} \right)^2 \prod_{k=1, k \neq i}^m \left(\left(\frac{i\epsilon}{m} \right)^2 - \left(\frac{k\epsilon}{m} \right)^2 \right)} \\ &= \frac{1 - 2\epsilon^{2m} \left(\frac{i}{m} \right) \prod_{k=1, k \neq i}^m \left(\left(\frac{i}{m} \right)^2 - \left(\frac{k}{m} \right)^2 \right)}{1 + 2\epsilon^{2m} \left(\frac{i}{m} \right) \prod_{k=1, k \neq i}^m \left(\left(\frac{i}{m} \right)^2 - \left(\frac{k}{m} \right)^2 \right)}. \end{aligned}$$

Thus $0 < \frac{dP}{dx_0} \left(\frac{i\epsilon}{m} \right) < 1$ (resp. > 1) if $m - i$ is even (resp. odd), that is, Γ_i is hyperbolic and stable (resp. unstable) if $m - i$ is even (resp. odd). The proof of Proposition 5.2 is finished. \square

Proof of Proposition 5.3. Obviously, $Z_\epsilon^g = Z_0$ for $\epsilon = 0$. The study of the bifurcated crossing limit cycles is extremely similar to the proof of Proposition 5.2. So we neglect some details. In fact, comparing the vector fields $Z_\epsilon^f = (X_\epsilon^f, Y_\epsilon^f)$ and $Z_\epsilon^g = (X_\epsilon^g, Y_\epsilon^g)$, we see $X_\epsilon^f = X_\epsilon^g$, so that we get the same upper Poincaré map

$$P_U(x_0, \epsilon) = -x_0 \quad \text{for } 0 < x_0 < \varpi_u(\epsilon) \text{ and } 0 < \epsilon < \epsilon_1. \quad (5.13)$$

as defined in (5.7). Here $\varpi_u(\epsilon)$ is given in (5.6). Besides, Y_ϵ^f and Y_ϵ^g have the same expression except that the function f is replaced by g . With the replacement, O is an invisible fold point of Y_ϵ^g , and Y_ϵ^g has $(0, -\epsilon/(b - \epsilon))$ as an equilibrium, which is of focus type if $b - \epsilon < 0$ and a saddle if $b - \epsilon > 0$. Therefore, carrying out a similar argument to Step 2 in the proof of Proposition 5.2, we can choose some $\epsilon_2 > 0$ and define a lower Poincaré map $P_L(x_1, \epsilon)$ for $\tilde{\varpi}_l(\epsilon) < x_1 < 0$ and $0 < \epsilon < \epsilon_2$, where

$$\tilde{\varpi}_l(\epsilon) = \begin{cases} -1 & \text{when } (0, -\epsilon/(b - \epsilon)) \text{ is of focus type, i.e., } b - \epsilon < 0, \\ \tilde{x}_s = -\frac{\epsilon}{\sqrt{b - \epsilon}} + O(\epsilon^2) & \text{when } (0, -\epsilon/(b - \epsilon)) \text{ is a saddle, i.e., } b - \epsilon > 0, \end{cases}$$

and $(\tilde{x}_s, 0)$ is the intersection between the stable manifold of $(0, -\epsilon/(b - \epsilon))$ and the negative x -axis. Notice that $\varpi_l(\epsilon)$ defined in (5.11) and $\tilde{\varpi}_l(\epsilon)$ are the same in the sense of neglecting the higher order terms. Since the first integral of Y_ϵ^g is

$$H_Y^g(x, y) = \frac{1}{2}x^2 + \epsilon g(x, \epsilon) - \frac{b - \epsilon}{2}y^2 - \epsilon y,$$

$P_L(x_1, \epsilon)$ satisfies

$$\frac{1}{2}x_1^2 + \epsilon g(x_1, \epsilon) = \frac{1}{2}P_L(x_1, \epsilon)^2 + \epsilon g(P_L(x_1, \epsilon), \epsilon). \quad (5.14)$$

The above analysis allows us to define a full Poincaré map $P(x_0, \epsilon) = P_L(P_U(x_0, \epsilon), \epsilon)$ for $x_0 \in \tilde{I}(\epsilon)$ and $0 < \epsilon < \epsilon_0$, where

$$\tilde{I}(\epsilon) = (0, \min\{-\tilde{\varpi}_l(\epsilon), \varpi_u(\epsilon)\}), \quad \epsilon_0 = \min\{\epsilon_1, \epsilon_2\}.$$

Hence, a fixed point of $P(x_0, \epsilon)$ in the interval $\tilde{I}(\epsilon)$ corresponds to a crossing periodic orbit of Z_ϵ^g . Furthermore, from (5.13) and (5.14) it follows that $P(x_0, \epsilon)$ satisfies

$$\frac{1}{2}x_0^2 + \epsilon g(-x_0, \epsilon) = \frac{1}{2}P(x_0, \epsilon)^2 + \epsilon g(P(x_0, \epsilon), \epsilon),$$

i.e.,

$$\frac{1}{2}x_0^2 = \frac{1}{2}P(x_0, \epsilon)^2 + \epsilon e^{-1/P(x_0, \epsilon)} \sin\left(\frac{\pi \epsilon}{P(x_0, \epsilon)}\right). \quad (5.15)$$

Now we study the fixed points of $P(x_0, \epsilon)$ in $\tilde{I}(\epsilon)$. Since $0 < |a| \leq 1/2$ and $0 < |b| \leq 1/2$ as assumed in Proposition 5.3, $\min\{-\tilde{\omega}_l(\epsilon), \tilde{\omega}_u(\epsilon)\} > \sqrt{2}\epsilon + O(\epsilon^2)$ for $0 < \epsilon < \epsilon_0$, so that $\epsilon/i \in \tilde{I}(\epsilon)$ for $i \in \mathbb{N}^+$ and $0 < \epsilon < \epsilon_0$. Here ϵ_0 can be reduced if necessary. As a consequence, by (5.15) we get that x_0 is a fixed point of $P(x_0, \epsilon)$ in $\tilde{I}(\epsilon)$ if and only if $x_0 = \epsilon/i$, $i \in \mathbb{N}^+$. This means that Z_ϵ^g has infinitely many nested crossing limit cycles. Moreover, these crossing limit cycles intersect the positive x -axis at $(\epsilon/i, 0)$, $i \in \mathbb{N}^+$. Using the first integrals, we get that these crossing limit cycles obey the algebraic curves Θ_i^+ and Θ_i^- defined in Proposition 5.3, $i \in \mathbb{N}^+$.

Finally, computing the derivative of $P(x_0, \epsilon)$ with respect to x_0 for (5.15), we get

$$\frac{dP}{dx_0}\left(\frac{\epsilon}{i}\right) = \frac{\epsilon/i}{\epsilon/i - \pi i^2 e^{-i/\epsilon} \cos(\pi i)}.$$

So $0 < \frac{dP}{dx_0}\left(\frac{\epsilon}{i}\right) < 1$ (resp. > 1) if i is odd (resp. even), which implies that Θ_i is hyperbolic and stable (resp. unstable) if i is odd (resp. even). This ends the proof of Proposition 5.3. \square

References

- [1] M. di Bernardo, C. J. Budd, A. R. Champneys, P. Kowalczyk, *Piecewise-Smooth Dynamical systems: Theory and Applications*, Applied Mathematical Sciences, Vol.163 (Springer Verlag, London), 2008.
- [2] C. A. Buzzi, T. Carvalho, R. D. Euzébio, On Poincaré-Bendixson theorem and non-trivial minimal sets in planar nonsmooth vector fields, *Publ. Mat.* **62** (2018), 113-131.
- [3] C. A. Buzzi, T. de Carvalho, M. A. Teixeira, Birth of limit cycles bifurcating from a nonsmooth center, *J. Math. Pures Appl.* **102** (2014), 36-47.
- [4] Q. Cao, M. Wiercigroch, E. E. Pavlovskaya, C. Grebogi, J. M. T. Thompson, Archetypal oscillator for smooth and discontinuous dynamics, *Phys. Rev. E* **74** (2006), 046218.
- [5] T. Carvalho, J. L. Cardoso, D. J. Tonon, Canonical forms for codimension one planar piecewise smooth vector fields with sliding region, *J. Dyn. Diff. Equat.* **30** (2018), 1899-1920.
- [6] T. Carvalho, D. J. Tonon, Normal forms for codimension one planar piecewise smooth vector fields, *Int. J. Bifurc. Chaos* **24** (2014), 1450090.
- [7] J. Castillo, J. Llibre, F. Verduzco, The pseudo-Hopf bifurcation for planar discontinuous piecewise linear differential systems, *Nonlinear Dyn.* **90** (2017), 1829-1840.
- [8] H. Chen, S. Duan, Y. Tang, J. Xie, Global dynamics of a mechanical system with dry friction, *J. Differential Equa.* **265** (2018), 5490-5519.
- [9] X. Chen, V. G. Romanovski, W. Zhang, Degenerate Hopf bifurcations in a family of FF-type switching systems, *J. Math. Anal. Appl.* **432** (2015), 1058-1076.
- [10] X. Chen, W. Zhang, Normal form of planar switching systems, *Disc. Cont. Dyn. Syst.* **36** (2016), 6715-6736.
- [11] B. Coll, A. Gasull, R. Prohens, Degenerate Hopf bifurcation in discontinuous planar systems, *J. Math. Anal. Appl.* **253** (2001), 671-690.

- [12] L. P. C. da Cruz, D. D. Novaes, J. Torregrosa, New lower bound for the Hilbert number in piecewise quadratic differential systems, *J. Differential Equa.* **266** (2019), 4170-4203.
- [13] A. F. Filippov, *Differential Equations with Discontinuous Righthand Sides*, Kluwer Academic Publishers, Dordrecht, 1988.
- [14] F. Giannakopoulos, K. Pliete, Planar systems of piecewise linear differential equations with a line of discontinuity, *Nonlinearity* **14** (2001), 1611-1632.
- [15] P. Glendinning, Classification of boundary equilibrium bifurcations in planar Filippov systems, *Chaos* **26** (2016), 013108.
- [16] M. Guardia, T. M. Seara, M. A. Teixeira, Generic bifurcations of low codimension of planar Filippov systems, *J. Differential Equa.* **250** (2011), 1967-2023.
- [17] J. K. Hale, *Ordinary Differential Equations*, Kreiger, 1980.
- [18] M. Han, W. Zhang, On Hopf bifurcation in non-smooth planar systems, *J. Differential Equa.* **248** (2010), 2399-2416.
- [19] P. Hartman, On the local linearization of differential equations, *Proc. Amer. Math. Soc.* **14** (1963), 568-573.
- [20] P. Hartman, *Ordinary Differential Equations*, John Wiley & Sons, New York, 1964.
- [21] K. U. Kristiansen, S. J. Hogan, Regularizations of two-fold bifurcations in planar piecewise smooth systems using blowup, *SIAM J. Appl. Dyn. Syst.* **14** (2015), 1731-1786.
- [22] T. Küpper, S. Moritz, Generalized Hopf bifurcation for non-smooth planar systems, *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **359** (2001), 2483-2496.
- [23] Yu. A. Kuznetsov, *Elements of Applied Bifurcation Theory* (Second Edition), Springer, New York, 1998.
- [24] Yu. A. Kuznetsov, S. Rinaldi, A. Gragnani, One parameter bifurcations in planar Filippov systems, *Int. J. Bifurc. Chaos* **13** (2003), 2157-2188.
- [25] F. Liang, M. Han, Degenerate Hopf bifurcation in nonsmooth planar systems, *Int. J. Bifurc. Chaos* **22** (2012), 1250057.
- [26] C. B. Reves, J. Larrosa, T. M. Seara, Regularization around a generic codimension one fold-fold singularity, *J. Differential Equa.* **265** (2018), 1761-1838.
- [27] S. Tang, J. Liang, Y. Xiao, R. A. Cheke, Sliding bifurcations of Filippov two stage pest control models with economic thresholds, *SIAM J. Appl. Math.* **72** (2012), 1061-1080.
- [28] A. Wang, Y. Xiao, A Filippov system describing media effects on the spread of infectious diseases, *Nonlinear Anal.: Hybrid Syst.* **11** (2014), 84-97.
- [29] L. Wei, X. Zhang, Normal form and limit cycle bifurcation of piecewise smooth differential systems with a center, *J. Differential Equa.* **261** (2016), 1399-1428.
- [30] J. Yang, M. Han, On Hopf bifurcations of piecewise planar Hamiltonian systems, *J. Differential Equa.* **250** (2011), 1026-1051.
- [31] Z. Zhang, T. Ding, W. Huang, Z. Dong, *Qualitative Theory of Differential Equations*, Science Publisher, 1985 (in Chinese); Transl. Math. Monogr., vol. 101, Amer. Math. Soc., Providence, RI, 1992.
- [32] Y. Zou, T. Küpper, W. J. Beyn, Generalized Hopf bifurcation for planar Filippov systems continuous at the origin, *J. Nonlinear Sci.* **16** (2006), 159-177.