

STABILITY OF SET-VALUED GENERALIZED ADDITIVE CAUCHY FUNCTIONAL EQUATION

GANG LU*, MINGZHU MU, YUANFENG JIN*, AND CHOONKIL PARK

ABSTRACT. The main purpose of this paper is to determine the solution of generalized convex set-valued mappings satisfying certain functional equation. Some conclusions of stability of set-valued functional equations are obtained.

1. INTRODUCTION AND PRELIMINARIES

The goal of this paper is to characterize set-valued solutions of a generalized additive Cauchy functional equation. In 1992, Rassias and Tabor [15] asked whether the functional

$$f(ax + by + c) = Af(x) + Bf(y) + C$$

with $abAB \neq 0$ is stable in the sense of Hyers, Ulam, and Rassias. And the functional equation is called *generalized additive Cauchy functional equation*.

Badea [1] answered this question of Rassias and Tabor for the case when $c = C = 0$, $a = A$ and $b = B$. As Gajda [2] extended the result of Badea [1] also modified the result of the previous theorem by using a similar method. Jung [4] investigated the stability problem in complex Banach space for a generalized additive Cauchy functional equation

$$f\left(x_0 + \sum_{j=1}^m a_j x_j\right) = \sum_{i=1}^m b_i f\left(\sum_{j=1}^m a_{ij} x_j\right).$$

An interesting question concerning set-valued mappings that satisfy some functional inclusions is the existence of solutions verifying certain properties. In the stability theory of functional equations, it is important to find solutions of set-valued mappings that are also solutions of certain functional equations. And more results were proved for set-valued mappings satisfying general additive Cauchy of the type

$$f(ax + by + c) \subseteq Af(x) + Bf(y) + C \quad (1.1)$$

or

$$Af(x) + Bf(y) \subseteq f(ax + by + c) + C \quad (1.2)$$

where $f : X \rightarrow 2^Y$, X and Y are real vector spaces, $a, b, A, B \in \mathbb{R}$, $c \in X$ and $C \in 2^Y$. By 2^Y we denote the collection of nonempty subsets of Y . Nikodam and Popa [8] considered the general solution of set-valued mappings satisfying linear relation, which

2010 *Mathematics Subject Classification.* Primary 54C60, 39B52, 47H04, 49J54.

Key words and phrases. Hyers-Ulam stability; additive set-valued functional equation; closed and convex subset.

*Corresponding authors.

can be regarded as a generalization of the additive single-valued functional equation with $a = A, b = B$ and $C = c = 0$. Lu *et al.* [5, 10] investigated the approximation of some set-valued functional equations with $C = c = 0$. Popa [11] determined conditions for which a set-valued map that verifies a relation of the form (1.1) admits a unique solution $f : X \rightarrow Y$ that satisfies

$$f(ax + by + c) \subseteq Af(x) + Bf(y),$$

for all $x, y \in X$. The case of equality of form (1.1) was also studied in [12]. The similar results [3, 9] were obtained in the case of (1.2).

Assume that Y is a topological vector space satisfying the T_0 -separation axiom. For real numbers s, t and sets $A, B \subset Y$ we put $sA + tB := \{y \in Y; y = sa + tb, a \in A, b \in B\}$. Suppose that the space 2^Y of all subsets of Y is endowed with the Hausdorff topology (see [14]). A set-valued function $F : X \rightarrow 2^Y$ is said to be additive if it satisfies the Cauchy functional equation $F(x_1 + x_2) = F(x_1) + F(x_2)$, $x_1, x_2 \in X$, where X is a semigroup.

In this paper, we characterize set-valued solutions of the following functional equation

$$f(ax + by + c) = AG(x) + BH(y) + C \quad (1.3)$$

for all $x, y \in X$, where a, b, A and B are positive real numbers, c and C are fixed bounded convex sets.

The family of all closed and convex subsets of Y will be denoted by $CC(Y)$, and the sets of all real numbers, rational numbers and positive integers are denoted by $\mathbb{R}, \mathbb{Q}, \mathbb{N}$, respectively.

Lemma 1.1. [6] *Let λ and μ be real numbers. If A and B are nonempty subsets of a real vector space X , then*

$$\begin{aligned} \lambda(A + B) &= \lambda A + \lambda B, \\ (\lambda + \mu)A &\subseteq \lambda A + \mu B. \end{aligned}$$

Moreover, if A is a convex set and $\lambda, \mu \geq 0$, then we have

$$(\lambda + \mu)A = \lambda A + \mu A.$$

Lemma 1.2. [13] *Let A, B be subsets of Y and assume that B is closed and convex. If there exists a bounded and nonempty set $C \subset Y$ such that $A + C \subset B + C$, then $A \subset B$.*

The following lemmas are rather known and can be easily verified. The proofs of them can be found in [6, 7].

Lemma 1.3. *If $\{A_n\}_{n \in \mathbb{N}}$ and $\{B_n\}_{n \in \mathbb{N}}$ are decreasing sequences of compact subsets of Y , then $\bigcap_{n \in \mathbb{N}} (A_n + B_n) = \bigcap_{n \in \mathbb{N}} A_n + \bigcap_{n \in \mathbb{N}} B_n$.*

Lemma 1.4. *If $\{A_n\}_{n \in \mathbb{N}}$ is a decreasing sequence of compact subsets of Y , then $A_n \rightarrow \bigcap_{n \in \mathbb{N}} A_n$.*

Lemma 1.5. *If A is a bounded subset of Y and $(s_n)_{n \in \mathbb{N}}$ is a real sequence converging to an $s \in \mathbb{R}$, then $s_n A \rightarrow sA$.*

Lemma 1.6. *If $A_n \rightarrow A$ and $B_n \rightarrow B$, then $A_n + B_n \rightarrow A + B$.*

Lemma 1.7. *If $A_n \rightarrow A$ and $A_n \rightarrow B$, then $clA = clB$.*

2. SET-VALUED SOLUTION OF THE PEXIDER FUNCTIONAL EQUATION

In this section, we give the solution of the following set-valued functional equation

$$F(ax + by + c) = AG(x) + BH(y) + C \quad (2.1)$$

for all $x, y \in X$.

Theorem 2.1. *Assume that $(X, +)$ is a vector space and Y is a T_0 topological vector space. If set-valued functions $F : X \rightarrow CC(Y)$, $G : X \rightarrow CC(Y)$ and $H : X \rightarrow CC(Y)$ satisfy the functional equation*

$$F(ax + by + c) = AG(x) + BH(y) + C \quad (2.2)$$

for all $x, y \in X$, where a, b, A and B are positive real numbers, c and C are fixed bounded convex sets. Then there exist an additive set-valued mapping $F_0 : X \rightarrow CC(Y)$ and sets $\alpha, \beta \in CC(Y)$ such that

$$F(x + c) = F_0\left(\frac{x}{a}\right) + K, \quad AG(x) = F_0(x) + A\alpha \quad \text{and} \quad BH(x) = F_0\left(\frac{bx}{a}\right) + B\beta$$

for all $x \in X$.

Proof. First, assume that $0 \in G(0)$ and $0 \in H(0)$. Then, for all $x, y \in X$, we have

$$\begin{aligned} AG(x) + BH(0) + C &= F(2ax + c) = F\left(ax + b\frac{ax}{b} + c\right) \\ &= AG(x) + BH\left(\frac{ax}{b}\right) + C \\ &\subset AG(x) + AG(0) + BH\left(\frac{ax}{b}\right) + C = AG(x) + F(ax + c) \\ &= AG(x) + AG(x) + BH(0) + C. \end{aligned}$$

By Lemma 1.2, we get $G(2x) \subset 2G(x)$, which implies that the sequence $(2^{-n}G(2^n x))_{n \in \mathbb{N}}$ is decreasing. Put $F_0(x) := \bigcap_{n \in \mathbb{N}} 2^{-n}G(2^n x)$, $x \in X$. It is clear that $F_0(x) \in CC(Y)$ for all $x \in X$. Similarly, we get $H(2x) \subset 2H(x)$ for all $x \in X$. Then

$$AG(x) + BH(0) + C = F(ax + c) = AG(0) + BH\left(\frac{ax}{b}\right) + C.$$

By Lemma 1.3, we obtain that $AF_0(x) = B \bigcap 2^{-n}H\left(\frac{2^n ax}{b}\right)$ for all $x \in X$. From (2.2), we get $F(a2^n x + c) = AG(2^n x) + BH(0) + C$, $n \in \mathbb{N}$ and so

$$AF_0(x) = \bigcap_{n \in \mathbb{N}} 2^{-n}F(a2^n x + c)$$

for all $x \in X$.

Hence, using once more Lemma 1.3, we get

$$\begin{aligned}
AF_0(x_1 + x_2) &= \bigcap_{n \in \mathbb{N}} 2^{-n} F(a2^n x_1 + a2^n x_2 + c) \\
&= \bigcap_{n \in \mathbb{N}} 2^{-n} \left(AG(2^n x_1) + BH\left(\frac{2^n a x_2}{b}\right) + C \right) \\
&= \bigcap_{n \in \mathbb{N}} 2^{-n} AG(2^n x_1) + \bigcap_{n \in \mathbb{N}} 2^{-n} BH\left(\frac{2^n a x_2}{b}\right) + \bigcap_{n \in \mathbb{N}} 2^{-n} C \\
&= AF_0(x_1) + AF_0(x_2), x_1, x_2 \in X,
\end{aligned}$$

which means that the set-valued function F_0 is additive.

Now observe that

$$F(nax + c) + (n - 1)AG(0) = F(ax + c) + (n - 1)AG(x) \quad (2.3)$$

for all $x \in X$ and $n \in \mathbb{N}$. Indeed, for $n = 1$ the equality is trivial. Assume that it holds for a natural number k . Then, in virtue of (2.2), we obtain

$$\begin{aligned}
F((k + 1)ax + c) + kAG(0) &= AG(x) + BH\left(\frac{akx}{b}\right) + C + kAG(0) \\
&= AG(x) + AG(0) + BH\left(\frac{akx}{b}\right) + C + kAG(0) - AG(0) \\
&= AG(x) + F(akx + c) + (k - 1)AG(0) \\
&= AkG(x) + F(ax + c)
\end{aligned}$$

which proves that (2.3) holds for $n = k + 1$. Thus, by induction, it holds for all $n \in \mathbb{N}$. In particular, we have

$$F(2^n ax + c) + (2^n - 1)AG(0) = F(ax + c) + (2^n - 1)AG(x),$$

and so

$$2^{-n} F(2^n ax + c) + (1 - 2^{-n})AG(0) = 2^{-n} F(ax + c) + (1 - 2^{-n})AG(x)$$

for all $x \in X$. By Lemma 1.4, $2^{-n} F(2^n ax + c) \rightarrow \bigcap_{n \in \mathbb{N}} 2^{-n} F(2^n ax + c) = F_0(x)$.

On the other hand, by Lemma 1.5, $1 - 2^{-n} G(0) \rightarrow G(0)$, $2^{-n} F(ax + c) \rightarrow \{0\}$ and $(1 - 2^{-n})G(x) \rightarrow G(x)$. Thus, using Lemmas 1.6 and 1.7, we get $cl[F_0(x) + AG(0)] = clAG(x)$, whence $AG(x) = F_0(x) + AG(0)$ for all $x \in X$. Similarly, we can obtain $BH(x) = F_0\left(\frac{bx}{a}\right) + BH(0)$, $x \in X$. Let $\alpha := G(0)$ and $\beta := H(0)$. Then $AG(x) = F_0(x) + A\alpha$ and $BH(x) = F_0\left(\frac{bx}{a}\right) + B\beta$ for all $x \in X$. Moreover $F(ax + c) = AG(x) + BH(0) + C = F_0(x) + AG(0) + BH(0) + C = F_0(x) + K$, $x \in X$, where $K = A\alpha + B\beta + F_0(0) + C$. This finishes our proof in the case that $0 \in G(0)$ and $0 \in H(0)$.

In the opposite case, fix arbitrarily points $a \in G(0)$ and $b \in H(0)$, and consider the set-valued mappings $F_1, G_1, H_1 : X \rightarrow CC(Y)$ defined by $G_1(x) := G(x) - a$ and $H_1 := H(x) - b$, $x \in X$. These set-valued mappings satisfy the equation (2.2) and moreover $0 \in G_1(0)$ and $0 \in H_1(0)$. Therefore, by what we have discussed previously, we can get the same result. This completes the proof. \square

CONCLUSION

We have determined the solution of generalized convex set-valued mappings satisfying certain functional equation. Some conclusions of stability of set-valued functional equations have been obtained.

DECLARATIONS

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The authors equally conceived of the study, participated in its design and coordination, drafted the manuscript, participated in the sequence alignment, and read and approved the final manuscript.

Funding

This work was supported by National Natural Science Foundation of China (No. 11761074), Project of Jilin Science and Technology Development for Leading Talent of Science and Technology Innovation in Middle and Young and Team Project(No.20200301053RQ), and the scientific research project of Guangzhou College of Technology and Business in 2020 (No. KA202032).

REFERENCES

- [1] C. Badea, *On the Hyers-Ulam stability of mappings: the direct method*. In: *Stability of Mappings of Hyers-Ulam Type* (edited by Th. M. Rassias and J. Tabor), Hadronic Press, Palm Harbor, Florida, 1994, pp. 7–13.
- [2] Z. Gajda, *On stability of the Cauchy equation on semigroups*, Aequationes Math. **36** (1988), 76–79.
- [3] D. Inoan and D. Popa, *On selections of generalized convex set-valued maps*, Aequationes Math. **88** (2014), 267–276.
- [4] S. Jung, *On modified Hyers-Ulam-Rassias stability of a generalized Cauchy functional equation*, Nonlinear Stud. **5** (1998), 59–67.
- [5] G. Lu and C. Park, *Hyers-Ulam stability of additive set-valued functional equations*, Appl. Math. Lett. **24** (2011), 1312–1316.
- [6] K. Nikodem, *On Jensen's functional equation for set-valued functions*, Radovi Mat. **3** (1987), 23–33.
- [7] K. Nikodem, *Set-valued solutions of the Pexider functional equation*, Funkcialaj Ekvacioj **31** (1988), 227–231.
- [8] K. Nikodem and D. Popa, *On single-valuedness of set-valued maps satisfying linear inclusions*, Banach J. Math. Anal. **3** (2009), 44–51.
- [9] K. Nikodem and D. Popa, *On selections of general linear inclusions*, Publ. Math. Debrecen **75** (2009), 239–249.
- [10] C. Park, D. O'Regan and R. Saadati, *Stability of some set-valued functional equations*, Appl. Math. Lett. **24** (2011), 1910–1914.
- [11] D. Popa, *A stability result for a general linear inclusion*, Nonlinear Funct. Anal. Appl. **3** (2004), 405–414.

- [12] D. Popa and N. Vornicescu, *Locally compact set-valued solutions for the general linear equation*, Aequationes Math. **67** (2004), 205–215.
- [13] H. Rådström, *An embedding theorem for space of convex sets*, Proc. Am. Math. Soc. **3** (1952), 165–169.
- [14] H. Rådström, *One-parameter semigroups of subsets of a real linear space*, Ark. Mat. **4** (1960), 87–97.
- [15] Th. M. Rassias and J. Tabor, *What is left of Hyers-Ulam stability?*, J. Nat. Geom. **1** (1992), 65–69.

GANG LYU

DIVISION OF FOUNDATIONAL TEACHING, GUANGZHOU COLLEGE OF TECHNOLOGY AND BUSINESS,
GUANGZHOU 510850, P.R. CHINA

Email address: lvgang@ybu.edu.cn

MINGZHU MU

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, SHENYANG UNIVERSITY OF TECHNOLOGY,
SHENYANG 110870, P.R. CHINA

Email address: 28138389@qq.com

YUANFENG JIN

DEPARTMENT OF MATHEMATICS, YANBIAN UNIVERSITY, YANJI 133001, P.R. CHINA

Email address: yfkim@ybu.edu.cn

CHOONKIL PARK

RESEARCH INSTITUTE FOR NATURAL SCIENCES, HANYANG UNIVERSITY, SEOUL 04763, KOREA

Email address: baak@hanyang.ac.kr