

THE CAUCHY PROBLEM FOR THE L^2 -CRITICAL GENERALIZED ZAKHAROV-KUZNETSOV EQUATION IN DIMENSION 3

FELIPE LINARES AND JOÃO P.G. RAMOS

ABSTRACT. We prove local well-posedness for the L^2 critical generalized Zakharov-Kuznetsov equation in H^s , $s \in (3/4, 1)$. We also prove that the equation is “almost well-posedness” for initial data $u_0 \in H^s$, $s \in [1, 2)$, in the sense that the solution belongs to a certain intersection $C([0, T]: H^s(\mathbb{R}^3)) \cap X_T^s$ and is unique within that class, where we can ensure continuity of the data-to-solution map in an only slightly larger space. We also prove that solutions satisfy the expected conservation of L^2 -mass for the whole $s \in (3/4, 2)$ range, and energy for $s \in (1, 2)$. By a limiting argument, this implies, in particular, global existence for small initial data in H^1 . Finally, we study the question of almost everywhere (a.e.) convergence of solutions of the initial value problem to initial data.

1. INTRODUCTION

In this paper we will study the initial value problem (IVP) associated to the L^2 critical generalized Zakharov-Kuznetsov equation in dimension 3, that is,

$$(1.1) \quad \begin{cases} \partial_t u + \partial_x \Delta u + \partial_x (u^{7/3}) = 0 & \text{on } \mathbb{R}^3 \times \mathbb{R}; \\ u(x, 0) = u_0(x) & \text{on } \mathbb{R}^3. \end{cases}$$

where u is a real function and Δ denotes the Laplace operator in space variables.

The equation above is related to the family of the well-known generalized Zakharov-Kuznetsov equation

$$(1.2) \quad \partial_t u + \partial_x \Delta u + u^k \partial_x u = 0, \quad \text{on } \mathbb{R}^d \times \mathbb{R}, \quad k \in \mathbb{Z}^+,$$

where $d = 2, 3$.

The equation above, for $k = 1$, arises in the context of plasma physics, where it was formally derived by Zakharov and Kuznetsov [39] as a long wave small-amplitude limit of the Euler-Poisson system in the “cold plasma” approximation. This formal long-wave limit was rigorously justified by Lannes, Linares and Saut in [28] (see also [23] for derivation in a different context).

We also notice that the generalized Zakharov-Kuznetsov equation (1.2) is not completely integrable, but it has a Hamiltonian structure and possesses three invariants, namely,

$$I(t) = \int_{\mathbb{R}^d} u(x, t) dx = I(0),$$

$$M(t) = \int_{\mathbb{R}^d} u^2(x, t) dx = M(0),$$

1991 *Mathematics Subject Classification.* Primary: 35Q53. Secondary: 35B05.

Key words and phrases. Zakharov-Kuznetsov equation, well-posedness, low regularity.

and

$$E(t) = \int_{\mathbb{R}^d} \left(|\nabla u|^2 - \frac{2u^{k+1}}{(k+1)(k+2)} \right) dx = E(0),$$

which are well known to be important in order to obtain a priori estimates that allows one to extend solutions of the IVP globally.

In addition, a scaling argument suggests that in order to obtain local well-posedness in $H^s(\mathbb{R}^d)$ for the IVP associated to equation (1.2) it is required to have $s \geq s_k = \frac{d}{2} - \frac{2}{k}$.

There is an extensive literature addressing the issue of local well-posedness of the IVP associated to the equation (1.2). For $k = 1$ and 2D, it starts with the work of Faminskii [14], where the local theory for initial data in the Sobolev spaces $H^s(\mathbb{R}^2)$, $s \geq 1$, is established. The method of proof used there combine smoothing estimates and a contraction mapping principle. We mention also some subsequent extensions obtained by Linares and Pastor in [29], refining this method with inspiration in the results for the Korteweg-de Vries equation

$$(1.3) \quad \partial_t u + \partial_x^3 u + u \partial_x u = 0, \quad \text{on } \mathbb{R} \times \mathbb{R}$$

by Kenig, Ponce and Vega [25]. A major improvement was obtained by Molinet and Pilod [34] and Grünrock and Herr [22], in both cases using the Fourier restriction method, where the authors prove local well-posedness in $H^s(\mathbb{R}^2)$, $s > 1/2$. We notice that the aforementioned results allow to extend the local solutions globally in $H^s(\mathbb{R}^2)$, $s \geq 1$. Recently, Kinoshita [26] obtained the best possible result attainable with the method of contraction principle. He proves a local theory in $H^s(\mathbb{R}^2)$, $s > -1/4$, by implementing sharp bilinear estimates and using a rather refined analysis. Kinoshita's result implies global well-posedness in $H^s(\mathbb{R}^2)$, $s \geq 0$. In the 2D case, the scale argument suggests local well-posedness results for (1.2) should hold for data in $H^s(\mathbb{R}^2)$, with $s > s_k = 1 - 2/k$. In that regard, sharp local results were obtained by Ribaud and Vento [35] for $k \geq 4$. In [21], Grünrock proved the local well-posedness for (1.2) in the case $k = 3$ in $H^s(\mathbb{R}^2)$, $s > 1/3$. For the particular nonlinearity $k = 2$, also called modified Zakharov-Kuznetsov equation (mZK), for which $L^2(\mathbb{R}^2)$ is the critical space suggested by the scale, Kinoshita [27] showed that the best possible result is $\dot{H}^s(\mathbb{R}^2)$, $s \geq 1/4$, complementing the Result by Ribaud and Vento [35] that proves local well-posedness in $H^s(\mathbb{R}^2)$, $s > 1/4$. In addition to that, recently Bhattacharya, Farah and Roudenko [1] proved global well-posedness for the same modified Zakharov-Kuznetsov equation in 2D if $s > 3/4$. For more related results see also [3], [30], and [15].

Regarding the 3D problem and $k = 1$, Ribaud and Vento [36] showed local well-posedness in $H^s(\mathbb{R}^3)$, $s > 1$ (see also [32]). Global well-posedness was proved by Molinet and Pilod [34] in the same spaces. Herr and Kinoshita in [24] establish an optimal local well-posedness result by using Picard iteration method, for data in $H^s(\mathbb{R}^3)$, $s > -1/2$ and so obtaining global well-posedness in $H^s(\mathbb{R}^3)$, $s \geq 0$. For $k \geq 3$, results by Grünrock [21] establish local well-posedness for data in $H^s(\mathbb{R}^3)$ and $s > 3/2 - 2/k$. The case $k = 2$ was particularly addressed by both Grünrock [20] and Kinoshita [27], where both authors show that the IVP (1.2) is locally well-posed in $H^s(\mathbb{R}^3)$ for $s > 1/2$. Kinoshita additionally

shows that we can push the former result up to the endpoint, and global existence holds in the whole $s \geq 1/2$ range.

We notice that in the 3D case the scale of the equation in (1.2) does not attain L^2 for any power $k \in \mathbb{Z}^+$, differently from the case in 2D. The natural question that arises is whether one can have a meaningful equation generalizing the ZK equation where we can expect local well-posedness in L^2 . Using the scale and letting k in (1.2) vary through the reals we find the equation (1.1), where we consider real initial data u_0 , for which we investigate the *real solutions* of (1.1).

The equation in (1.1) is called L^2 -critical not only because of L^2 being the largest resolution space for the IVP (1.1), but also due to the fact that the nonlinearity is the “borderline” case that imposes restrictions on the size of the data in L^2 in order to obtain global solutions in $H^1(\mathbb{R}^3)$. In this case, the energy and the mass are invariant (See (1.4), (1.5)). In addition, in the study of the existence and stability of ground state solutions to the equation in (1.2) (see (4.9) below) it is known that for $k < 4/3$ the ground states are stable and are unstable if $k > 4/3$ (see [9]). The case $k = 4/3$ is conjectured to be unstable.

Recently, Farah, Holmes, Roudenko and Yang in [17] showed that the solutions of the IVP associated to the 2D L^2 -critical equation in (1.2) blow-up in finite or infinite time. For ground states theory in the 2D case see [16] and [8].

The facts above motivate our interest in the study of the IVP (1.1). Our main goal here is to establish local and global well-posedness for the IVP (1.1). Nevertheless, the techniques above described do not apply directly in the analysis we will implement to prove the local theory.

Our main result is stated as follows.

Theorem 1.1. *Let $u_0 \in H^s$, $s \in (3/4, 1)$. Then there are function spaces X_T^s so that the IVP (1.1) has a unique solution*

$$u \in C([0, T]: H^s(\mathbb{R}^3)) \cap X_T^s,$$

where $T = T(\|u_0\|_s) > 0$. Moreover, the map $u_0 \mapsto u$ from $H^s(\mathbb{R}^d)$ to $C([0, T]: H^s(\mathbb{R}^3)) \cap X_T^s$ is locally Lipschitz continuous.

To obtain this result we first employ the methods introduced by Ribaud and Vento [35, 36], used to deal with the Zakharov-Kuznetsov equation in dimension 3 and the k -generalized Zakharov Kuznetsov equation, $k \in \mathbb{N}, k \geq 2$, in dimension 2.

In order to deal with the fact that our exponent lies strictly between 1 and 2, we need to come up with a method which takes into consideration both the approach for $k = 1$ [35] as well as that for $k = 2$ [20]. To handle the fact that the nonlinearity is not polynomial, we need to introduce an almost-paraproduct decomposition of $u^{7/3}$, together with a nonlinear lemma (see Lemma 3.1), in order to deal with that contribution. This nonlinear lemma can be interpreted as an anisotropic version of Proposition 2.3 in [10]. This allows us to employ the Picard iteration method in order to deal with $s \in (3/4, 1)$.

In the estimates defining the spaces X_T^s to solve the IVP (1.1) we include a maximal function estimate which is not enough to help us get further regularity but just $s = 1^-$

at most. To overcome that problem we prove an “almost well-posedness” result, in the following sense. For $s \in [1, 2)$, we modify the proof of the nonlinear Lemma 3.1 in order to obtain a priori estimates on the H^s norm of a solution $u(t)$ of (1.1). This is achieved by, instead of using the gradient in the analysis of low frequencies, applying the Laplacian. Unfortunately, this only gives us a priori bounds, and the Duhamel integral operator would only be $1/3$ -Hölder continuous with that method, which makes us use a different method than a direct Picard iteration approach. The way out is to consider solutions with smooth enough initial data and lower regularity level, and use the a priori estimate we have in our hands to upgrade this solution by proving it is actually smoother. In the end, we use approximations to the original initial data in H^s and functional analysis considerations to prove that the solution is, in fact, *strong*, and it belongs to the same kind of Banach spaces measuring regularity we used for the Picard iteration approach.

Theorem 1.2. *Let $u_0 \in H^s(\mathbb{R}^3)$, with $s \in [1, 2)$. Then there is $T = T(\|u_0\|_s)$ so that the solution u given by Lemma 4.3 is the unique that belongs to $L^\infty([0, T]: H^s(\mathbb{R}^3)) \cap \tilde{X}_T^s$, where we define $\tilde{X}_T^s = \bigcap_{3/4 < r < s} X_T^r$ and endow it with the norm $\|\cdot\|_{\tilde{X}_T^s}^2 = \int_{5/6}^s \|\cdot\|_{X_T^r}^2 dr$. Moreover, if $u_0, v_0 \in H^s$ satisfy that $\|u_0 - v_0\|_{H^s}$ is sufficiently small, then so is $\|u - v\|_{\tilde{X}_t^s}$.*

This result still does not give well-posedness in its total strength. Fortunately, as a consequence of Theorem 1.2, we are still able to obtain that the solution as stated belongs to the same kind of spaces as in the $s \in (3/4, 1)$ case.

Corollary 1.3. *Let $u_0 \in H^s(\mathbb{R}^3)$, $s \in [1, 2)$ be as before. Then the solution u to the IVP (1.1) belongs to X_T^s . In particular, $u \in C([0, T]: H^s(\mathbb{R}^3)) \cap X_T^s$.*

Remark 1.4. *In order to put this result in perspective, notice that Theorem 1.1 guarantees the data-to-solution map in the $s \in (3/4, 1)$ context is continuous, while in the remaining range with our current methods we can only guarantee that the solution does not lose regularity and forms a continuous curve in $H^s(\mathbb{R}^3)$.*

Remark 1.5. *Although the range $s \in (3/4, 2)$ looks reasonably far away from the scaling index $s_c = 0$, we mention that this endpoint cannot be included in the well-posedness theory, as least not if we demand that the data-to-solution $u_0 \mapsto u$ map is uniformly continuous. In Proposition 4.4 it is shown that the data-to-solution map is not uniformly continuous in $H^s(\mathbb{R}^3)$, $s \leq 0$.*

Next we would like to extend our solution globally in $H^1(\mathbb{R}^3)$. To do so we use the smooth solutions found in order to establish the previous theorem. We prove that those solutions allow us to justify the conservation of mass and energy satisfied by the flow of the equation. More precisely, our next main result reads as follows.

Theorem 1.6. *Let $u_0 \in H^s$, $s \in (3/4, 2)$, and denote by $u \in C([0, T]: H^s(\mathbb{R}^3)) \cap X_T^s$ the unique solution to the IVP (1.1) given by Theorem 1.2. The following assertions hold:*

- (i) *For each $t \in [0, T]$, where $T = T(\|u_0\|_s)$ is given by Theorem 1.2, the quantity*

$$(1.4) \quad M(u(t)) = \|u(t)\|_2^2$$

is preserved by the flow of the solution of (1.1).

(ii) If $s \in (1, 2)$, then, for each $t \in [0, T]$, the energy

$$(1.5) \quad E(u(t)) = \int_{\mathbb{R}^3} |\nabla u(t)|^2 dx dy - \frac{3}{5} \int_{\mathbb{R}^3} u(x, y, t)^{10/3} dx dy$$

is preserved by the flow of the solution of (1.1).

As a consequence of this theorem we obtain our second main result which is the global existence of solutions of the IVP (1.1) in H^1 :

Theorem 1.7. *Let $u_0 \in H^1(\mathbb{R}^3)$ and $u \in C([0, T]: H^1) \cap X_T^1$ the solution to (1.1), as obtained in Theorem 1.2 and Corollary 1.3. Then, if $\|u_0\|_2$ is sufficiently small, we have that the solution $u \in (L^\infty \cap C)(\mathbb{R}; H^1(\mathbb{R}^3))$.*

Finally, we are concerned with the question of almost everywhere (a.e.) convergence of solutions of the IVP (1.1) to initial data. We recall the problem proposed by Carleson in [5] in the context of Schrödinger operators: for which $s \in \mathbb{R}$ does it hold that if $u_0 \in H^s(\mathbb{R}^n)$, then

$$(1.6) \quad \lim_{t \rightarrow 0} e^{it\Delta} u_0(x) = u_0(x) \quad \text{for a.e. } x?$$

He showed that a.e. convergence holds for $u_0 \in H^{1/4}(\mathbb{R}^n)$, $n = 1$. Since then the study of this problem (linear and nonlinear) has gained a lot of attention, and inspired by the recent work of Compaan, Lucà and Staffilani [7], we prove that for each initial datum $u_0 \in H^s(\mathbb{R}^3)$, $s > \frac{3}{4}$, we have *pointwise convergence* of the flow to u_0 .

Theorem 1.8. *Let $s \in (3/4, 2)$. Then, for each initial datum $u_0 \in H^s(\mathbb{R}^d)$, the unique solution $u \in C([0, T]: H^s(\mathbb{R}^3))$ to the IVP (1.1) given by Theorems 1.1 and 1.2 converges pointwise to u_0 ; i.e.,*

$$\lim_{t \rightarrow 0} u(x, t) = u_0(x), \quad \text{for a.e. } x \in \mathbb{R}^3.$$

We use Grünrock's sharp $L_{x,y}^4$ maximal estimate together with an adaptation of our previous techniques to obtain such a result.

Organization. Before moving on to the proofs of our main results, we describe how the paper is organized.

In Section 2, we define the functional spaces we will work, and present linear estimates useful to define these spaces. In Section 3, we prove Theorem 1.1. In Section 4, we prove the “almost well-posedness result, Theorem 1.2. Finally, in Section 5, we prove the global result (Theorem 1.7), the conserved quantities (Theorem 1.6) and the Pointwise convergence (Theorem 1.8).

2. PRELIMINARIES

2.1. Notation and basic definitions. We use the modified Vinogradov equation $A \lesssim B$ several times to indicate that there is an absolute constant $C > 0$ so that $A \leq C \cdot B$. We also use the original Vinogradov equation $A \ll B$ to denote that there is a (relatively) large constant C with the property $A \leq C \cdot B$. We also use several times the notation $(-\Delta)^{s/2} f = \langle \nabla \rangle^s f = \mathcal{F}^{-1}((1 + |\xi|^2 + |\eta|^2)^{s/2} \hat{f})$.

Throughout this manuscript, we will work with the inhomogeneous Sobolev spaces H^s , which are the closure of Schwartz functions with respect to the norm $\|u\|_{H^s}^2 = \int_{\mathbb{R}^3} |\widehat{u}(\xi)|^2 (1 + |\xi|)^{2s} d\xi$. Of course, these function spaces are nested, as can be trivially seen from the definition.

We will also use several times the notation

$$\Delta_j(f) = \mathcal{F}^{-1}(\varphi_j \cdot \widehat{f}), \quad j > 0,$$

where φ is a smooth function supported in the annulus $\{z \in \mathbb{R}^3: |z| \in [1, 2]\}$, equal to one on $\{z \in \mathbb{R}^3: |z| \in [5/4, 7/4]\}$, $\varphi_j(z) = \varphi(z/2^j)$, and so that

$$\Delta_0(f) + \sum_{j>0} \Delta_j(f) = f,$$

where we define $\Delta_0(f) := \sum_{j \leq 0} \mathcal{F}^{-1}(\varphi_j \widehat{f})$. For any $j \gg 1$, the support properties of our φ imply that

$$\tilde{\Delta}_j := \Delta_{j-1} + \Delta_j + \Delta_{j+1}$$

satisfies that $\tilde{\Delta}_j \circ \Delta_j = \Delta_j$. We will use this identity implicitly throughout the text.

These operators are useful due to the following Littlewood–Paley characterization of Sobolev spaces (see, e.g., [19, Theorem 1.3.6]): $f \in H^s(\mathbb{R}^3)$ if and only if

$$|f|_s = \|\Delta_0(f)\|_2 + \left(\sum_{j>0} 2^{2sj} \|\Delta_j(f)\|_2^2 \right)^{1/2} < +\infty.$$

Moreover, the left-hand side above defines an equivalent norm to the usual $\|\cdot\|_{H^s}$ norm. With aid of the dyadic decomposition outlined above, we define the main spaces we will use to prove well-posedness. Define, for $u \in \mathcal{S}(\mathbb{R}^4)$, the following norms:

$$(2.1) \quad \|u\|_{X_T^s(\text{Sobolev})} = \|\Delta_0(u)\|_{L_T^\infty L_{x,y}^2} + \left(\sum_{j>0} 2^{2sj} \|\Delta_j(u)\|_{L_T^\infty L_{x,y}^2}^2 \right)^{1/2},$$

$$(2.2) \quad \|u\|_{X_T^s(\text{smooth})} = \|\Delta_0(u)\|_{L_x^\infty L_{y,T}^2} + \left(\sum_{j>0} 2^{2(s+1)j} \|\Delta_j(u)\|_{L_x^\infty L_{y,T}^2}^2 \right)^{1/2},$$

$$(2.3) \quad \|u\|_{X_T^s(\text{max})} = \|\Delta_0(u)\|_{L_x^2 L_{y,T}^\infty} + \left(\sum_{j>0} 2^{2(s-1-\varepsilon)j} \|\Delta_j(u)\|_{L_x^2 L_{y,T}^\infty}^2 \right)^{1/2},$$

$$(2.4) \quad \|u\|_{X_T^s(\text{Strichartz})} = \|\Delta_0(u)\|_{L_{x,y,T}^4} + \left(\sum_{j>0} 2^{2sj} \|\Delta_j(u)\|_{L_{x,y,T}^4}^2 \right)^{1/2}.$$

We therefore define the spaces X_T^s as the closure of $\mathcal{S}(\mathbb{R}^4)$ under the norm

$$(2.5) \quad \|u\|_{X_T^s} = \|u\|_{X_T^s(\text{Sobolev})} + \|u\|_{X_T^s(\text{smooth})} + \|u\|_{X_T^s(\text{max})} + \|u\|_{X_T^s(\text{Strichartz})}.$$

These are the main spaces which will help us prove well-posedness in our setting. We note that sometimes we will use the fact that, for $k > 0$, we may write

$$\|\Delta_k(u)\|_{X_T^s} \sim \|\Delta_k(u)\|_{Y_T^s},$$

where we let

$$\|u\|_{Y_T^s} = \|u\|_{L_T^\infty H_{x,y}^s} + \|\langle \nabla \rangle^{s+1} u\|_{L_x^\infty L_{y,T}^2} + \|\langle \nabla \rangle^{s-1+} u\|_{L_x^2 L_{y,T}^\infty} + \|\langle \nabla \rangle^s u\|_{L_{x,y,T}^4}.$$

2.2. Linear Estimates. Let

$$(2.6) \quad \begin{cases} \partial_t u + \partial_x \Delta u = 0 & \text{on } \mathbb{R}^3 \times \mathbb{R}; \\ u(x, 0) = u_0(x) & \text{on } \mathbb{R}^3, \end{cases}$$

be the linear part of (1.1). We denote the solution to this problem by $U(t)u_0$ defined by

$$\widehat{U(t)u_0}(\xi, \eta) = e^{it\xi(\xi^2 + |\eta|^2)} \widehat{u_0}(\xi, \eta), \quad \xi \in \mathbb{R}, \eta \in \mathbb{R}^2.$$

By Plancherel's theorem, this is an unitary group on L^2 . Besides that, the following estimates hold:

Proposition 2.1 (Kato Smoothing). *For any $u_0 \in L^2(\mathbb{R}^3)$, it holds that*

$$\|\nabla U(t)u_0\|_{L_x^\infty L_{y,T}^2} \lesssim \|u_0\|_{L^2}.$$

Proof. See, for instance, [36, Proposition 3.1]. □

Proposition 2.2 (Maximal Estimate). *For any $T \in (0, 1)$, it holds that*

$$\|U(t)u_0\|_{L_x^2 L_{y,T}^\infty} \lesssim \|u_0\|_{H^s},$$

for all $s > 1$ and all $u_0 \in \mathcal{S}(\mathbb{R}^3)$. In particular, for the endpoint $s = 1$, it holds that

$$\|U(t)\Delta_k u_0\|_{L_x^2 L_{y,T}^\infty} \lesssim k^2 \|\Delta_k u_0\|_{H^1},$$

for all $k \geq 1$.

Proof. See Propositions 3.2 and 3.3 in [36]. □

Proposition 2.3 ($L^2 - L^4$ Strichartz estimate). *For all $u_0 \in L^2(\mathbb{R}^3)$, it holds that*

$$\|U(t)u_0\|_{L_{x,y,t}^4} \lesssim \|u_0\|_{L^2}.$$

Proof. This is a direct consequence of the observations made in [20]. Alternatively, see also [21, Lemma 3]. □

As a consequence of those estimates, we have the following retarded estimates for the group $U(t)$:

Proposition 2.4 ([36], Propositions 3.5–3.7). *Let $f \in \mathcal{S}(\mathbb{R}^4)$. It holds that*

(i)

$$\left\| \nabla \int_0^t U(t-t')f(t') dt' \right\|_{L_T^\infty L_{x,y}^2} \lesssim \|f\|_{L_x^1 L_{y,T}^2};$$

(ii)

$$\left\| \nabla^2 \int_0^t U(t-t')f(t') dt' \right\|_{L_x^\infty L_{y,T}^2} \lesssim \|f\|_{L_x^1 L_{y,T}^2};$$

(iii)

$$\left\| \int_0^t U(t-t') \Delta_k f(t') dt' \right\|_{L_x^2 L_{y,T}^\infty} \lesssim_\varepsilon 2^{\varepsilon k} \|\Delta_k f\|_{L_x^1 L_{y,T}^2}.$$

Proposition 2.5 (Retarded $L^2 - L^4$ Strichartz). *Let $f \in \mathcal{S}(\mathbb{R}^4)$. It holds that*

$$\left\| \nabla \int_0^t U(t-t') f(t') dt' \right\|_{L_{x,y,t}^4} \lesssim \|f\|_{L_x^1 L_{y,t}^2}.$$

Proof. Proposition 2.3 together with the dual version of Proposition 2.1 implies that

$$\left\| \nabla \int_{-\infty}^\infty U(t-t') f(t') dt' \right\|_{L_{x,y,t}^4} \lesssim \|f\|_{L_x^1 L_{y,t}^2}.$$

Using Lemma B.3, part (i) in [2], which is an anisotropic version of the famous Christ–Kiselev lemma (see [6]), gives us the result. \square

3. THE CAUCHY PROBLEM FOR (1.1) : THE CASE $s \in (3/4, 1)$

As mentioned in the introduction, for the easier $s \in (3/4, 1)$ range we are able to employ the Picard iteration method. Indeed, we let

$$\Gamma(u) = U(t)u_0 + \int_0^t U(t-t') \partial_x(u^{7/3})(t') dt'.$$

We wish to prove that Γ preserves the metric space $E_a(T) = \{u \in X_T^s, \|u\|_{X_T^s} \leq a\}$ for some $a > 0$, and that, for such a , it also defines a contraction there. In order to prove the first fact, we need to prove that

$$(3.1) \quad \|U(t)u_0\|_{X_T^s} \lesssim C_s \|u_0\|_{H^s}.$$

This is a relatively simple computation using the fact that U is unitary and Propositions 2.1, 2.2 and 2.3, and thus we will omit it. Therefore, we are left with the integral term $\int_0^t U(t-t') \partial_x(u^{7/3})(t') dt'$ to handle. As bounding that term is essentially as difficult as bounding the difference

$$\int_0^t U(t-t') \partial_x(u^{7/3} - v^{7/3})(t') dt',$$

which naturally arises when trying to prove that Γ is a contraction, we prove only the latter.

On the other hand, we observe that the retarded Proposition 2.4(i), 2.4(ii), 2.4(iii) and 2.5 imply, by the same token as before, that

$$(3.2) \quad \left\| \int_0^t U(t-t') \partial_x(u^{7/3} - v^{7/3})(t') dt' \right\|_{X_T^s} \lesssim \left\| 2^{sk} \|\Delta_k(u^{7/3} - v^{7/3})\|_{L_x^1 L_{y,T}^2} \right\|_{\ell^2(\mathbb{N})}.$$

Until this point, we have followed essentially the same lines as in [35, 36]. The first point we have to change from the approach undertaken in those references is in the decomposition of the nonlinear part. In the case of the Zakharov–Kuznetsov equation, Ribaud and Vento’s approach involves writing

$$u^2 = \lim_{j \rightarrow \infty} (P_j u)^2 = (P_0 u)^2 + \sum_{j \geq 0} \Delta_{j+1} u (P_{j+1} u + P_j u).$$

Of course, such a clean formula is only available when the exponent is an integer. Our approach will be similar, yet being careful with the fact that we have a fractional nonlinearity.

Lemma 3.1. *Let $u, v \in \mathcal{S}(\mathbb{R}^4)$ and $k \geq 1$. It holds that $\|\Delta_k(u^{7/3} - v^{7/3})\|_{L_x^1 L_{y,T}^2}$ is bounded by an absolute constant times*

$$(3.3) \quad \begin{aligned} & \left(\|u\|_L^{4/3} + \|v\|_L^{4/3} \right) \sum_{m \geq k} \|\Delta_m(u - v)\|_{L_{x,y,T}^{p_1}} \\ & + \left(\|u\|_L^{1/3} + \|v\|_L^{1/3} \right) \|u - v\|_L \times \sum_{m \geq 0} \min(1, (k - m)_+ 2^{m-k}) \|\Delta_m u\|_{L_{x,y,T}^{p_1}}. \end{aligned}$$

Here, we abbreviate $L = L_x^{4p'_1/3} L_{y,T}^{4\tilde{p}_1/3}$, where q', \tilde{q} are defined for $q \geq 2$ so that $\frac{1}{q} + \frac{1}{q'} = 1$, $\frac{1}{q} + \frac{1}{\tilde{q}} = 1/2$.

Proof. We start by writing

$$\begin{aligned} \Delta_k(u^{7/3} - v^{7/3}) &= \sum_{m \geq 0} [\Delta_k((P_{m+1}u)^{7/3} - (P_{m+1}v)^{7/3}) - \Delta_k((P_m u)^{7/3} - (P_m v)^{7/3})] \\ &= \sum_{m \geq 0} \Delta_k(J_m(u, v)), \end{aligned}$$

where we define $J_m(u, v)$ to be

$$\begin{aligned} & P_{m+1}(u - v) \int_0^1 ((P_{m+1}u) + \theta(P_{m+1}(v - u)))^{4/3} \\ & - P_m(u - v) \int_0^1 ((P_m u) + \theta(P_m(v - u)))^{4/3} d\theta. \end{aligned}$$

We modify this expression once more: it can be rewritten as

$$\begin{aligned} & \Delta_{m+1}(u - v) \int_0^1 ((P_{m+1}u) + \theta(P_{m+1}(v - u)))^{4/3} d\theta \\ & - P_m(u - v) \int_0^1 [((P_m u) + \theta(P_m(v - u)))^{4/3} - ((P_{m+1}u) + \theta(P_{m+1}(v - u)))^{4/3}] d\theta. \end{aligned}$$

Finally, the last expression can be written in the form

$$\begin{aligned} & =: \Delta_{m+1}(u - v) \tilde{J}_m(u, v) + cP_m(u - v) \int_0^1 [\Delta_{m+1}u + \theta\Delta_{m+1}(u - v)] \\ & \left(\int_0^1 [((P_m u) + \theta(P_m(v - u))) + s(\Delta_{m+1}u + \theta\Delta_{m+1}(u - v))]^{1/3} ds \right) d\theta. \end{aligned}$$

By virtue of Minkowski's inequality, Young's convolution inequality and Hölder's inequality, we get that, for $m \geq k$,

$$\begin{aligned} & \|\Delta_k(J_m(u, v))\|_{L_x^1 L_{y,T}^2} \\ & \lesssim \|\Delta_{m+1}(u - v)\|_{L_{x,y,T}^{p_1}} \left(\|u\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}}^{4/3} + \|v\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}}^{4/3} \right) \\ & + \|\Delta_m u\|_{L_{x,y,T}^{p_1}} \left(\|u\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}}^{1/3} + \|v\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}}^{1/3} \right) \|u - v\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}}. \end{aligned}$$

Now, for $m < k$, we gain an exponential factor by introducing a gradient to take advantage of frequency localization around 2^k . Young's inequality several times plus Hölder's inequality then imply

$$\begin{aligned} & \|\Delta_k(J_m(u, v))\|_{L_x^1 L_{y,T}^2} \\ & \lesssim 2^{-k} \|\nabla[(P_{m+1}u)^{7/3} - (P_{m+1}v)^{7/3}] - \nabla[(P_m u)^{7/3} - (P_m v)^{7/3}]\|_{L_x^1 L_{y,T}^2} \\ & \lesssim 2^{-k} \sum_{i=0,1} \| |P_{m+i}(u - v)| (|\nabla P_{m+i}u| + |\nabla P_{m+i}v|) (|P_{m+i}u|^{1/3} + |P_{m+i}v|^{1/3}) \|_{L_x^1 L_{y,T}^2}. \end{aligned}$$

This is clearly bounded by an absolute constant times

$$2^{-k} \left(\sum_{m' \leq m+1} 2^{m'} \|\Delta_{m'} u\|_{L_{x,y,T}^{p_1}} \right) \|u - v\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}} \left(\|u\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}}^{1/3} + \|v\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}}^{1/3} \right).$$

Together with the bound for high frequencies this implies the lemma. \square

We note that this lemma is heavily inspired by Proposition 2.3 in [10], which is a clean version of such a result in the recent literature.

3.1. Analysis of high frequencies. We now look at the frequencies $\geq k$ in the bound given by Lemma 3.1. More precisely, we wish to estimate

$$(3.4) \quad \|w\|_{L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}} \quad \text{and} \quad \left\| 2^{sk} \sum_{m>k} \|\Delta_m w\|_{L_{T,x,y}^{p_1}} \right\|_{\ell^2(\mathbb{N})}$$

in terms of the norm $\|w\|_{X_T^s}$. By interpolating the first and fourth terms in the sum defining $\|\Delta_k w\|_{Y_T^s}$, it holds that

$$(3.5) \quad 2^{sk} \|\Delta_k w\|_{L_T^q L_{x,y}^p} \lesssim \|\Delta_k w\|_{Y_T^s},$$

whenever we have $\frac{1}{q} = \frac{\theta}{4}$, $\frac{1}{p} = \frac{1}{2} - \frac{\theta}{4}$. Therefore,

$$2^{sj} \|\Delta_j w\|_{L_{T,x,y}^{p_1}} \lesssim T^\delta 2^{sj} \|\Delta_j w\|_{L_T^{q_1} L_{x,y}^{p_1}} \lesssim T^\delta \|\Delta_j w\|_{Y_T^s},$$

for some $\delta > 0$. In order to take care of $\|w\|_{L_x^{4(p_1)'/3} L_{y,T}^{4(\tilde{p}_1)'/3}}$, we first bound it by

$$\sum_{j \geq 0} \|\Delta_j w\|_{L_x^{4(p_1)'/3} L_{y,T}^{4(\tilde{p}_1)'/3}}.$$

Now, interpolating the second and third terms defining Y_T^s , we get

$$(3.6) \quad 2^{(s+2\theta-1-\varepsilon)j} \|(\Delta_j w)\|_{L_x^{q_2} L_y^{4\tilde{p}_1/3}} \lesssim \|\Delta_j w\|_{Y_T^s},$$

where $\frac{3}{2\tilde{p}_1} = \theta$, and thus $q_2 = \frac{8p_1}{p_1+6}$. Bernstein's inequality then implies that

$$\|\Delta_j w\|_{L_x^{4(p_1)'/3} L_y^{4(\tilde{p}_1)'/3}} \lesssim 2^{\left(\frac{12-5p_1}{8p_1}\right)j} \|\Delta_j w\|_{L_x^{q_2} L_y^{4\tilde{p}_1/3}}.$$

This implies that

$$\|w\|_{L_x^{4(p_1)'/3} L_y^{4(\tilde{p}_1)'/3}} \lesssim \sum_{j \geq 0} 2^{\left[\left(\frac{12-5p_1}{8p_1}\right) - (s+2\theta-1-\varepsilon)\right]j} \|w\|_{Y_T^s}.$$

The sum above converges for $s > 1 - 2\theta - \frac{12-5p_1}{8p_1} = \frac{12+p_1}{8p_1}$. Notice that the argument above for using Bernstein's inequality implies, in particular, that the same strategy works as long as $p_1 < \frac{12}{5}$. Therefore, setting $p_1 = \frac{12}{5} - \gamma$ above with $\gamma > 0$ sufficiently small and running the argument, we will see that the second equation in (3.4) is bounded pointwise by

$$T^\delta \|(1_{j \leq 0} 2^{-sj}) * \|(\Delta_j w)\|_{Y_T^s}\|_{\ell^2(\mathbb{N})} \times \|w\|_{X_T^s}^{4/3} \lesssim T^\delta \|w\|_{X_T^s}^{7/3},$$

by the discrete version of Young's convolution inequality, as long as $s > \frac{3}{4}$.

3.2. Analysis of low frequencies. We need now to bound

$$(3.7) \quad \left\| 2^{sk} \sum_{k>m} (k-m)_+ 2^{-(k-m)} \|\Delta_m u\|_{L_{T,x,y}^{p_1}} \right\|_{\ell^2(\mathbb{N})}.$$

As we saw in the analysis for the high frequencies,

$$\|\Delta_m u\|_{L_{T,x,y}^{p_1}} \lesssim 2^{-sm} T^\delta \|\Delta_m u\|_{Y_T^s}.$$

Thus we get that

$$(3.8) \quad \begin{aligned} 2^{sk} \sum_{k>m} (k-m) 2^{-(k-m)} \|\Delta_m u\|_{L_{T,x,y}^{p_1}} &\lesssim T^\delta \sum_{k>m} (k-m) 2^{(s-1)(k-m)} \|\Delta_m u\|_{Y_T^s} \\ &= T^\delta \left[(1_{j \geq 0} j 2^{(s-1)j}) * \|\Delta_j u\|_{Y_T^s} \right] (k). \end{aligned}$$

Again by the discrete version of Young's inequality, we get that the $\ell^2(\mathbb{N})$ norm of the expression on the left hand side of (3.8) is bounded by

$$T^\delta \|u\|_{X_T^s} \|(1_{j \geq 0} j 2^{(s-1)j})\|_{\ell^1(\mathbb{N})} \lesssim T^\delta \|u\|_{X_T^s},$$

as long as $s < 1$. Therefore, we get from (3.3) that

$$\left\| \int_0^t U(t-t') \partial_x (u^{7/3})(t') dt' \right\|_{X_T^s} \lesssim T^\delta \|u\|_{X_T^s}^{7/3},$$

given that $s \in (3/4, 1)$.

Therefore, together with the estimate (3.3), we obtain

$$(3.9) \quad \left\| 2^{sk} \|\Delta_k (u^{7/3} - v^{7/3})\|_{L_x^1 L_y^2} \right\|_{\ell^2(\mathbb{N})} \lesssim T^\delta \|u - v\|_{X_T^s} \left(\|u\|_{X_T^s}^{4/3} + \|v\|_{X_T^s}^{4/3} \right).$$

3.3. Conclusion. The Duhamel formulation of (1.1) implies that the operator

$$\Gamma(u) = U(t)u_0 + \int_0^t U(t-t')\partial_x(u^{7/3})(t') dt'$$

is a contraction on $E_a(T)$, where we pick $a = 2C_s\|u_0\|_{H^s}$ the upper bound in (3.1) and $T \sim \|u_0\|_s^{-\beta}$, for some $\beta > 0$. Therefore, the fixed point theorem yields us that there is a solution to such a problem in $C([0, T]: H^s(\mathbb{R}^3)) \cap X_T^s$ with $s \in (3/4, 1)$, and it is unique. Moreover, standard considerations imply that the data-to-solution map is Lipschitz in that case, which concludes the proof of well-posedness.

4. THE CAUCHY PROBLEM FOR (1.1): THE CASE $s \in [1, 2)$.

4.1. An a priori bound. The gain of the form 2^{m-k} for low frequencies given by Lemma 3.1 is enough to allow us to prove local well-posedness for $s \in (3/4, 1)$, but it is clear from the analysis of low frequencies in §3.2 above that it breaks down for $s = 1$. In order to fix that, we will need the following version of Lemma 3.1.

Lemma 4.1. *Let $u \in \mathcal{S}(\mathbb{R}^4)$ and $k \geq 1$. It holds that $\|\Delta_k(u^{7/3})\|_{L_x^1 L_{y,T}^2}$ is bounded by an absolute constant times*

$$(4.1) \quad \|u\|_{X_T^s}^{4/3} \times \left(\sum_{m \geq k} \|\Delta_m(u)\|_{L_{x,y,T}^{p_1}} + \sum_{m \leq k} (k-m)_+ 2^{\min(2, \frac{1}{4} + s^-)(m-k)} \|\Delta_m u\|_{L_{x,y,T}^{p_1}} \right).$$

Here, we abbreviate $L = L_x^{4p_1'/3} L_{y,T}^{4\tilde{p}_1'/3}$, where q', \tilde{q}' are defined for $q \geq 2$ so that $\frac{1}{q} + \frac{1}{q'} = 1$, $\frac{1}{\tilde{q}} + \frac{1}{\tilde{q}'} = \frac{1}{2}$, and $s \in (1 - \gamma, +\infty)$ for some sufficiently small $\gamma > 0$.

Proof. The outline of the proof is exactly the same as in that of Lemma 3.1. If we write $\Delta_k(u^{7/3}) = \sum_{m \geq 0} \Delta_k(J_m u)$ as before, the same analysis as before yields that, for $m \geq k$,

$$\|\Delta_k(J_m u)\|_{L_x^1 L_{y,T}^2} \lesssim \|u\|_L^{4/3} \sum_{m \geq k} \|\Delta_m u\|_{L_{x,y,T}^{p_1}}.$$

On the other hand, for the frequencies $m < k$, we write

$$\begin{aligned} \|\Delta_k(J_m u)\|_{L_x^1 L_{y,T}^2} &\lesssim 2^{-2k} \sum_{i=0,1} \|\nabla^2(P_{m+i}u)^{7/3}\|_{L_x^1 L_{y,T}^2} \\ &\lesssim 2^{-2k} \sum_{i=0,1} \left(\|(\nabla^2 P_{m+i}u)(P_{m+i}u)^{4/3}\|_{L_x^1 L_{y,T}^2} + \| |\nabla P_{m+i}u|^2 (P_{m+i}u)^{1/3} \|_{L_x^1 L_{y,T}^2} \right). \\ &=: I_1(k, m) + I_2(k, m). \end{aligned}$$

In order to analyze the first term, we use Hölder's inequality in conjunction with Young's convolution and Bernstein's inequality again. This implies that

$$I_1(k, m) \lesssim \left(\sum_{m' \leq m} 2^{2(m'-k)} \|\Delta_{m'} u\|_{L_{x,y,T}^{p_1}} \right) \|u\|_L^{4/3}.$$

Using Fubini shows that this term can be absorbed into the right hand side of (4.1). Now, for the second term, we use Hölder's inequality for $L_x^{p_1'} L_{y,T}^{\tilde{p}_1'}$ with $(\nabla P_{m+i}u)(P_{m+i}u)^{1/3}$ and

$L_{x,y,T}^{p_1}$ with $\nabla P_{m+i}u$. Further uses of triangle inequality and Bernstein's inequality yield

$$(4.2) \quad I_2(k, m) \lesssim 2^{-2k} \left(\sum_{m' \leq m} 2^{m'} \|\Delta_{m'} u\|_{L_{x,y,T}^{p_1}} \right) \|(\nabla P_m u)(P_m u)^{1/3}\|_{L_{x^1}^{p'_1} L_{y,T}^{p_1}}.$$

Another use of Hölder's inequality shows that the right-hand side above is bounded by

$$C 2^{-2k} \left(\sum_{m' \leq m} 2^{m'} \|\Delta_{m'} u\|_{L_{x,y,T}^{p_1}} \right) \|u\|_L^{1/3} \|\nabla P_m u\|_{L_x^{4(p_1)'/3} L_{y,T}^{4\tilde{p}_1/3}}.$$

We focus on the last factor above. An analysis identical to the one employed in §3.1 implies that

$$\|\nabla P_m u\|_{L_x^{4(p_1)'/3} L_{y,T}^{4\tilde{p}_1/3}} \lesssim \sum_{m' \leq m} 2^{\left[\left(\frac{12-5p_1}{8p_1}\right) - (s+2\theta-1-\varepsilon)+1\right]m'} \|\Delta_{m'} u\|_{Y_T^s}.$$

with $\theta = \frac{3}{2p_1}$. Choosing p_1 sufficiently close to $12/5$, the right hand side above is bounded by

$$\sum_{m' \leq m} 2^{\left(\frac{3}{4}+(1-s)+\varepsilon\right)m'} \|\Delta_{m'} u\|_{Y_T^s} \lesssim 2^{\left(\frac{3}{4}+(1-s)+\varepsilon\right)k} \|u\|_{X_T^s}.$$

for any $s \in (1 - \gamma, +\infty)$, $\gamma > 0$ sufficiently small. Using (4.2), the upper bound above and Fubini,

$$\sum_{m \in \mathbb{N}, m \leq k} I_2(k, m) \lesssim \left(\sum_{m' \leq k} (k - m') 2^{\left(\frac{1}{4}+s-\varepsilon\right)(m'-k)} \|\Delta_{m'} u\|_{L_{x,y,T}^{p_1}} \right) \|u\|_{X_T^s}^{4/3}.$$

As this estimate can clearly be absorbed into the right-hand side of (4.1), the proof of the lemma is done. \square

We are now in a position to prove the following a priori bound:

Lemma 4.2. *Suppose $u \in X_T^s$ satisfies $\|u\|_{X_T^s} \leq 2C_s \|u_0\|_{H^s}$ and $s \in (3/4, 2)$. Then there is $T' = T'(\|u_0\|_s) < T$ so that*

$$\|\Gamma^j u\|_{X_{T''}^s} \leq 2C_s \|u_0\|_s,$$

for all $T'' \in [0, T']$, where $\{\Gamma^k u\}_{k \geq 0}$ denotes the orbit of the function u under the action of Γ .

Proof. We notice that the result holds trivially true for $s \in (3/4, 1)$ by the considerations in the previous section, and therefore we assume $s \in [1, 2)$.

We write the Duhamel formula defining Γ :

$$(4.3) \quad \Gamma u(t) = U(t)u_0 + \int_0^t U(t-t') \partial_x (u^{7/3})(t') dt'.$$

Applying the $X_{T''}^s$ norm on both sides, we see that

$$\|\Gamma u\|_{X_{T''}^s} \lesssim C_s \|u_0\|_s + \left\| 2^{sk} \|\Delta_k (u^{7/3})\|_{L_x^1 L_{y,T''}^2} \right\|_{\ell^2(\mathbb{N})}.$$

Due to Lemma 4.1 and the bounds from §3.2, we can bound this last expression by

$$C_s \|u_0\|_s + (T'')^\delta \|u\|_{X_{T''}^{s, 4/3}} \left\| 2^{sk} \left(\sum_{m \in \mathbb{N}} \min \left(1, (k-m)_+ 2^{\min(2, \frac{1}{4} + s^-)(m-k)} \right) \|\Delta_m u\|_{L_{T''}^q L_{x,y}^{p_1}} \right) \right\|_{\ell^2(\mathbb{N})}.$$

Again using that $\|\Delta_m u\|_{L_T^q L^{p_1} x, y} \lesssim 2^{-sm} \|\Delta_m u\|_{X_{T''}^s}$ we get that the $\ell^2(\mathbb{N})$ norm is, in turn, bounded by

$$(T'')^\delta \|u\|_{X_{T''}^{s, 4/3}} \left(\|\|\Delta_j u\|_{X_{T''}^s} * (1_{j \leq 0} 2^{sj})\|_{\ell^2(\mathbb{N})} + \|\|\Delta_j u\|_{X_{T''}^s} * (1_{j \geq 0} j 2^{(s - \min(2, 1/4 + s^-)j)}\|_{\ell^2(\mathbb{N})} \right).$$

Now, Young's convolution inequality allows us to bound the expression above by

$$(T'')^\delta \|u\|_{X_{T''}^s}^{7/3}$$

if $s < 2$. Indeed, if $s < 7/4$, then $\min(2, 1/4 + s^-) = 1 + s^-$, and thus $s - 1/4 - s^- = -1/4^+$. If $2 > s \geq 7/4$, we will just have $s - 2 < 0$ in the exponent of the term being convolved.

Therefore, if $T' = T'(\|u_0\|_s)$ is sufficiently small, as $\|u\|_{X_T^s} \leq 2C_s \|u_0\|_{H^s}$, it holds that $\|\Gamma u\|_{X_T^s} \leq 2C_s \|u_0\|_{H^s}$ as well. The assertion of the lemma follows then by iterating this result. \square

4.2. Weak well-posedness for $s \in [1, 2)$. The following lemma states an ‘almost well-posedness’ for the remaining range. It will be useful in order to prove the persistence of regularity and uniqueness for initial data in H^s , $s \in [1, 2)$.

Lemma 4.3. *Let $u_0 \in H^s(\mathbb{R}^3)$, with $s \in [1, 2)$. Then there is $T = T(\|u_0\|_s)$ and a unique solution u to (1.1) so that $u \in C([0, T]: H^r(\mathbb{R}^3)) \cap X_T^r$, for each $r \in (3/4, s)$.*

Proof. For $s \in [1, 2)$, let $\tilde{s} \in (3/4, 1)$ be a fixed parameter. Let $T = T(\|u_0\|_{\tilde{s}}, \|u_0\|_s)$ be sufficiently small, and $w \in X_T^{\tilde{s}}$ satisfy that $\|w\|_{X_T^{\tilde{s}}} \leq 2 \min(C_{\tilde{s}}, C_s) \|u_0\|_{\tilde{s}}$. The fact that the norms $\|\cdot\|_{X_T^s}$ are increasing with s and that we have proved Lipschitz well-posedness for $\tilde{s} \in (3/4, 1)$ implies that

$$\|\Gamma^j w - \Gamma^k w\|_{X_T^{\tilde{s}}} \rightarrow 0 \text{ as } \min(k, j) \rightarrow \infty.$$

By Lemma 4.2, we know that

$$\|\Gamma^j w - \Gamma^k w\|_{X_T^s} \leq 4C_s \|u_0\|_s.$$

By interpolating between the norms of $X_T^{\tilde{s}}$ and X_T^s , we obtain that

$$\|\Gamma^j w - \Gamma^k w\|_{X_T^r} \rightarrow 0 \text{ as } \min(j, k) \rightarrow \infty$$

for all $r \in [\tilde{s}, s)$. This implies that the sequence $\{\Gamma^j w\}_{j \geq 0}$ converges in X_T^r , and thus it converges to a fixed point of Γ , as the proof of Lemma 4.2 also shows that Γ is $1/3$ -Hölder continuous on X_T^r , $r \in [1, 2)$.

Let this fixed point be $w \in X_T^s$. The properties of Γ imply that $w \in C([0, T]: H^r) \cap X_T^r$. By the definition of Γ , this w satisfies the Duhamel formulation of (1.1), and thus it is a solution to (1.1) with initial data in $H^r(\mathbb{R}^3)$.

If $w_1, w_2 \in C([0, T]: H^r) \cap X_T^r$ are solutions to (1.1) for some $T > 0$, then, by the fact that the spaces H^r and X_T^r are nested in $r \in \mathbb{R}$, they are also strong solutions to (1.1)

in $C([0, T]: H^{\tilde{s}}) \cap X_{T'}^{\tilde{s}}$, for some $\tilde{s} \in (3/4, 1)$. As we know that uniqueness holds for that case, it also holds for $r \in [1, 2)$. This finishes the proof. \square

As a direct consequence, we are in position to prove the almost well-posedness result for $s \in [1, 2)$ stated in Theorem 1.2.

Proof of Theorem 1.2. Start by considering the set of mollified initial data $u_{0,\varepsilon} = \varphi_\varepsilon * u_0$ and u_ε the solution given by Lemma 4.3. By that result, it holds that u_ε belongs to $C([0, T_{\varepsilon,\delta}]: H^{s+\delta}) \cap X_{T_{\varepsilon,\delta}}^{s+\delta}$, where $T_{\varepsilon,\delta} = T(\varepsilon^{-\delta} \|u_0\|_s)$. Indeed, it holds that

$$(4.4) \quad \|u_\varepsilon\|_{X_{T'}^{s+\delta}} \leq 2C_s \|u_{0,\varepsilon}\|_{s+\delta}$$

whenever $T' < T(\|u_{0,\varepsilon}\|_{s+\delta})$. On the other hand, as $\|u_{0,\varepsilon}\|_{s+\delta} \lesssim \varepsilon^{-\delta} \|u_0\|_s$ and the T from Lemma 4.3 is nonincreasing on the norm of $\|u_0\|_s$, the conclusion follows. Furthermore, notice that for $\delta \sim \varepsilon$, we get that $T_{\varepsilon,\delta} \sim T(\|u_0\|_s)$ is *independent* of ε , like the upper bound in (4.4).

On the other hand, for $\tilde{s} \in (3/4, 1)$, the considerations in §3 imply that

$$(4.5) \quad \|u_{\varepsilon_1} - u_{\varepsilon_2}\|_{X_{T''}^{\tilde{s}}} \leq 2C_{\tilde{s}} \|u_{0,\varepsilon_1} - u_{0,\varepsilon_2}\|_{\tilde{s}},$$

whenever $T'' < \tilde{T}(\|u_0\|_{\tilde{s}})$. By (4.4) and the fact that the X_T^s -norms are increasing in $s \in \mathbb{R}$, it holds that $\{u_\varepsilon\}_{\varepsilon>0}$ is a bounded sequence in $X_{T'}^s, T' = T'(\|u_0\|_s)$. Interpolating (4.4) together with (4.5) then implies that $\{u_\varepsilon\}_{\varepsilon>0}$ is a Cauchy sequence in $X_{T''}^r$, for all $r \in (3/4, s)$ and all $T'' < \min(T_0(\|u_0\|_s), \tilde{T}(\|u_0\|_{\tilde{s}}))$.

By uniqueness of the limit, the solution u to the IVP (1.1) belongs to all $X_{T''}^r$ for such r and T'' . Notice, moreover, that the upper bound on the $X_{T''}^r$ -norm of the solution is uniformly bounded for such r and T'' . This readily implies that $u \in \tilde{X}_T^s$.

Also, notice that $\{u_\varepsilon\}_{\varepsilon>0}$ is a bounded sequence in $C([0, T_0]: H^s)$, and by the Banach-Alaoglu theorem, we can suppose (passing to a subsequence if necessary) that $u_\varepsilon \rightharpoonup u$ in $L^2([0, T]: H^s(\mathbb{R}^3))$. By properties of weakly convergent sequences and the uniform boundedness of u_ε in $C([0, T]: H^s)$, we can easily conclude that $u \in L^\infty([0, T_0]: H^s(\mathbb{R}^3))$.

Finally, the continuity of the solution with respect to initial data in the \tilde{X}_T^s norm can be proved as follows: let $u_0, v_0 \in H^s(\mathbb{R}^3)$, $s \in [1, 2)$, and let $u, v \in L^\infty([0, \delta]: H^s) \cap \tilde{X}_\delta^s$, for some common time $\delta > 0$. Then

$$(4.6) \quad \|u - v\|_{X_\delta^r} \leq \|u - u_\varepsilon\|_{X_\delta^r} + \|u_\varepsilon - v_\varepsilon\|_{X_\delta^r} + \|v - v_\varepsilon\|_{X_\delta^r}$$

holds trivially by triangle's inequality. Fix then $\eta > 0$ and let $r \in [3/4 + \eta, s - \eta]$. By the fact that $(u_\varepsilon, v_\varepsilon) \rightarrow (u, v)$ in $X_\delta^r \times X_\delta^r$, and that we can quantify the rate of such a convergence by the interpolation of (4.4) and (4.5), it holds that we can make the first and third terms in (4.6) uniformly small (with respect to $r \in [3/4 + \eta, s - \eta]$) by choosing $\varepsilon > 0$ sufficiently small.

On the other hand, for fixed $\varepsilon > 0$, the same interpolation argument shows that the middle term $\|u_\varepsilon - v_\varepsilon\|_{X_\delta^r}$ can be bounded in such an interval of values of r by some positive power of $\|u_{0,\varepsilon} - v_{0,\varepsilon}\|_{H^r} \lesssim C \|u_0 - v_0\|_{H^r}$. If v_0 and u_0 are close, then so is the middle term

of (4.6). Therefore,

$$\int_{5/6}^s \|u - v\|_{X_\delta^r}^2 dr \lesssim (s - \eta - 5/6) \times \delta + \eta \times \sup_{r \in [s-\eta, s]} \left(\|u\|_{X_\delta^r}^2 + \|v\|_{X_\delta^r}^2 \right).$$

By the uniform boundedness of the X_δ^r norms asserted above, the right-hand side in this last equation can be made arbitrarily small. \square

Proof of Corollary 1.3. We first prove a useful property of weak convergence:

Claim: If $u_\varepsilon \rightharpoonup u$ in $L^2([0, T]: H^s(\mathbb{R}^3))$, then $\Delta_k(u_\varepsilon) \rightharpoonup \Delta_k(u)$ in the same space. Indeed, let $v \in L^2([0, T]: H^{-s}(\mathbb{R}^3))$ be fixed. Then

$$\int_0^T \langle v(t), \Delta_k u_\varepsilon(t) \rangle_{H^{-s}, H^s} dt = \int_0^T \langle \Delta_k v(t), u_\varepsilon(t) \rangle_{H^{-s}, H^s} dt.$$

As $\Delta_k v \in L^2([0, T]: H^{-s}(\mathbb{R}^3))$, the right-hand side of the last equation still converges to

$$\int_0^T \langle \Delta_k v(t), u(t) \rangle_{H^{-s}, H^s} dt = \int_0^T \langle v(t), \Delta_k u(t) \rangle_{H^{-s}, H^s} dt$$

as $\varepsilon \rightarrow 0$. This concludes the proof of this claim.

Therefore, fix $K \in \mathbb{N}$. By passing to subsequences a finite number of times, we can suppose that, besides the weak convergence property given by the claim above, we have that $\Delta_k(u_\varepsilon) \rightarrow \Delta_k(u)$ for almost every $(x, y, t) \in \mathbb{R}^3 \times [0, T]$, $k = 1, 2, \dots, K$. In order to bound $\|u\|_{X_T^s}$, we need to bound terms of the form

$$(4.7) \quad \|\Delta_k(u)\|_{L_x^\infty L_{y,T}^2}, \|\Delta_k(u)\|_{L_x^2 L_{y,T}^\infty}, \|\Delta_k(u)\|_{L_T^\infty L_{x,y}^2}, \text{ and } \|\Delta_k(u)\|_{L_{x,y,T}^4}.$$

The easiest to handle is the last term in (4.7). Indeed,

$$\int_{\mathbb{R}^3 \times [0, T]} |\Delta_k(u)(x, y, t)|^4 dx dy dt \leq \liminf_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^3 \times [0, T]} |\Delta_k(u_\varepsilon)(x, y, t)|^4 dx dy dt$$

follows directly from Fatou's lemma and the asserted pointwise convergence from before. The strategy to bound the other terms is similar, but this time we need a substitute for the L^∞ norms appearing. Recall, for that purpose, that the normalized norms $\|\cdot\|_{\mathbb{L}^N(I, dm)} = \frac{1}{m(I)^{1/N}} \|\cdot\|_{L^N(I, dm)}$ are bounded by the L^∞ -norm on I . Moreover, if $f \in L^\infty$, then $\|f\|_{\mathbb{L}^N} \nearrow \|f\|_\infty$ as $N \rightarrow \infty$. Inspired by that, we can prove, for instance, that for fixed $R > 0, N \gg 1$,

$$(4.8) \quad \left(\int_{\mathbb{R}} \left(\int_{B_R^2 \times [0, T]} |\Delta_k(u)(x, y, t)|^N dy dt \right)^{2/N} dx \right)^{1/2}$$

is bounded – by the same Fatou argument as before – by

$$\liminf_{\varepsilon \rightarrow 0} \left(\int_{\mathbb{R}} \left(\int_{B_R^2 \times [0, T]} |\Delta_k(u_\varepsilon)(x, y, t)|^N dy dt \right)^{2/N} dx \right)^{1/2} \leq \liminf_{\varepsilon \rightarrow 0} \|\Delta_k(u_\varepsilon)\|_{L_x^2 L_{y,T}^\infty}.$$

On the other hand, taking first $N \rightarrow \infty$ in (4.8) and using the fact that L^N -averages converge monotonically to the supremum and monotone convergence, then taking $R \rightarrow \infty$

and using monotone convergence again, we see that (4.8) converges to $\|\Delta_k(u)\|_{L_x^2 L_{y,T}^\infty}$. As the other terms in (4.7) can be bounded in similar (in fact, simpler) ways, we have that, for each fixed $K \in \mathbb{N}$, the bound

$$\|P_K u\|_{X_T^s} \leq \limsup_{\varepsilon \rightarrow 0} \|u_\varepsilon\|_{X_T^s} \leq 2C_s \|u_0\|_s$$

holds, for $T = T(\|u_0\|_s)$ sufficiently small, given in Theorem 1.2. Taking $K \rightarrow \infty$ above then implies that $u \in X_T^s$, $\|u\|_{X_T^s} \leq 2C_s \|u_0\|_s$. Writing the Duhamel formulation of (1.1) that u satisfies as

$$u(t) - U(t)u_0 = \int_0^t U(t-t') \partial_x(u^{7/3})(t') dt'$$

and applying the X_T^s -norm on both sides and the estimates from Lemma 4.2, we get continuity of $u(t)$ at $t = 0$. By the group property, continuity at all other values of $t \in [0, T]$ follow in the same way. \square

Although the range $s \in (3/4, 2)$ looks reasonably far away from the scaling index $s_c = 0$, we mention that this endpoint cannot be included in the well-posedness theory, as least not if we demand that the data-to-solution $u_0 \mapsto u$ map is uniformly continuous. This is the content of the following proposition.

Proposition 4.4. *Let $s \leq 0$. Then the IVP (1.1) is ill-posed for initial data in $H^s(\mathbb{R}^3)$, in the sense that the data-to-solution map is not uniformly continuous.*

Proof. We only sketch the proof, as the main outline is the same as in [29, Theorem 1.2].

Indeed, fix $\phi \in H^1(\mathbb{R}^3)$ a positive, radial solution to

$$(4.9) \quad \Delta \phi - \phi + \phi^{7/3} = 0.$$

The existence of such a function is a by-product of the theory developed by Weinstein [38]. Now, we let $\psi_c(r) = c^{3/4} \phi(c^{1/2}r)$, where we abuse notation and denote a radial function by its value at radius $r \geq 0$. The function

$$u_c(z, t) = \psi_c((x - ct)^2 + |y|^2), \quad z = (x, y) \in \mathbb{R}^3,$$

can be seen through an easy computation to satisfy the IVP (1.1) with initial datum $u_{0,c}(z) = c^{3/4} \phi(\sqrt{c}z)$. Moreover, it is easy to verify that the Fourier transform of the initial data $\widehat{u_{0,c}}(\xi, \eta) = c^{-3/4} \widehat{\phi}\left(\frac{\xi}{\sqrt{c}}, \frac{\eta}{\sqrt{c}}\right)$. A simple calculation then shows that

$$\lim_{n \rightarrow \infty} \langle u_{0,c_n}, u_{0,c_{n+1}} \rangle_{L^2} = \|\phi\|_{L^2}^2,$$

where $c_n = n$, $\forall n \geq 1$. As we also have that $\|u_{0,c_n}\|_{L^2} = \|\phi\|_{L^2}$, we conclude that

$$\|u_{0,c_n} - u_{0,c_{n+1}}\|_{L^2} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

On the other hand, a similar computation with the Fourier transform for fixed time $t > 0$ yields that

$$\langle u_{c_n}(t), u_{c_{n+1}}(t) \rangle_{L^2} = \left(\frac{n}{n+1}\right)^{3/4} \int_{\mathbb{R}^3} e^{2\pi i \xi \sqrt{n}} \widehat{\phi}(\xi, \eta) \overline{\widehat{\phi}}\left(\frac{\sqrt{n}\xi}{\sqrt{n+1}}, \frac{\sqrt{n}\eta}{\sqrt{n+1}}\right) d\xi d\eta.$$

An application of the Riemann–Lebesgue lemma implies that the expression above goes to 0 as $n \rightarrow \infty$. By the way we defined our solutions, we have conservation of L^2 norm,

and thus $\|u_{c_n}(t)\|_2 = \|\phi\|_2$. This implies that

$$\lim_{n \rightarrow \infty} \|u_{c_n}(t) - u_{c_{n+1}}(t)\|_{L^2} = \sqrt{2}\|\phi\|_2.$$

Thus the solution map is not uniformly continuous. \square

5. CONSERVATION LAWS AND GLOBAL EXISTENCE IN H^s

5.1. The auxiliary problem. In order to prove that the conservation laws (1.4) and (1.5), we are going to be interested in the following smoother version of (1.1):

$$(5.1) \quad \begin{cases} \partial_t u^\varepsilon + \partial_x \Delta u^\varepsilon + \partial_x (F_\varepsilon^{7/3}(u^\varepsilon)) = 0 & \text{on } \mathbb{R}^3 \times \mathbb{R}; \\ u^\varepsilon(x, 0) = u_0(x) & \text{on } \mathbb{R}^3, \end{cases}$$

where we define $F_\varepsilon^\alpha(u) = \eta_\varepsilon * (u * \eta_\varepsilon)^\alpha$, for some smooth, even, positive function $\eta : \mathbb{R}^d \rightarrow \mathbb{R}$. We notice that this approach towards verifying the validity of the conservation laws is far from new; see, for instance, the pioneering work of Ginibre and Velo [18], the quantification due to Weinstein [38] and the book by Tao [37] for further details on this method.

The first result about this equation is that solutions to (5.1) are smooth for smooth initial data, as the following proposition states:

Proposition 5.1. *For each $u_0 \in H^N(\mathbb{R}^3)$, $N \gg 1$, there exists $T = T(\varepsilon, N, \|u\|_{H^N})$ such that the IVP (5.1) is locally well-posed in*

$$C([0, T]: H^N(\mathbb{R}^3)) \cap L^\infty([0, T]: W^{1, \infty}(\mathbb{R}^3)).$$

Proof. The proof involves a standard parabolic regularization method. Let

$$(5.2) \quad \begin{cases} \partial_t w + \eta(\Delta^2 w) + \partial_x \Delta w + \partial_x (F_\varepsilon^{7/3}(w)) = 0 & \text{on } \mathbb{R}^3 \times \mathbb{R}; \\ w(x, 0) = u_0(x) & \text{on } \mathbb{R}^3, \end{cases}$$

where we take $\eta > 0$. The fact that solutions of (5.2) are smooth can be proved easily with the contraction principle, using the fact that we may write the Duhamel formulation of the equation as

$$w(t) = e^{-\eta t \Delta^2} u_0 + \int_0^t e^{-\eta(t-t') \Delta^2} (\partial_x \Delta w + \partial_x (F_\varepsilon^{7/3} w))(t') dt'$$

and simple H^s estimates for the group $\{e^{-t\eta \Delta^2}\}_{t \in \mathbb{R}}$ given on the Fourier side. For more details see, for instance, [11, Section 3.1] or even [14]. Of course, this method gives us a maximal existence time depending on $\eta > 0$. In order to remove this restriction, we apply Δ^k to (5.2) and integrate the resulting equation against $\Delta^k w$. We obtain that

$$\partial_t \|\Delta^k w(t)\|_2^2 = -2 \int (\Delta^k w) \partial_x (\Delta^{k+1} w) - 2\eta \int (\Delta^k w) (\Delta^2 w) - 2 \int \partial_x \Delta^k (F_\varepsilon^{7/3} w) (\Delta^k w).$$

The first term on the right hand side above is zero, and the second is non-positive. Therefore, we are left with bounding

$$(5.3) \quad \left| \int \partial_x \Delta^k (F_\varepsilon^{7/3} w) (\Delta^k w) \right|.$$

By the way we defined the smooth nonlinearity, we obtain that $\partial_x \Delta^k (F_\varepsilon^{7/3} w) = (\Delta^k(\eta_\varepsilon)) * \partial_x (w * \eta_\varepsilon)^{7/3}$. This implies that (5.3) is bounded by

$$C_k \varepsilon^{-2k-1} \|\Delta^k w\|_2 \|(w * \eta_\varepsilon)^{4/3} (\partial_x w) * \eta_\varepsilon\|_2 \lesssim C_k \varepsilon^{-2k-1} \|\Delta^k w\|_2 \|w\|_{H^1}^{7/3}.$$

Putting this together with what we had before, summing in $2k \leq N$ and using first the Gagliardo–Nirenberg interpolation inequality, and then the Cauchy–Schwarz inequality plus properties of the convolution, we obtain

$$\partial_t \|w(t)\|_{H^N}^2 \leq C_N(\varepsilon) \|w(t)\|_{H^N} \|w\|_{H^1}^{7/3} \leq C_N(\varepsilon) \|w(t)\|_{H^N}^2 \|w\|_{H^1}^{4/3}.$$

The standard Gronwall trick implies that

$$(5.4) \quad \|w(t)\|_{H^N} \leq \|u_0\|_{H^N} \cdot \exp \left(C_N(\varepsilon) \int_0^t \|w(t')\|_{H^1}^{4/3} dt' \right).$$

Notice that this last equation does not depend on η , and thus it implies that there is $T_0 = T_0(\|u_0\|_{H^N}, N, \varepsilon)$ but not depending on η so that $\|w(t)\|_{H^N} \leq 2\|u_0\|_{H^N}$ for $t \in [0, T_0]$. This observation implies that the maximum time T_N of solution of (5.2) so that $u \in C^1([0, T_N]: H^N)$ for $u_0 \in H^N$ does not depend on $\eta > 0$.

These estimates, together with the very structure of the equation (5.2), enables us to use the Bona-Smith argument [4]. This implies that, if $w^{\eta, \varepsilon}$ denotes the solution to (5.2), then taking $\eta \rightarrow 0$, $w^{\eta, \varepsilon} \rightarrow u^\varepsilon$ in H^N . Moreover, because of (5.4), the fact that solutions to (5.2) are smooth and the fact the maximal time does not depend on η , it holds that solutions to (5.1) are also smooth for smooth enough initial data, and the problem is also well-posed on high order Sobolev spaces. We omit the details, as they are essentially completely contained in the references previously mentioned. \square

We have proved that, for each fixed $\varepsilon > 0$, the solution of (5.1) preserves regularity. This will be useful for proving that the solutions to (1.1) satisfy the conserved quantities.

On the other hand, in order to do so, we need to prove that solutions of (5.1) are well-defined for the range of Sobolev spaces in which we are working. This is the content of the next result.

Theorem 5.2. *Let $u_0 \in H^s(\mathbb{R}^3)$. The following assertions hold:*

- (i) *If $s \in (3/4, 1)$, then there is $T = T(\|u_0\|_s)$ independent of ε so that (5.1) is well-posed in $C([0, T]: H^s) \cap X_T^s$ with initial data u_0 .*
- (ii) *If $s \in [1, 2)$, then there is $T = T(\|u_0\|_s, \varepsilon)$ so that (5.1) is well-posed in $C([0, T]: H^s) \cap X_T^s$ with initial data u_0 .*

In both cases, the data-to-solution map is Lipschitz continuous.

Proof of part (i). Define the map

$$\Gamma_\varepsilon(w) = U(t)u_0 + \int_0^t U(t-t') \partial_x (F_\varepsilon^{7/3} w)(t') dt'.$$

Exactly in the same way as we did for the IVP (1.1), it suffices to control

$$\left\| \int_0^t U(t-t') \partial_x (F_\varepsilon^{7/3} w_1 - F_\varepsilon^{7/3} w_2)(t') dt' \right\|_{X_T^s}$$

$$\lesssim \left\| 2^{sk} \|\Delta_k(\eta_\varepsilon * ((w_1 * \eta_\varepsilon)^{7/3} - (w_2 * \eta_\varepsilon)^{7/3}))\|_{L_x^1 L_{y,T}^2} \right\|_{\ell^2(\mathbb{N})}.$$

As convolution with η_ε commutes with Δ_k , an application of Young's convolution inequality shows that the right hand side above is bounded by

$$\left\| 2^{sk} \|\Delta_k((w_1 * \eta_\varepsilon)^{7/3} - (w_2 * \eta_\varepsilon)^{7/3})\|_{L_x^1 L_{y,T}^2} \right\|_{\ell^2(\mathbb{N})}.$$

By the analysis undertaken in §3, the quantity above is controlled by $CT^\delta \|w_1 - w_2\|_{X_T^s}$. Therefore, the same argument works for proving well-posedness in $C([0, T]: H^s) \cap X_T^s$, $s \in (3/4, 1)$, of the IVP (5.1). Notice that, as none of the constants involved depended on $\varepsilon > 0$, the existence time does *not* depend on $\varepsilon > 0$ for such range of s .

Proof of part (ii). We follow the same path to existence as before, but now we need an analogue of Lemma 4.1 for differences. We follow thus the outline of proof of Lemma 3.1. The $m \geq k$ part can be kept without any modification. On the other hand, in order to estimate the contributions of terms $\|\Delta_k(J_m^\varepsilon(w_1, w_2))\|_{L_x^1 L_{y,T}^2}$ with $m < k$, we notice that

$$\begin{aligned} & \nabla^2(\eta_\varepsilon * [(P_m(w_1 * \eta_\varepsilon))^{7/3} - (P_m(w_2 * \eta_\varepsilon))^{7/3}]) \\ &= (\nabla \eta_\varepsilon) * \left[\nabla[(P_m(w_1 * \eta_\varepsilon))^{7/3} - (P_m(w_2 * \eta_\varepsilon))^{7/3}] \right]. \end{aligned}$$

Therefore, employing the same techniques as in both of those cases, the low-frequency contribution $\Delta_k(J_m^\varepsilon(w_1, w_2))$ can be bounded by

$$\frac{C}{\varepsilon} (\|w_1\|_L^{1/3} + \|w_2\|_L^{1/3}) \cdot \|w_1 - w_2\|_L \times \left(\sum_{m < k} (k - m)_+ 2^{2(m-k)} \|\Delta_m u\|_{L_{x,y,T}^{p_1}} \right).$$

The rest of the proof can be carried out in the exact same way as before. \square

Notice that redoing the proof of Lemmata 3.1 in 4.1 yields only a $\|u - v\|_{X_T^s}^{1/3}$ factor, and thus the operator Γ from §3 is only 1/3-Hölder continuous. For that reason we need to work with the smooth versions (5.1) and then hope to be able to pass to the limit. The next corollary is a major step in that direction.

Corollary 5.3. *Let $u_0 \in H^s(\mathbb{R}^3)$ for $s \in [1, 2)$. Denote the solution to (5.1) obtained in Theorem 5.2 by u^ε . Then u^ε extends to a time $T = T(\|u_0\|_s)$ independent of $\varepsilon > 0$.*

Proof. Using Theorem 5.2, part (ii), we know that there is T_ε so that $u^\varepsilon \in C([0, T_\varepsilon]: H^s) \cap X_{T_\varepsilon}^s$. On the other hand, it is not hard to verify that Theorem 1.2 and its proof generalize almost verbatim to solutions of (5.1) without that the time T for which the solution belongs to $L^\infty([0, T]: H^s(\mathbb{R}^3))$ depends on $\varepsilon > 0$. Therefore, by iterating Theorem 5.2 part (ii), we conclude the desired assertion. \square

5.2. Proof of Theorem 1.6. Finally, we prove that the L^2 norm (1.4) is preserved by the flow of (1.1) for all $s \in (3/4, 2)$, and that the energy (1.5) is conserved for $s \in (1, 2)$.

More specifically, we already know by the results in §3 and Lemma 4.3 and Theorem 1.2 that, whenever $u_0 \in H^s(\mathbb{R}^3)$, then there exists a solution $u \in C([0, T]: H^s(\mathbb{R}^3)) \cap X_T^s$, $T = T(\|u_0\|_s)$, and this solution is unique with such properties. We will prove that

Lemma 5.4. *Let u, u^ε be defined as before. Then, for each $s \in (3/4, 1)$, there is $T = T(\|u_0\|_s)$ so that $u^\varepsilon \rightarrow u$ in X_T^s as $\varepsilon \rightarrow 0$.*

Proof. Using the Duhamel formulation of (1.1) and (5.1), we can conclude that

$$\|u^\varepsilon - u\|_{X_T^s} \lesssim \left\| \int_0^t U(t-t') \partial_x (F_\varepsilon^{7/3}(u^\varepsilon) - u^{7/3})(t') dt' \right\|_{X_T^s}.$$

Now, the estimate (3.2), the second term on the right-hand side above is bounded by

$$(5.5) \quad \left\| 2^{sk} \|\Delta_k(\eta_\varepsilon * ((u^\varepsilon * \eta_\varepsilon)^{7/3} - (u * \eta_\varepsilon)^{7/3}))\|_{L_x^1 L_{y,T}^2} \right\|_{\ell^2(\mathbb{N})}$$

$$(5.6) \quad + \left\| 2^{sk} \|\Delta_k(\eta_\varepsilon * ((u * \eta_\varepsilon)^{7/3} - u^{7/3}))\|_{L_x^1 L_{y,T}^2} \right\|_{\ell^2(\mathbb{N})}$$

$$(5.7) \quad + \left\| 2^{sk} \|\Delta_k(\eta_\varepsilon * (u^{7/3}) - u^{7/3})\|_{L_x^1 L_{y,T}^2} \right\|_{\ell^2(\mathbb{N})}.$$

The first term (5.5) is clearly bounded by $CT^\delta \|u^\varepsilon - u\|_{X_T^s}$ by the methods from before, while the (5.6) is bounded by $CT^\delta \|(u * \eta_\varepsilon) - u\|_{X_T^s}$, which converges to zero by approximate identity properties, and (5.7) converges to zero by the fact that the summand within the $\ell^2(\mathbb{N})$ -norm is pointwise bounded by $2 \times 2^{sk} \|\Delta_k(u^{7/3})\|_{X_T^s}$, and by the properties of approximate identities converges pointwise to zero, which enables us to use the dominated convergence theorem.

This implies that $u^\varepsilon \rightarrow u$ in $X_T^s \cap C([0, T]: H^s(\mathbb{R}^3))$, $s \in (3/4, 1)$, as desired. \square

Next, we prove the conservation law for solutions of (5.1):

Lemma 5.5. *Let $u_0 \in H^s(\mathbb{R}^3)$, $s \in (3/4, 2)$, and u^ε be the solution to (5.1) given by the previous methods. The following assertions hold:*

(1) *The quantity*

$$(5.8) \quad M(u^\varepsilon(t)) = \|u^\varepsilon(t)\|_2^2$$

is constant for $t \in [0, T]$.

(2) *If, moreover, $s \in [1, 2)$, we have that the quantity*

$$(5.9) \quad E_\varepsilon(u^\varepsilon(t)) = \|\nabla u^\varepsilon(t)\|_2^2 - \frac{3}{5} \|(u^\varepsilon(t) * \eta_\varepsilon)\|_{10/3}^{10/3}$$

is constant for $t \in [0, T]$.

Proof. We first note that the proposition holds whenever we are dealing with *smooth* solutions to (5.1). Indeed, for (1), multiplying that equation by u^ε and integrating on \mathbb{R}^3 yields, together with a couple of integrations by parts, that the derivative of (5.8) is zero, which yields the claim. For (2), we integrate (5.1) against $\Delta u^\varepsilon + F_\varepsilon^{7/3}(u^\varepsilon)$ and integrations by parts in the same way as previously prescribed yields the result.

In order to pass to low regularity, we consider the sets

$$S_1 = \{t \in [0, T]: \|u^\varepsilon(t)\|_2^2 = \|u_0\|_2^2\}$$

and

$$S_2 = \left\{ t \in [0, T]: \|\nabla u^\varepsilon(t)\|_2^2 - \frac{3}{5}\|u^\varepsilon(t) * \eta_\varepsilon\|_{10/3}^{10/3} = \|\nabla u_0\|_2^2 - \frac{3}{5}\|u_0\|_{10/3}^{10/3} \right\}.$$

As we know that $u^\varepsilon \in C([0, T]: H^s)$ whenever $s \in (3/4, 2)$, then the expression defining (5.8) can be shown to be continuous with t , and thus $S_1 \subset [0, T]$ is closed. All we have to do it to show that it is also open. But this is a standard connectivity argument: if $t_1 \in S_1 \cap (0, T)$, then considering the IVP (5.1) with initial value $u^\varepsilon(t_1) * \varphi_\delta$ shows that the formal manipulations that we need in order to conclude that (5.8) hold rigorously for this new IVP on a neighbourhood of t_1 , independently of δ , by (5.4). Taking a limit as $\delta \rightarrow 0$ implies the desired result.

Similarly for S_2 , the expression in (5.9) is continuous in time for $t \in [0, T]$, by Theorem 5.2, Corollary 5.3 and the Gagliardo–Nirenberg interpolation inequality. As the set $S_2 \subset [0, T]$ is closed, we just need to conclude it is open once more. But this follows for $s \in [1, 2)$ by well-posedness of (5.1) in that range, together with an approximation argument as before. We skip the details. \square

Proof of Theorem 1.6. Part (i): Let first $s \in (3/4, 1)$. Then, it holds that, whenever $u_0 \in H^s$,

$$\|u^\varepsilon(t)\|_2 = \|u_0\|_2^2, \quad \forall t \in [0, T].$$

By Theorem 5.2, the existence time T above depends only on the H^s -norm of the initial datum. Now, by Lemma 5.4, we have that, for such T ,

$$\|u^\varepsilon - u\|_{X_T^s} \rightarrow 0 \Rightarrow \|u_0\|_2^2 = \|u^\varepsilon(t)\|_2^2 \rightarrow \|u(t)\|_2^2 \text{ as } \varepsilon \rightarrow 0.$$

This finishes the proof in this case. For $s \in [1, 2)$, we simply use that H^s are nested, and thus we can reduce to the previous case.

Part (ii): Assume first, by mollification, that $u_0 \in H^3(\mathbb{R}^3)$. We know, from Theorem 1.2 applied to solutions of (1.1) and (5.1), Corollary 1.3 and Lemma 5.4, that

$$\|u^\varepsilon - u\|_{X_T^s} \lesssim 4C_s \|u_0\|_s,$$

whenever $s \in [1, 2)$, and

$$\|u^\varepsilon - u\|_{X_T^{\tilde{s}}} \rightarrow 0, \quad \tilde{s} \in (3/4, 1),$$

as $\varepsilon \rightarrow 0$, and $T = T(\|u_0\|_s)$ is sufficiently small. Interpolating between these two estimates implies that $u^\varepsilon \rightarrow u$ in $X_T^s \cap C([0, T]: H^s)$ whenever $s \in [1, 2)$. Therefore, whenever $s \in [1, 2)$, we see that

$$E_\varepsilon(u_0) = E_\varepsilon(u^\varepsilon(t)) \rightarrow E(u(t)) \text{ as } \varepsilon \rightarrow 0 \text{ for } t \in [0, T]$$

follows from the aforementioned convergence and the Gagliardo–Nirenberg interpolation inequality. The left-hand side above, however, converges by approximate identity properties to $E(u_0)$.

Now, for general $u_0 \in H^s(\mathbb{R}^3)$, $s \in (1, 2)$, we solve (1.1) for initial data $u_{0,\varepsilon} = u_0 * \eta_\varepsilon$. Using the notation from Theorem 1.2, it holds that

$$(5.10) \quad E(u_{0,\varepsilon}) = E(u_\varepsilon(t)), \quad \forall t \in [0, T].$$

On the other hand, from Theorem 1.2 itself and properties of approximate identities, the left-hand side of (5.10) converges to $E(u_0)$, while the right-hand side converges to $E(u(t))$, for each $t \in [0, T]$ as long as $s \in (1, 2)$. This concludes the proof of the result. \square

5.3. Proof of Theorem 1.7. In order to prove Theorem 1.7, we need to work once more with the approximations $u_{0,\varepsilon}$ to our initial data. Theorem 1.6 shows that the solutions u_ε satisfy the conservation of energy (1.5) for $t \in [0, T], T = T(\|u_{0,\varepsilon}\|_1) \gtrsim T(\|u_0\|_1)$. Therefore, for $t \in [0, T]$,

$$\|\nabla u_\varepsilon(t)\|_2^2 \leq \|u_0\|_1^2 + \frac{3}{5}\|u_\varepsilon(t)\|_{10/3}^{10/3} \leq \|u_0\|_1^2 + \frac{3}{5}C_{opt}^{10/3}\|\nabla u_\varepsilon(t)\|_2^2\|u_0\|_2^{4/3},$$

where C_{opt} denotes the best constant in the Gagliardo–Nirenberg interpolation inequality

$$\|w\|_{10/3} \leq C_{opt}\|\nabla w\|_2^{3/5}\|w\|_2^{2/5}.$$

Therefore, if $\|u_0\|_2 < (\frac{5}{3})^{3/4} \frac{1}{C_{opt}^{5/2}}$, using conservation of (1.4) for u_ε , we get that

$$(5.11) \quad \|u_\varepsilon(t)\|_{H^1} \leq C_{\|u_0\|_2, \|u_0\|_{H^1}},$$

for all times t in which u_ε is well-defined. This proves already that, by reiterating Theorem 1.2 and Corollary 1.3, $u_\varepsilon \in (C \cap L^\infty)(\mathbb{R}; H^1(\mathbb{R}^3))$. In particular, by Lemma 4.2, there is another universal constant $\tilde{C}_{\|u_0\|_2, \|u_0\|_{H^1}}$ so that, for each $t_0 \in \mathbb{R}$,

$$(5.12) \quad \|u_\varepsilon(\cdot + t_0)\|_{X_T^1} \leq \tilde{C}_{\|u_0\|_2, \|u_0\|_{H^1}},$$

for each T sufficiently small depending only on $\|u_0\|_2, \|u_0\|_{H^1}$. Applying (5.11) and the standard functional analysis arguments mentioned before, we get that the solution u extends to all the real line as a function in $L^\infty(\mathbb{R}; H^s)$. In addition to it, by Corollary 1.3 applied to (1.1) with initial data $u(t_0)$ iteratively, we get that

$$\|u(\cdot + t_0)\|_{X_T^1} \leq \tilde{C}_{\|u_0\|_2, \|u_0\|_{H^1}}.$$

This implies, in particular, that $u \in C([t_0, t_0 + T]; H^1(\mathbb{R}^3))$, and as $t_0 \in \mathbb{R}$ was arbitrary, it holds that $u \in (L^\infty \cap C)(\mathbb{R}; H^1(\mathbb{R}^3))$, as desired.

We remark that, by the results in [38] (see also [31, Chapter 6]), we have that $\frac{1}{C_{opt}} = \frac{\|\phi\|_2^{2/5}}{(5/3)^{3/10}}$, where $\phi \in H^1(\mathbb{R}^3)$ is some positive, radial solution to

$$(5.13) \quad \Delta\phi - \phi + \phi^{7/3} = 0.$$

Therefore, the above arguments work whenever

$$\|u_0\|_2 < \|\phi\|_2.$$

This gives us an explicit quantification of how small the initial data has to be in order to have global existence results.

5.4. Consequences of the analysis of (5.1): Pointwise convergence to the initial data. At last, we prove that for each $u_0 \in H^s, s > \frac{3}{4}$, the flow of (1.1) converges *pointwise* to u_0 as $t \rightarrow 0$. In order to do so, we employ classical ideas from the theory of maximal functions, which have been recently employed in the context of nonlinear Schrödinger equations by Compaan, Lucà and Staffilani [7]. See also [12, 13] for previous work by

Erdoğan and Tzirakis on convergence results for the KdV equation and the nonlinear Schrödinger in one dimension.

We begin with a standard lemma translating maximal estimates into pointwise consequences. Here we note that we will work with (5.1), but we pick carefully the function η and the initial data. In fact, we can choose it so that $\eta_\varepsilon * f(z) = P_{1/\varepsilon}f(z)$, with $P_N = \sum_{j \leq N} \Delta_j$ defined as before, and we let $u_{0,\varepsilon} = \eta_\varepsilon * u_0$ be the initial data we work with. We will abuse notation and still denote by u^ε the solution to this new version of (5.1).

Notice that this specification does not change any of the previously discussed properties of the solutions u^ε to (5.1).

Lemma 5.6. *Suppose that, for $u_0 \in H^s(\mathbb{R}^3)$, we have that*

$$\|u^\varepsilon - u\|_{L^4_{x,y}L^\infty_T} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Then it holds that $u(z, t) \rightarrow u_0(z)$ as $t \rightarrow 0$ for almost every $z \in \mathbb{R}^3$.

Proof. The proof is analogous to that of [33, Lemma 3.8], so we omit it. \square

Proof of Theorem 1.8. We focus on proving the asserted convergence from Lemma 5.6. Indeed, an analogous strategy to that of Proposition 2.4 (iii) implies

$$\left\| \int_0^t U(t-t') \partial_x \Delta_k f(t') dt' \right\|_{L^4_{x,y}L^\infty_T} \lesssim 2^{(3/4)+k} \|\Delta_k f\|_{L^1_x L^2_{y,T}}.$$

By the Duhamel formulations of (1.1) and (5.1), we get that $\|u - u^\varepsilon\|_{L^4_{x,y}L^\infty_T}$ is bounded by

$$(5.14) \quad C_s \|(I - P_{1/\varepsilon})u\|_{H^s} + \left\| 2^{sj} \|\Delta_j (\eta_\varepsilon * (u^\varepsilon * \eta_\varepsilon)^{7/3} - u^{7/3})\|_{L^1_x L^2_{y,T}} \right\|_{\ell^2_j},$$

where $s > \frac{3}{4}$. By using the same decomposition as in Lemma 5.4, the second term in (5.14) is bounded by the sum

$$(5.5) + (5.6) + (5.7) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0,$$

where we argue exactly as in Lemma 5.4, now keeping in mind that the correction term $\|(I - P_{1/\varepsilon})u_0\|_{H^s} \rightarrow 0$ as $\varepsilon \rightarrow 0$ if $u_0 \in H^s(\mathbb{R}^3)$. This concludes the proof, and by Lemma 5.6 we deduce the desired pointwise convergence of the flow. \square

ACKNOWLEDGEMENTS

F. L. was partially supported by CNPq and FAPERJ, Brazil. J.P.G.R. acknowledges financial support from CNPq, Brazil.

REFERENCES

- [1] D. Bhattacharya, L.G. Farah, and S. Roudenko, Global well-posedness for low regularity data in the 2d modified Zakharov-Kuznetsov equation. arXiv:1906.05822.
- [2] W. Baoxiang, H. Lijia and H. Chunyan, *Global well-posedness and scattering for the derivative nonlinear Schrödinger equation with small rough data*. Ann. Inst. H. Poincaré - AN. **26** (2009), 2253–2281.

- [3] H. A. Biagioni and F. Linares, *Well-posedness results for the modified Zakharov-Kuznetsov equation*. Nonlinear equations: methods, models and applications (Bergamo, 2001), volume 54 of Progr. Nonlinear Differential Equations Appl., pages 181–189. Birkhäuser, Basel, 2003
- [4] J. L. Bona and R. Smith, *The initial-value problem for the Korteweg-de Vries equation*. Philos. Trans. Roy. Soc. London Ser. A, **278** (1975), 555–601.
- [5] L. Carleson. *Some analytic problems related to statistical mechanics*. Lecture Notes in Math No. 779, Springer-Verlag. Berlin and New York. 1979, pp. 9–45.
- [6] M. Christ and A. Kiselev, *Maximal functions associated to filtrations*. J. Func. Anal. **179** (2001), n. 2, 409–425.
- [7] E. Compaan, R. Lucà and G. Staffilani, *Pointwise Convergence of the Schrödinger flow*. Int. Math. Res. Notices (in press).
- [8] R. Côte, C. Muñoz, D. Pilod, and G. Simpson, *Asymptotic Stability of high-dimensional Zakharov-Kuznetsov solitons*. Arch. Ration. Mech. Anal. **220** (2016), no. 2, 639-710.
- [9] de Bouard, A. *Stability and instability of some nonlinear dispersive solitary waves in higher dimension*. Proc. Roy. Soc. Edinburgh Sect. A **126** (1996), no. 1, 89–112.
- [10] Y. Deng and P. Germain, *Growth of solutions to the NLS on irrational tori*. Int. Math. Res. Notices, **72** (2019), n. 9, 2919–2950.
- [11] M. B. Erdoğan and N. Tzirakis, *Dispersive Partial Differential Equations: Well-posedness and applications*. London Mathematical Society Student Texts **86**, Cambridge University Press, 2016.
- [12] M.B. Erdoğan and N. Tzirakis, *Global smoothing for the periodic KdV evolution*. Int. Math. Res. Notices **20** (2013), 4589–4614.
- [13] M.B. Erdoğan and N. Tzirakis, *Talbot effect for the cubic non-linear Schrödinger equation on the torus*. Math. Res. Lett. **20** (2013), n. 6, 1081–1090.
- [14] A. V. Faminskii, *The Cauchy problem for the Zakharov-Kuznetsov equation*. Differential' nye Uravneniya **31** (1995), n. 6, 1070–1081.
- [15] L.G. Farah, F. Linares and A. Pastor, *A note on the 2D generalized Zakharov-Kuznetsov equation: Local, global, and scattering results*. J. Diff. Eqs. **253** (2012), n. 8, 2558–2571.
- [16] L.G. Farah, J. Holmer, S. Roudenko, and Kai Yang, *Instability of solitons in the 2d cubic Zakharov-Kuznetsov equation*. preprint, arXiv: 1711.05907.
- [17] L.G. Farah, J. Holmer, S. Roudenko, and Kai Yang, *Blow-up in finite or infinite time of the 2D cubic Zakharov-Kuznetsov equation*. preprint, arXiv:1810.05121.
- [18] J. Ginibre and G. Velo, *On a Class of Nonlinear Schrödinger Equations. I. The Cauchy Problem, General Case*. J. Func. Anal **32** (1979), 1–32.
- [19] L. Grafakos, *Modern Fourier Analysis (third edition)*. Graduate Texts in Mathematics, **250**, Springer-Verlag, 2015.
- [20] A. Grünrock, *A Remark on the modified Zakharov-Kuznetsov equation in three space dimensions*. Math. Res. Lett. **21** (2014), n. 1, 127–131.
- [21] A. Grünrock, *On the generalized Zakharov-Kuznetsov equation at critical regularity*, preprint, arXiv:1509.09146.
- [22] A. Grünrock and S. Herr, *The Fourier restriction norm method for the Zakharov-Kuznetsov equation*. Disc. Cont. Din. Syst. **34** (2014), n. 5, 2061–2068.
- [23] D. Han-Kwan, *From Vlasov-Poisson to Korteweg-de Vries and Zakharov-Kuznetsov*. Comm. Math. Phys. **324** (2013), 961–993.
- [24] S. Herr and S. Kinoshita, *Subcritical well-posedness results for the Zakharov-Kuznetsov equation in dimension three and higher*. arXiv preprint: arXiv:2001.09047.
- [25] C.E. Kenig, G. Ponce, L. Vega, *Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle*, Commun. Pure Appl. Math. **46** (1993), 527–620.
- [26] S. Kinoshita, *Global Well-posedness for the Cauchy problem of the Zakharov-Kuznetsov equation in 2D*. arXiv preprint: arXiv:1905.01490.

- [27] S. Kinoshita, *Well-posedness for the Cauchy problem of the modified Zakharov-Kuznetsov equation*. arXiv preprint: arXiv:1911.13265.
- [28] D. Lannes, F. Linares and J.-C. Saut, *The Cauchy problem for the Euler-Poisson system and derivation of the Zakharov-Kuznetsov equation*. Prog. Non-linear Diff. Eqs Appl. **84** (2013), 181–213.
- [29] F. Linares and A. Pastor, *Well-posedness for the two-dimensional modified Zakharov-Kuznetsov equation*. SIAM J. Math. Anal. **41** (2009), No. 4, 1323–1339.
- [30] F. Linares and A. Pastor, *Local and global well-posedness for the 2D generalized Zakharov-Kuznetsov equation*. J. Funct. Anal., 260(4) (2011), 106–1085.
- [31] F. Linares and G. Ponce, *Introduction to Nonlinear Dispersive Equations (second edition)*. Universitext, Springer-Verlag, 2015.
- [32] F. Linares and J.-C. Saut, *The Cauchy problem for the 3D Zakharov-Kuznetsov equation*. Disc. Cont. Dyn. Systems - A **24** (2009), n. 2, 547–565.
- [33] F. Linares and J.P.G. Ramos, *Maximal function estimates and local well-posedness for the generalized Zakharov-Kuznetsov equation*. Preprint.
- [34] L. Molinet and D. Pilod, *Bilinear Strichartz estimates for the Zakharov-Kuznetsov equation and applications*. Annales de l'Institut Henri Poincaré (C) Non Linear Analysis **32** (2015), n. 2, 347–371.
- [35] F. Ribaud and S. Vento, *A Note on the Cauchy problem for the 2D generalized Zakharov-Kuznetsov equations*. C. R. Acad. Sci. Paris, Ser. I, **350** (2012), 499–503.
- [36] F. Ribaud and S. Vento, *Well-posedness results for the three-dimensional Zakharov-Kuznetsov equation*. SIAM J. Math. Anal. **44** (2012), n. 4, 2289–2304.
- [37] T. Tao, *Nonlinear Dispersive Equations: Local and Global Analysis*. NSF-CBMF Regional Conference Series in Mathematics, Number 106.
- [38] M. Weinstein, *Nonlinear Schrödinger Equations and Sharp Interpolation Estimates*. Commun. Math. Phys. **87** (1983), 567–576.
- [39] V. E. Zakharov and E. A. Kuznetsov, *On three dimensional solitons*. Sov. Phys. JETP. **39** (1974) 285–286.

(F. Linares) IMPA, INSTITUTO MATEMÁTICA PURA E APLICADA, ESTRADA DONA CASTORINA 110, 22460-320, RIO DE JANEIRO, RJ, BRAZIL

E-mail address: `linares@impa.br`

(J.P. Ramos) IMPA, INSTITUTO MATEMÁTICA PURA E APLICADA, ESTRADA DONA CASTORINA 110, 22460-320, RIO DE JANEIRO, RJ, BRAZIL

E-mail address: `joaopgramos95@gmail.com`