

APPLICATIONS OF ALGEBRAIC COMBINATORICS TO ALGEBRAIC GEOMETRY

DAVID KAZHDAN AND TAMAR ZIEGLER

Dedicated to the memory of Tonny Springer

ABSTRACT. We formulate a number of new results in Algebraic Geometry and outline their derivation from Theorem 2.12 which belongs to Algebraic Combinatorics.

1. INTRODUCTION

In this paper we present a derivation of a number of results in Algebraic Geometry from a theorem (Theorem 2.12) in Algebraic Combinatorics and its extension (Theorem 8.3) to the case of p -adic rings. In the first half of the article we outline the derivations of results in Algebraic Geometry relying on results of [10] and [11]. Afterwards we present a proof of an extension of Theorem 2.12 to the p -adic realm and an application of this result to Algebraic Geometry.

1.1. Basic definitions. We will denote finite fields by k and general fields by K . It will be important to distinguish between an algebraic variety \mathbf{X} defined over a field K and the set $X := \mathbf{X}(K)$ of K -points of \mathbf{X} . For a polynomial $P : V \rightarrow K$ we write $\mathbf{X}_P := P^{-1}(0) \subset \mathbf{V}$ and $X_P := P^{-1}(0) \subset V$. So $X_P = \mathbf{X}_P(K)$.

In this paper we heavily use the notion of rank of polynomials introduced in a paper by Schmidt [20], which characterizes complexity of polynomials.¹

Definition 1.1 (Schmidt rank). Let P be a polynomial of degree d on a K -vector space V . We define the *rank* $r_K(P)$ as the minimal number r such that P can be written in the form $P = \sum_{i=1}^r Q_i R_i$, where Q_i, R_i are K -polynomials on V of degrees $< d$.

When there is no confusion we will omit the subscript K from $r_K(P)$.

Example 1.2. If $P : V \rightarrow K$ is a non-degenerate quadratic form then $\dim(V)/2 \leq r(P) \leq 3\dim(V)/2$.

We show that morphisms $\mathbf{P} : \mathbf{V} \rightarrow \mathbf{A}^c$ defined by polynomials of sufficiently high rank and degree less than characteristic of K possess a number of nice properties. In particular we show that these morphisms are flat and that their

The second author is supported by ERC grant ErgComNum 682150.

¹This notion of complexity was also introduced later in the work of Ananyan and Hochster [1] where it is called *strength*.

fibers are complete intersections with rational singularities. In the case of fields of small characteristic we have to replace the rank $r(P)$ by the non-classical rank $r_{nc}(P)$ defined in 2.3.

For polynomials over finite fields there is also a notion of an analytic rank.

Definition 1.3 (Analytic rank). Let $k = \mathbb{F}_q$ and P be a polynomial of degree d on a k -vector space V . Fix a non trivial additive character $\psi : k \rightarrow \mathbb{C}^*$ and define the *analytic rank* $a_k(P) := -\log_q |q^{-\dim V} \sum_{v \in V} \psi(P(v))|$.

To simplify notations we assume throughout most of the introduction that $V = V_1 \times \cdots \times V_d$ and that polynomials $P : V \rightarrow K$ are multilinear. In this case the analytic rank does not depend on a choice a non-trivial additive character.

Most of the results in this paper are based on the following Theorem relating rank and analytic rank for multilinear polynomials over finite fields (see [3, 19]):

Fact 1.4. *There exist a function $\gamma_d(s)$, $\lim_{s \rightarrow \infty} \gamma_d(s) = \infty$, such that for any finite field k , and degree d multilinear polynomial P we have $a_k(P) \geq \gamma_d(r_k(P))$.*

1.2. Basic conjectures and questions. Before describing our results we formulate some basic questions and conjectures regarding the notions of rank defined above.

Conjecture 1.5 (d). For any $d \geq 2$ there exists $\kappa_d > 0$ such that for any multilinear polynomial P we have $r_K(P) \leq \kappa_d r_{\bar{K}}(P)$ where \bar{K} is the algebraic closure of K .

Remark 1.6.

- (1) It is easy to see that Conjecture 1.5 holds for $d = 2$ with $\kappa_2 = 1/2$.
- (2) Since in the case when $d = 3$ the Schmidt rank is equal to the *slice* rank the validity of Conjecture 1.5(3) follows from Theorem 2.5 and Proposition 4.9 in [6].

Conjecture 1.7 (d). For any $d \geq 2$ there exists $\epsilon_d > 0$ such that $a_{\mathbb{F}_q}(P) \geq \epsilon_d r_{\mathbb{F}_q}(P)$ for any multilinear polynomial of degree d .

Remark 1.8.

- (1) The inequality $a_{\mathbb{F}_q}(P) \leq r_{\mathbb{F}_q}(P)$ is known. See [13, 16].
- (2) It is easy to see that Conjecture 1.7 holds for $d = 2$ with $\epsilon_2 = 1/2$.

Let K be an algebraically close field, and $(P_s)_{s \in S}$ be a family of polynomials. We define $U \subset S$ as the subset of points $t \in S$ such that $r(P_t) = \max_{s \in S} r(P_s)$. It is easy to see that this subset is constructible.

Question 1.9. Is the subset $U \subset S$ locally closed?

1.3. Main results. We formulate in the introduction special cases of the main results. To simplify notations we assume that K is an algebraically closed field of characteristic $> d = \deg P$. For the validity of these results we do not have to assume that polynomials P are multilinear.

Theorem 1.10. *For any $d \geq 2$ there exists $r(d)$ such that varieties $\mathbf{X}_P \subset \mathbf{V}$ are geometrically irreducible normal varieties with rational singularities.*

For a formulation of two other results we need a couple of definitions.

Definition 1.11. A polynomial P of degree d on a K -vector space K^m is m -universal if for any $Q \in K[x_1, \dots, x_m]$ of degrees $\leq d$ there exists an affine map $\phi : K^m \rightarrow V$ such that $Q = \phi^*(P)$.

Theorem 1.12 (Universal). *There exists a function $R(d, m)$ such that any polynomial P of degree d and rank $\geq R(d, m)$ is m -universal.*

Definition 1.13. Let V be a K -vector space and X a subset of V . A function $f : X \rightarrow K$ is weakly polynomial of degree $\leq a$, if for any affine subspace $L \subset X$ the restriction of f on L is a polynomial of degree $\leq a$.

Theorem 1.14. *For any $a, d \geq 1$ there exists $r(a, d)$ such that the following holds.*

For any polynomial $P : V \rightarrow K$ of degree d and rank $\geq r(a, d)$, any weakly polynomial function of degree $\leq a$ on $X_P \subset V$ is a restriction of a polynomial of degree $\leq a$ on V .

The last result we state is for polynomials over finite fields \mathbb{F}_q .

Theorem 1.15. *There exists a function $\rho(d) > 0$ such that for any finite field \mathbb{F}_q of characteristic $> d$ and polynomials $P, Q \in \mathbb{F}_q[x_1, \dots, x_N]$ such that $\deg(P) = d$, $r_{\mathbb{F}_q}(P) \geq \rho(d)$, $\deg(Q) \leq q/d$ and such that $Q(\bar{x}) = 0$ for all $\bar{x} \in \mathbb{F}_q^N$ such that $P(\bar{x}) = 0$, we have $Q = PR$ for some $R \in \mathbb{F}_q[x_1, \dots, x_N]$.*

In the paper all the above mentioned results are extended to the case when we replace a polynomial P by a collection $\bar{P} = (P_1, \dots, P_c)$ of polynomials on V .

1.4. Acknowledgement. We thank U. Hrushovski for his help with simplifying the proof of Theorem 7.2.

2. THE MAIN TOOL

We start with defining the Schimdt rank for a family of polynomials.

Definition 2.1 (Schmidt rank). (1) Let P be a polynomial of degree d on a K -vector space V . We define the rank $r_K(P)$ as the minimal number r such that P can be written in the form $P = \sum_{i=1}^r Q_i R_i$, where Q_i, R_i are polynomials on V of degrees $< d$.

(2) For a collection of polynomials $\bar{P} = (P_i)_{1 \leq i \leq c}$ on V we define the rank $r_k(\bar{P})$ as the minimal rank of polynomials $P_{\bar{a}} := \sum_{i=1}^c a_i P_i$, $\bar{a} \in K^c \setminus \{0\}$.

Remark 2.2. (1) We will often consider \bar{P} as a morphism from \mathbf{V} to \mathbf{A}^c .

(2) When there is no confusion we write $r(\bar{P})$ instead of $r_k(\bar{P})$.

When the characteristic of the field is smaller then the maximal degree in \bar{P} we need to replace the notion of rank with non-classical rank.

Definition 2.3. Let P be a polynomial of degree d on a K -vector space V .

- (1) We denote by $\tilde{P} : V^d \rightarrow k$ the symmetric multilinear form given by $\tilde{P}(h_1, \dots, h_d) := \Delta_{h_1} \dots \Delta_{h_d} P : V^d \rightarrow k$, where $\Delta_h P(x) := P(x + h) - P(x)$. Note that \tilde{P} only depends on the top-degree part of P .
- (2) We define the *non-classical rank* (*nc-rank*) $r_{nc}(P)$ to be the rank of \tilde{P} .
- (3) For a collection $\bar{P} = (P_i)_{1 \leq i \leq c}$ of polynomials on V we define the nc-rank $r_{nc}(\bar{P})$ as the minimal nc-rank of polynomials $P_{\bar{a}} := \sum_{i=1}^c a_i P_i$, $\bar{a} \in k^c \setminus \{0\}$.

Remark 2.4. (1) If $\text{char}(K) > d$ then $r(P) \sim r_{nc}(P)$ (see [10]).

- (2) In low characteristic it can happen that P is of high rank while \tilde{P} is of low rank.

Example 2.5. Let K be a field of characteristic 2, $V = \mathbf{A}^n$ and $P(x_1, \dots, x_n) = \sum_{1 < i < j < k < l \leq n} x_i x_j x_k x_l$ is of rank $\sim n$, but of nc-rank 3, (see [21]).

Definition 2.6. Let $k := \mathbb{F}_q$, V be a k -vector space and $\bar{P} : V \rightarrow k^c$ be a map. We define

- (1) $F_{\bar{t}}(\bar{P}) := \{v \in V \mid \bar{P}(v) = \bar{t}\}$, $\bar{t} \in k^c$.
- (2) $\nu_{\bar{P}} : k^c \rightarrow \mathbb{C}$ is the function given by $\nu_{\bar{P}}(\bar{t}) := |F_{\bar{t}}(\bar{P})|/q^{\dim(V)-c}$.
- (3) A map \bar{P} is *s-uniform* if $|\nu_{\bar{P}}(\bar{t}) - 1| \leq q^{-s}$ for all $\bar{t} \in k^c$.

Lemma 2.7. Let $\bar{P} = (P_1, \dots, P_c)$, $P_i : V \rightarrow k$ be a collection of maps such that all the maps $P_{\bar{a}} := \sum_{i=1}^c a_i P_i$, $\bar{a} \in k^c \setminus \{0\}$ are *s-uniform*. Then the map \bar{P} is $(s - c)$ -uniform.

Proof. Let A be a finite commutative group written additively and Ξ the group of characters $\chi : A \rightarrow \mathbb{C}^*$ of A . For a function $f : A \rightarrow \mathbb{C}$ we define the Fourier transform $\hat{f}(\chi) : \Xi \rightarrow \mathbb{C}$ by $\hat{f}(\chi) = |A|^{-1} \sum_{a \in A} \chi(a) f(a)$. The next statement is the formula for the inverse Fourier transform.

Fact 2.8. $\sum_{\chi \in \Xi} \hat{f}(\chi) \chi(-a_0) = f(a_0)$ for any $a_0 \in A$.

Let $A = k^c$. We fix a non-trivial additive character $\psi : k \rightarrow \mathbb{C}^*$ and associate with any $\bar{a} \in k^c$ a character $\chi_{\bar{a}} : k^c \rightarrow \mathbb{C}^*$ by $\chi_{\bar{a}}(\bar{t}) = \psi(\langle \bar{a}, \bar{t} \rangle)$ where $\langle \cdot, \cdot \rangle : k^c \times k^c \rightarrow k$ is the natural pairing. The map $\bar{a} \rightarrow \chi_{\bar{a}}$ is a group isomorphism between A and the group Ξ of characters of A .

The Fourier transform of the function ν is given by

$$\hat{\nu}_{\bar{P}}(\chi_{\bar{a}}) = q^{-c} \sum_{\bar{t} \in k^c} \nu_{\bar{P}}(\bar{t}) \chi(\bar{t}) = q^{-\dim(V)} \sum_{v \in V} \psi(P_{\bar{a}}(v))$$

Since, by the assumptions of the Lemma, for any $\bar{a} \neq 0$ the map $P_{\bar{a}}$ is *s-uniform* we see that $|\hat{\nu}_{\bar{P}}(\chi_{\bar{a}})| \leq q^{-s}$ for $\bar{a} \neq 0$.

As follows from Claim 2.8 we have

$$\nu_{\bar{P}}(\bar{t}) = \sum_{\bar{a} \in k^c} \psi_{\bar{a}}(-\bar{t}) \hat{\nu}_{\bar{P}}(\bar{a}) = 1 + \sum_{\bar{a} \in k^c \setminus \{0\}} \psi_{\bar{a}}(-\bar{t}) \hat{\nu}_{\bar{P}}(\bar{a})$$

for all $\bar{t} \in k^c$. So $|\nu_{\bar{P}}(\bar{t}) - 1| \leq q^{-(s-c)}$. \square

The following result is from Algebraic Combinatorics (see [3], [19]).

Fact 2.9. *There exist a function $\alpha_d(s)$, $\lim_{s \rightarrow \infty} \alpha_d(s) = \infty$, such that all polynomials $P : V \rightarrow k$ of degree d and $r_{nc}(P) \geq \alpha_d(s)$ are s -uniform.*

Remark 2.10. (1) Fact 2.9 is the corner stone of our work.

- (2) Note that $\alpha_d(s)$ is independent of the finite field k and of the (dimension of the) k -vector space V .
- (3) A stronger form of this result showing the existence of explicitly definable functions $e(d), \gamma(d)$ such that one can take $\alpha_d(s) = e(d)s^{\gamma(d)}$ is proven in [19] and [8]. This improvement has a number of important applications in Algebraic Combinatorics. One can ask whether it is possible to replace $\gamma(d)$ by 1.

Definition 2.11. For $\bar{d} = (d_1, \dots, d_c)$ we write $r_{\bar{d}} := \alpha_d(c+2)$.

The following result, to which we often refer, follows immediately from Lemma 2.7 and Claim 2.9.

Theorem 2.12 (Uniform). *Let $\bar{P} = (P_1, \dots, P_c)$ be a collection of polynomials of degrees $\leq d$ and of nc-rank $r \geq r_{\bar{d}}$. Then $|\nu_{\bar{P}}(\bar{t}) - q^{-c}| \leq q^{-(c+2)}$ for all $\bar{t} \in k^c$.*

3. IRREDUCIBILITY OF FIBERS

In this section we show a derivation of the following result from Theorem 2.12.

Theorem 3.1. (1) *For any field K and a collection $\bar{P} = (P_1, \dots, P_c)$, of polynomials $P_i \in K[x_1, \dots, x_n]$ of degrees $\bar{d} = (d_1, \dots, d_c)$ and nc-rank over the algebraic closure of K is greater than $r_{\bar{d}}$, all the fibers $\mathbf{F}_{\bar{t}}(\bar{P}) := \bar{\mathbf{P}}^{-1}(\bar{t}) \subset \mathbf{A}^n$, $\bar{t} \in K^c$ are non-empty reduced, geometrically irreducible complete intersections.*

- (2) *The schemes $\mathbf{F}_{\bar{t}}(\bar{P})$ are reduced.*
- (3) *The schemes $\mathbf{F}_{\bar{t}}(\bar{P})$ are normal.*
- (4) *If $K = \mathbb{R}$ and all degrees $d_i, 1 \leq i \leq c$ are odd, then $\mathbf{F}_{\bar{t}}(\bar{P})(\mathbb{R}) \neq \emptyset, \bar{t} \in \mathbb{R}^c$.*

Remark 3.2. For fields of characteristic zero Theorem 3.1 follows from [20]. We include proofs to present our technique in details in the simplest case.

Proof. We first prove part (1). As is well known (see Krull's principal theorem, [17]) any irreducible component \mathbf{Y} of $\mathbf{F}_{\bar{t}}(\bar{P})$ is of dimension $\geq (n - c)$. So it is sufficient to show that the varieties $\mathbf{F}_{\bar{t}}(\bar{P})$ are irreducible and of dimension $\leq (n - c)$.

We first consider the case when K is a finite field when we can use the following result (see [15]).

Let $k := \mathbb{F}_q$, $k_l := \mathbb{F}_{q^l}$. Let \mathbf{X} be an m -dimensional algebraic variety defined over k and $c(\mathbf{X})$ be the number of irreducible components of \mathbf{X} of dimension m (considered as a variety over the algebraic closure \bar{k} of k). We define $\tau_l(\mathbf{X}) := \frac{|\mathbf{X}(k_l)|}{q^{ml}}$, for $l \geq 1$.

Claim 3.3. There exists $u \geq 1$ such that $\lim_{l \rightarrow \infty} \tau_{lu}(\mathbf{X}) = c(\mathbf{X})$.

For a proof of Theorem 3.1 in the case when $K = \mathbf{F}_q$ it is sufficient to observe that Claim 3.3 and Theorem 2.12 imply that $\dim(\mathbf{F}_{\bar{t}}(\bar{P})) = n - c$ and $c(\mathbf{F}_{\bar{t}}(\bar{P})) = 1$.

Now we consider the case when K is an algebraic closure of \mathbf{F}_q . Since $K = \bigcup \mathbf{F}_{q^n}$ we may assume that $\bar{t} \in \mathbf{F}_{q^n}^c$. So $\dim(\mathbf{F}_{\bar{t}}(\bar{P})) = n - c$ and $c(\mathbf{F}_{\bar{t}}(\bar{P})) = 1$. Therefore Theorem 3.1 is proven in the case when K is an algebraic closure of a finite field.

We start the reduction of the general case of Theorem 3.1 to the case when K is an algebraic closure of a finite field with a reformulation.

Definition 3.4. Let $\bar{d} = (d_1, \dots, d_c)$ and $n \geq 1$. A field K has property $\star(n, \bar{d})$ if for any collection $\bar{P} = (P_i)_{i=1}^c$ of polynomials $P_i \in K[x_1, \dots, x_n]$ of degrees d_i and $nc\text{-rank} \geq r_{\bar{d}}$, all the varieties $\mathbb{F}_{\bar{t}}$, $\bar{t} \in K^c$ are irreducible of dimension $\dim(V) - c$, where $r_{\bar{d}}$ is as in Definition 2.11.

Claim 3.5 (n). All fields have the property $\star(n, \bar{d})$ for any $n \geq 1$, $\bar{d} = (d_1, \dots, d_c)$.

Proof. It is clear that it is sufficient to prove this claim for algebraic closed fields. Our proof of Claim 3.5(n) uses the following result from Model theory (see [18]).

Fact 3.6. *Let T be the theory of algebraically closed fields. Then any first order property in T which is true for all algebraic closures of finite fields is true for all algebraically closed fields.*

Since $\star(n, \bar{d})$ is a first order property in T , Claim 3.5(n), $n \geq 1$ are proven. \square

Since the constant $r(\bar{d})$ does not depend on n , the validity of Claim 3.5(n) for all $n \geq 1$, implies the validity of the part (1) of Theorem 3.1.

Our proof of parts (2) and (3) is based on the following statement (see [14]).

Fact 3.7. *There exists a function $\tilde{r}(r, \bar{d})$ such that for any collection \bar{P} of degree \bar{d} and $nc\text{-rank} \geq \tilde{r}(r, \bar{d})$ there exist $v_i, w_i \in V$, $1 \leq i \leq c$ such that the collection $\bar{Q} = (\bar{P}, P_i/\partial_{v_i}, \partial P_i/\partial_{w_i})_{1 \leq i \leq c}$ is of $nc\text{-rank} \geq r$.*

(2) Since (as follows from (1)) varieties $\mathbf{F}_{\bar{t}}(\bar{P})$ are irreducible complete intersections, to show that schemes $\mathbf{F}_{\bar{t}}(\bar{P})$ are reduced it is sufficient to show that they are reduced at the generic point, and to show the normality of $\mathbf{F}_{\bar{t}}(\bar{P})$ it is sufficient to show that they are non singular in codimension 1.

To show that a variety $\mathbf{F}_{\bar{t}}(\bar{P})$ is reduced at the generic point it is sufficient to show the existence of vectors $v_1, \dots, v_c \in V$ such that $\partial P_i / \partial v_i(x) \neq 0, x \in \mathbf{F}_{\bar{t}}(\bar{P})$.

For $\bar{v} \in V^c$ we define the collection $\bar{R}(\bar{v})$ by $\bar{R}(\bar{v}) = (\bar{P}, \partial P_i / \partial v_i)_{1 \leq i \leq c}$. Let \bar{d}' be the degree of the collection \bar{R} . As follows from Claim 3.7 in the case when nc -rank of \bar{P} is greater than $\tilde{r}(r_{\bar{d}}, \bar{d})$ we can find $\bar{v} \in V^c$ such the nc -rank of $\bar{R}(\bar{v})$ is greater than $r_{\bar{d}}$. We fix such \bar{v} and write $\bar{R} := \bar{R}(\bar{v})$. We consider \bar{R} as a morphism $\bar{R} : \mathbf{V} \rightarrow \mathbf{A}^c \times \mathbf{A}^c$.

As follows from the part (1) we have $\dim(\mathbf{R}_{\bar{t}, \bar{s}}) = \dim(V) - 2c$ for all $\bar{t}, \bar{s} \in \mathbf{A}^c$.

Let $\mathbf{Z} \subset \mathbf{A}^c$ be the subvariety of $\bar{s} = (s_1, \dots, s_c)$ such that $s_1 s_2 \dots s_c = 0$ and $\mathbf{Y} = \bar{R}^{-1}(\bar{t} \times \mathbf{Z}) \subset \mathbf{F}_{\bar{t}}$. Since $\dim(\mathbf{R}_{\bar{t}, \bar{s}}) = \dim(V) - 2c$ for all $\bar{t}, \bar{s} \in \mathbf{A}^c$ we see that $\dim(\mathbf{Y}) = \dim(V) - c - 1 < \dim(\mathbf{F}_{\bar{t}})$. Since the scheme $\mathbf{F}_{\bar{t}}$ is reduced at any $x \in \mathbf{F}_{\bar{t}} \setminus \mathbf{Y}$ we see that the scheme $\mathbf{F}_{\bar{t}}$ is reduced.

The proof of (3) is completely analogous. We choose $v_i, w_i \in V, 1 \leq i \leq c$ such that the rank of the collection $\bar{Q} = (\bar{P}, \partial P_i / \partial v_i, \partial P_i / \partial w_i)_{1 \leq i \leq c}$ is greater than $\tilde{r}(r_{\bar{d}}, \bar{d})$ where \bar{d}' be the degree of the collection \bar{Q} . We consider \bar{Q} as a morphism $\bar{Q} : \mathbf{V} \rightarrow \mathbf{A}^c \times \mathbf{A}^c \times \mathbf{A}^c$. As follows from (1) the fibers of the morphism \bar{Q} are of dimension $\dim(V) - 3c$.

Let $\mathbf{Y}' := \bar{Q}^{-1}(\bar{t} \times \mathbf{Z} \times \mathbf{Z}) \subset \mathbf{F}_{\bar{t}}$. Since $\dim(\mathbf{R}_{\bar{t}, \bar{s}}) = \dim(V) - 2c$ for all $\bar{t}, \bar{s} \in \mathbf{A}^c$ we see that $\dim(\mathbf{Y}') = \dim(V) - c - 2 = \dim(\mathbf{F}_{\bar{t}}) - 2$. Since the scheme $\mathbf{F}_{\bar{t}}$ is smooth at any $x \in \mathbf{F}_{\bar{t}} \setminus \mathbf{Y}'$ we see that the scheme $\mathbf{F}_{\bar{t}}$ is normal.

For the proof of part (4) we fix a generic subspace $\mathbf{W} \subset \mathbf{V}$ of dimension c . Since $\dim(\mathbf{F}_{\bar{t}}(\bar{P})) = n - c$ it follows from the Bézout theorem that $|\mathbf{F}_{\bar{t}}(\bar{P}) \cap \mathbf{W}(\mathbb{C})| = \prod_i d_i$ is odd. Therefore there exists a point $x \in \mathbf{F}_{\bar{t}}(\bar{P}) \cap \mathbf{W}(\mathbb{C})$ which is invariant under complex conjugation. But then $x \in \mathbf{F}_{\bar{t}}(\bar{P}) \cap \mathbf{W}(\mathbb{R})$. \square

Corollary 3.8. *For any infinite field K , K -vector spaces $V_i, 1 \leq i \leq d$ and a collection $\bar{P} = (P_1, \dots, P_c)$ of polylinear functions $P_i : V_1 \times \dots \times V_d \rightarrow K$ of degrees $\bar{d} = (d_1, \dots, d_c)$ and $\text{rank} \geq r_{\bar{d}}$, we have $\mathbf{F}_{\bar{t}}(\bar{P})(K) \neq \emptyset$ for $\bar{t} \in K^c$.*

Proof. We fix $\bar{t} \in K^c$ and for $w \in \mathbf{W} := \mathbf{V}_2 \times \dots \times \mathbf{V}_d$ define $\mathbf{Z}(w) = \{v_1 \in V_1 \mid (v_1, w) \in \mathbf{F}_{\bar{t}}(\bar{P})\}$. Let $\mathbf{Y} \subset \mathbf{W}$ be the subvariety of $w \in \mathbf{W}$ such that $\dim(\mathbf{Z}(w)) < \dim(V_1) - c$. As follows from Theorem 3.1 the closure $\bar{\mathbf{Y}} \subset \mathbf{W}$ of \mathbf{Y} is a proper closed subset. Since K is infinite we have $(\mathbf{W} \setminus \bar{\mathbf{Y}})(K) \neq \emptyset$. For any $w \in \mathbf{W} \setminus \bar{\mathbf{Y}}$ the subset $\mathbf{Z}(w)$ is defined by a system of linear equations in V_1 . Since $\mathbf{Z}(w) \neq \emptyset$ we see that $\mathbf{Z}(w)(K) \neq \emptyset$. \square

4. UNIVERSALITY OF HIGH RANK POLYNOMIAL MAPPINGS

Definition 4.1. A collection $\bar{P} = (P_i)_{1 \leq i \leq c}$ of degree $\bar{d} = (d_1, \dots, d_c)$ of polynomials on a K -vector space V is *m-universal* if for any collection $\bar{Q} = (Q_i)_{1 \leq i \leq c}$ of polynomials $Q_i \in K[x_1, \dots, x_m]$ of degrees $\leq d_i$ there exists an affine map $\phi : K^m \rightarrow V$ such that $Q_i = \phi^*(P_i)$ for all $1 \leq i \leq c$.

Theorem 4.2 (Universal). *There exists a function $R(\bar{d}, m)$ such that any collection of polynomials $\bar{P} = (P_1, \dots, P_c)$ of degrees $\bar{d} = (d_1, \dots, d_c)$ and $\text{nc-rank} \geq R(\bar{d}, m)$ is m -universal, where K is a field which is either finite or algebraically closed.*

Proof. To simplify notations we only outline the proof in the case when $c = 1$. Let Φ be the vector space of affine maps $\phi : K^m \rightarrow V$ and L be the vector space of polynomials $Q \in K[x_1, \dots, x_m]$ of degree $\leq d$. Choose a basis $\lambda_j, j \in J$ of the dual space to L . For any polynomial P of degree d on V we define the map $\bar{R}_P : \Phi \rightarrow K^J$ by $\bar{R}_P(\phi) = (\lambda_j(\phi^*(P)))_{j \in J}$. We have to show the surjectivity of the map \bar{R}_P . This surjectivity follows immediately from the following result (see Claim 3.11 in [10]), and Fact 2.9.

Fact 4.3. *For any r there exists $h(r, d, m)$ such that the nc-rank of \bar{R}_P is $\geq r$ for any polynomial P on V of $\text{nc-rank} \geq h(r, d, m)$.*

□

Corollary 4.4. *If $K = \mathbb{R}$ and all the degrees $d_i, 1 \leq i \leq c$ in a collection \bar{P} are odd then under the conditions of Theorem 4.2 the collection \bar{P} is m -universal.*

Proof. In this case the polynomials in $\bar{R}_{\bar{P}}$ are of the same degrees and Corollary 4.4 follows from the part (2) of Theorem 3.1. □

5. THE UNIVERSALITY FOR NUMBER FIELDS

Let K be a number field.

Theorem 5.1. *For any $m, \bar{d} = (d_1, \dots, d_c)$ there exists $r(\bar{d}, m)$ such that any collection $\bar{P} : V \rightarrow K^c$ of degree \bar{d} and $\text{rank} \geq r(\bar{d}, m)$ is m -universal if either K is totally imaginary or all degrees d_i are odd or the collection \bar{P} consists of polylinear functions.*

Proof. Let Φ be the space of affine maps from \mathbf{A}^m to V . The condition $\phi^*(\bar{P}) = \bar{Q}$ is equivalent to a system $\bar{G} = \bar{G}(\bar{P}, \bar{Q})$ of $M = \sum_{i=1}^c \binom{m+d_i}{d_i}$ equations of degrees $\leq d = \max_{1 \leq i \leq c} d_i$ on $\phi \in \Phi$. Let $\mathbf{X} \subset \Phi$ be the subvariety defined by the system \bar{G} and $\mathbf{X}_{\text{sing}} \subset \mathbf{X}$ be the singular locus.

Let $D = d^M$, $E = (D - 1)D^{(2^D)} + 1$. As follows from Claim 3.10 in [10] there exist \tilde{r} such the inequality $r_{\mathbb{C}}(P) \geq \tilde{r}$ implies $r_{\mathbb{C}}(\bar{G}) \geq D!(E + M)$. By Proposition III_C of [20] we know that $\text{codim}_{\mathbf{X}}(\mathbf{X}_{\text{sing}}) > E + M$. Since the field K is infinite there exists a K -subspace $\mathbf{W} \subset \Phi$ of codimension $E + M + 1$ such that the intersection $\mathbf{Z} = \mathbf{X} \cap \mathbf{W}$ is non-singular and geometrically irreducible. Since $\dim(\mathbf{W}) \geq M + E$ and $\text{codim}_{\Phi}(\mathbf{X}) \leq M$ we have $\dim(\mathbf{Z}) \geq E$. Since $\deg(\mathbf{Z}) = \deg(\mathbf{X}) \leq d^M$ we see that \mathbf{Z} satisfies the conditions of Theorem 4.2 in [22]. Now Theorem 5.1 follows from Theorem 1.1 in [5], part (2) of Theorem 3.1 and Corollary 3.8. □

6. WEAKLY POLYNOMIAL FUNCTIONS

We start the next topic with a couple of definitions.

- Definition 6.1.** (1) Let K be a field, V be a K -vector space and X a subset of V . A function $f : X \rightarrow K$ is *weakly polynomial* of degree $\leq a$, if for any affine subspace $L \subset X$ the restriction of f on L is a polynomial of degree $\leq a$.
- (2) A subset $X \subset V$ satisfies the condition \star_a if any weakly polynomial function of degree $\leq a$ on X is a restriction of a polynomial function of degree $\leq a$ on V .

The following example demonstrates the existence of cubic surfaces $X \subset K^2$ which do not have the property \star_1 for any field $K \neq \mathbb{F}_2$.

Example 6.2. Let $V = K^2$, $Q = xy(x - y)$, and let X be the variety defined by Q . Then $X = X_0 \cup X_1 \cup X_2$ where $X_0 = \{v \in V | x = 0\}$, $X_1 = \{v \in V | y = 0\}$, $X_2 = \{v \in V | x = y\}$. The function $f : X \rightarrow K$ such that $f(x, 0) = f(0, y) = 0$, $f(x, x) = x$ is weakly linear but one can not extend it to a linear function on V .

Definition 6.3. A field K is *e-admissible* for $e \geq 1$ if K^\star contains the subgroup μ_e of roots of order e .

Remark 6.4. To simplify notations we define $C := \mu_e$.

Theorem 6.5 (Extension). *There exists an $S = S(a, d)$ such that any hypersurface $Y \subset V$ of degree d and $nc\text{-rank} \geq S$ satisfies the condition \star_a if K is an ad -admissible field which is either finite field or is algebraically closed or is a number field.*

- Remark 6.6.** (1) The main difficulty in a proof of Theorem 6.5 is the non-uniqueness of an extension of f to a polynomial on V in the case when $a > d$.
- (2) An analogous statement is true for weakly polynomial functions on subsets $X_{\bar{P}}$ where \bar{P} is a collection of a sufficiently high nc -rank.

Proof. We fix the degree d of P . The proof consists of two steps. We first construct a collection $\mathbf{X}_m \subset \mathbf{V}_m$ of hypersurfaces of degree d and $nc\text{-rank} \geq m$ defined over \mathbb{Z} such that for all ad -admissible fields the subsets $X_m := \mathbf{X}_m(K) \subset V_m$ satisfy the condition \star_a . In the second step we derive the general case of Theorem 6.5 from this special case.

6.1. The first step. We denote by $x^i, 1 \leq i \leq d$ the coordinate functions on \mathbb{A}^d .

Definition 6.7.

- (1) $\mathbf{W} := \mathbf{A}^d$ and $\mu : \mathbf{W} \rightarrow \mathbf{A}$ is the product $\mu(x^1, \dots, x^d) := \prod_{j=1}^d x^j$.
- (2) $\mathbf{V}_m := \mathbf{W}^m$ and $Q_m(w_1, \dots, w_m) := \sum_{i=1}^m \mu(w_i)$,

$$(3) \mathbf{X}_m = \mathbf{X}_{Q_m}.$$

Proposition 6.8.

- (1) $r_{nc}(Q_m) \geq m$.
- (2) For any *ad*-admissible field K the subvariety $\mathbf{X}_m(K) \subset \mathbf{V}_m(K)$ has the property \star_a .

Proof. The proof of the inequality $r_{nc}(Q_m) \geq m$ is easy. In the case when $K = \mathbb{C}$ the result follows immediately from Lemma 16.1 of [20].

To outline a proof of the second statement we introduce a number of definitions. We fix m and write X instead of X_m . Since our field K is *ad*-admissible the group K^\star contains a finite subgroup C isomorphic to \mathbb{Z}_{ad} .

Definition 6.9.

- (1) $W_1 := \{(x^1, \dots, x^d) \in W \mid x^i = 1, i \geq 2\}$.
- (2) $L := W_1^m \cap X$.
- (3) $H \subset C^d \subset (K^\star)^d$ is the kernel of the product map $\mu_C : C^d \rightarrow C$. The group H acts on W by $(c^1, \dots, c^d)(x^1, \dots, x^d) = (c^1 x^1, \dots, c^d x^d)$.
- (4) Θ is the group of characters $\theta : H^m \rightarrow K^\star$.
- (5) We write elements of V_m in the form $v = (w_1, \dots, w_m)$, $1 \leq i \leq m$, $w_i \in W$. The group H^m acts on $X_m \subset V_m$ by $(h_1, \dots, h_m)(w_1, \dots, w_m) = (h_1 w_1, \dots, h_m w_m)$.
- (6) $\mathcal{P}_a^w(X) \subset k[X]$ is the subspace of weakly polynomial functions of degree $\leq a$ on X .
- (7) $\mathcal{P}_a(X) \subset \mathcal{P}_a^w(X)$ is the subspace of functions $f : X \rightarrow k$ which are restrictions of polynomial functions on V of degree $\leq a$.
- (8) For $\theta \in \Theta$, denote by $\mathcal{P}_a^w(X)^\theta \subset \mathcal{P}_a^w(X)$, $\mathcal{P}_a(X)^\theta \subset \mathcal{P}_a(X)$, the subspaces θ -eigenfunctions.

Since $C \subset K^\star$ we have direct sum decompositions $\mathcal{P}_a^w(X) = \oplus_{\theta \in \Theta} \mathcal{P}_a^w(X)^\theta$ and $\mathcal{P}_a(X) = \oplus_{\theta \in \Theta} \mathcal{P}_a(X)^\theta$. Therefore for a proof of Proposition 6.8 it is sufficient to show the equality $\mathcal{P}_a^w(X)^\theta = \mathcal{P}_a(X)^\theta$ for all $\theta \in \Theta$.

Fix $f \in \mathcal{P}_a^w(X)^\theta$. Since $L \subset V$ is a linear subspace the restriction $f|_L$ extends to a polynomial on V . So (after the subtraction of a polynomial) we may assume that $f|_L \equiv 0$. As shown in [10] Section 4.2, any weakly admissible function $f \in \mathcal{P}_a^w(X)^\theta$ vanishing on L is identically 0. So $f \in \mathcal{P}_a(X)^\theta$. \square

6.2. The second step. The proof of the general case of Theorem 6.5 is based on the following result.

Proposition 6.10. *There exists a function $r(d, a)$ such that the following holds. Let K be a field which is either finite or algebraically closed, V a K -vector space, P a polynomial of degree d and $W \subset V$ an affine subspace such that the *nc*-rank of the restriction of P on W is $\geq r(d, a)$. Then any weakly polynomial function f on X of degree $\leq a$ such that $f|_{X \cap W}$ extends to a polynomial on W of degree $\leq a$ is a restriction of a polynomial of degree $\leq a$ on V .*

Proof. After subtracting a polynomial from f we may assume that $f|_{X \cap W} \equiv 0$. Using induction on the codimension of W we reduce the Proposition to the case when $W \subset V$ is a hyperplane. We fix a direct sum decomposition $V = W \oplus K$ and denote by $t : V \rightarrow K$ the projection. Our proof is by induction on a .

The function $g' := f/t$ is defined on $X \setminus X \cap W$. We start with a construction of an extension of g' to a function g on X . Given a point $y \in X \cap W$ consider the set \mathcal{L} of lines $L \subset X$ such that $L \cap W = \{y\}$. Since f is weakly polynomial, the restriction f_L is a polynomial $p_L(t)$ vanishing at 0. We define $g_L(y)$ as the value of $p_L(t)/t$ at 0. It is clear that the following two results (see Proposition 4.33 of [10]) imply the validity of Proposition 6.10.

Claim 6.11. Under the conditions of Proposition 6.10, there exist $g(y) \in K$ such that

- (1) $|\mathcal{L}^-|/|\mathcal{L}| < 1/q$ if $K = \mathbb{F}_q$ and
- (2) $\dim(\mathcal{L}^-) < \dim(\mathcal{L})$ if K is algebraically closed

where $\mathcal{L}^- = \{L \in \mathcal{L} \mid g_L(y) \neq g(y)\}$.

We extend g' to a function on X whose values on $y \in X \cap W$ are equal to $g(y)$.

Claim 6.12. The function $g : X \rightarrow K$ is weakly polynomial of degree $a - 1$.

□

Now we can finish the proof of Theorem 6.5. Fix m such that $r_{nc}(Q_m) \geq r(d, a)$. Let $R(d, m)$ be as Theorem 4.2 or in Theorem 5.1 if K is a number field. We claim that for any admissible field K which is either finite or algebraically closed, any hypersurface $Y \subset V$ of degree d and nc -rank $\geq R(d, m)$ satisfies \star_a . Indeed, let f be a weakly polynomial function on $X = X_P$ of degree $\leq a$ where $P : V \rightarrow K$ is a polynomial of degree d of nc -rank $\geq R(d, m)$. Since $r_{nc}(P) \geq R(d, m)$ there exists an affine map $\phi : K^m \rightarrow V$ such that $P \circ \phi = Q_m$. It is clear that the function $f \circ \phi$ is a weakly polynomial function on X_m of degree $\leq a$. Therefore it follows from Proposition 6.8 that the restriction of f on $\text{Im}(\phi) \cap X$ extends to a polynomial on $\text{Im}(\phi)$. It follows now from Proposition 6.10 that f extends to a polynomial on V .

□

7. NULLSTELLENSATZ

Let K be a field and V be a finite dimensional K -vector space. We denote by \mathbf{V} the corresponding K -scheme, and by $\mathcal{P}(V)$ the algebra of polynomial functions on \mathbf{V} defined over K .

For a finite collection $\bar{P} = (P_1, \dots, P_c)$ of polynomials on \mathbf{V} we denote by $J(\bar{P})$ the ideal in $\mathcal{P}(V)$ generated by these polynomials, and by $\mathbf{X}_{\bar{P}}$ the subscheme of \mathbf{V} defined by this ideal.

Given a polynomial $R \in \mathcal{P}(\mathbf{V})$, we would like to find out whether it belongs to the ideal $J(\bar{P})$. It is clear that the following condition is necessary for the inclusion $R \in J(\bar{P})$.

(N) $R(x) = 0$ for all K -points $x \in \mathbf{X}_{\bar{P}}(K)$.

Claim 7.1 (Nullstellensatz). Suppose that the field K is algebraically closed and that the ideal $J(\bar{P})$ is radical. Then any polynomial R satisfying the condition (N) lies in $J(\bar{P})$.

We will show that the analogous result hold for $k = \mathbb{F}_q$ if $\mathbf{X}_{\mathbf{P}}$ is of high nc -rank.

From now on we fix a degree vector $\bar{d} = (d_1, \dots, d_c)$ and write $D := \prod_{i=1}^c d_i$. We denote by $\mathcal{P}_{\bar{d}}(\mathbf{V})$ the space of \bar{d} -families of polynomials $\mathbf{P} = (P_i)_{i=1}^c$ on V such that $\deg(P_i) \leq d_i$.

Theorem 7.2. *There exists an bound $r(\bar{d}) > 0$ such that for any finite field $k = \mathbb{F}_q$, $a < q/D$ and any collection \bar{P} of degrees \bar{d} and nc -rank $\geq r(\bar{d})$ the following holds. Any polynomial Q on V of degree a vanishing on $\mathbf{X}_{\bar{P}}(k)$ belongs to the ideal $J(\mathbf{P})$.*

Proof. Our proof is based on the following *rough bound* (see [7]).

Lemma 7.3. *Let $\bar{P} = (P_i)_{i=1}^c \subset \mathbb{F}_q[x_1, \dots, x_n]$ be a collection of polynomials of degrees d_i , $1 \leq i \leq c$ such that the variety $\mathbf{Y} := \mathbf{X}_{\bar{P}} \subset \mathbf{A}^n$ is of dimension $n - c$. Then $|\mathbf{Y}(\mathbb{F}_q)| \leq q^{n-c} D$ where $D := \prod_{i=1}^c d_i$.*

For the convenience we reproduce the proof of this result.

Proof. Let F be the algebraic closure of \mathbb{F}_q . Then $\mathbf{Y}(\mathbb{F}_q)$ is the intersection of $\mathbf{Y}(F)$ with hypersurfaces \mathbf{Y}_j , $1 \leq j \leq n$ defined by the equations $h_j(x_1, \dots, x_n) = 0$ where $h_j(x_1, \dots, x_n) = x_j^q - x_j$.

Let H_1, \dots, H_{n-c} be generic linear combinations of the h_j with algebraically independent coefficients from an transcendental extension F' of F and $\mathbf{Z}_1, \dots, \mathbf{Z}_{n-c} \subset \mathbf{A}^n$ be the corresponding hypersurfaces.

Intersect successively \mathbf{Y} with $\mathbf{Z}_1, \mathbf{Z}_2, \dots, \mathbf{Z}_{n-c}$. Inductively we see that for each $j \leq n - c$, each component C of the intersection $\mathbf{Y} \cap \mathbf{Z}_1 \cap \dots \cap \mathbf{Z}_j$ has dimension $n - c - j$. Indeed, passing from j to $j + 1$ for $j < n - c$ we have $\dim(C) = n - c - j > 0$. So not all the functions h_i vanish on C . Hence by the genericity of the choice of linear combinations $\{H_j\}$ we see that H_{j+1} does not vanish on C and therefore $\mathbf{Z}_{j+1} \cap C$ is of pure dimension $n - c - j - 1$. Thus the intersection $\mathbf{Y} \cap \mathbf{Z}_1 \cap \dots \cap \mathbf{Z}_{n-c}$ has dimension 0. This concludes the induction step.

By Bézout's theorem we see that $|\mathbf{Y} \cap \mathbf{Z}_1 \cap \dots \cap \mathbf{Z}_{n-c}| \leq q^{n-c} D$. Since $\mathbf{Y}(\mathbb{F}_q) = \mathbf{Y}(\mathbb{F}_q) \cap \mathbf{Z}_1(\mathbb{F}_q) \cap \dots \cap \mathbf{Z}_n(\mathbb{F}_q) \subset \mathbf{X}(\mathbb{F}_q) \cap \mathbf{Y}_1(\mathbb{F}_q) \cap \dots \cap \mathbf{Y}_{n-c}(\mathbb{F}_q)$ we see that $|\mathbf{Y}(\mathbb{F}_q)| \leq q^{n-c} D$. \square

Now we can finish the proof of Theorem 7.2. Let $R \in \mathbb{F}_q[x_1, \dots, x_n]$ be a polynomial of degree a vanishing on the set $\mathbf{X}_R(\mathbb{F}_q)$. Suppose that R does not lie in the ideal generated by the P_i , $1 \leq i \leq c$. Let $\mathbf{Y} := \mathbf{X}_{\bar{P}}$. Since R vanishes on $\mathbf{X}_{\bar{P}}(\mathbb{F}_q)$ we have $\mathbf{Y}(\mathbb{F}_q) = \mathbf{X}_{\bar{P}}(\mathbb{F}_q)$.

Since $\mathbf{X}_{\bar{P}}$ is irreducible we have $\dim(\mathbf{Y}) = n - c - 1$. As follows from Lemma 7.3 we have the upper bound $|\mathbf{Y}(\mathbb{F}_q)| \leq aDq^{n-c-1}$. On the other hand as follows from Theorem 2.12 there exists an bound $r(\bar{d}) > 0$ such that the condition $r_{nc}(\bar{P}) \geq r(\bar{d})$ implies the inequality $|\mathbf{X}_{\bar{P}}(\mathbb{F}_q)| > q^{n-c} \frac{q-1}{q}$, which is a contradiction to $q > aD$. \square

8. p -ADIC BIAS-RANK

For the last two applications we formulate and prove an analogue of Claim 2.9 for local non-archimedian fields of characteristic zero.

Let K/\mathbb{Q}_p be a finite extension of degree u , $\mathcal{O} \subset K$ the ring of integers, $\mathfrak{p} \subset \mathcal{O}$ the maximal ideal and $q = |\mathcal{O}/\mathfrak{p}|$. For $l \geq 1$ we write $A_l = \mathcal{O}/\mathfrak{p}^l$.

Definition 8.1. (1) For a map $P : A_l^n \rightarrow A_l$ and $a \in A_l$ we write $\nu(a) := q^{(1-n)l} |P^{-1}(a)|$.
 (2) A map P is s -uniform if $|\nu(a) - 1| \leq q^{-s}$.
 (3) We denote by Ξ_l the group of additive characters $\chi : A_l \rightarrow \mathbb{C}^\times$.
 (4) For $\chi \in \Xi_l$ we denote by $d(\chi)$ the smallest number d such that $\chi|_{\mathfrak{p}^d A_l} \equiv 1$.
 (5) $\Xi_l^d \subset \Xi_l$ is the subset of characters χ such that $d(\chi) = d$.
 (6) For $\chi \in \Xi$ we define $b(P; \chi) := q^{-nl} \sum_{v \in A_l^n} \chi(P(v)) = q^{-nl} \sum_{a \in A_l} \nu(a) \chi(a)$.
 (7) For $P : A_l^n \rightarrow A_l$ of degree d , we denote $\tilde{P} : (A_l^n)^d \rightarrow A_l$ the multilinear form $\tilde{P}(h_1, \dots, h_d) = \Delta_{h_d} \dots \Delta_{h_1} P$, where $\Delta_h P(x) = P(x+h) - P(x)$.
 (8) We write \tilde{P} for the reduction of $\tilde{P} \bmod \mathfrak{p}$.

Claim 8.2. If $|b(P; \chi)| < q^{-sd(\chi)}$ for all $\chi \in \Xi$ then P is $(s-2)$ -uniform.

Proof. The proof is completely analogous to the proof of Lemma 2.7. \square

Theorem 8.3. *There exists a function $r(d, s, u)$ such that the following holds: for any polynomial $P : A_l^n \rightarrow A_l$ of degree d such that the reduction $\hat{P} : (\mathbb{F}_q^n)^d \rightarrow \mathbb{F}_q$ of \tilde{P} is of rank $\geq r(d, s, u)$, we have $|b(P; \chi)| < q^{-sd(\chi)}$ for all $\chi \in \Xi_d$.*

To simplify notations we assume that $K = \mathbb{Q}_p$. The proof in the general case is completely analogous. We denote $V_l = A_l^n$, and $e(x) = e^{2\pi i x}$. We introduce the Gowers uniformity norms: if G is a finite abelian group $f : G \rightarrow \mathbb{C}$ we define

$$\|f\|_{U_m}^{2^m} = \mathbb{E}_{x, h_1, \dots, h_m \in G} \partial_{h_m} \dots \partial_{h_1} f(x)$$

where $\partial_h f(x) = f(x+h) - \overline{f(x)}$. Note that $\partial_h \chi(P(x)) = \chi(\Delta_h P(x))$

Our proof of Theorem 8.3, which is by induction in d, l is based on the following stronger result.

Proposition 8.4 (d,l). *For any $s > 0$, there exists $r = r^B(d, s)$ such that for any polynomial S of degree $< d$, any m , if \hat{R} is of rank $> r$ then*

$$|\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l + S(x)/\mathfrak{p}^m)| < q^{-sl}.$$

Remark 8.5. The constant $r = r^B(d, s)$ does not depend on l .

Proof. We start with the following result.

Lemma 8.6. *Let $r = \alpha_d(2^{2d-1}ds)$ be as in Fact 2.9. Then Proposition 8.4(d,l) holds for $l \leq d$.*

Proof. It suffices to show that for any $1 \leq l \leq d$, for any polynomial $R : V_l \rightarrow A_l$ of degree d such that the rank of \hat{R} is $> r$, a polynomial $S : V_l^d$ of degree $< d$, and any m we have

$$|\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l + S(x)/\mathfrak{p}^m)| < q^{-ds}.$$

Suppose that

$$|\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l + S(x)/\mathfrak{p}^m)| \geq q^{-ds}.$$

for some $1 < l \leq d, m \geq 0$. We observe that by the CS inequality we have

$$\begin{aligned} |\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l + S(x)/\mathfrak{p}^m)|^2 &\leq \mathbb{E}_{y \in V_{l-1}^d} |\mathbb{E}_{x_1 \in V_{l-1}} e(\tilde{R}(x_1, y)/\mathfrak{p}^l + S(x_1, y)/\mathfrak{p}^m)|^2 \\ &= \mathbb{E}_{y \in V_{l-1}^d} \mathbb{E}_{x_1, t \in V_{l-1}} e((\tilde{R}(x_1 + t, y) - \tilde{R}(x_1, y))/\mathfrak{p}^l + (S(x_1 + t, y) - S(x_1, y))/\mathfrak{p}^m). \end{aligned}$$

whiich is equal to

$$\mathbb{E}_{y \in V_{l-1}^d} \mathbb{E}_{x_1, t \in V_{l-1}} e(\tilde{R}(t, y)/\mathfrak{p}^l + (S(x_1 + t, y) - S(x_1, y))/\mathfrak{p}^m)$$

Applying CS inequality $d - 1$ more times in the coordinates x_2, \dots, x_d we obtain (since S is of degree $< d$):

$$(\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l))^{1/2^d} \geq q^{-ds}.$$

In other words

$$\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l) \geq q^{-2^d ds}.$$

We rewrite the LHS as

$$\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l) = \mathbb{E}_x \mathbb{E}_{y \in V_{l-1}^d} e(\tilde{R}(x + \mathfrak{p}y)/\mathfrak{p}^l).$$

We observe that by the CS inequality we have

$$\begin{aligned} |\mathbb{E}_x \mathbb{E}_{y \in V_{l-1}^d} e(\tilde{R}(x + \mathfrak{p}y)/\mathfrak{p}^l)|^2 &\leq \mathbb{E}_x \mathbb{E}_{y' \in V_{l-1}^{d-1}} |\mathbb{E}_{y_1 \in V_{l-1}} e(\tilde{R}(x_1 + \mathfrak{p}y_1, x' + \mathfrak{p}y')/\mathfrak{p}^l)|^2 \\ &= \mathbb{E}_x \mathbb{E}_{y' \in V_{l-1}^{d-1}} \mathbb{E}_{y_1, h_1 \in V_{l-1}} e((\tilde{R}(x_1 + \mathfrak{p}(y_1 + h_1), x' + \mathfrak{p}y') - \tilde{R}(x_1 + \mathfrak{p}y_1, x' + \mathfrak{p}y'))/\mathfrak{p}^l) \end{aligned}$$

Which is equal to

$$\mathbb{E}_x \mathbb{E}_{y' \in V_{l-1}^{d-1}} \mathbb{E}_{y_1, h_1 \in V_{l-1}} e((\tilde{R}(\mathfrak{p}h_1, x' + \mathfrak{p}y')/\mathfrak{p}^l) = \mathbb{E}_x \mathbb{E}_{y' \in V_{l-1}^{d-1}} \mathbb{E}_{y_1, h_1 \in V_{l-1}} e((\tilde{R}(h_1, x' + \mathfrak{p}y')/\mathfrak{p}^{l-1}).$$

Applying the CS inequality $l - 2$ more times we see that

$$\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l) \leq [\mathbb{E}_x \mathbb{E}_{h_i \in V_{l-1}} e(\tilde{R}(h_1, \dots, h_{l-1}, x_l, \dots, x_d)/\mathfrak{p})]^{1/2^{l-1}},$$

and therefore

$$\mathbb{E}_{x \in V_1^d} e(\hat{R}(x_1, \dots, x_d)/\mathfrak{p}) \geq q^{-2^{d+l-1}ds}.$$

Thus \hat{R} is of rank $\leq \alpha_d(2^{d+l-1}ds)$. \square

We prove Proposition 8.4(d,l) by induction on d, l . Let P be such that \hat{P} is of rank $> c(d, s)$ where

$$c(d, s) = \max \{ \alpha_d(2^{2d-1}ds), \alpha_d(2^d(r^B(d-1, 2s+1) + 2ds+1)) \}.$$

By Lemma 8.6, Proposition 8.4(d,l) holds when $l \leq d$. So we assume $l > d$.

Suppose that Proposition 8.4(d',l') holds if either $d' < d$ or $d' = d$ and $l' < l$. If Proposition 8.4(d,l) fails then there exists a degree d polynomial R such that $\tilde{R} : V_l^d \rightarrow A_l$ is of rank $> c(d, s)$ and some polynomial $S : V_l^d \rightarrow A_l$ of degree $< d$ and $m \in \mathbb{N}$ such that:

$$|\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l + S(x)/\mathfrak{p}^m)| \geq 1/q^{ls}.$$

From this it follows that

$$\begin{aligned} \frac{1}{q^{2ls}} &\leq |\mathbb{E}_{x \in V_l^d} e(\tilde{R}(x)/\mathfrak{p}^l + S(x)/\mathfrak{p}^m)|^2 \\ &= \mathbb{E}_{x, y \in V_l^d} e((\tilde{R}(x+y) - \tilde{R}(x))/\mathfrak{p}^l + (S(x+y) - S(x))/\mathfrak{p}^m) \\ &= \mathbb{E}_{t \in V_1^d} \mathbb{E}_{y \in V_l^d: y \equiv t(\mathfrak{p})} \mathbb{E}_{x \in V_l^d} e((\tilde{R}(x+y) - \tilde{R}(x))/\mathfrak{p}^l + (S(x+y) - S(x))/\mathfrak{p}^m) \end{aligned}$$

Fix t and consider the inner average:

$$a_t := \mathbb{E}_{y \in V_l^d: y \equiv t(\mathfrak{p})} \mathbb{E}_{x \in V_l^d} e((\tilde{R}(x+y) - \tilde{R}(x))/\mathfrak{p}^l + (S(x+y) - S(x))/\mathfrak{p}^m)$$

By shifting $x \rightarrow x + y'$ and averaging we see that

$$\begin{aligned} a_t &= \mathbb{E}_{x \in V_l^d} \mathbb{E}_{y, y' \in V_l^d: y \equiv t(\mathfrak{p}), y' \equiv 0(\mathfrak{p})} \\ &\quad e((\tilde{R}(x+y'+y) - \tilde{R}(x+y'))/\mathfrak{p}^l + (S(x+y'+y) - S(x+y'))/\mathfrak{p}^m) \\ &= \mathbb{E}_{x \in V_l^d} |\mathbb{E}_{y \equiv t(\mathfrak{p})} e((\tilde{R}(x+y))/\mathfrak{p}^l + S(x+y)/\mathfrak{p}^m)|^2 \end{aligned}$$

Lemma 8.7. $|a_t| \leq \frac{1}{q^{2(l-d)s}}$

Proof.

$$\begin{aligned} &|\mathbb{E}_{y \equiv t(\mathfrak{p})} e((\tilde{R}(x+y))/\mathfrak{p}^l + S(x+y)/\mathfrak{p}^m)| \\ &= |\mathbb{E}_{y \in V_l^d, y \equiv 0(\mathfrak{p})} e((\tilde{R}(x+t+y))/\mathfrak{p}^l + S(x+t+y)/\mathfrak{p}^m)| \\ &= |\mathbb{E}_{y \in V_{l-1}^d} e((\tilde{R}(x+t+\mathfrak{p}y))/\mathfrak{p}^l + S(x+t+\mathfrak{p}y)/\mathfrak{p}^m)| \\ &= |\mathbb{E}_{y \in V_{l-1}^d} e(\frac{\tilde{R}(y)}{\mathfrak{p}^{l-d}} + \frac{S(y)}{\mathfrak{p}^{m-d}} + \frac{S_{x,t}(\mathfrak{p}y)}{\mathfrak{p}^m} + \frac{T_{x,t}(\mathfrak{p}y)}{\mathfrak{p}^l})| \end{aligned}$$

(if $d > m$ the second term does not exist) where $T_{x,t}(\mathfrak{p}y)$ is of degree $\leq (d-1)$ in y . By the induction hypothesis the latter is $\leq \frac{1}{q^{(l-d)s}}$. \square

Claim 8.8. Let $b_t \geq 0$ be such that $\mathbb{E}_{t \in E} b_t \geq \frac{1}{q^{2ls}}$ and $b_t \leq \frac{1}{q^{2(l-d)s}}$ then for $\frac{1}{2q^{2ds}}|E|$ many t we have that $b_t \geq \frac{1}{2q^{2ls}}$.

Proof. Let F be the set where $b_t \geq \frac{1}{2q^{2ls}}$. We have that

$$\frac{1}{q^{2ls}}|E| \leq \sum_{t \in E} |a_t| \leq \sum_{t \in F} \frac{1}{q^{2(l-d)s}} + \sum_{t \in E-F} \frac{1}{2q^{2ls}}$$

Rearranging we get

$$|E| \leq |F|(q^{2ds} - 1/2) + |E|\frac{1}{2},$$

so that $|F| \geq \frac{|E|}{2(q^{2ds}-1/2)}$. \square

Applying this Claim to $b_t := |a_t|$ we see that for $\geq \frac{1}{2q^{2ds}}|V_1^d|$ many $t \in V_1^d$ we have that

$$|\mathbb{E}_{y \in V_l^d: y \equiv t(\mathfrak{p})} \mathbb{E}_{x \in V_l^d} e((\tilde{R}(x+y) - \tilde{R}(x))/\mathfrak{p}^l + (S(x+y) - S(x))/\mathfrak{p}^m)| \geq \frac{1}{2q^{2ls}} \geq \frac{1}{q^{l(2s+1)}}.$$

Now $Q_y = \Delta_y R$ is of degree $< d$ so by the induction on the degree \hat{Q}_y is of rank $< r^B(d-1, 2s+1)$.

This implies that for $\geq \frac{1}{2q^{2ds}}|V_1^d|$ many $t \in V_1^d$, if $y \equiv t(\mathfrak{p})$ then

$$\|\mathbb{E}_{x \in V_1^d} e((\hat{R}(x+y) - \hat{R}(x))/\mathfrak{p})\|_{U_{d-1}} \geq \frac{1}{q^{r^B(d-1, 2s+1)}}$$

But this now implies that

$$\mathbb{E}_{t \in V_1^d} \|\mathbb{E}_{y \in V_l^d: y \equiv t(\mathfrak{p})} \mathbb{E}_{x \in V_l^d} e((\hat{R}(x+y) - \hat{R}(x))/\mathfrak{p})\|_{U_{d-1}} \geq \frac{1}{q^{r^B(d-1, 2s+1)} q^{2ds+1}}$$

But LHS is bounded by

$$\|e(\hat{R}(x)/\mathfrak{p})\|_{U_d}^2 \leq \|e(\hat{R}(x)/\mathfrak{p})\|_{U_d} \leq |\mathbb{E}_{x \in V_1} e(\hat{R}(x)/\mathfrak{p})|^{1/2^d}$$

and we are given that \hat{R} is of rank $> \alpha_d(2^d(r^B(d-1, 2s+1) + 2ds+1))$, contradiction. \square

9. RATIONAL SINGULARITIES

Definition 9.1. Let \mathbf{X} be a normal irreducible variety over a field of characteristic zero and $a : \tilde{\mathbf{X}} \rightarrow \mathbf{X}$ a resolution of singularities. We say that \mathbf{X} has rational singularities if $R^i a_*(\mathcal{O}_{\tilde{\mathbf{X}}}) = \{0\}$ for $i > 0$.

Remark 9.2. This property of \mathbf{X} does not depend on a choice of a resolution $a : \tilde{\mathbf{X}} \rightarrow \mathbf{X}$.

Theorem 9.3. *There exists a function $r(\bar{d}), \bar{d} = (d_1, \dots, d_c)$ such that the following holds.*

Let $\bar{P} = (P_i)_{i=1}^c \subset \mathbb{C}[x_1, \dots, x_n]$ be a collection of (not necessarily homogeneous) polynomials of degrees $d_i, 1 \leq i \leq c$ of rank $\geq r(\bar{d})$. Then the variety $\mathbf{X}_{\bar{P}}$ has rational singularities.

Proof. To simplify the exposition we assume that $c = 1$. Thus $\bar{P} = (P)$ where $P \in \mathbb{C}[x_1, \dots, x_n]$ is a homogeneous polynomial of degree d .

We first consider the case when $P \in K[x_1, \dots, x_n]$ where K/\mathbb{Q} is a finite extension. In this case there exists an infinite set S of prime ideals in \mathcal{O}_K such that

- (1) for any $\pi \in S$ the completion $\mathcal{O}_{K\pi}$ of \mathcal{O}_K at π is isomorphic to \mathbb{Z}_p where $p = \text{char}(\mathcal{O}_K/\pi)$,
- (2) $P \in \mathcal{O}_{K\pi}[x_1, \dots, x_n]$ and
- (3) the reduction $\hat{P} \in \mathbb{F}_p[x_1, \dots, x_n]$ is of rank $r(P)$ over the algebraic closure of \mathbb{F}_p .

We fix $\pi \in S$ such that $p > \deg(P)$. As follows from Theorem A of [2] it is sufficient to establish the inequalities $|\mathbf{X}(\mathbb{Z}/p^m\mathbb{Z}) - p^{m(n-1)}| \leq p^{m(n-1)-1/2}, m \geq 1$.

We see that for number fields the validity of Theorem 9.3 in the case when $P \in K[x_1, \dots, x_n]$ follows immediately from Theorem 8.3.

We show now how to derive the general case of Theorem 9.3 from the case when $P \in K[x_1, \dots, x_n]$ where K/\mathbb{Q} is a finite extension.

Definition 9.4. Let $a : \mathbf{X} \rightarrow \mathbf{Y}$ be a morphism between complex algebraic varieties such that all the fibers X_y are normal and geometrically irreducible. We write $Y = \mathbf{Y}(\mathbb{C})$ and denote by $Y_a \subset Y$ the subset of points y such that the fiber X_y has rational singularities.

Lemma 9.5. *If a projective morphism a is defined over \mathbb{Q} then the subset $Y_a \subset Y$ is also defined over \mathbb{Q} .*

Proof. The proof is by induction in $d = \dim(Y)$. Suppose that the Lemma is known in the case when the dimension of the base is $< d$. Let t be the generic point of \mathbf{Y} and \mathbf{X}_t the fiber of \mathbf{X} over t . Fix a resolution $\tilde{b} : \tilde{\mathbf{X}}_t \rightarrow \mathbf{X}_t$ over the field $k(t)$ of rational functions on Y . Then there exists a non-empty open subset $U \subset \mathbf{Y}$ such that $b := \tilde{b}|_{(a \circ \tilde{b})^{-1}U}$ is a resolution of $X_U := a^{-1}(U)$. By definition $Y_a \cap U = \{u \in U \mid R^i b_*(\mathcal{O}_{\tilde{X}})_u = \{0\}\}$ for all $i > 0$. Since the sheaves $R^i b_*(\mathcal{O}_{\tilde{X}})$ are coherent we see that the subset $Y_a \cap U$ is defined over \mathbb{Q} . On the other hand, the inductive assumption implies that $Y_a \cap (Y \setminus U)$ is defined over \mathbb{Q} . \square

Now we can finish a proof of Theorem 9.3 in the general case. Consider the trivial fibration $\hat{a} : \mathbf{A}^n \times \mathbf{Y} \rightarrow \mathbf{Y}$ where $\mathbf{Y} \subset \mathbf{P}_d^n$ is the constructible subset of polynomials of degree d on \mathbf{A}^n and of rank $\geq r(d)$. Let $\mathbf{X} \subset \mathbf{A}^n \times \mathbf{Y}$ be the hypersurface such that $\hat{a}^{-1}(P) \cap \mathbf{X} = \mathbf{X}_P$ and $a : \mathbf{X} \rightarrow \mathbf{Y}$ be the restriction of \hat{a} onto \mathbf{X} . As follows from Theorem 3.1 and Proposition III_C of [20] all fibers

of a are irreducible and normal. For a proof of Theorem 9.3 we have to show that $Y_a = Y$. The validity of Theorem 9.3 in the case when $P \in K[x_1, \dots, x_n]$, where K/\mathbb{Q} is any finite extension, shows that any point of Y defined over a finite extension K of \mathbb{Q} belongs to Y_a . Since the subset Y_a of Y is defined over \mathbb{Q} we see that $Y_a = Y$. \square

REFERENCES

- [1] Ananyan, T., Hochster, M. *Small subalgebras of polynomial rings and Stillman's Conjecture*. J. Amer. Math. Soc. 33 (2020), 291-309.
- [2] Aizenbud, A., Avni, N. *Counting points of schemes over finite rings and counting representations of arithmetic lattices*. Duke Math. J. 167 (2018), no. 14, 2721-2743.
- [3] Bhowmick A., Lovett S. *Bias vs structure of polynomials in large fields, and applications in effective algebraic geometry and coding theory*. arXiv:1506.02047
- [4] Bik, A., Draisma, J., Eggermont, R. *Polynomials and tensors of bounded strength*. arXiv 1805.01816
- [5] Browning, T., Heath-Brown, D. *Forms in many variables and differing degrees*. J. Eur. Math. Soc. (JEMS) 19 (2017), no. 2, 357 - 394
- [6] Dersken, H. *The G -stable rank for tensors* arxiv.org/abs/2002.08435.
- [7] Hrushovski, E. *The Elementary Theory of the Frobenius Automorphisms* arXiv: 0406514
- [8] Janzer, O. *Polynomial bound for the partition rank vs the analytic rank of tensors* arXiv:1902.11207
- [9] Kazhdan, D., Ziegler, T. *Extending linear and quadratic functions from high rank varieties*. arXiv:1712.01335
- [10] Kazhdan, D., Ziegler, T. *Properties of high rank subvarieties of affine spaces*. To appear GAFA. arXiv:1902.00767.
- [11] Kazhdan, D., Ziegler, T. *Polynomials as splines* Sel. Math. New Ser. (2019) 25; 31.
- [12] Kazhdan, D., Ziegler, T. *On ranks of polynomials*. Algebr. Represent. Theory 21 (2018), no. 5, 1017–1021
- [13] Kazhdan, D., Ziegler, T. *Approximate cohomology*. Sel. Math. New Ser. (2018) 24: 499.
- [14] Kazhdan, D., Ziegler, T. *On the codimension of the singular locus*. ArXiv 1907.11750.
- [15] Lang, S., Weil A. *Number of points of varieties in finite fields*. Amer. J. Math. 76 (1954).
- [16] Lovett, S. *The Analytic rank of tensors and its applications*. Discrete Analysis 2019:7
- [17] Matsumura, H. *Commutative ring theory*. Cambridge Studies in Advanced Mathematics, 8. Cambridge University Press, Cambridge, 1989.
- [18] Marker, D. *Model Theory : An Introduction* Graduate Texts in Mathematics, Vol. 217, 2002
- [19] Milićević, L. *Polynomial bound for partition rank in terms of analytic rank* Geom. Funct. Anal. 29 (2019), no. 5, 1503-1530.
- [20] Schmidt, W. M. *The density of integer points on homogeneous varieties*. Acta Math. 154 (1985), no. 3-4, 243-296.
- [21] Tao, T.; Ziegler, T. *The inverse conjecture for the Gowers norm over finite fields in low characteristic*, Ann. Comb. 16 (1) (2012) 121-188.
- [22] Wooley, T. *Solvable points on smooth projective varieties*. Monatsh. Math. 180 (2016), no. 2, 391–403.