

Maximum and comparison principles for degenerate elliptic systems and some applications ^{*†}

Edir Junior Ferreira Leite [‡]

*Departamento de Matemática, Universidade Federal de Viçosa,
CCE, 36570-900, Viçosa, MG, Brazil*

Abstract

In this paper we develop a detailed study on maximum and comparison principles related to the following nonlinear eigenvalue problem

$$\begin{cases} -\Delta_p u = \lambda a(x)|v|^{\beta_1-1}v & \text{in } \Omega; \\ -\Delta_q v = \mu b(x)|u|^{\beta_2-1}u & \text{in } \Omega; \\ u = v = 0 & \text{on } \partial\Omega, \end{cases}$$

where $p, q \in (1, \infty)$, $\beta_1, \beta_2 > 0$ satisfy $\beta_1\beta_2 = (p-1)(q-1)$, $\Omega \subset \mathbb{R}^n$ is a bounded domain with C^2 -boundary, $a, b \in L^\infty(\Omega)$ are given functions, both assumed to be strictly positive on compact subsets of Ω , and Δ_p and Δ_q are quasilinear elliptic operators, stand for p -Laplacian and q -Laplacian, respectively. We classify all couples $(\lambda, \mu) \in \mathbb{R}^2$ such that both the (weak and strong) maximum and comparison principles corresponding to the above system hold in Ω . Explicit lower bounds for principal eigenvalues of this system in terms of the measure of Ω are also proved. As application, given $\lambda, \mu \geq 0$ we measure explicitly how small has to be $|\Omega|$ so that weak and strong maximum principles associated to the above problem hold in Ω .

1 Introduction and statements

In this paper we study maximum and comparison principles related to the following system:

$$\begin{cases} -\Delta_p u = \lambda a(x)|v|^{\beta_1-1}v & \text{in } \Omega; \\ -\Delta_q v = \mu b(x)|u|^{\beta_2-1}u & \text{in } \Omega; \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (1)$$

^{*}AMS Subject Classification 2010: 35B50; 35B51; 35J70; 35J92

[†]Key words: Maximum principles, Comparison principles, Lower bound of eigenvalues

[‡]E-mail addresses: edirjrleite@ufv.br (E.J.F. Leite)

where $\Omega \subset \mathbb{R}^n$ is a bounded domain with C^2 -boundary (not necessarily connected), $p, q \in (1, \infty)$, $a, b \in L^\infty(\Omega)$ are given functions satisfying

$$\operatorname{ess\,inf}_{x \in \Omega} a(x) > 0 \quad \text{and} \quad \operatorname{ess\,inf}_{x \in \Omega} b(x) > 0,$$

$\beta_1, \beta_2 > 0$ with $\beta_1 \beta_2 = (p-1)(q-1)$, and $(\lambda, \mu) \in \mathbb{R}^2$. The p -Laplacian is defined by

$$\Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u)$$

for any $u \in W_0^{1,p}(\Omega)$ with values $\Delta_p u \in W^{-1, \frac{p}{p-1}}(\Omega)$, the dual space of $W_0^{1,p}(\Omega)$.

Existence, nonexistence and uniqueness of nontrivial solutions to the system (1) have been widely investigated during the three last decades for $p = q = 2$ and, more generally, for $p, q \in (1, \infty)$. For $p = q = 2$, we refer for instance to [8], [15], [18], [24], [34] and [37], where in particular notions of sub-superlinearity, sub-supercriticality and criticality have been introduced. Still in the first part, the eigenvalue problem, i.e., $\beta_1 \beta_2 = 1$, was completely studied in [35]. For $p, q \in (1, \infty)$, we refer to [9] when $\beta_1 \beta_2 > (p-1)(q-1)$ and [11] when $\beta_1 \beta_2 = (p-1)(q-1)$.

The connection between principal eigenvalues and maximum principles have been investigated in [1, 2, 3, 7, 32] for cooperative systems and in [38] for non-cooperative systems (see also [30] for a more complete discussion) and more recently in [26], where system (1) was analyzed in the special case when $p = q = 2$, however, instead of Δ , a general second order elliptic operator was considered. For the nonlocal context, we refer to [27].

Here we extend the results of [26] for $p, q > 1$, that is, we establish the connection between principal spectral curves for systems (1) and maximum and comparison principles related. For this, we shall present a bit of notation. Note that, given any $f \in L^\infty(\Omega)$, there exists a unique weak solution $u \in W_0^{1,p}(\Omega)$ of the classical problem

$$\begin{cases} -\Delta_p u &= f(x) & \text{in } \Omega; \\ u &= 0 & \text{on } \partial\Omega. \end{cases} \quad (2)$$

Notice that, $u \in C^{1,\alpha}(\overline{\Omega})$ for some $\alpha \in (0, 1)$ (see [16, 22, 29, 40]). We denote $X := [C_0^1(\overline{\Omega})]^2$, $X_+ := \{(u, v) \in X : u \geq 0 \text{ and } v \geq 0 \text{ in } \Omega\}$, and $\overset{\circ}{X}_+$ is the topological interior of X_+ in X . Note that, $\overset{\circ}{X}_+$ is nonempty and characterized by $(u, v) \in \overset{\circ}{X}_+$ if, and only if, $(u, v) \in X$ satisfies:

$$u, v > 0 \text{ in } \Omega \quad \text{and} \quad \frac{\partial u}{\partial \nu}, \frac{\partial v}{\partial \nu} < 0 \text{ on } \partial\Omega,$$

where $\nu \equiv \nu(x_0)$ denotes the exterior unit normal to $\partial\Omega$ at $x_0 \in \partial\Omega$ (see [11]).

As is well known, the operator Δ_p satisfies the weak maximum principle, that is, for any weak solution $u \in W_0^{1,p}(\Omega)$ to

$$\begin{cases} -\Delta_p u &= f(x) & \text{in } \Omega; \\ u &\geq 0 & \text{on } \partial\Omega, \end{cases}$$

with $f \in L^\infty(\Omega)$ and $f \geq 0$ in Ω , one has $u \geq 0$ in Ω and Δ_p also satisfies the strong maximum principle, i.e., moreover $u > 0$ in Ω whenever $f \not\equiv 0$ in Ω (see [22], [39] and [41]).

Let (u, v) in $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$. The weak formulation of the system (1) is given by

$$\lambda \int_{\Omega} a(x) |v|^{\beta_1-1} v w dx = \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla w dx, \quad \forall w \in C_0^1(\Omega)$$

and

$$\mu \int_{\Omega} b(x) |u|^{\beta_2-1} u w dx = \int_{\Omega} |\nabla v|^{q-2} \nabla v \nabla w dx, \quad \forall w \in C_0^1(\Omega).$$

A couple $(\lambda, \mu) \in \mathbb{R}_+^* \times \mathbb{R}_+^* = (0, \infty)^2$ is said to be an eigenvalue of (1) if the system admits a nontrivial weak solution (φ, ψ) in $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ which is called an eigenfunction associated to (λ, μ) . We say that (λ, μ) is a principal eigenvalue if admits a positive eigenfunction (φ, ψ) ; i.e., φ and ψ are positive in Ω . We also say that (λ, μ) is simple in $\overset{\circ}{X}_+$ if for any eigenfunctions $(\varphi, \psi), (\tilde{\varphi}, \tilde{\psi}) \in \overset{\circ}{X}_+$, there exists $\rho > 0$ such that $\tilde{\varphi} = \rho\varphi$ and $\tilde{\psi} = \rho\mu^{\frac{1}{\beta_2}}\psi$ in Ω .

The existence of principal eigenvalues of (1) and some of their qualitative properties were treated in Cuesta and Takác [11]. Namely, they proved that the set formed by these principal eigenvalues is given by the following smooth curve

$$\mathcal{C}_1 := \left\{ (\lambda, \mu) \in (\mathbb{R}_+^*)^2 : \lambda^{\frac{1}{\sqrt{\beta_1(p-1)}}} \mu^{\frac{1}{\sqrt{\beta_2(q-1)}}} = \Lambda' \right\},$$

for some $\Lambda' > 0$, which satisfies:

- (i) $(\lambda, \mu) \in \mathbb{R}_+ \times \mathbb{R}_+$ is a principal eigenvalue of the system (1) if, and only if, $(\lambda, \mu) \in \mathcal{C}_1$;
- (ii) The curve \mathcal{C}_1 is simple in $\overset{\circ}{X}_+$, that is, (λ, μ) is simple in $\overset{\circ}{X}_+$ for all $(\lambda, \mu) \in \mathcal{C}_1$;
- (iii) Let $(\varphi, \psi) \in X$ be an eigenfunction corresponding to $(\lambda, \mu) \in \mathcal{C}_1$. Therefore either $(\varphi, \psi) \in \overset{\circ}{X}_+$ or $(-\varphi, -\psi) \in \overset{\circ}{X}_+$.

Recently, in [28] the author established another properties satisfied by principal curve \mathcal{C}_1 such as local isolation, monotonicity on Ω and monotonicity and continuity of the principal eigenvalue with respect to the weight functions a and b .

By weak maximum principle, denoted by **(WMP)**, we mean that for any weak solution $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ of the system

$$\begin{cases} -\Delta_p u = \lambda a(x) |v|^{\beta_1-1} v + f(x) & \text{in } \Omega; \\ -\Delta_q v = \mu b(x) |u|^{\beta_2-1} u + g(x) & \text{in } \Omega; \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (3)$$

with $f, g \in L^\infty(\Omega)$ and $f, g \geq 0$ in Ω , verifies $u, v \geq 0$ in Ω . Besides, if at least, u or v is positive in Ω whenever $f + g \not\equiv 0$ in Ω , we say that the strong maximum principle, denoted by **(SMP)**, corresponding to (1) holds in Ω . When $\lambda, \mu > 0$, **(SMP)** can be rephrased as $u, v > 0$ in Ω

whenever $f + g \not\equiv 0$ in Ω . In the case that $u > 0$ ($v > 0$) in Ω , we get $\frac{\partial}{\partial \nu} u(x_0) < 0$ ($\frac{\partial}{\partial \nu} v(x_0) < 0$) for all $x_0 \in \partial\Omega$.

We are ready to classify completely in terms of the principal curve \mathcal{C}_1 the set of couples $(\lambda, \mu) \in \mathbb{R}^2$ such that **(WMP)** and **(SMP)** hold in Ω .

Namely:

Theorem 1.1. *Let $(\lambda, \mu) \in \mathbb{R}^2$ and \mathcal{R}_1 be the open region in the first quadrant below \mathcal{C}_1 . The following assertions are equivalent:*

- (i) $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$;
- (ii) **(WMP)** corresponding to (1) holds in Ω ;
- (iii) **(SMP)** corresponding to (1) holds in Ω .

Note that the sets

$$\mathcal{R}_1 = \left\{ (\lambda, \mu) \in (\mathbb{R}_+^*)^2 : \lambda^{\frac{1}{\sqrt{\beta_1(p-1)}}} \mu^{\frac{1}{\sqrt{\beta_2(q-1)}}} < \Lambda' \right\}$$

and $\overline{\mathcal{R}_1} \setminus \mathcal{C}_1 = \mathcal{R}_1 \cup \{(\lambda, 0) : \lambda \geq 0\} \cup \{(0, \mu) : \mu \geq 0\}$ are unbounded and are depicted in Figure 1.




Figure 1: The principal curve \mathcal{C}_1 .

We notice that weak and strong comparison principles, denoted respectively by **(WCP)** and **(SCP)**, are very important tools for establish the uniqueness and positivity of solutions for elliptic problems, among others, to certain counterparts of (1). We say that **(WCP)** holds in

Ω if, for any weak solutions (u, v) and (z, w) in $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ to the following systems, respectively:

$$\begin{cases} -\Delta_p u = \lambda a(x)|v|^{\beta_1-1}v + f_1(x) & \text{in } \Omega; \\ -\Delta_q v = \mu b(x)|u|^{\beta_2-1}u + g_1(x) & \text{in } \Omega; \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (4)$$

$$\begin{cases} -\Delta_p z = \lambda a(x)|w|^{\beta_1-1}w + f_2(x) & \text{in } \Omega; \\ -\Delta_q w = \mu b(x)|z|^{\beta_2-1}z + g_2(x) & \text{in } \Omega; \\ z = w = 0 & \text{on } \partial\Omega, \end{cases} \quad (5)$$

with $f_1, f_2, g_1, g_2 \in L^\infty(\Omega)$ and $0 \leq f_1 \leq f_2$ and $0 \leq g_1 \leq g_2$ in Ω , one has $u \leq z$ and $v \leq w$ in Ω . If, in addition, at least, $u < z$ or $v < w$ in Ω whenever $f_1 + g_1 \not\equiv f_2 + g_2$ in Ω , we say that **(SCP)** corresponding to (1) holds in Ω . When $\lambda, \mu > 0$, **(SCP)** in Ω can be rewritten as $u < z$ and $v < w$ in Ω whenever $f_1 + g_1 \not\equiv f_2 + g_2$ in Ω . In the case that $u < z$ ($v < w$) in Ω , we clearly have $\frac{\partial}{\partial \nu} u(x_0) > \frac{\partial}{\partial \nu} z(x_0)$ ($\frac{\partial}{\partial \nu} v(x_0) > \frac{\partial}{\partial \nu} w(x_0)$) for all $x_0 \in \partial\Omega$.

Our next theorem characterizes completely the **(WCP)** and **(SCP)** corresponding to (1) in terms of the smooth curve \mathcal{C}_1 . Precisely:

Theorem 1.2. *Let \mathcal{R}_1 be as in Theorem 1.1 and $(\lambda, \mu) \in \mathbb{R}^2$. The following assertions are equivalent:*

- (i) $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$;
- (ii) **(WCP)** corresponding to (1) holds in Ω ;
- (iii) **(SCP)** corresponding to (1) holds in Ω .

The validity of weak and strong comparison principles for problems involving the p -Laplacian operator is usually very delicate (see [10, 11, 12, 13, 14, 19, 20, 21, 22, 23, 36, 39]). For example, the operator Δ_p satisfies the strong comparison principle only under the additional condition zero Dirichlet boundary values on $\partial\Omega$. So, we need assume the same condition on the **(WMP)**, **(SMP)**, **(WCP)** and **(SCP)** associated to the problem (1). Moreover, we prove the **(WCP)** and **(SCP)** under the following additional condition: $f_1 \geq 0$ and $g_1 \geq 0$ in Ω , when $(\lambda, \mu) \in \mathcal{R}_1$.

Now, we characterize when such **(WMP)** (or **(SMP)**) corresponding to (1) is satisfied in domains Ω of small Lebesgue measure. For this, we shall obtain an explicit lower estimate of Λ' in terms of the Lebesgue measure of Ω .

In Theorem 2.6 of [5], Berestycki, Nirenberg and Varadhan established a lower estimate for principal eigenvalues corresponding to the problem, in the scalar context, involving linear second order elliptic operators. In Theorem 5.1 of [31], López-Gómez obtained an explicit lower estimate for such principal eigenvalues. Later, in Theorem 10.1 of [6] Cano-Casanova and López-Gómez extended this result for mixed boundary conditions. The key tool for the proof of the Theorem 2.6 of [5] is an ABP estimate for second order uniformly elliptic operators while in proofs of the Theorems 5.1 of [31] and 10.1 of [6] is used in a crucial way the celebrated Faber-Krahn inequality

of Faber [17] and Krahn [25]. Recently, in [28] the author obtained an explicit lower bounds for principal eigenvalues of the system (1) via Faber-Krahn inequality for the first eigenvalue of $-\Delta_p$. Here, we will use an ABP estimate related to p -Laplacian operator with explicit constant (see Theorem 3 of [4]). Namely:

Theorem 1.3. *Let $a, b \in L^\infty(\Omega) \cap C(\Omega)$ and \mathcal{C}_1 be the principal curve associated to (1). Then*

$$\Lambda' \geq \frac{|\Omega|^{-\left(\frac{1}{ns} + \frac{1}{nr}\right)}}{c(n, p)^{\frac{\beta_2}{s}} c(n, q)^{\frac{\beta_1}{r}} d^{\frac{\beta_2}{s} + \frac{\beta_1}{r}} \|a\|_{L^\infty(\Omega)}^{\frac{1}{r}} \|b\|_{L^\infty(\Omega)}^{\frac{1}{s}}}, \quad (6)$$

where $d := \text{diam}(\Omega)$, $r := \sqrt{\beta_1(p-1)}$, $s := \sqrt{\beta_2(q-1)}$, $|\cdot|$ stands for the Lebesgue measure of \mathbb{R}^n ,

$$c(n, p) := \left(n \min\{1, p-1\} |B_1|^{\frac{1}{n}} \right)^{-\frac{1}{p-1}}$$

and B_1 is the unit ball of \mathbb{R}^n . In particular,

$$\lim_{|\Omega| \downarrow 0} \Lambda' = +\infty.$$

Note that we get an explicit lower estimate of Λ' in terms of the Lebesgue measure of Ω , diameter of Ω , explicit constants $c(n, p), c(n, q)$ and the weighted functions $a, b \in L^\infty(\Omega) \cap C(\Omega)$. As an interesting consequence of Theorems 1.1 and 1.3, we obtain the following characterization of maximum principles:

Theorem 1.4. *Let $r, s, c(n, p)$ and $c(n, q)$ be as in Theorem 1.3 and $a, b \in L^\infty(\Omega) \cap C(\Omega)$. Define $d := \text{diam}(\Omega)$ and*

$$\eta := \frac{1}{\left[\lambda^{\frac{1}{r}} \mu^{\frac{1}{s}} \left(c(n, p)^{\frac{\beta_2}{s}} c(n, q)^{\frac{\beta_1}{r}} d^{\frac{\beta_2}{s} + \frac{\beta_1}{r}} \|a\|_{L^\infty(\Omega)}^{\frac{1}{r}} \|b\|_{L^\infty(\Omega)}^{\frac{1}{s}} \right) \right]^{\frac{nrs}{r+s}}}.$$

The following assertions are equivalent:

- (i) $\lambda \geq 0$ and $\mu \geq 0$;
- (ii) **(WMP)** corresponding to (1) holds in Ω provided that $|\Omega| < \eta$;
- (iii) **(SMP)** corresponding to (1) holds in Ω provided that $|\Omega| < \eta$.

Finally, as application of Theorems 1.1 and 1.2, we present a characterization in terms of the principal curve \mathcal{C}_1 the set of couples $(\lambda, \mu) \in \mathbb{R}^2$ such that the system (3) admits a unique weak nonnegative solution (u, v) for any pair $(f, g) \in (L^\infty(\Omega))^2$ of nonnegative functions. Precisely, we have:

Theorem 1.5. *Let \mathcal{R}_1 be as in Theorem 1.1 and $(\lambda, \mu) \in \mathbb{R}^2$. Then, the couple $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$ if, and only if, the system (3) admits a unique weak solution $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ and this satisfies $u, v \geq 0$ in Ω for any pair $(f, g) \in (L^\infty(\Omega))^2$ of nonnegative functions. Moreover, $(u, v) \in (C^{1,\alpha}(\overline{\Omega}))^2$ for some $\alpha \in (0, 1)$.*

The rest of paper is organized into five sections. In Section 2 we show Theorem 1.1 by mean of maximum and comparison principles and Hopf's lemma related to the p -Laplacian operator. In Section 3 we characterize weak and strong comparison principles associated to the system (1) stated in Theorem 1.2 by using Theorem 1.1 as a key tool. In Section 4 we get an explicit lower estimate of Λ' in terms of the Lebesgue measure of Ω stated Theorem 1.3. The characterization of maximum principle in domains Ω of small Lebesgue measure stated in Theorem 1.4 is established in Section 5. Finally, in Section 6 we prove Theorem 1.5 by using Theorems 1.1 and 1.2.

2 Proof of Theorem 1.1

In this section we prove Theorem 1.1, which classify completely the couples $(\lambda, \mu) \in \mathbb{R}^2$ such that **(WMP)** and **(SMP)** hold in Ω . Notice that it suffices to show only that (i) \Leftrightarrow (ii). In this case, we clearly have (ii) \Leftrightarrow (iii). In fact, it is obvious that **(SMP)** in Ω implies **(WMP)** in Ω . Conversely, assume that **(WMP)** holds in Ω and let (u, v) be a weak solution of the system (3) with $f, g \in L^\infty(\Omega)$, $f, g \geq 0$ in Ω and $f + g \not\equiv 0$ in Ω . Thus, $u, v \geq 0$ in Ω and, by (i), we obtain $\lambda, \mu \geq 0$. Then, the conclusion of **(SMP)** follows, since Δ_p satisfies the strong maximum principle (see [22, 39, 41]).

In order to proof that **(WMP)** in Ω leads to $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$, assume instead that $(\lambda, \mu) \notin \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$. Let $(\lambda, \mu) \in \mathcal{C}_1$ and $(\tilde{\varphi}, \tilde{\psi})$ be a positive eigenfunction associated to (λ, μ) . Then, $(-\tilde{\varphi}, -\tilde{\psi})$ is a negative eigenfunction associated to (λ, μ) and so **(WMP)** fails in Ω .

Assume now that $(\lambda, \mu) \in \mathbb{R}^2$ is a fixed couple outside of $\overline{\mathcal{R}_1}$. If $(\lambda, \mu) \in (\mathbb{R}_+^*)^2$, we get $\lambda > \lambda_1$ and $\mu > \mu_1$, where (λ_1, μ_1) is a principal eigenvalue of (1) with $\frac{\mu}{\lambda} = \frac{\mu_1}{\lambda_1}$. Denote by (φ, ψ) a positive eigenfunction corresponding to (λ_1, μ_1) . Thus, we derive

$$\begin{cases} -\Delta_p(-\varphi) - \lambda a(x)|-\psi|^{\beta_1-1}(-\psi) &= -\lambda_1 a(x)\psi^{\beta_1} + \lambda a(x)\psi^{\beta_1} \\ &= (\lambda - \lambda_1)a(x)\psi^{\beta_1} \geq (\neq) 0 & \text{in } \Omega; \\ -\Delta_q(-\psi) - \mu b(x)|-\varphi|^{\beta_2-1}(-\varphi) &= -\mu_1 b(x)\varphi^{\beta_2} + \mu b(x)\varphi^{\beta_2} \\ &= (\mu - \mu_1)b(x)\varphi^{\beta_2} \geq (\neq) 0 & \text{in } \Omega \end{cases}$$

and $-\varphi = 0 = -\psi$ on $\partial\Omega$. Since, $-\varphi, -\psi < 0$ in Ω , **(WMP)** doesn't hold in Ω .

Now, suppose that $\lambda < 0$. Then, there exists $(\lambda_1, \mu_1) \in \mathcal{C}_1$ with $\lambda_1 > 0$ small enough (and so $\mu_1 > 0$ large enough) so that $\lambda < -\lambda_1$ and $\mu > -\mu_1$. Then, we get

$$\begin{cases} -\Delta_p(-\varphi) - \lambda a(x)\psi^{\beta_1} &= -\lambda_1 a(x)\psi^{\beta_1} - \lambda a(x)\psi^{\beta_1} \\ &= -(\lambda + \lambda_1)a(x)\psi^{\beta_1} \geq (\neq) 0 & \text{in } \Omega; \\ -\Delta_q\psi - \mu b(x)|-\varphi|^{\beta_2-1}(-\varphi) &= \mu_1 b(x)\varphi^{\beta_2} + \mu b(x)\varphi^{\beta_2} \\ &= (\mu + \mu_1)b(x)\varphi^{\beta_2} \geq (\neq) 0 & \text{in } \Omega \end{cases}$$

and $-\varphi = 0 = \psi$ on $\partial\Omega$. However, $-\varphi < 0$ in Ω and so **(WMP)** fails in Ω .

For the remaining case $\lambda \geq 0$ and $\mu < 0$, there exists $(\lambda_1, \mu_1) \in \mathcal{C}_1$ with $\lambda_1 > 0$ large enough (and so $\mu_1 > 0$ small enough) so that $\lambda > -\lambda_1$ and $\mu < -\mu_1$. Therefore,

$$\begin{cases} -\Delta_p\varphi - \lambda a(x)|-\psi|^{\beta_1-1}(-\psi) &= \lambda_1 a(x)\psi^{\beta_1} + \lambda a(x)\psi^{\beta_1} \\ &= (\lambda + \lambda_1)a(x)\psi^{\beta_1} \geq (\neq) 0 & \text{in } \Omega; \\ -\Delta_q(-\psi) - \mu b(x)\varphi^{\beta_2} &= -\mu_1 b(x)\varphi^{\beta_2} - \mu b(x)\varphi^{\beta_2} \\ &= -(\mu + \mu_1)b(x)\varphi^{\beta_2} \geq (\neq) 0 & \text{in } \Omega \end{cases}$$

and $\varphi = 0 = -\psi$ on $\partial\Omega$. But, $-\psi < 0$ in Ω and so again **(WMP)** fails in Ω .

Conversely, we next show that **(WMP)** holds in Ω for any couple $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$. Since Δ_p and Δ_q satisfies weak maximum principle in Ω , we have **(WMP)** holds in Ω if either $\lambda = 0$ and $\mu \geq 0$ or $\lambda \geq 0$ and $\mu = 0$. Now let $(\lambda, \mu) \in \mathcal{R}_1$. Let (u, v) be a weak solution of the system (3). Note that $\lambda < \lambda_1$ and $\mu < \mu_1$, where (λ_1, μ_1) is a principal eigenvalue of (1) with $\frac{\mu}{\lambda} = \frac{\mu_1}{\lambda_1}$. Let (φ, ψ) be a positive eigenfunction associated to (λ_1, μ_1) . Arguing by contradiction, assume that u or v is negative somewhere in Ω . Then, by strong comparison principle and Hopf's lemma for the p -Laplacian (see [22, 39, 41]), there exists some $\gamma > 0$ such that

$$-u \leq \gamma\varphi \text{ and } -v \leq \gamma^\omega\psi \text{ in } \Omega,$$

where $\omega := \frac{p-1}{\beta_1}$. Let $\bar{\gamma}$ be the minimum of such γ' s. Thus, $\bar{\gamma} > 0$. Since $\lambda < \lambda_1$ and $\mu < \mu_1$, we derive

$$\begin{cases} -\Delta_p(-u) \leq \lambda a(x)|-v|^{\beta_1-1}(-v) \leq \lambda a(x)(\bar{\gamma}^\omega\psi)^{\beta_1} \leq (\neq) \lambda_1 a(x)(\bar{\gamma}^\omega\psi)^{\beta_1} = -\Delta_p(\bar{\gamma}\varphi) & \text{in } \Omega; \\ -\Delta_q(-v) \leq \mu b(x)|-u|^{\beta_2-1}(-u) \leq \mu b(x)(\bar{\gamma}\varphi)^{\beta_2} \leq (\neq) \mu_1 b(x)(\bar{\gamma}\varphi)^{\beta_2} = -\Delta_q(\bar{\gamma}^\omega\psi) & \text{in } \Omega \end{cases}$$

and $\bar{\gamma}\varphi = -u = \bar{\gamma}^\omega\psi = -v = 0$ on $\partial\Omega$. It follows from the strong comparison principle to each above equation (see Theorem A.1 of [11]) that $-u < \bar{\gamma}\varphi$ and $-v < \bar{\gamma}^\omega\psi$ in Ω . Thus, we can find $0 < \varepsilon < 1$ such that $-u \leq \varepsilon\bar{\gamma}\varphi$ and $-v \leq (\varepsilon\bar{\gamma})^\omega\psi$ in Ω , contradicting the definition of $\bar{\gamma}$. Then, $u, v \geq 0$ in Ω . Hence, we complete the wished proof of theorem. ■

3 Proof of Theorem 1.2

In this section we establish the characterization of **(WCP)** and **(SCP)** in terms of the principal curve \mathcal{C}_1 as stated in Theorem 1.2. Using strong comparison principle (see Theorem A.1 of [11]) and arguing in a similar way as in the proof of Theorem 1.1, we see that it suffices to show only that (i) \Leftrightarrow (ii).

(ii) \Rightarrow (i) Taking $u, v \equiv 0$ in Ω (and so $f_1, f_2 \equiv 0$ in Ω), we have **(WCP)** in Ω implies **(WMP)** in Ω . Then, by Theorem 1.1, **(WCP)** in Ω implies $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$.

(i) \Rightarrow (ii) Consider $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$. Since Δ_p satisfies weak maximum and comparison principles in Ω , the conclusion is direct in the cases that $\lambda = 0$ or $\mu = 0$. Thus, it suffices to consider $(\lambda, \mu) \in \mathcal{R}_1$. In this case, since (u, v) is a weak solution of the problem (4), by Theorem 1.1, if $f_1, g_1 \equiv 0$ in Ω we have $u, v \equiv 0$ in Ω and if $f_1 + g_1 \not\equiv 0$ in Ω , we get $u, v > 0$ in Ω . Note that, if $f_1, g_1 \equiv 0$ in Ω , then the conclusion follows readily from **(WMP)** in Ω . Assume then $f_1 + g_1 \not\equiv 0$ in Ω (and so $u, v > 0$ in Ω). Thus, we also have $f_2 + g_2 \not\equiv 0$ in Ω . Therefore, since (z, w) is a weak solution of (5), by **(SMP)**, we have $z, w > 0$ in Ω .

To finish the proof of **(WCP)**, it suffices to show that $u \leq z$ in Ω . This follows directly by using the fact that Δ_q satisfies weak comparison principle in Ω (see [39]). Assume by contradiction that $u > z$ somewhere in Ω . In this case, the set $\Gamma := \{\gamma > 0 : z > \gamma u \text{ and } w > \gamma^\omega v \text{ in } \Omega\}$, where $\omega := \frac{p-1}{\beta_1}$, is nonempty by Hopf's Lemma (see [41]) and is also upper bounded. Set $\bar{\gamma} := \sup \Gamma > 0$. Notice that $z \geq \bar{\gamma}u$ and $w \geq \bar{\gamma}^\omega v$ in Ω . Note also that the statement of contradiction leads to $\bar{\gamma} < 1$. Thus, since $f_1 + g_1 \not\equiv 0$ and $f_2 + g_2 \not\equiv 0$ in Ω , we derive

$$\begin{cases} -\Delta_p(\bar{\gamma}u) = \lambda a(x)(\bar{\gamma}^\omega v)^{\beta_1} + \bar{\gamma}^{p-1}f_1 \leq (\neq) \lambda a(x)w^{\beta_1} + f_2 = -\Delta_p(z) & \text{in } \Omega; \\ -\Delta_q(\bar{\gamma}^\omega v) = \mu b(x)(\bar{\gamma}u)^{\beta_2} + \bar{\gamma}^{\beta_2}g_1 \leq (\neq) \mu b(x)z^{\beta_2} + g_2 = -\Delta_q(w) & \text{in } \Omega \end{cases}$$

and $\bar{\gamma}u = z = \bar{\gamma}^\omega v = w = 0$ on $\partial\Omega$. It follows from the strong comparison principle to each above equation (see Theorem A.1 of [11]) that $z > \bar{\gamma}u$ and $w > \bar{\gamma}^\omega v$ in Ω . Thus, we can find $0 < \varepsilon < 1$ such that $z \geq (\bar{\gamma} + \varepsilon)u$ and $w \geq (\bar{\gamma} + \varepsilon)^\omega v$ in Ω , contradicting the definition of $\bar{\gamma}$. This concludes the desired proof. ■

4 Proof of Theorem 1.3

Let $a, b \in L^\infty(\Omega) \cap C(\Omega)$, $(\lambda_1, \mu_1) \in \mathcal{C}_1$ and $(\varphi, \psi) \in X$ be a positive eigenfunction of the problem (1) corresponding to (λ_1, μ_1) . Since $\lambda_1 a \varphi^{\beta_1}, \mu_1 b \psi^{\beta_2} \in C(\Omega)$, by Theorem 1.8 of [33], we have (φ, ψ) is a viscosity subsolution of the system (1). Thus, applying the ABP estimate for the p -Laplacian (see Theorem 3 of [4]) to the first equation of (1), we have

$$\|\varphi\|_{L^\infty(\Omega)} = \sup_{\Omega} \varphi \leq c(n, p) d \lambda_1^{\frac{1}{p-1}} \|a\|_{L^\infty(\Omega)}^{\frac{1}{p-1}} \|\psi\|_{L^\infty(\Omega)}^{\frac{\beta_1}{p-1}} |\Omega|^{\frac{1}{n(p-1)}} \quad (7)$$

and for the q -Laplacian to the second equation of (1), we obtain

$$\|\psi\|_{L^\infty(\Omega)} \leq c(n, q) d \mu_1^{\frac{1}{q-1}} \|b\|_{L^\infty(\Omega)}^{\frac{1}{q-1}} \|\varphi\|_{L^\infty(\Omega)}^{\frac{\beta_2}{q-1}} |\Omega|^{\frac{1}{n(q-1)}}. \quad (8)$$

Therefore, joining inequalities (7) and (8) and using that

$$\beta_1 \beta_2 = (p-1)(q-1) \quad \text{and} \quad \lambda_1^{\frac{1}{p}} \mu_1^{\frac{1}{q}} = \Lambda',$$

we finally derive (6) and conclude the proof. ■

5 Proof of Theorem 1.4

By Theorem 1.1, the necessity of (i) and the equivalence between **(WMP)** and **(SMP)** follow directly. Thus, it suffices to prove that the assertion (i) implies (ii).

Assume that $\lambda \geq 0$ and $\mu \geq 0$. If either $\lambda = 0$ or $\mu = 0$, then by Theorem 1.1, the desired **(WMP)** follow.

Finally, assume now that $\lambda > 0$ and $\mu > 0$. We consider the positive constant η given by

$$\eta := \frac{1}{\left[\lambda^{\frac{1}{r}} \mu^{\frac{1}{s}} \left(c(n, p)^{\frac{\beta_2}{s}} c(n, q)^{\frac{\beta_1}{r}} d^{\frac{\beta_2}{s} + \frac{\beta_1}{r}} \|a\|_{L^\infty(\Omega)}^{\frac{1}{r}} \|b\|_{L^\infty(\Omega)}^{\frac{1}{s}} \right) \right]^{\frac{nrs}{r+s}}},$$

where $d := \text{diam}(\Omega)$, $r := \sqrt{\beta_1(p-1)}$, $s := \sqrt{\beta_2(q-1)}$ and $c(n, p)$ and $c(n, q)$ are the explicit constants of ABP estimate for the p -Laplacian and q -Laplacian, respectively. Then, by using the estimate (6) of Theorem 1.3, we obtain

$$\Lambda' \geq \frac{|\Omega|^{-\left(\frac{1}{ns} + \frac{1}{nr}\right)}}{c(n, p)^{\frac{\beta_2}{s}} c(n, q)^{\frac{\beta_1}{r}} d^{\frac{\beta_2}{s} + \frac{\beta_1}{r}} \|a\|_{L^\infty(\Omega)}^{\frac{1}{r}} \|b\|_{L^\infty(\Omega)}^{\frac{1}{s}}} > \lambda^{\frac{1}{r}} \mu^{\frac{1}{s}}$$

whenever $|\Omega| < \eta$. Therefore, we get $(\lambda, \mu) \in \mathcal{R}_1$ for such domains and so, by Theorem 1.1 the assertion (ii) holds. This concludes the proof of theorem. ■

6 Proof of Theorem 1.5

Let $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$ and $f, g \in L^\infty(\Omega)$ such that $f, g \geq 0$ in Ω . If either $\lambda = 0$ or $\mu = 0$, then clearly, by existence and uniqueness of weak solution for the problem (2), the system (3) admits a unique weak solution $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$. Applying the weak maximum principle to each equation of (3), we get $u, v \geq 0$ in Ω .

Assume now that $(\lambda, \mu) \in \mathcal{R}_1$. Then, by Theorem 3.1 of [11], the system (3) admits a unique weak solution $(u, v) \in X_+$. Let $(z, w) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ be a weak solution of the system (3). So, by **(WCP)**, $u = z$ and $v = w$ in Ω . Therefore, the system (3) admits a unique weak solution in $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ and this is nonnegative.

Conversely, assume that $(\lambda, \mu) \in \mathbb{R}^2$ and the problem (3) admits a unique weak solution $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ and that this satisfies $u, v \geq 0$ in Ω for any pair $(f, g) \in (L^\infty(\Omega))^2$ of nonnegative functions. Thus, **(WMP)** associated to (1) holds in Ω and so by Theorem 1.1, we have $(\lambda, \mu) \in \overline{\mathcal{R}_1} \setminus \mathcal{C}_1$. This ends the proof. ■

References

- [1] B. Alziary, J. Fleckinger, M.-H. Lécureux - *Principal eigenvalue and Maximum principle for some elliptic systems defined on general domains with refined Dirichlet boundary condition*, Comm. Math. Anal. 7 (2009), 1-11.
- [2] H. Amann - *Maximum principles and principal eigenvalues*, in *10 Mathematical Essays on Approximation in Analysis and Topology*, (J. Ferrera, J. López-Gómez and F. R. Ruiz del Portal eds.), pp. 1-60, Elsevier, Amsterdam, 2005.
- [3] I. Anton, J. López-Gómez - *Principal eigenvalue and maximum principle for cooperative periodic-parabolic systems*, Nonlinear Analysis 178 (2019), 152-189.
- [4] R. Argiolas, F. Charro, I. Peral - *On the Aleksandrov-Bakel'man-Pucci Estimate for Some Elliptic and Parabolic Nonlinear Operators*, Arch. Rational Mech. Anal. 202 (2011), 875-917.
- [5] H. Berestycki, L. Nirenberg, S.R.S. Varadhan - *The principal eigenvalue and maximum principle for second order elliptic operators in general domains*, Comm. pure Appl. Math. (1994), 47-92.
- [6] S. Cano-Casanova, J. López-Gómez - *Properties of the principal eigenvalues of a general class of nonclassical mixed boundary value problems*, J. Diff. Eq. 178 (2002), 123-211.
- [7] P.A. Caudevilla, J. López-Gómez - *Asymptotic behaviour of principal eigenvalues for a class of cooperative systems*, J. Diff. Eq. 244 (2008), 1093-1113.
- [8] Ph. Clément, D. G. de Figueiredo, E. Mitidieri - *Positive solutions of semilinear elliptic systems*, Comm. in PDE 17 (1992), 923-940.
- [9] Ph. Clément, R.F. Manásevich, E. Mitidieri - *Positive solutions for a quasilinear system via blow up*, Comm. P.D.E., 18 (1993), 2071-2106.
- [10] M. Cuesta, P. Takác - *A strong comparison principle for the Dirichlet p -Laplacian*, in *Reaction Diffusion Systems*, (G. Caristi and E. Mitidieri, Eds.). Lecture Notes in Pure and Applied Mathematics, 194. Marcel Dekker (1997).
- [11] M. Cuesta, P. Takác - *Nonlinear eigenvalue problems for degenerate elliptic systems*, Differ. Integral Equations 23 (2010), 1117-1138.
- [12] L. Damascelli - *Comparison theorems for some quasilinear degenerate elliptic operators and applications to symmetry and monotonicity results*, Ann. Inst. H. Poincaré. Analyse non linéaire 15 (1998), 493-516.
- [13] L. Damascelli, B. Sciunzi - *Regularity, monotonicity and symmetry of positive solutions of m -Laplace equations*, J. Differential Equations 206 (2004), 483-515.

- [14] L. Damascelli, B. Sciunzi - *Harnack inequalities, maximum and comparison principles, and regularity of positive solutions of m -Laplace equations*, Calc. Var. Partial Differential Equations 25 (2006), 139-159.
- [15] D. G. de Figueiredo, P. L. Felmer - *On superquadratic elliptic systems*, Trans. Amer. Math. Soc. 343 (1994), 99-116.
- [16] DE. DiBenedetto - *$C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations*, Nonlinear Anal. 7 (1983), 827-850.
- [17] G. Faber - *Beweis, dass unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt (in German)*, Sitzungsberichte der mathematisch-physikalischen Klasse der Bayerischen Akademie der Wissenschaften zu München Jahrgang (1923), 169-172.
- [18] P. Felmer, S. Martínez - *Existence and uniqueness of positive solutions to certain differential systems*, Adv. Differential Equations 4 (1998), 575-593.
- [19] J. Fleckinger-Pellé, J. P. Gossez, P. Takác, F. de Thélin - *Nonexistence of solutions and an anti-maximum principle for cooperative systems with the p -Laplacian*, Math. Nachr. 194 (1998), 49-78.
- [20] J. Fleckinger-Pellé, J. Hernández, P. Takác, F. de Thélin - *Uniqueness and positivity for solutions of equations with the p -Laplacian*. Reaction diffusion systems (Trieste, 1995), 141-155, Lecture Notes in Pure and Appl. Math., 194, Dekker, New York, 1998.
- [21] J. Fleckinger-Pellé, P. Takác - *Uniqueness of positive solutions for nonlinear cooperative systems with the p -Laplacian*, Indiana Univ. Math. J. 43 (1994), 1227-1253.
- [22] J. García-Melián, J. C. Sabina de Lis - *Maximum and comparison principles for operators involving the p -Laplacian*, J. Math. Anal. Appl. 218 (1998), 49-65.
- [23] M. Guedda, L. Veron - *Quasilinear elliptic equations involving critical Sobolev exponents*, Nonlinear Anal. 13 (1989), 879-902.
- [24] J. Hulshof, R. C. A. M. van der Vorst - *Differential systems with strongly indefinite variational structure*, J. Funct. Analysis, 114 (1993), 32-58.
- [25] E. Krahn - *Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises*, Math. Ann. 91 (1925), 97-100.
- [26] E. J. F. Leite, M. Montenegro - *Maximum and comparison principles to Lane-Emden systems*, Journal of the London Mathematical Society 101 (2020), 23-42.
- [27] E. J. F. Leite, M. Montenegro - *Principal curves to nonlocal Lane-Emden systems and related maximum principles*, Calc. Var. Partial Differential Equations 59 (2020), 118.

- [28] E. J. F. Leite - *On the principal eigenvalues of the degenerate elliptic systems*, Electronic Journal of Qualitative Theory of Differential Equations 40 (2020), 1-15.
- [29] G. Lieberman - *Boundary regularity for solutions of degenerate elliptic equations*, Non-linear Anal. 12 (1988), 1203-1219.
- [30] J. López-Gómez - *Linear Second Order Elliptic Operators*, World Scientific, Singapore, 2013.
- [31] J. López-Gómez - *The maximum principle and the existence of principal eigenvalue for some linear weighted boundary value problems*, J. Diff. Eq. 127 (1996), 263-294.
- [32] J. López-Gómez, M. Molina-Meyer - *The maximum principle for cooperative weakly coupled elliptic systems and some applications*, Diff. Int. Equations 7 (1994), 383-398.
- [33] M. Medina, P. Ochoa - *On viscosity and weak solutions for non-homogeneous p -Laplace equations*, Adv. Nonlinear Anal. 8 (2019), 468-481.
- [34] E. Mitidieri - *A Rellich type identity and applications*, Comm. Partial Differential Equations 18 (1993), 125-151.
- [35] M. Montenegro - *The construction of principal spectra curves for Lane-Emden systems and applications*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 29 (2000), 193-229.
- [36] P. Pucci, J. Serrin - *The maximum principle*, Birkhäuser Verlag, Basel, 2007.
- [37] J. Serrin, H. Zou - *Existence of positive entire solutions of elliptic Hamiltonian systems*, Comm. Partial Differential Equations 23 (1998), 577-599.
- [38] G. Sweers - *Strong positivity in $C(\overline{\Omega})$ for elliptic systems*, Math. Z. 209 (1992), 251-271.
- [39] P. Tolksdorf - *On the Dirichlet problem for quasilinear equations in domains with conical boundary points*, Comm. P.D.E., 8 (1983), 773-817.
- [40] P. Tolksdorf - *Regularity for a more general class of quasilinear elliptic equations*, J. Differential Equations, 51 (1984), 126-150.
- [41] J. L. Vázquez - *A strong maximum principle for some quasilinear elliptic equations*, Appl. Math. Optim., 12 (1984), 191-202.