arXiv:2005.12698v1 [math.CA] 23 May 2020

BÉZIER VARIANT OF GENERALIZED BERNSTEIN-DURRMEYER TYPE OPERATORS

Karunesh Kumar Singh
Department of Applied Sciences and Humanities
Institute of Engineering and Technology
Lucknow-226021(Uttar Pradesh), India
Email: kksiitr.singh@gmail.com
and
Asha Ram Gairola
Department of Mathematics, Doon University
Dehradun-248001 (Uttarakhand), India
Email: ashagairola@gmail.com

Abstract. In this paper, we define Bézier variant of generalized Bernstein-Durrmeyer type operators of second order, introduced by Ana et al. Then, we find an error estimate in terms of terms of Ditzian Totik modulus of smoothness. Next, we study rate of approximation for larger class function of bounded variation.

1. Introduction

For $C[0,1]$, the space of continuous functions defined on $[0,1]$, and let $n \in \mathbb{N}$, the Bernstein polynomial operators $B_n(f) : C[0,1] \longrightarrow C[0,1]$ are given by

$$B_n(f;x) := \sum_{k=0}^{n} p_{n,k}(x) f\left(\frac{k}{n}\right), \quad x \in [0,1],$$

where the basis functions for the above polynomials $p_{n,k}(x) = \binom{n}{k} x^k (1-x)^{n-k}$ for $0 \leq k \leq n$.The fundamental polynomials satisfy following recurrence relation:

$$p_{n,k}(x) = (1-x)p_{n-1,k}(x) + xp_{n-1,k-1}(x), 0 < k < n. \tag{1.1}$$

The Bernstein polynomial operators have some interesting properties eg. preservation of convexity, Lipschitz constant and monotonicity etc. Although $B_n(f;x)$ are not suitable to approximate integrable function as such. In order to approximate bounded and integrable functions on $[0,1]$, Durrmeyer [5] and Lupaş [12] introduced its integral modifications as follows:

$$D_n(f;x) := (n+1) \int_0^1 \sum_{k=0}^{n} p_{n,k}(x) p_{n,k}(u) f(u)\, \mathrm{d}u. \tag{1.2}$$

AMS Subject Classification: 41A25, 41A28, 41A36
Key Words. Linear positive operators, Rate of approximation, Modulus of continuity, Bezier variant of operators, Ditzian Totik modulus of smoothness.

These operators are obtained by a suitable modification of Bernstein polynomial operators, wherein the values $f(k/n)$ are replaced by the average values

$$(n+1)\int_0^1 p_{n,k}(u)f(u)\,\mathrm{d}u$$

of f in the interval $[0,1]$. The operators $D_n(f;x)$ have been extensively studied by Derriennic [2] and other authors ([3],[5],[7]),[9], [14]. It is seen that the convergence of operators $D_n(f;x)$ to $f(x)$ is uniform. Also, it turns out that the order of approximation by the operators $D_n(f;x)$ is at best, $O(n^{-1})$ however smooth the function may be.

It is known that ([8]) linear positive operators e.g. Bernstein polynomials operators, Baskakov operators, integral variants of Bernstein operators and others are saturated by order n^{-1}. Consequently, it is not possible to improve the rate of convergence by these operators except for linear functions. Recently a new approach to improve the rate of approximation by decomposition of the its weight function $p_{n,k}(x)$ was recently introduced by Khosravian-Arab et al.[11]. The authors of [11] applied this technique to the Bernstein polynomials operators and were improved the order of approximation from linear order $O(n^{-1})$ to the quadratic order $O(n^{-2})$ and cubic order $O(n^{-3})$ also. Subsequently, this technique was also applied to the Bernstein-Durrmeyer operators $D_n(f;x)$ by Ana Maria et al. in [1].

For $f \in L_B[0,1]$, Acu et al. [1] introduced generalized Bernstein Durrmeyer polynomial operators of order II as follows:

$$D_n^{M,2}(f;x) := (n+1)\sum_{k=0}^{n} p_{n,k}^{M,2}(x)\int_0^1 p_{n,k}(u)f(u)\,\mathrm{d}u, n \geq 3 \tag{1.3}$$

where

$$p_{n,k}^{M,2}(x) = g(x,n)p_{n-2,k}(x) + h(x,n)p_{n-2,k-1}(x) + g(1-x,n)p_{n-2,k-2}(x), \tag{1.4}$$

and $p_{n-2,k}(x) = 0$ if $k < 0$ and $k > n-2$ and the function $g(x,n) = g_2(n)x^2 + g_1(n)x + g_0(n)$ and $h(x,n) = h_0(n)x(1-x)$, where $g_i(n)$, $h_i(n)$ are the unknown sequences to be determined suitably by imposing the identities $D_n^{M,2}(e_i;x) = e_i(x)$ for $i = 0,1$ and for $i = 2$. So we get the condition $2g_2(n) - h_0(n) = 0$, $g_2(n) + 2b_0(n) + b_1(n) = 1$. Particularly, if we take $g_2(n) = g_0(n) = 1$, $g_1(n) = -2$ and $h_0(n) = 2$. Then the fundamental polynomials (1.4) for the operators $D_n^{M,2}(f;x)$ is reduced to fundamental polynomials for Bernstein polynomials in (1.1). Moreover, if we assume sequences such that $g_0(n) = \frac{3}{2}, g_1(n) = -n$ and $g_2(n) = n-2$ and $h_0(n) = 2(n-2)$. Then we have,

Lemma 1. [1] *We have*

$$D_n^{M,2}((u-x);x) = 0,$$

$$D_n^{M,2}((u-x)^2;x) = \frac{20x(1-x)}{(n+2)(n+3)} - \frac{3}{(n+2)(n+3)}.$$

Remark 1. *By an application of lemma 1, we have*

$$D_n^{M,2}((u-x)^2;x) \leq \frac{C}{n+2}\varphi^2(x), x \in [0,1], \tag{1.5}$$

where $\varphi^2(x) = x(1-x)$ *and* C *is positive constant.*

Very recently, Kajla and Acar [10] has applied this method to $\alpha-$ Bernstein operators. Motivated to study of various variants of these operators, we define the Bézier Variant of the operator $D_n^{M,2}(f;x)$ for a larger class, namely the class of functions of bounded variation. For $\mu \geq 1$ and $f \in L_B[0,1]$, we define Bézier variant of the operator $D_n^{M,2}(f;x)$ as

$$D_{n,\mu}^{M,2}(f;x) := (n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x)\int_0^1 p_{n,k}(u)f(u)\,\mathrm{d}u, \tag{1.6}$$

where

$$Q_{n,k}^{(\mu)}(x) = (J_{n,k}(x))^{\mu} - (J_{n,k+1}(x))^{\mu}, J_{n,k}(x) = \sum_{j=k}^{n} K_{n,j}(x).$$

For $\mu = 1$, this family of Bézier operators yield the operators $D_{n,\mu}^{M,2}(f;x)$ studied by Acu et al. [1]. Let

$$W_{n,\mu}(x,u) = (n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x)p_{n,k}(u).$$

Then $D_{n,\mu}^{M,2}(f;x) := \int_0^1 W_{n,\mu}(x,u)f(u)\mathrm{d}u$.

Lemma 2. *If* $f \in C[0,1]$ *then,* $\|D_n^{M,2}(f)\| \leq \|f\|$, *where* $\|.\|$ *denotes the sup-norm on* $[0,1]$.

Lemma 3. *If* $f \in C[0,1]$ *then,* $\|D_{n,\mu}^{M,2}(f)\| \leq \mu\|f\|$.

Proof. Using the inequality $|a^{\mu} - b^{\mu}| \leq \mu|a-b|$ with $0 \leq a, b \leq 1, \mu \geq 1$ and definition of $D_{n,\mu}^{M,2}(f)$, we get for $\mu \geq 1$

$$0 < [(J_{n,k}(x))^{\mu} - (J_{n,k+1}(x))^{\mu}] \leq \mu[J_{n,k}(x) - J_{n,k+1}(x)] = \mu K_{n,k}(x) \tag{1.7}$$

Applying $D_{n,\mu}^{M,2}(f;x)$ and Lemma 3, we get

$$\|D_{n,\mu}^{M,2}(f)\| \leq \mu\|D_n^{M,2}(f)\| \leq \mu\|f\|.$$

□

2. Direct result

Now, we derive a direct result for Bézier operators in terms of Ditzian Totik modulus of smoothness $\omega_{\varphi}(f,t)$.

Theorem 1. *Let $\varphi(x) = \sqrt{x(1-x)}$, $\mu \geq 1$ and $x \in [0,1]$. If $f \in C[0,1]$ then*

$$|D_{n,\mu}^{M,2}(f;x) - f(x)| \leq C\omega_{\varphi}\left(f, \sqrt{\frac{1}{n+2}}\right),$$

where C is any absolute constant.

Proof. We know that $g(t) = g(x) + \int_x^t g'(u)\mathrm{d}u$. So

$$|D_{n,\mu}^{M,2}(g;x) - g(x)| = \left|D_{n,\mu}^{M,2}\left(\int_x^t g'(u)\mathrm{d}u, x\right)\right|. \tag{2.1}$$

For any $x, t \in (0,1)$, we have

$$\left|\int_x^t g'(u)\mathrm{d}u\right| \leq \|\varphi g'\|\left|\int_x^t \frac{1}{\varphi(u)}\mathrm{d}u\right|. \tag{2.2}$$

Therefore,

$$\begin{aligned}
\left|\int_x^t \frac{1}{\varphi(u)}\mathrm{d}u\right| &= \left|\int_x^t \frac{1}{\sqrt{u(1-u)}}\mathrm{d}u\right| \\
&\leq \left|\int_x^t \left(\frac{1}{\sqrt{u}} + \frac{1}{\sqrt{1-u}}\right)\mathrm{d}u\right| \\
&\leq 2\left(|\sqrt{t} - \sqrt{x}| + |\sqrt{1-t} - \sqrt{1-x}|\right) \\
&= 2|t-x|\left(\frac{1}{\sqrt{t}+\sqrt{x}} + \frac{1}{\sqrt{1-t}+\sqrt{1-x}}\right) \\
&< 2|t-x|\left(\frac{1}{\sqrt{x}} + \frac{1}{\sqrt{1-x}}\right) \\
&< \frac{2\sqrt{2}|t-x|}{\varphi(x)}. \qquad (2.3)
\end{aligned}$$

Combining (2.1)-(2.3) and using Cauchy-Schwarz inequality, we have that

$$\begin{aligned}
|D_{n,\mu}^{M,2}(g;x) - g(x)| &\leq 2\sqrt{2}\|\varphi g'\|\varphi^{-1}(x)D_{n,\mu}^{M,2}(|t-x|;x) \\
&\leq 2\sqrt{2}\|\varphi g'\|\varphi^{-1}(x)\left(D_{n,\mu}^{M,2}((t-x)^2;x)\right)^{1/2}.
\end{aligned}$$

Now using inequality (1.5), we obtain

$$|D_{n,\mu}^{M,2}(g;x) - g(x)| < C\sqrt{\frac{1}{(n+2)}}\|\varphi g'\|. \tag{2.4}$$

Using Lemma 1 and inequality (2.4), we obtain

$$\begin{aligned} |D_{n,\mu}^{M,2}(f;x)-f| &\leq |D_{n,\mu}^{M,2}(f-g;x)|+|f-g|+|D_{n,\mu}^{M,1}(g;x)-g(x)| \\ &\leq C\left(\|f-g\|+\sqrt{\frac{1}{(n+2)}}\|\varphi g'\|\right). \end{aligned} \tag{2.5}$$

Taking infimum on both sides of (2.5) over all $g \in W_{\phi}^{2}$, we reach to

$$|D_{n,\mu}^{M,2}(f;x)-f| \leq CK_{\varphi}\left(f,\sqrt{\frac{1}{(n+2)}}\right)$$

Using relation $K_{\varphi}(f,t) \sim \omega_{\varphi}(f,t)$, we get the required result. □

Ozarslan and Aktuglu [13] is considered the Lipschitz-type space with two parameters $\alpha_1 \geq 0, \alpha_2 > 0$, which is defined as

$$Lip_{M}^{(\alpha_1,\alpha_2)}(\zeta)=\left\{f \in C[0,1] : |f(t)-f(x)| \leq C_0 \frac{|t-x|^{\zeta}}{(t+\alpha_1 x^2+\alpha_2 x)^{\zeta/2}} : t \in [0,1], x \in (0,1)\right\},$$

where $0 < \zeta \leq 1$, and C_0 is any absolute constant.

Theorem 2. *Let $f \in Lip_{M}^{(\alpha_1,\alpha_2)}(\zeta)$. Then for all $x \in (0,1]$, there holds:*

$$|D_{n,\mu}^{M,2}(f;x)-f| \leq C\left(\frac{\mu\varphi^2(x)}{(n+2)(\alpha_1 x^2+\alpha_2 x)}\right)^{\frac{\zeta}{2}},$$

where C is any absolute constant.

Proof. Using Lemma 1 and (1.7) and Hölder's inequality with $p = \frac{2}{\zeta}$ and $p = \frac{2}{2-\zeta}$, we get

$$
\begin{aligned}
&|D_{n,\mu}^{M,2}(f;x) - f(x)| \\
&\leq (n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x) \int_0^1 p_{n,k}(u)|f(u)-f(x)|\,\mathrm{d}u \\
&\leq (n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x) \left(\int_0^1 p_{n,k}(u)|f(u)-f(x)|^{\frac{2}{\zeta}}\,\mathrm{d}u\right)^{\frac{\zeta}{2}} \\
&\leq \left[(n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x) \int_0^1 p_{n,k}(u)|f(u)-f(x)|^{\frac{2}{\zeta}}\,\mathrm{d}u\right]^{\frac{\zeta}{2}} \times \\
&\left[(n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x) \int_0^1 p_{n,k}(u)\,\mathrm{d}u\right]^{\frac{2-\zeta}{2}} \\
&= \left[(n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x) \int_0^1 p_{n,k}(u)|f(u)-f(x)|^{\frac{2}{\zeta}}\,\mathrm{d}u\right]^{\frac{\zeta}{2}} \\
&\leq C\left((n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x) \int_0^1 p_{n,k}(u)\frac{(u-x)^2}{(u+\alpha_1 x^2+\alpha_2 x)}\mathrm{d}u\right)^{\frac{\zeta}{2}} \\
&\leq \frac{C}{(\alpha_1 x^2+\alpha_2 x)^{\frac{\zeta}{2}}}\left((n+1)\sum_{k=0}^{n} Q_{n,k}^{(\mu)}(x) \int_0^1 p_{n,k}(u)(u-x)^2\mathrm{d}u\right)^{\frac{\zeta}{2}} \\
&\leq \frac{C}{(\alpha_1 x^2+\alpha_2 x)^{\frac{\zeta}{2}}}[D_{n,\mu}^{M,2}((u-x)^2;x)]^{\frac{\zeta}{2}} \\
&\leq C\left(\frac{\mu\varphi^{(}x)}{(n+2)(\alpha_1 x^2+\alpha_2 x)}\right)^{\frac{\zeta}{2}}.
\end{aligned}
$$

Hence, the desired result follows. □

3. Rate of convergence

If we define

$$\kappa_{n,\mu}(x,y) = \int_0^y W_{n,k}(x,t)\mathrm{d}t.$$

Then, it is obvious that $\kappa_{n,\mu}(x,1) = 1$.

Lemma 4. *Let $x \in (0,1)$ and $C > 2$ then for sufficiently large n we have*

$$\kappa_{n,\mu}(x,y) = \int_0^y W_{n,\mu}(x,t)dt \leq \frac{C\mu x(1-x)}{n(x-y)^2}, 0 < y < x$$

$$1 - \kappa_{n,\mu}(x,z) = \int_z^1 W_{n,\mu}(x,t)dt \leq \frac{C\mu x(1-x)}{n(z-x)^2}, x < z < 1.$$

Remark 2. *We have*

$$\begin{aligned}\kappa_{n,\mu}(x,y) &= \int_0^y W_{n,\mu}(x,t)dt \leq \int_0^y W_{n,\mu}(x,t)\frac{(t-x)^2}{(y-x)^2}dt \\ &= \frac{D^{M,2}_{n,\mu}((t-x)^2;x)}{(y-x)^2} \leq \frac{\mu D^{M,2}_{n}((t-x)^2;x)}{(y-x)^2} \leq \frac{\mu}{n+2}.\frac{\varphi^2(x)}{(y-x)^2}.\end{aligned}$$

Now, we discuss the approximation properties of functions having derivative of bounded variation on $[0,1]$. Let $\mathbb{B}[0,1]$ denote the class of differentiable functions g defined on $[0,1]$, whose derivative g' is of bounded variation on $[0,1]$. The functions $g \in \mathbb{B}[0,1]$ is expressed as $g(x) = \int_0^x h(t)dt + g(0)$, where $h \in \mathbb{B}[0,1]$, i.e., h is a function of bounded variation on $[0,1]$.

Theorem 3. *Let $f \in \mathbb{B}[0,1]$ then for $\mu \geq 1, 0 < x < 1$ a and sufficiently large n we have*

$$\begin{aligned}\left|D^{M,2}_{n,\mu}(f;x) - f(x)\right| \leq{}& \left(\frac{1}{\mu+1}|f'(x+) + \mu f'(x-)| + |f'(x+) - f'(x-)|\right)\frac{\mu}{n+2}\varphi^2(x) \\ &+ \frac{\mu}{n+2}.\frac{\varphi^2(x)}{x^2}\sum_{k=1}^{\sqrt{n}}\left(\bigvee_{x-\frac{x}{k}}^{x}(f')_x\right) + \frac{x}{\sqrt{n}}\left(\bigvee_{x}^{x+\frac{1-x}{\sqrt{n}}}(f')_x\right)\left(\bigvee_{x-\frac{x}{k}}^{x}(f')_x\right) \\ &+ \frac{\mu}{n+2}.\frac{\varphi^2(x)}{1-x}\sum_{k=1}^{\sqrt{n}}\left(\bigvee_{x}^{x+\frac{1-x}{k}}(f')_x\right) + \frac{1-x}{\sqrt{n}}\left(\bigvee_{x}^{x+\frac{1-x}{\sqrt{n}}}(f')_x\right).\end{aligned}$$

$$(f')_x(t) = \begin{cases} f'(t) - f'(x-), & 0 \leq t < x \\ 0, & t = x \\ f'(t) - f'(x+), & x < t \leq 1. \end{cases}$$

Proof. As we know that $D_{n,\mu}^{M,2}(1;x)=1$. Therefore, we have

$$\begin{aligned} D_{n,\mu}^{M,2}(f;x)-f(x) &= \int_0^1 W_{n,\mu}(x,u)(f(u)-f(x))\,\mathrm{d}u \\ &= \int_0^1 W_{n,\mu}(x,u)\int_x^u f'(t)\,dt\,\mathrm{d}u \end{aligned} \tag{3.1}$$

Since $f\in\mathbb{B}[0,1]$, we may write

$$\begin{aligned} f'(t) &= \frac{f'(x+)+\mu f'(x-)}{\mu+1}+(f')_x(t)+\frac{f'(x+)-f'(x-)}{2}\left(\mathrm{sign}(t-x)+\frac{\mu-1}{\mu+1}\right) \\ &+ \delta_x(t)\left(f'(t)-\frac{f'(x+)+\mu f'(x-)}{2}\right) \end{aligned} \tag{3.2}$$

where

$$\mathrm{sign}(t)=\begin{cases} 1, & t>0 \\ 0, & t=0 \\ -1, & t<0. \end{cases}$$

and

$$\delta_x(t)=\begin{cases} 1, & t=x \\ 0, & t\neq x. \end{cases}$$

Putting the value of $f'(t)$ form (3.2) in 3.1, we get estimates corresponding to four terms of (3.2), say K_1,K_2,K_3 and K_4 respectively. Obviously,

$$K_4=\int_0^1\left(\int_x^u\left(f'(t)-\frac{f'(x+)+\mu f'(x-)}{2}\right)\delta_x(t)dt\right)W_{n,\mu}(x,u)\mathrm{d}u=0.$$

Now, Using Cauchy's Schwarz inequality and Lemma 4, we obtain

$$\begin{aligned} K_1=\int_0^1\left(\int_x^u\frac{f'(x+)+\mu f'(x-)}{\mu+1}dt\right)W_{n,\mu}(x,u)\mathrm{d}u &= \frac{f'(x+)+\mu f'(x-)}{\mu+1}\int_0^1(u-x)W_{n,\mu}(x,u)\mathrm{d}u \\ &= \frac{f'(x+)+\mu f'(x-)}{\mu+1}D_{n,\mu}^{M,1}((u-x);x) \\ &\leq \frac{1}{\sqrt{n+2}}\frac{f'(x+)+\mu f'(x-)}{\mu+1}\varphi(x). \end{aligned}$$

Next, again using Cauchy's Schwarz inequality and Lemma 4, we obtain

$$\begin{aligned}
K_3 &= \int_0^1 \left(\int_x^u \left(\frac{f'(x+) - f'(x-)}{2} \right) \left(\operatorname{sign}(t-x) + \frac{\mu-1}{\mu+1} \right) dt \right) W_{n,\mu}(x,u)\mathrm{d}u \\
&= \left(\frac{f'(x+) - f'(x-)}{2} \right) \Bigg[- \int_0^x \left(\int_u^x \left(\operatorname{sign}(t-x) + \frac{\mu-1}{\mu+1} \right) dt \right) W_{n,\mu}(x,u)\mathrm{d}u \\
&+ \int_x^1 \left(\int_x^u \left(\operatorname{sign}_\mu(t-x) + \frac{\mu-1}{\mu+1} \right) dt \right) W_{n,\mu}(x,u)\mathrm{d}u \Bigg] \\
&\leq |f'(x+) - f'(x-)| \int_0^1 |u-x| W_{n,\mu}(x,u)\mathrm{d}u \\
&\leq |f'(x+) - f'(x-)|\, D_{n,\mu}^{M,1}(|u-x|;x) \\
&\leq \frac{1}{\sqrt{n+2}} |f'(x+) - f'(x-)|\, \varphi(x).
\end{aligned}$$

Now, we estimate K_2 as follows:

$$\begin{aligned}
K_2 = \int_0^1 \left(\int_x^u (f')_x(t)dt \right) W_{n,\mu}(x,u)\mathrm{d}u &= \int_0^x \left(\int_x^u (f')_x(t)dt \right) W_{n,\mu}(x,u)\mathrm{d}u \\
&+ \int_x^1 \left(\int_x^u (f')_x(t)dt \right) W_{n,\mu}(x,u)\mathrm{d}u \\
&= K_5 + K_6, \text{say}.
\end{aligned}$$

Using Lemma 4 and definition of $\kappa_{n,\mu}(x,u)$, we may write

$$K_5 = \int_0^x \left(\int_x^u (f')_x(t)dt \right) \frac{d}{\mathrm{d}u} \kappa_{n,\mu}(x,u)\mathrm{d}u$$

Integrating by parts, we obtain

$$\begin{aligned}|K_5| &\leq \int_0^x |(f')_x(u)|\kappa_{n,\mu}(x,u)\mathrm{d}u \\ &\leq \int_0^{x-x/\sqrt{n}} |(f')_x(u)|\kappa_{n,\mu}(x,u)\mathrm{d}u + \int_{x-x/\sqrt{n}}^x |(f')_x(u)|\kappa_{n,\mu}(x,u)\mathrm{d}u \\ &= K_7 + K_8, \text{say.}\end{aligned}$$

In view of facts $(f')_x(x) = 0$ and $\kappa_{n,\mu}(x,u) \leq 1$, we get

$$\begin{aligned}K_8 &= \int_0^x |(f')_x(u) - (f')_x(x)|\kappa_{n,\mu}(x,u)\mathrm{d}u \\ &\leq \int_{x-x/\sqrt{n}}^x \left(\bigvee_u^x (f')_x\right)\mathrm{d}u \\ &\leq \left(\bigvee_u^x (f')_x\right)\int_{x-x/\sqrt{n}}^x \mathrm{d}u = \frac{x}{\sqrt{n}}\left(\bigvee_u^x (f')_x\right).\end{aligned}$$

Using Lemma 4, definition of $\kappa_{n,\mu}(x,u)$, and transformation $u = x - \frac{x}{t}$ we may write

$$\begin{aligned}K_7 &\leq \frac{\mu}{n+2}\varphi^2(x)\int_0^{x-x/\sqrt{n}} |(f')_x(u) - (f')_x(x)|\frac{\mathrm{d}u}{(u-x)^2} \\ &\leq \frac{\mu}{n+2}\varphi^2(x)\int_0^{x-x/\sqrt{n}} \left(\bigvee_u^x (f')_x\right)\frac{\mathrm{d}u}{(u-x)^2} \\ &\leq \frac{\mu}{n+2}\frac{\varphi^2(x)}{x^2}\int_1^{\sqrt{n}} \left(\bigvee_{x-\frac{x}{t}}^x (f')_x\right)dt \\ &\leq \frac{\mu}{n+2}\frac{\varphi^2(x)}{x^2}\sum_{k=1}^{|\sqrt{n}|}\left(\bigvee_{x-\frac{x}{t}}^x (f')_x\right).\end{aligned}$$

Combining the estimates of I_7 and I_8, we have

$$|K_5| \leq \frac{\mu}{n+2}\frac{\varphi^2(x)}{x^2}\sum_{k=1}^{|\sqrt{n}|}\left(\bigvee_{x-\frac{x}{t}}^x (f')_x\right) + \frac{x}{\sqrt{n}}\left(\bigvee_u^x (f')_x\right)$$

In order to estimate K_6, we use integration by parts, Lemma 4 and transformation $z = x + \frac{1-x}{\sqrt{n}}$. Therefore,we proceed as follows:

$$
\begin{aligned}
|K_6| &= \left| \int_x^1 \left(\int_x^u (f')_x(t)dt \right) W_{n,\mu}(x,u)\mathrm{d}u \right| \\
&= \left| \int_x^z \left(\int_x^u (f')_x(t)dt \right) \frac{\partial}{\partial u}(1-\kappa_{n,\mu}(x,u))\mathrm{d}u + \int_z^1 \left(\int_x^u (f')_x(t)dt \right) \frac{\partial}{\partial u}(1-\kappa_{n,\mu}(x,u))\mathrm{d}u \right| \\
&= \left| \left[\int_x^u (f')_x(t)dt(1-\kappa_{n,\mu}(x,u)) \right]_x^z - \int_x^z (f')_x(u)(1-\kappa_{n,\mu}(x,u))\mathrm{d}u \right. \\
&\quad \left. + \left[\int_x^u (f')_x(t)dt(1-\kappa_{n,\mu}(x,u)) \right]_z^1 - \int_z^1 (f')_x(u)(1-\kappa_{n,\mu}(x,u))\mathrm{d}u \right| \\
&= \left| \int_x^z (f')_x(u)(1-\kappa_{n,\mu}(x,u))\mathrm{d}u + \int_z^1 (f')_x(u)(1-\kappa_{n,\mu}(x,u))\mathrm{d}u \right| \\
&\leq \frac{\mu}{n+2}.\varphi^2(x) \int_z^1 \left(\bigvee_x^u (f')_x \right) (u-x)^{-2}\,\mathrm{d}u + \int_x^z \left(\bigvee_x^u (f')_x \right) \mathrm{d}u \\
&\leq \frac{\mu}{n+2}.\varphi^2(x) \int_{x+\frac{1-x}{\sqrt{n}}}^1 \left(\bigvee_x^u (f')_x \right) (u-x)^{-2}\,\mathrm{d}u + \frac{1-x}{\sqrt{n}} \left(\bigvee_x^{x+\frac{1-x}{\sqrt{n}}} (f')_x \right).
\end{aligned}
$$

Now, substituting $t = \frac{1-x}{t-x}$, we get

$$
\begin{aligned}
|K_6| &\leq \frac{\mu}{n+2}.\varphi^2(x) \int_1^{\sqrt{n}} \left(\bigvee_x^{x+\frac{1-x}{t}} (f')_x \right) (1-x)^{-1}\,dt + \frac{1-x}{\sqrt{n}} \left(\bigvee_x^{x+\frac{1-x}{\sqrt{n}}} (f')_x \right) \\
&\leq \frac{\mu}{n+2} \frac{\varphi^2(x)}{1-x} \sum_{k=1}^{\sqrt{n}} \left(\bigvee_x^{x+\frac{1-x}{k}} (f')_x \right) + \frac{1-x}{\sqrt{n}} \left(\bigvee_x^{x+\frac{1-x}{\sqrt{n}}} (f')_x \right).
\end{aligned}
$$

Combining the estimates of $K_1 - K_8$, we get the desired result. Hence the proof follows. □

References

[1] A. M. Acu, V. Gupta and G. Tachev, Better numerical approximation by Durrmeyer type operators, Results Math. (2019) 74-90.

[2] M.M. Derriennic, Sur l'approximation de fonctions intégrables sur [0, 1] par des polynômes de Bernstein modifies, J. Approx. Theory, 31(1981), 325-343.
[3] Z. Ditzian and K. Ivanov, Bernstein-type operators and their derivatives, J. Approx. Theory 56(1) (1989), 72–90.
[4] Z. Ditzian and V. Totik, Moduli of Smoothness, Springer-Verlag, New York, (1987).
[5] J. L. Durrmeyer, Une formule d'inversion de la transformée de Laplace: applications à la théorie des moments, Thése de 3e cycle, Paris, 1967.
[6] H. H. Gonska, On approximation of continuously differentiable functions by positive linear operators, Bull. Austral. Math. Soc., Vol. 27 (1983), 73–81.
[7] H. H. Gonska and X. L. Zhou, A global inverse theorem on simultaneous approximation by Bernstein-Durrmeyer operators, J. Approx. Theory 67(3) (1991), 284–302.
[8] V. Gupta and R.P. Agarwal, Convergence Estimates In Approximation Theory, Springer, New York (2014).
[9] V. Gupta, A.-J. L.-Moreno and J.-M. L.-Palacios, On simultaneous approximation of the Bernstein Durrmeyer operators, Appl. Math. Comput. 213(1) (2009), 112-120.
[10] A. Kajla and T. Acar, Modified $\alpha-$Bernstein operators with better approximation properties, Ann. Funct. Anal. 10(4) (2019), 570–582.
[11] H. K.-Arab, M. Dehghan and M. R. Eslahchi, A new approach to improve the order of approximation of the Bernstein operators: Theory and applications, Numer. Algorithms 77(1) (2018), 111–150.
[12] A. Lupaş, Die Folge der Betaoperatoren, Dissertation, Universität Stuttgart, (1972).
[13] M. A. Özarslan and H. Aktuğlu, Local approximation for certain King type operators, Filomat 27(1) (2013), 173–181.
[14] H.M. Srivastava and V. Gupta, A certain family of summation-integral type operators, Math. Comput. Modelling, 37(2003), 1307–1315.