Grassmann angle formulas and identities

André L. G. Mandolesi *

December 31, 2021

Abstract

Grassmann angles improve upon similar concepts of angle between subspaces that measure volume contraction in orthogonal projections, working for real or complex subspaces, and being more efficient when dimensions are different. Their relations with contractions, inner and exterior products of multivectors are used to obtain formulas for computing these or similar angles in terms of arbitrary bases, and various identities for the angles with certain families of subspaces. These include generalizations of the Pythagorean trigonometric identity $\cos^2 \theta + \sin^2 \theta = 1$ for high dimensional and complex subspaces, which are connected to generalized Pythagorean theorems for volumes, quantum probabilities and Clifford geometric product.

Keywords: Grassmann angle, angle between subspaces, Pythagorean identity, Grassmann algebra, exterior algebra.

MSC: 15A75, 51M05

1 Introduction

Measuring the separation between subspaces is important in many areas, from geometry and linear algebra to statistics [10], operator perturbation theory [14], data mining [12], etc. In high dimensions, no single number fully describes this separation, and different concepts are used: gap, principal angles, minimal angle, Friedrichs angle, and others. An often used angle concept [5, 7, 9, 11] combines principal angles in a way that shows how volumes contract when orthogonally projected between the subspaces. Research on it has been focused on real spaces, but many applications require complex ones.

The relation between angle and volume contraction changes significantly in the complex case. Taking this into account, in [15] we defined a Grassmann angle that works well in both cases, with only a few adjustments. It is based on that same angle, but has a subtly important difference, which makes it easier to use with subspaces of distinct dimensions: it is asymmetric, reflecting the

^{*}Instituto de Matemática e Estatística, Universidade Federal da Bahia, Av. Adhemar de Barros s/n, 40170-110, Salvador - BA, Brazil. ORCID 0000-0002-5329-7034. E-mail: andre.mandolesi@ufba.br

dimensional asymmetry between the subspaces in a way that leads to better properties and more general results. This angle is intrinsically connected with Grassmann algebra, and has found interesting applications in the geometry of Grassmannians, Clifford algebra, quantum theory, etc.

In [16] we expressed contractions, inner, exterior and Clifford products of blades (simple multivectors) in terms of Grassmann angles. In this article we use the products to get formulas for the angles in terms of arbitrary bases, and identities relating the angles with the subspaces of an orthogonal partition, with coordinate subspaces of orthogonal bases, and others. Some of them generalize the Pythagorean trigonometric identity $\cos^2 \theta + \sin^2 \theta = 1$ for high dimensional and complex subspaces, being related to real and complex Pythagorean theorems for volumes [17] and quantum probabilities [18], and giving a geometric interpretation for an algebraic property of the Clifford product. Some of our results correspond, in the real case, to known ones, but our methods provide simpler proofs, while also extending them to the complex case.

Section 2 presents concepts and results which will be needed. We obtain formulas for computing Grassmann angles in section 3, generalized Pythagorean identities in section 4, and other useful identities in section 5.

2 Preliminaries

Here we review results we will use. See [15, 16] for proofs and more details.

In this article, X is a n-dimensional vector space over \mathbb{R} (real case) or \mathbb{C} (complex case), with inner product $\langle \cdot, \cdot \rangle$ (Hermitian product in the complex case, with conjugate-linearity in the first argument). For subspaces $V, W \subset X$, $\operatorname{Proj}_{W} : X \to W$ and $\operatorname{Proj}_{W}^{V} : V \to W$ are orthogonal projections. A *line* is a 1-dimensional subspace.

2.1 Grassmann algebra and partial orthogonality

In the Grassmann algebra ΛX [19, 24], a *p*-blade is a simple multivector $\nu = v_1 \wedge \ldots \wedge v_p \in \Lambda^p X$ of grade p, where $v_1, \ldots, v_p \in X$. If $\nu \neq 0$, it represents the *p*-dimensional subspace $V = \operatorname{span}(v_1, \ldots, v_p)$, and $\Lambda^p V = \operatorname{span}(\nu)$. A scalar $\nu \in \Lambda^0 X$ is a 0-blade, representing $\{0\}$.

The inner product of $\nu = v_1 \wedge \ldots \wedge v_p$ and $\omega = w_1 \wedge \ldots \wedge w_p$,

$$\langle \nu, \omega \rangle = \det (\langle v_i, w_j \rangle) = \begin{vmatrix} \langle v_1, w_1 \rangle & \cdots & \langle v_1, w_p \rangle \\ \vdots & \ddots & \vdots \\ \langle v_p, w_1 \rangle & \cdots & \langle v_p, w_p \rangle \end{vmatrix},$$

is extended linearly (sesquilinearly, in the complex case), with blades of distinct grades being orthogonal, and $\langle \nu, \omega \rangle = \bar{\nu}\omega$ for $\nu, \omega \in \Lambda^0 X$. In the real case, the norm $\|\nu\| = \sqrt{\langle \nu, \nu \rangle}$ gives the *p*-dimensional volume of the parallelotope spanned by v_1, \ldots, v_p . In the complex case, $\|\nu\|^2$ gives the 2*p*-dimensional volume of the parallelotope spanned by v_1, iv_1, \ldots, v_p , iv_p .

Given a subspace $W \subset X$, let $P = \operatorname{Proj}_W$. The orthogonal projection of $\nu = v_1 \wedge \ldots \wedge v_p$ on $\Lambda W \subset \Lambda X$ is $P\nu = Pv_1 \wedge \ldots \wedge Pv_p$.

Orthogonality in $\Lambda^p X$ corresponds to a weaker orthogonality concept in X.

Definition. For subspaces $V, W \subset X$, V is *partially orthogonal* to W ($V \perp W$) if there is a nonzero $v \in V$ such that $\langle v, w \rangle = 0$ for all $w \in W$.

Proposition 2.1. Let $V, W \subset X$ and $U \subset V$ be subspaces, with V represented by a blade ν , and $P = \operatorname{Proj}_W$. Then:

- i) $V \perp W \Leftrightarrow \dim P(V) < \dim V$.
- ii) If $V \perp W$ then $P\nu = 0$, otherwise $P\nu$ represents P(V).
- iii) If $V \not\perp W$ then $U \not\perp W$.

Proposition 2.2. Let $V, W \subset X$ be nonzero subspaces, and $p = \dim V$. Then $V \perp W \Leftrightarrow \Lambda^p V \perp \Lambda^p W$.

2.2 Coordinate decomposition

Definition. For integers $1 \le p \le q$, let

$$\mathcal{I}_{p}^{q} = \{(i_{1}, \dots, i_{p}) \in \mathbb{N}^{p} : 1 \leq i_{1} < \dots < i_{p} \leq q\}.$$

For any multi-index $I = (i_1, \ldots, i_p) \in \mathcal{I}_p^q$, we write $|I| = i_1 + \ldots + i_p$ and, if p < q, $\hat{I} = (1, \ldots, \hat{i_1}, \ldots, \hat{i_p}, \ldots, q) \in \mathcal{I}_{q-p}^q$, where each $\hat{i_k}$ indicates that index has been removed. Also, let $\mathcal{I}_0^q = \{0\}$, and for $I \in \mathcal{I}_0^q$ let |I| = 0 and $\hat{I} = (1, \ldots, q) \in \mathcal{I}_q^q$. For $I \in \mathcal{I}_q^q$ let $\hat{I} = 0 \in \mathcal{I}_0^q$.

Definition. Given a basis $\beta = (w_1, \ldots, w_q)$ of $W \subset X$, and $1 \leq p \leq q$, the coordinate *p*-subspaces for β are the $\binom{q}{p}$ subspaces

$$W_I = \operatorname{span}(w_{i_1}, \ldots, w_{i_n}),$$

for $I = (i_1, \ldots, i_p) \in \mathcal{I}_p^q$, which are represented by the *coordinate p-blades* of β ,

$$\omega_I = w_{i_1} \wedge \ldots \wedge w_{i_p} \in \Lambda^p W_I. \tag{1}$$

For $I \in \mathcal{I}_0^q$ we have the coordinate 0-subspace $W_I = \{0\}$ and $\omega_I = 1 \in \Lambda^0 W_I$.

When β is orthonormal, $\{\omega_I\}_{I \in \mathcal{I}_n^q}$ is an orthonormal basis of $\Lambda^p W$.

Definition. Given a decomposed nonzero blade $\omega = w_1 \wedge \ldots \wedge w_q \in \Lambda^q X$, take the basis $\beta = (w_1, \ldots, w_q)$ of its subspace. For any $0 \leq p \leq q$ and $I \in \mathcal{I}_p^q$, the coordinate decomposition of ω (w.r.t. I and β) is

$$\omega = \sigma_I \,\omega_I \wedge \omega_{\hat{I}},\tag{2}$$

with ω_I and $\omega_{\hat{I}}$ as in (1), and $\sigma_I = (-1)^{|I| + \frac{p(p+1)}{2}}$.

2.3 Principal angles and vectors

Definition. The Euclidean angle $\theta_{v,w} \in [0,\pi]$ between nonzero vectors $v, w \in X$ is given by $\cos \theta_{v,w} = \frac{\operatorname{Re}\langle v,w \rangle}{\|v\| \|w\|}$. In the complex case, there is also a Hermitian angle $\gamma_{v,w} \in [0,\frac{\pi}{2}]$ defined by $\cos \gamma_{v,w} = \frac{|\langle v,w \rangle|}{\|v\| \|w\|}$.

The Hermitian angle is the Euclidean angle between v and $\operatorname{span}_{\mathbb{C}}(w)$.

A list of principal angles [4, 13] is necessary to fully describe the relative position of high dimensional subspaces.

Definition. Let $V, W \subset X$ be nonzero subspaces, $p = \dim V$, $q = \dim W$ and $m = \min\{p, q\}$. Orthonormal bases (e_1, \ldots, e_p) of V and (f_1, \ldots, f_q) of W are associated principal bases, formed by principal vectors, with principal angles $0 \le \theta_1 \le \ldots \le \theta_m \le \frac{\pi}{2}$, if

$$\langle e_i, f_j \rangle = \delta_{ij} \cos \theta_i.$$
 (3)

A singular value decomposition [4, 6] gives such bases: for $P = \operatorname{Proj}_{W}^{V}$, the e_i 's and f_i 's are orthonormal eigenvectors of P^*P and PP^* , respectively, and the $\cos \theta_i$'s are square roots of the eigenvalues of P^*P , if $p \leq q$, or PP^* otherwise. The θ_i 's are uniquely defined, but the e_i 's and f_i 's are not. P is given in principal bases by a $q \times p$ diagonal matrix formed with the $\cos \theta_i$'s, as

$$Pe_{i} = \begin{cases} f_{i} \cdot \cos \theta_{i} & \text{if } 1 \leq i \leq m, \\ 0 & \text{if } i > m. \end{cases}$$

$$\tag{4}$$

A geometric interpretation of principal angles is that the unit sphere of V projects to an ellipsoid in W with semi-axes of lengths $\cos \theta_i$, for $1 \le i \le m$ (in the complex case there are 2 semi-axes for each i). They can also be described recursively with a minimization condition: e_1 and f_1 form the smallest angle θ_1 between nonzero vectors of V and W; in their orthogonal complements we obtain e_2 , f_2 and θ_2 in the same way; and so on.

2.4 Grassmann angles and blade products

Grassmann angles were introduced in [15], and related to blade products in [16].

Definition. Let $V, W \subset X$ be subspaces, ν be a nonzero blade representing V, and $P = \operatorname{Proj}_W$. The *Grassmann angle* $\Theta_{V,W} \in [0, \frac{\pi}{2}]$ is given by

$$\cos \Theta_{V,W} = \frac{\|P\nu\|}{\|\nu\|}.$$
(5)

As blade norms (squared, in the complex case) give volumes, $\Theta_{V,W}$ tells us how volumes in V contract when orthogonally projected on W. In simple cases, where there is an unambiguous concept of angle between subspaces, $\Theta_{V,W}$ coincides with it, as when V is a line, or V and W are planes in \mathbb{R}^3 .

The main difference between Grassmann angles and similar ones [5, 7, 9, 11] is its asymmetry: $\Theta_{V,W} = \frac{\pi}{2}$ if dim $V > \dim W$, so, in general, $\Theta_{V,W} \neq \Theta_{W,V}$

when dimensions are different. It reflects the dimensional asymmetry between the subspaces, leading to better and more general results. For example, the contraction and exterior product formulas in proposition 2.6, or theorems 3.5 and 3.7, only hold without any restrictions thanks to this asymmetry.

Grassmann angles work well in complex spaces, and this makes them useful for applications in areas like quantum theory. Differences due to volumes being given, in this case, by squared norms, have important implications [17, 18].

These angles have many useful properties, some of which are listed below.

Proposition 2.3. Let $V, W \subset X$ be subspaces, with principal angles $\theta_1, \ldots, \theta_m$, where $m = \min\{p, q\}$ for $p = \dim V$ and $q = \dim W$, and let $P = \operatorname{Proj}_W^V$.

- i) $\Theta_{V,W} = \frac{\pi}{2} \iff V \perp W.$
- ii) If $V = \operatorname{span}(v)$ and $W = \operatorname{span}(w)$ for nonzero $v, w \in X$ then $\Theta_{V,W} = \min\{\theta_{v,w}, \pi \theta_{v,w}\}$ in the real case, and $\Theta_{V,W} = \gamma_{v,w}$ in the complex one.
- *iii)* If V is a line and $v \in V$ then $||Pv|| = ||v|| \cos \Theta_{V,W}$.
- iv) vol $P(S) = \text{vol } S \cdot \cos \Theta_{V,W}$ (cos² $\Theta_{V,W}$ in complex case), where $S \subset V$ is a parallelotope and vol is the p-dimensional volume (2p in complex case).
- v) $\cos^2 \Theta_{V,W} = \det(\bar{\mathbf{P}}^T \mathbf{P})$, where \mathbf{P} is a matrix for P in orthonormal bases of V and W.
- vi) If p > q then $\Theta_{V,W} = \frac{\pi}{2}$, otherwise $\cos \Theta_{V,W} = \prod_{i=1}^{m} \cos \theta_i$.
- vii) If p = q then $\Theta_{V,W} = \Theta_{W,V}$.
- viii) $\Theta_{T(V),T(W)} = \Theta_{V,W}$ for any orthogonal (unitary, in the complex case) transformation $T: X \to X$.
- ix) If V' and W' are the orthogonal complements of $V \cap W$ in V and W, respectively, then $\Theta_{V,W} = \Theta_{V',W'}$.
- x) $\Theta_{V,W} = \Theta_{W^{\perp},V^{\perp}}$.

The Grassmann angle with an orthogonal complement has extra properties which grant it a special name and notation.

Definition. The complementary Grassmann angle $\Theta_{V,W}^{\perp} \in [0, \frac{\pi}{2}]$ of subspaces $V, W \subset X$ is $\Theta_{V,W}^{\perp} = \Theta_{V,W^{\perp}}$.

In general, this is not the usual complement, i.e. $\Theta_{V,W}^{\perp} \neq \frac{\pi}{2} - \Theta_{V,W}$.

Proposition 2.4. Let $V, W \subset X$ be subspaces.

- $i) \ \Theta_{V,W}^{\perp} = 0 \ \Leftrightarrow \ V \perp W.$
- *ii)* $\Theta_{VW}^{\perp} = \frac{\pi}{2} \Leftrightarrow V \cap W \neq \{0\}.$
- iii) If $\theta_1, \ldots, \theta_m$ are the principal angles then $\cos \Theta_{V,W}^{\perp} = \prod_{i=1}^m \sin \theta_i$.

- iv) If V is a line then $\Theta_{V,W}^{\perp} = \frac{\pi}{2} \Theta_{V,W}$.
- v) $\Theta_{V,W}^{\perp} = \Theta_{W,V}^{\perp}$.

In section 3 we get a simpler way to prove (iii) than the one in [15].

Definition. The *(left) contraction* of $\nu \in \Lambda^p X$ on $\omega \in \Lambda^q X$ is the unique $\nu \lrcorner \omega \in \Lambda^{q-p} X$ such that, for all $\mu \in \Lambda^{q-p} X$,

$$\langle \mu, \nu \lrcorner \, \omega \rangle = \langle \nu \land \mu, \omega \rangle. \tag{6}$$

This contraction coincides with the inner product when p = q, and it is asymmetric, with $\nu \lrcorner \omega = 0$ if p > q. It differs from the one used in Clifford geometric algebra [3] by a reversion.

Proposition 2.5. For $\nu \in \Lambda^p X$ and any blade $\omega \in \Lambda^q X$, with $p \leq q$,

$$\nu \lrcorner \, \omega = \sum_{I \in \mathcal{I}_p^q} \sigma_I \, \langle \nu, \omega_I \rangle \, \omega_{\hat{I}}, \tag{7}$$

where σ_I , ω_I and $\omega_{\hat{I}}$ are as in (2) for any decomposition $\omega = w_1 \wedge \ldots \wedge w_q$.

Proposition 2.6. Let $\nu, \omega \in \Lambda X$ be blades representing $V, W \subset X$, respectively.

- i) $|\langle \nu, \omega \rangle| = \|\nu\| \|\omega\| \cos \Theta_{V,W}$, if ν and ω have equal grades.
- *ii)* $\|\nu \lrcorner \omega\| = \|\nu\| \|\omega\| \cos \Theta_{V,W}$.
- *iii)* $\|\nu \wedge \omega\| = \|\nu\| \|\omega\| \cos \Theta_{VW}^{\perp}$.

With a Grassmann angle $\Theta_{V,W}$ for oriented subspaces (with orientations of $\nu, \omega \in \Lambda^p X$) given by $\cos \Theta_{V,W} = \frac{\langle \nu, \omega \rangle}{|\langle \nu, \omega \rangle|} \cos \Theta_{V,W}$ if $\langle \nu, \omega \rangle \neq 0$, otherwise $\Theta_{V,W} = \frac{\pi}{2}$, we also have

$$\langle \nu, \omega \rangle = \|\nu\| \|\omega\| \cos \Theta_{V,W}. \tag{8}$$

3 Formulas for Grassmann angles

Here we give formulas for computing Grassmann and complementary Grassmann angles in terms of arbitrary bases, thus generalizing proposition 2.3v. They can be adapted for use with similar angles [5, 7, 9, 11], with some dimensional restrictions (dim $V \leq \dim W$ in theorem 3.5, dim $V \leq \dim W^{\perp}$ in theorem 3.7) as these other angles are symmetric.

The first formula, for subspaces of same dimension, follows from proposition 2.6*i*. A similar result, for the real case, appears in [5].

Theorem 3.1. Given bases (v_1, \ldots, v_p) and (w_1, \ldots, w_p) of p-dimensional subspaces $V, W \subset X$, let $A = (\langle w_i, w_j \rangle), B = (\langle w_i, v_j \rangle)$ and $D = (\langle v_i, v_j \rangle)$. Then

$$\cos^2 \Theta_{V,W} = \frac{|\det B|^2}{\det A \cdot \det D}$$

Example 3.2. In \mathbb{C}^3 , let $V = \operatorname{span}(v_1, v_2)$ and $W = \operatorname{span}(w_1, w_2)$ with $v_1 = (1, -\xi, 0), v_2 = (0, \xi, -\xi^2), w_1 = (1, 0, 0)$ and $w_2 = (0, \xi, 0),$ where $\xi = e^{i\frac{2\pi}{3}}$. The theorem gives $\Theta_{V,W} = \operatorname{arccos} \frac{\sqrt{3}}{3}$. Since $v = (\xi, \xi^2, -2) \in V$ and $w = (1, \xi, 0) \in W$ are orthogonal to $V \cap W = \operatorname{span}(v_1)$, proposition 2.3*ii* and *ix* give $\Theta_{V,W} = \Theta_{\operatorname{span}(v),\operatorname{span}(w)} = \gamma_{v,w}$, and the Hermitian angle formula confirms the result.

The next formulas require some determinant identities.

Proposition 3.3 (Laplace's Expansion [22, p.80]). Given a $q \times q$ matrix M and a multi-index $J \in \mathcal{I}_p^q$, with $1 \leq p < q$,

$$\det M = \sum_{I \in \mathcal{I}_p^q} (-1)^{|I| + |J|} \det M_{I,J} \cdot \det M_{\hat{I},\hat{J}},$$

where $M_{I,J}$ is the $p \times p$ submatrix formed by entries with row indices in I and column indices in J, and $M_{\hat{I},\hat{J}}$ is the $(q-p) \times (q-p)$ submatrix formed by entries with row indices not in I and column indices not in J.

Proposition 3.4 (Schur's determinant identity [1]). Let $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ be a $(q + p) \times (q + p)$ matrix, partitioned into $q \times q$, $q \times p$, $p \times q$ and $p \times p$ matrices A, B, C and D, respectively. If A is invertible then

$$\det M = \det A \cdot \det(D - CA^{-1}B).$$
(9)

Likewise, if D is invertible then

$$\det M = \det D \cdot \det(A - BD^{-1}C).$$
⁽¹⁰⁾

Proof. Follows by decomposing M into block triangular matrices, as $M = \begin{pmatrix} A & 0_{q \times p} \\ C & \mathbb{1}_{p \times p} \end{pmatrix} \begin{pmatrix} \mathbb{1}_{q \times q} & A^{-1}B \\ 0_{p \times q} & D - CA^{-1}B \end{pmatrix}$ or $M = \begin{pmatrix} A - BD^{-1}C & BD^{-1} \\ 0_{p \times q} & \mathbb{1}_{p \times p} \end{pmatrix} \begin{pmatrix} \mathbb{1}_{q \times q} & 0_{q \times p} \\ C & D \end{pmatrix}$.

We now get a formula for distinct dimensions, simpler than one given in [7] for a similar angle (which corrects another formula from [23]).

Theorem 3.5. Given bases (v_1, \ldots, v_p) of V and (w_1, \ldots, w_q) of W, let $A = (\langle w_i, w_j \rangle), B = (\langle w_i, v_j \rangle), and D = (\langle v_i, v_j \rangle)$. Then

$$\cos^2 \Theta_{V,W} = \frac{\det(\bar{B}^T A^{-1} B)}{\det D}$$

Proof. If p > q then $\Theta_{V,W} = \frac{\pi}{2}$, and the determinant of $\bar{B}^T A^{-1} B$ vanishes as it is a $p \times p$ matrix with rank at most q.

If $p \leq q$, applying proposition 3.3, with J = (q + 1, ..., q + p), to the $(q+p) \times (q+p)$ block matrix $M = \begin{pmatrix} A & B \\ \bar{B}^T & 0_{p \times p} \end{pmatrix}$, we get

$$\det M = \sum_{I \in \mathcal{I}_p^p} (-1)^{|I| + pq + \frac{p(p+1)}{2}} \cdot \det B_I \cdot \det N_{\hat{I}},$$

where B_I is the $p \times p$ submatrix of M formed by lines of B with indices in I, and $N_{\hat{I}} = \begin{pmatrix} A_{\hat{I}} \\ \bar{B}^T \end{pmatrix}$ is its $q \times q$ complementary submatrix, formed by lines of Awith indices not in I and all of \bar{B}^T .

For $\nu = v_1 \wedge \ldots \wedge v_p$ and $\omega = w_1 \wedge \ldots \wedge w_q$ we have, by (6) and (7),

$$\|\nu_{\lrcorner}\omega\|^{2} = \langle \nu \wedge (\nu_{\lrcorner}\omega), \omega \rangle = \sum_{I \in \mathcal{I}_{p}^{q}} \sigma_{I} \langle \omega_{I}, \nu \rangle \langle \nu \wedge \omega_{\hat{I}}, \omega \rangle.$$

Since det $B_I = \langle \omega_I, \nu \rangle$, det $N_{\hat{I}} = \langle \omega_{\hat{I}} \wedge \nu, \omega \rangle = (-1)^{pq+p} \langle \nu \wedge \omega_{\hat{I}}, \omega \rangle$ and $\sigma_I = (-1)^{|I| + \frac{p(p+1)}{2}}$, we obtain $\|\nu \lrcorner \omega\|^2 = (-1)^p$ det M. Proposition 2.6*ii* then gives $\cos^2 \Theta_{V,W} = \frac{(-1)^p \det M}{\det D \det A}$, and the result follows from (9).

Example 3.6. In \mathbb{R}^4 , let v = (1, 0, 1, 0), $w_1 = (0, 1, 1, 0)$, $w_2 = (1, 2, 2, -1)$, $V = \operatorname{span}(v)$ and $W = \operatorname{span}(w_1, w_2)$. Then $A = \begin{pmatrix} 2 & 4 \\ 4 & 10 \end{pmatrix}$, $B = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$ and D = (2), and the theorem gives $\Theta_{V,W} = 45^\circ$, as one can verify by projecting v on W. Switching the roles of V and W, we now have A = (2), $B = (1 \ 3)$, $D = \begin{pmatrix} 2 & 4 \\ 4 & 10 \end{pmatrix}$ and $\Theta_{W,V} = 90^\circ$, which is correct since dim $W > \dim V$.

We now get formulas for the complementary Grassmann angle.

Theorem 3.7. Given bases (v_1, \ldots, v_p) of V and (w_1, \ldots, w_q) of W, let $A = (\langle w_i, w_j \rangle), B = (\langle w_i, v_j \rangle), and D = (\langle v_i, v_j \rangle)$. Then

$$\cos^2 \Theta_{V,W}^{\perp} = \frac{\det(A - BD^{-1}\bar{B}^T)}{\det A}.$$
(11)

Proof. Let $\nu = v_1 \wedge \ldots \wedge v_p$ and $\omega = w_1 \wedge \ldots \wedge w_q$. The result is obtained applying proposition 2.6*iii* to $\omega \wedge \nu$, and using (10) with $M = \begin{pmatrix} A & B \\ B^T & D \end{pmatrix}$.

Corollary 3.8. If **P** is a matrix representing Proj_W^V in orthonormal bases of V and W then

$$\cos^2 \Theta_{V,W}^{\perp} = \det(\mathbb{1}_{q \times q} - \mathbf{P}\bar{\mathbf{P}}^T).$$
(12)

This gives an easy way to prove proposition 2.4*iii*, as in principal bases **P** is a diagonal matrix formed with the $\cos \theta_i$'s.

Example 3.9. In example 3.6, (11) gives $\Theta_{V,W}^{\perp} = 45^{\circ}$, in agreement with proposition 2.4*iv*. The same formula also gives $\Theta_{W,V}^{\perp} = 45^{\circ}$, as expected by proposition 2.4*v*. Direct calculations show the principal angles of V and W^{\perp} are 0° and 45°, as are those of W and V^{\perp} , confirming the results.

Example 3.10. In example 3.2, using (11) with the bases (v_1, v_2) and (w_1, w_2) , or (12) with the orthonormal bases $(\frac{v_1}{\sqrt{2}}, \frac{v}{\sqrt{6}})$ and $(\frac{v_1}{\sqrt{2}}, \frac{w}{\sqrt{2}})$, we get $\Theta_{V,W}^{\perp} = 90^{\circ}$, as expected by proposition 2.4*ii*, since $V \cap W \neq \{0\}$.




Figure 1: Pythagorean identities for subspaces of equal dimensions.

4 Generalized Pythagorean identities

The Pythagorean trigonometric identity $\cos^2 \theta + \sin^2 \theta = 1$ can be written as $\cos^2 \theta_x + \cos^2 \theta_y = 1$, with θ_x and θ_y being angles a line in \mathbb{R}^2 makes with the axes. We give generalizations for Grassmann angles which, with proposition 2.3*iv*, lead to real and complex Pythagorean theorems for volumes [17]. Some correspond to known results in real spaces, which are now extended to the complex case, with important implications for quantum theory [18]. We also get a geometric interpretation for a property of the Clifford product.

The first identity relates the Grassmann angles of a (real or complex) line with all subspaces of an orthogonal partition of X.

Theorem 4.1. Given an orthogonal partition $X = W_1 \oplus \cdots \oplus W_k$ and a line $L \subset X$,

$$\sum_{i=1}^{k} \cos^2 \Theta_{L,W_i} = 1.$$
 (13)

Proof. Given a nonzero $v \in L$, as $||v||^2 = \sum_i ||\operatorname{Proj}_{W_i} v||^2$ the result follows from proposition 2.3*iii*.

Example 4.2. If θ_x , θ_y and θ_z are the angles between a line in \mathbb{R}^3 and the axes (fig. 1a), then $\cos^2 \theta_x + \cos^2 \theta_y + \cos^2 \theta_z = 1$.

This is a known identity for direction cosines and, like other examples we give, is only meant to illustrate the theorem in \mathbb{R}^3 . The relevance of our result lies mainly in the complex case, where it has important connections with quantum theory.

Example 4.3. If X is the complex Hilbert space of a quantum system, L is the complex line of a quantum state vector ψ , and the W_i 's are the eigenspaces of

a quantum observable, the probability of getting result *i* when measuring ψ is $p_i = \|\operatorname{Proj}_{W_i}\psi\|^2/\|\psi\|^2$ [2]. So, by proposition 2.3*iii*, $p_i = \cos^2 \Theta_{L,W_i}$, and (13) reflects the fact that the total probability is 1.

By proposition 2.3iv, $\cos^2 \Theta_{L,W_i}$ measures area contraction in orthogonal projections of the complex line L. This is explored in [18] to get a new interpretation for quantum probabilities and derive the Born rule.

The next identities relate Grassmann angles of a (real or complex) subspace with coordinate subspaces of an orthogonal basis of X.

Theorem 4.4. If $V \subset X$ is a p-dimensional subspace, and $n = \dim X$,

$$\sum_{I \in \mathcal{I}_p^n} \cos^2 \Theta_{V, W_I} = 1$$

where the W_I 's are the coordinate p-subspaces of an orthogonal basis of X.

Proof. Without loss of generality, assume the basis is orthonormal, so its coordinate *p*-blades ω_I form an orthonormal basis of $\Lambda^p X$. For an unit blade $\nu \in \Lambda^p V \subset \Lambda^p X$ we have $\sum_I |\langle \nu, \omega_I \rangle|^2 = 1$, and the result follows from proposition 2.6*i*.

A similar result for the real case, in terms of products of cosines of principal angles, appears in [20]. The theorem extends to affine subspaces, as in the next example, which is a dual of example 4.2 via proposition 2.3x.

Example 4.5. If θ_{xy} , θ_{xz} and θ_{yz} are the angles a plane in \mathbb{R}^3 makes with the coordinate planes (fig. 1b) then $\cos^2 \theta_{xy} + \cos^2 \theta_{xz} + \cos^2 \theta_{yz} = 1$.

Example 4.6. Let ξ, w_1, w_2 and V be as in example 3.2, $w_3 = (0, 0, \xi^2)$ and $W_{ij} = \operatorname{span}(w_i, w_j)$. As the unitary transformation given by $T = \begin{pmatrix} 0 & 0 & \xi \\ \xi & 0 & 0 \\ 0 & \xi & 0 \end{pmatrix}$ maps $W_{12} \mapsto W_{23}, W_{23} \mapsto W_{13}$, and preserves V, proposition 2.3*viii* gives $\Theta_{V,W_{12}} = \Theta_{V,W_{23}} = \Theta_{V,W_{13}}$. Since W_{12}, W_{13} and W_{23} are the coordinate 2-subspaces of the orthogonal basis (w_1, w_2, w_3) of \mathbb{C}^3 , theorem 4.4 gives $\cos \Theta_{V,W_{ij}} = \frac{\sqrt{3}}{3}$, in agreement with that example.

A formula relating the geometric product of Clifford algebra [8] to Grassmann angles, given in [16], implies $||AB||^2 = ||A||^2 ||B||^2 \sum_J \cos^2 \Theta_{V,Y_J}$, where A and B are p-blades representing subspaces V and W, and the Y_J 's are all coordinate p-subspaces of a certain orthogonal basis of Y = V + W. This gives a geometric interpretation for a simple yet important property of this product. Those in proposition 2.6 are submultiplicative for blades because they correspond to projections on single subspaces. The geometric product, on the other hand, involves projections on all Y_J 's, and with theorem 4.4 we see this is what allows ||AB|| = ||A|| ||B||.

Extending a result of [21], we also have identities for Grassmann angles with coordinate subspaces of a dimension different from V.

Theorem 4.7. Let $V \subset X$ be a p-dimensional subspace, $0 \le q \le n = \dim X$, and the W_I 's be the coordinate q-subspaces of an orthogonal basis of X.

i) If
$$p \le q$$
 then $\sum_{I \in \mathcal{I}_q^n} \cos^2 \Theta_{V,W_I} = \binom{n-p}{n-q}$

ii) If
$$p > q$$
 then $\sum_{I \in \mathcal{I}_q^n} \cos^2 \Theta_{W_I,V} = \binom{p}{q}$

Proof. We can assume $p, q \neq 0$ and that the basis is orthonormal. So, for $0 \leq r \leq n$ and with ω_I 's as in (1), $\{\omega_I\}_{I \in \mathcal{I}_r^n}$ and $\{\omega_{\hat{I}}\}_{I \in \mathcal{I}_r^n}$ are orthonormal bases of $\Lambda^r X$ and $\Lambda^{n-r} X$, respectively.

(i) For an unit blade $\nu \in \Lambda^p V$ and $I = (i_1, \ldots, i_q) \in \mathcal{I}_q^n$ we have, by (7),

$$u \lrcorner \omega_{I} = \sum_{J \in \mathcal{I}_{p}^{q}} \sigma_{J} \langle \nu, (\omega_{I})_{J} \rangle (\omega_{I})_{\hat{J}},$$

where $(\omega_I)_J = w_{i_{j_1}} \wedge \ldots \wedge w_{i_{j_p}}$ for $J = (j_1, \ldots, j_p)$, and likewise for $(\omega_I)_{\hat{J}}$. As the $(\omega_I)_{\hat{J}}$'s are orthonormal, proposition 2.6*ii* gives

$$\cos^2 \Theta_{V,W_I} = \|\nu \lrcorner \omega_I\|^2 = \sum_{J \in \mathcal{I}_p^q} |\langle \nu, (\omega_I)_J \rangle|^2,$$

and therefore

$$\sum_{I \in \mathcal{I}_q^n} \cos^2 \Theta_{V,W_I} = \sum_{I \in \mathcal{I}_q^n} \sum_{J \in \mathcal{I}_p^q} |\langle \nu, (\omega_I)_J \rangle|^2$$
$$= \frac{\binom{n}{q}\binom{q}{p}}{\binom{n}{p}} \sum_{K \in \mathcal{I}_p^n} |\langle \nu, \omega_K \rangle|^2 = \binom{n-p}{n-q} ||\nu||^2,$$

where the binomial coefficients account for the number of times each ω_K appears as a $(\omega_I)_J$ in the double summation.

(*ii*) For each $I \in \mathcal{I}_q^n$, proposition 2.3*x* gives $\Theta_{W_I,V} = \Theta_{V^{\perp},W_I^{\perp}}$. As $W_I^{\perp} = W_{\hat{I}}$ for $\hat{I} \in \mathcal{I}_{n-q}^n$, and dim $V^{\perp} = n - p < n - q = \dim W_I^{\perp}$, the result follows from the previous case.

The following examples are again duals of each other.

Example 4.8. If θ_{xy}, θ_{xz} and θ_{yz} are the angles a line in \mathbb{R}^3 makes with the coordinate planes (fig. 2a) then $\cos^2 \theta_{xy} + \cos^2 \theta_{xz} + \cos^2 \theta_{yz} = 2$.

Example 4.9. If θ_x, θ_y and θ_z are the angles between the axes and a plane in \mathbb{R}^3 (fig. 2b) then $\cos^2 \theta_x + \cos^2 \theta_y + \cos^2 \theta_z = 2$.




Figure 2: Pythagorean identities for subspaces of different dimensions.

5 Other identities

Using the Grassmann angle $\Theta_{V,W}$ for oriented subspaces, we have:

Theorem 5.1. Given p-dimensional oriented subspaces $V, W \subset X$,

$$\cos \Theta_{V,W} = \sum_{I \in \mathcal{I}_p^n} \cos \Theta_{V,X_I} \cos \Theta_{W,X_I},$$

where the X_I 's are the coordinate p-subspaces of an orthogonal basis of X, with orientations given by the corresponding coordinate p-blades.

Proof. The result follows by decomposing unit blades $\nu \in \Lambda^p V$ and $\omega \in \Lambda^p W$ (with the orientations of V and W) in the orthonormal basis of $\Lambda^p X$ formed with the normalized coordinate p-blades, and applying (8).

This gives an inequality for $\cos \Theta_{V,W} = |\cos \Theta_{V,W}|$, like one from [21].

Corollary 5.2. $\cos \Theta_{V,W} \leq \sum_{I} \cos \Theta_{V,X_{I}} \cos \Theta_{W,X_{I}}$.

Example 5.3. $\cos \theta = \cos \alpha_x \cos \beta_x + \cos \alpha_y \cos \beta_y + \cos \alpha_z \cos \beta_z$ for the angle $\theta \in [0, \pi]$ between 2 oriented lines in \mathbb{R}^3 forming angles $\alpha_x, \alpha_y, \alpha_z$ and $\beta_x, \beta_y, \beta_z$ (all in $[0, \pi]$) with the positive axes.

The last identities will require some preparation.

Definition. A coordinate subspace of a principal basis β of V w.r.t. W is a principal subspace¹ (of V w.r.t. W, for β). Two or more subspaces of V are coprincipal (w.r.t. W) if they are principal for the same β .

Note that a subspace being principal depends on both V and W, even if they are left implicit. Also, $\{0\}$ is always principal.

¹Some authors use 'principal subspace' for span (e_i, f_i) , where $e_i \in V$ and $f_i \in W$ are principal vectors corresponding to the same principal angle θ_i .

Lemma 5.4. Let $V, W \subset X$ be nonzero subspaces and $U \subset V$ be any subspace. If $U \perp W$ then U is principal w.r.t. W.

Proof. By (3), the union of an orthonormal basis of U and a principal basis of $U^{\perp} \cap V$ w.r.t. W gives a principal basis of V w.r.t. W.

Lemma 5.5. Let $V, W \subset X$ be nonzero subspaces, with associated principal bases β_V and β_W , $U \subset V$ be any subspace, and $P = \operatorname{Proj}_W$. Then:

- i) U is principal for $\beta_V \Leftrightarrow U^{\perp} \cap V$ is principal for β_V .
- ii) U is principal for $\beta_V \Rightarrow P(U)$ is principal for β_W . The converse holds if $V \not \perp W$.

Proof. (i) If U is spanned by some vectors of the orthogonal basis β_V , $U^{\perp} \cap V$ is spanned by the others. (ii) Follows from (4), and the converse from proposition 2.1 as well.

Proposition 5.6. Let $V, W \subset X$ be nonzero subspaces, $U \subset V$ be any subspace and $P = \operatorname{Proj}_W$. Then U is principal $\Leftrightarrow P(U) \perp P(U^{\perp} \cap V)$.

Proof. (\Rightarrow) Follows from (4). (\Leftarrow) By lemmas 5.4 and 5.5*i*, we can assume U, $U^{\perp} \cap V$, P(U) and $P(U^{\perp} \cap V)$ are not {0}. As $P(U) \perp P(U^{\perp} \cap V)$ implies $U \perp P(U^{\perp} \cap V)$ and $P(U) \perp U^{\perp} \cap V$, from associated principal bases of U and P(U), and of $U^{\perp} \cap V$ and $P(U^{\perp} \cap V)$, we form principal bases for V and P(V).

Proposition 5.7. Let $V, W \subset X$ be nonzero subspaces, $V_1, V_2 \subset V$ be distinct r-dimensional coprincipal subspaces w.r.t. W, and $P = \operatorname{Proj}_W$. Then:

- i) $V_1 \perp V_2$.
- ii) If $V_1 \not\perp W$ then $P(V_1) \perp P(V_2)$.
- *iii)* $\langle \nu_1, \nu_2 \rangle = 0$ and $\langle P\nu_1, P\nu_2 \rangle = 0$ for any $\nu_1 \in \Lambda^r V_1$, $\nu_2 \in \Lambda^r V_2$.

Proof. (i) V_1 has an element e of the principal basis which V_2 does not. (ii) If $V_1 \not\perp W$ then $Pe \neq 0$, and by proposition 5.6 $P(\text{span}(e)) \perp P(V_2)$. (iii) By proposition 2.2, $\Lambda^r V_1 \perp \Lambda^r V_2$ and, if $V_1 \not\perp W$, $\Lambda^r P(V_1) \perp \Lambda^r P(V_2)$. If $V_1 \not\perp W$ then $P\nu_1 = 0$, by proposition 2.1*ii*.

Theorem 5.8. Given nonzero subspaces $V, W \subset X$ and $U \subset V$, let $r = \dim U$ and $p = \dim V$. Then

$$\cos^2 \Theta_{U,W} = \sum_{I \in \mathcal{I}_r^p} \cos^2 \Theta_{U,V_I} \cdot \cos^2 \Theta_{V_I,W},$$

where the V_I 's are coordinate r-subspaces of a principal basis β of V w.r.t. W.




Figure 3: $\cos^2 \beta = \cos^2 \alpha + \sin^2 \alpha \cos^2 \theta$

Proof. The coordinate r-blades $\nu_I \in \Lambda^r V_I$ of β form an orthonormal basis of $\Lambda^r V$, so $P\mu = \sum_I \langle \nu_I, \mu \rangle P\nu_I$ for an unit $\mu \in \Lambda^r U$ and $P = \operatorname{Proj}_W$. By proposition 5.7 the $P\nu_I$'s are mutually orthogonal, so $\|P\mu\|^2 = \sum_I |\langle \mu, \nu_I \rangle|^2 \|P\nu_I\|^2$. The result follows from proposition 2.6*i* and (5).

By theorem 4.4 $\sum_{I \in \mathcal{I}_r^p} \cos^2 \Theta_{U,V_I} = 1$, so this result means $\cos^2 \Theta_{U,W}$ is a weighted average of the $\cos^2 \Theta_{V_I,W}$'s, with weights given by the $\cos^2 \Theta_{U,V_I}$'s.

Example 5.9. Given planes $V, W \subset \mathbb{R}^3$ and a line $U \subset V$, let $\alpha = \Theta_{U,V \cap W}$, $\beta = \Theta_{U,W}$ and $\theta = \Theta_{V,W}$ (fig. 3). As $V \cap W$ and $(V \cap W)^{\perp} \cap V$ are principal lines of V w.r.t. W, we have $\cos^2 \beta = \cos^2 \alpha \cdot 1 + \sin^2 \alpha \cdot \cos^2 \theta$.

Definition. A partition $V = \bigoplus_i V_i$ is *principal* w.r.t. W if the V_i 's are coprincipal subspaces of V w.r.t. W.

Any principal partition is an orthogonal partition. Note that some subspaces of a partition can be $\{0\}$.

Proposition 5.10. Let $V, W \subset X$ be nonzero subspaces, $P = \operatorname{Proj}_W$, and $V = \bigoplus_i V_i$ be an orthogonal partition. The following are equivalent:

- i) $V = \bigoplus_i V_i$ is a principal partition w.r.t. W.
- ii) $P(V) = \bigoplus_i P(V_i)$ is a principal partition w.r.t. V.
- iii) $P(V) = \bigoplus_i P(V_i)$ is an orthogonal partition.

Proof. $(i \Rightarrow ii)$ The $P(V_i)$'s are pairwise disjoint by proposition 5.6, and coprincipal by lemma 5.5*ii.* $(ii \Rightarrow iii)$ Immediate. $(iii \Rightarrow i)$ As the $P(V_i)$'s are mutually orthogonal, $V_i \perp P(V_j)$ if $i \neq j$. As the V_i 's are also mutually orthogonal, by (3) the union of principal bases of theirs w.r.t. W gives a principal basis of V. \Box

In [15] we got $\cos \Theta_{V,W} = \prod_i \cos \Theta_{V_i,W}$ for a principal partition $V = \bigoplus_i V_i$. We now generalize this for orthogonal partitions and get a partial converse.

Theorem 5.11. Let $V_1, V_2, W \subset X$ be subspaces and $P = \operatorname{Proj}_W$. If $V_1 \perp V_2$,

 $\cos \Theta_{V_1 \oplus V_2, W} = \cos \Theta_{V_1, W} \cdot \cos \Theta_{V_2, W} \cdot \cos \Theta_{P(V_1), P(V_2)}^{\perp}.$

Proof. By propositions 2.1*iii* and 2.3*i*, we can assume $V_1 \not\perp W$ and $V_2 \not\perp W$. As $V_1 \perp V_2$, if ν_1 and ν_2 are unit blades representing them, $\nu_1 \wedge \nu_2$ is an unit blade representing $V_1 \oplus V_2$. By (5), $\cos \Theta_{V_1 \oplus V_2, W} = \|P(\nu_1 \wedge \nu_2)\| = \|(P\nu_1) \wedge (P\nu_2)\|$, and the result follows from propositions 2.1*ii* and 2.6*iii*.

Corollary 5.12. Let $V, W \subset X$ be subspaces and $P = \operatorname{Proj}_W$. For an orthogonal partition $V = \bigoplus_{i=1}^k V_i$,

$$\cos \Theta_{V,W} = \prod_{i=1}^k \cos \Theta_{V_i,W} \cdot \prod_{i=1}^{k-1} \cos \Theta_{P(V_i),P(V_{i+1}\oplus\ldots\oplus V_k)}^{\perp}.$$

Proposition 5.13. For nonzero subspaces $V, W \subset X$ with $V \not\perp W$, a partition $V = \bigoplus_i V_i$ is principal w.r.t. W if, and only if, it is orthogonal and $\cos \Theta_{V,W} = \prod_i \cos \Theta_{V_i,W}$.

Proof. Corollary 5.12 and propositions 2.4i and 5.10 give the converse.

References

- R. A. Brualdi and H. Schneider, Determinantal identities: Gauss, Schur, Cauchy, Sylvester, Kronecker, Jacobi, Binet, Laplace, Muir, and Cayley, Linear Algebra Appl. 52 (1983), 769–791.
- [2] C. Cohen-Tannoudji, B. Diu, and F. Laloe, *Quantum mechanics*, vol. 1, Wiley-VCH, 2019.
- [3] L. Dorst, *The inner products of geometric algebra*, Applications of Geometric Algebra in Computer Science and Engineering (L. Dorst, C. Doran, and J. Lasenby, eds.), Birkhäuser, 2002, pp. 35–46.
- [4] A. Galántai and C. J. Hegedűs, Jordan's principal angles in complex vector spaces, Numer. Linear Algebra Appl. 13 (2006), no. 7, 589–598.
- [5] H. Gluck, Higher curvatures of curves in Euclidean space, II, Amer. Math. Monthly 74 (1967), no. 9, 1049–1056.
- [6] G. H. Golub and C. F. Van Loan, *Matrix computations*, Johns Hopkins University Press, 2013.
- [7] H. Gunawan, O. Neswan, and W. Setya-Budhi, A formula for angles between subspaces of inner product spaces, Beitr. Algebra Geom. 46 (2005), no. 2, 311–320.
- [8] D. Hestenes and G. Sobczyk, *Clifford algebra to geometric calculus*, D. Reidel, Dordrecht, 1984.
- [9] E. Hitzer, Angles between subspaces computed in Clifford algebra, AIP Conference Proceedings, vol. 1281, AIP, 2010, pp. 1476–1479.

- [10] H. Hotelling, Relations between two sets of variates, Biometrika 28 (1936), no. 3/4, 321–377.
- [11] S. Jiang, Angles between Euclidean subspaces, Geom. Dedicata 63 (1996), 113-121.
- [12] Y. Jiao, X. Shen, G. Li, and Y. Gu, Subspace principal angle preserving property of Gaussian random projection, 2018 IEEE Data Science Workshop (DSW), IEEE, 2018, pp. 115–119.
- [13] C. Jordan, Essai sur la géométrie à n dimensions, Bull. Soc. Math. France 3 (1875), 103–174.
- [14] T. Kato, Perturbation theory for linear operators, Springer-Verlag, Berlin, 1995.
- [15] A. L. G. Mandolesi, Grassmann angles between real or complex subspaces, arXiv:1910.00147 [math.GM] (2019).
- [16] _____, Blade products in Grassmann and Clifford algebras, arXiv:1910.07327 [math.GM] (2020).
- [17] _____, Projection factors and generalized real and complex Pythagorean theorems, Adv. Appl. Clifford Al. **30** (2020), no. 43.
- [18] _____, Quantum fractionalism: the Born rule as a consequence of the complex Pythagorean theorem, Phys. Lett. A 384 (2020), no. 28, 126725.
- [19] M. Marcus, *Finite dimensional multilinear algebra*, vol. 2, Marcel Dekker Inc., New York, 1975.
- [20] J. Miao and A. Ben-Israel, On principal angles between subspaces in Rⁿ, Linear Algebra Appl. 171 (1992), 81–98.
- [21] _____, Product cosines of angles between subspaces, Linear Algebra Appl. 237/238 (1996), 71–81.
- [22] T. Muir and W. H. Metzler, A treatise on the theory of determinants, Dover Phoenix editions, Dover Publications, 2003.
- [23] I. B. Risteski and K. G. Trencevski, Principal values and principal subspaces of two subspaces of vector spaces with inner product, Beitr. Algebra Geom. 42 (2001), no. 1, 289–300.
- [24] T. Yokonuma, Tensor spaces and exterior algebra, Translations of Mathematical Monographs, American Mathematical Society, 1992.