

Capped vertex with descendants for zero dimensional A_∞ quiver varieties

H. Dinkins and A. Smirnov

Abstract

In this paper, we study the capped vertex functions associated to certain zero-dimensional type- A Nakajima quiver varieties. The insertion of descendants into the vertex functions can be expressed by the Macdonald operators, which leads to explicit combinatorial formulas for the capped vertex functions.

We determine the monodromy of the vertex functions and show that it coincides with the elliptic R -matrix of symplectic dual variety. We apply our results to give the vertex functions and the characters of the tautological bundles on the quiver varieties formed from arbitrary stability conditions.

1 Introduction

Our main objects of study in this paper are certain K -theoretic enumerative invariants of Nakajima quiver varieties known as vertex functions, see Section 7 in [Oko15] and Section 2 below. Let X be a Nakajima quiver variety, see [Gin12], [Nak98], or Section 2 in [MO12] for an introduction. Vertex functions for X come in two flavors, capped and uncapped. The vertex functions are defined by an equivariant count of quasimaps to X , and the type of vertex function is determined by whether the moduli space of quasimaps is considered with a nonsingular condition or a relative condition, see Section 2 below.

One can consider either type of vertex function with descendants inserted, and these give a collection of natural classes in the K -theory of X . Starting with a vector bundle on X and an evaluation map on the appropriate moduli space of quasimaps, we can pullback the vector bundle under the evaluation

map to obtain a class on the quasimap moduli space. The insertion of a descendant into a vertex function refers to the quasimap count obtained by tensoring the structure sheaf of the quasimap moduli space with a class obtained in this way. The K -theory of Nakajima quiver varieties is generated by a collection of tautological vector bundles, one for each vertex of the quiver, and any of these gives rise to a descendant that can be inserted into a vertex function.

For quiver varieties arising from type- A quivers, there are known procedures for computing the vertex functions as a power series in the Kähler parameters, see Section 1 in [AO17], Section 4.5 in [PSZ16], and Section 2.4 [DS19b]. In this paper, we restrict our attention to the capped and uncapped vertex functions with descendants for zero-dimensional type- A quiver varieties. Such varieties are indexed by partitions, and we denote them by X_λ for a partition λ .

Our main result is that the insertion of descendants into the uncapped vertex function can be realized by the action of the difference operators of the trigonometric Ruijsenaars-Schneider model (also known as the Macdonald difference operators). Let $\mathbf{V}_\lambda(\mathbf{z})$ be the uncapped vertex function for X_λ , and let $\mathbf{V}_\lambda^{(\tau)}(\mathbf{z})$ be the uncapped vertex function with descendant τ , then the Theorem 3 reads:

$$T(\tau)\mathbf{V}_\lambda(\mathbf{z}) = \mathbf{V}_\lambda^{(\tau)}(\mathbf{z}) \quad (1)$$

where $T(\tau)$ is the Macdonald difference operator associated with τ .

In [DS19b], we proved

$$\mathbf{V}_\lambda(\mathbf{z}) = \prod_{\square \in \lambda} \prod_{i=0}^{\infty} \frac{1 - \hbar z_\square q^i}{1 - z_\square q^i} \quad (2)$$

where z_\square denotes a certain monomial in the Kähler parameters depending on the box \square in the Young diagram for λ , see (7) below.

This result allows us to explicitly compute the vertex functions with descendant insertions. For instance, if \mathcal{V}_n is the tautological vector bundle on X_λ corresponding to n th vertex of the quiver and $\tau_{n,r} = \bigwedge^r \mathcal{V}_n$, then from (1) and (2), we obtain the following rational function for the capped vertex function with descendant $\tau_{n,r}$:

$$\hat{\mathbf{V}}_\lambda^{(\tau_{n,r})}(\mathbf{z}) = \sum_{\substack{I \subset \mathcal{C}_\lambda(n) \\ |I|=r}} \prod_{\substack{\square \in I \\ \square' \in \mathcal{C}_\lambda(n) \setminus I}} \frac{\hbar \zeta_{\square'} - \zeta_\square}{\zeta_{\square'} - \zeta_\square} \prod_{\square \in I} \prod_{\square' \in S_\lambda(\square)} \frac{1 - z_{\square'}}{1 - \hbar z_{\square'}}$$

where $C_\lambda(n)$ and $S_\lambda(\square)$ denote certain subsets of boxes in λ , and ζ_\square are monomials in ζ_i , related to the Kähler parameters by $z_i = \frac{\zeta_{i-1}}{\zeta_i}$.

In Sections 5 and 6, we give an application of our formula to quiver varieties with non-canonical stability conditions. In general, Nakajima quiver varieties are defined as geometric invariant theory quotients and thus depend on a choice of stability condition $\theta \in \text{Lie}_{\mathbb{R}}(\mathbf{K})$, where \mathbf{K} is the Kähler torus $\mathbf{K} := (\mathbb{C}^\times)^{|I|}$ and I is the vertex set of the quiver. The varieties X_λ are determined by the positive stability condition, see (3) below. In the literature, explicit computations involving Nakajima quiver varieties almost always consider only the positive and negative stability conditions.

As the stability condition varies, the varieties obtained from them change only when crossing certain hyperplanes. This gives rise to a collection of cones in $\text{Lie}_{\mathbb{R}}(\mathbf{K})$. The toric compactification $\bar{\mathbf{K}}$ of \mathbf{K} given by the fan generated by these cones is known as the Kähler moduli space.

The vertex function of X_λ is the solution of a q -difference equation (see [OS16]), and it is expected that the vertex functions of the varieties given by the same dimension data as X_λ , but with different stability conditions, solve the same q -difference equation. Furthermore, the vertex function corresponding to a choice of stability condition gives a solution of this q -difference equation holomorphic in a neighborhood of the limit point on the Kähler moduli space corresponding to this stability condition.

By studying the explicit form of the q -difference equation in Section 5, it is straightforward to give a formula for the solution holomorphic in a neighborhood of an arbitrary limit point of the Kähler moduli space. For the reasons explained above, such solutions are expected to coincide with the vertex functions of the quiver variety with the appropriate stability condition. As further evidence of this expected correspondence, we examine the monodromy of the q -difference equation and verify that this agrees, up to a constant, with the elliptic R-matrix of the symplectic dual variety, see [AO16].

Putting all this together, we start with the capped vertex function with descendant τ for the variety with positive stability condition, and examine the limit as the Kähler parameters approach a limit point corresponding to a general stability condition θ . This provides us with the character of τ on the quiver variety with identical dimension data as X_λ and arbitrary stability parameter θ .

Acknowledgments

A. Smirnov was supported by the Russian Science Foundation under grant 19-11-00062. The authors would also like to thank Peter Koroteev for pointing out the connection between the vertex functions of the cotangent bundle of the flag variety and the varieties studied here.

2 Quasimaps and Vertex Functions

2.1

Let λ be a Young diagram rotated by 45° as in Figure 1. Let $v_i \in \mathbb{N}$, $i \in \mathbb{Z}$ denote the number of boxes in the i th vertical column, as oriented in the Figure. We assume that $i = 0$ corresponds to the column which contains the corner box of λ . Let $\mathbf{v} = (v_i)$ and $\mathbf{w}_i = \delta_{i,0}$. Let




Figure 1: The partition $\lambda = (5, 4, 3, 2)$ rotated by 45° and $\mathbf{v} = (v_i) = (\dots, 0, 0, 1, 1, 2, 2, 3, 2, 2, 1, 0, 0, \dots)$.

$$X_\lambda := \mathcal{M}(\mathbf{v}, \mathbf{w}) \tag{3}$$

denote the A_∞ Nakajima quiver variety defined by these data, with stability condition given by the character

$$\theta_+ : (g_i) \rightarrow \prod_i \det(g_i).$$

We will refer to this stability condition as the *canonical* or *positive* stability condition. One of the goals of this paper is to analyze the enumerative invariants (vertex functions) of quiver varieties arising for generic $\theta = (\theta_1, \dots, \theta_{|I|})$ stability conditions corresponding to characters

$$\theta : (g_i) \rightarrow \prod_i \det(g_i)^{\theta_i}.$$

Throughout, we will assume that the A_∞ quiver is oriented with arrows pointing to the right. Nakajima quiver varieties come equipped with a natural action of a torus \mathbb{T} and a collection of \mathbb{T} -equivariant vector bundles which we call \mathcal{V}_i , $i \in \mathbb{Z}$. We denote by \hbar the weight of the \mathbb{T} -module $\mathbb{C}\omega$, where ω is the symplectic form on X_λ .

2.2

To define the vertex functions, we need to study moduli spaces of stable quasimaps from \mathbb{P}^1 to a Nakajima quiver variety X , as introduced in [CKM14]. We review the main objects of study in the case of Nakajima quiver varieties, see Section 6 of [Oko15] and Section 2 of [PSZ16].

The definition of a quasimap to a variety X requires a presentation of X as a geometric invariant theory quotient. For a Nakajima quiver variety arising from a quiver Q with vertex set I and dimensions \mathbf{v}, \mathbf{w} , this takes the form

$$X := \mu^{-1}(0) //_{\theta} G_{\mathbf{v}} = \mu^{-1}(0)^{\theta-s} / G_{\mathbf{v}}$$

where μ is the moment map associated to the $G_{\mathbf{v}} := \prod_{i \in I} GL(\mathbf{v}_i)$ action on $T^*Rep_Q(\mathbf{v}, \mathbf{w})$, the cotangent bundle of the space of framed representations of Q and $\mu^{-1}(0)^{\theta-s}$ is the intersection of $\mu^{-1}(0)$ with the stable points defined by a choice of θ ([Gin12]).

In the context of the varieties X_λ , this data looks as follows. The dimension vectors correspond to vector spaces V_i and the space $T^*Rep_Q(\mathbf{v}, \mathbf{w})$ consists of 4-tuples (A, B, I, J) so that

$$A = \bigoplus_{i \in \mathbb{Z}} A_i, \quad B = \bigoplus_{i \in \mathbb{Z}} B_i, \quad A_i \in \text{Hom}(V_i, V_{i+1}), \quad B_i \in \text{Hom}(V_{i+1}, V_i).$$

and

$$I \in V_0 \quad J \in V_0^*$$

The moment map is $\mu(A, B, I, J) = [A, B] + IJ$ and the tuple (A, B, I, J) is θ_+ -stable if and only if

$$\mathbb{C}\langle A, B \rangle I = \bigoplus_{i \in \mathbb{Z}} V_i \quad (4)$$

where $\mathbb{C}\langle A, B \rangle$ denotes the ring of (noncommutative) polynomials in A, B .

For generic choice of θ the condition (4) should be substituted by more complicated conditions described in Proposition 5.1.5 in [Gin12].

2.3

We recall some facts about quasimaps to a quiver variety X . For more details, see [CKM14] and [Oko15] Sections 4-6.

Definition 1. A stable genus zero quasimap to X relative to p_1, \dots, p_m is given by the following data

$$(C, p'_1, \dots, p'_m, P, f, \pi)$$

where

- C is a genus zero connected curve with at worst nodal singularities and the p'_1, \dots, p'_m are nonsingular points of C .
- P is a principal G_v bundle over C .
- f is a section of the bundle $P \times_{G_v} T^*Rep_Q(v, w)$ satisfying $\mu = 0$.
- $\pi : C \rightarrow D$ is a regular map.

satisfying the following conditions

1. There is a distinguished component C_0 of C so that π restricts to an isomorphism $\pi : C_0 \cong D$ and $\pi(C \setminus C_0)$ is zero dimensional (possibly empty).
2. $\pi(p'_i) = p_i$.
3. $f(p)$ is stable in the sense of (4) for all but a finite set of points disjoint from p'_1, \dots, p'_m and the nodes of C .

4. The line bundle $\omega_{\tilde{C}} \left(\sum_i p'_i + \sum_j q_j \right) \otimes \mathcal{L}_\theta^\epsilon$ is ample for every rational $\epsilon > 0$, where $\mathcal{L}_\theta = P \times_{G_v} \mathbb{C}_\theta$, \tilde{C} is the closure of $C \setminus C_0$, q_j are the nodes of \tilde{C} , and \mathbb{C}_θ is the one dimensional G_v -module defined by the stability condition θ .




Figure 2: An example of the domain of a relative quasimap with four marked points. A chain of rational curves is attached to each point p_i , and condition 4 implies that the last component of each chain has a marked point p'_i . The map π collapses each chain to a single point.

Definition 2. A relative quasimap $(C, p'_1, \dots, p'_m, P, f, \pi)$ is nonsingular at $p \in C$ if $f(p)$ is stable in the sense of (4). In this case, $f(p)$ gives a point in the quiver variety.

Definition 3. The degree of a quasimap $(C, p'_1, \dots, p'_m, P, f, \pi)$ is the tuple $\mathbf{d} = (d_i)_{i \in \mathbb{Z}}$ where d_i is the degree of the rank v_i vector bundle $P \times_{G_v} V_i \rightarrow C$.

Theorem 1. ([CKM14] Theorem 7.2.2) The stack $\mathbf{QM}_{\text{relative } p_1, \dots, p_m}^{\mathbf{d}}$ parameterizing the data of stable genus zero quasimaps to X is a Deligne-Mumford stack of finite type with a perfect obstruction theory.

Definition 4. Let $\mathbf{QM}_{\text{nonsing } p_1, \dots, p_m}^{\mathbf{d}}$ be the stack parameterizing the data of degree \mathbf{d} quasimaps to X relative to p_1, \dots, p_m such that $C \cong D \cong \mathbb{P}^1$. For such a quasimap, most of the conditions in Definition 1 become trivially satisfied.

Restricting the obstruction theory of $\mathbf{QM}_{\text{relative } p_1, \dots, p_m}^d$ gives a perfect obstruction theory on $\mathbf{QM}_{\text{nonsing } p_1, \dots, p_m}^d$. The symmetrized virtual structure sheaf on such a space will be denoted by $\widehat{\mathcal{O}}_{\text{vir}}^d$, with the context determining exactly which quasimap space we are considering.

Given a quasimap $(C, p'_1, \dots, p'_m, P, f, \pi)$ and $p \in C$, there is an evaluation map to the quotient stack:

$$\text{ev}_p(C, p'_1, \dots, p'_m, P, f, \pi) = f(p) \in [\mu^{-1}(0)/G_{\mathbf{v}}]$$

Given a Schur functor τ in the tautological bundles on X_λ , let τ_{stack} be the associated K -theory class on $[\mu^{-1}(0)/G_{\mathbf{v}}]$. Then we can define an induced K -theory class on $\mathbf{QM}_{\text{relative } p_1, \dots, p_m}^d$:

$$\tau|_p := \text{ev}_p^*(\tau_{\text{stack}}) \quad (5)$$

2.4

The action of the torus \mathbb{T} on a quiver variety X and of \mathbb{C}_q^\times on \mathbb{P}^1 induce an action of $\mathbb{T} \times \mathbb{C}_q^\times$ on quasimaps to X . Let $p_1 = 0$ and $p_2 = \infty$ in \mathbb{P}^1 . In what follows, we will denote $\mathbf{z}^d = \prod_i z_i^{d_i}$ and use this notation to keep track of the degree of quasimaps. The variables z_i are known as the Kähler parameters, and are characters of the Kähler torus

$$\mathbf{K} := (\mathbb{C}^\times)^{|I|}$$

where I is the vertex set of the quiver.

The evaluation maps on relative quasimaps are proper ([Oko15] Section 7.4), and thus we can make the following definition.

Definition 5. The capped vertex function with descendant τ inserted at p_1 is the formal power series

$$\widehat{\mathbf{V}}^{(\tau)}(\mathbf{z}) = \sum_{\mathbf{d}} \text{ev}_{p_2, *}(\widehat{\mathcal{O}}_{\text{vir}}^d \otimes \tau|_{p_1}, \mathbf{QM}_{\text{relative } p_2}^d) \mathbf{z}^d \in K_{\mathbb{T}}(X)[[\mathbf{z}]]$$

where $\widehat{\mathcal{O}}_{\text{vir}}^d$ is the symmetrized virtual structure sheaf on $\mathbf{QM}_{\text{relative } p_2}^d$.

While the evaluation map ev_{p_2} on $\mathbf{QM}_{\text{nonsing } p_2}^d$ is not proper, the restriction to the \mathbb{C}_q^\times -fixed locus

$$\text{ev}_{p_2} : (\mathbf{QM}_{\text{nonsing } p_2}^d)^{\mathbb{C}_q^\times} \rightarrow X$$

is ([Oko15] Section 7.2). Using equivariant localization, we can thus make the following definition.

Definition 6. The bare vertex function with descendant τ inserted at p_1 is the formal power series

$$\mathbf{V}^{(\tau)}(\mathbf{z}) = \sum_d \text{ev}_{p_2,*}(\widehat{\mathcal{O}}_{\text{vir}}^d \otimes \tau|_{p_1}, \text{QM}_{\text{nonsing } p_2}^d) \mathbf{z}^d \in K_{\mathbb{T} \times \mathbb{C}_q^\times}(X)_{\text{loc}}[[\mathbf{z}]]$$

where $\widehat{\mathcal{O}}_{\text{vir}}^d$ is the symmetrized virtual structure sheaf on $\text{QM}_{\text{nonsing } p_2}^d$.

In what follows, we will omit the superscript (τ) in the bare vertex function when $\tau = 1$.

2.5

Definition 7. The capping operator is the formal series

$$\Psi(\mathbf{z}) = \sum_d \text{ev}_{p_1,*} \otimes \text{ev}_{p_2,*}(\widehat{\mathcal{O}}_{\text{vir}}^d, \text{QM}_{\text{nonsing } p_2}^d \text{ relative } p_1) \mathbf{z}^d \in K_{\mathbb{T}}^{\otimes 2}(X)_{\text{loc}}[[\mathbf{z}]]$$

where $\widehat{\mathcal{O}}_{\text{vir}}^d$ denotes the symmetrized virtual structure sheaf on $\text{QM}_{\text{nonsing } p_2}^d \text{ relative } p_1$

The standard pairing on equivariant K -theory

$$(\mathcal{F}, \mathcal{G}) = \chi(\mathcal{F} \otimes \mathcal{G})$$

allows us to interpret $\Psi(\mathbf{z})$ as a linear map

$$\Phi(\mathbf{z}) : K_{\mathbb{T}}(X)_{\text{loc}}[[\mathbf{z}]] \rightarrow K_{\mathbb{T}}(X)_{\text{loc}}[[\mathbf{z}]]$$

We have the following theorem:

Theorem 2. ([Oko15] Section 7.4) The capping operator satisfies the equation

$$\widehat{\mathbf{V}}^{(\tau)}(\mathbf{z}) = \Psi(\mathbf{z}) \mathbf{V}^{(\tau)}(\mathbf{z})$$

2.6

In the simplest situation of the zero-dimensional quiver varieties X_λ , which we consider in the present paper, we have $K_{\mathbb{T}}(X_\lambda)_{loc} = \mathbb{Q}(\hbar, q)$. Thus, in this case all the functions defined above are power series in the Kähler parameters with some rational coefficients in \hbar, q :

$$\hat{\mathbf{V}}^{(\tau)}(\mathbf{z}), \mathbf{V}^{(\tau)}(\mathbf{z}), \Psi(\mathbf{z}) \in \mathbb{Q}(\hbar, q)[[\mathbf{z}]]$$

Let us denote $g(\mathbf{z}) = \hat{\mathbf{V}}^{(1)}(\mathbf{z})$. With this notation, it follows from the previous theorem that for X_λ we have

$$\Psi(\mathbf{z}) = \frac{g(\mathbf{z})}{\mathbf{V}^{(1)}(\mathbf{z})}$$

and thus the capped vertex with descendent τ has the form:

$$\hat{\mathbf{V}}^{(\tau)}(\mathbf{z}) = g(\mathbf{z}) \frac{\mathbf{V}^{(\tau)}(\mathbf{z})}{\mathbf{V}^{(1)}(\mathbf{z})}$$

We see that $g(\mathbf{z})$ appears as a normalization prefactor in the formulas for the capped vertex. Thus, it will be convenient to redefine the capped vertex function by normalizing as $\hat{\mathbf{V}}^{(\tau)}(\mathbf{z}) \rightarrow \hat{\mathbf{V}}^{(\tau)}(\mathbf{z})/g(\mathbf{z})$.

Let us note here that the function $g(\mathbf{z})$ can be computed explicitly. It coincides with the multiplicative identity of the quasimap quantum K-theory ring, see Section 3.2 [PSZ16]. In the case of X_λ , it is given by the gluing matrix \mathbf{G} . It can be shown [KS] that the gluing matrix of X equals the zero slope K -theoretic R -matrix of the symplectic dual variety $X^!$. The K -theoretic version of Proposition 6 then gives:

$$g(\mathbf{z}) = \mathbf{G} = \prod_{\square \in \lambda} \frac{1 - z_\square}{1 - \hbar z_\square}.$$

3 Vertex Functions for X_λ

3.1

Fix a partition λ , We do not distinguish between a partition and its Young diagram. Let $\mathbf{v} = (v_i)$ be as in Section 2.1. We define the following functions for $\square \in \lambda$. Let $c_\lambda(\square)$ denote the content of \square , or the horizontal coordinate

of \square when the Young diagram of λ is rotated as in Figure 3. We assume the corner box has content 0. Let $C_\lambda(n)$ be the set of all boxes of content n .

Let $h_\lambda(\square)$ denote the height of \square in λ , normalized so that the heights of the boxes with content i take all the values between 1 and \mathbf{v}_i . Let $a_\lambda(\square)$ and $l_\lambda(\square)$ denote the length of the arm and leg (not including \square) based at \square , respectively. If \square has rectangular coordinates (i, j) , then

$$l_\lambda(\square) = \lambda_i - j \quad \text{and} \quad a_\lambda(\square) = \lambda'_j - i$$

where λ' is the transpose of λ .

Define

$$\zeta_\square = \begin{cases} \left(\frac{\hbar}{q}\right)^{h_\lambda(\square)} \zeta_{c(\square)+a_\lambda(\square)} & \text{if } c(\square) \geq 0 \\ \left(\frac{q}{\hbar}\right)^{h_\lambda(\square)} \zeta_{c(\square)-l_\lambda(\square)-1} & \text{if } c(\square) < 0 \end{cases}$$

where ζ_i are a collection of variables related to the Kähler parameters by $z_i = \frac{\zeta_{i-1}}{\zeta_i}$. Define the difference operator

$$p_i f(\zeta_i) := f(q\zeta_i)$$

where f is some function of ζ_i . Then define

$$p_\square = \begin{cases} \prod_{i=c(\square)+a_\lambda(\square)}^{\prod_{i=c(\square)+\mathbf{v}_{c(\square)}-h_\lambda(\square)} p_i^{-1}} & \text{if } c(\square) \geq 0 \\ \prod_{i=c(\square)+l_\lambda(\square)-1}^{\prod_{i=c(\square)-\mathbf{v}_{c(\square)}+h_\lambda(\square)-1} p_i} & \text{if } c(\square) < 0 \end{cases} \quad (6)$$

See Figure 3 for an example.

3.2

Fix $n, r \in \mathbb{Z}$ so that $1 \leq r \leq \mathbf{v}_n$.

We define the following difference operator:

$$T_\lambda^{n,r} = \hbar^{r(r-1)/2+\beta(n)} \sum_{\substack{I \subset C_\lambda(n) \\ |I|=r}} \prod_{\substack{\square \in I \\ \square' \in C_\lambda(n) \setminus I}} \frac{\hbar \zeta_{\square'} - \zeta_\square}{\zeta_{\square'} - \zeta_\square} \prod_{i \in I} p_\square$$

where $\beta(n) = |n|$ if $n < 0$ and 0 otherwise. We note that up to normalization and relabeling of the variables, the operators $T_\lambda^{n,r}$ are exactly the Macdonald difference operators, also known as the difference operators of the trigonometric Ruijsenaars-Schneider model, see [Mac79] and [Kor18]. In particular the q -difference operators $T_\lambda^{n,r}$ commute with each other. We denote by

$$\text{RS} = \mathbb{Q}[T_\lambda^{n,r}]_{n,r \in \mathbb{Z}}$$

the commutative ring they generate.




Figure 3: The values of ζ_{\square} are shown inside the boxes of $C_{\lambda}(0)$. If \square is the box with content 0 and height 1, then $a_{\lambda}(\square) = 3$ and $l_{\lambda}(\square) = 4$. Also, $p_{\square} = \prod_{i=0+3-1}^{0+3} p_i^{-1} = p_2^{-1} p_3^{-1}$.

3.3

Let us recall that the descendent insertions of the vertex functions are labeled by elements (5) of the ring:

$$K([\mu^{-1}(0)/G_{\mathbf{v}}]) \cong \text{char}(G_{\mathbf{v}}) \quad \text{where} \quad G_{\mathbf{v}} = \prod_{i \in \mathbb{Z}} GL(\mathbf{v}_i)$$

Let us denote by V_n the \mathbf{v}_n -dimensional fundamental representation of $GL(\mathbf{v}_n)$. Then this ring is generated by the classes $\tau_{n,r} := \bigwedge^r V_n$. Let us define a homomorphism

$$T : \text{char}(G_{\mathbf{v}}) \rightarrow \text{RS}$$

by $T(\tau_{n,r}) = T_{\lambda}^{n,r}$. Applying the substitution $z_i = \frac{\zeta_{i-1}}{\zeta_i}$, the elements of RS act as q -difference operators on the vertex functions. Our main result involves expressing the insertion of a descendent as the action of such operators:

Theorem 3. The insertion of descendent τ into the bare vertex function can be expressed as

$$T(\tau) \mathbf{V}_{\lambda}(\mathbf{z}) = \mathbf{V}_{\lambda}^{(\tau)}(\mathbf{z})$$

Clearly, to prove the theorem it is enough to show that

$$T_{\lambda}^{n,r} \mathbf{V}_{\lambda}(\mathbf{z}) = \mathbf{V}_{\lambda}^{(\tau_{n,r})}(\mathbf{z}).$$

3.4

Before giving the proof of Theorem 3, we explore an important consequence. Define

$$z_{\square} := \prod_{\square' \in H_{\lambda}(\square)} \widehat{z}_{c(\square')}$$

where the shifted parameters \widehat{z}_i are

$$\widehat{z}_i := \left(\frac{\hbar}{q}\right)^{\sigma_{\lambda}(i)} z_i \quad \text{where} \quad \sigma_{\lambda}(i) := \begin{cases} \mathbf{v}_{i-1} - \mathbf{v}_i & \text{if } i \neq 0 \\ \mathbf{v}_{i-1} - \mathbf{v}_i + 1 & \text{if } i = 0 \end{cases}$$

and $H_{\lambda}(\square)$ denotes the set of boxes in the hook based at \square in λ . See Figure 4.




Figure 4: The shaded boxes are an example of a hook in $\lambda = (5, 4, 3, 2)$. If \square is the box at the base of the hook shown, then $z_{\square} = z_0 \left(\frac{\hbar}{q} z_1\right) z_2 \left(\frac{\hbar}{q} z_3\right)$ and $\zeta_{\square} = \frac{\hbar}{q} \zeta_3$.

In [DS19b], the authors prove the following formula for $\mathbf{V}_{\lambda}(z)$:

Theorem 4. ([DS19b] Theorem 1)

$$\mathbf{V}_{\lambda}(z) = \prod_{\square \in \lambda} \prod_{i=0}^{\infty} \frac{1 - \hbar z_{\square} q^i}{1 - z_{\square} q^i}. \quad (7)$$

Note that z_{\square} can be expressed in terms of the ζ_i as

$$z_{\square} = \frac{\zeta_{c_{\lambda}(\square)-l_{\lambda}(\square)-1}}{\zeta_{c_{\lambda}(\square)+a_{\lambda}(\square)}} \prod_{i=c_{\lambda}(\square)-l_{\lambda}(\square)}^{c_{\lambda}(\square)+a_{\lambda}(\square)} \left(\frac{\hbar}{q}\right)^{\sigma_{\lambda}(i)} \quad (8)$$

Using our notation above, the set of all boxes with the same content as a given box $\square \in \lambda$ is denoted $C_{\lambda}(c_{\lambda}(\square))$. There is a minimal rectangular Young diagram $\mu \subset \lambda$ containing $C_{\lambda}(c_{\lambda}(\square))$. Explicitly, if $c_{\lambda}(\square) = i < 0$, then $\mu = (\mu_1, \dots, \mu_{v_i})$ where $\mu_j = -i + v_i$. If $c_{\lambda}(\square) = i \geq 0$, then $\mu = (\mu_1, \dots, \mu_{i+v_i})$ where $\mu_j = v_i$. With this notation, we define the slice inside λ through \square as:

$$S_{\lambda}(\square) := \begin{cases} \text{row containing } \square \subset \mu & \text{if } c_{\lambda}(\square) \geq 0 \\ \text{column containing } \square \subset \mu & \text{if } c_{\lambda}(\square) < 0 \end{cases}$$

where the row and column are understood in the ‘‘rectangular’’ sense, see Figure 5.




Figure 5: Two slices inside λ are shown. The red boxes make up the slice through the box with content 1 and height 1. The blue boxes make up the slice through the box with content -2 and height 1.

Combining Theorem 3 with (7), we obtain the following corollary, giving an explicit combinatorial formula for the capped vertex with descendant.

Corollary 1.

$$\hat{\mathbf{V}}_{\lambda}^{(\tau_{n,r})}(\mathbf{z}) = \hbar^{r(r-1)/2+\beta(n)} \sum_{\substack{I \subset C_{\lambda}(n) \\ |I|=r}} \prod_{\substack{\square \in I \\ \square' \notin I}} \frac{\hbar \zeta_{\square'} - \zeta_{\square}}{\zeta_{\square'} - \zeta_{\square}} \prod_{\square \in I} \prod_{\square' \in S_{\lambda}(\square)} \frac{1 - z_{\square'}}{1 - \hbar z_{\square'}}$$

Proof. By Theorem 2, we know that

$$\hat{\mathbf{V}}_{\lambda}^{(\tau_{n,r})}(\mathbf{z}) = \frac{\mathbf{V}_{\lambda}^{(\tau_{n,r})}(\mathbf{z})}{\mathbf{V}_{\lambda}(\mathbf{z})}$$

Applying Theorem 3, we obtain

$$\hat{\mathbf{V}}_{\lambda}^{(\tau_{n,r})}(\mathbf{z}) = \frac{T_{\lambda}^{n,r} \mathbf{V}_{\lambda}(\mathbf{z})}{\mathbf{V}_{\lambda}(\mathbf{z})}$$

From (6) and (8), we see that the operator p_{\square} has the following effect on $z_{\square'}$:

$$p_{\square}(z_{\square'}) = \begin{cases} q z_{\square'}, & \square' \in S_{\lambda}(\square) \\ z_{\square'}, & \square' \notin S_{\lambda}(\square) \end{cases}$$

From (7), we see that the bare vertex function transforms under the scaling of a fixed z_{\square} by q as follows:

$$\mathbf{V}_{\lambda}(\mathbf{z}) \Big|_{z_{\square}=qz_{\square}} = \frac{1 - z_{\square}}{1 - \hbar z_{\square}} \mathbf{V}_{\lambda}(\mathbf{z})$$

Putting all this together, we have

$$\begin{aligned} \hat{\mathbf{V}}_{\lambda}^{(\tau_{n,r})}(\mathbf{z}) &= \frac{T_{\lambda}^{n,r} \mathbf{V}_{\lambda}(\mathbf{z})}{\mathbf{V}_{\lambda}(\mathbf{z})} \\ &= \frac{1}{\mathbf{V}_{\lambda}(\mathbf{z})} \left(\hbar^{r(r-1)/2+\beta(n)} \sum_{\substack{I \subset C_{\lambda}(n) \\ |I|=r}} \prod_{\substack{\square \in I \\ \square' \notin I}} \frac{\hbar \zeta_{\square'} - \zeta_{\square}}{\zeta_{\square'} - \zeta_{\square}} \prod_{\square \in I} \prod_{\square' \in S_{\lambda}(\square)} \frac{1 - z_{\square'}}{1 - \hbar z_{\square'}} \right) \mathbf{V}_{\lambda}(\mathbf{z}) \\ &= \hbar^{r(r-1)/2+\beta(n)} \sum_{\substack{I \subset C_{\lambda}(n) \\ |I|=r}} \prod_{\substack{\square \in I \\ \square' \notin I}} \frac{\hbar \zeta_{\square'} - \zeta_{\square}}{\zeta_{\square'} - \zeta_{\square}} \prod_{\square \in I} \prod_{\square' \in S_{\lambda}(\square)} \frac{1 - z_{\square'}}{1 - \hbar z_{\square'}} \end{aligned}$$

□

We note that for $|q| < 1$, $\hat{\mathbf{V}}_{\lambda}^{(\tau_{n,r})}(\mathbf{z})$ converges to a rational function of the Kähler parameters. Rationality of descendant insertions in the case of the cohomology of the Hilbert scheme of points in \mathbb{C}^2 was established in [PP13]. It is expected that the same is true in K -theory for general Nakajima quiver varieties, see [AO17].

4 Proof of Theorem 3

4.1

The vertex functions for type A Nakajima quiver varieties can be described by natural integral representations of Mellin-Barnes type see Section 1.1.5-1.1.6 in [AO17]. These integral representations were investigated for the quiver varieties isomorphic to cotangent bundles over Grassmannians and partial flag varieties in [PSZ16; Kor+17; Kor18]. We explain below what this description looks like.

Let X be a Nakajima quiver variety arising from a type A_n quiver with vertex set I , dimension vector \mathbf{v} , and framing dimension vector \mathbf{w} .

For a character $w_1 + \dots + w_m \in K_{\mathbb{T}}(X)$ we denote

$$\Phi(w_1 + \dots + w_m) = \varphi(w_1) \dots \varphi(w_m), \quad \varphi(w) := \prod_{n=0}^{\infty} (1 - wq^n).$$

and extend this definition by linearity to polynomials in $K_{\mathbb{T}}(X)$ with negative coefficients. Let \mathcal{P} be the bundle over X associated to the virtual G -module

$$\bigoplus_{i \in I} \text{Hom}(W_i, V_i) + \bigoplus_{i \rightarrow j} \text{Hom}(V_i, V_j) - \bigoplus_{i \in I} \text{Hom}(V_i, V_i) \quad (9)$$

where $i \rightarrow j$ denotes the sum over the arrows of the quiver.

If $x_{i,1}, \dots, x_{i,v_i}$ denote the Grothendieck roots of i -th tautological bundle (i.e. the bundle over X associated to the G -module V_i) and $a_{i,j}$ denote the equivariant parameters associated to the framings for $j = 1, \dots, w_i$, then

$$\begin{aligned} \mathcal{P} = \sum_{i \in I} \left(\sum_{j=1}^{v_i} x_{i,j}^{-1} \right) \left(\sum_{j=1}^{v_{i+1}} x_{i+1,j} \right) + \sum_{i \in I} \left(\sum_{j=1}^{w_i} a_{i,j}^{-1} \right) \left(\sum_{j=1}^{v_i} x_{i,j} \right) \\ - \sum_{i \in I} \left(\sum_{j=1}^{v_i} x_{i,j}^{-1} \right) \left(\sum_{j=1}^{v_i} x_{i,j} \right) \in K_{\mathbb{T}}(X). \end{aligned}$$

We abbreviate the set of Grothendieck roots of the tautological bundles by \mathbf{x} and define the following formal expression:

$$\mathbf{e}(\mathbf{x}, \mathbf{z}) := \exp \left(\frac{1}{\ln q} \sum_{i \in I} \ln(z_i) \ln(\det \mathcal{V}_i) \right) = \exp \left(\frac{1}{\ln q} \sum_{i \in I} \sum_{j=1}^{v_i} \ln(z_i) \ln(x_{i,j}) \right)$$

For $t \in K_{\mathbb{T}}(X)$ and $p \in X^{\mathbb{T}}$, let $t|_p \in K_{\mathbb{T}}(p)$ be the restriction of t . Then the restriction of the vertex function to a fixed point $p \in X^{\mathbb{T}}$ is

$$\mathbf{V}_p^{(\tau)}(\mathbf{a}, \mathbf{z}) = \frac{1}{\Phi\left((q - \hbar)\mathcal{P}|_p\right)\mathbf{e}(\mathbf{x}|_p, \mathbf{z})} \tilde{\mathbf{V}}_p^{(\tau)}(\mathbf{a}, \mathbf{z}), \quad (10)$$

where

$$\tilde{\mathbf{V}}_p^{(\tau)}(\mathbf{a}, \mathbf{z}) = \prod_{i,j} \int_0^{x_{i,j}|_p} d_q x_{i,j} \Phi\left((q - \hbar)\mathcal{P}\right)\mathbf{e}(\mathbf{x}, \mathbf{z})\tau(\mathbf{x})$$

and the integral symbols denote the Jackson q -integral over all Grothendieck roots:

$$\int_0^a d_q x f(x) := \sum_{n=0}^{\infty} f(aq^n),$$

For an indeterminate x , we define the q -Pochhammer symbol by

$$(x)_d := \frac{\varphi(x)}{\varphi(xq^d)}$$

We note that (10) is a power series in z_i with coefficients given by combinations of q -Pochhammer symbols in the equivariant parameters.

Equation (10) arises when one analyzes the torus fixed points on the quasimap moduli space and computes the vertex function using K -theoretic equivariant localization.

4.2

The cotangent bundle of the full flag variety can be described as a Nakajima quiver variety corresponding to the A_{n-1} quiver with dimension vectors $\mathbf{v} = (n-1, n-2, \dots, 0)$ and $\mathbf{w} = (n, 0, \dots, 0)$. In [Kor18], Koroteev proves that the vertex function of the cotangent bundle of the full flag variety restricted to an appropriate fixed point is an eigenvector of the tRS operators, with eigenvalues given by the elementary symmetric functions of the equivariant parameters.

More generally, we can allow some redundant information which disappears upon taking the quotient. We start with a partition λ with associated dimension \mathbf{v} as before, and some integer n which will correspond to the content of a set of boxes in λ . If $n \geq 0$, we consider the quiver variety with

dimensions given by the number of boxes in the Young diagram strictly to the right of the n th column. If $n < 0$, we obtain our dimensions from the portion of the partition strictly to the left of the n th column. The number of boxes with content n gives the framing dimension.

The corresponding quiver variety is canonically isomorphic to the cotangent bundle of the variety parameterizing full flags inside a v_n -dimensional vector space. The vertex functions of the quiver varieties differ, as the vertex functions are sensitive to the way in which the quotient is taken. Because of this difference, we refer to the flag variety obtained by a partition λ and choice of n as a “redundant flag variety.”

For an example of this “partition truncation”, see Figure 6. In this notation, the usual quiver variety description of the cotangent bundle of the full flag variety can be obtained from square partitions.




Figure 6: Obtaining the dimension information for a redundant flag variety from the partition $(5, 4, 3, 2, 1)$ with $n = 0$. The framing corresponds to the square node.

With this in mind, we can make sense of ζ_{\square} as before for \square inside a truncated partition.

Let

$$T_r = \hbar^{r(r-1)/2} \sum_{\substack{I \subset C_\lambda(n) \\ |I|=r}} \prod_{\substack{\square \in I \\ \square' \notin I}} \frac{\hbar \zeta_{\square'} - \zeta_\square}{\zeta_{\square'} - \zeta_\square} \prod_{\square \in I} \tilde{p}_\square$$

where $\tilde{p}_\square = \frac{a_{n-h_\lambda(\square)}}{\hbar^{n-h_\lambda(\square)-1}} p_\square$.

As before, with the change of variables $z_i = \frac{\zeta_{i-1}}{\zeta_i}$, the operators T_r act on the vertex functions of the redundant flag variety. In our notation, Koroteev's theorem can be generalized to the following:

Theorem 5. ([Kor18] Theorem 2.6) Fix λ and n . Let $\mathbf{V}_p(\mathbf{a}, \mathbf{z})$ be the bare vertex function of the associated redundant flag variety restricted to the fixed point p at which the weights of the tautological bundle \mathcal{V}_i are $\{a_j : 1 \leq j \leq \mathbf{v}_i\}$ for all i . Then

$$T_r \mathbf{V}_p(\mathbf{a}, \mathbf{z}) = e_r(\mathbf{a}) \mathbf{V}_p(\mathbf{a}, \mathbf{z})$$

where $e_r(\mathbf{a})$ denotes the r th elementary symmetric function in the equivariant parameters.

4.3

Let λ be a partition so that the associated dimension vector is $\mathbf{v} = (\mathbf{v}_{-r}, \dots, \mathbf{v}_s)$. Fix $n \in \mathbb{Z}$. For definiteness, we assume $n \geq 0$. Let $\mathbf{V}_p(\mathbf{a}, \mathbf{z})$ be as in Theorem 5. In other words, $\mathbf{V}_p(\mathbf{a}, \mathbf{z})$ is the restriction to a fixed point of the vertex function of the quiver variety with dimension $(\mathbf{v}_{n+1}, \dots, \mathbf{v}_s)$ and framing dimension $(\mathbf{v}_n, 0, \dots, 0)$.

Note 1. The proof of Theorem 3 involves interpreting certain terms in the power series for $\mathbf{V}_\lambda(\mathbf{z})$ as vertex functions for redundant flag varieties with specialized equivariant parameters. For definiteness, we will assume throughout that $n \geq 0$. If $n < 0$, the same arguments given below can be modified in the obvious manner.

Lemma 1. Specializing the equivariant parameters to $a_i = \hbar^{i-1} q^{d_{n,i}}$ and relabeling the Kähler parameters in $\mathbf{V}_p(\mathbf{a}, \mathbf{z})$, we have

$$\mathbf{V}_\lambda(\mathbf{z}) = \sum_{d_{i,j}} \Psi \prod_{i=-r}^n \prod_{j=1}^{\mathbf{v}_i} z_i^{d_{i,j}} \prod_{i=n+1}^s \prod_{j=1}^{\mathbf{v}_i} z_i^{d_{n,j}} \mathbf{V}_p(\mathbf{a}, z_{n+1}, \dots, z_s) \Big|_{a_i = \hbar^{i-1} q^{d_{n,i}}}$$

where Ψ represents φ function terms in (10) that do not depend on $d_{i,j}$ for $i > n$.

Proof. From (10) vertex function for X_λ is

$$\begin{aligned}
\mathbf{V}_\lambda(\mathbf{z}) &= \sum_{d_{i,j}} \prod_{j=1}^{v_0} \frac{\varphi(\hbar^j)}{\varphi(\hbar^j q^{d_{0,j}})} \frac{\varphi(q \hbar^{j-1} q^{d_{0,j}})}{\varphi(q \hbar^{j-1})} \\
&\prod_{i < 0} \prod_{j=1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi(\hbar^{k-j})}{\varphi(\hbar^{k-j} q^{d_{i+1,k-d_{i,j}}})} \frac{\varphi(q \hbar^{k-j-1} q^{d_{i+1,k-d_{i,j}}})}{\varphi(q \hbar^{k-j-1})} \\
&\prod_{i \geq 0} \prod_{j=1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi(\hbar^{k-j+1})}{\varphi(\hbar^{k-j+1} q^{d_{i+1,k-d_{i,j}}})} \frac{\varphi(q \hbar^{k-j} q^{d_{i+1,k-d_{i,j}}})}{\varphi(q \hbar^{k-j})} \\
&\prod_{i \in \mathbb{Z}} \prod_{j,k=1}^{v_i} \frac{\varphi(q \hbar^{k-j})}{\varphi(q \hbar^{k-j} q^{d_{i,k-d_{i,j}}})} \frac{\varphi(\hbar^{k-j+1} q^{d_{i,k-d_{i,j}}})}{\varphi(\hbar^{k-j+1})} \mathbf{z}^{\mathbf{d}} \tag{11}
\end{aligned}$$

where each $d_{i,j}$ is summed from 0 to ∞ .

Separating the terms corresponding to the i th column for $i > n$, we have

$$\begin{aligned}
\mathbf{V}_\lambda(\mathbf{z}) &= \sum_{d_{i,j}} \Psi \prod_{i=-r}^n \prod_{j=1}^{v_i} z_i^{d_{i,j}} \prod_{j=1}^{v_n} \prod_{k=1}^{v_{n+1}} \frac{\varphi(q \hbar^{i-j} q^{d_{n+1,k-d_{n,j}}})}{\varphi(\hbar \hbar^{i-j} q^{d_{n+1,k-d_{n,j}}})} \\
&\prod_{i=n+1}^{s-1} \prod_{j=1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi(q \hbar^{k-j} q^{d_{i+1,k-d_{i,j}}})}{\varphi(\hbar \hbar^{k-j} q^{d_{i+1,k-d_{i,j}}})} \\
&\prod_{i=n+1}^s \prod_{j,k=1}^{v_i} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{i,k-d_{i,j}}})}{\varphi(q \hbar^{k-j} q^{d_{i,k-d_{i,j}}})} \prod_{i=n+1}^s \prod_{j=1}^{v_i} z_i^{d_{i,j}}
\end{aligned}$$

where Ψ represents some φ function terms, which do not depend on $d_{i,j}$ for $i > n$.

On the other hand, from (10) the vertex function of the redundant flag variety arising from a partition λ and integer n at the fixed point p has the form

$$\begin{aligned}
\mathbf{V}_p(\mathbf{a}, z_{n+1}, \dots, z_s) &= \sum_{f_{i,j}} \prod_{j=1}^{v_n} \prod_{k=1}^{v_{n+1}} \frac{\varphi\left(q \frac{a_k}{a_j} q^{f_{n+1,k}}\right)}{\varphi\left(q \frac{a_k}{a_j}\right)} \frac{\varphi\left(\hbar \frac{a_k}{a_j}\right)}{\varphi\left(\hbar \frac{a_k}{a_j} q^{f_{n+1,k}}\right)} \\
&\quad \prod_{i=n+1}^{s-1} \prod_{j=1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi\left(q \frac{a_k}{a_j} q^{f_{i+1,k}-f_{i,j}}\right)}{\varphi\left(q \frac{a_k}{a_j}\right)} \frac{\varphi\left(\hbar \frac{a_k}{a_j}\right)}{\varphi\left(\hbar \frac{a_k}{a_j} q^{f_{i+1,k}-f_{i,j}}\right)} \\
&\quad \prod_{i=n+1}^s \prod_{j,k=1}^{v_i} \frac{\varphi\left(\hbar \frac{a_k}{a_j} q^{f_{i,k}-f_{i,j}}\right)}{\varphi\left(\hbar \frac{a_k}{a_j}\right)} \frac{\varphi\left(q \frac{a_k}{a_j}\right)}{\varphi\left(q \frac{a_k}{a_j} q^{f_{i,k}-f_{i,j}}\right)} \prod_{i=n+1}^s \prod_{j=1}^{v_i} z_i^{f_{i,j}}
\end{aligned}$$

We reindex the summation as follows. First, replace $f_{n,j}$ by $d_{n,j}$. Second, substitute $f_{i,j}$ by $d_{i,j} - d_{n,j}$ for $i > n$. By examining the terms in $\mathbf{V}_\lambda(\mathbf{z})$, we see that a term in the sum is only nonzero if and only if the $d_{i,j}$ give a set of interlacing partitions (see [DS19b] Proposition 7). In particular, we must have $d_{i,j} - d_{n,j} \geq 0$ and so the sum for $\mathbf{V}_p(\mathbf{a}, z_{n+1}, \dots, z_s)$ can be reindexed as

$$\begin{aligned}
\mathbf{V}_p(\mathbf{a}, z_{n+1}, \dots, z_s) &= \sum_{d_{i,j}} \prod_{j=1}^{v_n} \prod_{k=1}^{v_{n+1}} \frac{\varphi\left(q \frac{a_k}{a_j} q^{d_{n+1,k}-d_{n,k}}\right)}{\varphi\left(\hbar \frac{a_k}{a_j} q^{d_{n+1,k}-d_{n,k}}\right)} \frac{\varphi\left(\hbar \frac{a_k}{a_j}\right)}{\varphi\left(q \frac{a_k}{a_j}\right)} \\
&\quad \prod_{i=n+1}^{s-1} \prod_{j=1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi\left(q \frac{a_k}{a_j} q^{d_{i+1,k}-d_{n,k}-(d_{i,j}-d_{n,j})}\right)}{\varphi\left(\hbar \frac{a_k}{a_j} q^{d_{i+1,k}-d_{n,k}-(d_{i,j}-d_{n,j})}\right)} \frac{\varphi\left(\hbar \frac{a_k}{a_j}\right)}{\varphi\left(q \frac{a_k}{a_j}\right)} \\
&\quad \prod_{i=n+1}^s \prod_{j,k=1}^{v_i} \frac{\varphi\left(\hbar \frac{a_k}{a_j} q^{d_{i,k}-d_{n,k}-(d_{i,j}-d_{n,j})}\right)}{\varphi\left(q \frac{a_k}{a_j} q^{d_{i,k}-d_{n,k}-(d_{i,j}-d_{n,j})}\right)} \frac{\varphi\left(q \frac{a_k}{a_j}\right)}{\varphi\left(\hbar \frac{a_k}{a_j}\right)} \prod_{i=n+1}^s \prod_{j=1}^{v_i} z_i^{d_{i,j}-d_{n,j}}
\end{aligned}$$

Next, we substitute $a_i = \hbar^{i-1} q^{d_{n,i}}$ and obtain

$$\begin{aligned}
& \mathbf{V}_p(\mathbf{a}, z_{n+1}, \dots, z_s) \Big|_{a_i = \hbar^{i-1} q^{d_{n,i}}} \\
&= \sum_{d_{i,j}} \prod_{j=1}^{v_n} \prod_{k=1}^{v_{n+1}} \frac{\varphi(q \hbar^{k-j} q^{d_{n+1,k} - d_{n,j}})}{\varphi(\hbar \hbar^{k-j} q^{d_{n+1,k} - d_{n,j}})} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \\
& \prod_{i=n+1}^{s-1} \prod_{j=1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi(q \hbar^{k-j} q^{d_{i+1,k} - d_{i,j}})}{\varphi(\hbar \hbar^{k-j} q^{d_{i+1,k} - d_{i,j}})} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \\
& \prod_{i=n+1}^s \prod_{j,k=1}^{v_i} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{i,k} - d_{i,j}})}{\varphi(q \hbar^{k-j} q^{d_{i,k} - d_{i,j}})} \frac{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \prod_{i=n+1}^s \prod_{j=1}^{v_i} z_i^{d_{i,j} - d_{n,j}}
\end{aligned}$$

We simplify some of the terms in the above summation:

$$\begin{aligned}
& \prod_{j=1}^{v_n} \prod_{k=1}^{v_{n+1}} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \prod_{i=n+1}^{s-1} \prod_{j=1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \\
& \prod_{i=n+1}^s \prod_{j,k=1}^{v_i} \frac{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \\
&= \prod_{i=n}^{s-1} \prod_{j=1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \prod_{i=n+1}^s \prod_{j,k=1}^{v_i} \frac{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \\
&= \prod_{i=n}^{s-1} \prod_{j=v_{i+1}+1}^{v_i} \prod_{k=1}^{v_{i+1}} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \\
&= \prod_j \prod_{k=1}^{v_{j+1}} \frac{\varphi(\hbar \hbar^{k-j} q^{d_{n,k} - d_{n,j}})}{\varphi(q \hbar^{k-j} q^{d_{n,k} - d_{n,j}})} \\
& \quad v_j - v_{j+1} = 1
\end{aligned}$$

From (10), it is easy to see that all of these φ function terms appear in Ψ . Thus we have shown that

$$\mathbf{V}_\lambda(z) = \sum_{d_{i,j}} \Psi' \prod_{i=-r}^n \prod_{j=1}^{v_i} z_i^{d_{i,j}} \prod_{i=n+1}^s \prod_{j=1}^{v_i} z_i^{d_{n,j}} \mathbf{V}_p(\mathbf{a}, z_{n+1}, \dots, z_s) \Big|_{a_i = \hbar^{i-1} q^{d_{n,i}}}$$

where Ψ' is a product of φ function terms, none of which depend on $d_{i,j}$ for $i > n$. This proves the lemma. \square

4.4 Conclusion of the proof

Let $\square \in \lambda$ be the box with content n and height m . Substituting $z_i = \frac{\zeta_{i-1}}{\zeta_i}$ and applying p_\square to Lemma 1 we have

$$\begin{aligned} p_\square \mathbf{V}_\lambda(\mathbf{z}) &= \sum_{d_{i,j}} \Psi \prod_{i=-r}^n \prod_{j=1}^{v_i} \left(\frac{\zeta_{i-1}}{\zeta_i} \right)^{d_{i,j}} \prod_{i=n+1}^s \prod_{j=1}^{v_i} \left(\frac{\zeta_{i-1}}{\zeta_i} \right)^{d_{n,j}} q^{d_{n,m}} p_\square \mathbf{V}_p(\mathbf{a}, \zeta) \Big|_{a_i = \hbar^{i-1} q^{d_{n,i}}} \\ &= \sum_{d_{i,j}} \Psi \prod_{i=-r}^n \prod_{j=1}^{v_i} \left(\frac{\zeta_{i-1}}{\zeta_i} \right)^{d_{i,j}} \prod_{i=n+1}^s \prod_{j=1}^{v_i} \left(\frac{\zeta_{i-1}}{\zeta_i} \right)^{d_{n,j}} (\tilde{p}_\square \mathbf{V}_p(\mathbf{a}, \zeta)) \Big|_{a_i = \hbar^{i-1} q^{d_{n,i}}} \end{aligned}$$

And so

$$\begin{aligned} T_\lambda^{n,r} \mathbf{V}_\lambda(\mathbf{z}) &= \sum_{d_{i,j}} \Psi \prod_{i=-r}^n \prod_{j=1}^{v_i} \left(\frac{\zeta_{i-1}}{\zeta_i} \right)^{d_{i,j}} \prod_{i=n+1}^s \prod_{j=1}^{v_i} \left(\frac{\zeta_{i-1}}{\zeta_i} \right)^{d_{n,j}} T_r \mathbf{V}_p(\mathbf{a}, \zeta) \Big|_{a_i = \hbar^{i-1} q^{d_{n,i}}} \\ &= \sum_{d_{i,j}} \Psi \prod_{i=-r}^n \prod_{j=1}^{v_i} \left(\frac{\zeta_{i-1}}{\zeta_i} \right)^{d_{i,j}} \\ &\quad \prod_{i=n+1}^s \prod_{j=1}^{v_i} \left(\frac{\zeta_{i-1}}{\zeta_i} \right)^{d_{n,j}} e_r(q^{d_{n,1}}, \dots, \hbar^{v_n-1} q^{d_{n,v_n}}) \mathbf{V}_p(\mathbf{a}, \zeta) \Big|_{a_i = \hbar^{i-1} q^{d_{n,i}}} \\ &= \mathbf{V}_\lambda^{(\tau_{n,r})}(\mathbf{z}) \end{aligned}$$

which concludes the proof.

5 Monodromy of vertex functions

5.1

Let \mathbf{K} be a torus with coordinates $\mathbf{z} = (z_1, \dots, z_n)$. Let $\sigma = (\sigma_1, \dots, \sigma_n) \in \text{cochar}_{\mathbb{R}}(\mathbf{K})$ be a cocharacter. We denote $\mathbf{z}q^\sigma = (z_1 q^{\sigma_1}, \dots, z_n q^{\sigma_n})$ for some $q \in \mathbb{C}^\times$ with $|q| < 1$.

Let us consider scalar q -difference equations (qde) of the form:

$$\Psi(\mathbf{z}q^\sigma) = M_\sigma(\mathbf{z})\Psi(\mathbf{z}), \quad (12)$$

where $\Psi(\mathbf{z})$ denotes a \mathbb{C} -valued function on \mathbf{K} and $M_\sigma(\mathbf{z}) \in \mathbb{Q}(z_1, \dots, z_n)$ satisfies

$$\lim_{q \rightarrow 0} M_\sigma(q^\delta) \neq 0, \infty \quad (13)$$

for generic $\delta \in \text{cochar}_{\mathbb{R}}(\mathbf{K})$. Clearly, the limits above will not change if we scale the cocharacter $\delta \rightarrow l\delta$ for some real $l > 0$. This provides a decomposition

$$\text{Lie}_{\mathbb{R}}(\mathbf{K}) \supset \coprod \mathfrak{C}$$

into a set of cones for which the limits (13) remain the same. The cones \mathfrak{C} are sometimes called *asymptotic zones* of the qde (12).

Definition 8. We say that a function $F(z)$ is analytic in an asymptotic zone \mathfrak{C} if it is given by a power series

$$F(z) = \sum_{\langle d, \mathfrak{C} \rangle > 0} c_d z^d \quad (14)$$

with non-zero radius of convergence. Here, $\langle \cdot, \cdot \rangle$ denotes the natural pairing on characters and cocharacters.

It is convenient to view the asymptotic zones \mathfrak{C} as “infinities” in certain toric compactification $\bar{\mathbf{K}}$ of \mathbf{K} . The closure of each chamber \mathfrak{C} is a strongly convex rational polyhedral cone, and the set of such cones generates a fan Δ . The toric variety $\bar{\mathbf{K}}$ associated to this fan contains \mathbf{K} as a subvariety. See Figure 7 below for an example. The chambers \mathfrak{C} then correspond to certain points $0_{\mathfrak{C}} \in \bar{\mathbf{K}}$. A function $F(z)$ is analytic in an asymptotic zone \mathfrak{C} when (14) is the Taylor series of a function on $\bar{\mathbf{K}}$ holomorphic in a non-zero neighborhood of $0_{\mathfrak{C}}$.




Figure 7: An arrangement of chambers in $\text{Lie}_{\mathbb{R}}(\mathbf{K})$ and the corresponding toric compactification $\bar{\mathbf{K}}$.

We will use the following notation

$$M_{\sigma}(0_{\mathfrak{C}}) := \lim_{q \rightarrow 0} M_{\sigma}(q^{\delta}), \quad \delta \in \mathfrak{C}.$$

Proposition 1. For a chamber \mathfrak{C} there exists a unique solution of (12) of the form:

$$\Psi_{\mathfrak{C}}(\mathbf{z}) = \mathbf{z}^{\frac{\ln(a^\sigma)}{\ln(q)}} F_{\mathfrak{C}}(\mathbf{z}) \quad (15)$$

where $F_{\mathfrak{C}}(\mathbf{z})$ is holomorphic near $0_{\mathfrak{C}}$ with $F_{\mathfrak{C}}(0_{\mathfrak{C}}) = 1$ and $a^\sigma = M_\sigma(0_{\mathfrak{C}})$.

Proof. Substituting (15) to (12) we see that it is consistent near $0_{\mathfrak{C}}$ and coefficients $c_{\mathbf{d}}$ in (14) are fixed uniquely by $c_0 = 1$. \square

We call $F_{\mathfrak{C}}(\mathbf{z})$ the *analytic part* of the solution $\Psi_{\mathfrak{C}}(\mathbf{z})$.

Definition 9. The q -periodic function

$$R_{\mathfrak{C}_1 \leftarrow \mathfrak{C}_2}(\mathbf{z}) = \Psi_{\mathfrak{C}_1}(\mathbf{z}) \Psi_{\mathfrak{C}_2}^{-1}(\mathbf{z})$$

is called the monodromy of the qde from asymptotic zone \mathfrak{C}_2 to zone \mathfrak{C}_1 .

5.2

Now, let $\mathbf{K} = (\mathbb{C}^\times)^{|\lambda|}$ be a Kähler torus of X_λ . From (7) we have:

Proposition 2. The vertex function of X_λ satisfies the q -difference equation:

$$\mathbf{V}_\lambda(z_\square q) = \frac{1 - z_\square}{1 - z_\square \hbar} \mathbf{V}_\lambda(z_\square) \quad (16)$$

This is a special case of the qde's associated with quiver varieties discussed in [OS16].

Each z_\square corresponds to a \mathbf{K} -character σ_\square . The complement of the hyperplanes σ_\square^\perp

$$\mathrm{Lie}_{\mathbb{R}}(\mathbf{K}) \setminus \{\sigma_\square^\perp : \square \in \lambda\} = \coprod \mathfrak{C} \quad (17)$$

is the union of chambers defining asymptotic zones of (16). The corresponding compactification $\bar{\mathbf{K}}$ is sometimes called *Kähler moduli space* of the quiver variety X_λ . The vertex function (7) is the unique solution of the qde (16) for the chamber defining the positive stability condition:

$$\mathfrak{C}_+ = \{\theta \in \mathrm{Lie}_{\mathbb{R}}(\mathbf{K}) : \langle \theta, \sigma_\square \rangle > 0, \square \in \lambda\}.$$

The solutions for other chambers are easy to describe.

Proposition 3. The solution of (16) corresponding to a chamber \mathfrak{C} equals:

$$\Psi_{\lambda, \mathfrak{C}}(\mathbf{z}) = z^{\frac{\ln(\hbar_{\mathfrak{C}})}{\ln(q)}} \prod_{\substack{\square \in \lambda, \\ \langle \sigma_{\square}, \mathfrak{C} \rangle > 0}} \prod_{i=0}^{\infty} \frac{1 - \hbar z_{\square} q^i}{1 - z_{\square} q^i} \prod_{\substack{\square \in \lambda, \\ \langle \sigma_{\square}, \mathfrak{C} \rangle < 0}} \prod_{i=1}^{\infty} \frac{1 - z_{\square}^{-1} q^i}{1 - z_{\square}^{-1} \hbar^{-1} q^i} \quad (18)$$

where $z^{\frac{\ln(\hbar_{\mathfrak{C}})}{\ln(q)}}$ denotes the weight determined by the transformations

$$z_{\square} \rightarrow z_{\square} q \quad \Longrightarrow \quad z^{\frac{\ln(\hbar_{\mathfrak{C}})}{\ln(q)}} \rightarrow z^{\frac{\ln(\hbar_{\mathfrak{C}})}{\ln(q)}} \hbar^{p_{\mathfrak{C}}}$$

where $p_{\mathfrak{C}} = |\{\square \in \lambda : \langle \sigma_{\square}, \mathfrak{C} \rangle < 0\}| \in \mathbb{Z}$.

Proof. It is elementary to check that this function solves (16). It is also clear that the analytic part of $\Psi_{\lambda, \mathfrak{C}}(\mathbf{z})$ is holomorphic near $0_{\mathfrak{C}} \in \overline{K}$ \square

5.3

Let us discuss the geometric meaning of the solutions described in Proposition 3. Recall that to define a Nakajima quiver variety one needs to specify a *stability* condition for the geometric invariant theory quotient. The stability condition is specified by a choice of $\theta \in \text{Lie}_{\mathbb{R}}(\mathbf{K})$. The corresponding quiver variety changes (by a symplectic flop) when θ crosses certain hyperplanes in $\text{Lie}_{\mathbb{R}}(\mathbf{K})$. The complement of these hyperplanes divides the space into a set of chambers \mathfrak{C} . The quiver variety obtained from a choice of cocharacter depends only on the chamber that contains it.

The vertex function for a quiver variety $X_{\mathfrak{C}}$ formed by a stability condition from a chamber \mathfrak{C} is given by a power series over the degrees of effective curves, which are given by \mathfrak{C} :

$$\mathbf{V}_{X_{\mathfrak{C}}}(\mathbf{z}) = 1 + \sum_{\langle \mathbf{d}, \mathfrak{C} \rangle > 0} c_{\mathbf{d}} \mathbf{z}^{\mathbf{d}}$$

The relation between the vertex functions for different stability conditions is described by the following idea:

Conjecture 1. The quantum difference equations for quiver varieties are invariant under a change of the stability condition.

In the case of equivariant cohomology, this conjecture was proven in [MO12].

This conjecture implies that the vertex functions for $X_{\mathfrak{C}}$ are solutions of *the same qde (independent of \mathfrak{C}), analytic near different points of the Kähler moduli space.*

Let $X_{\lambda, \mathfrak{C}}$ be the quiver variety obtained from the dimension data given by λ with the stability condition \mathfrak{C} .

Corollary 2. If Conjecture 1 holds then the analytic part of (18) is the vertex function of the quiver variety $X_{\lambda, \mathfrak{C}}$.

In the rest of this paper we assume that Conjecture 1 holds.

5.4

Proposition 4. The monodromy of (16) from \mathfrak{C}_2 to \mathfrak{C}_1 equals:

$$R_{\mathfrak{C}_1 \leftarrow \mathfrak{C}_2}(\mathbf{z}) = (-\hbar^{1/2})^{p_{\mathfrak{C}_1} - p_{\mathfrak{C}_2}} \mathbf{z}^{\frac{\ln(\hbar \mathfrak{e}_1)}{\ln(q)} - \frac{\ln(\hbar \mathfrak{e}_2)}{\ln(q)}} \frac{\prod_{\substack{\square \in \lambda \\ \langle \sigma_{\square}, \mathfrak{C}_1 \rangle < 0}} \vartheta(z_{\square}) \prod_{\substack{\square \in \lambda \\ \langle \sigma_{\square}, \mathfrak{C}_1 \rangle > 0}} \vartheta(z_{\square} \hbar)}{\prod_{\substack{\square \in \lambda \\ \langle \sigma_{\square}, \mathfrak{C}_2 \rangle < 0}} \vartheta(z_{\square}) \prod_{\substack{\square \in \lambda \\ \langle \sigma_{\square}, \mathfrak{C}_2 \rangle > 0}} \vartheta(z_{\square} \hbar)}$$

where

$$\vartheta(z) = (z^{1/2} - z^{-1/2}) \prod_{i=1}^{\infty} (1 - zq^i)(1 - z^{-1}q^i)$$

denotes the odd Jacobi theta function.

Proof. By definition

$$R_{\mathfrak{C}_1 \leftarrow \mathfrak{C}_2}(\mathbf{z}) = \frac{\Psi_{\lambda, \mathfrak{C}_1}(\mathbf{z})}{\Psi_{\lambda, \mathfrak{C}_2}(\mathbf{z})}$$

and the result follows immediately from (18). \square

By Theorem 3, the descendent vertex function differs from vertex function with trivial descendent insertion by a rational function. Taking the analytic parts of (18) we obtain:

Proposition 5.

$$\mathbf{V}_{\lambda, \mathfrak{C}_1}^{(\tau)}(\mathbf{z}) = \mathcal{R}_{\mathfrak{C}_1 \leftarrow \mathfrak{C}_2}(\mathbf{z}) \mathbf{V}_{\lambda, \mathfrak{C}_2}^{(\tau)}(\mathbf{z})$$

where

$$\mathcal{R}_{\mathfrak{e}_1, \mathfrak{e}_2}(\mathbf{z}) = (-\hbar^{1/2})^{p_{\mathfrak{e}_1} - p_{\mathfrak{e}_2}} \frac{\prod_{\substack{\square \in \lambda \\ \langle \sigma_{\square}, \mathfrak{e}_1 \rangle < 0}} \vartheta(z_{\square}) \prod_{\substack{\square \in \lambda \\ \langle \sigma_{\square}, \mathfrak{e}_1 \rangle > 0}} \vartheta(z_{\square} \hbar)}{\prod_{\substack{\square \in \lambda \\ \langle \sigma_{\square}, \mathfrak{e}_2 \rangle < 0}} \vartheta(z_{\square}) \prod_{\substack{\square \in \lambda \\ \langle \sigma_{\square}, \mathfrak{e}_2 \rangle > 0}} \vartheta(z_{\square} \hbar)}. \quad (19)$$

5.5

Let X_{λ}^{\dagger} denote the symplectic dual variety of X_{λ} . As explained in Section 4.7 of [DS19b], $X_{\lambda}^{\dagger} \cong T^*\mathbb{C}^{|\lambda|}$ equipped with an action of the torus $\mathbb{K} \times \mathbb{C}_{\hbar}^{\times}$, where the second factor acts by scaling the symplectic form with weight \hbar . The \mathbb{K} -fixed set of X_{λ}^{\dagger} consists of a single point p (the origin of $\mathbb{C}^{|\lambda|}$). The character of the tangent space was computed in Proposition 4.9.1 of [DS19b]:

$$\text{char}(T_p X_{\lambda}^{\dagger}) = \sum_{\square \in \lambda} z_{\square} + z_{\square}^{-1} \hbar^{-1}. \quad (20)$$

Thus the chambers (17) are the equivariant chambers of the symplectic dual variety¹. In view of this observation the previous proposition can be reformulated in the language of the elliptic stable envelopes [AO16].

Proposition 6.

$$\mathcal{R}_{\mathfrak{e}_1 \leftarrow \mathfrak{e}_2}(\mathbf{z}) = \hbar^{(p_{\mathfrak{e}_1} - p_{\mathfrak{e}_2})/2} \mathcal{R}_{\mathfrak{e}_1, \mathfrak{e}_2}^{X_{\lambda}^{\dagger}}(\mathbf{z}) \quad (21)$$

where $\mathcal{R}_{\mathfrak{e}_1, \mathfrak{e}_2}^{X_{\lambda}^{\dagger}}(\mathbf{z})$ is the elliptic R-matrix of the symplectic dual variety X_{λ}^{\dagger} :

$$\mathcal{R}_{\mathfrak{e}_1, \mathfrak{e}_2}^{X_{\lambda}^{\dagger}}(\mathbf{z}) := \text{Stab}_{\mathfrak{e}_2}^{-1} \circ \text{Stab}_{\mathfrak{e}_1},$$

and $\text{Stab}_{\mathfrak{e}}$ denotes the elliptic stable envelope of X_{λ}^{\dagger} for an equivariant chamber \mathfrak{e} .

Proof. The elliptic stable envelope of X_{λ}^{\dagger} is defined by a set of axioms, see Section 3 in [AO16]. In the case of a finite fixed point set these conditions were explained in Section 2.13 of [Smi19]. The diagonal restrictions of the

¹By definition, these chambers are connected components of the complement of hyperplanes $w^{\perp} \subset \text{Lie}_{\mathbb{R}}(\mathbb{K})$ where w runs over the weights appearing in normal bundles to fixed components, see Section 9.1.2 in [Oko15]

elliptic stable envelope are described there by formula (21). For $X_\lambda^!$ there is only one fixed point and the character of the tangent space is given by (20). Thus,

$$\text{Stab}_{\mathfrak{C}} = \prod_{\substack{\square \in \lambda \\ \langle \sigma_\square, \mathfrak{C} \rangle < 0}} \vartheta(z_\square) \prod_{\substack{\square \in \lambda \\ \langle \sigma_\square, \mathfrak{C} \rangle > 0}} \vartheta\left(\frac{1}{z_\square \hbar}\right) = (-1)^{p_{\mathfrak{C}}} \prod_{\substack{\square \in \lambda \\ \langle \sigma_\square, \mathfrak{C} \rangle < 0}} \vartheta(z_\square) \prod_{\substack{\square \in \lambda \\ \langle \sigma_\square, \mathfrak{C} \rangle > 0}} \vartheta(z_\square \hbar)$$

and the proposition follows from (19). \square

This result generalizes the relation between vertex functions of zero-dimensional varieties and characters of tangent spaces of symplectic dual varieties which we discussed in [DS19a].

6 Characters of tautological bundles over X_λ

6.1

Let $\tau \in \text{char}(G_v)$ which we understand as a symmetric polynomial $\tau(\mathbf{x})$ in the Grothendieck roots $\mathbf{x} = \{x_\square\}_{\square \in \lambda}$ of the tautological bundles.

The associated tautological bundle over the quiver variety $X_{\lambda, \mathfrak{C}}$ defines a Laurent polynomial:

$$\tau(\mathbf{x}_{\mathfrak{C}}) \in K_{\mathbb{C}_\hbar^\times}(X_{\lambda, \mathfrak{C}}) = \mathbb{Q}[\hbar^{\pm 1}]$$

where $\mathbf{x}_{\mathfrak{C}} = \{\hbar^{m_{\mathfrak{C}}(\square)}\}_{\square \in \lambda}$ for some $m_{\mathfrak{C}}(\square) \in \mathbb{Z}$.

For the positive stability condition θ_+ the integers $m_{\mathfrak{C}}(\square)$ are easy to compute, see Section 2.6 in [DS19b]. For a general \mathfrak{C} , we can analyze the stability conditions as described in Proposition 5.1.5 of [Gin12]. This is, however, an indirect description, and it is not obvious how to compute the characters $\mathbf{x}_{\mathfrak{C}}$ in this approach. In this section, we derive an explicit combinatorial formula for $\mathbf{x}_{\mathfrak{C}}$ from the properties of vertex functions discussed previously.

6.2

The computation of $\mathbf{x}_{\mathfrak{C}}$ is based on the following two simple results.

Proposition 7. Let $\hat{\mathbf{V}}_{\lambda, \mathfrak{C}_1}^{(\tau)}(\mathbf{z})$ and $\hat{\mathbf{V}}_{\lambda, \mathfrak{C}_2}^{(\tau)}(\mathbf{z})$ be capped vertex functions of the quiver varieties $X_{\lambda, \mathfrak{C}_1}$, and $X_{\lambda, \mathfrak{C}_2}$, respectively. Then

$$\hat{\mathbf{V}}_{\lambda, \mathfrak{C}_1}^{(\tau)}(\mathbf{z}) = \hat{\mathbf{V}}_{\lambda, \mathfrak{C}_2}^{(\tau)}(\mathbf{z}) \in \mathbb{Q}(\mathbf{z}, q, \hbar)$$

Proof. From the Proposition 5 we see that:

$$\hat{\mathbf{V}}_{\lambda, \mathfrak{e}_1}^{(\tau)}(\mathbf{z}) = \frac{\mathbf{V}_{\lambda, \mathfrak{e}_1}^{(\tau)}(\mathbf{z})}{\mathbf{V}_{\lambda, \mathfrak{e}_1}^{(1)}(\mathbf{z})} = \frac{\mathbf{V}_{\lambda, \mathfrak{e}_2}^{(\tau)}(\mathbf{z})}{\mathbf{V}_{\lambda, \mathfrak{e}_2}^{(1)}(\mathbf{z})} = \hat{\mathbf{V}}_{\lambda, \mathfrak{e}_2}^{(\tau)}(\mathbf{z})$$

For the positive stability condition the capped vertex function is described by Theorem 3 and is obviously rational. \square

Proposition 8. The capped vertex function $\hat{\mathbf{V}}_{\lambda, \mathfrak{e}}^{(\tau)}(\mathbf{z})$ has the following expansion:

$$\hat{\mathbf{V}}_{\lambda, \mathfrak{e}}^{(\tau)}(\mathbf{z}) = \tau(\mathbf{x}_{\mathfrak{e}}) + \sum_{\langle \mathbf{d}, \mathfrak{e} \rangle > 0} c_{\mathbf{d}} \mathbf{z}^{\mathbf{d}}$$

where $c_{\mathbf{d}} \in \mathbb{Q}(q, \hbar)$.

Proof. By definition, the capped vertex function is a power series over degrees of quasimaps in the effective cone determined by \mathfrak{e} :

$$\hat{\mathbf{V}}_{\lambda, \mathfrak{e}}^{(\tau)}(\mathbf{z}) = \sum_{\langle \mathbf{d}, \mathfrak{e} \rangle \geq 0} c_{\mathbf{d}} \mathbf{z}^{\mathbf{d}}$$

The quasimaps of degree zero are trivial which means $\mathbf{QM}_{\text{nonsing } p_2}^0 = X_{\lambda}$, and thus the degree zero coefficient in this expansion is $\tau(\mathbf{x}_{\mathfrak{e}}) \in K_{\mathbb{T}}(X_{\lambda})$. \square

Corollary 3. The capped vertex function for $X_{\lambda, \mathfrak{e}_+}$ where \mathfrak{e}_+ is the chamber corresponding to θ_+ satisfies

$$\tau(\mathbf{x}_{\mathfrak{e}}) = \hat{\mathbf{V}}_{\lambda}^{(\tau)}(0_{\mathfrak{e}}).$$

Proof. This follows immediately from Proposition 7 and Proposition 8. \square

6.3

Let σ_{\square} be as before. Let $\sigma_{\square, \square'}$ be the \mathbb{K} -character corresponding to $\frac{\zeta_{\square}}{\zeta_{\square'}}$. We define $m_{\mathfrak{e}}(\square) \in \mathbb{Z}$ by

$$m_{\mathfrak{e}}(\square) = |\{\square' \in C_{\lambda}(\square) : \square' \neq \square, \langle \sigma_{\square, \square'}, \mathfrak{e} \rangle > 0\}| - |\{\square' \in S_{\lambda}(\square) : \langle \sigma_{\square'}, \mathfrak{e} \rangle < 0\}|.$$

Theorem 6.

$$\tau(\mathbf{x}_{\mathfrak{e}}) = \tau(\{\hbar^{m_{\mathfrak{e}}(\square)}\}).$$

Proof. The character $\tau(\mathbf{x}_{\mathfrak{C}})$ is given by substitution of the Grothendieck roots by some monomials $x_{\square} \rightarrow \hbar^{m_{\mathfrak{C}}(\square)}$. As $\tau(\mathbf{x})$ are symmetric in the Grothendieck roots of the tautological bundles, it is enough to prove the proposition for the n -th tautological bundles corresponding to the polynomials:

$$\tau_n(\mathbf{x}) = \sum_{\square \in C_{\lambda}(n)} x_{\square}.$$

By definition, for $\theta \in \mathfrak{C}$

$$\hat{\mathbf{V}}_{\lambda}^{(\tau_n(\mathbf{x}))}(0_{\mathfrak{C}}) = \lim_{w \rightarrow 0} \hat{\mathbf{V}}_{\lambda}^{(\tau_n(\mathbf{x}))}(\theta(w))$$

where w denotes coordinate on \mathbb{C}^{\times} and $\hat{\mathbf{V}}_{\lambda}^{(\tau_n(\mathbf{x}))}(\mathbf{z})$ denotes the capped vertex function for $X_{\lambda, \mathfrak{C}_+}$. By Corollary 1, we have

$$\lim_{w \rightarrow 0} \hat{\mathbf{V}}_{\lambda}^{(\tau_n(\mathbf{x}))}(\theta(w)) = \sum_{\square \in C_{\lambda}(n)} \prod_{\substack{\square' \in C_{\lambda}(n) \\ \square' \neq \square \\ \frac{\zeta_{\square}}{\zeta_{\square'}} \rightarrow 0}} \hbar \prod_{\substack{\square' \in S_{\lambda}(\square) \\ z_{\square'} \rightarrow \infty}} \frac{1}{\hbar}$$

where $z_{\square'} \rightarrow \infty$, $\frac{\zeta_{\square}}{\zeta_{\square'}} \rightarrow 0$ describes the behaviour of these weights as $w \rightarrow 0$. These limits depend only on \mathfrak{C} and these conditions are equivalent to $\langle \sigma_{\square'}, \mathfrak{C} \rangle < 0$ and $\langle \sigma_{\square, \square'}, \mathfrak{C} \rangle > 0$ respectively. \square

References

- [AO16] Mina Aganagic and Andrei Okounkov. *Elliptic stable envelopes*. Apr. 2016. arXiv: 1604.00423v4 [math.AG].
- [AO17] Mina Aganagic and Andrei Okounkov. “Quasimap counts and Bethe eigenfunctions”. In: *Mosc. Math. J.* 17 (2017), pp. 565–600.
- [CKM14] Ionuț Ciocan-Fontanine, Bumsig Kim, and Davesh Maulik. “Stable quasimaps to GIT quotients”. In: *J. Geom. Phys.* 75 (2014), pp. 17–47.
- [DS19a] Hunter Dinkins and Andrey Smirnov. “Characters of tangent spaces at torus fixed points and 3d-mirror symmetry”. In: *Lett. Math. Phys.* (Aug. 2019), to appear. arXiv: 1908.01199v2 [math.AG].

- [DS19b] Hunter Dinkins and Andrey Smirnov. “Quasimaps to zero-dimensional A_∞ -quiver varieties”. In: *Int. Math. Res. Not. IMRN* (Dec. 2019), to appear. arXiv: 1912.04834 [math.AG].
- [Gin12] Victor Ginzburg. “Lectures on Nakajima’s quiver varieties”. In: *Geometric methods in representation theory. I*. Vol. 24. Sémin. Congr. Soc. Math. France, Paris, 2012, pp. 145–219.
- [Kor+17] Peter Koroteev et al. *Quantum K-theory of Quiver Varieties and Many-Body Systems*. May 2017. arXiv: 1705.10419 [math.AG].
- [Kor18] Peter Koroteev. *A-type Quiver Varieties and ADHM Moduli Spaces*. May 2018. arXiv: 1805.00986 [math.AG].
- [KS] Yakov Kononov and Andrey Smirnov. *Pursuing quantum difference equations II, in preparation*.
- [Mac79] I. G. Macdonald. *Symmetric functions and Hall polynomials*. English. Clarendon Press ; Oxford University Press Oxford : New York, 1979. ISBN: 0198535309.
- [MO12] Davesh Maulik and Andrei Okounkov. “Quantum Groups and Quantum Cohomology”. In: *Astérisque* 408 (Nov. 2012).
- [Nak98] Hiraku Nakajima. “Quiver varieties and Kac-Moody algebras”. In: *Duke Math. J.* 91.3 (1998), pp. 515–560.
- [Oko15] Andrei Okounkov. *Lectures on K-theoretic computations in enumerative geometry*. 2015. arXiv: 1512.07363 [math.AG].
- [OS16] Andrei Okounkov and Andrey Smirnov. *Quantum difference equation for Nakajima varieties*. 2016. arXiv: 1602.09007 [math-ph].
- [PP13] R. Pandharipande and A. Pixton. “Descendents on local curves: rationality”. In: *Compos. Math.* 149.1 (2013), pp. 81–124.
- [PSZ16] Petr Pushkar, Andrey Smirnov, and Anton Zeitlin. “Baxter Q-operator from quantum K-theory”. In: *Adv. Math.* 360 (Dec. 2016).
- [Smi19] Andrey Smirnov. “Elliptic stable envelope for Hilbert scheme of points in the plane”. In: *Selecta Math.* 26 (Dec. 2019).

Hunter Dinkins
Department of Mathematics,
University of North Carolina at Chapel Hill,
Chapel Hill, NC 27599-3250, USA
hdinkins@live.unc.edu

Andrey Smirnov
Department of Mathematics,
University of North Carolina at Chapel Hill,
Chapel Hill, NC 27599-3250, USA
Steklov Mathematical Institute
of Russian Academy of Sciences,
Gubkina str. 8, Moscow, 119991, Russia
asmirnov@email.unc.edu