

Quantum Photovoltaic Cells Driven by Photon Pulses

Sangchul Oh,^{1,*} Jungjun Park,² and Hyunchul Nha³

¹*Qatar Environment and Energy Research Institute,*

Hamad Bin Khalifa University, Qatar Foundation, P.O.Box 5825, Doha, Qatar

²*Korea Institute for Advanced Study, 85 Hoegiro, Dongdaemun-gu Seoul 02455, Republic of Korea*

³*Texas A&M University at Qatar, P.O.Box 23874, Education City, Doha, Qatar*

(Dated: March 14, 2024)

We investigate the quantum thermodynamics of two quantum systems, a two-level system and a four-level quantum photocell, each driven by photon pulses as a quantum heat engine. We set these systems to be in thermal contact only with a cold reservoir while the heat (energy) source, conventionally given from a hot thermal reservoir, is supplied by a sequence of photon pulses. The dynamics of each system is governed by a coherent interaction due to photon pulses in terms of the Jaynes-Cummings Hamiltonian together with the system-bath interaction described by the Lindblad master equation. We calculate the thermodynamic quantities for the two-level system and the quantum photocell including the change in system energy, power delivered by photon pulses, power output to an external load, heat dissipated to a cold bath, and entropy production. We thereby demonstrate how a quantum photocell in the cold bath can operate as a continuum quantum heat engine with the sequence of photon pulses continuously applied. We specifically introduce the power efficiency of the quantum photocell in terms of the ratio of output power delivered to an external load with current and voltage to the input power delivered by the photon pulse. Our study indicates a possibility that a quantum system driven by external fields can act as an efficient quantum heat engine under non-equilibrium thermodynamics.

I. INTRODUCTION

Thermodynamics deals with the evolution of systems, usually in contact with reservoirs, describing the dynamics under universal laws independent of microscopic details. Among its four laws, the second law dictates the total entropy of a closed system can never decrease over time and that the closed system spontaneously evolves toward the state with maximum entropy. One of the possible statements about the second law of thermodynamics is to set the upper bound on the efficiency of heat engines. Heat engines convert heat energy, which typically flows from a hot source to a cold sink, to mechanical energy or chemical energy. The efficiency of energy conversion is defined by the ratio of the work output to the amount of heat energy input. The ultimate efficiency of the heat engine is known in equilibrium thermodynamics to be determined only by temperatures of hot and cold heat baths, T_h and T_c , respectively, i.e. $\eta = 1 - T_c/T_h$, the so-called the Carnot limit.

Photovoltaic cells (or solar cells) and photosynthesis, just like classical heat engines, convert photon energy from the sun into electric energy and chemical energy, respectively. The upper limit of the efficiency of p - n junction solar cells with an energy bandgap is known as the Shockley-Queisser limit [1]. The key assumptions in deriving the Shockley-Queisser limit are (i) photons with energies less than the bandgap are not utilized, (ii) a photon with energy greater than the bandgap produces only one electron-hole pair and (iii) only the radiative recombination of electron-hole pairs is considered. While

the non-radiative loss may be minimized by the manufacturing technology, radiative recombination is the intrinsic energy loss governed by the law of physics. Assuming the sun and the solar cell are described as black-bodies with temperature $T_s = 6000$ K and $T_c = 300$ K, respectively, the maximum efficiency is about 30% for a solar cell with a bandgap of 1.137 eV [1]. Shockley and Queisser [1] also showed that the maximum efficiency of a single p - n junction solar cell would be approximately 44% around 1.137 eV if there is no radiative recombination loss.

Recently, many theoretical studies have suggested that noise-induced quantum coherence [2, 3], Fano-induced coherence [4] or delocalized quantum states of interacting dipoles [5–8] can reduce the radiative recombination loss of a solar cell, thus enhancing the efficiency of solar cells. The same idea is applied to the photosynthetic complex [9–12]. Most of these studies employ the donor-acceptor quantum photocell model where the donor is in thermal equilibrium with a hot bath, i.e., the sun at 5800 K and the acceptor is at room temperature. The photocell operating while continuously contacting both heat reservoirs is called a continuous quantum heat engine [13, 14]. A typical example of the continuous quantum heat engine is Scovil and Schulz-Dubois' three-level masers whose efficiency achieves the Carnot efficiency [15]. By solving the master equation for the quantum photocell, it was shown that the noise-induced quantum coherence or the dark state of the donor may enhance the power output. However, it was not clear whether the efficiency of the photocell could be enhanced by the quantum effect. Some works assumed that the mean photon number of the hot thermal bath could be $\bar{n} = 60000$, but the mean photon number of the sun at energy 1.8 eV as a black body at 5800 K is only about

* soh@hbku.edu.qa

$\bar{n} = 0.037$ [16, 17]. To address this issue, Ref. [16] introduced the pumping term and showed that the dark state could enhance power output but not its efficiency.

In this paper, we explore another form of quantum heat engine. We consider two quantum systems, a two-level system and a donor-acceptor quantum photocell, and investigate their quantum dynamics under coherent driving and system-bath interaction. Each quantum system is in thermal contact only with a cold reservoir but not with a hot reservoir. Instead, they are driven by a sequence of photon pulses that supply input energy to the systems, which is conventionally done by a hot reservoir. The photon pulses represent the stream of energy source to the system and may thus remove the unrealistic assumption of high mean photon number of the sun by the previous works. We solve the time-dependent Markov Lindblad master equation and investigate the thermodynamic quantities such as the change in energy, the heat dissipation to the cold bath, the power delivered by the photon pulse and the entropy generation. Specifically we introduce the power efficiency of the quantum photocell in terms of the ratio of output power delivered to an external load to input power delivered by the photon pulses.

This paper is organized as follows. In Sec. II, we review briefly the quantum dynamics and quantum thermodynamics of an open quantum system based on the master equation approach. In Sec. III, we examine a two-level system in a cold bath driven by photon pulses, and examine how the energy, heat current, and entropy change. In Sec. IV a quantum photovoltaic cell with donor and acceptor driven by repeated photon pulses is considered. We calculate the quantum thermodynamic quantities and the power output by the sequence of the photon pulses together with engine efficiency. Finally, in Sec. V we summarize our results with some discussion.

II. QUANTUM THERMODYNAMICS OF OPEN QUANTUM SYSTEMS

We start with a brief review of quantum thermodynamics of an open quantum system which exchanges energy and entropy with its environment. The equations presented in this section will be applied to those examples in next two sections III and IV. As usual, we assume the Born-Markov approximations: a weak interaction between an open quantum system and the environment, and the extremely short correlation time of the environment, i.e., no memory effect. The density operator $\rho(t)$ of the quantum system with a slowly varying time-dependent Hamiltonian obeys the Lindblad-Gorini-Kossakowski-Sudarshan (LGKS) master equation [18–22]

$$\frac{d}{dt}\rho(t) = \mathcal{L}[\rho(t)] = -\frac{i}{\hbar}[H_S(t), \rho(t)] + \mathcal{D}[\rho(t)], \quad (1)$$

where $H_S(t) = H_0 + H_1(t)$ is the Hamiltonian of the system. Here H_0 represents a time-independent un-

perturbed Hamiltonian and $H_1(t)$ an external time-dependent perturbation. The decoherence and the dissipation of the open quantum system due to environmental interaction are described by the non-unitary operator

$$\mathcal{D}[\rho] = \sum_k \left(2L_k \rho L_k^\dagger - L_k^\dagger L_k \rho - \rho L_k^\dagger L_k \right), \quad (2)$$

where L_k are the Lindblad operators determined according to the type of interaction.

From the solution $\rho(t)$ of Eq. (1), one can calculate the quantum thermodynamic quantities. The first law of classical thermodynamics states the energy conservation, $dE = \delta Q + \delta W$. The time-dependent internal energy of the system is given by $E(t) = \text{tr} \{ \rho(t) H_S(t) \}$. Its derivative with respect to time gives rise to the first law of quantum thermodynamics [23–25]

$$\frac{d}{dt}E(t) = \dot{Q}(t) + \dot{W}(t) = J(t) + P(t). \quad (3)$$

Here $J(t)$ is the heat current from the environment into the system

$$J(t) \equiv \dot{Q}(t) = \text{tr} \left(\frac{d\rho(t)}{dt} H_S(t) \right), \quad (4)$$

and $P(t)$ is the power delivered to system by external forces,

$$P(t) \equiv \dot{W}(t) = \text{tr} \left(\rho(t) \frac{dH_S}{dt} \right). \quad (5)$$

Since H_0 is the time-independent Hamiltonian, the power can be written as $P(t) = \text{tr} \left(\rho(t) \frac{dH_1(t)}{dt} \right)$. The change in energy of the system for finite time can be obtained by integrating the heat current and the power as

$$\Delta E(t) = Q(t) + W(t) = \int_0^t J(s) ds + \int_0^t P(s) ds. \quad (6)$$

The second law of thermodynamics describes the irreversibility of dynamics, where the entropy plays a key role. The von Neumann entropy $S(t)$ of the system in the state $\rho(t)$ is given by

$$S(t) = -\text{tr} \{ \rho(t) \log \rho(t) \}. \quad (7)$$

The thermodynamic entropy \mathcal{S} is written as $\mathcal{S} = k_B S(t)$ with the Boltzmann constant k_B . The net change in the entropy dS_{net} (entropy production) of the whole system+reservoir can be written in terms of the entropy change of the system, dS , and the entropy flow due to heat from environment to system, dS_e , as

$$dS_{\text{net}} = dS - dS_e. \quad (8)$$

The change in the entropy of the system, dS , over time is written as

$$\frac{dS}{dt} = -\text{tr} \{ \dot{\rho}(t) \log \rho(t) \} \quad (9a)$$

$$= -\text{tr} \{ \mathcal{L}[\rho(t)] \log \rho(t) \}, \quad (9b)$$

where $\text{tr}\{\dot{\rho}(t)\} = 0$ and the quantum Markov master equation (1) are used. The entropy flow dS_e per unit time from the environment into the system is written as

$$\frac{dS_e}{dt} \equiv J_S = \beta \dot{Q}(t) \quad (10a)$$

$$= \beta \text{tr}\{\dot{\rho} H_S(t)\} = \beta \text{tr}\{\mathcal{L}[\rho(t)] H_S(t)\}, \quad (10b)$$

where $\beta = 1/k_B T$ and T is the temperature of the environment. The net entropy production rate $\sigma(t)$ of the system is given by

$$\sigma(t) = \frac{dS_{\text{net}}}{dt} = \dot{S}(t) - \beta \dot{Q}(t) \geq 0, \quad (11)$$

where $\sigma(t) \geq 0$ comes from the Spohn inequality [23–26]. Eq. (11) may be written as

$$\sigma \equiv -\frac{d}{dt} S(\rho(t) \parallel \rho_{\text{ss}}) \geq 0, \quad (12)$$

where $S(\rho(t) \parallel \rho_{\text{ss}}) \equiv \text{tr}[\rho(t)(\log \rho(t) - \log \rho_{\text{ss}})]$ is the relative entropy of $\rho(t)$ with respect to the stationary state ρ_{ss} , for example, the canonical state of the system, $\rho_{\text{ss}} = \rho_{\beta} = e^{-\beta H_s(t)}/Z$. This is called the second law of non-equilibrium quantum thermodynamics in the weak coupling limit.

III. A TWO-LEVEL SYSTEM DRIVEN BY PHOTON PULSES

As an application of quantum thermodynamics of open quantum systems presented in Sec. II, we first consider a two-level quantum system which is in contact with a cold bath at temperature T_c and driven by repeated photon pulses, as depicted in Fig. 1. The hot thermal bath supplying energy does not have a direct contact with the quantum system. Its role is here replaced by a sequence of photon pulses to the two-level system. We examine how the two-level system absorbs and dissipates energy and generates entropy during this process in order to gain insight into the nonequilibrium dynamics due to photon pulses.

The unperturbed Hamiltonian of the two-level system with energy levels E_0 and E_1 may be written as

$$H_0 = -\frac{\hbar\omega_0}{2} \sigma_z, \quad (13)$$

where $\omega_0 = (E_1 - E_0)/\hbar$ and $\sigma_z = |0\rangle\langle 0| - |1\rangle\langle 1|$. The interaction between a two-level system and incoming photon pulses is described by the Jaynes-Cummings Hamiltonian [27]

$$H_1(t) = i\hbar [g^*(t)\sqrt{\gamma}\sigma_- - g(t)\sqrt{\gamma}\sigma_+], \quad (14)$$

where $\sigma_+ = |1\rangle\langle 0|$ and $\sigma_- = |0\rangle\langle 1|$ are the raising and the lowering operators, respectively. We set $E_1 - E_0 = 1$ eV. Here γ is the Weisskopf-Wigner spontaneous decay rate

$$\gamma = \frac{1}{4\pi\epsilon_0} \frac{4\omega_0^3 d_{01}^2}{3\hbar c^3}, \quad (15)$$




Figure 1. A two-level system with energy levels E_0 and E_1 in contact with a cold thermal bath at T_c is driven by Gaussian photon pulses serving as an energy source in our work.

where d_{01} is the transition dipole moment between the two states $|0\rangle$ and $|1\rangle$. A typical value of γ for an atom or a quantum dot for visible light emission is in the order of nano seconds corresponding to μeV , while the energy at visible frequencies is about eV, i.e., femto seconds. As a numerical calculation becomes demanding with a big difference between these time-scales, we use for our study the values of parameters as listed in Table I. We consider the photon pulses given at peak times t_i as $g(t) = \alpha \sum_i \xi(t; t_i)$ with coherent states having average photon number $\langle n \rangle = |\alpha|^2$ and a Gaussian pulse shape [28–30]

$$\xi(t; t_i) \equiv \left(\frac{\Omega^2}{2\pi}\right)^{1/4} \exp\left[-\frac{\Omega^2(t-t_i)^2}{4} - i\omega_0 t\right], \quad (16)$$

Here $1/\Omega$ is the pulse bandwidth.

Under the Born-Markov approximation, the interaction of a two-level system and the thermal photon bath is recast to the dissipative operator \mathcal{D} acting on the density matrix of the system [21, 31]

$$\begin{aligned} \mathcal{D}_C[\rho] = & \frac{\gamma}{2} (\bar{n}_c + 1) (2\sigma_- \rho \sigma_+ - \sigma_+ \sigma_- \rho + \rho \sigma_+ \sigma_-) \\ & + \frac{\gamma}{2} \bar{n}_c (2\sigma_+ \rho \sigma_- - \sigma_- \sigma_+ \rho + \rho \sigma_- \sigma_+). \end{aligned} \quad (17)$$

Here \bar{n}_c is the mean photon number of the cold bath at the frequency ω_0 in thermal equilibrium of temperature T_c

$$\bar{n}_c = \frac{1}{e^{\hbar\omega_0/k_B T_c} - 1}. \quad (18)$$

As noted in Refs. [31, 32], at optical frequencies and room temperature, the mean photon number \bar{n} is very small and negligible while it has a finite value at microwave frequencies and the room temperature. For example, with $\hbar\omega_0 = \text{eV}$ and $T_c = 300$ K one obtains $\bar{n} \approx 6.5 \times 10^{-31}$. At the optical frequencies, $\hbar\omega_0 = 1.8$ eV and the temperature of the sun as a black body, $T_s = 5800$ K, the mean photon number is $\bar{n} \approx 0.0317$ [16, 32]. With this in mind, Eq. (17) reduces to

$$\mathcal{D}_C[\rho] \approx \frac{\gamma}{2} (2\sigma_- \rho \sigma_+ - \sigma_+ \sigma_- \rho + \rho \sigma_+ \sigma_-). \quad (19)$$

With Eqs. (13), (17), and (14), the Lindblad equation for the two-level system, in contact with the cold thermal

Table I. Typical parameters used in this work

Energy gap of the two-level system	$E_1 - E_0 = \hbar\omega_0 = 1.0$ eV
Energy gap of the donor of the quantum photocell	$E_1 - E_0 = \hbar\omega_0 = 1.8$ eV
Energy gap of the acceptor of the quantum photocell	$E_2 - E_3 = 1.6$ eV
Weisskopf-Winger constant	$\hbar\gamma = 1.24$ μ eV ~ 1.0 meV
Phonon decay constant	$\hbar\gamma_{12} = \hbar\gamma_{30} = 12 \sim 120$ meV
Photon number of a pulse	$\langle n \rangle = \alpha ^2 = 1$ or 10
Temperature of the cold bath	$T_c = 300$ K
Width of a Gaussian pulse	$\Omega = \omega_0/4\pi$




Figure 2. (a) The density matrix elements of the two-level system and the sequence of Gaussian photon pulses $g(t)$ are plotted over time. (b) The rate of energy change $\frac{dE(t)}{dt}$ and the power $P(t)$, (c) the heat current $J(t)$ are calculated as functions of time. (d) The energy $E(t)$, the work $W(t)$, the heat transfer $Q(t)$ and the system entropy $S(t)$ are calculated as functions of time. (e) The rate of system entropy change and the entropy production are calculated over time. The parameters are taken as $\langle n \rangle = 1$, $\gamma = 10^{-2}\omega_0$, $\Omega = \omega_0/4\pi$, and $\hbar\omega_0 = 1$ eV.

bath and driven by a Gaussian photon pulse, is given by

$$\frac{d}{dt}\rho(t) = -\frac{i}{\hbar}[H_0 + H_1(t), \rho(t)] + \mathcal{D}_C[\rho(t)]. \quad (20)$$

The quantum dynamics and the quantum thermodynam-




Figure 3. (a) The density matrix elements of the two-level system and the sequence of Gaussian photon pulses $g(t)$ are plotted over time. (b) The rate of energy change $\frac{dE(t)}{dt}$ and the power $P(t)$, (c) the heat current $J(t)$ are calculated as functions of time. (d) The energy $E(t)$, the work $W(t)$, the heat transfer $Q(t)$ and the system entropy $S(t)$ are calculated as functions of time. (e) The rate of system entropy change and the entropy production are calculated over time. The parameters are $\langle n \rangle = 10$, $\gamma = 10^{-2}\omega_0$, $\Omega = \omega_0/4\pi$, and $\hbar\omega_0 = 1$ eV.

ics of the two level system are investigated by solving Eq. (20) numerically using the Runge-Kutta method. The parameters used in numerical simulation are shown in Table I.

Fig. 2 and Fig. 3 describe the thermodynamic quanti-




Figure 4. (a) The density matrix elements of the two-level system and the sequence of Gaussian photon pulses $g(t)$, (b) rate of energy change $\frac{dE(t)}{dt}$ and power $P(t)$, (c) heat current $J(t)$, (d) energy $E(t)$, work $W(t)$, heat $Q(t)$ and system entropy $S(t)$, (e) rate of system entropy change and entropy production, all as functions of time. This describes the case of regularly spaced sequence of photon pulses. Parameters: $\langle n \rangle = 1$, $\gamma = 10^{-2}\omega_0$, $\Omega = \omega_0/4\pi$, and $\hbar\omega_0 = 1 \text{ rmeV}$.

ties of the system when a sequence of Gaussian photon pulses are applied at regular interval ($g(t)$): green curves in (a)) with $\langle n \rangle = 1$ and $\langle n \rangle = 10$, respectively. Fig. 2 (a) shows the time-evolution of the density matrix elements and the sequence of Gaussian photon pulses, which is first applied around the peak time $\omega_0 t/2\pi = 50$ with $\langle n \rangle = 1$. The initial state of the two level system is assumed to be in a superposed state $|\psi(0)\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$. As shown in Fig. 2 (a), the superposed state decays to the ground state, i.e., the system becomes in thermal equilibrium with the cold thermal bath before the photon pulse is applied. When the Gaussian photon pulse is first applied around $\omega_0 t/2\pi = 50$, the system gets excited and then becomes decayed into the ground state after the pulse is gone. This process is repeated according to each Gaussian pulse.

With regard to the first law of quantum thermodynamics, Fig. 2 (b) and (c) plot the rate of energy change, the power and the heat current. The heat current $J(t)$ from the environment to the system is always negative. This means the excited state of the two-level system releases its energy to the cold bath. In contrast, the power $P(t)$




Figure 5. (a) The density matrix elements of the two-level system and the sequence of Gaussian photon pulses $g(t)$, (b) rate of energy change $\frac{dE(t)}{dt}$ and power $P(t)$, (c) heat current $J(t)$, (d) energy $E(t)$, work $W(t)$, heat $Q(t)$ and system entropy $S(t)$, (e) rate of system entropy change and entropy production, all as functions of time. This describes the case of irregularly spaced sequence of photon pulses. Parameters: $\langle n \rangle = 1$, $\gamma = 10^{-2}\omega_0$, $\Omega = \omega_0/4\pi$, and $\hbar\omega_0 = 1 \text{ eV}$.

and $\dot{E}(t)$ oscillates out of phase while the photon pulse is applied. Fig. 2 (d) shows the entropy $S(t)$ of the two-level system together with its energy $E(t)$, the work done $W(t)$ on the system and the heat transfer $Q(t)$. Fig. 2 (e) shows the entropy production $\sigma(t) = \dot{S}(t) - \beta\dot{Q}$ as a function of time in relation to the second law of thermodynamics. The entropy production $\sigma(t)$ is always positive confirming the second law.

In Fig. 3, we see more oscillatory behaviors in the quantities due to a stronger photon pulse with $\langle n \rangle = 10$ than those with $\langle n \rangle = 1$ in Fig. 2. Nevertheless, the overall trend is similar to that explained above for Fig. 2.

We now examine how the quantum thermodynamic quantities depend on the temporal shape of Gaussian pulse sequence. In Figs. 4 and 5, we plot the same quantities as those in Figs. 2 and 3, but compare two cases, that is, regularly spaced (Fig. 4) and irregularly spaced (Fig. 5) sequence of Gaussian pulses with the same mean number $\langle n \rangle = |\alpha|^2 = 1$. As described by the curve $g(t)$, the peak times t_i 's in Eq. (16) are regularly (not regularly) spaced in the left (right) panel. In both cases, the overall trend of the thermodynamic quantities are

similar to that explained for Fig. 2 while the actual response of the system does depend on the temporal shape of the pulse sequence. Remarkably, we see that the output power $P(t)$ (blue curve in (b)) and the accumulated work $W(t)$ (blue curve in (d)) depend on the temporal shape of the incoming pulses even with the same $|\alpha|^2 = 1$, which can have implications for practical photocell operation. In particular, we find that the case of regular sequencing of pulses yield a higher value of work.

IV. QUANTUM PHOTOCELL DRIVEN BY PHOTON PULSES

For a quantum heat engine, let us now consider a quantum photovoltaics cell driven by photon pulses, as shown in Fig 6. The quantum photocell we consider is a 4-level quantum system composed of a donor and an acceptor. In 1959, Scovil and Du-Bois [15] consider the 3-level system as the simplest quantum heat engine where one part of the 3-level system is in thermal equilibrium with a hot bath and the other part with a cold bath. Many previous studies [2–10] took a similar assumption that the donor of the quantum photocell is in contact with the hot bath, i.e., the sun, and the acceptor is in thermal contact with the cold bath. In contrast to the previous works, we assume that the quantum photocell is in thermal contact only with the cold bath. In our previous work [16], the pumping term was introduced in the Lindblad master equation to describe the energy flow from the hot bath. Here the input energy is supplied by the sequence of incoming photon pulses.

The cyclic operation of the quantum photocell can be performed with the sequence as follows: (i) The donor absorbs incoming photons and the electron becomes excited with the transition from the ground state $|0\rangle$ to the excited state $|1\rangle$. (ii) The phonon vibration makes the excited electron at the donor transfer to the acceptor state $|2\rangle$. (iii) The acceptor is coupled to an external load and the current flow (electric work) is represented by the transition decay from the state $|2\rangle$ to the state $|3\rangle$. (iv) The electron in the state $|3\rangle$ of the acceptor returns to the ground state $|0\rangle$ of the donor by a vibrational or non-radiative decay.

The unperturbed Hamiltonian of the quantum photocell with 4-levels is written as

$$H_0 = -E_0|0\rangle\langle 0| - E_1|1\rangle\langle 1| - E_2|2\rangle\langle 2| - E_3|3\rangle\langle 3|. \quad (21)$$

Similar to Eq. (14), the interaction of the donor of the photocell with the incoming photon pulses is again described by the Jaynes-Cummings Hamiltonian

$$H_1(t) = i\hbar [g^*(t)\sqrt{\gamma}\sigma_- - g(t)\sqrt{\gamma}\sigma_+], \quad (22)$$

where $\sigma_+ = |1\rangle\langle 0|$ and $\sigma_- = |0\rangle\langle 1|$. Same as Eq. (17), the interaction of the donor of the quantum photocell with the cold thermal bath is represented by the Lindblad




Figure 6. Schematic diagram of a donor-acceptor photocell. γ_{01} is the spontaneous decay due to the coupling with the cold thermal bath. γ_{21} and γ_{03} are the transfer rate between the donor and the acceptor. Γ stands for the external load or an electrical resistance.

operator,

$$\mathcal{D}_c[\rho] = \frac{\gamma_{01}}{2} (\bar{n}_c + 1) (2\sigma_- \rho \sigma_+ - \sigma_+ \sigma_- \rho + \rho \sigma_+ \sigma_-) + \frac{\gamma_{01}}{2} \bar{n}_c (2\sigma_+ \rho \sigma_- - \sigma_- \sigma_+ \rho + \rho \sigma_- \sigma_+). \quad (23)$$

The electron transfer between the states $|1\rangle$ and $|2\rangle$ and that between the state $|3\rangle$ and $|0\rangle$ are described by the Lindblad operator $\mathcal{D}_{\text{ph}}[\rho] = \mathcal{D}_{\text{ph}}^{(1,2)}[\rho] + \mathcal{D}_{\text{ph}}^{(3,0)}[\rho]$, where

$$\mathcal{D}_{\text{ph}}^{(i,j)}[\rho] = \frac{\gamma_{ij}}{2} (\bar{n}_{\text{ph}} + 1) (2L_{ij}\rho L_{ij}^\dagger - L_{ij}^\dagger L_{ij}\rho + \rho L_{ij}^\dagger L_{ij}) + \frac{\gamma_{ij}}{2} \bar{n}_{\text{ph}} (2L_{ij}^\dagger \rho L_{ij} - L_{ij} L_{ij}^\dagger \rho + \rho L_{ij} L_{ij}^\dagger). \quad (24)$$

Here γ_{12} and γ_{30} represent the transition rate between $|1\rangle$ and $|2\rangle$ and between $|3\rangle$ and $|0\rangle$, respectively. $L_{ij} = |i\rangle\langle j|$ and $L_{ij}^\dagger = |j\rangle\langle i|$ are the lowering and raising operators, respectively. \bar{n}_{ph} is the phonon occupation number at $\hbar\omega = E_1 - E_2 = E_3 - E_0$ and $T_c = 300$ K. The work done by the quantum photocell to the external load is described by the ohmic dissipation

$$\mathcal{D}_{\text{ohm}}[\rho] = \frac{\Gamma}{2} (2L_3\rho L_3^\dagger - L_3^\dagger L_3\rho + \rho L_3^\dagger L_3), \quad (25)$$

where $L_3 = |3\rangle\langle 2|$. Here Γ represent the conductance of the external load and may be changed from zero corresponding to the open circuit and to a big number representing the short-circuit of the quantum photocell. With Eqs. (21), (22), (23), (24), (25), the LGKS equation for the quantum photocell is written as

$$\frac{d}{dt}\rho(t) = -\frac{i}{\hbar} [H_0 + H_1(t), \rho(t)] + \mathcal{D}_c[\rho] + \mathcal{D}_{\text{ph}}^{(1,2)}[\rho] + \mathcal{D}_{\text{ph}}^{(3,0)}[\rho] + \mathcal{D}_{\text{ohm}}[\rho]. \quad (26)$$

Since the quantum photocell has no direct interaction Hamiltonian between the donor and the acceptor, the Hamiltonian $H_S(t) = H_0 + H_1(t)$ of the quantum photocell can be written as the sum of the time-dependent donor Hamiltonian H_D and the time-independent acceptor Hamiltonian H_A ,

$$H_S(t) = H_D(t) + H_A, \quad (27)$$

where $H_D(t) \equiv -E_0|0\rangle\langle 0| - E_1|1\rangle\langle 1| + H_1(t)$ and $H_A \equiv -E_2|2\rangle\langle 2| - E_3|3\rangle\langle 3|$. This partition makes it possible to express some quantum thermodynamic quantities as the sum of the donor and acceptor parts. The energy of the quantum photocell is given by the sum of the energies of the donor and acceptor

$$E(t) = \text{tr} \{ \rho(t) H_S(t) \} = E_D(t) + E_A(t), \quad (28)$$

where the donor energy $E_D(t)$ and the acceptor energy $E_A(t)$ are given by

$$E_D(t) = \text{tr}_D \{ \rho_D(t) H_D(t) \}, \quad (29a)$$

$$E_A(t) = \text{tr}_A \{ \rho_A(t) H_A \}, \quad (29b)$$

respectively. Here $\rho_A = \text{tr}_D \{ \rho \}$ and $\rho_D = \text{tr}_A \{ \rho \}$ are the density operators of the donor and acceptor, respectively. Since the photon pulse delivers the power only to the donor, the power $P(t)$ is given by the power of the donor

$$P(t) = \text{tr} \{ \rho(t) \dot{H}_S(t) \} = \text{tr}_D \{ \rho_D \dot{H}_1(t) \} \equiv P_D(t). \quad (30)$$

The heat dissipation occurs at the donor and acceptor. Thus the heat current $J(t)$ is written as the sum of the two parts

$$\begin{aligned} J(t) &= \text{tr}_D \{ \dot{\rho}_D(t) H_D(t) \} + \text{tr}_A \{ \dot{\rho}_A(t) H_A \} \\ &= J_D(t) + J_A(t). \end{aligned} \quad (31)$$

Here the heat current of the acceptor J_A may be associated with the power delivered by the quantum photocell to the external load. Finally, the entropy of the quantum photocell $S(t) = \text{tr} \{ \rho \log \rho \}$ can be written as the sum of the entropies of the donor and acceptor, $S_D(t) = -\text{tr}_D \{ \rho_D \log \rho_D \}$ and $S_A(t) = -\text{tr}_A \{ \rho_A \log \rho_A \}$, too. This is because there is no coherent interaction between the donor and the acceptor so the whole density operator of the system has a structure $\rho = \rho_D \oplus \rho_A$.

We calculate the current through the external load as

$$I = e\Gamma \cdot \rho_{22}, \quad (32)$$

and the voltage across the external load

$$eV = E_2 - E_3 + k_B T \log \left(\frac{\rho_{22}}{\rho_{33}} \right). \quad (33)$$

The latter comes from the relation $\rho_{22}/\rho_{33} = \exp(-(E_2 - E_3 - eV)/k_B T)$. The electric power delivered to the external load by the photocell is written as $P_{\text{out}} = I(t) \cdot V(t)$ which depends on the external conductance Γ . Now that




Figure 7. (a) The diagonal matrix elements of the density operator of the photocell and the pulse profiles are plotted as a function of time. (b) The changes in the energy of the donor and acceptor, $(dE/dt)_D = E'_D$ and $(dE/dt)_A = E'_A$, the heat currents of the donor and acceptor, J_D and J_A , are plotted as a function of time. (c) The entropy of the quantum photocell, $S(t)$, the entropy of the donor, $S_D(t)$, and the entropy of the acceptor, $S_A(t)$, are calculated as a function of time. The power $P_D(t)$ delivered to the donor by the photon pulses, the current $I(t)$, the voltage $V(t)$, and the efficiency η are plotted as a function of time. Parameters: $\langle n \rangle = 10$, $\gamma = 10^{-3}\omega_0$, $\gamma = 10^{-2}\omega_0$, $\Gamma = 0.1\omega_0$, $\Omega = \omega_0/4\pi$, and $\hbar\omega_0 = E_1 - E_0 = 1.8 \text{ eV}$.

we have the power delivered by the photon pulse, Eq. (30) and the electric power output $P_{\text{out}}(t)$, we can define the power efficiency of the quantum photocell as

$$\eta_p = \frac{P_{\text{out}}(t)}{P_D(t)}. \quad (34)$$

We solve numerically the LGKS Eq. (26) using the Runge-Kutta method for different sequences of photon pulses to obtain the quantum thermodynamic quantities. Fig. 7 (a) plots the population of each level of the quantum photocell over time when the sequences of the Gaussian pulses are applied one immediately after another almost in a continuum limit. Fig. 7 (b) shows the power $P_D(t)$ delivered to the donor by the photon, the heat dissipation at the donor and the acceptor, $J_D(t)$ and $J_A(t)$, the change in energies of the donor and acceptor, $\dot{E}_D(t)$ and $\dot{E}_A(t)$. Fig. 7 (c) show the total entropy $S(t)$ of the system and the entropies of the donor and the acceptor, $S_D(t)$ and $S_A(t)$. Our numerical calculation confirms that the total entropy is the sum of those of




Figure 8. From discrete mode to the continuous mode operation by changing the interval of the pulses. Parameters: $\langle n \rangle = 1$, $\gamma = 10^{-3}\omega_0$, $\gamma = 10^{-2}\omega_0$ and $\Omega = \omega_0/4\pi$.

the donor and the acceptor, $S(t) = S_D(t) + S_A(t)$, as explained before.

Fig. 7 (d) depicts the the current $I(t)$, the voltage $V(t)$, the electric power output $P_{\text{out}}(t)$, the power delivered by the photon $P_D(t)$, and the power efficiency η . From the figures, we see that these quantities initially show an oscillatory behavior then become saturated in the long time limit. In particular, the asymptotic value of power efficiency is as high as $\eta_p \sim 0.36$, which can also be interpreted as the work efficiency, i.e. work output divided by energy input, when η_p is constant.

There are different types of quantum heat engines like continuous engine, two-stroke engine and four-stroke engine. Many other studies considered the quantum photocell as the continuous heat engine where the donor is in thermal contact with the hot reservoir and the acceptor is in the cold bath. Fig. 7 demonstrates the photocell as a continuous heat engine, which is not in contact with a hot bath, but is supplied input energy by photon pulses. In our case, we have a flexibility of engineering the input photon pulses as desired. In Fig. ??, we further compare the power efficiency between two cases. Fig. ?? (a) is the case where the photon pulses are applied at a finite time interval (discrete mode operation). On the other hand, Fig. ?? (b) is the case where the photon pulses are

applied almost continuously. Both have the same energy parameter $\langle n \rangle = 1$ of incoming Gaussian pulses. In the discrete mode, we see an oscillatory behavior of power efficiency between 0.2 and 0.6. In the continuum mode, the power efficiency does not oscillate but asymptotically approaches the value $\eta_p \sim 0.36$.

Let us compare the two cases in Fig. 7 and Fig. ?? (b) both dealing with the continuum limit with different energies $\langle n \rangle = 10$ and $\langle n \rangle = 1$, respectively. It is interesting to find that the efficiency turns out to be the same $\eta_p \sim 0.36$ regardless of the pulse energy of our consideration. Of course, the transient behaviors are different as the high-energy case shows an oscillatory behavior while the low-energy case does not. Aside from details, we see that our model of heat engine offers a possibility to make an efficient quantum engine with a proper design.

V. SUMMARY

In this paper, we studied quantum thermodynamics of two open quantum systems, the two-level system and the quantum photovoltaic model, driven by the Gaussian photon pulses. By solving the master equation with the time-dependent Hamiltonian of the Gaussian photon pulses, we calculated quantum thermodynamic quantities. For the two-level system in the cold bath, we examined the first law of quantum thermodynamics, which relates the energy change of the system, the heat current, and the power. We also illustrated the second law of thermodynamics by confirming that the entropy production is positive.

More importantly, we investigated the quantum photovoltaic cell in the cold bath driven by the sequence of the Gaussian photon pulses. The power efficiency of the quantum photocell was considered as the ratio of the output power delivered to the external load by the photocell to the input power delivered by the photon pulses. We showed that the quantum photocell as a heat engine can operate both in the discrete stroke mode and in the continuous stroke mode by changing the sequence of the photon pulses.

Our model of quantum heat engine based on a driven quantum system in contact with a single bath seems worthwhile to further investigate. In our work we showed that an efficiency as high as $\eta_p = 0.36$ can be achieved, which should be further explored in a broad range of system parameters. There are some meaningful directions to consider. One is to study how the dark state or the quantum coherence can further enhance the performance of the photocell. We also note that recently Chan *et al.* [30] studied the quantum dynamics of excitons by absorption of single photons in photosynthetic light-harvesting complexes. It would be interesting how the photosynthetic light-harvesting complexes behave when the photon pulses are applied. Moreover, while we considered the Gaussian photon pulses in the current work, other

photon pulses, for example, hyperbolic secant, rectangular, or symmetric exponential pulses may be tested to come up with an optimal design [29]. An open problem is how to mimic the thermal photon from the hot ther-

mal bath and to incorporate the thermal photons into the simulation. The quantum photocells considered here can be simulated on quantum computers [33, 34].

-
- [1] William Shockley and Hans J. Queisser, “Detailed balance limit of efficiency of p-n junction solar cells,” *Journal of Applied Physics* **32**, 510–519 (1961).
- [2] Marlan O. Scully, “Quantum photocell: Using quantum coherence to reduce radiative recombination and increase efficiency,” *Phys. Rev. Lett.* **104**, 207701 (2010).
- [3] Marlan O. Scully, Kimberly R. Chapin, Konstantin E. Dorfman, Moochan Barnabas Kim, and Anatoly Svidzinsky, “Quantum heat engine power can be increased by noise-induced coherence,” *Proceedings of the National Academy of Sciences* **108**, 15097–15100 (2011).
- [4] Anatoly A. Svidzinsky, Konstantin E. Dorfman, and Marlan O. Scully, “Enhancing photovoltaic power by fano-induced coherence,” *Phys. Rev. A* **84**, 053818 (2011).
- [5] C. Creatore, M. A. Parker, S. Emmott, and A. W. Chin, “Efficient biologically inspired photocell enhanced by delocalized quantum states,” *Phys. Rev. Lett.* **111**, 253601 (2013).
- [6] Yiteng Zhang, Sangchul Oh, Fahhad H. Alharbi, Gregory S. Engel, and Sabre Kais, “Delocalized quantum states enhance photocell efficiency,” *Phys. Chem. Chem. Phys.* **17**, 5743–5750 (2015).
- [7] Amir Fruchtmann, Rafael Gómez-Bombarelli, Brendon W. Lovett, and Erik M. Gauger, “Photocell optimization using dark state protection,” *Phys. Rev. Lett.* **117**, 203603 (2016).
- [8] K. D. B. Higgins, B. W. Lovett, and E. M. Gauger, “Quantum-enhanced capture of photons using optical ratchet states,” *The Journal of Physical Chemistry C* **121**, 20714–20719 (2017).
- [9] Konstantin E. Dorfman, Dmitri V. Voronine, Shaul Mukamel, and Marlan O. Scully, “Photosynthetic reaction center as a quantum heat engine,” *Proceedings of the National Academy of Sciences* **110**, 2746–2751 (2013).
- [10] N. Killoran, S. F. Huelga, and M. B. Plenio, “Enhancing light-harvesting power with coherent vibrational interactions: A quantum heat engine picture,” *The Journal of Chemical Physics* **143**, 155102 (2015).
- [11] Yasuhiro Yamada, Youhei Yamaji, and Masatoshi Imada, “Exciton lifetime paradoxically enhanced by dissipation and decoherence: Toward efficient energy conversion of a solar cell,” *Phys. Rev. Lett.* **115**, 197701 (2015).
- [12] Richard Stones, Hoda Hossein-Nejad, Rienk van Grondelle, and Alexandra Olaya-Castro, “On the performance of a photosystem ii reaction centre-based photocell,” *Chem. Sci.* **8**, 6871–6880 (2017).
- [13] Ronnie Kosloff and Amikam Levy, “Quantum heat engines and refrigerators: Continuous devices,” *Annual Review of Physical Chemistry* **65**, 365–393 (2014), pMID: 24689798.
- [14] Raam Uzdin, Amikam Levy, and Ronnie Kosloff, “Equivalence of quantum heat machines, and quantum-thermodynamic signatures,” *Phys. Rev. X* **5**, 031044 (2015).
- [15] H. E. D. Scovil and E. O. Schulz-DuBois, “Three-level masers as heat engines,” *Phys. Rev. Lett.* **2**, 262–263 (1959).
- [16] Sangchul Oh, “Efficiency and power enhancement of solar cells by dark states,” *Physics Letters A* **383**, 125857 (2019).
- [17] Stefano Tomasi and Ivan Kassal, “Classification of coherent enhancements of light-harvesting processes,” *The Journal of Physical Chemistry Letters* **11**, 2348–2355 (2020), pMID: 32119554.
- [18] G. Lindblad, “On the generators of quantum dynamical semigroups,” *Communications in Mathematical Physics* **48**, 119–130 (1976).
- [19] Vittorio Gorini, Andrzej Kossakowski, and E. C. G. Sudarshan, “Completely positive dynamical semigroups of n-level systems,” *Journal of Mathematical Physics* **17**, 821–825 (1976).
- [20] R. Alicki and K. Lendi, *Quantum Dynamical Semigroup and Applications* (Springer-Verlag, Berlin, 1984).
- [21] Heinz-Peter Breuer and Francesco Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, New York, 2002).
- [22] Ángel Rivas and Susana F. Huelga, *Open Quantum Systems*, 1st ed. (Springer-Verlag Berlin Heidelberg, 2012).
- [23] Herbert Spohn, “Entropy production for quantum dynamical semigroups,” *Journal of Mathematical Physics* **19**, 1227–1230 (1978).
- [24] Herbert Spohn and Joel L. Lebowitz, “Irreversible thermodynamics for quantum systems weakly coupled to thermal reservoirs,” in *Advances in Chemical Physics*, Vol. XXXVIII, edited by Stuart A. Rice (John Wiley & Sons, Ltd, 1978) pp. 109–142.
- [25] R. Alicki, “The quantum open system as a model of the heat engine,” *Journal of Physics A: Mathematical and General* **12**, L103–L107 (1979).
- [26] Siddhartha Das, Sumeet Khatri, George Siopsis, and Mark M. Wilde, “Fundamental limits on quantum dynamics based on entropy change,” *Journal of Mathematical Physics* **59**, 012205 (2018).
- [27] E. T. Jaynes and F. W. Cummings, “Comparison of quantum and semiclassical radiation theories with application to the beam maser,” *Proceedings of the IEEE* **51**, 89–109 (1963).
- [28] Rodney Loudon, *The Quantum Theory of Light* (Clarendon Press, Oxford, 1973).
- [29] Yimin Wang, Jiří Minář, Lana Sheridan, and Valerio Scarani, “Efficient excitation of a two-level atom by a single photon in a propagating mode,” *Phys. Rev. A* **83**, 063842 (2011).
- [30] Herman C H Chan, Omar E Gamel, Graham R Fleming, and K Birgitta Whaley, “Single-photon absorption by single photosynthetic light-harvesting complexes,” *Journal of Physics B: Atomic, Molecular and Optical Physics* **51**, 054002 (2018).

- [31] Howard Carmichael, *An Open Systems Approach to Quantum Optics* (Springer-Verlag, Berlin, 1993).
- [32] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics* (Cambridge University Press, New York, 1995).
- [33] Anton Potočnik, Arno Bargerbos, Florian A. Y. N. Schröder, Saeed A. Khan, Michele C. Collodo, Simone Gasparinetti, Yves Salathé, Celestino Creatore, Christopher Eichler, Hakan E. Türeci, Alex W. Chin, and Andreas Wallraff, “Studying light-harvesting models with superconducting circuits,” *Nature Communications* **9**, 904 (2018).
- [34] Gleb Maslennikov, Shiqian Ding, Roland Hablützel, Jaren Gan, Alexandre Roulet, Stefan Nimmrichter, Jibo Dai, Valerio Scarani, and Dzmitry Matsukevich, “Quantum absorption refrigerator with trapped ions,” *Nature Communications* **10**, 202 (2019).