

The Navier-Stokes equation with time quasi-periodic external force: existence and stability of quasi-periodic solutions

Riccardo Montalto

Abstract. We prove the existence of small amplitude, time-quasi-periodic solutions (invariant tori) for the incompressible Navier-Stokes equation on the d -dimensional torus \mathbb{T}^d , with a small, quasi-periodic in time external force. We also show that they are orbitally and asymptotically stable in H^s (for s large enough). More precisely, for any initial datum which is close to the invariant torus, there exists a unique global in time solution which stays close to the invariant torus for all times. Moreover, the solution converges asymptotically to the invariant torus for $t \rightarrow +\infty$, with an exponential rate of convergence $O(e^{-\alpha t})$ for any arbitrary $\alpha \in (0, 1)$.

Keywords: Fluid dynamics, Navier-Stokes equation, quasi-periodic solutions, asymptotic and orbital stability.

MSC 2010: 37K55, 35Q30, 76D05.

Contents

1	Introduction	2
2	Functional spaces	7
2.1	Leray projector and some elementary properties of the Navier-Stokes equation	9
3	Construction of quasi-periodic solutions	10
3.1	Proof of Theorem 1.1	13
4	Orbital and asymptotic stability	14
4.1	Dispersive estimates for the heat propagator	15
4.2	Proof of Proposition 4.1	19
4.3	Proof of Theorem 1.2	21
A	appendix	21

1 Introduction

We consider the Navier-Stokes equation for an incompressible fluid on the d -dimensional torus \mathbb{T}^d , $d \geq 2$, $\mathbb{T} := \mathbb{R}/2\pi\mathbb{Z}$,

$$\begin{cases} \partial_t u - \Delta u + u \cdot \nabla u + \nabla p = \varepsilon f(\omega t, x) \\ \operatorname{div} u = 0 \end{cases} \quad (1.1)$$

where $\varepsilon \in (0, 1)$ is a small parameter, the frequency $\omega = (\omega_1, \dots, \omega_\nu) \in \mathbb{R}^\nu$ is a ν -dimensional vector and $f : \mathbb{T}^\nu \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ is a smooth quasi-periodic external force. The unknowns of the problem are the velocity field $u = (u_1, \dots, u_d) : \mathbb{R} \times \mathbb{T}^d \rightarrow \mathbb{R}^d$, and the pressure $p : \mathbb{R} \times \mathbb{T}^d \rightarrow \mathbb{R}$. For convenience, we set the viscosity parameter in front of the laplacian equal to one. We assume that f has zero space-time average, namely

$$\int_{\mathbb{T}^\nu \times \mathbb{T}^d} f(\varphi, x) d\varphi dx = 0. \quad (1.2)$$

The purpose of the present paper is to show the existence and the stability of smooth quasi-periodic solutions of the equation (1.1). More precisely we show that if f is a sufficiently regular vector field satisfying (1.2), for ε sufficiently small and for $\omega \in \mathbb{R}^\nu$ diophantine, i.e.

$$\begin{aligned} |\omega \cdot \ell| &\geq \frac{\gamma}{|\ell|^\nu}, \quad \forall \ell \in \mathbb{Z}^\nu \setminus \{0\}, \\ \text{for some } \gamma &\in (0, 1), \end{aligned} \quad (1.3)$$

¹ then the equation (1.1) admits smooth quasi-periodic solutions (which are referred to also as invariant tori) $u_\omega(t, x) = U(\omega t, x)$, $p_\omega(t, x) = P(\omega t, x)$, $U : \mathbb{T}^\nu \times \mathbb{T}^d \rightarrow \mathbb{R}^d$, $P : \mathbb{T}^\nu \times \mathbb{T}^d \rightarrow \mathbb{R}$ of size $O(\varepsilon)$, oscillating with the same frequency $\omega \in \mathbb{R}^\nu$ of the forcing term. If the forcing term has zero-average in x , i.e.

$$\int_{\mathbb{T}^d} f(\varphi, x) dx = 0, \quad \forall \varphi \in \mathbb{T}^\nu \quad (1.4)$$

then the result holds for any frequency vector $\omega \in \mathbb{R}^\nu$, without requiring any non-resonance condition. Furthermore, we show also the orbital and the asymptotic stability of these quasi-periodic solutions in high Sobolev norms. More precisely, for any sufficiently regular initial datum which is δ -close to the invariant torus (w.r. to the H^s topology), the corresponding solution of (1.1) is global in time and it satisfies the following properties.

- **Orbital stability.** For all times $t \geq 0$, the distance in H^s between the solution and the invariant torus is of order $O(\delta)$.
- **Asymptotic stability.** The solution converges asymptotically to the invariant torus in high Sobolev norm $\|\cdot\|_{H_x^s}$ as $t \rightarrow +\infty$, with a rate of convergence which is exponential, i.e. $O(e^{-\alpha t})$, for any arbitrary $\alpha \in (0, 1)$.

¹It is well known that a.e. frequency in \mathbb{R}^ν (w.r. to the Lebesgue measure) is diophantine.

In order to state precisely our main results, we introduce some notations. For any vector $a = (a_1, \dots, a_p) \in \mathbb{R}^p$, we denote by $|a|$ its Euclidean norm, namely $|a| := \sqrt{a_1^2 + \dots + a_p^2}$. Let $d, n \in \mathbb{N}$ and a function $u \in L^2(\mathbb{T}^d, \mathbb{R}^n)$. Then $u(x)$ can be expanded in Fourier series

$$u(x) = \sum_{\xi \in \mathbb{Z}^d} \widehat{u}(\xi) e^{ix \cdot \xi}$$

where its Fourier coefficients $\widehat{u}(\xi)$ are defined by

$$\widehat{u}(\xi) := \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} u(x) e^{-ix \cdot \xi} dx, \quad \forall \xi \in \mathbb{Z}^d.$$

For any $s \geq 0$, we denote by $H^s(\mathbb{T}^d, \mathbb{R}^n)$ the standard Sobolev space of functions $u : \mathbb{T}^d \rightarrow \mathbb{R}^n$ equipped by the norm

$$\|u\|_{H_x^s} := \left(\sum_{\xi \in \mathbb{Z}^d} \langle \xi \rangle^{2s} |\widehat{u}(\xi)|^2 \right)^{\frac{1}{2}}, \quad \langle \xi \rangle := \max\{1, |\xi|\}. \quad (1.5)$$

We also define the Sobolev space of functions with zero average

$$H_0^s(\mathbb{T}^d, \mathbb{R}^n) := \left\{ u \in H^s(\mathbb{T}^d, \mathbb{R}^n) : \int_{\mathbb{T}^d} u(x) dx = 0 \right\}. \quad (1.6)$$

Moreover, given a Banach space $(X, \|\cdot\|_X)$ and an interval $\mathcal{I} \subseteq \mathbb{R}$, we denote by $\mathcal{C}_b^0(\mathcal{I}, X)$ the space of bounded, continuous functions $u : \mathcal{I} \rightarrow X$, equipped with the sup-norm

$$\|u\|_{\mathcal{C}_t^0 X} := \sup_{t \in \mathcal{I}} \|u(t)\|_X.$$

For any integer $k \geq 1$, $\mathcal{C}_b^k(\mathcal{I}, X)$ is the space of k -times differentiable functions $u : \mathcal{I} \rightarrow X$ with continuous and bounded derivatives equipped with the norm

$$\|u\|_{\mathcal{C}_t^k X} := \max_{n \leq k} \|\partial_t^n u\|_{\mathcal{C}_t^0 X}.$$

In a similar way we define the spaces $\mathcal{C}^0(\mathbb{T}^\nu, X)$, $\mathcal{C}^k(\mathbb{T}^\nu, X)$, $k \geq 1$ and the corresponding norms $\|\cdot\|_{\mathcal{C}_\varphi^0 X}$, $\|\cdot\|_{\mathcal{C}_\varphi^k X}$ (where \mathbb{T}^ν is the ν -dimensional torus). We also denote by $\mathcal{C}^N(\mathbb{T}^\nu \times \mathbb{T}^d, \mathbb{R}^d)$ the space of N -times continuously differentiable functions $\mathbb{T}^\nu \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ equipped with the standard \mathcal{C}^N norm $\|\cdot\|_{\mathcal{C}^N}$.

Notation. Throughout the whole paper, the notation $A \lesssim B$ means that there exists a constant C which can depend on the number of frequencies ν , the dimension of the torus d , the constant γ appearing in the diophantine condition (1.3) and on the \mathcal{C}^N norm of the forcing term $\|f\|_{\mathcal{C}^N}$. Given n positive real numbers $s_1, \dots, s_n > 0$, we write $A \lesssim_{s_1, \dots, s_n} B$ if there exists a constant $C = C(s_1, \dots, s_n) > 0$ (eventually depending also on $d, \nu, \gamma, \|f\|_{\mathcal{C}^N}$) such that $A \leq CB$.

We are now ready to state the main results of our paper.

Theorem 1.1 (Existence of quasi-periodic solutions). *Let $s > d/2 + 1$, $N > \frac{3\nu}{2} + s + 2$, $\omega \in \mathbb{R}^\nu$ diophantine (see (1.3)) and assume that the forcing term f is in $\mathcal{C}^N(\mathbb{T}^\nu \times \mathbb{T}^d, \mathbb{R}^d)$ and it satisfies (1.2). Then there exists $\varepsilon_0 = \varepsilon_0(f, s, d, \nu) \in (0, 1)$ small enough and a constant $C = C(f, s, d, \nu) > 0$ large enough such that for any $\varepsilon \in (0, \varepsilon_0)$ there exist $U \in \mathcal{C}^1(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^d))$, $P \in \mathcal{C}^0(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}))$ satisfying*

$$\int_{\mathbb{T}^\nu \times \mathbb{T}^d} U(\varphi, x) d\varphi dx = 0, \quad \int_{\mathbb{T}^d} P(\varphi, x) dx = 0, \quad \forall \varphi \in \mathbb{T}^\nu$$

such that $(u_\omega(t, x), p_\omega(t, x)) := (U(\omega t, x), P(\omega t, x))$ solves the Navier-Stokes equation (1.1) and

$$\|U\|_{\mathcal{C}_\varphi^1 H_x^s}, \|P\|_{\mathcal{C}_\varphi^0 H_x^s} \leq C\varepsilon.$$

If the forcing term f has zero space average, i.e. it satisfies (1.4), then the same statement holds for any frequency vector $\omega \in \mathbb{R}^\nu$ and $U(\varphi, x)$ satisfies

$$\int_{\mathbb{T}^d} U(\varphi, x) dx = 0, \quad \forall \varphi \in \mathbb{T}^\nu.$$

Theorem 1.2 (Stability). *Let $\alpha \in (0, 1)$, $s > d/2 + 1$, $N > \frac{3\nu}{2} + s + 2$, u_ω, p_ω be given in Theorem 1.1. Then there exists $\delta = \delta(f, s, \alpha, d, \nu) \in (0, 1)$ small enough and a constant $C = C(f, s, \alpha, d, \nu) > 0$ large enough such that for $\varepsilon \leq \delta$ and for any initial datum $u_0 \in H^s(\mathbb{T}^d, \mathbb{R}^d)$ satisfying*

$$\|u_0 - u_\omega(0, \cdot)\|_{H_x^s} \leq \delta, \quad \int_{\mathbb{T}^d} (u_0(x) - u_\omega(0, x)) dx = 0$$

there exists a unique global classical solution (u, p) of the Navier-Stokes equation (1.1) with initial datum $u(0, x) = u_0(x)$ which satisfies

$$\begin{aligned} u &\in \mathcal{C}_b^0([0, +\infty), H^s(\mathbb{T}^d, \mathbb{R}^d)) \cap \mathcal{C}_b^1([0, +\infty), H^{s-2}(\mathbb{T}^d, \mathbb{R}^d)), \quad p \in \mathcal{C}_b^0([0, +\infty), H_0^s(\mathbb{T}^d, \mathbb{R})), \\ \int_{\mathbb{T}^d} (u(t, x) - u_\omega(t, x)) dx &= 0, \quad \forall t \geq 0, \\ \|u(t, \cdot) - u_\omega(t, \cdot)\|_{H_x^s}, \|\partial_t u(t, \cdot) - \partial_t u_\omega(t, \cdot)\|_{H_x^{s-2}}, \|p(t, \cdot) - p_\omega(t, \cdot)\|_{H_x^s} &\leq C\delta e^{-\alpha t} \end{aligned}$$

for any $t \geq 0$.

The investigation of the Navier-Stokes equation with time periodic external force dates back to Serrin [40], Yudovich [41], Lions [30], Prodi [36] and Prouse [37]. In these papers the authors proved the existence of weak periodic solutions on bounded domains, oscillating with the same frequency of the external force. The existence of weak quasi-periodic solutions in dimension two has been proved by Prouse [38]. More recently these results have been extended to unbounded domains by Maremonti [27], Maremonti-Padula [28], Salvi [39] and then by Galdi [19], [20], Galdi-Silvestre [21], Galdi-Kyed [22] and Kyed [32]. We point out that in some of the aforementioned results, no smallness assumptions on the forcing term are needed and therefore, the periodic solutions obtained are not small in size, see for

instance [40], [41], [36], [37], [38], [28], [39]. The asymptotic stability of periodic solutions (also referred to as *attainability property*) has been also investigated in [27], [28], but it is only proved w.r. to the L^2 -norm and the rate of convergence provided is $O(t^{-\eta})$ for some constant $\eta > 0$. More recently Galdi and Hishida [23] proved the asymptotic stability for the Navier-Stokes equation with a traslation velocity term, by using the Lorentz spaces and they provided a rate of convergence which is essentially $O(t^{-\frac{1}{2}+\varepsilon})$. In the present paper we consider the Navier-Stokes equation on the d -dimensional torus with a small, quasi-periodic in time external force. We show the existence of smooth quasi-periodic solutions (which are also referred to as invariant tori) of small amplitude and we prove their orbital and asymptotic stability in H^s for s large enough (at least larger than $d/2 + 1$). Furthermore the rate of convergence to the invariant torus, in H^s , for $t \rightarrow +\infty$ is of order $O(e^{-\alpha t})$ for any arbitrary $\alpha \in (0, 1)$. To the best of our knowledge, this is the first result of this kind.

It is also worth to mention that the existence of quasi-periodic solutions, that is also referred to as KAM (Kolmogorov-Arnold-Moser) theory, for dispersive and hyperbolic-type PDEs is a more difficult matter, due to the presence of the so-called *small divisors problem*. The existence of time-periodic and quasi-periodic solutions of PDEs started in the late 1980s with the pioneering papers of Kuksin [33], Wayne [42] and Craig-Wayne [13], see also [34], [31] for generalizations to PDEs with unbounded nonlinearities. We refer to the recent review [7] for a complete list of references.

Many PDEs arising from fluid dynamics like the water waves equations or the Euler equation are fully nonlinear or quasi-linear equations (the nonlinear part contains as many derivatives as the linear part). The breakthrough idea, based on pseudo-differential calculus and micro-local analysis, in order to deal with these kind of PDEs has been introduced by Iooss, Plotnikov and Toland [25] in the problem of finding periodic solutions for the water waves equation. The methods developed in [25], combined with a *KAM-normal form* procedure have been used to develop a general method for PDEs in one dimension, which allows to construct *quasi-periodic* solutions of quasilinear and fully nonlinear PDEs, see [2], [17], [11], [1] and references therein. The extension of KAM theory to higher space dimension $d > 1$ is a difficult matter due to the presence of very strong resonance-phenomena, often related to high multiplicity of eigenvalues. The fist breakthrough results in this directions (for equations with perturbations which do not contain derivatives) have been obtained by Eliasson and Kuksin [16] and by Bourgain [12] (see also Berti-Bolle [8], [9], Geng-Xu-You [24], Procesi-Procesi [35], Berti-Corsi-Procesi [10].)

Extending KAM theory to PDEs with unbounded perturbations in higher space dimension is one of the main open problems in the field. Up to now, this has been achieved only in few examples, see [29], [15], [18], [6], [4], [5] and recently on the 3D Euler equation [3] which is the most meaningful physical example.

For the Navier-Stokes equation, unlike in the aforementioned papers on KAM for PDEs, the existence of quasi-periodic solutions is not a small divisors problem and it can be done by using a classical fixed point argument. This is due to the fact that the Navier-Stokes equation is a parabolic PDE and the presence of dissipation avoids the small divisors. In the same spirit, it is also worth to mention [14], in which the authors investigate quasi-periodic solutions of nonlinear wave equations with dissipation. We also point out that the present paper is the first example in which the stability of invariant tori, in high Sobolev

norms, is proved for all times (and it is even an asymptotic stability). This is possible since the presence of the dissipation allows to prove strong time-decay estimates from which one deduces orbital and asymptotic stability. In the framework of dispersive and hyperbolic PDEs, the orbital stability of invariant tori is usually proved only for *large times* by using normal form techniques. The first result in this direction has been proved in [26]. In the remaining part of the introduction, we sketch the main points of our proof.

As we already explained above, the absence of small divisors is due to the fact that the Navier-Stokes equation is a parabolic PDE. More precisely, this fact is related to invertibility properties of the linear operator $L_\omega := \omega \cdot \partial_\varphi - \Delta$ (where $\omega \cdot \partial_\varphi := \sum_{i=0}^\nu \omega_i \partial_{\varphi_i}$) acting on Sobolev spaces of functions $u(\varphi, x)$, $(\varphi, x) \in \mathbb{T}^\nu \times \mathbb{T}^d$ with zero average w.r. to x . Since the eigenvalues of L_ω are $i\omega \cdot \ell + |j|^2$, $\ell \in \mathbb{Z}^\nu$, $j \in \mathbb{Z}^d \setminus \{0\}$, the inverse of L_ω *gains two space derivatives*, see Lemma 3.2. This is sufficient to perform a fixed point argument on the map Φ defined in (3.13) from which one deduces the existence of smooth quasi-periodic solutions of small amplitude. The asymptotic and orbital stability of quasi-periodic solutions (which are constructed in Section 3) are proved in Section 4. More precisely we show that for any initial datum u_0 which is δ -close to the quasi-periodic solution $u_\omega(0, x)$ in H^s norm (and such that $u_0 - u_\omega(0, \cdot)$ has zero average), there exists a unique classical solution (u, p) such that

$$\|u(t, \cdot) - u_\omega(t, \cdot)\|_{H_x^s} = O(\delta e^{-\alpha t}), \quad \|p(t, \cdot) - p_\omega(t, \cdot)\|_{H_x^s} = O(\delta e^{-\alpha t}), \quad \alpha \in (0, 1)$$

for any $t \geq 0$. This is exactly the content of Theorem 1.2, which easily follows from Proposition 4.1. This Proposition is proved also by a fixed point argument on the nonlinear map Φ defined in (4.31) in weighted Sobolev spaces \mathcal{E}_s (see (4.14)), defined by the norm

$$\|u\|_{\mathcal{E}_s} := \sup_{t \geq 0} e^{\alpha t} \|u(t, \cdot)\|_{H_x^s}$$

where $\alpha \in (0, 1)$ is a fixed constant. The fixed point argument relies on some *dispersive-type estimates* for the heat propagator $e^{t\Delta}$, which are proved in Section 4.1. The key estimates are the following.

1. For any $u_0 \in H^{s-1}(\mathbb{T}^d, \mathbb{R}^d)$ with zero average and for any $n \in \mathbb{N}$, $\alpha \in (0, 1)$, $t > 0$, one has

$$\|e^{t\Delta} u_0\|_{H_x^s} \leq C(n, \alpha) t^{-\frac{n}{2}} e^{-\alpha t} \|u_0\|_{H_x^{s-1}} \quad (1.7)$$

for some constant $C(n, \alpha) > 0$ (see Lemma 4.2). This estimate states that the heat propagator *gains one-space derivative* and exponential decay in time $e^{-\alpha t} t^{-\frac{n}{2}}$. Note that, without gain of derivatives on u_0 , the exponential decay is stronger, namely e^{-t} , see Lemma 4.2-(i).

2. For any $f \in \mathcal{E}_{s-1}$

$$\left\| \int_0^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} \leq C(\alpha) e^{-\alpha t} \|f\|_{\mathcal{E}_{s-1}} \quad (1.8)$$

for some constant $C(\alpha) > 0$ (see Proposition 4.5). This estimate states that the integral term which usually appears in the Duhamel formula (see (4.31)) gains one space derivative w.r. to $f(t, x)$ and keeps the same exponential decay in time as $f(t, x)$.

We also remark that the constants $C(n, \alpha)$, $C(\alpha)$ appearing in the estimates (1.7), (1.8) tend to ∞ when $\alpha \rightarrow 1$. This is the reason why it is not possible to get a decay $O(e^{-t})$ in the asymptotic stability estimate provided in Theorem 1.2.

The latter two estimates allow to show in Proposition 4.9 that the map Φ defined in (4.31) is a contraction. The proof of Theorem 1.2 is then easily concluded in Section 4.3.

As a concluding remark, it is also worth to mention that our methods does not cover the zero viscosity limit $\mu \rightarrow 0$, where μ is the usual viscosity parameter in front of the laplacian (that we set for convenience equal to one). Indeed some constants in our estimates become infinity when $\mu \rightarrow 0$. Actually, it would be very interesting to study the *singular perturbation problem* for $\mu \rightarrow 0$ and to see if one is able to recover the quasi-periodic solutions of the Euler equation constructed in [3].

ACKNOWLEDGEMENTS. The author is supported by INDAM-GNFM. The author warmly thanks Dario Bambusi, Luca Franzoi, Thomas Kappeler and Alberto Maspero for their feedbacks.

2 Functional spaces

In this section we collect some standard technical tools which will be used in the proof of our results.

For $u = (u_1, \dots, u_n) \in H^s(\mathbb{T}^d, \mathbb{R}^n)$,

$$\|u\|_{H_x^s} \simeq \max_{i=1, \dots, n} \|u_i\|_{H_x^s}. \quad (2.1)$$

The following standard algebra lemma holds.

Lemma 2.1. *Let $s > d/2$ and $u, v \in H^s(\mathbb{T}^d, \mathbb{R}^n)$. Then $u \cdot v \in H^s(\mathbb{T}^d, \mathbb{R})$ (where \cdot denotes the standard scalar product on \mathbb{R}^n) and $\|u \cdot v\|_{H_x^s} \lesssim_s \|u\|_{H_x^s} \|v\|_{H_x^s}$.*

We also consider functions

$$\mathbb{T}^\nu \rightarrow L^2(\mathbb{T}^d, \mathbb{R}^n), \quad \varphi \mapsto u(\varphi, \cdot)$$

which are in $L^2(\mathbb{T}^\nu, L^2(\mathbb{T}^d, \mathbb{R}^n))$. We can write the Fourier series of a function

$$u \in L^2(\mathbb{T}^\nu, L^2(\mathbb{T}^d, \mathbb{R}^n)) \text{ as}$$

$$u(\varphi, \cdot) = \sum_{\ell \in \mathbb{Z}^\nu} \hat{u}(\ell, \cdot) e^{i\ell \cdot \varphi} \quad (2.2)$$

where

$$\hat{u}(\ell, \cdot) := \frac{1}{(2\pi)^\nu} \int_{\mathbb{T}^\nu} u(\varphi, \cdot) e^{-i\ell \cdot \varphi} d\varphi \in L^2(\mathbb{T}^d, \mathbb{R}^n), \quad \ell \in \mathbb{Z}^\nu. \quad (2.3)$$

By expanding also the function $\hat{u}(\ell, \cdot)$ in Fourier series, we get

$$\begin{aligned} \hat{u}(\ell, x) &= \sum_{j \in \mathbb{Z}^d} \hat{u}(\ell, j) e^{ij \cdot x}, \\ \hat{u}(\ell, j) &:= \frac{1}{(2\pi)^{\nu+d}} \int_{\mathbb{T}^{\nu+d}} u(\varphi, x) e^{-i\ell \cdot \varphi} e^{-ij \cdot x} d\varphi dx, \quad (\ell, j) \in \mathbb{Z}^\nu \times \mathbb{Z}^d \end{aligned} \quad (2.4)$$

and hence we can write

$$u(\varphi, x) = \sum_{\ell \in \mathbb{Z}^\nu} \sum_{j \in \mathbb{Z}^d} \widehat{u}(\ell, j) e^{i\ell \cdot \varphi} e^{ij \cdot x}. \quad (2.5)$$

For any $\sigma, s \geq 0$, we define the Sobolev space $H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^n))$ as the space of functions $u \in L^2(\mathbb{T}^\nu, L^2(\mathbb{T}^d, \mathbb{R}^n))$ equipped by the norm

$$\|u\|_{\sigma, s} \equiv \|u\|_{H_\varphi^\sigma H_x^s} := \left(\sum_{\ell \in \mathbb{Z}^\nu} \langle \ell \rangle^{2\sigma} \|\widehat{u}(\ell)\|_{H_x^s}^2 \right)^{\frac{1}{2}} = \left(\sum_{\ell \in \mathbb{Z}^\nu} \sum_{j \in \mathbb{Z}^d} \langle \ell \rangle^{2\sigma} \langle j \rangle^{2s} |\widehat{u}(\ell, j)|^2 \right)^{\frac{1}{2}}. \quad (2.6)$$

Similarly to (2.1), one has that for $u = (u_1, \dots, u_n) \in H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^n))$

$$\|u\|_{\sigma, s} \simeq \max_{i=1, \dots, n} \|u_i\|_{\sigma, s}. \quad (2.7)$$

If $\sigma > \nu/2$, then

$$\begin{aligned} H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^n)) & \text{ is compactly embedded in } C^0(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^n)), \\ \text{and } \|u\|_{C_\varphi^0 H_x^s} & \lesssim_\sigma \|u\|_{H_\varphi^\sigma H_x^s}. \end{aligned} \quad (2.8)$$

Moreover, the following standard algebra property holds.

Lemma 2.2. *Let $\sigma > \frac{\nu}{2}$, $s > \frac{d}{2}$, $u, v \in H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^n))$. Then $u \cdot v \in H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^n))$ and $\|u \cdot v\|_{\sigma, s} \lesssim_{\sigma, s} \|u\|_{\sigma, s} \|v\|_{\sigma, s}$.*

For any $u \in L^2(\mathbb{T}^d, \mathbb{R}^n)$ we define the orthogonal projections π_0 and π_0^\perp as

$$\pi_0 u := \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} u(x) dx = \widehat{u}(0) \quad \text{and} \quad \pi_0^\perp u := u - \pi_0 u = \sum_{\xi \in \mathbb{Z}^d \setminus \{0\}} \widehat{u}(\xi) e^{ix \cdot \xi}. \quad (2.9)$$

According to (2.9), (2.5), every function $u \in L^2(\mathbb{T}^\nu, L^2(\mathbb{T}^d, \mathbb{R}^n))$ can be decomposed as

$$\begin{aligned} u(\varphi, x) &= u_0(\varphi) + u_\perp(\varphi, x), \\ u_0(\varphi) &:= \pi_0 u(\varphi) = \sum_{\ell \in \mathbb{Z}^\nu} \widehat{u}(\ell, 0) e^{i\ell \cdot \varphi}, \\ u_\perp(\varphi, x) &:= \pi_0^\perp u(\varphi, x) = \sum_{\ell \in \mathbb{Z}^\nu} \sum_{j \in \mathbb{Z}^d \setminus \{0\}} \widehat{u}(\ell, j) e^{i\ell \cdot \varphi} e^{ij \cdot x}. \end{aligned} \quad (2.10)$$

Clearly if $u \in H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^n))$, $\sigma, s \geq 0$, then

$$\begin{aligned} u_0 &\in H^\sigma(\mathbb{T}^\nu, \mathbb{R}^d) \quad \text{and} \quad \|u_0\|_\sigma \leq \|u\|_{\sigma, 0} \leq \|u\|_{\sigma, s}, \\ u_\perp &\in H^\sigma(\mathbb{T}^\nu, H_0^s(\mathbb{T}^d, \mathbb{R}^n)) \quad \text{and} \quad \|u_\perp\|_{\sigma, s} \leq \|u\|_{\sigma, s}, \\ \|u\|_{\sigma, s} &= \|u_0\|_\sigma + \|u_\perp\|_{\sigma, s}. \end{aligned} \quad (2.11)$$

We also prove the following lemma that we shall apply in Section 4.

Lemma 2.3. *Let $\sigma > \nu/2$, $U \in H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^n))$ and $\omega \in \mathbb{R}^\nu$. Defining $u_\omega(t, x) := U(\omega t, x)$, $(t, x) \in \mathbb{R} \times \mathbb{T}^d$, one has that $u_\omega \in \mathcal{C}_b^0(\mathbb{R}, H^s(\mathbb{T}^d, \mathbb{R}^n))$ and $\|u_\omega\|_{\mathcal{C}_t^0 H_x^s} \lesssim_\sigma \|U\|_{\sigma, s}$.*

Proof. By the Sobolev embedding (2.8), and using that the map $\mathbb{R} \rightarrow \mathbb{T}^d$, $t \mapsto \omega t$ is continuous, one has that $u_\omega \in \mathcal{C}_b^0(\mathbb{R}, H^s(\mathbb{T}^d, \mathbb{R}^n))$ and

$$\|u_\omega\|_{\mathcal{C}_t^0 H_x^s} \leq \|U\|_{\mathcal{C}_\varphi^0 H_x^s} \lesssim_\sigma \|U\|_{H_\varphi^\sigma H_x^s}.$$

□

2.1 Leray projector and some elementary properties of the Navier-Stokes equation

We introduce the space of zero-divergence vector fields

$$\mathcal{D}_0(\mathbb{T}^d) := \left\{ u \in L^2(\mathbb{T}^d, \mathbb{R}^d) : \operatorname{div}(u) = 0 \right\} \quad (2.12)$$

where clearly the divergence has to be interpreted in a distributional sense. The L^2 -orthogonal projector on this subspace of $L^2(\mathbb{T}^d, \mathbb{R}^d)$ is called the *Leray* projector and its explicit formula is given by

$$\begin{aligned} \mathfrak{L} : L^2(\mathbb{T}^d, \mathbb{R}^d) &\rightarrow \mathcal{D}_0(\mathbb{T}^d), \\ \mathfrak{L}(u) &:= u + \nabla(-\Delta)^{-1} \operatorname{div}(u) \end{aligned} \quad (2.13)$$

where the inverse of the laplacian (on the space of zero average functions) $(-\Delta)^{-1}$ is defined by

$$(-\Delta)^{-1} u(x) := \sum_{\xi \in \mathbb{Z}^d \setminus \{0\}} \frac{1}{|\xi|^2} \widehat{u}(\xi) e^{ix \cdot \xi}. \quad (2.14)$$

By expanding in Fourier series, the Leray projector \mathfrak{L} can be written as

$$\mathfrak{L}(u)(x) = u(x) + \sum_{\xi \in \mathbb{Z}^d \setminus \{0\}} \frac{\xi}{|\xi|^2} \xi \cdot \widehat{u}(\xi) e^{ix \cdot \xi}. \quad (2.15)$$

By the latter formula, one immediately deduces some elementary properties of the Leray projector \mathfrak{L} . One has

$$\int_{\mathbb{T}^d} \mathfrak{L}(u)(x) dx = \int_{\mathbb{T}^d} u(x) dx, \quad \forall u \in L^2(\mathbb{T}^d, \mathbb{R}^d) \quad (2.16)$$

and for any Fourier multiplier Λ , $\Lambda u(x) = \sum_{\xi \in \mathbb{Z}^d} \Lambda(\xi) \widehat{u}(\xi) e^{ix \cdot \xi}$, the commutator

$$[\mathfrak{L}, \Lambda] = \mathfrak{L}\Lambda - \Lambda\mathfrak{L} = 0. \quad (2.17)$$

Moreover

$$\begin{aligned} \|\mathfrak{L}(u)\|_{H_x^s} &\lesssim \|u\|_{H_x^s}, \quad \forall u \in H^s(\mathbb{T}^d, \mathbb{R}^d), \\ \|\mathfrak{L}(u)\|_{\sigma, s} &\lesssim \|u\|_{\sigma, s}, \quad \forall u \in H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^d)). \end{aligned} \quad (2.18)$$

For later purposes, we now prove the following Lemma.

Lemma 2.4. (i) Let $u, v \in H^1(\mathbb{T}^d, \mathbb{R}^d)$ and assume that $\operatorname{div}(u) = 0$, then $u \cdot \nabla v$, $\mathfrak{L}(u \cdot \nabla v)$ have zero average.

(ii) Let $\sigma > \nu/2$, $s > d/2$, $u \in H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^d))$, $v \in H^\sigma(\mathbb{T}^\nu, H^{s+1}(\mathbb{T}^d, \mathbb{R}^d))$. Then $u \cdot \nabla v \in H^\sigma(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^d))$ and $\|u \cdot \nabla v\|_{\sigma, s} \lesssim_{\sigma, s} \|u\|_{\sigma, s} \|v\|_{\sigma, s+1}$.

Proof. PROOF OF (i). By integrating by parts,

$$\int_{\mathbb{T}^d} \mathfrak{L}(u \cdot \nabla v) dx \stackrel{(2.16)}{=} \int_{\mathbb{T}^d} u \cdot \nabla v dx = - \int_{\mathbb{T}^d} \operatorname{div}(u) v dx = 0.$$

PROOF OF (ii). For $u = (u_1, \dots, u_d)$, $v = (v_1, \dots, v_d)$, the vector field $u \cdot \nabla v$ is given by

$$u \cdot \nabla v = (u \cdot \nabla v_1, u \cdot \nabla v_2, \dots, u \cdot \nabla v_d).$$

Then the claimed statement follows by (2.7) and the algebra Lemma 2.2. \square

3 Construction of quasi-periodic solutions

We look for quasi periodic solutions $u_\omega(t, x)$, $p_\omega(t, x)$ of the equation (1.1), oscillating with frequency $\omega = (\omega_1, \dots, \omega_\nu) \in \mathbb{R}^\nu$, namely we look for $u_\omega(t, x) := U(\omega t, x)$, $p_\omega(t, x) := P(\omega t, x)$ where $U : \mathbb{T}^\nu \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ and $P : \mathbb{T}^\nu \times \mathbb{T}^d \rightarrow \mathbb{R}$ are smooth functions. This leads to solve a functional equation for $U(\varphi, x)$, $P(\varphi, x)$ of the form

$$\begin{cases} \omega \cdot \partial_\varphi U - \Delta U + U \cdot \nabla U + \nabla P = \varepsilon f(\varphi, x) \\ \operatorname{div} U = 0. \end{cases} \quad (3.1)$$

If we take the divergence of the latter equation, one gets

$$\Delta P = \operatorname{div}(\varepsilon f - U \cdot \nabla U) \quad (3.2)$$

and by projecting on the space of zero divergence vector fields, one gets a closed equation for U of the form

$$\omega \cdot \partial_\varphi U - \Delta U + \mathfrak{L}(U \cdot \nabla U) = \varepsilon \mathfrak{L}(f), \quad U(\varphi, \cdot) \in \mathcal{D}_0(\mathbb{T}^d) \quad (3.3)$$

where we recall the definitions (2.12), (2.13). According to the splitting (2.10) and by applying the projectors π_0, π_0^\perp to the equation (3.3) one gets the decoupled equations

$$\omega \cdot \partial_\varphi U_0(\varphi) = \varepsilon f_0(\varphi) \quad (3.4)$$

and

$$\omega \cdot \partial_\varphi U_\perp - \Delta U_\perp + \mathfrak{L}(U_\perp \cdot \nabla U_\perp) = \varepsilon \mathfrak{L}(f_\perp). \quad (3.5)$$

Then, since ω is diophantine (see (1.3)) and using that

$$\int_{\mathbb{T}^\nu} f_0(\varphi) d\varphi = \int_{\mathbb{T}^\nu \times \mathbb{T}^d} f(\varphi, x) d\varphi dx \stackrel{(1.2)}{=} 0,$$

($\widehat{f}(0, 0) = 0$) the averaged equation (3.4) can be solved explicitly by setting

$$U_0(\varphi) := (\omega \cdot \partial_\varphi)^{-1} f_0(\varphi) = \sum_{\ell \in \mathbb{Z}^\nu \setminus \{0\}} \frac{\widehat{f}(\ell, 0)}{i\omega \cdot \ell} e^{i\ell \cdot \varphi}. \quad (3.6)$$

By (2.11) and using (1.3), one gets the estimate

$$\|U_0\|_\sigma \leq \varepsilon \gamma^{-1} \|f_0\|_{\sigma+\nu} \leq \varepsilon \gamma^{-1} \|f\|_{\sigma+\nu, 0}. \quad (3.7)$$

Remark 3.1 (Non resonance conditions). *The diophantine condition (1.3) on the frequency vector ω is used only to solve the averaged equation (3.4). In order to solve the equation (3.5) on the space of zero average functions (w.r. to x) no resonance conditions are required.*

We now solve the equation (3.5) by means of a fixed point argument. To this aim, we need to analyze some invertibility properties of the linear operator

$$L_\omega := \omega \cdot \partial_\varphi - \Delta. \quad (3.8)$$

Lemma 3.2 (Invertibility of L_ω). *Let $\sigma, s \geq 0$, $g \in H^\sigma(\mathbb{T}^\nu, H_0^s(\mathbb{T}^d, \mathbb{R}^d))$ and assume that g has zero divergence. Then there exists a unique $u := L_\omega^{-1} g \in H^\sigma(\mathbb{T}^\nu, H_0^{s+2}(\mathbb{T}^d, \mathbb{R}^d))$ with zero divergence which solves the equation $L_\omega u = g$. Moreover*

$$\|u\|_{\sigma, s+2} \leq \|g\|_{\sigma, s}. \quad (3.9)$$

Proof. By (2.5), we can write

$$L_\omega u(\varphi, x) = \sum_{\ell \in \mathbb{Z}^\nu} \sum_{j \in \mathbb{Z}^d \setminus \{0\}} (i\omega \cdot \ell + |j|^2) \widehat{u}(\ell, j) e^{i\ell \cdot \varphi} e^{ij \cdot x}.$$

Note that since $j \neq 0$, one has that

$$|i\omega \cdot \ell + |j|^2| = \sqrt{|\omega \cdot \ell|^2 + |j|^4} \geq |j|^2. \quad (3.10)$$

Hence, the equation $L_\omega u = g$ admits the unique solution with zero space average given by

$$u(\varphi, x) := L_\omega^{-1} g(\varphi, x) = \sum_{\ell \in \mathbb{Z}^\nu} \sum_{j \in \mathbb{Z}^d \setminus \{0\}} \frac{\widehat{g}(\ell, j)}{i\omega \cdot \ell + |j|^2} e^{i\ell \cdot \varphi} e^{ij \cdot x} \quad (3.11)$$

Clearly if $\operatorname{div}(g) = 0$ and then also $\operatorname{div}(u) = 0$. We now estimate $\|u\|_{\sigma, s+2}$. According to (2.6), (3.11), one has

$$\begin{aligned} \|u\|_{\sigma, s+2}^2 &= \sum_{\ell \in \mathbb{Z}^\nu} \sum_{j \in \mathbb{Z}^d \setminus \{0\}} \langle \ell \rangle^{2\sigma} \langle j \rangle^{2(s+2)} \frac{|\widehat{g}(\ell, j)|^2}{|i\omega \cdot \ell + |j|^2|^2} \\ &\stackrel{(3.10)}{\leq} \sum_{\ell \in \mathbb{Z}^\nu} \sum_{j \in \mathbb{Z}^d \setminus \{0\}} \langle \ell \rangle^{2\sigma} |j|^{2(s+2)} |j|^{-4} |\widehat{g}(\ell, j)|^2 = \|g\|_{\sigma, s}^2 \end{aligned}$$

which proves the claimed statement. \square

We now implement the fixed point argument for the equation (3.5) (to simplify notations we write U instead of U_\perp). For any $\sigma, s, R \geq 0$, we define the ball

$$\mathcal{B}_{\sigma,s}(R) := \left\{ U \in H^\sigma(\mathbb{T}^\nu, H_0^s(\mathbb{T}^d, \mathbb{R}^d)) : \operatorname{div}(U) = 0, \quad \|U\|_{\sigma,s} \leq R \right\}. \quad (3.12)$$

and we define the nonlinear operator

$$\Phi(U) := L_\omega^{-1} \mathfrak{L}(\varepsilon f - U \cdot \nabla U), \quad U \in \mathcal{B}_{\sigma,s}(R). \quad (3.13)$$

The following Proposition holds.

Proposition 3.3 (Contraction for Φ). *Let $\sigma > \nu/2$, $s > d/2 + 1$, $f \in \mathcal{C}^N(\mathbb{T}^\nu \times \mathbb{T}^d, \mathbb{R}^d)$, $N > \sigma + s - 2$. Then there exists a constant $C_* = C_*(f, \sigma, s) > 0$ large enough and $\varepsilon_0 = \varepsilon_0(f, \sigma, s) \in (0, 1)$ small enough, such that for any $\varepsilon \in (0, \varepsilon_0)$, the map $\Phi : \mathcal{B}_{\sigma,s}(C_*\varepsilon) \rightarrow \mathcal{B}_{\sigma,s}(C_*\varepsilon)$ is a contraction.*

Proof. Let $U \in \mathcal{B}_{\sigma,s}(C_*\varepsilon)$. We apply Lemmata 2.4-(i), 3.2 from which one immediately deduces that

$$\int_{\mathbb{T}^3} \Phi(U) dx = 0, \quad \operatorname{div}(\Phi(U)) = 0. \quad (3.14)$$

Moreover

$$\begin{aligned} \|\Phi(U)\|_{\sigma,s} &= \left\| L_\omega^{-1} \mathfrak{L}(\varepsilon f - U \cdot \nabla U) \right\|_{\sigma,s} \stackrel{(2.18), (3.9)}{\lesssim} \left\| \varepsilon f - U \cdot \nabla U \right\|_{\sigma,s-2} \\ &\lesssim \varepsilon \|f\|_{\sigma,s-2} + \|U \cdot \nabla U\|_{\sigma,s-1}. \end{aligned}$$

Note that since $f \in \mathcal{C}^N$ with $N > \sigma + s - 2$, one has that $\|f\|_{\sigma,s-2} \lesssim \|f\|_{\mathcal{C}^N}$. In view of Lemma 2.4-(ii), using that $\sigma > \nu/2$, $s - 1 > d/2$, one gets that

$$\|\Phi(U)\|_{\sigma,s} \leq C(f, s, \sigma)(\varepsilon + \|U\|_{\sigma,s-1}\|U\|_{\sigma,s}) \leq C(f, s, \sigma)(\varepsilon + \|U\|_{\sigma,s}^2)$$

for some constant $C(f, s, \sigma) > 0$. Using that $\|U\|_{\sigma,s} \leq C_*\varepsilon$, one gets that

$$\|\Phi(U)\|_{\sigma,s} \leq C(f, s, \sigma)\varepsilon + C(f, s, \sigma)C_*^2\varepsilon^2 \leq C_*\varepsilon$$

provided

$$C_* \geq 2C(f, s, \sigma) \quad \text{and} \quad \varepsilon \leq \frac{1}{2C(f, s, \sigma)C_*}.$$

Hence $\Phi : \mathcal{B}_{\sigma,s}(C_*\varepsilon) \rightarrow \mathcal{B}_{\sigma,s}(C_*\varepsilon)$. Now let $U_1, U_2 \in \mathcal{B}_{\sigma,s}(C_*\varepsilon)$ and we estimate

$$\Phi(U_1) - \Phi(U_2) = L_\omega^{-1} \mathfrak{L}(U_1 \cdot \nabla U_1 - U_2 \cdot \nabla U_2).$$

One has that

$$\begin{aligned} \|\Phi(U_1) - \Phi(U_2)\|_{\sigma,s} &\leq \left\| L_\omega^{-1} \mathfrak{L}((U_1 - U_2) \cdot \nabla U_1) \right\|_{\sigma,s} + \left\| L_\omega^{-1} \mathfrak{L}(U_2 \cdot \nabla(U_1 - U_2)) \right\|_{\sigma,s} \\ &\stackrel{(2.18), (3.9), \text{Lemma 2.4}}{\lesssim_{s,\sigma}} \|U_1 - U_2\|_{\sigma,s} \|U_1\|_{\sigma,s} + \|U_1 - U_2\|_{\sigma,s} \|U_2\|_{\sigma,s} \\ &\leq C(s, \sigma)(\|U_1\|_{\sigma,s} + \|U_2\|_{\sigma,s}) \|U_1 - U_2\|_{\sigma,s} \end{aligned}$$

for some constant $C(s, \sigma) > 0$. Since $U_1, U_2 \in \mathcal{B}_{\sigma, s}(C_*\varepsilon)$, one then has that

$$\|\Phi(U_1) - \Phi(U_2)\|_{\sigma, s} \leq 2C(s, \sigma)C_*\varepsilon\|U_1 - U_2\|_{\sigma, s} \leq \frac{1}{2}\|U_1 - U_2\|_{\sigma, s}$$

provided $\varepsilon \leq \frac{1}{4C(s, \sigma)C_*}$. Hence Φ is a contraction. \square

3.1 Proof of Theorem 1.1

Proposition 3.3 implies that for $\sigma > \nu/2$, $s > \frac{d}{2} + 1$, there exists a unique $U_\perp \in H^\sigma(\mathbb{T}^\nu, H_0^s(\mathbb{T}^d, \mathbb{R}^d))$, $\|U_\perp\|_{\sigma, s} \lesssim_{\sigma, s} \varepsilon$ which is a fixed point of the map Φ defined in (3.13). We fix $\sigma := \nu/2 + 2$ and $N > \frac{3\nu}{2} + s + 2$. By the Sobolev embedding property (2.8), since $\sigma - 1 > \nu/2$, one gets that

$$U_\perp \in \mathcal{C}_b^1(\mathbb{T}^\nu, H_0^s(\mathbb{T}^d, \mathbb{R}^d)), \quad \|U_\perp\|_{\mathcal{C}_\varphi^1 H_x^s} \lesssim_s \varepsilon \quad (3.15)$$

and U_\perp is a classical solution of the equation (3.5). Similarly, by recalling (3.4), (3.6), (3.7), one gets that

$$U_0 \in \mathcal{C}^1(\mathbb{T}^\nu, \mathbb{R}^d), \quad \|U_0\|_{\mathcal{C}_\varphi^1} \leq \varepsilon\gamma^{-1}\|f\|_{\frac{3\nu}{2}+2, 0} \leq \varepsilon\gamma^{-1}\|f\|_{\mathcal{C}^N}, \quad \int_{\mathbb{T}^\nu} U_0(\varphi) d\varphi = 0 \quad (3.16)$$

and U_0 is a classical solution of the equation (3.4). Hence $U = U_0 + U_\perp \in \mathcal{C}^1(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^d))$ is a classical solution of (3.3) and it satisfies $\int_{\mathbb{T}^\nu \times \mathbb{T}^d} U(\varphi, x) d\varphi dx = 0$. The unique solution with zero average in x of the equation (3.2) is given by

$$P := (-\Delta)^{-1} \operatorname{div}(U \cdot \nabla U - \varepsilon f).$$

Hence, $P \in \mathcal{C}_b^0(\mathbb{T}^\nu, H_0^s(\mathbb{T}^d, \mathbb{R}^d))$ and

$$\begin{aligned} \|P\|_{\mathcal{C}_\varphi^0 H_x^s} &\stackrel{(2.8), \sigma=\nu/2+2}{\lesssim} \|P\|_{\sigma, s} \lesssim_{\sigma, s} \varepsilon\|f\|_{\sigma, s-1} + \|U \cdot \nabla U\|_{\sigma, s-1} \\ &\stackrel{\text{Lemma 2.4}}{\lesssim_{\sigma, s}} \varepsilon\|f\|_{\sigma, s-1} + \|U\|_{\sigma, s}^2. \end{aligned}$$

The claimed estimate on P then follows since $\|f\|_{\sigma, s-1} \lesssim \|f\|_{\mathcal{C}^N}$, $\|U\|_{\sigma, s} \leq C_*\varepsilon$. Note that if f has zero average in x , one has that

$$f_0(\varphi) = \pi_0 f(\varphi) = \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d} f(\varphi, x) dx = 0, \quad \forall \varphi \in \mathbb{T}^\nu.$$

The equation (3.4) reduces to $\omega \cdot \partial_\varphi U_0 = 0$. Hence the only solution $U = U_0 + U_\perp$ of (3.3) with zero average in x is the one where we choose $U_0 = 0$ and hence $U = U_\perp$. The claimed statement has then been proved.

4 Orbital and asymptotic stability

We now want to study the Cauchy problem for the equation (1.1) for initial data which are close to the quasi-periodic solution (u_ω, p_ω) , where

$$u_\omega(t, x) := U(\omega t, x), \quad p_\omega(t, x) := P(\omega t, x) \quad (4.1)$$

and the periodic functions $U \in \mathcal{C}^1(\mathbb{T}^\nu, H^s(\mathbb{T}^d, \mathbb{R}^d))$, $P \in \mathcal{C}^0(\mathbb{T}^\nu, H_0^s(\mathbb{T}^d, \mathbb{R}^d))$ are given by Theorem 1.1. We then look for solutions which are perturbations of the quasi-periodic ones (u_ω, p_ω) , namely we look for solutions of the form

$$u(t, x) = u_\omega(t, x) + v(t, x), \quad p(t, x) = p_\omega(t, x) + q(t, x). \quad (4.2)$$

Plugging the latter ansatz into the equation (1.1), one obtains an equation for $v(t, x)$, $q(t, x)$ of the form

$$\begin{cases} \partial_t v - \Delta v + u_\omega \cdot \nabla v + v \cdot \nabla u_\omega + v \cdot \nabla v + \nabla q = 0 \\ \operatorname{div}(v) = 0. \end{cases} \quad (4.3)$$

If we take the divergence in the latter equation we get the equation for the pressure $q(t, x)$

$$-\Delta q = \operatorname{div}\left(u_\omega \cdot \nabla v + v \cdot \nabla u_\omega + v \cdot \nabla v\right). \quad (4.4)$$

By using the Leray projector defined in (2.13), we then get a closed equation for v of the form

$$\begin{cases} \partial_t v - \Delta v + \mathfrak{L}\left(u_\omega \cdot \nabla v + v \cdot \nabla u_\omega + v \cdot \nabla v\right) = 0 \\ \operatorname{div}(v) = 0. \end{cases} \quad (4.5)$$

We prove the following

Proposition 4.1. *Let $s > d/2 + 1$, $\alpha \in (0, 1)$. Then there exists $\delta = \delta(s, \alpha, d, \nu) \in (0, 1)$ small enough and $C = C(s, \alpha, d, \nu) > 0$ large enough, such that for any $\varepsilon \in (0, \delta)$ and for any initial datum $v_0 \in H_0^s(\mathbb{T}^d, \mathbb{R}^d)$ with $\|v_0\|_{H_x^s} \leq \delta$, there exists a unique global classical solution*

$$v \in \mathcal{C}_b^0([0, +\infty), H_0^s(\mathbb{T}^d, \mathbb{R}^d)) \cap \mathcal{C}_b^1([0, +\infty), H_0^{s-2}(\mathbb{T}^d, \mathbb{R}^d)) \quad (4.6)$$

of the equation (4.5) which satisfies

$$\|v(t, \cdot)\|_{H_x^s}, \|\partial_t v(t, \cdot)\|_{H_x^{s-2}} \leq C\delta e^{-\alpha t}, \quad \forall t \geq 0. \quad (4.7)$$

The Proposition above will be proved by a fixed point argument in some weighted Sobolev spaces which take care of the decay in time of the solutions we are looking for. In the next section we shall exploit some decay estimates of the linear heat propagator which will be used in the proof of our result.

4.1 Dispersive estimates for the heat propagator

In this section we analyze some properties of the heat propagator. We recall that the heat propagator is defined as follows. Consider the Cauchy problem for the heat equation

$$\begin{cases} \partial_t u - \Delta u = 0 \\ u(0, x) = u_0(x), \end{cases} \quad u_0 \in H_0^s(\mathbb{T}^d, \mathbb{R}^d). \quad (4.8)$$

It is well known that there exists a unique solution

$$u \in \mathcal{C}_b^0([0, +\infty), H_0^s(\mathbb{T}^d, \mathbb{R}^d)) \cap \mathcal{C}_b^1([0, +\infty), H_0^{s-2}(\mathbb{T}^d, \mathbb{R}^d))$$

which can be written as $u(t, x) := e^{t\Delta} u_0(x)$, namely

$$u(t, x) = e^{t\Delta} u_0(x) = \sum_{\xi \in \mathbb{Z}^d \setminus \{0\}} e^{-t|\xi|^2} \widehat{u}_0(\xi) e^{ix \cdot \xi}. \quad (4.9)$$

Lemma 4.2. (i) Let $u_0 \in H_0^s(\mathbb{T}^d, \mathbb{R}^d)$. Then

$$\|e^{t\Delta} u_0\|_{H_x^s} \leq e^{-t} \|u_0\|_{H_x^s}, \quad \forall t \geq 0. \quad (4.10)$$

(ii) Let $u_0 \in H_0^{s-1}(\mathbb{T}^d, \mathbb{R}^d)$. Then, for any integer $n \geq 1$ and for any $\alpha \in (0, 1)$,

$$\|e^{t\Delta} u_0\|_{H_x^s} \lesssim_n t^{-\frac{n}{2}} (1 - \alpha)^{-\frac{n}{2}} e^{-\alpha t} \|u_0\|_{H_x^{s-n}} \lesssim_n t^{-\frac{n}{2}} (1 - \alpha)^{-\frac{n}{2}} e^{-\alpha t} \|u_0\|_{H_x^{s-1}}, \quad \forall t > 0. \quad (4.11)$$

Proof. The item (i) follows by (4.9), using that $e^{-t|\xi|^2} \leq e^{-t}$ for any $t \geq 0$, $\xi \in \mathbb{Z}^d \setminus \{0\}$, since $|\xi|^2 \geq 1$. We now prove the item (ii). Let $n \in \mathbb{N}$, $\alpha \in (0, 1)$. One has

$$\begin{aligned} \|e^{t\Delta} u_0\|_{H_x^s}^2 &= \sum_{\xi \in \mathbb{Z}^d \setminus \{0\}} |\xi|^{2s} e^{-2t|\xi|^2} |\widehat{u}_0(\xi)|^2 \\ &= \sum_{\xi \in \mathbb{Z}^d \setminus \{0\}} e^{-2\alpha t|\xi|^2} |\xi|^{2(s-n)} |\xi|^{2n} e^{-2(1-\alpha)t|\xi|^2} |\widehat{u}_0(\xi)|^2. \end{aligned} \quad (4.12)$$

Using that for any $\xi \in \mathbb{Z}^d \setminus \{0\}$, $t \geq 0$, $e^{-2\alpha t|\xi|^2} \leq e^{-2\alpha t}$, by (4.12), one gets that

$$\|e^{t\Delta} u_0\|_{H_x^s}^2 \leq e^{-2\alpha t} \sum_{\xi \in \mathbb{Z}^d \setminus \{0\}} |\xi|^{2(s-n)} |\xi|^{2n} e^{-2(1-\alpha)t|\xi|^2} |\widehat{u}_0(\xi)|^2 \quad (4.13)$$

By Lemma A.1 (applied with $\zeta = 2(1 - \alpha)t$), one has that

$$\sup_{\xi \in \mathbb{Z}^d \setminus \{0\}} |\xi|^{2n} e^{-2(1-\alpha)t|\xi|^2} \leq \sup_{y \geq 0} y^n e^{-2(1-\alpha)ty} \leq \frac{C(n)}{(1 - \alpha)^{nt^n}}$$

for some constant $C(n) > 0$. Therefore by (4.13), one gets that

$$\begin{aligned} \|e^{t\Delta} u_0\|_{H_x^s}^2 &\lesssim_n t^{-n} e^{-2\alpha t} (1 - \alpha)^{-n} \sum_{\xi \in \mathbb{Z}^d \setminus \{0\}} |\xi|^{2(s-n)} |\widehat{u}_0(\xi)|^2 \\ &\lesssim_n t^{-n} e^{-2\alpha t} (1 - \alpha)^{-n} \|u_0\|_{H_x^{s-n}}^2. \end{aligned}$$

The second inequality in (4.11) clearly follows since $\|\cdot\|_{H_x^{s-n}} \leq \|\cdot\|_{H_x^{s-1}}$ for $n \geq 1$. \square

We fix $\alpha \in (0, 1)$ and for any $s \geq 0$, we define the space

$$\mathcal{E}_s := \left\{ u \in \mathcal{C}_b^0([0, +\infty), H_0^s(\mathbb{T}^d, \mathbb{R}^d)) : \|u\|_{\mathcal{E}_s} := \sup_{t \geq 0} e^{\alpha t} \|u(t)\|_{H_x^s} \right\}. \quad (4.14)$$

Clearly

$$\|\cdot\|_{\mathcal{E}_s} \leq \|\cdot\|_{\mathcal{E}_{s'}}, \quad \forall s \leq s'. \quad (4.15)$$

The following elementary lemma holds:

Lemma 4.3. (i) *Let $u \in \mathcal{E}_s$. Then*

$$\begin{aligned} \|u\|_{\mathcal{C}_t^0 H_x^s} &\lesssim \|u\|_{\mathcal{E}_s} \quad \text{and} \quad \|\mathfrak{L}(u)\|_{\mathcal{E}_s} \lesssim \|u\|_{\mathcal{E}_s}, \\ \|u(t)\|_{H_x^s} &\leq e^{-\alpha t} \|u\|_{\mathcal{E}_s}, \quad \forall t \geq 0. \end{aligned} \quad (4.16)$$

(ii) *Let $s > d/2$, $u \in \mathcal{E}_s$, $v \in \mathcal{C}_b^0([0, +\infty), H^{s+1}(\mathbb{T}^d, \mathbb{R}^d))$, $\operatorname{div}(u) = 0$. Then the product $u \cdot \nabla v \in \mathcal{E}_s$ and*

$$\|u \cdot \nabla v\|_{\mathcal{E}_s} \lesssim_s \|u\|_{\mathcal{E}_s} \|v\|_{\mathcal{C}_t^0 H_x^{s+1}}. \quad (4.17)$$

(iii) *Let $s > d/2$, $u \in \mathcal{C}_b^0([0, +\infty), H^s(\mathbb{T}^d, \mathbb{R}^d))$, $\operatorname{div}(u) = 0$ and $v \in \mathcal{E}_{s+1}$. Then the product $u \cdot \nabla v \in \mathcal{E}_s$ and*

$$\|u \cdot \nabla v\|_{\mathcal{E}_s} \lesssim_s \|u\|_{\mathcal{C}_t^0 H_x^s} \|v\|_{\mathcal{E}_{s+1}}. \quad (4.18)$$

(iv) *Let $s > d/2$, $u \in \mathcal{E}_s$, $\operatorname{div}(u) = 0$, $v \in \mathcal{E}_{s+1}$. Then $u \cdot \nabla v \in \mathcal{E}_s$ and*

$$\|u \cdot \nabla v\|_{\mathcal{E}_s} \lesssim_s \|u\|_{\mathcal{E}_s} \|v\|_{\mathcal{E}_{s+1}}. \quad (4.19)$$

Proof. The item (i) is very elementary and it follows in a straightforward way by the definition (4.14) and by recalling the estimate (2.18) on the Leray projector \mathfrak{L} . We prove the item (ii). By Lemma 2.4-(i), one has that $u(t) \cdot \nabla v(t)$ has zero average in x . Moreover

$$u \cdot \nabla v = \left(u \cdot \nabla v_1, \dots, u \cdot \nabla v_d \right)$$

therefore, since $s > d/2$, by Lemma 2.1 and using (2.1) one has that for any $t \in [0, +\infty)$ $\|u(t) \cdot \nabla v(t)\|_{H_x^s} \lesssim_s \|u(t)\|_{H_x^s} \|v(t)\|_{H_x^{s+1}}$ implying that

$$\begin{aligned} e^{\alpha t} \|u(t) \cdot \nabla v(t)\|_{H_x^s} &\lesssim_s e^{\alpha t} \|u(t)\|_{H_x^s} \|v(t)\|_{H_x^{s+1}} \lesssim_s \left(\sup_{t \geq 0} e^{\alpha t} \|u(t)\|_{H_x^s} \right) \left(\sup_{t \geq 0} \|v(t)\|_{H_x^{s+1}} \right) \\ &\lesssim_s \|u\|_{\mathcal{E}_s} \|v\|_{\mathcal{C}_t^0 H_x^{s+1}}. \end{aligned} \quad (4.20)$$

Passing to the supremum over $t \geq 0$ in the left hand side of (4.20), we get the claimed statement. The item (iii) follows by similar arguments and the item (iv) follows by applying items (i) and (ii). \square

We now prove some estimates for the heat propagator $e^{t\Delta}$ in the space \mathcal{E}_s .

Lemma 4.4. *Let $s \geq 0$, $u_0 \in H_0^s(\mathbb{T}^d, \mathbb{R}^d)$. Then $\|e^{t\Delta} u_0\|_{\mathcal{E}_s} \lesssim_\alpha \|u_0\|_{H_x^s}$.*

Proof. We have to estimate uniformly w.r. to $t \geq 0$, the quantity $e^{\alpha t} \|e^{t\Delta} u_0\|_{H_x^s}$. For $t \in [0, 1]$, since $e^{\alpha t} \leq e^\alpha \stackrel{\alpha \leq 1}{\leq} e$ and by applying Lemma 4.2-(i), one gets that

$$e^{\alpha t} \|e^{t\Delta} u_0\|_{H_x^s} \lesssim \|e^{t\Delta} u_0\|_{H_x^s} \lesssim \|u_0\|_{H_x^s}, \quad \forall t \in [0, 1]. \quad (4.21)$$

For $t > 1$, by applying Lemma 4.2-(ii) (for $n = 1$), one gets that

$$e^{\alpha t} \|e^{t\Delta} u_0\|_{H_x^s} \lesssim_\alpha t^{-\frac{1}{2}} \|u_0\|_{H_x^{s-1}} \lesssim_\alpha \|u_0\|_{H_x^s}, \quad \forall t > 1. \quad (4.22)$$

Hence the claimed statement follows by (4.21), (4.22) passing to the supremum over $t \geq 0$. \square

The main result of this section is the following Proposition

Proposition 4.5. *Let $s \geq 1$, $f \in \mathcal{E}_{s-1}$ and define*

$$u(t) \equiv u(t, \cdot) := \int_0^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau. \quad (4.23)$$

Then $u \in \mathcal{E}_s$ and

$$\|u\|_{\mathcal{E}_s} \lesssim_\alpha \|f\|_{\mathcal{E}_{s-1}}. \quad (4.24)$$

The proof is split in several steps. The first step is to estimate the integral in (4.23) for any $t \in [0, 1]$.

Lemma 4.6. *Let $t \in [0, 1]$, $f \in \mathcal{E}_{s-1}$ and u defined by (4.23). Then*

$$\|u(t)\|_{H_x^s} \lesssim_\alpha \|f\|_{\mathcal{C}_t^0 H_x^{s-1}}.$$

Proof. Let $t \in [0, 1]$. Then

$$\begin{aligned} \|u(t)\|_{H_x^s} &\leq \int_0^t \left\| e^{(t-\tau)\Delta} f(\tau, \cdot) \right\|_{H_x^s} d\tau \stackrel{(4.11)}{\lesssim_\alpha} \int_0^t e^{-\frac{t-\tau}{2}} \frac{\|f(\tau, \cdot)\|_{H_x^{s-1}}}{\sqrt{t-\tau}} d\tau \\ &\stackrel{e^{-\frac{t-\tau}{2}} \leq 1}{\lesssim_\alpha} \int_0^t \frac{1}{\sqrt{t-\tau}} d\tau \|f\|_{\mathcal{C}_t^0 H_x^{s-1}}. \end{aligned} \quad (4.25)$$

By making the change of variables $z = t - \tau$, one gets that

$$\int_0^t \frac{1}{\sqrt{t-\tau}} d\tau = \int_0^t \frac{1}{\sqrt{z}} dz \stackrel{t \leq 1}{\leq} \int_0^1 \frac{1}{\sqrt{z}} dz = 2$$

and hence in view of (4.25), one gets $\|u(t)\|_{H_x^s} \lesssim_\alpha \|f\|_{\mathcal{C}_t^0 H_x^{s-1}}$ for any $t \in [0, 1]$, which is the claimed statement. \square

For $t > 1$, we split the integral term in (4.23) as

$$\int_0^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau = \int_0^{t-\frac{1}{2}} e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau + \int_{t-\frac{1}{2}}^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau$$

and we estimate separately the two terms in the latter formula. More precisely the first term is estimated in Lemma 4.7 and the second one in Lemma 4.8.

Lemma 4.7. *Let $t > 1$. Then*

$$\left\| \int_0^{t-\frac{1}{2}} e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} \lesssim_\alpha e^{-\alpha t} \|f\|_{\mathcal{E}_{s-1}}$$

Proof. Let $t > 1$. One then has

$$\begin{aligned} \left\| \int_0^{t-\frac{1}{2}} e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} &\leq \int_0^{t-\frac{1}{2}} \left\| e^{(t-\tau)\Delta} f(\tau, \cdot) \right\|_{H_x^s} d\tau \\ &\stackrel{(4.11)}{\lesssim_{n,\alpha}} \int_0^{t-\frac{1}{2}} e^{-\alpha(t-\tau)} \frac{1}{(t-\tau)^{\frac{n}{2}}} \|f(\tau, \cdot)\|_{H_x^{s-1}} d\tau \\ &\lesssim_{n,\alpha} e^{-\alpha t} \int_0^{t-\frac{1}{2}} \frac{1}{(t-\tau)^{\frac{n}{2}}} e^{\alpha\tau} \|f(\tau, \cdot)\|_{H_x^{s-1}} d\tau \\ &\lesssim_{n,\alpha} e^{-\alpha t} \left(\int_0^{t-\frac{1}{2}} \frac{d\tau}{(t-\tau)^{\frac{n}{2}}} \right) \sup_{\tau \geq 0} e^{\alpha\tau} \|f(\tau, \cdot)\|_{H_x^{s-1}}. \end{aligned} \tag{4.26}$$

By choosing $n = 4$ and by making the change of variables $z = t - \tau$, one gets that

$$\int_0^{t-\frac{1}{2}} \frac{d\tau}{(t-\tau)^{\frac{n}{2}}} = \int_0^{t-\frac{1}{2}} \frac{d\tau}{(t-\tau)^2} = \int_{\frac{1}{2}}^t \frac{dz}{z^2} \leq \int_{\frac{1}{2}}^{+\infty} \frac{dz}{z^2} < \infty.$$

The latter estimate, together with the estimate (4.26) imply the claimed statement. \square

Lemma 4.8. *Let $t > 1$. Then*

$$\left\| \int_{t-\frac{1}{2}}^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} \lesssim_\alpha e^{-\alpha t} \|f\|_{\mathcal{E}_{s-1}}.$$

Proof. One has

$$\begin{aligned} \left\| \int_{t-\frac{1}{2}}^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} &\leq \int_{t-\frac{1}{2}}^t \left\| e^{(t-\tau)\Delta} f(\tau, \cdot) \right\|_{H_x^s} d\tau \\ &\stackrel{(4.11)}{\lesssim_\alpha} e^{-\alpha t} \int_{t-\frac{1}{2}}^t \frac{1}{\sqrt{t-\tau}} e^{\alpha\tau} \|f(\tau, \cdot)\|_{H_x^{s-1}} d\tau \\ &\lesssim_\alpha e^{-\alpha t} \int_{t-\frac{1}{2}}^t \frac{d\tau}{\sqrt{t-\tau}} \left(\sup_{\tau \geq 0} e^{\alpha\tau} \|f(\tau, \cdot)\|_{H_x^{s-1}} \right). \end{aligned} \tag{4.27}$$

By making the change of variables $z = t - \tau$, one gets

$$\int_{t-\frac{1}{2}}^t \frac{d\tau}{\sqrt{t-\tau}} = \int_0^{\frac{1}{2}} \frac{dz}{\sqrt{z}} = \sqrt{2}.$$

The latter estimate, together with (4.27) imply the claimed statement. \square

PROOF OF PROPOSITION 4.5.

Proof. For any $t \in [0, 1]$, since $e^{\alpha t} \leq e^\alpha \stackrel{\alpha < 1}{\leq} e$, by Lemma 4.6, one has that

$$\begin{aligned} e^{\alpha t} \left\| \int_0^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} &\lesssim \left\| \int_0^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} \\ &\stackrel{(4.16)}{\lesssim_\alpha} \|f\|_{\mathcal{C}_t^0 H_x^{s-1}} \lesssim_\alpha \|f\|_{\mathcal{E}_{s-1}}. \end{aligned} \quad (4.28)$$

For any $t > 1$, by applying Lemmata 4.7, 4.8, one gets

$$\begin{aligned} e^{\alpha t} \left\| \int_0^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} &\leq e^{\alpha t} \left\| \int_0^{t-\frac{1}{2}} e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} \\ &\quad + e^{\alpha t} \left\| \int_{t-\frac{1}{2}}^t e^{(t-\tau)\Delta} f(\tau, \cdot) d\tau \right\|_{H_x^s} \\ &\lesssim_\alpha \|f\|_{\mathcal{E}_{s-1}}. \end{aligned} \quad (4.29)$$

The claimed statement then follows by (4.28), (4.29) and passing to the supremum over $t \in [0, +\infty)$. \square

4.2 Proof of Proposition 4.1

The Proposition 4.1 is proved by a fixed point argument. For any $\delta > 0$ and $s \geq 0$, we define the ball

$$\mathcal{B}_s(\delta) := \left\{ v \in \mathcal{E}_s : \operatorname{div}(v) = 0, \quad \|v\|_{\mathcal{E}_s} \leq \delta \right\} \quad (4.30)$$

and for any $v \in \mathcal{B}_s(\delta)$, we define the map

$$\begin{aligned} \Phi(v) &:= e^{t\Delta} v_0 + \int_0^t e^{(t-\tau)\Delta} \mathcal{N}(v)(\tau, \cdot) d\tau, \\ \mathcal{N}(v) &:= -\mathfrak{L}\left(u_\omega \cdot \nabla v + v \cdot \nabla u_\omega + v \cdot \nabla v\right). \end{aligned} \quad (4.31)$$

Proposition 4.9. *Let $s > d/2 + 1$, $\alpha \in (0, 1)$. Then there exists $\delta = \delta(s, \alpha, \nu, d) \in (0, 1)$ small enough such that for any $\varepsilon \in (0, \delta)$, $\Phi : \mathcal{B}_s(\delta) \rightarrow \mathcal{B}_s(\delta)$ is a contraction.*

Proof. Since v_0 has zero divergence and zero average, then clearly by (4.9), $\operatorname{div}(e^{t\Delta} v_0) = 0$, $\int_{\mathbb{T}^3} e^{t\Delta} v_0(x) dx = 0$. Now let $v(t, x)$ be a function with zero average and zero divergence. Clearly $\operatorname{div}(\mathcal{N}(v)) = 0$ since in the definition of $\mathcal{N}(v)$ in (4.31), there is the Leray projector. Moreover using that $\operatorname{div}(v) = \operatorname{div}(u_\omega) = 0$, by Lemma 2.4-(i), one gets $\mathcal{N}(v)$ has zero average and then by (4.9) also $\int_0^t e^{(t-\tau)\Delta} \mathcal{N}(v)(\tau, \cdot) d\tau$ has zero average. Hence, we have shown that $\operatorname{div}(\Phi(v)) = 0$ and $\int_{\mathbb{T}^d} \Phi(v) dx = 0$. Let now $\|v\|_{\mathcal{E}_s} \leq \delta$. We estimate $\|\Phi(v)\|_{\mathcal{E}_s}$. By recalling (4.31), Lemma 4.4, Proposition 4.5 and Lemma 4.3, one has

$$\begin{aligned} \|\Phi(v)\|_{\mathcal{E}_s} &\stackrel{(4.31)}{\leq} \|e^{t\Delta} v_0\|_{\mathcal{E}_s} + \left\| \int_0^t e^{(t-\tau)\Delta} \mathcal{N}(v)(\tau, \cdot) d\tau \right\|_{\mathcal{E}_s} \\ &\lesssim_\alpha \|v_0\|_{H_x^s} + \|\mathcal{N}(v)\|_{\mathcal{E}_{s-1}} \\ &\lesssim_{s, \alpha} \|v_0\|_{H_x^s} + \|u_\omega\|_{\mathcal{C}_t^0 H_x^s} \|v\|_{\mathcal{E}_s} + \|v\|_{\mathcal{E}_s}^2. \end{aligned} \quad (4.32)$$

By the estimates of Theorem 1.1, by the definition (4.1) and by applying Lemma 2.3, one has

$$u_\omega \in \mathcal{C}_b^0(\mathbb{R}, H^s(\mathbb{T}^d, \mathbb{R}^d)) \quad \text{and} \quad \|u_\omega\|_{\mathcal{C}_t^0 H_x^s} \lesssim_s \varepsilon \lesssim_s^{\varepsilon \leq \delta} \delta. \quad (4.33)$$

Since $v \in \mathcal{B}_s(\delta)$, the estimate (4.32) implies that

$$\|\Phi(v)\|_{\mathcal{E}_s} \leq C(s, \alpha) \left(\|v_0\|_{H_x^s} + \delta^2 \right). \quad (4.34)$$

Hence $\|\Phi(v)\|_{\mathcal{E}_s} \leq \delta$ provided

$$C(s, \alpha) \|v_0\|_{H_x^s} \leq \frac{\delta}{2}, \quad C(s, \alpha) \delta \leq \frac{1}{2}.$$

These conditions are fulfilled by taking δ small enough and $\|v_0\|_{H_x^s} \ll \delta$. Hence $\Phi : \mathcal{B}_s(\delta) \rightarrow \mathcal{B}_s(\delta)$. Now let $v_1, v_2 \in \mathcal{B}_s(\delta)$. We need to estimate

$$\Phi(v_1) - \Phi(v_2) = \int_0^t e^{(t-\tau)\Delta} \left(\mathcal{N}(v_1) - \mathcal{N}(v_2) \right) (\tau, \cdot) d\tau. \quad (4.35)$$

By (4.31)

$$\begin{aligned} \|\mathcal{N}(v_1) - \mathcal{N}(v_2)\|_{\mathcal{E}_{s-1}} &\leq \left\| \mathfrak{L} \left(u_\omega \cdot \nabla(v_1 - v_2) \right) \right\|_{\mathcal{E}_{s-1}} + \left\| \mathfrak{L} \left((v_1 - v_2) \cdot \nabla u_\omega \right) \right\|_{\mathcal{E}_{s-1}} \\ &\quad + \left\| \mathfrak{L} \left((v_1 - v_2) \cdot \nabla v_1 \right) \right\|_{\mathcal{E}_{s-1}} + \left\| \mathfrak{L} \left(v_2 \cdot \nabla(v_1 - v_2) \right) \right\|_{\mathcal{E}_{s-1}} \\ &\stackrel{(4.16)}{\lesssim} \left\| u_\omega \cdot \nabla(v_1 - v_2) \right\|_{\mathcal{E}_{s-1}} + \left\| (v_1 - v_2) \cdot \nabla u_\omega \right\|_{\mathcal{E}_{s-1}} \\ &\quad + \left\| (v_1 - v_2) \cdot \nabla v_1 \right\|_{\mathcal{E}_{s-1}} + \left\| v_2 \cdot \nabla(v_1 - v_2) \right\|_{\mathcal{E}_{s-1}} \\ &\stackrel{(4.17), (4.18), (4.19)}{\lesssim_s} \left(\|u_\omega\|_{\mathcal{C}_t^0 H_x^s} + \|v_1\|_{\mathcal{E}_s} + \|v_2\|_{\mathcal{E}_s} \right) \|v_1 - v_2\|_{\mathcal{E}_s}. \end{aligned} \quad (4.36)$$

Hence, (4.33) and using that $v_1, v_2 \in \mathcal{B}_s(\delta)$ ($\|v_1\|_{\mathcal{E}_s}, \|v_2\|_{\mathcal{E}_s} \leq \delta$) and the estimate (4.36) imply that

$$\|\mathcal{N}(v_1) - \mathcal{N}(v_2)\|_{\mathcal{E}_{s-1}} \lesssim_s \delta \|v_1 - v_2\|_{\mathcal{E}_s}. \quad (4.37)$$

By (4.35), one gets that

$$\begin{aligned} \|\Phi(v_1) - \Phi(v_2)\|_{\mathcal{E}_s} &\stackrel{\text{Proposition 4.5}}{\lesssim_{s, \alpha}} \|\mathcal{N}(v_1) - \mathcal{N}(v_2)\|_{\mathcal{E}_{s-1}} \\ &\stackrel{(4.37)}{\leq} C(s, \alpha) \delta \|v_1 - v_2\|_{\mathcal{E}_s} \end{aligned} \quad (4.38)$$

for some constant $C(s, \alpha) > 0$. Therefore

$$\|\Phi(v_1) - \Phi(v_2)\|_{\mathcal{E}_s} \leq \frac{1}{2} \|v_1 - v_2\|_{\mathcal{E}_s}$$

provided $\delta \leq \frac{1}{2C(s, \alpha)}$. The claimed statement has then been proved. \square

PROOF OF PROPOSITION 4.1 CONCLUDED. By Proposition 4.9, using the contraction mapping theorem there exists a unique $v \in \mathcal{B}_s(\delta)$ which is a fixed point of the map Φ in (4.31). By the functional equation $v = \Phi(v)$, one deduces in a standard way that

$$v \in \mathcal{C}_b^1([0, +\infty), H_0^{s-2}(\mathbb{T}^d, \mathbb{R}^d))$$

and hence v is a classical solution of the equation (4.5). By (4.16) and using the trivial fact that $\|\Delta v\|_{\mathcal{E}_{s-2}} \leq \|v\|_{\mathcal{E}_s}$

$$\begin{aligned} \|\partial_t v\|_{\mathcal{E}_{s-2}} &\lesssim \|v\|_{\mathcal{E}_s} + \|u_\omega \cdot \nabla v\|_{\mathcal{E}_{s-2}} + \|v \cdot \nabla u_\omega\|_{\mathcal{E}_{s-2}} + \|v \cdot \nabla v\|_{\mathcal{E}_{s-2}} \\ &\stackrel{(4.17)-(4.19)}{\lesssim_s} \|v\|_{\mathcal{E}_s} \left(1 + \|u_\omega\|_{\mathcal{C}_t^0 H_x^s} + \|v\|_{\mathcal{E}_s}\right). \end{aligned} \quad (4.39)$$

Therefore using that $v \in \mathcal{B}_s(\delta)$ ($\|v\|_{\mathcal{E}_s} \leq \delta$) and by (4.33), $\|u_\omega\|_{\mathcal{C}_t^0 H_x^s} \lesssim_s \delta$, one gets, for δ small enough, the estimate $\|\partial_t v\|_{\mathcal{E}_{s-2}} \lesssim_s \delta$ and the claimed statement follows by recalling (4.16).

4.3 Proof of Theorem 1.2

In view of Proposition 4.1, it remains only to solve the equation (4.4) for the pressure $q(t, x)$. The only solution with zero average of this latter equation is given by

$$q := (-\Delta)^{-1} \left(\operatorname{div} \left(u_\omega \cdot \nabla v + v \cdot \nabla u_\omega + v \cdot \nabla v \right) \right). \quad (4.40)$$

Using that $\|(-\Delta)^{-1} \operatorname{div} a\|_{\mathcal{E}_s} \lesssim \|a\|_{\mathcal{E}_{s-1}}$ for any $a \in \mathcal{E}_s$, one gets the inequality

$$\|q\|_{\mathcal{E}_s} \lesssim \|u_\omega \cdot \nabla v\|_{\mathcal{E}_{s-1}} + \|v \cdot \nabla u_\omega\|_{\mathcal{E}_{s-1}} + \|v \cdot \nabla v\|_{\mathcal{E}_{s-1}}. \quad (4.41)$$

Hence arguing as in (4.39), one deduces the estimate $\|q\|_{\mathcal{E}_s} \lesssim_s \delta$. The claimed estimate on q then follows by recalling (4.16) and the proof is concluded (recall that by (4.2), $v = u - u_\omega$, $q = p - p_\omega$).

A appendix

Lemma A.1. *Let $n \in \mathbb{N}$, $\zeta > 0$ and $f : [0, +\infty] \rightarrow [0, +\infty]$ defined by $f(y) := y^n e^{-\zeta y}$. Then*

$$\max_{y \geq 0} f(y) = (n/\zeta)^n e^{-n}$$

Proof. One has that $f \geq 0$ and

$$f(0) = 0, \quad \lim_{y \rightarrow +\infty} f(y) = 0.$$

Moreover

$$f'(y) = y^{n-1} e^{-y} (n - \zeta y)$$

therefore f admits a global maximum at $y = n/\zeta$. This implies that

$$\max_{y \geq 0} f(y) = f(n/\zeta) = (n/\zeta)^n e^{-n}$$

and the lemma follows. \square

References

- [1] P. BALDI P., M. BERTI, E. HAUS, R. MONTALTO, *Time quasi-periodic gravity water waves in finite depth*, Inventiones Math. 214 (2), 739–911, 2018.
- [2] P. BALDI, M. BERTI, R. MONTALTO, *KAM for quasi-linear and fully nonlinear forced perturbations of Airy equation*, Math. Annalen 359, 471–536, 2014.
- [3] P. BALDI, R. MONTALTO. *Quasi-periodic incompressible Euler flows in 3D*. Preprint arXiv:2003.14313, 2020.
- [4] D. BAMBUSI, B. GREBERT, A. MASPERO, D. ROBERT. *Growth of Sobolev norms for abstract linear Schrödinger Equations*. To appear in JEMS. Preprint arXiv:1706.09708, 2017.
- [5] D. BAMBUSI, B. GREBERT, A. MASPERO, AND D. ROBERT. *Reducibility of the quantum Harmonic oscillator in d -dimensions with polynomial time dependent perturbation*. Analysis and PDEs, 11(3):775–799, 2018.
- [6] D. BAMBUSI, B. LANGELLA, R. MONTALTO. *Reducibility of non-resonant transport equation on \mathbb{T}^d with unbounded perturbations*. Ann. I. H. Poincaré 20, 1893–1929 (2019). <https://doi.org/10.1007/s00023-019-00795-2>.
- [7] M. BERTI, *KAM for PDEs*, Boll. Unione Mat. Ital. (2016) 9:115-142.
- [8] M. BERTI M. P. BOLLE. *Quasi-periodic solutions with Sobolev regularity of NLS on \mathbb{T}^d with a multiplicative potential*. Eur. Jour. Math. 15, 229-286 (2013).
- [9] M. BERTI, P. BOLLE. *Sobolev quasi-periodic solutions of multidimensional wave equations with a multiplicative potential*. Nonlinearity 25 (9), pp. 2579–2613, 2012.
- [10] M. BERTI, L. CORSI, M. PROCESI *An abstract Nash-Moser theorem and quasi-periodic solutions for NLW and NLS on compact Lie groups and homogeneous manifolds*. Comm. Math. Phys., vol. 334, n. 3, pp. 1413-1454, 2015.
- [11] M. BERTI, R. MONTALTO, *Quasi-periodic standing wave solutions of gravity capillary standing water waves*, Memoirs of the Amer. Math. Society. Vol. 263, Number 1273, 2019.
- [12] J. BOURGAIN, *Quasi-periodic solutions of Hamiltonian perturbations of 2D linear Schrödinger equations*, Ann. of Math. 148, 363–439 (1998).
- [13] WALTER CRAIG AND C. EUGENE WAYNE. *Newton’s method and periodic solutions of nonlinear wave equations*. Comm. Pure Appl. Math., 46 (11): 1409–1498, 1993.
- [14] R. CALLEJA, A. CELLETTI, L. CORSI, R. DE LA LLAVE. *Response solutions for quasi-periodically forced, dissipative wave equations*. SIAM J. Math Anal. 49 (2017), no.4, 3161–3207.

- [15] L. CORSI, R. MONTALTO. *Quasi-periodic solutions for the forced Kirchhoff equation on \mathbb{T}^d* . Nonlinearity 31,5075-5109 (2018). <https://doi.org/10.1088/1361-6544/aad6fe>
- [16] H.L. ELIASSEN AND S.B. KUKSIN. *On reducibility of Schrödinger equations with quasiperiodic in time potentials*. Comm. Math. Phys., 286(1):125–135, 2009.
- [17] R. FEOLA, M. PROCESI, *Quasi-periodic solutions for fully nonlinear forced reversible Schrödinger equations*, J. Diff. Eq., 259, no. 7, 3389–3447, 2015.
- [18] R. FEOLA, F. GIULIANI, R. MONTALTO, M. PROCESI. *Reducibility of first order linear operators on tori via Moser’s theorem*. Journal of Functional Analysis, 276 (2019) 932970.
- [19] G. P. GALDI. *Existence and uniqueness of time-periodic solutions to the Navier- Stokes equations in the whole plane*. Discrete Contin. Dyn. Syst., Ser. S, 6(5):1237-1257, 2013.
- [20] P. G. GALDI. *On Time-Periodic Flow of a Viscous Liquid past a Moving Cylinder*. Arch. Ration. Mech. Anal., 210(2):451-498, 2013.
- [21] G. P. GALDI AND A. L. SILVESTRE. *Existence of time-periodic solutions to the Navier-Stokes equations around a moving body*. Pac. J. Math., 223(2):251-267, 2006.
- [22] G. P. GALDI AND M. KYED. *Time-periodic solutions to the navier-Stokes equations in the three-dimensional whole-space with a drift term: Asymptotic profile at spatial infinity*. To appear in Contemporary Mathematics, 2016
- [23] G. P. GALDI, T. HISHIDA. *Attainability of Time-Periodic flow of a Viscous Liquid Past an Oscillating Body*. Preprint arXiv. 2001.07292 (2020)
- [24] J. GENG J, X. XU, J. YOU *An infinite dimensional KAM theorem and its application to the two dimensional cubic Schrödinger equation*. Adv. Math. 226, 5361-5402, 2011
- [25] G. IOOSS, P. I. PLOTNIKOV, AND J. F. TOLAND. *Standing waves on an infinitely deep perfect fluid under gravity*. Arch. Ration. Mech. Anal., 177(3):367–478, 2005.
- [26] J. LIU, H. CONG, X. YUAN. *Stability of KAM tori for nonlinear Schrödinger equation*. Mem. Amer. Math. Society, Vol. 239, No. 1134, 2016.
- [27] P. MAREMONTI. *Existence and stability of time-periodic solutions to the Navier-Stokes equations in the whole space*. Nonlinearity, 4(2):503-529, 1991. 2
- [28] P. MAREMONTI, M. PADULA. *Existence, uniqueness, and attainability of periodic solutions of the Navier-Stokes equations in exterior domains*. J. Math. Sci. (New York), 93 (1999), 719–746.
- [29] R. MONTALTO. *A reducibility result for a class of linear wave equations on \mathbb{T}^d* . Int. Math. Res. Notices, Vol. 2019, No. 6, pp. 1788–1862 doi: 10.1093/imrn/rnx167.

- [30] J. L. LIONS. *Quelques methodes de resolution des problemes aux limites non lineares*. Gounod and Gautier-Villars (1969) Paris.
- [31] J. LIU, X. YUAN, *A KAM Theorem for Hamiltonian Partial Differential Equations with Unbounded Perturbations*, Comm. Math. Phys, 307 (3) , 629–673, 2011.
- [32] M. KYED. *Existence and regularity of time-periodic solutions to the three-dimensional Navier-Stokes equations*. Nonlinearity, 27(12), 2909-2935, 2014.
- [33] S. B. KUKSIN. *Hamiltonian perturbations of infinite-dimensional linear systems with an imaginary spectrum*. Funct. Anal. Appl. 21:192–205, 1987.
- [34] S. KUKSIN, *A KAM theorem for equations of the Korteweg-de Vries type*, Rev. Math. Phys., 10, 3, 1–64, 1998.
- [35] C. PROCESI, M. PROCESI *A KAM algorithm for the completely resonant nonlinear Schrödinger equation*. Advances in Mathematics, volume 272, 399-470, 2015.
- [36] G. PRODI. *Qualche risultato riguardo alle equazioni di Navier-Stokes nel caso bidimensionale*. Rend. Sem. Mat. Univ. Padova, 30:1-15, 1960.
- [37] G. PROUSE. *Soluzioni periodiche dellequazione di Navier-Stokes*. Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8), 35:443-447, 1963.
- [38] G. PROUSE. *Soluzioni quasi-periodiche dellequazione differenziale di Navier-Stokes in due dimensioni*. Rend. Sem. Mat. Univ. Padova, 33:186-212, 1963
- [39] R. SALVI. *On the Existence of Periodic Weak Solutions on the Navier-Stokes Equations in Exterior Regions with Periodically Moving Boundaries*. In: Sequeira A. (eds) NavierStokes Equations and Related Nonlinear Problems. Springer, Boston, MA (1995).
- [40] J. SERRIN. *A note on the existence of periodic solutions of the Navier-Stokes equations*. Arch. Ration. Mech. Anal. 3, 120-122 (1959)
- [41] V.I. YUDOVICH. *Periodic motions of a viscous incompressible fluid*. Sov. Math. Dokl. 1, 168-172 (1960).
- [42] C. EUGENE WAYNE. *Periodic and quasi-periodic solutions of nonlinear wave equations via KAM theory*. Comm. Math. Phys., 127(3):479-528, 1990.

Riccardo Montalto

Dipartimento di Matematica “Federigo Enriques”
 Università degli Studi di Milano
 Via Cesare Saladini 50
 20133 Milano, Italy
 riccardo.montalto@unimi.it