

# Inflation in $f(R, h)$ theory formulated in the Palatini formalism

Nayan Das,<sup>a</sup> Sukanta Panda<sup>a</sup>

<sup>a</sup>Department of Physics, Indian Institute of Science Education and Research, Bhopal 462066, India

E-mail: [nayandas@iiserb.ac.in](mailto:nayandas@iiserb.ac.in), [sukanta@iiserb.ac.in](mailto:sukanta@iiserb.ac.in)

**Abstract.** A new model for inflation using modified gravity in the Palatini formalism is constructed. Here non-minimal coupling of scalar field  $h$  with the curvature  $R$  as a general function  $f(R, h)$  is considered. Explicit inflation models for some choices of  $f(R, h)$  are developed. By writing an equivalent scalar-tensor action for this model and going over to Einstein frame, slow roll parameters are constructed. There exists a large parameter space which satisfies values of  $n_s$  and limits on  $r$  compatible with planck 2018 data.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The Model</b>	<b>2</b>
<b>3</b>	<b>Slow roll parameters and Results</b>	<b>3</b>
<b>4</b>	<b>Conclusion</b>	<b>8</b>

---

## 1 Introduction

There are mainly two formulations in General Relativity (GR) popularly known as Palatini and Metric formalisms. Palatini formalism or first order formalism treats space time connections as independent variable[1–9], whereas in Metric formalism, these connections are not independent but derived from metric itself. But in GR these two approaches produce same Einstein equation. Hence dynamics are equivalent in both formalisms. This is not true for modified gravity models and models where fields are nonminimally coupled to gravity. In these cases both formalisms represent different physical situations [1–3].

Inflation [10–16] was first developed in the early 1980s to solve problems of standard Big Bang theory like horizon problem, fine-tuning problem etc. Quantum fluctuations also started during period of inflation which led to the cosmic microwave background(CMB) anisotropy and provided the seed for the formation of large scale structure of the universe. A more popular model for inflation is the Starobinsky model which is a pure gravity theory with an additional  $R^2$  term in the Einstein-Hilbert action. An equivalent scalar tensor theory of Starobinsky model has an additional scalar degree of freedom apart from two tensor degrees of freedom of Einstein-Hilbert action[17–19]. In Einstein frame, this theory is equivalent to usual scalar field model with a potential suitable for a valid inflation model which satisfies all the constraints from CMB data.

In a broad sense, the Starobinsky model falls under a general framework of  $f(R)$  gravity. The above analysis works best in metric formulation of gravity. However, in the Palatini formalism, no additional propagating degrees of freedom appears in  $f(R)$  gravity theory [3]. Because of this no inflation is possible in this scenario. Hence a scalar field needs to be added to this action to develop an Inflation model in Palatini formulation of gravity. In this line, first work appeared in [20] where nonminimal couplings of scalars are considered. Since then many works related to different inflationary potentials, preheating, reheating, postinflationary phases, dark matter has been done with many variants of action including a  $R^2$  term [21–63]. For an introduction to Palatini inflation models refer to [64] and references therein.

In this work, we formulate a palatini inflation model in  $f(R, h)$  theory. In metric formalism inflation model in  $f(R, h)$  gravity has been considered earlier in literature[65–67]. Our action is of the form similar to form obtained in [65]. But our approach here is to develop a model of inflation in palatini formalism. We show that such an action provide a favorable inflationary scenario satisfying Planck 2018 data[68].

This paper is organised as follows. In the next section, we introduce the action of our model. Then we express it in terms of equivalent scalar tensor action in Einstein frame. In

section 3, slow roll parameters are defined. Values of  $n_s$  and  $r$  are also calculated numerically and the results are presented. In section 4, we summarize our results.

## 2 The Model

We start with a general action of the form-

$$S = \int d^4x \sqrt{-g} \left[ \frac{1}{2} f(R, h) - \frac{1}{2} g^{\mu\nu} \partial_\mu h \partial_\nu h - V(h) \right] \quad (2.1)$$

where  $g_{\alpha\beta}$  is the metric,  $g$  is its determinant,  $f(R, h) = G(h)(R + \alpha R^2)$ ,  $R$  is the Ricci scalar defined as  $R = g^{\alpha\beta} R^\gamma_{\alpha\gamma\beta}(\Gamma, \partial\Gamma)$  in Palatini formalism and  $\alpha$  is a constant. Here we have chosen Planck mass to be unity. An equivalent action in terms of an auxiliary field  $\phi$  can be written as-

$$S = \int d^4x \sqrt{-g} \left[ \frac{1}{2} f(\phi, h) + \frac{1}{2} f'(\phi, h)(R - \phi) - \frac{1}{2} g^{\mu\nu} \partial_\mu h \partial_\nu h - V(h) \right] \quad (2.2)$$

where  $f'(\phi, h) = \frac{\partial f(\phi, h)}{\partial \phi}$ . Varying  $S$  with respect to  $\phi$  in equation (2.2), we get  $\phi = R$  if  $\frac{\partial^2 f(\phi, h)}{\partial \phi^2} \neq 0$  and using this result in equation (2.2), we recover equation (2.1). Rearranging equation (2.2), action becomes

$$S = \int d^4x \sqrt{-g} \left[ \frac{1}{2} f'(\phi, h) R - W(\phi, h) - \frac{1}{2} g^{\mu\nu} \partial_\mu h \partial_\nu h - V(h) \right] \quad (2.3)$$

where  $W(\phi, h) = \frac{1}{2} \phi f'(\phi, h) - \frac{1}{2} f(\phi, h)$ . By making a conformal transformation-

$$g_{\mu\nu} \rightarrow f'(\phi, h) g_{\mu\nu}, \quad (2.4)$$

action is obtained in Einstein frame as-

$$S = \int d^4x \sqrt{-g} \left[ \frac{1}{2} R - \frac{1}{2} \frac{\partial_\mu h \partial^\mu h}{f'(\phi, h)} - \frac{W(\phi, h) + V(h)}{f'(\phi, h)^2} \right] \quad (2.5)$$

Let us define a new potential  $\hat{V}(\phi, h)$  as-

$$\hat{V}(\phi, h) \equiv \frac{W(\phi, h) + V(h)}{f'(\phi, h)^2}. \quad (2.6)$$

Now for our choice of  $f(R, h) = G(h)(R + \alpha R^2)$ [65], the new potential can be written as-

$$\hat{V}(\phi, h) = \frac{1}{f'(\phi, h)^2} \left[ \frac{1}{8\alpha G(h)} [f'(\phi, h) - G(h)]^2 + V(h) \right]. \quad (2.7)$$

Varying (2.5) with respect to  $\phi$ , we get the constraint equation-

$$f'(\phi, h) = \frac{8\alpha V(h) + G(h)}{1 - 2\alpha \partial_\mu h \partial^\mu h}. \quad (2.8)$$

Inserting (2.8) to equation (2.7), we can eliminate  $\phi$ . Then again inserting (2.7) to (2.5) and rearranging and simplifying, we get-

$$S = \int d^4x \sqrt{-g} \left[ \frac{1}{2}R - \frac{1}{2(8\alpha V + G)} \partial^\mu h \partial_\mu h + \frac{\alpha}{2(8\alpha V + G)} (\partial^\mu h \partial_\mu h)^2 - \frac{V}{G(8\alpha V + G)} \right] \quad (2.9)$$

Now the last term can be defined as effective potential in the Einstein frame-

$$U \equiv \frac{V}{G(8\alpha V + G)} \quad (2.10)$$

In order to bring kinetic part of scalar field into that of canonical form, we introduce a new field  $\chi$  as-

$$\frac{dh}{d\chi} = \pm \sqrt{(8\alpha V + G)} \quad (2.11)$$

In terms of  $\chi$ , equation (2.9) becomes-

$$S = \int d^4x \sqrt{-g} \left[ \frac{1}{2}R - \frac{1}{2} \partial^\mu \chi \partial_\mu \chi + \frac{\alpha}{2} (8\alpha V + G) (\partial^\mu \chi \partial_\mu \chi)^2 - U \right] \quad (2.12)$$

Now we are ready to build up our inflation model, with this action. Next we define the slow roll parameters to estimate the observables which will decide the fate of our model. Here we have neglected contribution due to third term of the action to the inflation phase. This is a valid assumption so far as the slow roll inflation is concerned (see [64] and references therein). In this case then it's model of inflation driven by a scalar field with potential  $U$ <sup>1</sup>.

### 3 Slow roll parameters and Results

Before analysing dynamics of our model we define slow roll parameters (mainly  $\epsilon$  and  $\eta$ ) which are helpful to decide whether a model can describe inflation or not. They can be written as-

$$\begin{aligned} \epsilon &= \frac{1}{2} \left( \frac{dU}{d\chi} \right)^2 \\ \eta &= \frac{d^2U}{d\chi^2} \end{aligned} \quad (3.1)$$

The slow roll parameters  $\epsilon$  and  $\eta$  must be  $\ll 1$  during inflation phase of expansion. For the potential in equation (2.10), slow roll parameters become

---

<sup>1</sup>Background evolution of  $\chi$  field for a flat FRW metric with scale factor  $a$  and hubble parameter  $H = \frac{\dot{a}}{a}$  is-

$$a^2 \ddot{\chi} + 2a^3 H \dot{\chi} + 12aHf(\chi) (\dot{\chi})^2 + 3 \frac{df(\chi)}{d\chi} (\dot{\chi})^4 + a^4 U' = 0 \quad (2.13)$$

where  $f(\chi) = \frac{\alpha}{2}(8\alpha V + G)$  and  $\dot{\phantom{x}}$  and  $'$  represent derivatives with respect to conformal time and  $\chi$  respectively. Under slow-roll approximation, terms proportional to  $\dot{\chi}^2$  and  $\dot{\chi}^4$  can be neglected and equation (2.13) reduces to usual FRW equations for minimally coupled field.

$$\begin{aligned}\epsilon &= \frac{G\bar{\epsilon}}{(8\alpha V + G)} \left[ 1 - \left( \frac{8\alpha \frac{V}{G^2} G'}{\frac{V'}{V} - \frac{2G'}{G}} \right) \right]^2 \\ \eta &= \bar{\eta} \frac{G}{8\alpha V + G} - \left[ (-256\alpha^2 V^3 G'^2 + 8\alpha G V^2 (-11G''^2 + 8\alpha G' V' + 16\alpha V G'')) \right. \\ &\quad \left. + 2G^2 V (8\alpha G' V' + 24\alpha V G'') + G^3 (24\alpha V'^2 - 16\alpha V V'') \right] / (2G^2 V (8\alpha V + G))\end{aligned}\quad (3.2)$$

where  $\bar{\epsilon}$  and  $\bar{\eta}$  are slow roll parameters for  $\alpha = 0$  and  $'$  represents derivative with respect to  $h$ . Number of e-folds,  $N$  can be written as ( for unit planck mass)-

$$N = - \int_{h_i}^{h_f} dh \frac{1}{\sqrt{2\epsilon(h)(8\alpha V + G)}} \quad (3.3)$$

Inflation ends when  $\epsilon \simeq 1$  and this in turn determines the value of  $h_f$ . Putting the value of  $h_f$  and  $N$  (taken in between 60 and 70) in (3.3), value of  $h_i$  can be obtained. Putting this value to equations (3.1) and (3.2), value of  $\epsilon$  and  $\eta$  can be obtained respectively. Scalar spectral index and tensor-to-scalar ratio can then be calculated from the following equations-

$$\begin{aligned}r &= 16\epsilon \\ n_s &= 1 + 2\eta - 6\epsilon\end{aligned}\quad (3.4)$$

Now we estimate  $r$  and  $n_s$  for various cases. Here we restrict our analysis only for quadratic and quartic potentials. We choose two values of  $G = \gamma h^2$  and  $G = 1 + \gamma h^2$  and  $\gamma$  is a dimensionless constant.

**Case 1 :**  $V = \beta h^2$  and  $G = \gamma h^2$


For this case

$$\begin{aligned}\epsilon &= 2(8\alpha\beta + \gamma) \\ \eta &= 4(8\alpha\beta + \gamma)\end{aligned}\quad (3.5)$$


As neither  $\epsilon$  nor  $\eta$  depend on  $h$ , so it will lead to constant  $r$  and  $n_s$  for a particular choice of  $\alpha$ ,  $\beta$  and  $\gamma$ . In this case then it would be difficult to end the inflation and to decide the value of  $h_f$ . Therefore we will not discuss this case further here.

**Case 2 :**  $V = \beta h^4$  and  $G = \gamma h^2$


In this case,  $\epsilon$  and  $\eta$  depend only on  $h$  and  $\gamma$  for a particular value of  $N$ . The parameters  $\alpha$  and  $\beta$  fixes the value of the field  $\chi$  or  $h$ . In fig. 1, potential  $U$  vs  $\chi$  is plotted. In fig. 2 the relation between  $h$  vs  $\chi$  is plotted. The value of  $\chi$  changes discontinuously as  $h$  changes from negative to positive value. Here positive sign is taken in equation (2.11). Instead if negative sign is taken, all values of  $\chi$ 's will be negative. However, discontinuity of  $\chi$  is there for both positive and negative sign. For fig. 1 and fig. 2, the value of parameters  $\alpha = 0.5$ ,  $\beta = \frac{1}{4} \times 10^{-4}$  and  $\gamma = 0.02$  is chosen. For this particular set, the numerically estimated value of  $n_s$  and  $r$  are 0.967 and 0.023 respectively (we take  $N = 60$  here). Also the result for spectral index and tensor to scalar ratio is shown in fig. 3. Here  $\gamma$  is changing and  $\alpha = 5$ ,  $\beta = \frac{1}{4} \times 10^{-4}$  and negative sign is considered in equation (2.11). It clearly shows that there can be large parameter space available which can satisfy Planck constraints.



**Figure 1.** Potential as a function of canonical scalar field



**Figure 2.** Plot of  $\chi(h)$



**Figure 3.** Plot of  $r$  vs  $n_s$  for changing  $\gamma$

**Case 3 :**  $V = \beta h^4$  and  $G = 1 + \gamma h^2$

For case 3 and case 4,  $G(h)$  is taken as  $G = 1 + \gamma h^2$ . This makes the Lagrangian in (2.1) equivalent to the form  $\frac{1}{2}(R + \alpha R^2) + \frac{1}{2}\gamma h^2(R + \alpha R)$  plus usual scalar field term. Similar to case 2, the graphs of  $U$  vs  $\chi$  and  $h$  vs  $\chi$  are shown in fig 4 and fig. 5 respectively. Unlike to the previous case, here  $\chi(h)$  is continuous for all range of  $h$ . For fig 4 and fig 5, the value of parameters are  $\alpha = 10^{-4}$ ,  $\beta = 10^{-4}$  and  $\gamma = 0.03$ . For this particular value of parameters, we obtain  $n_s = 0.966$  and  $r = 0.017$ . The potential in equation (2.10) takes the form-


$$U = \frac{\beta h^4}{(1 + \gamma h^2)(8\alpha\beta h^4 + 1 + \gamma h^2)} \quad (3.6)$$

Dividing both numerator and denominator by  $h^6$ , we get -


$$U = \frac{\beta \frac{1}{h^2}}{(\frac{1}{h^2} + \gamma)(8\alpha\beta + \frac{1}{h^4} + \gamma \frac{1}{h^2})} \quad (3.7)$$

We can clearly observe that  $U \rightarrow 0$  as  $h \rightarrow \infty$ . Hence this potential will not give rise to plateau unlike the results for Lagrangian where  $h$  is coupled only with  $R$  [38–40, 45, 49, 69]. The results for  $n_s$  and  $r$  are shown in fig. 6, 7 and 8. In fig. 6,  $\beta = 10^{-4}$ ,  $\gamma = 10^{-2}$  and  $\alpha$  is


changing. In fig. 7,  $\alpha = 10^{-4}$ ,  $\gamma = 10^{-2}$  and  $\beta$  is changing. And in fig. 8,  $\alpha = 10^{-3}$ ,  $\beta = 10^{-4}$  and  $\gamma$  is changing. Here we consider positive sign in equation (2.11). Note that both 6 and 7 look same. This happens because in the expression of  $r$  and  $n_s$ ,  $\alpha$  and  $\beta$  always appear in pairs i.e. in the form of  $(\alpha\beta)$  or  $(\alpha\beta)^2$ . Similar to case 2, here also a large parameter space available which satisfies Planck constraints on  $r$  and  $n_s$ . It is observed from 8 that we can tune parameters of our model to obtain a very small value for  $r$ .




**Figure 4.** Potential as a function of canonical scalar field




**Figure 5.** Plot of  $\chi(h)$



**Figure 6.** Plot of  $r$  vs  $n_s$  for changing  $\alpha$



**Figure 7.** Plot of  $r$  vs  $n_s$  for changing  $\beta$




**Figure 8.** Plot of  $r$  vs  $n_s$  for changing  $\gamma$

**Case 4 :**  $V = \beta h^2$  and  $G = 1 + \gamma h^2$


Analogous to case 3, potential graph and  $h$  vs  $\chi$  are shown in fig 9 and fig 10 for  $\alpha = 0.1$ ,  $\beta = 0.01$  and  $\gamma = 10^{-4}$  which give  $n_s = 0.965$  and  $r = 0.04$ . Here, potential in equation (2.10) becomes -

$$U = \frac{\beta h^2}{(1 + \gamma h^2)(8\alpha\beta h^2 + 1 + \gamma h^2)} \quad (3.8)$$

As  $h \rightarrow \infty$ ,  $U \rightarrow 0$ . So again no plateau in the potential. The  $n_s$  and  $r$  graphs are plotted in fig. 11, 12 and 13. In fig. 11  $\beta = 0.01$ ,  $\gamma = 0.001$  and  $\alpha$  is changing. In fig. 12  $\alpha = 10^{-2}$ ,  $\gamma = 10^{-3}$  and  $\beta$  is varying. In fig. 13  $\alpha = 10^{-3}$ ,  $\beta = 0.01$  and  $\gamma$  is varying. Here we consider positive sign in equation (2.11). In this case as well, we find a large parameter space available which is compatible with Planck 2018 data.




**Figure 9.** Potential as a function of canonical scalar field




**Figure 10.** Plot of  $\chi(h)$






**Figure 11.** Plot of  $r$  vs  $n_s$  for changing  $\alpha$



**Figure 12.** Plot of  $r$  vs  $n_s$  for changing  $\beta$



**Figure 13.** Plot of  $r$  vs  $n_s$  for changing  $\gamma$

## 4 Conclusion

Inflation models are constructed in a modified gravity theory in Palatini formalism. Here our modified action contains a general non-minimal coupling of scalar field with  $R$  and  $R^2$  term separately. Unlike metric formalism, Palatini formalism does not introduce a new scalar degree of freedom in our theory. It turns out that  $R^2$  term in the action can be translated to a higher order kinetic term and a new potential term for scalar field in an equivalent scalar tensor set-up in Einstein frame. However, in slow roll inflation setting we can safely neglect the effect of higher order kinetic term in the action. The modified potential is responsible for the inflation. In this equivalent scalar tensor theory, we calculate scalar spectral index( $n_s$ ) and tensor to scalar ratio( $r$ ) for four different cases depending on the form of nonminimal coupling and potential of scalar field. We have varied three unknown parameters of our theory to extract the information about  $n_s$  and  $r$ . It is observed that in most of the cases we have a large parameter space which can match with the results of Planck 2018 constraints on  $n_s$  and  $r$ . Palatini formalism in modified gravity theories may provide a large class of models

for Inflation satisfying CMB data. It would be interesting to look at reheating in such a scenario. Also a more general action with nonminimal coupling of scalar field can be studied to explore the cosmology of very early universe. A dark energy picture of these models is worth investing in future.

## Acknowledgments

This work was partially funded by DST (Govt. of India), Grant No. SERB/PHY/2017041.

## References

- [1] T. P. Sotiriou, *f(R) gravity and scalar-tensor theory*, *Class. Quant. Grav.* **23** (2006) 5117–5128, [[gr-qc/0604028](#)].
- [2] T. P. Sotiriou and S. Liberati, *Metric-affine f(R) theories of gravity*, *Annals Phys.* **322** (2007) 935–966, [[gr-qc/0604006](#)].
- [3] T. P. Sotiriou and V. Faraoni, *f(R) Theories Of Gravity*, *Rev. Mod. Phys.* **82** (2010) 451–497, [[0805.1726](#)].
- [4] M. Borunda, B. Janssen and M. Bastero-Gil, *Palatini versus metric formulation in higher curvature gravity*, *JCAP* **11** (2008) 008, [[0804.4440](#)].
- [5] A. De Felice and S. Tsujikawa, *f(R) theories*, *Living Rev. Rel.* **13** (2010) 3, [[1002.4928](#)].
- [6] G. J. Olmo, *Palatini Approach to Modified Gravity: f(R) Theories and Beyond*, *Int. J. Mod. Phys. D* **20** (2011) 413–462, [[1101.3864](#)].
- [7] S. Capozziello and M. De Laurentis, *Extended Theories of Gravity*, *Phys. Rept.* **509** (2011) 167–321, [[1108.6266](#)].
- [8] T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, *Modified Gravity and Cosmology*, *Phys. Rept.* **513** (2012) 1–189, [[1106.2476](#)].
- [9] S. Nojiri, S. Odintsov and V. Oikonomou, *Modified Gravity Theories on a Nutshell: Inflation, Bounce and Late-time Evolution*, *Phys. Rept.* **692** (2017) 1–104, [[1705.11098](#)].
- [10] A. H. Guth, *Inflationary universe: A possible solution to the horizon and flatness problems*, *Phys. Rev. D* **23** (Jan, 1981) 347–356.
- [11] A. Starobinsky, *A new type of isotropic cosmological models without singularity*, *Physics Letters B* **91** (1980) 99 – 102.
- [12] A. Linde, *A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems*, *Physics Letters B* **108** (1982) 389 – 393.
- [13] A. Linde, *Chaotic inflation*, *Physics Letters B* **129** (1983) 177 – 181.
- [14] D. H. Lyth and A. Riotto, *Particle physics models of inflation and the cosmological density perturbation*, *Phys. Rept.* **314** (1999) 1–146, [[hep-ph/9807278](#)].
- [15] A. Riotto, *Inflation and the theory of cosmological perturbations*, *ICTP Lect. Notes Ser.* **14** (2003) 317–413, [[hep-ph/0210162](#)].
- [16] D. Baumann, *Inflation*, in *Physics of the large and the small, TASI 09, proceedings of the Theoretical Advanced Study Institute in Elementary Particle Physics, Boulder, Colorado, USA, 1-26 June 2009*, pp. 523–686, 2011, [0907.5424](#), DOI.
- [17] A. A. Starobinsky, *A New Type of Isotropic Cosmological Models Without Singularity*, *Adv. Ser. Astrophys. Cosmol.* **3** (1987) 130–133.

- [18] V. F. Mukhanov and G. V. Chibisov, *Quantum Fluctuations and a Nonsingular Universe*, *JETP Lett.* **33** (1981) 532–535.
- [19] A. A. Starobinsky, *The Perturbation Spectrum Evolving from a Nonsingular Initially De-Sitter Cosmology and the Microwave Background Anisotropy*, *Sov. Astron. Lett.* **9** (1983) 302.
- [20] F. Bauer and D. A. Demir, *Inflation with Non-Minimal Coupling: Metric versus Palatini Formulations*, *Phys. Lett. B* **665** (2008) 222–226, [[0803.2664](#)].
- [21] T. Koivisto and H. Kurki-Suonio, *Cosmological perturbations in the palatini formulation of modified gravity*, *Class. Quant. Grav.* **23** (2006) 2355–2369, [[astro-ph/0509422](#)].
- [22] N. Tamanini and C. R. Contaldi, *Inflationary Perturbations in Palatini Generalised Gravity*, *Phys. Rev. D* **83** (2011) 044018, [[1010.0689](#)].
- [23] F. Bauer and D. A. Demir, *Higgs-Palatini Inflation and Unitarity*, *Phys. Lett. B* **698** (2011) 425–429, [[1012.2900](#)].
- [24] K. Enqvist, T. Koivisto and G. Rigopoulos, *Non-metric chaotic inflation*, *JCAP* **05** (2012) 023, [[1107.3739](#)].
- [25] A. Borowiec, M. Kamionka, A. Kurek and M. Szydlowski, *Cosmic acceleration from modified gravity with Palatini formalism*, *JCAP* **02** (2012) 027, [[1109.3420](#)].
- [26] A. Stachowski, M. Szydlowski and A. Borowiec, *Starobinsky cosmological model in Palatini formalism*, *Eur. Phys. J. C* **77** (2017) 406, [[1608.03196](#)].
- [27] C. Fu, P. Wu and H. Yu, *Inflationary dynamics and preheating of the nonminimally coupled inflaton field in the metric and Palatini formalisms*, *Phys. Rev. D* **96** (2017) 103542, [[1801.04089](#)].
- [28] S. Rasanen and P. Wahlman, *Higgs inflation with loop corrections in the Palatini formulation*, *JCAP* **11** (2017) 047, [[1709.07853](#)].
- [29] T. Tenkanen, *Resurrecting Quadratic Inflation with a non-minimal coupling to gravity*, *JCAP* **12** (2017) 001, [[1710.02758](#)].
- [30] A. Racioppi, *Coleman-Weinberg linear inflation: metric vs. Palatini formulation*, *JCAP* **12** (2017) 041, [[1710.04853](#)].
- [31] T. Markkanen, T. Tenkanen, V. Vaskonen and H. Veermäe, *Quantum corrections to quartic inflation with a non-minimal coupling: metric vs. Palatini*, *JCAP* **03** (2018) 029, [[1712.04874](#)].
- [32] L. Järv, A. Racioppi and T. Tenkanen, *Palatini side of inflationary attractors*, *Phys. Rev. D* **97** (2018) 083513, [[1712.08471](#)].
- [33] S. Rasanen, *Higgs inflation in the Palatini formulation with kinetic terms for the metric*, [1811.09514](#).
- [34] A. Racioppi, *New universal attractor in nonminimally coupled gravity: Linear inflation*, *Phys. Rev. D* **97** (2018) 123514, [[1801.08810](#)].
- [35] P. Carrilho, D. Mulryne, J. Ronayne and T. Tenkanen, *Attractor Behaviour in Multifield Inflation*, *JCAP* **06** (2018) 032, [[1804.10489](#)].
- [36] V.-M. Enckell, K. Enqvist, S. Rasanen and E. Tomberg, *Higgs inflation at the hilltop*, *JCAP* **06** (2018) 005, [[1802.09299](#)].
- [37] F. Bombacigno and G. Montani, *Big bounce cosmology for Palatini  $R^2$  gravity with a Nieh–Yan term*, *Eur. Phys. J. C* **79** (2019) 405, [[1809.07563](#)].
- [38] V.-M. Enckell, K. Enqvist, S. Rasanen and L.-P. Wahlman, *Inflation with  $R^2$  term in the Palatini formalism*, *JCAP* **1902** (2019) 022, [[1810.05536](#)].
- [39] I. Antoniadis, A. Karam, A. Lykkas and K. Tamvakis, *Palatini inflation in models with an  $R^2$  term*, *JCAP* **11** (2018) 028, [[1810.10418](#)].

- [40] I. Antoniadis, A. Karam, A. Lykkas, T. Pappas and K. Tamvakis, *Rescuing Quartic and Natural Inflation in the Palatini Formalism*, *JCAP* **03** (2019) 005, [[1812.00847](#)].
- [41] S. Rasanen and E. Tomberg, *Planck scale black hole dark matter from Higgs inflation*, *JCAP* **01** (2019) 038, [[1810.12608](#)].
- [42] J. P. B. Almeida, N. Bernal, J. Rubio and T. Tenkanen, *Hidden Inflaton Dark Matter*, *JCAP* **03** (2019) 012, [[1811.09640](#)].
- [43] T. Takahashi and T. Tenkanen, *Towards distinguishing variants of non-minimal inflation*, *JCAP* **04** (2019) 035, [[1812.08492](#)].
- [44] K. Kannike, A. Kubarski, L. Marzola and A. Racioppi, *A minimal model of inflation and dark radiation*, *Phys. Lett. B* **792** (2019) 74–80, [[1810.12689](#)].
- [45] T. Tenkanen, *Minimal Higgs inflation with an  $R^2$  term in Palatini gravity*, *Phys. Rev. D* **99** (2019) 063528, [[1901.01794](#)].
- [46] K. Shimada, K. Aoki and K.-i. Maeda, *Metric-affine Gravity and Inflation*, *Phys. Rev. D* **99** (2019) 104020, [[1812.03420](#)].
- [47] J. Wu, G. Li, T. Harko and S.-D. Liang, *Palatini formulation of  $f(R, T)$  gravity theory, and its cosmological implications*, *Eur. Phys. J. C* **78** (2018) 430, [[1805.07419](#)].
- [48] A. Kozak and A. Borowiec, *Palatini frames in scalar–tensor theories of gravity*, *Eur. Phys. J. C* **79** (2019) 335, [[1808.05598](#)].
- [49] R. Jinno, K. Kaneta, K.-y. Oda and S. C. Park, *Hillclimbing inflation in metric and Palatini formulations*, *Phys. Lett. B* **791** (2019) 396–402, [[1812.11077](#)].
- [50] A. Edery and Y. Nakayama, *Palatini formulation of pure  $R^2$  gravity yields Einstein gravity with no massless scalar*, *Phys. Rev. D* **99** (2019) 124018, [[1902.07876](#)].
- [51] J. Rubio and E. S. Tomberg, *Preheating in Palatini Higgs inflation*, *JCAP* **04** (2019) 021, [[1902.10148](#)].
- [52] R. Jinno, M. Kubota, K.-y. Oda and S. C. Park, *Higgs inflation in metric and Palatini formalisms: Required suppression of higher dimensional operators*, *JCAP* **03** (2020) 063, [[1904.05699](#)].
- [53] M. Giovannini, *Post-inflationary phases stiffer than radiation and Palatini formulation*, *Class. Quant. Grav.* **36** (2019) 235017, [[1905.06182](#)].
- [54] T. Tenkanen and L. Visinelli, *Axion dark matter from Higgs inflation with an intermediate  $H_*$* , *JCAP* **08** (2019) 033, [[1906.11837](#)].
- [55] N. Bostan, *Quadratic, Higgs and hilltop potentials in the Palatini gravity*, [1908.09674](#).
- [56] T. Tenkanen, *Trans-Planckian censorship, inflation, and dark matter*, *Phys. Rev. D* **101** (2020) 063517, [[1910.00521](#)].
- [57] I. D. Gialamas and A. Lahanas, *Reheating in  $R^2$  Palatini inflationary models*, *Phys. Rev. D* **101** (2020) 084007, [[1911.11513](#)].
- [58] M. Shaposhnikov, A. Shkerin and S. Zell, *Quantum Effects in Palatini Higgs Inflation*, [2002.07105](#).
- [59] A. Lloyd-Stubbs and J. McDonald, *Sub-Planckian  $\phi^2$  Inflation in the Palatini Formulation of Gravity with an  $R^2$  term*, [2002.08324](#).
- [60] P. M. Sá, *Unified description of dark energy and dark matter within the generalized hybrid metric-Palatini theory of gravity*, [2002.09446](#).
- [61] I. Antoniadis, A. Lykkas and K. Tamvakis, *Constant-roll in the Palatini- $R^2$  models*, *JCAP* **04** (2020) 033, [[2002.12681](#)].

- [62] D. Ghilencea, *Palatini quadratic gravity: spontaneous breaking of gauged scale symmetry and inflation*, [2003.08516](#).
- [63] T. Takahashi, T. Tenkanen and S. Yokoyama, *Violation of slow-roll in non-minimal inflation*, [2003.10203](#).
- [64] T. Tenkanen, *Tracing the high energy theory of gravity: an introduction to Palatini inflation*, *Gen. Rel. Grav.* **52** (2020) 33, [[2001.10135](#)].
- [65] T. Rador,  *$f(R)$  Gravities a la Brans-Dicke*, *Phys. Lett. B* **652** (2007) 228–232, [[hep-th/0702081](#)].
- [66] A. de la Cruz-Dombriz, E. Elizalde, S. D. Odintsov and D. Sáez-Gómez, *Spotting deviations from  $R^2$  inflation*, *JCAP* **05** (2016) 060, [[1603.05537](#)].
- [67] J. Mathew, J. P. Johnson and S. Shankaranarayanan, *Inflation with  $f(R, \phi)$  in Jordan frame*, *Gen. Rel. Grav.* **50** (2018) 90, [[1705.07945](#)].
- [68] PLANCK collaboration, N. Aghanim et al., *Planck 2018 results. VI. Cosmological parameters*, [1807.06209](#).
- [69] T. Tenkanen and E. Tomberg, *Initial conditions for plateau inflation: a case study*, *JCAP* **04** (2020) 050, [[2002.02420](#)].