

PREPARED FOR SUBMISSION TO JHEP

On-Shell Perspectives on the Massless Limit of Massive Supergravity

Daniel J Burger^{a,b}, Nathan Moynihan^{a,b} & Jeff Murugan^a

^a*The Laboratory for Quantum Gravity & Strings (QGASLab)*

^b*High Energy Physics, Cosmology & Astrophysics Theory (HEPCAT) group ,
Department of Mathematics,
University Of Cape Town,
Private Bag, Rondebosch, 7701, South Africa
E-mail: burgerjdaan@gmail.com, nathantmoynihan@gmail.com,
jeff.murugan@uct.ac.za*

ABSTRACT: Massive gravity exhibits a famous discontinuity in its 2-point linearized amplitude for t-channel scattering of gravitational sources, in the $m \rightarrow 0$ limit. In essence, the source of this vDVZ discontinuity is in the failure of the zero-helicity mode to decouple in this limit. In [1], we showed how this result could be understood in the context of modern on-shell methods and, in particular, the BCFW construction. In this article, we provide a similar on-shell perspective to the equally interesting but lesser known discontinuity first discovered by Deser, Kay and Stelle in massive supergravity.

Contents

1	Introduction	1
2	The Supersymmetric vDVZ Discontinuity	3
3	$\mathcal{N} = 1$ Supersymmetric Discontinuity	5
3.1	Bold Notation and the Stückelberg Decomposition	5
3.2	Vector Multplet	6
3.3	Scalar Multplet	9
4	Supersymmetry Breaking and the Discontinuity	12
5	Discussion	14
A	Conventions and Notation	16
B	Some Miscellaneous Identities	17

1 Introduction

Endowing the graviton with a small but nonzero mass is an appealing idea for many reasons, the most significant of which is a credible explanation for the observed late-time acceleration of the Universe without the need to invoke exotic forms of matter and energy. However, since nothing in life is free, this approach is not without its own pathologies. Among these are a ghost mode and a, less scary but perhaps more famous, discontinuity in the 2-point function of the theory in the massless limit. The former was resolved in the recent seminal work of de Rham *et.al.* [2] while the latter boils down to a noncommutativity of limits. For this, there are two possibilities:

- turning off interactions does not necessarily commute with the massless limit. In other words, in order to resolve the vDVZ discontinuity, it is necessary to go beyond the linearised Fierz-Pauli action, leading to the famed Vainshtein screening mechanism of [3], or
- the massless limit does not commute with the limit of vanishing cosmological constant.

Either case will break Birkhoff’s theorem resulting in a vDVZ-like discontinuity. To clarify the situation surrounding this discontinuity, in [1] we attempted to re-phrase the discontinuity in the language of scattering amplitudes, largely because these modern on-shell methods are unencumbered by much of the baggage of the usual Lagrangian formulation of the problem. Recently, there have been several advances in our understanding of gravity by harnessing the on-shell scattering amplitudes paradigm [4–21], including approaches based on the double copy [22–67] (see [68] for a recent review), the *classical* double copy [14, 69–106] and those probing the on-shell structure of massive gravity [107, 108]. In [1], we utilised this modern approach to show how the vDVZ discontinuity manifests at the level of the scattering amplitudes but did not go quite far enough in disambiguating between the two sources above, mostly because computing massive on-shell amplitudes in gravity at higher (than linear) order was beyond the scope of that article. At the same time we were reminded that this massive to massless discontinuity of the spin-2 graviton is also shared by a spin- $\frac{3}{2}$ Rarita-Schwinger field coupled to a conserved vector-spinor current j^μ . This is not an unexpected result since, when the spin- $\frac{3}{2}$ field is coupled to the current, the supermatter interactions resulting from single-fermion interactions have precisely the form required by supersymmetry to complement single-graviton exchange between stress tensor sources [109]. Demonstrating this result using standard methods is a nontrivial exercise in supergravity manipulations [110]. Furthermore, we are motivated by the fact that, while all evidence so far suggest a massless graviton, the same is not true for the Rarita-Schwinger field. First, we would expect to have observed the gravitino were it massless. Secondly, various supersymmetry breaking mechanisms (such as the super-Higgs effect, or gravitationally induced SUSY breaking) have been shown to endow the Rarita-Schwinger particle with a non-zero mass [111–113].

In this article, we approach the spin- $\frac{3}{2}$ discontinuity from the point of view of on-shell massive amplitudes following the analysis in [1]. There it was shown that such a discontinuity can be observed in a purely on-shell manner by directly constructing the massive scattering amplitudes of the theory, taking their massless limit and comparing them with the constructed massless amplitudes. To be concrete, we consider scattering amplitudes in $\mathcal{N} = 1$ 4D supergravity, whose gauge multiplet consists entirely of a (spin-2) graviton and one Majorana (spin- $\frac{3}{2}$) spinor gravitino. This can be coupled to matter multiplets that preserve the supersymmetry, specifically an $\mathcal{N} = 1$ vector $(1, \frac{1}{2})$ multiplet consisting of a gauge-boson and gaugino, and an $\mathcal{N} = 1$ chiral $(\frac{1}{2}, 0)$ multiplet consisting of a spin- $\frac{1}{2}$ fermion and a complex scalar.

Giving the graviton a non-zero mass, increases the (on-shell) degrees of freedom

from two to five, introducing two vectors and one scalar. On the other hand, giving the gravitino mass results in two additional fermionic modes, corresponding to four on-shell degrees of freedom that are grouped by spin as $\{\pm\frac{3}{2}, \pm\frac{1}{2}\}$. In the case of the massive spin-2 action, the vector modes completely decouple, while the scalar modes couple to the trace of the matter stress energy tensor. In the massless limit then, any coupled matter that has a trace-free stress energy sector won't suffer a discontinuity, while any matter stress energy tensor with a non-zero trace will suffer one. From the on-shell scattering amplitude perspective, this same phenomena can be observed by computing the scattering of scalars via a massive graviton and comparing the massless limit of this amplitude with the same amplitude constructed from a massless graviton. The supersymmetric analog of this statement is that the fermionic modes of the gravitino couple to the Dirac gamma-trace of the current $\gamma_\mu j^\mu$ so in this case, we compare to amplitudes with non-zero gamma-trace.

2 The Supersymmetric vDVZ Discontinuity

To begin, we should first clarify what is the analogue of the vDVZ discontinuity in the supersymmetric context. Towards this end, we will utilise the Stückelberg formalism, appropriately adapted. Consider then, the free massive Rarita-Schwinger action

$$S = - \int d^4x \, e \left(\frac{1}{2} \bar{\Psi}_\mu^\alpha \gamma^{\mu\rho\nu} \partial_\rho \Psi_{\nu\alpha} - \frac{m}{2} \bar{\Psi}_\mu^\alpha \gamma^{\mu\nu} \Psi_{\nu\alpha} + \bar{\Psi}_\mu^\alpha j_\alpha^\mu \right) \equiv - \int d^4x \, e \, \mathcal{L}, \quad (2.1)$$

where α is a spinor index, μ, ν, ρ are Lorentz indices and e is, as usual, the determinant of the frame field $e_\mu^a(x)$. Our γ -conventions read

$$\gamma^{\mu\rho\nu} \equiv i\epsilon^{\mu\sigma\rho\nu} \gamma_5 \gamma_\sigma, \quad \gamma^{\mu\nu} \equiv \frac{i}{2} [\gamma^\mu, \gamma^\nu]. \quad (2.2)$$

Without the mass term, this action is invariant under

$$\Psi_\mu^\alpha \longrightarrow \Psi_\mu^\alpha + \partial_\mu \chi^\alpha, \quad \bar{\Psi}_\mu^\alpha \longrightarrow \bar{\Psi}_\mu^\alpha + \bar{\chi}^\alpha \overleftarrow{\partial}_\mu, \quad (2.3)$$

for some spinor χ^α . This symmetry is broken by the mass term but can be restored if we introduce the supercovariant derivative

$$D_\mu = \partial_\mu + \frac{1}{2} m \gamma_\mu, \quad (2.4)$$

in terms of which the Lagrangian becomes

$$\mathcal{L} = \frac{1}{2} \bar{\Psi}_\mu^\alpha \gamma^{\mu\rho\nu} D_\rho \Psi_{\nu\alpha} + \bar{\Psi}_\mu^\alpha j_\alpha^\mu, \quad (2.5)$$

after judicious use of the identity $\epsilon^{\mu\nu\rho\sigma} \gamma_5 \gamma_\nu = 2\gamma^\mu \gamma^{\rho\sigma}$. We can now introduce χ as a (spinorial) Stückelberg field via the transformation¹

$$\Psi_\mu^\alpha \longrightarrow \Psi_\mu^\alpha + \frac{1}{\sqrt{3}m} D_\mu \chi^\alpha. \quad (2.6)$$

Under this transformation, the Lagrangian density varies as

$$\delta\mathcal{L} = -\frac{\sqrt{3}m}{2} \left(\bar{\Psi}^\alpha \chi_\alpha + \bar{\chi}^\alpha \Psi_\alpha \right) - \bar{\chi}^\alpha (\not{\partial} + m) \chi_\alpha + \frac{1}{\sqrt{6}} \bar{\chi}^\alpha j_\alpha. \quad (2.7)$$

Subsequently, taking the massless limit does not lead to the original massless Lagrangian since,

$$\mathcal{L}^{massive} \Big|_{m \rightarrow 0} = \mathcal{L}^{massless} + \frac{1}{\sqrt{6}} \bar{\chi}^\alpha j_\alpha. \quad (2.8)$$

It is this that we identify as the SUSY equivalent of the vDVZ discontinuity, since any matter with a γ -traceless current will couple differently than matter with a non-vanishing γ -trace.

As alluded to in the introduction, we wish to couple to two chiral multiplets: a scalar multiplet $(\frac{1}{2}, 0)$ and a vector multiplet $(1, \frac{1}{2})$. The corresponding vector-spinor currents are, by Noethers theorem,

$$j_\alpha^\mu[\Phi, \psi] = [i\gamma^\nu \partial_\nu (\phi_1 - i\gamma_5 \phi_2) - m(\phi_1 + i\gamma_5 \phi_2)] \gamma^\mu \psi_\alpha \quad (2.9)$$

$$j_\alpha^\mu[A, \lambda] = \gamma^\rho \gamma^\nu \gamma^\mu \lambda_\alpha F_{\rho\nu}, \quad (2.10)$$

where $\Phi = \phi_1 + i\phi_2$ is a complex scalar, ψ_α a Majorana fermion (both with mass m_Φ), λ_α a massless photino and $F_{\rho\nu}$ the Maxwell tensor for photon A_ν . These are both conserved, i.e. that $\partial_\mu j_\alpha^\mu[\Phi, \psi] = \partial_\mu j_\alpha^\mu[A, \lambda] = 0$, however only one has a non-zero Dirac trace, e.g.

$$j_\alpha[\Phi, \psi] \neq 0, \quad j_\alpha[A, \lambda] = 0. \quad (2.11)$$

While this formulation is clearly off-shell, it will inform the on-shell investigation to which we now turn.

¹The factor of $\frac{1}{\sqrt{3}m}$ ensures a canonical fermionic kinetic term.

3 $\mathcal{N} = 1$ Supersymmetric Discontinuity

Let's start by thinking about the supermultiplets. To ensure that we preserve the correct symmetries we have to let all the particles in the multiplet have the same mass. To this end we will draw on the vector- and scalar-multiplet currents as they are stated in the previous section. In the interest of self-containment, many of the techniques and conventions used throughout this chapter will be set up in this first section.

3.1 Bold Notation and the Stückelberg Decomposition

As described in [6], we can significantly reduce the notational overhead that comes with incorporating $SU(2)$ little group indices in favour of bold-facing the massive particle spinors. Given our treatment of the action in section 2, it will be useful to perform a similar analysis at the level of the on-shell three-particle amplitudes. To this end, consider the coupling of a massless photon and massless fermion to a massive gravitino. The associated amplitude is given by

$$\mathcal{M}_3^{\{IJK\}}[1^{-1/2}, 2^{-1}, \mathbf{3}^{3/2}] = \frac{\kappa}{m} \langle \mathbf{13} \rangle \langle \mathbf{23} \rangle^2. \quad (3.1)$$

Simply unbolding this expression does *not* yield the correct result, since taking the massless limit tells us that the spin $-3/2$ mode of the gravitino diverges. Instead, we need to write it in a form where we *can* unbold, i.e.

$$\begin{aligned} \mathcal{M}_3^{\{IJK\}}[1^{-1/2}, 2^{-1}, \mathbf{3}^{3/2}] &= \frac{\kappa}{m^2} \langle 1 | p_2 | \mathbf{3} \rangle \langle \mathbf{23} \rangle^2 \\ &= -\kappa \frac{\langle 12 \rangle \langle \mathbf{23} \rangle^2 [\mathbf{32}]}{\langle \mathbf{23}^I \rangle [\mathbf{3}_I 2]}. \end{aligned} \quad (3.2)$$

Now, when we unbold, we find that the longitudinal spin $1/2$ mode of the gravitino is non-vanishing in the massless limit, since

$$\mathcal{M}_3^{\{IJK\}}[1^{-1/2}, 2^{-1}, \mathbf{3}^{3/2}] \Big|_{m \rightarrow 0} = -\frac{\kappa}{3} \langle 12 \rangle \langle 23 \rangle, \quad (3.3)$$

and with the inclusion of the relevant symmetry factor. This is consistent with the field theory treatment. On the other hand, we find that the $3/2$ mode does indeed survive, but for a different helicity choice of the fermion and photon

$$\mathcal{M}_3^{\{IJK\}}[1^{+1/2}, 2^{-1}, \mathbf{3}^{3/2}] = \kappa \frac{\langle \mathbf{23} \rangle^3}{\langle 12 \rangle} \longrightarrow \kappa \frac{\langle 23 \rangle^3}{\langle 12 \rangle}. \quad (3.4)$$

This implies that the act of expanding three-particle amplitudes into their various helicity components is the on-shell avatar of the Stückelberg decomposition, reflecting our result in section 2. This example manifests a subtlety in this approach; for the ‘unbolding’ of massive amplitudes to be physically meaningful, we will often have to make sure that we have teased out as much of the explicit mass dependence as possible, a point that will be important for what follows. Another, perhaps more easily digested, example is the three-particle amplitude of a massive vector and two massless scalars. This amplitude is treated in detail in [6] by making appropriate choices for the massive spinor indices to reveal the underlying helicities in the massless limit. The amplitude can be written in the generic form

$$\mathcal{A}^{\{IJ\}}[1^0, 2^0, \mathbf{3}^1] \propto \frac{\langle \mathbf{3} | p_1 p_2 | \mathbf{3} \rangle}{m^2} + \frac{\langle \mathbf{3} | p_1 | \mathbf{3} \rangle}{m} + \frac{[\mathbf{3} | p_1 p_2 | \mathbf{3}]}{m^2}. \quad (3.5)$$

Naively unbolding any of these terms to take the massless limit appears to be divergent. However, using the fact that $m^2 = \langle 12 \rangle [12]$ it can be rewritten as

$$\mathcal{A}^{\{IJ\}}[1^0, 2^0, \mathbf{3}^1] \propto \frac{\langle \mathbf{3} 1 \rangle \langle 2 \mathbf{3} \rangle}{\langle 12 \rangle} + m \frac{\langle \mathbf{3} 1 \rangle [1 \mathbf{3}]}{\langle 1 \mathbf{3}' \rangle [\mathbf{3}_I 1]} + \frac{[\mathbf{3} 1] [2 \mathbf{3}]}{[12]}. \quad (3.6)$$

Now that the mass factors in the denominators have been taken care of we can simply unbold and retain the term with the appropriate helicity. The first term is clearly the $h_3 = -1$ helicity mode, the last term the $h_3 = +1$ helicity mode and the scalar mode vanishes in the massless limit as expected. We see that we can simply extract the appropriate modes of the vector in the massless limit by finding the explicit mass dependence of a general form of the amplitude and then simply unbolding.

3.2 Vector Multiplet

We begin by computing the amplitudes involving an $\mathcal{N} = 1$ massless vector multiplet $(1, \frac{1}{2})$. This produces a three-particle vertex consisting of a gauge boson (a photon), a fermion (the photino) and a gravitino interacting with a coupling κ . We’ll require the propagating gravitino to be massive in order to study the effects of taking the massless limit. The associated four-particle diagram is given by Fig. 1.

We need only construct the left hand three-particle amplitude for all the possible helicity configurations. This is because the right hand three-particle amplitudes can be obtained by complex conjugation of eqs. (3.7) and (3.8), and making the replacements $p_1 \rightarrow p_3$, $p_2 \rightarrow p_4$. Throughout this article we will use the formalism developed in Ref. [6], including **bold** notation to highlight when a spinor represents a massive particle, suppressing the $SU(2)$ little group indices unless required for clarity. Using this formalism, we find that the possible three-particle amplitudes are




Figure 1. Vector multiplet four-particle

$$\mathcal{M}_3^{\{JKL\}}[1^{+1/2}, 2^{-1}, \mathbf{p}^{3/2}] = \frac{\kappa}{m^2} \langle 2\mathbf{p} \rangle^3 [12] = \kappa \frac{\langle 2\mathbf{p} \rangle^3}{\langle 12 \rangle}, \quad (3.7)$$

$$\mathcal{M}_3^{\{JKL\}}[1^{-1/2}, 2^{-1}, \mathbf{p}^{3/2}] = \frac{\kappa}{m} \langle 1\mathbf{p} \rangle \langle 2\mathbf{p} \rangle^2. \quad (3.8)$$

From this it is now simple to compute the two possible four-particle amplitudes by contracting the massive indices of the internal particle

$$\begin{aligned} \mathcal{M}_4[1^{+1/2}, 2^{-1}, 3^{-1/2}, 4^{+1}] &= \mathcal{M}^{\{IJK\}}[1^{+1/2}, 2^{-1}, \mathbf{p}^{3/2}] \frac{1}{p^2 + m^2} \tilde{\mathcal{M}}_{\{IJK\}}[3^{-1/2}, 4^{+1}, -\mathbf{p}^{3/2}] \\ &= -\frac{\kappa^2}{t + m^2} \frac{[14]}{[34]} \langle 2 | p_1 | 4 \rangle^2 \\ &= \frac{[14]}{[34]} \mathcal{M}_4[1^0, 2^{-1}, 3^0, 4^{+1}], \end{aligned} \quad (3.9)$$

where $\mathcal{M}_4[1^0, 2^{-1}, 3^0, 4^{+1}]$ is the massive graviton mediated scalar-photon amplitude [1]. This explicitly shows that the supersymmetric Ward identity is satisfied, as expected. This amplitude only has an explicit mass dependence in the propagator, and we can therefore easily take the massless limit. We want to compare the massless limit of the amplitude with the four-particle amplitude with an initially massless gravitino, which must be constructed from the three-particle amplitude

$$\mathcal{M}_3[1^{+1/2}, 2^{-1}, p^{-3/2}] = \kappa \frac{\langle 2p \rangle^3}{\langle 12 \rangle}. \quad (3.10)$$

Using this, we find that the massless four-particle amplitude is

$$\begin{aligned}\mathcal{M}_4[1^{+1/2}, 2^{-1}, 3^{-1/2}, 4^{+1}] \Big|_{m=0} &= \sum_{\pm} \mathcal{M}_3[1^{+1/2}, 2^{-1}, p^{\pm 3/2}] \frac{1}{p^2} \mathcal{M}_3[3^{-1/2}, 4^{+1}, -p^{\mp 3/2}] \\ &= -\frac{\kappa^2}{t} \frac{[14]}{[34]} \langle 2|p_1|4 \rangle^2,\end{aligned}\tag{3.11}$$

in agreement with eq. (3.9). Explicitly,

$$\mathcal{M}_4[1^{+1/2}, 2^{-1}, 3^{-1/2}, 4^{+1}] \Big|_{m \rightarrow 0} = \mathcal{M}_4[1^{+1/2}, 2^{-1}, 3^{-1/2}, 4^{+1}] \Big|_{m=0}.\tag{3.12}$$

If our only concern was reproducing the field-theory result in section 2 we could well stop at this point, noting that this is the only *chiral* amplitude possible for this field configuration. However, we do not wish to be led by the field theory construction, and so with no *a priori* reason to discard the other helicity possibility, we have include it. To construct the non-chiral four-particle amplitude, notice that the three-particle amplitude in eq.(3.8) is symmetric in two of its indices and, summing over the internal $SU(2)$ indices, allows us to write

$$\mathcal{M}_3^{\{JKL\}} \mathcal{M}_{3\{JKL\}} = \frac{1}{6} \mathcal{M}_3^{JKL} \mathcal{M}_{3\{JKL\}} = \frac{1}{3} \mathcal{M}_3^{JKL} (\mathcal{M}_{3JKL} + \mathcal{M}_{3KJL} + \mathcal{M}_{3LKJ}),$$

which in turn permits the four-particle amplitude to be written as

$$\begin{aligned}\mathcal{M}_4[1^{-1/2}, 2^{-1}, 3^{+1/2}, 4^{+1}] &= \mathcal{M}_3^{\{IJK\}}[1^{-1/2}, 2^{-1}, \mathbf{p}^{3/2}] \frac{1}{p^2 + m^2} \tilde{\mathcal{M}}_{3\{IJK\}}(3^{+1/2}, 4^{+1}, -\mathbf{p}^{3/2}) \\ &= \frac{\kappa^2}{3m^2(p^2 + m^2)} (\langle 1|p|3 \rangle \langle 2|p|4 \rangle^2 + 2 \langle 1|p|4 \rangle \langle 2|p|3 \rangle \langle 2|p|4 \rangle) \\ &= -\frac{\kappa^2}{3(p^2 + m^2)} \langle 12 \rangle [34] \langle 2|p|4 \rangle + \mathcal{O}(m^2),\end{aligned}\tag{3.13}$$

where, in the last line we have used the Schouten identity². Now we can easily take the massless limit to find

$$\begin{aligned}\mathcal{M}_4[1^{-1/2}, 2^{-1}, 3^{+1/2}, 4^{+1}] \Big|_{m \rightarrow 0} &= -\frac{\kappa^2}{3t} \langle 12 \rangle [34] \langle 2|p|4 \rangle \\ &= \frac{1}{3} \mathcal{M}_4[1^{-1/2}, 2^{-1}, 3^{+1/2}, 4^{+1}] \Big|_{m=0}\end{aligned}\tag{3.14}$$

²To see that the second piece of the amplitude is $\propto m^4$, multiply by $[12]^2/[12]^2$ and use conservation of momentum.

and where

$$\mathcal{M}_4[1^{-1/2}, 2^{-1}, 3^{+1/2}, 4^{+1}] \Big|_{m=0} = \mathcal{M}_3[1^{-1/2}, 2^{-1}, p^{+1/2}] \mathcal{M}_3[3^{+1/2}, 4^{+1}, -p^{-1/2}] / t.$$

From this it is clear that this helicity structure produces a contribution due to the spin- $\frac{1}{2}$ mode of the gravitino but with an overall factor of a $1/3$ when compared with its massless counterpart.

3.3 Scalar Multiplet

For the scalar multiplet, we need to consider a massive gravitino mediated interaction of a massive fermion and a massive scalar. For simplicity, we will take the masses of the latter two to be the same. The corresponding four-particle diagram is given in Fig. 2.




Figure 2. Scalar multiplet four-particle

Following Ref. [6] the all-massive three-particle amplitude can be constructed as

$$\begin{aligned} \mathcal{M}_3^{I\{JKL\}}[1^{1/2}, 2^0, p^{3/2}] &= g_1 \langle \mathbf{1} \mathbf{p} \rangle \langle \mathbf{p} \mathbf{p} \rangle \\ &+ g_2 (\langle \mathbf{1} \mathbf{p} \rangle \langle \mathbf{p} | p_1 p | \mathbf{p} \rangle + \langle \mathbf{p} \mathbf{p} \rangle (\langle \mathbf{1} | p_1 p | \mathbf{p} \rangle + \langle \mathbf{p} | p_1 p | \mathbf{1} \rangle)) \\ &+ g_3 (\langle \mathbf{1} | p_1 p | \mathbf{p} \rangle + \langle \mathbf{p} | p_1 p | \mathbf{1} \rangle) \langle \mathbf{p} | p_1 p | \mathbf{p} \rangle, \end{aligned} \quad (3.15)$$

where we have chosen to expand in a basis ($\mathcal{O}^{ab} = p_1^{\{a} p_b^{\} \}, \epsilon^{ab}$) and the coupling functions³ g_i have an undetermined mass dependence that will be fixed shortly. To simplify this, note that since the external $SU(2)$ indices of the gravitino have to be symmetrised, any term that has a factor $\langle \mathbf{p}^I \mathbf{p}^J \rangle = m \epsilon^{IJ}$ (where IJ are free indices) will vanish once symmetrised and can be ignored. Taking the external particle masses

³The ‘coupling functions’ are functions of particle mass and the gravitational coupling κ only, with the mass dependence needing to be fixed by e.g. demanding the correct high-energy behaviour

to be m_1 and the internal gravitino to have mass m , reduces the above amplitude to

$$\mathcal{M}_3^{I\{JKL\}}(\mathbf{1}^{1/2}, \mathbf{2}^0, \mathbf{p}^{3/2}) = m(-g_2 + g_3 m^2) \langle \mathbf{1p} \rangle \langle \mathbf{p} | p_1 | \mathbf{p} \rangle + 2m^2 m_1 g_3 [\mathbf{1p}] \langle \mathbf{p} | p_1 | \mathbf{p} \rangle. \quad (3.16)$$

In getting to this point we have made liberal use of the identities in appendix B. In order to fix the coupling functions g_i , we can compare these amplitudes with the ones derived in limit where one particle becomes massless. Taking the external mass to zero then, we find

$$\mathcal{M}_3^{I\{JKL\}}[\mathbf{1}^{1/2}, \mathbf{2}^0, \mathbf{p}^{3/2}] \Big|_{m_1 \rightarrow 0} = \begin{cases} \frac{\kappa}{m} \langle \mathbf{1p} \rangle \langle \mathbf{p} | p_1 | \mathbf{p} \rangle, & h_1 = -1/2 \\ \frac{\kappa}{m} [\mathbf{1p}] \langle \mathbf{p} | p_1 | \mathbf{p} \rangle, & h_1 = +1/2, \end{cases} \quad (3.17)$$

which implies that the following limits must hold

$$\lim_{m_1 \rightarrow 0} m(-g_2 + g_3 m^2) = \frac{\kappa}{m}, \quad \lim_{m_1 \rightarrow 0} 2m^2 m_1 g_3 = \frac{\kappa}{m}. \quad (3.18)$$

Next, to isolate any m_1 dependence, take $m \rightarrow 0$ so that

$$\mathcal{M}_3^{I\{JKL\}}[\mathbf{1}^{1/2}, \mathbf{2}^0, \mathbf{p}^{3/2}] \Big|_{m \rightarrow 0} \simeq \begin{cases} \kappa m_1 \langle \mathbf{1p} \rangle, & h_p = -1/2, \\ \kappa \frac{\langle \mathbf{1p} \rangle \langle p | p_1 | \xi \rangle}{[p\xi]}, & h_p = -3/2, \\ \kappa m_1 [\mathbf{1p}], & h_p = +1/2, \\ \kappa \frac{[\mathbf{1p}] \langle \xi | p_1 | p \rangle}{\langle \xi p \rangle}, & h_p = +3/2 \end{cases} \quad (3.19)$$

Now we can, for example, look at the $-3/2$ mode of the gravitino, choose $\xi = \eta$ and compare with the $I = J = K = 1$ amplitude,

$$\mathcal{M}_3^{I\{111\}}[\mathbf{1}^{1/2}, \mathbf{2}^0, \mathbf{p}^{3/2}] = \kappa \frac{\langle \mathbf{1p} \rangle \langle p | p_1 | \eta \rangle}{m} + \mathcal{O}(m). \quad (3.20)$$

Recognising that in the massless limit $m \rightarrow [p\eta]$, implies that the limits in eq. (3.18) must hold. In other words, the amplitude has no explicit m_1 -dependence (although it will have *implicit* dependence on m_1 to recover e.g. the $h_p = \pm 1/2$ amplitudes). Putting this together then, the three-particle amplitude must take the form

$$\mathcal{M}_3^{I\{JKL\}}[\mathbf{1}^{1/2}, \mathbf{2}^0, \mathbf{p}^{3/2}] = \frac{\kappa}{m} (\langle \mathbf{1p} \rangle + [\mathbf{1p}]) \langle \mathbf{p} | p_1 | \mathbf{p} \rangle. \quad (3.21)$$

With the three-particle amplitude in hand, we can now construct the four-particle amplitude. Given its lack of index symmetry however, the four-particle amplitude will have six distinct tensor structures and is particularly unwieldy. We will spare the reader the gory details. Suffice it to say that, with judicious use of the identities

in appendix B, it can be written as

$$\begin{aligned}\mathcal{M}_4^{I_1 I_3} [1^{1/2}, 2^0, 3^{1/2}, 4^0] &= \frac{\kappa^2}{6tm^2} (\langle 1|p|3\rangle - \langle 3|p|1\rangle) (-6m^2(2p_1 \cdot p_3) + 2(2p \cdot p_1)^2 \\ &\quad + 4m^2 m_1^2) - 4mm_1^2 (\langle 1|p|3\rangle + \langle 3|p|1\rangle) (2p \cdot p_1) \\ &\quad + 4m (\langle 13\rangle + [13]) (2p \cdot p_1)^2 + \mathcal{O}(m).\end{aligned}\quad (3.22)$$

Now, using the fact that $2p \cdot p_1 = -2p \cdot p_3 = -m^2$, we take the massless limit to get,

$$\mathcal{M}_4^{I_1 I_3} [1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m \rightarrow 0} = -\frac{\kappa^2}{t} (\langle 1|p|3\rangle - \langle 3|p|1\rangle) (2p_1 \cdot p_3 - \frac{2}{3}(m_1^2)). \quad (3.23)$$

This in turn needs to be compared to the four-particle amplitude with a massless gravitino exchange,

$$\begin{aligned}\mathcal{M}_4^{3/2 I_1 I_3} [1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m=0} &= \sum_{\pm 3/2} \mathcal{M}_3^{I_1 \{JKL\}} (1^{1/2}, 2^0, p^{\pm 3/2}) \frac{1}{t} \mathcal{M}_3^{I_3 \{JKL\}} (3^{1/2}, 4^0, -p^{\mp 3/2}) \\ &= \frac{\kappa^2}{t} (\langle 1|p|3\rangle \frac{\langle \zeta | p_3 p p_1 | \xi \rangle}{\langle \zeta | p | \xi \rangle} - \text{C.C.}),\end{aligned}\quad (3.24)$$

where ζ and ξ are reference spinors. To facilitate this comparison, we choose $\zeta = \xi$ and utilise conservation of momentum and the Schouten identity to write the amplitude as

$$\mathcal{M}_4^{3/2 I_1 I_3} [1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m=0} = -\frac{\kappa^2}{t} (\langle 1|p|3\rangle - \langle 3|p|1\rangle) (2p_1 \cdot p_3). \quad (3.25)$$

To make a concrete comparison of (3.23) with its massless propagator counterparts we also need to compute the four particle amplitude with a massless fermion exchange. Using the three particle amplitudes given in [6] this is straightforwardly computed as

$$\begin{aligned}\mathcal{M}_4^{1/2 I_1 I_3} [1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m=0} &= \sum_{\pm 1/2} \mathcal{M}_3^{I_1 \{JKL\}} (1^{1/2}, 2^0, p^{\pm 1/2}) \frac{1}{t} \mathcal{M}_3^{I_3 \{JKL\}} (3^{1/2}, 4^0, -p^{\mp 1/2}) \\ &= \frac{\kappa^2}{t} (\langle 1|p|3\rangle - \langle 3|p|1\rangle) (m_1^2).\end{aligned}\quad (3.26)$$

We are now in a position to write the amplitude (3.23) in terms of its massless counterparts (3.25) and (3.26) as

$$\mathcal{M}_4^{I_1 I_3} [1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m \rightarrow 0} = \mathcal{M}_4^{3/2 I_1 I_3} [1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m=0} + \frac{2}{3} \mathcal{M}_4^{1/2 I_1 I_3} [1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m=0}. \quad (3.27)$$

From this expression, it is evident that a discontinuity should be expected in

the scalar sector, in precisely the same form as from the field theory analysis. Furthermore, taking the external particle masses to zero and making a specific helicity choice shows that the four-particle amplitude satisfies the expected Ward identity

$$\mathcal{M}_4[1^{-1/2}, 2^0, 3^{+1/2}, 4^0] = \frac{\langle 12 \rangle}{\langle 32 \rangle} \mathcal{M}_4[1^0, 2^0, 3^0, 4^0], \quad (3.28)$$

where $\mathcal{M}_4[1^0, 2^0, 3^0, 4^0]$ is the massless limit of eq. 3.18 in Ref [1].

4 Supersymmetry Breaking and the Discontinuity

An interesting question that arises from our analysis above is whether this discontinuity persists below the supersymmetry breaking scale. Indeed, from the field theory perspective, none of the arguments for the existence of the discontinuity depend in any substantial way on whether the supersymmetry is broken or not, just that a massive spin- $\frac{3}{2}$ propagator couples to a current with $j \cdot \gamma = 0$ which is then compared to one for which $j \cdot \gamma \neq 0$. Our on-shell analysis however provides a simple test of this hypothesis by considering multiplets with distinct masses, *e.g.* a massive scalar and fermion with masses $m_s \neq m_f$.

The natural multiplets to consider in this case are,

1. the vector multiplet with a massless photon and a fermion of mass m_f and
2. the scalar multiplet with a massive scalar and massive fermion with respective masses $m_s \neq m_f$.

However there is a subtlety that requires some discussion. In the previous case with unbroken supersymmetry, the vector multiplet amplitudes only contain a single mass with which to constrain the coupling function, and while the scalar multiplet contains two distinct masses, there are well defined massless amplitudes to compare to (after taking appropriate massless limits), thereby allowing us to derive the correct mass structure. Breaking supersymmetry, on the other hand, results in even more masses that could form part of the coupling function. In this case, little group scaling and dimensional analysis alone are insufficient to constrain the masses.

For example, in the scalar multiplet where only the fermion is massive, according to the one-massive formula given in Ref. [6], the amplitude $\mathcal{M}_3[1^{1/2}, 2^0, p^{\pm 3/2}] = 0$. This is due to the requirement that $S + h_1 - h_2$ and $S + h_2 - h_1$ must both be positive, which is clearly not the case if $S = 1/2, h_1 = 0$ and $h_2 = \pm 3/2$. On the other hand, the fully massless amplitude *does* exist and is given by

$$\mathcal{M}_3[1^{-1/2}, 2^0, p^{-3/2}] = \kappa \frac{\langle 1p \rangle^2 \langle 2p \rangle}{\langle 12 \rangle}. \quad (4.1)$$

This is obviously not a limit of the one-massive case above, a fact that is also true if only the scalar is left massive. It seems then that the only consistent way to fix the mass dependence of the coupling function g_i is to demand that every amplitude has a well defined massless limit and discarding it if it doesn't. To constrain such coupling functions, let's consider the two-massive-one-massless three-particle amplitude, taking care to retain the gravitino as one of the massive particles. We will compare this to the amplitude with *only* the gravitino massive

$$\mathcal{M}_3^{\{IJK\}}[1^{-1/2}, 2^0, \mathbf{p}^{3/2}] = \frac{\kappa}{m} \langle 1\mathbf{p} \rangle \langle \mathbf{p} | p_1 | \mathbf{p} \rangle \quad (4.2)$$

since this does recover the massless amplitude in eq. (4.1). Demanding that we recover this amplitude from the two-massive amplitudes

$$\begin{aligned} \mathcal{M}_3^{I\{JKL\}}[1^{1/2}, 2^0, \mathbf{p}^{3/2}] &= \frac{\kappa}{m} (\langle 1\mathbf{p} \rangle \langle \mathbf{p} | p_1 | \mathbf{p} \rangle + [1\mathbf{p}] \langle \mathbf{p} | p_1 | \mathbf{p} \rangle) \\ \mathcal{M}_3^{\{JKL\}}[1^{-1/2}, 2^0, \mathbf{p}^{3/2}] &= \frac{\kappa}{m} \langle 1\mathbf{p} \rangle \langle \mathbf{p} | p_1 | \mathbf{p} \rangle, \end{aligned} \quad (4.3)$$

constrains the three-massive particle amplitude to the form,

$$\mathcal{M}_3^{I\{JKL\}}[1^{1/2}, 2^0, \mathbf{p}^{3/2}] = \frac{\kappa}{m} (\langle 1\mathbf{p} \rangle + [1\mathbf{p}]) \langle \mathbf{p} | p_1 | \mathbf{p} \rangle. \quad (4.4)$$

The construction of the four-particle amplitude now mirrors that outlined in the previous section exactly, with the result that

$$\begin{aligned} \mathcal{M}_4^{I_1 I_3}[1^{1/2}, 2^0, 3^{1/2}, 4^0] &= \frac{\kappa^2}{6tm^2} ((\langle 1 | p | 3 \rangle - \langle 3 | p | 1 \rangle) (-6m^2(2p_1 \cdot p_3) + 2(2p \cdot p_1)^2 \\ &\quad + 4m^2m_f^2) - 4mm_f^2(\langle 1 | p | 3 \rangle + \langle 3 | p | 1 \rangle)(2p \cdot p_1) \\ &\quad + 4m(\langle 13 \rangle + [13])(2p \cdot p_1)^2) + \mathcal{O}(m). \end{aligned} \quad (4.5)$$

The difference between this expression and (3.23) is that $2p \cdot p_1 = m^2 + m_f^2 - m_s^2 = -2p \cdot p_3$. Correspondingly, the amplitude (4.5) will have terms of order $\mathcal{O}(m^{-1})$ and therefore has no consistent massless limit. This appears to be an artefact of constructing massive three-particle amplitudes in this formalism, and specifically fixing the mass structure of the arbitrary coupling functions. If this inconsistency were somehow resolved, we would expect the mass factor in the denominator to have an additive form, $m + m_f - m_s$, which would allow the massless limit to be taken in the four-particle amplitude. That said, it is not clear to us how such a denominator could arise from a local quantum field theory with polynomial interactions. For example, an action containing a term of the form $\sim \phi \bar{\Psi}^\alpha \psi_\alpha$ will have at most a single mass in the denominator coming from the Rarita-Schwinger polarization. In this sense, a mass dependence of $\mathcal{O}(m^{-1})$ is to be expected and therefore the formalism does seemingly produce the correct amplitude, but one which

apparently doesn't have well defined massless limits. It is entirely plausible that this problem is cured by a high energy mechanism, e.g. the Higgs, which does in some circumstances restore the high energy limit via the goldstone equivalence theorem. The gravitino-goldstino equivalence theorem shows that spontaneously broken symmetries involving longitudinally polarized gravitinos can have a well defined high energy behaviour via the goldstone equivalence [114]. However, it is not clear to us that this will solve all of the issues raised in this section, and is at any rate beyond the scope of this article and so we leave this investigation to the future.

Amplitudes for the vector multiplet exhibit a similar inconsistency in that the amplitude with a single massive photino does not reduce to the required massless limit. Specifically,

$$\mathcal{M}_3^I[\mathbf{1}^{1/2}, 2^{-1}, p^{-3/2}] = \frac{\kappa}{m_f} \langle \mathbf{1}p \rangle \langle p2 \rangle^2 \quad (4.6)$$

cannot reproduce the amplitude $\mathcal{M}_3[1^{-1/2}, 2^{-1}, p^{-3/2}] = 0$ in the limit that $m_f \rightarrow 0$. Again we turn to the case where the gravitino is the only massive particle to fix the coupling function. The corresponding amplitude,

$$M^{I\{JKL\}}[\mathbf{1}^{1/2}, 2^{-1}, \mathbf{p}^{3/2}] = \frac{\kappa}{m} ([\mathbf{1}\mathbf{p}]\langle \mathbf{p}2 \rangle^2 + \langle \mathbf{1}\mathbf{p} \rangle \langle \mathbf{p}2 \rangle^2), \quad (4.7)$$

correctly reproduces all of the expected three-particle amplitudes in the relevant limits. From this, and with some straightforward but tedious simplification, we find the four-particle amplitude

$$\begin{aligned} \mathcal{M}_4^{I_1 I_3}[\mathbf{1}^{1/2}, 2^{-1}, \mathbf{3}^{1/2}, 4^{-1}] &= \frac{\kappa^2}{3tm^2} (\langle \mathbf{1}2 \rangle [\mathbf{3}4] (-m^2 - m_f^2 - 3u) + \langle \mathbf{2}3 \rangle [\mathbf{1}4] (3m_f^2 + 3m^2) \\ &\quad + (\langle \mathbf{1}3 \rangle + [\mathbf{1}3]) \langle 2|p|4 \rangle (3m_f - m) \langle 2|p|4 \rangle). \end{aligned} \quad (4.8)$$

Again there is no consistent massless limit to be taken and a similar argument as in the scalar multiplet follows. Clearly, this issue requires further analysis.

5 Discussion

Using the recently developed formalism for massive scattering amplitudes, we have shown in this article that the chiral scalar multiplet and the non-chiral vector multiplet of massive supergravity both produce discontinuous scattering amplitudes in the $m \rightarrow 0$ limit, reproducing on-shell the field theory result of [110]. That said, scattering amplitudes themselves are not observables. Indeed, the primary significance of the original vDVZ discontinuity of massive gravity lies in the empirical fact that the light bending angle in massive gravity deviates from observation by a factor of $\frac{3}{4}$, provided the amplitudes are normalised to recover

Newtonian gravity. This argument is obviously less relevant in the context of supergravity. However, our results establish the discontinuity at the amplitude level as an on-shell avatar of the Stückelberg decomposition.

Let's unpack this a little, beginning with our results from the supersymmetric case. In all the diagrams in this article, our conventions are that time moves upwards and external particles are considered outgoing. This results in a decided difference in the two amplitudes computed in the supersymmetric vector multiplet. The first that we computed in (3.9) describes the chiral multiplet. This amplitude does *not* have a discontinuity in the massless limit in that $\mathcal{M}_4[1^{+1/2}, 2^{-1}, 3^{-1/2}, 4^{+1}] \Big|_{m \rightarrow 0} = \mathcal{M}_4^{3/2}[1^{+1/2}, 2^{-1}, 3^{-1/2}, 4^{+1}] \Big|_{m=0}$. The next one that we computed, (3.13) describes a non-chiral multiplet. This amplitude does indeed exhibit a discontinuity in the massless limit, $\mathcal{M}_4[1^{-1/2}, 2^{-1}, 3^{+1/2}, 4^{+1}] \Big|_{m \rightarrow 0} = \frac{1}{3} \mathcal{M}_4^{1/2}[1^{-1/2}, 2^{-1}, 3^{+1/2}, 4^{+1}] \Big|_{m=0}$ ⁴. Guided by the fact that the field theory is chiral, we need only consider that piece. It is worth noting that if our only source of information came from on-shell amplitude methods we would necessarily have to include the non-chiral part, resulting in an ambiguity in the realization of discontinuity. An interesting problem for the future would be to understand how to project out the chiral part of the amplitude. Moving on to the scalar multiplet, we find that

$$\begin{aligned} \mathcal{M}_4^{I_1 I_3}[1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m \rightarrow 0} &= \mathcal{M}_4^{3/2 I_1 I_3}[1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m=0} \\ &+ \frac{2}{3} \mathcal{M}_4^{1/2 I_1 I_3}[1^{1/2}, 2^0, 3^{1/2}, 4^0] \Big|_{m=0}, \end{aligned}$$

manifesting the discontinuity anticipated from the field theory analysis.

Next, we attempted to see the discontinuity with supersymmetry broken, endowing the external particle species with distinct masses. Sadly, we were unable to express the relevant four-particle amplitudes in a form in which the massless limit can be cleanly taken. This is due to how the structure of the coupling functions in the three-particle amplitudes is determined. Specifically, the amplitudes can be organised such that they reveal a mass dependence of the order $\sim 1/m$ and is not finite in the massless limit while, for all of the massless limits to be well defined, we require a mass dependence of the form $1/(m + m_f - m_s)$. We remain unsatisfied with this puzzle but leave its resolution for future work. Another noteworthy point in the non-supersymmetric case is that the vector multiplet amplitude (4.8) contains

⁴The superscript on the amplitude here denotes the spin of the massless propagator.

terms that correspond to both the chiral and non-chiral pieces as a result of the now massive fermion.

The on-shell technology developed in [6] for massive particle scattering is both conceptually and computationally powerful, but like any new technology, it requires extensive beta-testing to iron out any bugs. This article details one such test. We set out to give an on-shell derivation of the spin- $\frac{3}{2}$ analogue of the famous vDVZ discontinuity of massive gravity, expecting the calculation to be clean and unambiguous. We were met with several subtleties, some of which we were able to resolve and some, like how to treat symmetry breaking and chirality, we remain puzzled by. Clearly though, there is still much work that remains to be done.

Acknowledgements

We would like to thank the anonymous referee of one of our previous papers for pointing us to this problem. JM is supported by the NRF of South Africa under grant CSUR 114599. DB and NM is supported by funding from the DST/NRF SARCHI in Physical Cosmology. Any opinion, finding and conclusion or recommendation expressed in this material is that of the authors and the NRF does not accept any liability in this regard.

A Conventions and Notation

In the interests of self-containment, we collect here some of our conventions for computing massive amplitudes that we use in the main text. Our convention for massive spinors can be summarized as:

- If no up-down massive spinor index pair is explicit on adjacent massive spinors they are considered to be contracted.
- For spinor indices we use the lower case Latin alphabet and for little-group indices we use the upper case Latin alphabet.
- Epsilon conventions and helicity basis:

$$\epsilon_{ab} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad \epsilon^{ab} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad \zeta^{-I} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \zeta^{+I} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- Mandelstam variables:

$$s = -(p_1 + p_4)^2, \quad u = -(p_1 + p_3)^2, \quad t = -(p_1 + p_2)^2$$

- Massless spinor relations:

$$p^{ab} = -|p\rangle^a [p]^b, \quad |-p\rangle = -|p\rangle, \quad |-p] = [p]$$

- Massive spinor relations:

$$\begin{aligned} p^{ab} &= |\mathbf{p}\rangle^{aI} {}_I[\mathbf{p}]^b = -|p\rangle^a [p]^b - |\eta_p\rangle^a [\eta_p]^b, \quad p^2 = \det(p^{ab}) = -m_p^2, \\ \langle p\eta_p \rangle &= [p\eta_p] = m_p, \quad \langle \mathbf{p} |_a^I p^{ab} = m [\mathbf{p}]^{bI}, \quad [\mathbf{p}]^{\dot{a}I} p_{\dot{a}b} = -m \langle \mathbf{p} |_b^I, \\ \langle i|pp|j \rangle &= -m_p^2 \langle ij \rangle, \quad {}^I\langle \mathbf{p}\mathbf{p} \rangle^J = m \epsilon^{IJ} {}_I[\mathbf{p}\mathbf{p}]_J = -m \epsilon_{IJ}, \\ \langle i\mathbf{p} \rangle^I {}_I\langle \mathbf{p} j \rangle &= m \langle ij \rangle, \quad [i\mathbf{p}]^I {}_I[\mathbf{p} j] = m [ij], \quad |\mathbf{p}\rangle^I {}_J[\mathbf{p}] = -|\mathbf{p}\rangle_I^J [\mathbf{p}] \end{aligned}$$

B Some Miscellaneous Identities

- Simplification of the general scalar multiplet amplitude where we symmetrize over the massive relevant massive indices

$$\begin{aligned} \langle \mathbf{p}^{\{I} \mathbf{p}^{J\}} \rangle &= m \epsilon^{\{IJ\}} = 0, \\ \langle \mathbf{1} | p_1 p | \mathbf{p} \rangle &= -m_f m [\mathbf{1}\mathbf{p}], \\ \langle \mathbf{p} | p_1 p | \mathbf{1} \rangle &= m m_f [\mathbf{1}\mathbf{p}] - (m^2 + m_f^2 - m_s^2) \langle \mathbf{1}\mathbf{p} \rangle, \\ \langle \mathbf{p} | p_1 p | \mathbf{p} \rangle &= -m \langle \mathbf{p} | p_1 | \mathbf{p} \rangle. \end{aligned} \tag{B.1}$$

- Here are some identities commonly used in the scalar multiplet amplitudes. We keep the masses of the three particles distinct for clarity. For the terms in the all massive three-particle amplitude that are not listed here, simply exchange

$1 \rightleftharpoons 3$ or where relevant complex conjugate.

$$\begin{aligned}
\langle \mathbf{p}^I | p_1 | \mathbf{p}^J | [\mathbf{p}_J | p_3 | \mathbf{p}_I] &= -m^2 (2p_1 \cdot p_3) \\
\langle \mathbf{p}^I | p_1 p p_3 | \mathbf{p}_I] &= m^2 (2p_1 \cdot p_3) + (2p \cdot p_1)(2p \cdot p_3) \\
\langle \mathbf{1} | p p_3 p p_1 p | \mathbf{3} \rangle &= m^2 \langle \mathbf{1} | p p_3 p_1 | \mathbf{3} \rangle + m^2 m_f \langle \mathbf{13} \rangle (2p \cdot p_1) + \langle \mathbf{1} | p | \mathbf{3} \rangle (2p \cdot p_1)(2p \cdot p_3) \\
\langle \mathbf{1} | p p_3 p p_1 | \mathbf{3} \rangle &= m^2 \langle \mathbf{1} | p_3 p_1 | \mathbf{3} \rangle - m_f \langle \mathbf{3} | p | \mathbf{1} \rangle (2p \cdot p_3) + \langle \mathbf{13} \rangle (2p \cdot p_3)^2 \\
\langle \mathbf{1} | p p_3 p_1 p | \mathbf{3} \rangle &= m^2 \langle \mathbf{1} | p_3 p_1 | \mathbf{3} \rangle - m_f \langle \mathbf{1} | p | \mathbf{3} \rangle (2p \cdot p_1) + \langle \mathbf{13} \rangle (2p \cdot p_1)^2 - m_f \langle \mathbf{3} | p | \mathbf{1} \rangle (2p \cdot p_1) \\
\langle \mathbf{1} | p_3 p p_1 p | \mathbf{3} \rangle &= -m^2 \langle \mathbf{1} | p_3 p_1 | \mathbf{3} \rangle + m_f \langle \mathbf{1} | p | \mathbf{3} \rangle (2p \cdot p_1) - \langle \mathbf{13} \rangle (2p \cdot p_1)^2 \\
\langle \mathbf{1} | p_3 p_1 p | \mathbf{3} \rangle &= -m_f^2 \langle \mathbf{3} | p | \mathbf{1} \rangle - m_f [\mathbf{13}] (2p \cdot p_3) + \langle \mathbf{1} | p | \mathbf{3} \rangle (2p_1 \cdot p_3) \\
\langle \mathbf{1} | p_3 p p_1 | \mathbf{3} \rangle &= m_f^2 \langle \mathbf{3} | p | \mathbf{1} \rangle - m_f \langle \mathbf{13} \rangle (2p \cdot p_1) - \langle \mathbf{1} | p | \mathbf{3} \rangle (2p_1 \cdot p_3) + m_f [\mathbf{13}] (2p \cdot p_3) \\
\langle \mathbf{1} | p p_3 p_1 | \mathbf{3} \rangle &= -m_f^2 \langle \mathbf{3} | p | \mathbf{1} \rangle + m_f \langle \mathbf{13} \rangle (2p \cdot p_1) + \langle \mathbf{1} | p | \mathbf{3} \rangle (2p_1 \cdot p_3)
\end{aligned} \tag{B.2}$$

References

- [1] N. Moynihan and J. Murugan, *Comments on scattering in massive gravity, vDVZ and BCFW*, *Class. Quant. Grav.* **35** (2018) 155005 [[1711.03956](#)].
- [2] C. de Rham, G. Gabadadze and A. J. Tolley, *Resummation of Massive Gravity*, *Phys. Rev. Lett.* **106** (2011) 231101 [[1011.1232](#)].
- [3] A. Vainshtein, *To the problem of nonvanishing gravitation mass*, *Physics Letters B* **39** (1972) 393 .
- [4] F. Cachazo and A. Guevara, *Leading Singularities and Classical Gravitational Scattering*, [1705.10262](#).
- [5] A. Guevara, *Holomorphic Classical Limit for Spin Effects in Gravitational and Electromagnetic Scattering*, *JHEP* **04** (2019) 033 [[1706.02314](#)].
- [6] N. Arkani-Hamed, T.-C. Huang and Y.-t. Huang, *Scattering Amplitudes For All Masses and Spins*, [1709.04891](#).
- [7] A. Guevara, A. Ochirov and J. Vines, *Scattering of Spinning Black Holes from Exponentiated Soft Factors*, [1812.06895](#).
- [8] M.-Z. Chung, Y.-T. Huang, J.-W. Kim and S. Lee, *The simplest massive S-matrix: from minimal coupling to Black Holes*, *JHEP* **04** (2019) 156 [[1812.08752](#)].
- [9] R. Carballo-Rubio, F. Di Filippo and N. Moynihan, *Taming higher-derivative interactions and bootstrapping gravity with soft theorems*, [1811.08192](#).
- [10] S. Caron-Huot and Z. Zahraee, *Integrability of Black Hole Orbits in Maximal Supergravity*, *JHEP* **07** (2019) 179 [[1810.04694](#)].
- [11] A. Guevara, A. Ochirov and J. Vines, *Black-hole scattering with general spin directions from minimal-coupling amplitudes*, [1906.10071](#).

- [12] W. T. Emond and N. Moynihan, *Scattering Amplitudes, Black Holes and Leading Singularities in Cubic Theories of Gravity*, [1905.08213](#).
- [13] N. Arkani-Hamed, Y.-t. Huang and D. O’Connell, *Kerr Black Holes as Elementary Particles*, [1906.10100](#).
- [14] N. Moynihan, *Kerr-Newman from Minimal Coupling*, *JHEP* **01** (2020) 014 [[1909.05217](#)].
- [15] Y.-T. Huang, U. Kol and D. O’Connell, *The Double Copy of Electric-Magnetic Duality*, [1911.06318](#).
- [16] M.-Z. Chung, Y.-T. Huang and J.-W. Kim, *Kerr-Newman stress-tensor from minimal coupling to all orders in spin*, [1911.12775](#).
- [17] A. Cristofoli, N. E. J. Bjerrum-Bohr, P. H. Damgaard and P. Vanhove, *On Post-Minkowskian Hamiltonians in General Relativity*, [1906.01579](#).
- [18] M.-Z. Chung, Y.-t. Huang and J.-W. Kim, *From quantized spins to rotating black holes*, [1908.08463](#).
- [19] N. E. J. Bjerrum-Bohr, A. Cristofoli and P. H. Damgaard, *Post-Minkowskian Scattering Angle in Einstein Gravity*, [1910.09366](#).
- [20] D. J. Burger, W. T. Emond and N. Moynihan, *Rotating Black Holes in Cubic Gravity*, *Phys. Rev. D* **101** (2020) 084009 [[1910.11618](#)].
- [21] N. Moynihan and J. Murugan, *On-Shell Electric-Magnetic Duality and the Dual Graviton*, [2002.11085](#).
- [22] Z. Bern, J. J. M. Carrasco and H. Johansson, *Perturbative Quantum Gravity as a Double Copy of Gauge Theory*, *Phys. Rev. Lett.* **105** (2010) 061602 [[1004.0476](#)].
- [23] Z. Bern, L. J. Dixon, D. C. Dunbar, M. Perelstein and J. S. Rozowsky, *On the relationship between Yang-Mills theory and gravity and its implication for ultraviolet divergences*, *Nucl. Phys. B* **530** (1998) 401 [[hep-th/9802162](#)].
- [24] M. B. Green, J. H. Schwarz and L. Brink, *$N=4$ Yang-Mills and $N=8$ Supergravity as Limits of String Theories*, *Nucl. Phys. B* **198** (1982) 474.
- [25] Z. Bern, J. S. Rozowsky and B. Yan, *Two loop four gluon amplitudes in $N=4$ superYang-Mills*, *Phys. Lett. B* **401** (1997) 273 [[hep-ph/9702424](#)].
- [26] J. J. M. Carrasco and H. Johansson, *Five-Point Amplitudes in $N=4$ Super-Yang-Mills Theory and $N=8$ Supergravity*, *Phys. Rev. D* **85** (2012) 025006 [[1106.4711](#)].
- [27] J. J. M. Carrasco, M. Chiodaroli, M. Gunaydin and R. Roiban, *One-loop four-point amplitudes in pure and matter-coupled $N = 4$ supergravity*, *JHEP* **03** (2013) 056 [[1212.1146](#)].
- [28] C. R. Mafra and O. Schlotterer, *The Structure of n -Point One-Loop Open Superstring Amplitudes*, *JHEP* **08** (2014) 099 [[1203.6215](#)].

- [29] R. H. Boels, R. S. Isermann, R. Monteiro and D. O’Connell, *Colour-Kinematics Duality for One-Loop Rational Amplitudes*, *JHEP* **04** (2013) 107 [[1301.4165](#)].
- [30] N. E. J. Bjerrum-Bohr, T. Dennen, R. Monteiro and D. O’Connell, *Integrand Oxidation and One-Loop Colour-Dual Numerators in $N=4$ Gauge Theory*, *JHEP* **07** (2013) 092 [[1303.2913](#)].
- [31] Z. Bern, S. Davies, T. Dennen, Y.-t. Huang and J. Nohle, *Color-Kinematics Duality for Pure Yang-Mills and Gravity at One and Two Loops*, *Phys. Rev.* **D92** (2015) 045041 [[1303.6605](#)].
- [32] Z. Bern, S. Davies and T. Dennen, *The Ultraviolet Structure of Half-Maximal Supergravity with Matter Multiplets at Two and Three Loops*, *Phys. Rev.* **D88** (2013) 065007 [[1305.4876](#)].
- [33] J. Nohle, *Color-Kinematics Duality in One-Loop Four-Gluon Amplitudes with Matter*, *Phys. Rev.* **D90** (2014) 025020 [[1309.7416](#)].
- [34] Z. Bern, S. Davies, T. Dennen, A. V. Smirnov and V. A. Smirnov, *Ultraviolet Properties of $N=4$ Supergravity at Four Loops*, *Phys. Rev. Lett.* **111** (2013) 231302 [[1309.2498](#)].
- [35] S. G. Naculich, H. Nastase and H. J. Schnitzer, *All-loop infrared-divergent behavior of most-subleading-color gauge-theory amplitudes*, *JHEP* **04** (2013) 114 [[1301.2234](#)].
- [36] C. R. Mafra and O. Schlotterer, *Towards one-loop SYM amplitudes from the pure spinor BRST cohomology*, *Fortsch. Phys.* **63** (2015) 105 [[1410.0668](#)].
- [37] Z. Bern, S. Davies and T. Dennen, *Enhanced ultraviolet cancellations in $\mathcal{N} = 5$ supergravity at four loops*, *Phys. Rev.* **D90** (2014) 105011 [[1409.3089](#)].
- [38] C. R. Mafra and O. Schlotterer, *Two-loop five-point amplitudes of super Yang-Mills and supergravity in pure spinor superspace*, *JHEP* **10** (2015) 124 [[1505.02746](#)].
- [39] S. He, R. Monteiro and O. Schlotterer, *String-inspired BCJ numerators for one-loop MHV amplitudes*, *JHEP* **01** (2016) 171 [[1507.06288](#)].
- [40] Z. Bern, S. Davies and J. Nohle, *Double-Copy Constructions and Unitarity Cuts*, *Phys. Rev.* **D93** (2016) 105015 [[1510.03448](#)].
- [41] G. Mogull and D. O’Connell, *Overcoming Obstacles to Colour-Kinematics Duality at Two Loops*, *JHEP* **12** (2015) 135 [[1511.06652](#)].
- [42] M. Chiodaroli, M. Gunaydin, H. Johansson and R. Roiban, *Spontaneously Broken Yang-Mills-Einstein Supergravities as Double Copies*, *JHEP* **06** (2017) 064 [[1511.01740](#)].
- [43] Z. Bern, J. J. M. Carrasco, W.-M. Chen, H. Johansson, R. Roiban and M. Zeng, *Five-loop four-point integrand of $N = 8$ supergravity as a generalized double copy*, *Phys. Rev.* **D96** (2017) 126012 [[1708.06807](#)].

- [44] H. Johansson and A. Ochirov, *Color-Kinematics Duality for QCD Amplitudes*, *JHEP* **01** (2016) 170 [[1507.00332](#)].
- [45] S. Oxburgh and C. D. White, *BCJ duality and the double copy in the soft limit*, *JHEP* **02** (2013) 127 [[1210.1110](#)].
- [46] Y. Geyer, R. Monteiro and R. Stark-Muchão, *Two-Loop Scattering Amplitudes: Double-Forward Limit and Colour-Kinematics Duality*, *JHEP* **12** (2019) 049 [[1908.05221](#)].
- [47] C. D. White, *Factorization Properties of Soft Graviton Amplitudes*, *JHEP* **05** (2011) 060 [[1103.2981](#)].
- [48] S. Melville, S. G. Naculich, H. J. Schnitzer and C. D. White, *Wilson line approach to gravity in the high energy limit*, *Phys. Rev.* **D89** (2014) 025009 [[1306.6019](#)].
- [49] A. Luna, S. Melville, S. G. Naculich and C. D. White, *Next-to-soft corrections to high energy scattering in QCD and gravity*, *JHEP* **01** (2017) 052 [[1611.02172](#)].
- [50] R. Saotome and R. Akhoury, *Relationship Between Gravity and Gauge Scattering in the High Energy Limit*, *JHEP* **01** (2013) 123 [[1210.8111](#)].
- [51] A. Sabio Vera, E. Serna Campillo and M. A. Vazquez-Mozo, *Color-Kinematics Duality and the Regge Limit of Inelastic Amplitudes*, *JHEP* **04** (2013) 086 [[1212.5103](#)].
- [52] H. Johansson, A. Sabio Vera, E. Serna Campillo and M. Á. Vázquez-Mozo, *Color-Kinematics Duality in Multi-Regge Kinematics and Dimensional Reduction*, *JHEP* **10** (2013) 215 [[1307.3106](#)].
- [53] H. Johansson, A. Sabio Vera, E. Serna Campillo and M. A. Vazquez-Mozo, *Color-kinematics duality and dimensional reduction for graviton emission in Regge limit*, in *International Workshop on Low X Physics (Israel 2013) Eilat, Israel, May 30-June 04, 2013*, 2013, [1310.1680](#).
- [54] T. Bargheer, S. He and T. McLoughlin, *New Relations for Three-Dimensional Supersymmetric Scattering Amplitudes*, *Phys. Rev. Lett.* **108** (2012) 231601 [[1203.0562](#)].
- [55] Y.-t. Huang and H. Johansson, *Equivalent $D=3$ Supergravity Amplitudes from Double Copies of Three-Algebra and Two-Algebra Gauge Theories*, *Phys. Rev. Lett.* **110** (2013) 171601 [[1210.2255](#)].
- [56] G. Chen and Y.-J. Du, *Amplitude Relations in Non-linear Sigma Model*, *JHEP* **01** (2014) 061 [[1311.1133](#)].
- [57] M. Chiodaroli, Q. Jin and R. Roiban, *Color/kinematics duality for general abelian orbifolds of $N=4$ super Yang-Mills theory*, *JHEP* **01** (2014) 152 [[1311.3600](#)].
- [58] H. Johansson and A. Ochirov, *Pure Gravities via Color-Kinematics Duality for Fundamental Matter*, *JHEP* **11** (2015) 046 [[1407.4772](#)].
- [59] H. Johansson and J. Nohle, *Conformal Gravity from Gauge Theory*, [1707.02965](#).

- [60] C. Cheung, C.-H. Shen and C. Wen, *Unifying Relations for Scattering Amplitudes*, *JHEP* **02** (2018) 095 [[1705.03025](#)].
- [61] M. Chiodaroli, M. Gunaydin, H. Johansson and R. Roiban, *Gauged Supergravities and Spontaneous Supersymmetry Breaking from the Double Copy Construction*, *Phys. Rev. Lett.* **120** (2018) 171601 [[1710.08796](#)].
- [62] G. Chen, H. Johansson, F. Teng and T. Wang, *On the kinematic algebra for BCJ numerators beyond the MHV sector*, *JHEP* **11** (2019) 055 [[1906.10683](#)].
- [63] J. Plefka, C. Shi and T. Wang, *Double copy of massive scalar QCD*, *Phys. Rev.* **D101** (2020) 066004 [[1911.06785](#)].
- [64] R. Aoude and A. Helset, *Soft Theorems and the KLT-Relation*, *JHEP* **04** (2020) 044 [[1911.09962](#)].
- [65] A. E. Lipstein and P. McFadden, *Double copy structure and the flat space limit of conformal correlators in even dimensions*, [1912.10046](#).
- [66] T. Adamo and A. Ilderton, *Classical and quantum double copy of back-reaction*, [2005.05807](#).
- [67] L. Borsten and S. Nagy, *The pure BRST Einstein-Hilbert Lagrangian from the double-copy to cubic order*, [2004.14945](#).
- [68] Z. Bern, J. J. Carrasco, M. Chiodaroli, H. Johansson and R. Roiban, *The Duality Between Color and Kinematics and its Applications*, [1909.01358](#).
- [69] R. Monteiro, D. O’Connell and C. D. White, *Black holes and the double copy*, *JHEP* **12** (2014) 056 [[1410.0239](#)].
- [70] A. Luna, R. Monteiro, D. O’Connell and C. D. White, *The classical double copy for Taub-NUT spacetime*, *Phys. Lett.* **B750** (2015) 272 [[1507.01869](#)].
- [71] A. Luna, R. Monteiro, I. Nicholson, D. O’Connell and C. D. White, *The double copy: Bremsstrahlung and accelerating black holes*, *JHEP* **06** (2016) 023 [[1603.05737](#)].
- [72] W. D. Goldberger and A. K. Ridgway, *Radiation and the classical double copy for color charges*, *Phys. Rev.* **D95** (2017) 125010 [[1611.03493](#)].
- [73] A. Anastasiou, L. Borsten, M. J. Duff, L. J. Hughes and S. Nagy, *Yang-Mills origin of gravitational symmetries*, *Phys. Rev. Lett.* **113** (2014) 231606 [[1408.4434](#)].
- [74] L. Borsten and M. J. Duff, *Gravity as the square of Yang-Mills?*, *Phys. Scripta* **90** (2015) 108012 [[1602.08267](#)].
- [75] A. Anastasiou, L. Borsten, M. J. Duff, M. J. Hughes, A. Marrani, S. Nagy et al., *Twin supergravities from Yang-Mills theory squared*, *Phys. Rev.* **D96** (2017) 026013 [[1610.07192](#)].
- [76] A. Anastasiou, L. Borsten, M. J. Duff, A. Marrani, S. Nagy and M. Zoccali, *Are all supergravity theories Yang-Mills squared?*, *Nucl. Phys.* **B934** (2018) 606 [[1707.03234](#)].

- [77] G. L. Cardoso, S. Nagy and S. Nampuri, *A double copy for $\mathcal{N} = 2$ supergravity: a linearised tale told on-shell*, *JHEP* **10** (2016) 127 [[1609.05022](#)].
- [78] L. Borsten, *$D = 6$, $\mathcal{N} = (2, 0)$ and $\mathcal{N} = (4, 0)$ theories*, *Phys. Rev.* **D97** (2018) 066014 [[1708.02573](#)].
- [79] A. Anastasiou, L. Borsten, M. J. Duff, A. Marrani, S. Nagy and M. Zoccali, *The Mile High Magic Pyramid*, [1711.08476](#).
- [80] A. Anastasiou, L. Borsten, M. J. Duff, S. Nagy and M. Zoccali, *Gravity as Gauge Theory Squared: A Ghost Story*, *Phys. Rev. Lett.* **121** (2018) 211601 [[1807.02486](#)].
- [81] G. Lopes Cardoso, G. Inverso, S. Nagy and S. Nampuri, *Comments on the double copy construction for gravitational theories*, *PoS CORFU2017* (2018) 177 [[1803.07670](#)].
- [82] W. D. Goldberger, S. G. Prabhu and J. O. Thompson, *Classical gluon and graviton radiation from the bi-adjoint scalar double copy*, *Phys. Rev.* **D96** (2017) 065009 [[1705.09263](#)].
- [83] W. D. Goldberger and A. K. Ridgway, *Bound states and the classical double copy*, *Phys. Rev.* **D97** (2018) 085019 [[1711.09493](#)].
- [84] W. D. Goldberger, J. Li and S. G. Prabhu, *Spinning particles, axion radiation, and the classical double copy*, *Phys. Rev.* **D97** (2018) 105018 [[1712.09250](#)].
- [85] A. Luna, R. Monteiro, I. Nicholson, A. Ochirov, D. O’Connell, N. Westerberg et al., *Perturbative spacetimes from Yang-Mills theory*, *JHEP* **04** (2017) 069 [[1611.07508](#)].
- [86] A. Luna, I. Nicholson, D. O’Connell and C. D. White, *Inelastic Black Hole Scattering from Charged Scalar Amplitudes*, *JHEP* **03** (2018) 044 [[1711.03901](#)].
- [87] C.-H. Shen, *Gravitational Radiation from Color-Kinematics Duality*, *JHEP* **11** (2018) 162 [[1806.07388](#)].
- [88] M. Levi, *Effective Field Theories of Post-Newtonian Gravity: A comprehensive review*, [1807.01699](#).
- [89] J. Plefka, J. Steinhoff and W. Wormsbecher, *Effective action of dilaton gravity as the classical double copy of Yang-Mills theory*, *Phys. Rev.* **D99** (2019) 024021 [[1807.09859](#)].
- [90] C. Cheung, I. Z. Rothstein and M. P. Solon, *From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion*, *Phys. Rev. Lett.* **121** (2018) 251101 [[1808.02489](#)].
- [91] M. Carrillo González, R. Penco and M. Trodden, *Radiation of scalar modes and the classical double copy*, *JHEP* **11** (2018) 065 [[1809.04611](#)].
- [92] R. Monteiro, I. Nicholson and D. O’Connell, *Spinor-helicity and the algebraic classification of higher-dimensional spacetimes*, *Class. Quant. Grav.* **36** (2019) 065006 [[1809.03906](#)].

- [93] J. Plefka, C. Shi, J. Steinhoff and T. Wang, *Breakdown of the classical double copy for the effective action of dilaton-gravity at NNLO*, *Phys. Rev.* **D100** (2019) 086006 [[1906.05875](#)].
- [94] B. Maybee, D. O’Connell and J. Vines, *Observables and amplitudes for spinning particles and black holes*, [1906.09260](#).
- [95] H. Johansson and A. Ochirov, *Double copy for massive quantum particles with spin*, *JHEP* **09** (2019) 040 [[1906.12292](#)].
- [96] A. P. V. and A. Manu, *Classical double copy from Color Kinematics duality: A proof in the soft limit*, *Phys. Rev.* **D101** (2020) 046014 [[1907.10021](#)].
- [97] M. Carrillo González, R. Penco and M. Trodden, *Shift symmetries, soft limits, and the double copy beyond leading order*, [1908.07531](#).
- [98] Y. F. Bautista and A. Guevara, *On the Double Copy for Spinning Matter*, [1908.11349](#).
- [99] I. Bah, R. Dempsey and P. Weck, *Kerr-Schild Double Copy and Complex Worldlines*, *JHEP* **02** (2020) 180 [[1910.04197](#)].
- [100] M. Carrillo González, B. Melcher, K. Ratliff, S. Watson and C. D. White, *The classical double copy in three spacetime dimensions*, *JHEP* **07** (2019) 167 [[1904.11001](#)].
- [101] W. D. Goldberger and J. Li, *Strings, extended objects, and the classical double copy*, *JHEP* **02** (2020) 092 [[1912.01650](#)].
- [102] K. Kim, K. Lee, R. Monteiro, I. Nicholson and D. Peinador Veiga, *The Classical Double Copy of a Point Charge*, *JHEP* **02** (2020) 046 [[1912.02177](#)].
- [103] A. Banerjee, E. Colgáin, J. A. Rosabal and H. Yavartanoo, *Ehlers as EM duality in the double copy*, [1912.02597](#).
- [104] A. Luna, S. Nagy and C. White, *The convolutional double copy: a case study with a point*, [2004.11254](#).
- [105] L. Alfonsi, C. D. White and S. Wikeley, *Topology and Wilson lines: global aspects of the double copy*, [2004.07181](#).
- [106] N. Bahjat-Abbas, R. Stark-Muchão and C. D. White, *Monopoles, shockwaves and the classical double copy*, *JHEP* **04** (2020) 102 [[2001.09918](#)].
- [107] A. Momeni, J. Rumbutis and A. J. Tolley, *Massive Gravity from Double Copy*, [2004.07853](#).
- [108] L. A. Johnson, C. R. T. Jones and S. Paranjape, *Constraints on a Massive Double-Copy and Applications to Massive Gravity*, [2004.12948](#).
- [109] S. Deser and A. Waldron, *(Dis)continuities of massless limits in spin 3/2 mediated interactions and cosmological supergravity*, *Phys. Lett.* **B501** (2001) 134 [[hep-th/0012014](#)].

- [110] S. Deser, J. H. Kay and K. S. Stelle, *Hamiltonian Formulation of Supergravity*, *Phys. Rev.* **D16** (1977) 2448.
- [111] S. Deser and B. Zumino, *Broken Supersymmetry and Supergravity*, *Phys. Rev. Lett.* **38** (1977) 1433.
- [112] E. Cremmer, B. Julia, J. Scherk, P. van Nieuwenhuizen, S. Ferrara and L. Girardello, *Super-higgs effect in supergravity with general scalar interactions*, *Phys. Lett.* **79B** (1978) 231.
- [113] E. Cremmer, P. Fayet and L. Girardello, *Gravity Induced Supersymmetry Breaking and Low-Energy Mass Spectrum*, *Phys. Lett.* **122B** (1983) 41.
- [114] R. Casalbuoni, S. De Curtis, D. Dominici, F. Feruglio and R. Gatto, *A GRAVITINO - GOLDSTINO HIGH-ENERGY EQUIVALENCE THEOREM*, *Phys. Lett.* **B215** (1988) 313.