

# Barbero-Immirzi Value from Experiment

Leonid Perlov,  
Department of Physics, University of Massachusetts, Boston, USA  
leonid.perlov@umb.edu

July 13, 2020

## Abstract

We consider GR as a limit case of the Scalar-Tensor theory with Barbero-Immirzi field when the field tends to a constant. We use Shapiro time delay experimental limit of  $1/w = (2.1 \pm 2.3)10^{-5}$  provided by the Cassini spacecraft to find the Barbero-Immirzi parameter value.

## 1 Barbero-Immirzi Value from Experimental Data

This is a short note on obtaining Barbero-Immirzi value from the experimental data by using the PPN (Parametric Post Newtonian) framework. The Barbero-Immirzi parameter appears in the Holst action as a coefficient of the topological term that vanishes due to the Bianchi identity.

$$S = \frac{1}{16\pi G} \int d^4x \, ee_I^\mu e_J^\nu (R^{IJ}_{\mu\nu} - \frac{\beta}{2} \epsilon^{IJ}_{KL} R^{KL}_{\mu\nu}) \quad (1)$$

Thus, it is impossible to obtain its value from the classical GR theory. In this paper we consider GR as a limit of the Scalar-Tensor theory with Barbero-Immirzi field and non-zero torsion, when the field tends to a constant. The Barbero-Immirzi field theory was introduced and studied in [1], and further investigated in [2]. Its action [1] is as follows:

$$S = \frac{1}{16\pi G} \int \sqrt{-g} \, d^4x \left[ R - \frac{3}{2} \left( \frac{1}{1 + \beta^2} \right) \partial_\mu \beta \partial^\mu \beta \right] \quad (2)$$

$R$  - a torsion-free Riemann scalar.

We can see from this formula that in the limit  $\beta = \text{Constant}$  this Scalar-Tensor theory becomes GR with Hilbert-Palatini action. The Scalar-Tensor theories are rare types of theories besides GR that are still in agreement with the experimental data [3], [6]. Moreover, in the limit, when a Scalar-Tensor theory PPN parameter  $w(\phi)$  tends to infinity, the theory becomes pure GR. The experimental data provided by Cassini in 2003 [5] put the following bound on  $w(\phi) > 10^5 / (2.1 \pm 2.3)$ . The more recent (2012) change in orbital period caused by dipolar gravitational radiation of the pulsar-white dwarf binary PSR J1738+0333 [4] improved this bound by the factor of 2:  $w(\phi) >$

$10^5/(2(2.1 \pm 2.3))$ , however only for non Brans-Dicke ( $w$  is not constant) theories. Therefore, in this paper we will use only Cassini's limit. It's worth mentioning that back in 1972 the experimental bound was much lower [6]:  $w(\phi) > 6$ , and it seems that the experimental data will keep pushing it further up to become GR. The bound limit we use in this paper comes from Shapiro time delay data provided in 2003 by the Cassini spacecraft on its way to Saturn [5], [3] with the result  $\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$ , where  $w(\phi) = 1 + \frac{\gamma}{\gamma-1} = 10^5/(2.1 \pm 2.3)$ . For the graph showing various available experimental data we refer to [4] Figure 7. The bounds provided by the most recent binary pulsars are not competitive with the solar-system Shapiro time delay measurement provided by Cassini due to the near equality of the star masses suppressing dipole radiation [3].

The PPN Scalar-Tensor theory action written in "Einstein frame" contains a scalar field  $\phi(x)$  and a coupling function  $w(\phi)$  [3]:

$$S = \frac{1}{16\pi G} \int \sqrt{-g} d^4x \left[ R - \frac{3 + 2w(\phi)}{2\phi^2} \partial_\mu \phi \partial^\mu \phi \right] \quad (3)$$

By comparing (2) with (3) we see that (2) is a particular case of (3) when the scalar field  $\phi(x)$  is a Barbero-Immirzi field  $\beta(x)$ :  $\phi(x) = \beta(x)$ . By equating the second terms coefficients in (2) and (3) we obtain:

$$\frac{3}{2(1 + \beta^2)} = \frac{3 + 2w(\beta)}{2\beta^2} \quad (4)$$

solving for  $\beta$ :

$$\beta^2 = -1 - \frac{3}{2w} \quad (5)$$

By using Cassini spacecraft bound:  $1/w(\beta) = (2.1 \pm 2.3)10^{-5}$  we obtain:

$$\beta^2 = -1 - \frac{3}{2}(2.1 \pm 2.3)10^{-5} \quad (6)$$

or

$$\beta = \pm i(1 \pm 0.008) \quad (7)$$

It is expected that the experimental  $w(\beta)$  bound will keep growing as it has been doing since 1960, starting from the value 6, reaching hundreds at the end of 1970th, and becoming Cassini's  $10^5/(2.1 \pm 2.3)$  today. The ongoing BepiColombo mission to Mercury, launched in 2018, will improve factor of 4 the Cassini's result in measuring PPN  $\gamma$  in July 2022, when it will experience the first solar conjunction [7].

To conclude there are two separate limits depending on the same parameter  $w(\beta)$  measured from experiment. The bigger  $w(\beta)$  the closer this Scalar-Tensor theory is to GR. At the same time the bigger  $w(\beta)$  the closer Barbero-Immirzi value is to  $\pm i$ , pushing it to the self-dual Ashtekar case. Today its value is  $\beta = \pm i(1 \pm 0.008)$ .

### Acknowledgment

I am very grateful to Michael Bukatin for reviewing this note and for his supportive enthusiastic spirit.

## References

- [1] G Calcagni, S Mercuri, "Barbero–Immirzi field in canonical formalism of pure gravity", *Phys. Rev. D* 79, 084004 (2009)
- [2] F. Bombacigno, G. Montani, "Implications of the Holst term in a  $f(R)$  theory with torsion", *Phys. Rev. D* 99, 064016 (2019)
- [3] C. Will "Theory and experiment in gravitational physics", Cambridge University Press, Cambridge, (2018)
- [4] P.C.C. Freire, N. Wex, G. Esposito-Farese, J.P.W. Verbiest, M. Bailes, B. A. Jacoby, M. Kramer, I. H. Stairs, J. Antoniadis, G.H. Janssen, "The relativistic pulsar–white dwarf binary PSR J1738+0333 – II. The most stringent test of scalar–tensor gravity", *Monthly Notices of the Royal Astronomical Society*, V 423, Issue 4, (2012)
- [5] B. Bertotti, P Tortora, "A test of general relativity using radio links with the Cassini spacecraft", *Nature* volume 425, pages 374–376(2003)
- [6] C. Will "Theoretical tools of experimental gravitation", Cambridge University Press, Cambridge, (1972)
- [7] L. Imperi, Lucano Iass, "The determination of the post-Newtonian parameter  $\gamma$  during the cruise phase of BepiColombo", *Class Quant. Grav.* Volume 34, Number 7, (2017)
- [8] J. F. Barbero, "Real Ashtekar variables for Lorentzian signature space-times", *Phys. Rev. D* 51, 5507 (1995)