

Revisiting the Black Hole Thermodynamics in Horndeski Gravity

F. F. Santos^{1*}

¹*Departamento de Física, Universidade Federal da Paraíba,
Caixa Postal 5008, 58051-970, João Pessoa, Paraíba, Brazil.*

In this article, we investigate the thermodynamics of an asymptotically AdS black hole in Horndeski's gravity. In our investigation, we considered the black hole solutions asymptotically AdS in (3+1)-dimensions, which is a solution of Horndeski's theory. In the canonical Ensemble, the relationship between Horndeski's parameters provides results that are in accordance with Hawking-Page's description of thermodynamics of black holes in asymptotically Anti-de Sitter space.

arXiv:2005.14154v2 [hep-th] 23 Jun 2020

*Electronic address: fabiano.ffa23@gmail.com

I. INTRODUCTION

Since its discovery in 1974, Horndeski's gravity has aroused great interest ranging from applications in the field of cosmology to black holes [1–4]. However, a case of particular interest is the study of static black hole solutions [5, 6] with asymptotically anti-de Sitter behavior. This Anti-de Sitter (AdS) spacetime played an extremely important role in the holographic duality known as AdS/CFT [7, 8]. In recent years, the gauge/gravity duality has offered remarkable support for the dynamics of systems strongly coupled to condensed matter systems [9, 10] in the study of the universal limits of transport coefficients, including viscosity/entropy radius [11, 12], which is conjectured based on the "bottom up" models. Some examples of violation of this limit were presented by [13–15], however, in [13, 14] considerations of the AdS/CFT correspondence were presented in the context of Horndeski's gravity.

However, the recent discovery of event GW170817 of the merger of the two neutron stars, has imposed strong restrictions on sectors of Horndeski's gravity [16] one of these sectors is the Jhon sector of Horndeski's gravity, this sector derives from a special subclass known as theories F4 [17, 18]. Horndeski's gravity [1–4], is characterized by a single theory of scalar tensors with second order field equations and second order energy moment tensor [1, 2, 5, 19–21]. The Lagrangians associated with this theory produce second order equations of motion [20, 22–24]. In addition, the theory includes four arbitrary functions of the scalar field and its kinetic term [17, 18, 25–27]. Within the scope of the construction of black holes this theory has shown that it is possible to work on the development of Hawking-Page phase transitions at a critical temperature [20, 21, 28]. However, in the context of high energy particle collisions, we have the creation of tiny black holes for the primitive Universe, this can leave stable remains instead of completely evaporating as a result of the Hawking radiation [29]. In addition, at scales comparable to quantum gravity near TeV, the LHC can produce a black hole every second. Black holes are well understood objects, when their mass exceeds Planck's fundamental mass $M_p \sim \text{TeV}$. In this case, if the mass of the black hole approaches that of Planck ($M_{BH} \approx M_p$), we have that black holes become "sticky" [30] and its properties end up becoming complex in this regime. This obstacle can be ignored by taking semi-classical arguments, that is, when we are in the regime where $M_{BH} \gg M_p$.

Despite the wide variety of studies involving black holes [4, 20, 21, 28] Hawking provided

that black holes for an AdS geometry have a minimum temperature, which occurs when its size is in the order of the characteristic radius of the AdS space [31]. On the other hand, for larger black holes their deviation from temperature measured at infinity to red is large. This fact showed that these black holes have a specific positive heat and can be in stable equilibrium with thermal radiation at a fixed temperature. In addition, another implication is that the canonical Ensemble exists for the space asymptotically AdS [9, 31, 32].

Recent investigations regarding black hole thermodynamics have been matched in [20, 33] where studies involving a gravitational soliton were considered as a background to investigate finite Euclidean action. In addition, in [33] thermodynamics for BTZ-like black holes was discussed by both the Euclidean and Cardy methods. On the other hand, we can explore the thermodynamics of black holes through a very elegant and robust argument, in which we have that the horizons correspond to thermally mixed states [9, 34]. However, for a semi-classical regime, the volume theory partition function (accompanied by asymptotic boundary conditions of the AdS, must provide an equivalence with the large field theory partition function) as an integral path over the metrics. As addressed by [16] one of the reasons that put many corrected cosmological models to the test is the fact that the vast majority of them produce massive gravitons primarily. One of these models that we have already commented on is the model involving the Jhon sector term of Horndeski's gravity. In this way, so that we can imitate the field theory calculations, looking for euclidean saddle points in bulk of the theory, we have the integral of the path, the action must contain a Gibbons-Hawking contour term for the Jhon sector of Horndeski's gravity to provide the correct variational problem (Dirichlet) and also a boundary counter-term for having the condition of finite action, this term has already been presented by [35].

In our studies we will analyze the thermodynamics of a black hole in asymptotically Anti-de Sitter space in Horndeski Gravity. The point $\gamma = -\alpha/\Lambda$, provide us with new features [36], which will be useful for probing the contour conditions of Dirichlet for the scalar field in the contour. Thus, we can analyze the thermodynamics of black holes, which is provided by the partition function. This work is summarized as follows: In Section II, we present Horndeski's Theory. In Section III, we address the issue of finding the ansatz black hole. In Section IV, we explore the canonical set and analyze thermodynamic quantities. Finally, in the section V, we present our conclusions.

II. HORNDESKI THEORY

In this session, we will match the Jhon sector of Horndeski's gravity [17, 18, 27]. This model is expressed by the following action

$$S[g_{\mu\nu}, \phi] = \int d^4x \sqrt{-g} \mathcal{L} + S_m. \quad (1)$$

$$\mathcal{L} = \kappa(R - 2\Lambda) - \frac{1}{2}(\alpha g_{\mu\nu} - \gamma G_{\mu\nu}) \nabla^\mu \phi \nabla^\nu \phi \quad (2)$$

Where we define a new field $\phi' \equiv \psi$ and $\kappa = (16\pi G)^{-1}$. S_m is used to describe a perfect fluid. In this action, the field has a dimension of $(mass)^2$ and the parameters α and γ control the strength of the kinetic couplings, α is dimensionless and γ has a dimension of $(mass)^{-2}$. Another characteristic of the term $\nabla^\mu \phi \nabla^\nu \phi$ is that it is a second-order covariant derived from the scalar field. Furthermore, action is invariable under the displacement symmetry $\phi \rightarrow \phi + constant$ and under the discrete transformation $\phi \rightarrow -\phi$. Thus, the Einstein-Horndeski field equations can be written formally by varying the action (12) $\delta S[g_{\mu\nu}, \phi]$ providing

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{1}{2\kappa} T_{\mu\nu}, \quad (3)$$

where $T_{\mu\nu} = \alpha T_{\mu\nu}^{(1)} + \gamma T_{\mu\nu}^{(2)}$ the energy-momentum tensors $T_{\mu\nu}^{(1)}$ and $T_{\mu\nu}^{(2)}$ take the following form

$$\begin{aligned} T_{\mu\nu}^{(1)} &= \nabla_\mu \phi \nabla_\nu \phi - \frac{1}{2} g_{\mu\nu} \nabla_\lambda \phi \nabla^\lambda \phi \\ T_{\mu\nu}^{(2)} &= \frac{1}{2} \nabla_\mu \phi \nabla_\nu \phi R - 2 \nabla_\lambda \phi \nabla_{(\mu} \phi R_{\nu)}^\lambda - \nabla^\lambda \phi \nabla^\rho \phi R_{\mu\lambda\nu\rho} \\ &\quad - (\nabla_\mu \nabla^\lambda \phi)(\nabla_\nu \nabla_\lambda \phi) + (\nabla_\mu \nabla_\nu \phi) \square \phi + \frac{1}{2} G_{\mu\nu} (\nabla \phi)^2 \\ &\quad - g_{\mu\nu} \left[-\frac{1}{2} (\nabla^\lambda \nabla^\rho \phi)(\nabla_\lambda \nabla_\rho \phi) + \frac{1}{2} (\square \phi)^2 - (\nabla_\lambda \phi \nabla_\rho \phi) R^{\lambda\rho} \right]. \end{aligned}$$

And the scalar field equation is given by

$$\nabla_\mu [(\alpha g^{\mu\nu} - \gamma G^{\mu\nu}) \nabla_\nu \phi] = 0. \quad (4)$$

III. BLACK HOLE SOLUTIONS

In our case, for Einstein-Horndeski gravity, we consider the following *Ansatz* for a four-dimensional black hole of the form

$$ds^2 = \frac{L^2}{r^2} \left(-f(r)dt^2 + dx^2 + dy^2 + \frac{dr^2}{f(r)} \right). \quad (5)$$

The spherically symmetrical static configurations of certain Galileons have an invariance of change, these were first proposed to admit a no-hair theorem [14, 19]. In addition, the no-hair theorem for Galileons requires that the square of the conserved current $J^\mu = (\alpha g^{\mu\nu} - \gamma G^{\mu\nu})\nabla_\nu\phi$, defined in (4), should not diverge on the horizon. However, we can escape the no-hair theorem [19, 21], for this we need to impose that the conserved current's radial component disappears in an identical way without restricting the radial dependence of the scalar field [19]:

$$\alpha g_{rr} - \gamma G_{rr} = 0. \quad (6)$$

As $\phi'(r) \equiv \psi(r)$, we can easily see that this condition wipes out $\psi^2(r)$, regardless of its behavior on the horizon. The metric function $f(r)$ can be found using the equation (6). It can be shown that the equation (4) is satisfied by the following solution

$$f(r) = \frac{\alpha L^2}{3\gamma} - \left(\frac{r}{r_+} \right)^3, \quad (7)$$

$$\psi^2(r) = -\frac{2L^2\kappa(\alpha + \gamma\Lambda)}{\alpha\gamma r^2} \frac{1}{f(r)}. \quad (8)$$

The Einstein-Horndeski field equations (3) and (4) are satisfied by the equations (7) and (8). The scalar field given by the equation (8) is real for $\alpha > 0$ and $\gamma < 0$. As discussed by [20], $\alpha/(3\gamma) = L^{-2}$ is defined as an effective radius of the AdS L [20] where the solutions can be asymptotically dS or AdS for $\alpha/\gamma < 0$ or $\alpha/\gamma > 0$, respectively. We can notice that the equation (7) with $\alpha/(3\gamma) = L^{-2}$, we have that for $f \rightarrow 1$ with $r \rightarrow 0$ it is a spacetime asymptotically AdS. Thus, the further we advance in spacetime, entering the infrared (IR) region for large values of r , we find a horizon in $r = r_+$ and thus both $g_{tt}(r)$ and $\psi^2(r)$, the latter in action (12) ensures that this is a genuine vacuum solution. Furthermore, we have that the surface located at $r = r_+$ is infinitely shifted to red in relation to an asymptotic

observer. On the other hand, we can see by looking at the equation (8), which $(\alpha + \gamma\Lambda) > 0$, indicates ghost freedom. Thus, stability requires that $(\alpha + \gamma\Lambda)$ is not negative, this leads to an interval of the form $-\infty < \gamma \leq \alpha/(-\Lambda)$ [36].

In fact, we have that the appearance of a black hole, which has a flat horizon \mathfrak{R}^2 , leads us to a physics in IR that corresponds to placing the invariant theory of scale at a finite temperature [9]. So, we have that the temperature of the black hole is given by

$$T(r_+) = \frac{3}{4\pi r_+} \quad (9)$$

The minimum temperature value in $r_c = L/\sqrt{3}$, can be written as

$$T_c = \frac{3\sqrt{3}}{4\pi L} = \frac{1}{4\pi} \sqrt{\frac{\alpha}{\gamma}} \quad (10)$$

This minimum occurs when its size is in the order of the characteristic radius of the anti-de Sitter space, which at $\gamma = -\alpha/\Lambda$ [31].

IV. THE CANONICAL ENSEMBLE

However, for geometry that represents a state of finite density, free thermodynamic energy can be obtained by evaluating Euclidean action as discussed by [9]. In fact, as already well established, the path integral must include a Gibbons-Hawking boundary term to provide the correct variational problem (Dirichlet) and a constant boundary counter-term to make the action finite. Thus, for the action (12), we have that the terms of Gibbons-Hawking and counter-term are expressed by S_{bdry} and S_{ct} , which are included in the total action given by the following equation

$$\begin{aligned} S &= S_N + S_{bdry} + S_{bdry}^{mat} + S_{ct} \\ &= \int_N d^4x \sqrt{-g} \mathcal{L}_H + 2\kappa \int_{bdry} d^3x \sqrt{-h} \mathcal{L}_{bdry} + 2\kappa \int_Q d^3x \mathcal{L}_{mat} + S_{ct} \end{aligned} \quad (11)$$

$$\mathcal{L}_H = \kappa(R - 2\Lambda) - \frac{1}{2}(\alpha g_{\mu\nu} - \gamma G_{\mu\nu}) \nabla^\mu \phi \nabla^\nu \phi \quad (12)$$

$$\mathcal{L}_{bdry} = K + \frac{\gamma}{4} (\nabla_\mu \phi \nabla_\nu \phi n^\mu n^\nu - (\nabla \phi)^2) K + \frac{\gamma}{4} \nabla_\mu \phi \nabla_\nu \phi K^{\mu\nu} \quad (13)$$

$$\mathcal{L}_{ct} = c_0 + c_1 R + c_2 R^{ij} R_{ij} + c_3 R^2 + b_1 (\partial_i \phi \partial^i \phi)^2 + \dots \quad (14)$$

Where \mathcal{L}_{mat} is a Lagrangian of possible fields of matter at the border and \mathcal{L}_{bdry} corresponds to the terms with dependency on γ for the Horndeski gravity described for \mathcal{L}_H [35]. In addition, we have that $K_{\mu\nu} = h_{\mu}^{\beta}\nabla_{\beta}n_{\nu}$ is the extrinsic curvature and its line is given by the contraction $K = h^{\mu\nu}K_{\mu\nu}$ and h is the metric induced at the border. In addition, S_{ct} is the necessary contour counter-term for the asymptotic AdS spacetime. Furthermore, the scalar field must fall fast enough near the limit (that is, it reaches zero or a constant), close to the boundary [9]. However, close to the $\psi(r \rightarrow 0) \rightarrow \infty$ boundary, but one way for the scalar field to fall fast enough is at the point $\gamma = -\alpha/\Lambda$, providing $\phi(r \rightarrow 0) = constant$. Thus, we have that the action (11) is reduced to form

$$S_E = - \int d^4x \sqrt{g} [\kappa(R - 2\Lambda)] - 2\kappa \int_{r \rightarrow 0} d^3x \sqrt{h} \left[K - \frac{2}{L} \right] \quad (15)$$

We can work with the Canonical Ensemble where the free energy is given by $F = -T \ln(Z)$, Z is the partition function defined by $Z = e^{-S_E}$ e S_E is the Euclidean action. Through the relationship $\ln(Z) = -S_E$, we have $F = TS_E$ [9], however, using the equations (9,10), we can write the following thermodynamic relationship:

$$\ln(Z) = \sigma \frac{T^2}{T_c^2}, \quad (16)$$

$$F = -\sigma \frac{T^3}{T_c^2} \quad (17)$$

Where $\sigma \equiv \frac{3V_2}{4G}$ and $M_p = G^{-1/2}$ is the mass of Planck in units in which $\hbar = c = 1$ [31]. Now, we can construct the thermal states in the anti-de Sitter space, periodically identifying the coordinate of the imaginary time τ with the period $\beta = 1/T$. Thus, we can find the expected value of energy, entropy and specific heat as

$$\langle E \rangle = - \frac{\partial \ln(Z)}{\partial \beta}, \quad (18)$$

$$S = - \frac{\partial F}{\partial T}, \quad (19)$$

$$c = T \frac{\partial S}{\partial T}. \quad (20)$$




FIG. 1: The figure shows the behavior of the equations (17), (18), (19) and (20), respectively, where $\langle E \rangle > 0$ (top left) blue line increases to $T > T_c$ and the red line decreases to $T < T_c$. The positive specific heat $c > 0$ blue line increases to $T > T_c$ and the red line decreases to $T < T_c$ (upper right). Entropy in the positive region $S > 0$ increases to $T > T_c$ and on the red line it decreases to $T < T_c$ (bottom-left). The fact that $c > 0$ implies a higher value of the black hole mass, gives us that it is at least locally stable. The (bottom left) shows the free energy $F < 0$ blue line for $T > T_c$ and red line $F \rightarrow 0$ for $T < T_c$ both cases indicate that the system has overall stability.




FIG. 2: The free energy for large black holes (blue line) and small (red line) for the values of $\gamma > 0$ and $\gamma < 0$, respectively.

According to figure 1, we have that the values of $\langle E \rangle > 0$ and $c > 0$ are in according to [31] indicating a higher mass value of the black hole, which is at least locally stable. In addition, the free energy $F < 0$ indicates that the system has global stability [28], but for that free energy value it provides $c > 0$. In fact, large black holes (LBH) have negative free energy. However, we can see that $T > T_c$ is an adequate approximation. This approximation can be related to the fact that $\alpha \sim 0$, that small value of α , these considerations in the equation (10) provide that T_c is very small, giving us enough support to consider $T > T_c$. In addition, $T < T_c$ occurs when $\gamma \sim 0$, as long as T_c is very large, providing $T < T_c$. For $\gamma < 0$ in the equation (10), the free energy becomes positive, which characterizes a small black hole (SBH). The figure 2 shows the behavior of free energy for $\gamma > 0$ and $\gamma < 0$, which is a phase transition in this scenario, given by the small black hole and large black hole (SBH -LBH) [41]. We can see that $\langle E \rangle$ and S satisfy the thermodynamic relationship $F = \langle E \rangle - TS$.

V. CONCLUSION

In this article, we show that an asymptotically AdS black hole is a solution to Horndeski's gravity. An interesting fact was that at the point $\gamma = -\alpha/\Lambda$, we explored the canonical Ensemble, where free energy and other thermodynamic quantities are calculated in terms of the temperature T/T_c . The free energy of the black hole, its entropy and internal energy satisfy the thermodynamic relationship of $F = \langle E \rangle - TS$ where S obeys the famous law of Hawking [13, 32, 37–40]. Thus, together with specific heat, it plays an important role in the investigation of thermal stability. As the free energy is negative $F < 0$ we have global stability, as shown in the figure 1 with $c > 0$, showing that the system has local stability [31]. However, for $\gamma < 0$, free energy implies local stability. In the cases $\gamma > 0$ and $\gamma < 0$ figure 2, we have a phase transition between the small black hole and the large black hole (SBH-LBH) [28, 41], which also implies a one-phase transition between the dS and the AdS space, which is in accordance with $\alpha/\gamma < 0$ and $\alpha/\gamma > 0$, respectively, where the dS case suggests the creation of tiny black holes at the beginning of the Universe, which can leave stable remains instead of completely evaporating as a result of Hawking radiation [29], for a phase with large black holes.

However, as the energy $\langle E \rangle$ increases to $\alpha \sim 0$, as shown in the figure 1, this means that the black hole "captured" a new mass energy (matter or radiation goes beyond the surface

surrounding the black hole, which we call the horizon), where the sector of matter is given only by the kinetic term associated with the scalar field. On the other hand, if the energy $\langle E \rangle \sim 0$ decreases to $\gamma \sim 0$, see figure 1, it suggests that the only mechanism proposed to decrease the mass of a black hole is the evaporation of Hawking radiation, which is a process of quantum origin and arises from the creation of pairs of particles from the vacuum energy in the horizon region.

Acknowledgments

We would like to thank CNPq and CAPES for partial financial support. I would also like to thank Mokhtar Hassine for fruitful discussions at the end of this work.

-
- [1] L. Heisenberg, *A systematic approach to generalisations of General Relativity and their cosmological implications*, Phys. Rept. **796**, 1 (2019), [arXiv:1807.01725 [gr-qc]].
 - [2] G. W. Horndeski, *Second-order scalar-tensor field equations in a four-dimensional space*, Int. J. Theor. Phys. **10**, 363 (1974).
 - [3] J. P. Bruneton, M. Rinaldi, A. Kanfon, A. Hees, S. Schlogel and A. Fuzfa, *Fab Four: When John and George play gravitation and cosmology*, Adv. Astron. **2012**, 430694 (2012), [arXiv:1203.4446 [gr-qc]].
 - [4] F. A. Brito and F. F. Dos Santos, *Braneworlds in Horndeski gravity*, arXiv:1810.08196 [hep-th].
 - [5] A. Cisterna and C. Erices, *Asymptotically locally AdS and flat black holes in the presence of an electric field in the Horndeski scenario*, Phys. Rev. D **89**, 084038 (2014), [arXiv:1401.4479 [gr-qc]].
 - [6] A. Cisterna, T. Delsate, L. Ducobu and M. Rinaldi, *Slowly rotating neutron stars in the nonminimal derivative coupling sector of Horndeski gravity*, Phys. Rev. D **93**, no. 8, 084046 (2016), [arXiv:1602.06939 [gr-qc]].
 - [7] G. W. Gibbons, *Anti-de-Sitter spacetime and its uses*, arXiv:1110.1206 [hep-th].
 - [8] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, Int. J. Theor. Phys. **38**, 1113 (1999) [Adv. Theor. Math. Phys. **2**, 231 (1998)], [hep-th/9711200].

- [9] S. A. Hartnoll, *Lectures on holographic methods for condensed matter physics*, *Class. Quant. Grav.* **26**, 224002 (2009), [arXiv:0903.3246 [hep-th]].
- [10] S. Sachdev, *What can gauge-gravity duality teach us about condensed matter physics?*, *Ann. Rev. Condensed Matter Phys.* **3**, 9 (2012), [arXiv:1108.1197 [cond-mat.str-el]].
- [11] P. Kovtun, D. T. Son and A. O. Starinets, *Holography and hydrodynamics: Diffusion on stretched horizons*, *JHEP* **0310**, 064 (2003), [hep-th/0309213].
- [12] P. Kovtun, D. T. Son and A. O. Starinets, *Viscosity in strongly interacting quantum field theories from black hole physics*, *Phys. Rev. Lett.* **94**, 111601 (2005), [hep-th/0405231].
- [13] X. H. Feng, H. S. Liu, H. Lü and C. N. Pope, *Black Hole Entropy and Viscosity Bound in Horndeski Gravity*, *JHEP* **1511**, 176 (2015), [arXiv:1509.07142 [hep-th]].
- [14] F. A. Brito and F. F. Santos, *Black branes in asymptotically Lifshitz spacetime and viscosity/entropy ratios in Horndeski gravity*, arXiv:1901.06770 [hep-th].
- [15] M. Sadeghi, *Black Brane Solution in Rastall AdS Massive Gravity and Viscosity Bound*, *Mod. Phys. Lett. A* **33**, no. 37, 1850220 (2018), [arXiv:1809.08698 [hep-th]].
- [16] V. Oikonomou and F. Fronimos, *Reviving non-Minimal Horndeski-like Theories after GW170817: Kinetic Coupling Corrected Einstein-Gauss-Bonnet Inflation*, [arXiv:2006.05512 [gr-qc]].
- [17] C. Charmousis, E. J. Copeland, A. Padilla and P. M. Saffin, *General second order scalar-tensor theory, self tuning, and the Fab Four*, *Phys. Rev. Lett.* **108**, 051101 (2012), [arXiv:1106.2000 [hep-th]].
- [18] C. Charmousis, E. J. Copeland, A. Padilla and P. M. Saffin, *Self-tuning and the derivation of a class of scalar-tensor theories*, *Phys. Rev. D* **85**, 104040 (2012), [arXiv:1112.4866 [hep-th]].
- [19] M. Bravo-Gaete and M. Hassaine, *Lifshitz black holes with a time-dependent scalar field in a Horndeski theory*, *Phys. Rev. D* **89**, 104028 (2014), [arXiv:1312.7736 [hep-th]].
- [20] A. Anabalón, A. Cisterna and J. Oliva, *Asymptotically locally AdS and flat black holes in Horndeski theory*, *Phys. Rev. D* **89**, 084050 (2014), [arXiv:1312.3597 [gr-qc]].
- [21] M. Rinaldi, *Black holes with non-minimal derivative coupling*, *Phys. Rev. D* **86**, 084048 (2012), [arXiv:1208.0103 [gr-qc]].
- [22] C. Deffayet, X. Gao, D. A. Steer and G. Zahariade, *From k-essence to generalised Galileons*, *Phys. Rev. D* **84**, 064039 (2011), [arXiv:1103.3260 [hep-th]].
- [23] M. Rinaldi, *Mimicking dark matter in Horndeski gravity*, *Phys. Dark Univ.* **16**, 14 (2017),

- [arXiv:1608.03839 [gr-qc]].
- [24] A. Cisterna, M. Hassaine, J. Oliva and M. Rinaldi, *Axionic black branes in the k-essence sector of the Horndeski model*, Phys. Rev. D **96**, no. 12, 124033 (2017), [arXiv:1708.07194 [hep-th]].
- [25] J. Gleyzes, D. Langlois, F. Piazza and F. Vernizzi, *Essential Building Blocks of Dark Energy*, JCAP **1308**, 025 (2013), [arXiv:1304.4840 [hep-th]].
- [26] M. Zumalacárregui and J. García-Bellido, *Transforming gravity: from derivative couplings to matter to second-order scalar-tensor theories beyond the Horndeski Lagrangian*, Phys. Rev. D **89**, 064046 (2014), [arXiv:1308.4685 [gr-qc]].
- [27] A. A. Starobinsky, S. V. Sushkov and M. S. Volkov, *The screening Horndeski cosmologies*, JCAP **1606**, no. 06, 007 (2016), [arXiv:1604.06085 [hep-th]].
- [28] M. Rostami, J. Sadeghi, S. Miraboutalebi, B. Pourhassan and A. A. Masoudi, *Phase transition of modified Horndeski gravity with new method*, arXiv:1909.01799 [gr-qc].
- [29] T. Nakama and J. Yokoyama, *Micro black holes formed in the early Universe and their cosmological implications*, Phys. Rev. D **99**, no. 6, 061303 (2019), [arXiv:1811.05049 [gr-qc]].
- [30] S. Dimopoulos and G. L. Landsberg, *Black holes at the LHC*, Phys. Rev. Lett. **87**, 161602 (2001), [hep-ph/0106295].
- [31] S. W. Hawking and D. N. Page, *Thermodynamics of Black Holes in anti-De Sitter Space*, Commun. Math. Phys. **87**, 577 (1983).
- [32] Y. C. Ong, *Hawking Evaporation Time Scale of Topological Black Holes in Anti-de Sitter Spacetime*, Nucl. Phys. B **903**, 387 (2016), [arXiv:1507.07845 [gr-qc]].
- [33] O. Baake, M. Bravo-Gaete and M. Hassaine, *Spinning black holes for generalized scalar tensor theories in three dimensions*, [arXiv:2005.10869 [hep-th]].
- [34] L. Susskind and J. Lindesay, *An introduction to black holes, information and the string theory revolution: The holographic universe*,
- [35] Y. Z. Li, H. Lu and H. Y. Zhang, *Scale Invariance vs. Conformal Invariance: Holographic Two-Point Functions in Horndeski Gravity*, Eur. Phys. J. C **79**, no. 7, 592 (2019), [arXiv:1812.05123 [hep-th]].
- [36] W. J. Jiang, H. S. Liu, H. Lu and C. N. Pope, *DC Conductivities with Momentum Dissipation in Horndeski Theories*, JHEP **1707**, 084 (2017), [arXiv:1703.00922 [hep-th]].
- [37] D. N. Page, *Finite upper bound for the Hawking decay time of an arbitrarily large black hole in anti-de Sitter spacetime*, Phys. Rev. D **97**, no. 2, 024004 (2018), [arXiv:1507.02682 [hep-th]].

- [38] H. Xu and M. H. Yung, *Black hole evaporation in Lovelock gravity with diverse dimensions*, Phys. Lett. B **794**, 77 (2019), [arXiv:1904.06503 [gr-qc]].
- [39] H. Xu and M. H. Yung, *Black hole evaporation in Conformal (Weyl) gravity*, Phys. Lett. B **793**, 97 (2019), [arXiv:1811.07309 [gr-qc]].
- [40] Y. C. Ong, *The Persistence of the Large Volumes in Black Holes*, Gen. Rel. Grav. **47**, no. 8, 88 (2015), [arXiv:1503.08245 [gr-qc]].
- [41] N. K. A., C. L. A. Rizwan, A. K. M. and M. S. Ali, *Photon Orbits and Thermodynamic Phase Transition of Regular AdS Black Holes*, arXiv:1912.11909 [gr-qc].