

Super-Penrose process for extremal rotating neutral white holes

O. B. Zaslavskii

Department of Physics and Technology,

Kharkov V.N. Karazin National University,

4 Svoboda Square, Kharkov 61022, Ukraine and

Institute of Mathematics and Mechanics, Kazan Federal University,

*18 Kremlyovskaya St., Kazan 420008, Russia**

We consider collision of two particles 1 and 2 near the horizon of the extremal rotating axially symmetric neutral generic black hole producing particles 3 and 4. We discuss the scenario in which both particles 3 and 4 fall into a black hole and move in a white hole region. If particle 1 is fine-tuned, the energy $E_{c.m.}$ in the centre of mass grows unbounded (the Bañados-Silk-West effect). Then, particle 3 can, in principle, reach a flat infinity in another universe. If not only $E_{c.m.}$ but also the corresponding Killing energy E is unbounded, this gives a so-called super-Penrose process (SPP). We show that the SPP is indeed possible. Thus white holes turn out to be potential sources of high energy fluxes that transfers from one universe to another. This generalizes recent observations made by Patil and Harada for the Kerr metric. We analyze two different regimes of the process on different scales.

PACS numbers: 04.70.Bw, 97.60.Lf

I. INTRODUCTION

During last decade a lot of work has been made for investigation of properties of high energy processes near black holes. This was stimulated by the paper [1], where it was found that collision of two particles near rapidly rotating black holes can lead to unbounded energies $E_{c.m.}$ in the centre of mass frame. This was called the Bañados-Silk-West (BSW) effect, after the authors' names. After its publication, it turned out, that there are also

*Electronic address: zaslav@ukr.net

earlier works [2], [3] in which near-horizon particle collisions in the Kerr metric were investigated. Meanwhile, a typical process considered there, includes head-on collision between two arbitrary particles, where particle 1 arrives from infinity while particle 2 comes from the horizon (see eq. 2.57 of [3]). But as far as particle 2 is concerned, this is nothing else than a typical behavior of a particle near a white hole. If such a region is allowed in the complete space-time, the effect of unbounded $E_{c.m.}$ for head-on collisions exists even in the Schwarzschild metric [4], [5]. Thus white holes can be an alternative to black ones as a source of high energy collisions.

More important question is whether it is possible to gain not only unbounded $E_{c.m.}$ but also unbounded conserved Killing energies E since it is the latter quantity which can be measured in the Earth laboratory, at least in principle. The collisions in the Schwarzschild background are useless for this purpose since energy cannot be extracted at all. For such an extraction, the existence of negative energies and ergoregion are required that makes it possible the Penrose process [6] or its collisional analogue [7]. If the energy gain is unbounded, this is called the super-Penrose process (SPP). For black holes, the energy gain is finite, so the SPP is impossible for them (see [8] and references therein).

In this context, there is a scenario with participation of white holes, different from those in [2] - [5]. Now, both particles collide near the black hole horizon in "our" part of Universe but afterwards the products of reaction leave it. Passing through the horizon, they appear inside a white whole region and, eventually, transfer energy to another universe. Or, vice versa, collision in another universe can give rise to high energy in our one. If such a process is possible, this would give astrophysical realization of high energy transfer with white holes as a source that was suggested earlier [9], [10].

The concrete process of this kind in the Kerr background was considered recently in [11]. The authors showed that the conserved energy of produced particles can be as large as one like. In other words, the SPP is possible. Our aim is to extend consideration to generic rotating axially symmetric stationary white holes. In doing so, we exploit the approach that, in our view, is simpler and was already used for examination of the energy extraction from generic black holes of the aforementioned type including the Kerr metric [12], [13]. In particular, we do not use transformation between the three frames (center of mass, locally non-rotating and stationary ones) and work in the original frame.

Although there are reasons to believe that white holes are unstable (see Sec. 15 of [14]),

motivation for consideration of such objects stems from different roots. (i) High energy process, if they are confirmed, can themselves contribute to the instability of white holes, so they are important for elucidation of the fate of such objects. (ii) The complete theory of the BSW effect should take into consideration all possible configurations and scenarios, at least for better understanding the phenomenon.

We use the system of units in which the fundamental constants $G = c = 1$.

II. BASIC EQUATIONS

Let us consider the metric

$$ds^2 = -N^2 dt^2 + \frac{dr^2}{A} + g_\phi(d\phi - \omega dt)^2 + g_\theta d\theta^2, \quad (1)$$

where $g_\phi \equiv g_{\phi\phi}$, $g_\theta \equiv g_{\theta\theta}$, all coefficients do not depend on t and ϕ . For a given energy E , angular momentum L and mass m the equation of motion in the equatorial plane read

$$m\dot{t} = \frac{X}{N^2}, \quad (2)$$

$$X = E - \omega L, \quad (3)$$

$$m\dot{\phi} = \frac{L}{g_\phi} \quad (4)$$

$$m\dot{r} = \sigma P, \quad \sigma = \pm 1, \quad (5)$$

$$P = \sqrt{X^2 - \tilde{m}^2 N^2}, \quad (6)$$

$$\tilde{m}^2 = m^2 + \frac{L^2}{g_\phi}, \quad (7)$$

dot denotes differentiation with respect to the proper time τ . To simplify formulas, we assume that in the equatorial plane $A = N^2$. Otherwise, we can always achieve this equality by redefining the radial coordinate. The forward-in-time condition requires

$$X \geq 0. \quad (8)$$

In what follows, we will use the standard classification of particles. If $X_H = 0$, a particle is called critical. If $X_H = O(1)$, it is called usual. If $X_H = O(N_c)$, it is called near-critical. Here, subscripts "H" and "c" refer to the quantities calculated on the horizon and the point of collision, respectively.

For the near-critical particle, we use presentation

$$L = \frac{E}{\omega_H}(1 + \delta) \quad (9)$$

exploited in [13]. Here,

$$\delta = C_1 N_c \quad (10)$$

is a small quantity for collisions near the horizon, $C_1 = O(1)$ is a constant.

Near the horizon, we assume the Taylor expansion that for the extremal case reads [15]

$$\omega = \omega_H - B_1 N + O(N^2). \quad (11)$$

Then, we have the following approximate expressions there.

The critical particle:

$$X = \frac{b}{h} E N + O(N^2), \quad (12)$$

$$P = N \sqrt{E^2 \left(\frac{b^2 - 1}{h^2} \right) - \frac{1}{h^2} - m^2}. \quad (13)$$

A usual particle:

$$X = X_H + B_1 L N + O(N^2), \quad (14)$$

$$X_H = E - \omega_H L, \quad (15)$$

$$P = X + O(N^2). \quad (16)$$

The near-critical particle:

$$X = E \left(\frac{b}{h} - C_1 \right) N \quad (17)$$

$$P = N \sqrt{E^2 \left[\left(\frac{b}{h} - C_1 \right)^2 - \frac{1}{h^2} \right] - m^2 + O(N^2)}. \quad (18)$$

We introduced notations $b = B_1 \sqrt{g_H}$, $h = \omega_H \sqrt{g_H}$.

III. SCENARIO OF COLLISION

In the point of collision we assume the validity of conservation laws for the energy and angular momentum:

$$E_0 \equiv E_1 + E_2 = E_3 + E_4, \quad (19)$$

$$L_0 = L_1 + L_2 = L_3 + L_4. \quad (20)$$

It follows from (19), (20) that

$$X_0 \equiv X_1 + X_2 = X_3 + X_4. \quad (21)$$

There is also the conservation law for the radial momentum:

$$\sigma_1 P_1 + \sigma_2 P_2 = \sigma_3 P_3 + \sigma_4 P_4. \quad (22)$$

We assume that particles 1 and 2 fall from infinity, so $\sigma_1 = \sigma_2 = -1$. We are interested in high energy processes in which $E_{c.m.}$ is unbounded since this is the necessary condition for E to be unbounded as well [16], [17]. To this end, we choose particle 1 to be the critical, particle 2 being usual since this gives rise to the unbounded $E_{c.m.}$ [1], [18]. Then, one of particles (say, 3) is near-critical and the other one (4) is usual [12], [13]. All possible scenarios can be described by two parameters - the sign of C_1 and the value of σ_3 immediately after collision (OUT for $\sigma_3 = +1$ and IN for $\sigma_3 = -1$). As a result, we have 4 scenarios OUT+, OUT-, IN+, IN-. The first three were already analyzed in [13], where IN- was rejected since it corresponds to both particles falling into a black hole. However, now it is this scenario which we focus on. It corresponds to high energy propagation in the white hole region (see below). Thus we have $\sigma_3 = \sigma_4 = -1$. We must analyze the process under discussion for $N_c \rightarrow 0$ on the basis of the conservation law (22). In doing so, we follow the lines of Ref. [13] applying the corresponding approach to the case that was not considered there.

IV. LOWER BOUNDS ON ENERGY

If we collect the terms of the zeroth and first order in N_c and take into account the approximate expressions (12) - (18), we obtain

$$F = -\sqrt{E_3^2 \left[\left(\frac{b}{h} - C_1 \right)^2 - \frac{1}{h^2} \right]} - m_3^2, \quad (23)$$

where

$$F \equiv A + E_3 \left(C_1 - \frac{b}{h} \right), \quad (24)$$

$$A = \frac{E_1 b - \sqrt{E_1^2 (b^2 - 1) - m_1^2 h^2}}{h}, \quad (25)$$

$$C_1 = \frac{b}{h} - \frac{A^2 + m_3^2 + \frac{E_3^2}{h^2}}{2E_3A}, \quad (26)$$

$$F = \frac{A^2 - m_3^2 - \frac{E_3^2}{h^2}}{2A}. \quad (27)$$

We are interested in scenario IN-. Then, $C_1 < 0$ gives us

$$E_3^2 - 2E_3hA_1b + h^2(A^2 + m_3^2) > 0, \quad (28)$$

that can be rewritten as

$$(E_3 - \lambda_+)(E_3 - \lambda_-) > 0, \quad (29)$$

$$\lambda_{\pm} = h[A_1b \pm \sqrt{A^2(b^2 - 1) - m_3^2}]. \quad (30)$$

The condition $F < 0$ gives us

$$E_3^2 > h^2(A^2 - m_3^2) \equiv \lambda_0^2. \quad (31)$$

If λ_{\pm} are real, both bounds $E_3 > \lambda_+$ and $E_3 > \lambda_0$ are quite compatible with each other. If λ_{\pm} are complex, (28) and (31) are mutually consistent as well. Thus there is no upper bound on E_3 and the SPP is possible.

As far as particle 4 is concerned, it has $E_4 < 0$. To obey the forward-in-time condition (8), it must have $L_4 = -|L_4| < 0$. Then, $X_4 = |L_4|\omega - |E_4|$. Assuming that there is a flat infinity, where $\omega \rightarrow 0$, we see that particle 4 either falls into singularity or oscillates between turning points r_1 and r_2 . In doing so, it can intersect the horizons, thus appearing in new "universes" due to a potentially rich space-time structure inside similarly to what takes place for the Kerr metric [19]. However, under a rather weak and reasonable restrictions on the properties of the metric, it cannot have more than 1 turning point in the outer region, so the situation when $r_+ < r_1 \leq r \leq r_2$ is impossible. This was shown for the Kerr metric in [20] and generalized in [21]. For more information about trajectories of particles 3 and 4, the metric should be specified.

The above treatment changes only slightly if we consider the Schnittman process [22] when the critical particle 1 does not come from infinity but moves from the horizon. Then, instead of (25), we should take $A = \frac{E_1b + \sqrt{E_1^2(b^2 - 1) - m_1^2h^2}}{h}$.

V. SUPERENERGETIC PARTICLES

In the above treatment, we tacitly assumed that all energies and angular momenta are finite and do not grow unbounded when $N_c \rightarrow 0$. The only place where N_c appear in the relation between them are equalities (9), (10), where it gives only small corrections. The above approximate expressions for particle characteristics (12) - (18) take into account this circumstances. In particular, for a usual particle, $X = O(1)$, the second term in the radical in (6) has the order N_c^2 . For a near-critical one, both terms in P have the order $O(N_c)$. Meanwhile, it turns out that there exists self-consistent scenario, in which

$$L_3 = \frac{l_3}{\sqrt{N_c}}, \quad (32)$$

where l_3 is some coefficient not containing N_c . For small N_c , $L_4 = L_0 - L_3 \approx -\frac{l_3}{\sqrt{N_c}}$.

It was found in [23], where it was pointed out that it corresponds to falling both particles in a black hole, so it was put aside since we were interested in particles returning to infinity. But now, it is this case that came into play. Therefore, we take advantage of formulas already derived in [23] but exploit them in a new context - see eqs. (35), (36) below. If (32) is satisfied, the previous consideration fails and the conservation law (22) is to be analyzed anew. Now, for particles 3 and 4 the second term in (6) gives a small correction (whereas for finite L_3 both terms for particle 3 would have the same order), so

$$P_{3,4} \approx \sqrt{X_{3,4}^2 - N_c \frac{l_{3,4}^2}{(g_\phi)_H}} \approx X_{3,4} - \frac{N_c}{2X_{3,4}} \frac{l_3^2}{(g_\phi)_H}. \quad (33)$$

Then, taking into account (12) - (16) for particles 1 and 2, (19) - (21) and discarding the terms $O(N_c^2)$ and higher, one can show after algebraic manipulations that the following equation holds:

$$\frac{l_3^2}{2(g_\phi)_H} \left(\frac{1}{X_3} + \frac{1}{X_4} \right) = A. \quad (34)$$

Using again (21), one can obtain

$$(X_{3,4})_c \approx \frac{(X_0)_c}{2} (1 \mp \sqrt{1-b}), \quad (35)$$

where

$$b \equiv \frac{2l_3^2}{(g_\phi)_H X_0 A}. \quad (36)$$

It is implied that $b < 1$.

It follows from definition (3) that now

$$E_3 = (X_3)_H + \omega_H L_3 \approx (X_3)_c + \omega_H \frac{l_3}{\sqrt{N_c}}. \quad (37)$$

Thus we have two usual particles which come down into a black hole. This is contrasted with the standard case when particle 3 is near-critical and returns to infinity.

We see that there are two energy scales for the SPP. On the first scale, E_3 can be as large as we like but with reservation that $E_3 \ll \frac{\text{const}}{\sqrt{N_c}}$. On the second scale, $E_3 \sim \frac{1}{\sqrt{N_c}}$.

In doing so, $E_4 = E_0 - E_3$ is negative having the same order $N_c^{-1/2}$. Discussion about the properties of the trajectories of such particles from Section IV applies now as well.

VI. CONCLUSIONS

Thus we showed that particle collision on our side of Universe (near the black hole horizon) can lead to high energy fluxes on the other side. If, vice versa, collision occurs in "another world", we can detect its consequences in our one. We did not resort to the transformation between the original stationary frame and the center of mass one. We would like to stress that high energy behavior is found for the energies E_3 that can be in principle detected in a laboratory. These results qualitatively agree with claims made in [11] for the Kerr metric.

It is instructive to compare the situation for static charged and neutral rotating black/white holes collecting the results of the present and previous works [24], [25].

	SPP	q_3	L_3
charged black holes	yes	large	arbitrary
charged white holes	yes	large	$O(N_c^{-1/2})$
neutral rotating black holes	no	0	arbitrary
neutral rotating white holes	yes	0	unbounded

Table 1. Conditions of the existence of the super-Penrose process.

We see that for the Reissner-Nordström metric, if we compare collisions near black and white holes, the situation is partially complementary to each other. For finite L_i (for example, with all $L_i = 0$), there is the SPP in the black hole case. However, it fails to exist near white holes. As shown in [25], only if particles with angular momenta $L_{3,4} = O(N_c^{-1/2})$ come into play, we obtain the SPP. Meanwhile, for the rotating case, the SPP does not exist for black holes at all. Instead, the white hole scenario opens new possibilities for the SPP,

in which E and L of particles at infinity are unbounded. This happens on two scales: on the first one E and L do not contain the parameter N_c^{-1} , on the second scale they have the order $O(N_c^{-1/2})$ similarly to the static charged case. The concrete properties of collisions are described by somewhat different formulas and the type of energetic particles are different: in the first case particle 3 is near-critical, in the second one it is usual. However, in both cases the conserved energy E_3 is unbounded.

The phenomenon under discussion is two-faced. On one hand, it shows that high energy processes are indeed possible due to white holes and poses anew the question about their potential role in nature. From the other hand, it poses also a question about backreaction of such collisions on the metric itself, including the fate of white holes.

Acknowledgments

This work was supported by the Russian Government Program of Competitive Growth of Kazan Federal University.

-
- [1] Bañados, M., Silk, J., West, S.M.: Kerr black holes as particle accelerators to arbitrarily high energy. *Phys. Rev. Lett.* **103**, 111102 (2009). arXiv:0909.0169
 - [2] Piran T., Shaham J., Katz J.: High efficiency of the Penrose mechanism for particle collisions. *Astrophys. J.* **196**, L107 (1975)
 - [3] T. Piran and J. Shaham.: “Upper bounds on collisional Penrose processes near rotating black-hole horizons, *Phys. Rev. D* **16**, 1615 (1977)
 - [4] Grib A. A., Pavlov Yu. V.: Are black holes totally black? *Gravit. Cosmol.* **21**, 13 (2015). arXiv:1410.5736
 - [5] Zaslavskii O. B.: On white holes as particle accelerators. *Gravit. Cosmol.* **24**, 92 (2018) arXiv:1707.07864
 - [6] Penrose R.: Gravitational Collapse: the role of general relativity. *Rivista del Nuovo Cimento. Numero Speciale I*, 252 (1969)
 - [7] Schnittman J. D.: The Collisional Penrose Process. *Gen. Relat. and Gravitation.* **50** (2018) 77. arXiv:1910.02800

- [8] Zaslavskii O. B.: Is the super-Penrose process possible near black holes? *Phys. Rev. D* **93** (2016), 024056 arXiv:1511.07501
- [9] Novikov I. D.: Delayed explosion of a part of the Friedmann Universe and quasars. *Astron. Zh.* **6**, 1075 (1964) [*Sov. Astronomy* **8**, 857 (1964)].
- [10] Narlikar J. V., Appa Rao K. M. V., Dadhich N. .: High energy radiation from white holes. *Nature* **251**, 590 (1974).
- [11] Patil M., Harada T.: Extremal Kerr white holes as a source of ultra high energy particles. arXiv:2004.12874
- [12] T. Harada, H. Nemoto, and U. Miyamoto.: Upper limits of particle emission from high-energy collision and reaction near a maximally rotating Kerr black hole. *Phys. Rev. D* **86**, 024 027 (2012). arXiv:1205.7088
- [13] Zaslavskii O. B.: Energetics of particle collisions near dirty rotating extremal black holes: Banados-Silk-West effect versus Penrose process. *Phys. Rev. D* **86**, 084030 (2012), arXiv:1205.4410
- [14] Frolov, V.P., Novikov, I.D.: *Physics of Black Holes*. Kluwer Academic, Dordrecht (1998)
- [15] Tanatarov I. V., Zaslavskii O. B.: Dirty rotating black holes: regularity conditions on stationary horizons. *Phys. Rev. D* **86**, 044019, (2012). arXiv:1206.2580
- [16] Patil, M., Harada, T., Nakao, K., Joshi, P.S., Kimura, M.: Infinite efficiency of collisional Penrose process: can over-spinning Kerr geometry be the source of ultra-high-energy cosmic rays and neutrinos? *Phys. Rev. D* **93**, 104015 (2016). arXiv:1510.08205
- [17] Tanatarov I. V., Zaslavskii O. B.: Collisional super-Penrose process and Wald inequalities. *Gen. Relativ. Gravit.* **49**, 119 (2017). arXiv:1611.05912
- [18] O.B. Zaslavskii.: Acceleration of particles as universal property of rotating black holes, *Phys. Rev. D* **82** (2010) 083004. arXiv:1007.3678
- [19] Carter B.: Global structure of the Kerr family of gravitational fields. *Phys. Rev.* **174**, 1559 (1968)
- [20] Grib A. A., Pavlov Yu. V., Vertogradov V. D.: Geodesics with negative energy in the ergosphere of rotating black holes. *Mod. Phys. Lett. A* **29**, 1450110 (2014). arXiv:1304.7360
- [21] O. B. Zaslavskii.: On geodesics with negative energies in the ergoregions of dirty black holes, *Mod. Phys. Lett. A* **30**, 1550055 (2015). arXiv:1412.1725
- [22] J. D. Schnittman.: Revised upper limit to energy extraction from a Kerr black hole, *Phys.*

- Rev. Lett. **113**, 261102 (2014) arXiv:1410.6446
- [23] O.B. Zaslavskii.: Center of mass energy of colliding electrically neutral particles and super-Penrose process Phys. Rev. D 100, 024050 (2019). arXiv:1904.04874
- [24] Zaslavskii O. B.: Energy extraction from extremal charged black holes due to the BSW effect. Phys. Rev. D **86**, 124039 (2012) arXiv:1207.5209
- [25] Zaslavskii O. B.: Super-Penrose process for extremal charged white holes. arXiv:2005.11090