

Constructing black holes in Einstein-Maxwell-scalar theory

Shuang Yu* Jianhui Qiu† Changjun Gao‡

*Key Laboratory of Computational Astrophysics,
National Astronomical Observatories,
Chinese Academy of Sciences, Beijing 100101, China and
School of Astronomy and Space Sciences,
University of Chinese Academy of Sciences,
No. 19A, Yuquan Road, Beijing 100049, China*

Abstract

Exact black hole solutions in the Einstein-Maxwell-scalar theory are constructed. They are the extensions of dilaton black holes in de Sitter or anti de Sitter universe. As a result, except for a scalar potential, a coupling function between the scalar field and the Maxwell invariant is present. Then the corresponding Smarr formula and the first law of thermodynamics are investigated.

PACS numbers: 04.50.Kd, 04.70.Dy

* yushuang@nao.cas.cn

† jhqiu@nao.cas.cn

‡ gaocj@bao.ac.cn

I. INTRODUCTION

The theory containing coupling between the scalar fields and gravity was firstly considered by Fisher who found a static and spherically symmetric solution of the Einstein massless scalar field equations [1]. Whereafter, variety of theories emerge with the introduction of scalar field, such as dimensional reductions, low-energy limit of string theories, various models of supergravity, etc. The low energy limit of the string theory has the Einstein action which is supplemented by fields, such as axion, gauge fields, and dilaton coupled in a nontrivial way to other fields. One of the most important theories of the low-energy limit of string theories is the Dilaton gravitation theory [2]. It is a very interesting theory because the dilaton field can naturally couples to gauge fields, and the properties of the black hole change drastically due to the dilaton field. The presence of the dilaton field can change the causal structure and asymptotic behavior of the spacetime as well as the thermodynamics and stability of the electrically charged black holes [3]. Therefore the dilaton black holes have attracted many attentions in recent decades.

On the other hand, anti de-Sitter space and conformal field theory correspondence (AdS/CFT) [4] is a significant method to unify quantum fields and gravitations. This correspondence between the gravity in an AdS spacetime and conformal field theory (CFT) leeching on to the boundary of the AdS spacetime suggests that physics of the gauge field defined on the AdS-boundary can be derived from gravitations in the bulk space and vice versa [5]. Stimulated by AdS/CFT, many works are devoted to find the asymptotically (anti)-de Sitter black hole solution in dilaton gravity with various self-interacting potential of dilaton. Exact solutions of charged dilaton black holes have been constructed by many authors. Some black holes are asymptotically flat [6–8] where the dilaton changes the causal structure of the black hole and leads to the curvature singularities at finite radii. Meanwhile, some solutions are asymptotically neither flat nor (anti)-de Sitter [9]. Finally, the asymptotically (anti)-de Sitter versions are found in [10–17]. These solutions reveal that the cosmological constant proposed in the Einstein theory is coupled to Liouville-type dilaton scalar potential. With the building of these asymptotically (anti)-de Sitter dilaton black holes, many interesting physical phenomena have been explored, such as the black hole thermodynamics [13, 18, 19], holographic thermalization [20], black hole phase transition in extend phase space [21] etc.

The purpose of this paper is, starting from the known dilaton black holes in de Sitter or anti-de Sitter universe, to construct new exact asymptotically (anti)-de Sitter black hole solutions in the Einstein-Maxwell-scalar theory which is the generalization of Einstein-Maxwell-dilaton theory. It is known that the Einstein-Maxwell-scalar theory can emerge naturally in physics, for example, in the contexts of Kaluza-Klein models [22], supergravity/string theory [23] and cosmology [24]. In the last two years, a remarkable phenomenon of spontaneous scalarisation of charged black holes is discovered by [25, 26]. It has motivated vast studies on various Einstein-Maxwell-scalar models (see [27] and references therein). This is another motivation for this research. Concretely, inspired by the expression of dilaton black holes in de Sitter or anti-de Sitter universe, we find two new exact black hole solutions in the Einstein-Maxwell-scalar theory. These solutions are the extensions of Reissner-Nordstrom-(anti)de Sitter and dilaton-(anti)de Sitter solutions. We subsequently investigate the thermodynamics.

The paper is organized as follows. In Sec. II, we derive the equations of motion in the Einstein-Maxwell-scalar theory. In Sec. III, we present the first electrically charged dilaton black hole in (anti)-de Sitter spacetime and give the analysis on its horizons. Sec. IV studied its thermodynamics which covers the buildings of Smarr formula and first law of thermodynamics. Then the second electrically charged dilaton black hole in (anti)-de Sitter spacetime with arbitrary coupling constant α is found in Sec. V and the corresponding analysis of thermodynamics are given in Sec. VI. Finally, we give the conclusion and discussion in Sec. VII. Throughout this paper, we adopt the system of units in which $G = c = \hbar = 1$ and the metric signature $(-, +, +, +)$.

II. THE EQUATIONS OF MOTION

The action of Einstein-Maxwell-scalar theory is given by

$$S = \int d^4x \sqrt{-g} [R - 2\nabla_\mu \phi \nabla^\mu \phi - K(\phi) F^2 - V(\phi)] , \quad (1)$$

where R is the Ricci scalar curvature, $F^2 \equiv F_{\mu\nu} F^{\mu\nu}$ comes from the Maxwell field, $K(\phi)$ is the coupling function between scalar field and Maxwell field. $V(\phi)$ is the scalar potential. When $K = 1$ and $V = 2\lambda$, it is the Einstein-Maxwell theory with cosmological constant λ . The theory gives the Reissner-Nordstrom-de Sitter solution. When $K = e^{2\phi}$ and $V = 0$, it

gives the dilaton black hole solution [6, 7]. When $K = e^{2\phi}$ and

$$V = \frac{\lambda}{3} (e^{2\phi} + 4 + e^{-2\phi}) , \quad (2)$$

it gives the dilaton black hole in de Sitter universe [10]. Except for these solutions, there are other important solutions with different K and V [11–17]. In principle, once the expressions of $K(\phi)$ and $V(\phi)$ are given, the gravity theory and the corresponding black hole solution are assigned. This is the usual way. But in this paper, we shall presume not $K(\phi)$ and $V(\phi)$, but the black hole solution in advance. Then $K(\phi)$ and $V(\phi)$ are subsequently determined. This is the solution-generating method [28]. We require that the parameters such as the mass M and electric charge Q don't appear in the coupling function $K(\phi)$ and the scalar potential $V(\phi)$. In the next, we derive the equations of motion.

Varying the action with respect to the metric, Maxwell, and dilaton fields, respectively, yields

$$R_{\mu\nu} = 2\nabla_\mu\phi\nabla_\nu\phi + \frac{1}{2}g_{\mu\nu}V + 2K \left(F_{\mu\alpha}F_\nu^\alpha - \frac{1}{4}g_{\mu\nu}F^2 \right) , \quad (3)$$

$$\nabla_\mu (KF^{\mu\nu}) = 0 , \quad (4)$$

$$\nabla_\mu\nabla^\mu\phi - \frac{1}{4}(V_{,\phi} + K_{,\phi}F^2) = 0 , \quad (5)$$

where “ $,\phi$ ” denotes the derivative with respect to ϕ . The metric for static and spherically symmetric black hole solution can always be written as

$$ds^2 = -U(r)dt^2 + U(r)^{-1}dr^2 + f(r)(d\theta^2 + \sin^2\theta d\phi^2) . \quad (6)$$

In this spacetime, the non-vanishing components of four-vector A_μ is uniquely $A_0(r)$. So the equations of motion turn out to be

$$2ff'' + 4f^2\phi'^2 - f'^2 = 0 , \quad (7)$$

$$(fKA_0')' = 0 , \quad (8)$$

$$2f^2U'' + 2fUf'' + 2fU'f' - Uf'^2 + 4Uf^2\phi'^2 - 4f^2KA_0'^2 + 2f^2V = 0 , \quad (9)$$

$$fU\phi'' + Uf'\phi' + fU'\phi' + \frac{1}{2}fK_{,\phi}A_0'^2 - \frac{1}{4}fV_{,\phi} = 0 . \quad (10)$$

Here prime denotes the derivative with respect to r . Observing these equations, we find six unknown functions. They are U , f , ϕ , A_0 , K , V . But we have only four equations of motion. Therefore, the equations of motion are not closed. In principle, one can presume any two of them in advance. Then the equations of motion are closed. In the next section, we shall bring forward U and f beforehand. Afterwards, the other four functions are worked out. By this way, we construct two black hole solutions.

III. THE FIRST SOLUTION

We recall that when $K = e^{2\phi}$ and $V = 0$, the theory gives the dilaton black hole solution with [6, 7]

$$\begin{aligned} U &= 1 - \frac{2M}{r}, & f &= r \left(r - \frac{Q^2}{M} \right), \\ \phi &= -\frac{1}{2} \ln \left(1 - \frac{Q^2}{Mr} \right). \end{aligned} \quad (11)$$

Here M and Q have the physical meaning of ADM mass and electric charge of the black hole, respectively. This spacetime is asymptotically flat. On the other hand, when $K = e^{2\phi}$ and V of Eq. (2), the theory gives the dilaton black hole in de Sitter universe with [10]

$$\begin{aligned} U &= 1 - \frac{2M}{r} - \frac{1}{3}\lambda f, & f &= r \left(r - \frac{Q^2}{M} \right), \\ \phi &= -\frac{1}{2} \ln \left(1 - \frac{Q^2}{Mr} \right). \end{aligned} \quad (12)$$

Here λ is a constant. When $Q = 0$, it goes back to the Schwarzschild-de Sitter solution. In order to construct a new solution, we observe the expression of U and find that the λ term is proportional to not r^2 , but f . Inspired by this fact, we presume the new solution

$$U = 1 - \frac{2M}{r} + \frac{\beta Q^2}{f} - \frac{1}{3}\lambda f, \quad (13)$$

$$f = r \left(r + \frac{\gamma Q^2}{M} \right). \quad (14)$$

Here β and γ are dimensionless constants. The β term is a new term and it is similar to the Q^2/r^2 term in Reissner-Nordstrom-de Sitter solution. Then substituting the equations (13) and (14) into the equations of motion (7,8,9,10), we obtain

$$\phi = -\frac{1}{2} \ln \left(1 + \frac{\gamma Q^2}{Mr} \right), \quad (15)$$

$$A_0 = \frac{\gamma Q}{r} - \frac{\beta Q}{2r} - \frac{\beta M Q}{2(Mr + \gamma Q^2)}, \quad (16)$$

$$K = \frac{2e^{2\phi}}{-2\gamma + \beta + \beta e^{4\phi}}, \quad (17)$$

$$V = \frac{\lambda}{3} (e^{2\phi} + 4 + e^{-2\phi}). \quad (18)$$

We notice that the scalar potential, Eq. (18) is exactly that in Eq. (2). Two dimensionless constants γ and β appear in the coupling function K . Up to this point, equations from Eq. (13) to Eq. (18) constitute the first black hole solution in this paper. When $\gamma = -1$, $\beta = 0$, the solution returns to Eq. (12). When $\gamma = 0$ and $\beta = 1$, the Reissner-Nordstrom-de Sitter solution is produced. We point out that in dilaton gravity, we have $K = e^{2\phi}$ which is a positive function. Therefore, we require $\gamma < 0$ and $\beta > 0$ in order that K is always positive.

In the following, we shall compute the dilaton charge D , the electric charge Q and the ADM mass M of the black hole. The scalar charge D can be computed by

$$D = \frac{1}{4\pi} \oint \nabla_\mu \phi \, dS^\mu = \frac{\gamma Q^2}{2M}. \quad (19)$$

Here the integration is taken over a spacelike surface enclosing the origin. The scalar charge is a conserved value and thus it does not depend on the choice of the surface. It is clear that the scalar charge is determined by the mass M , electric charge Q and the dimensionless constant γ .

The electric charge of the black hole is shown to be

$$Q = \frac{1}{4\pi} \oint K(\phi) \nabla_\mu A_0 \, dS^\mu. \quad (20)$$

The electric charge is also a conserved value and it does not depend on the choice of the surface.

The ADM mass [29–31] satisfies

$$M = \frac{1}{16\pi} \lim_{S^\mu \rightarrow i^0} \oint g^{\alpha\beta} (g_{\alpha\mu,\beta} - g_{\alpha\beta,\mu}) \, dS^\mu, \quad (21)$$

where i^0 is the spacelike infinity. Now we conclude that the scalar charge D , the electric charge Q and the ADM mass M do satisfy the corresponding definitions.

There is a curvature singularity at $r = -2D$ for this spacetime. As for the black hole horizons, we find that, under the condition that $M > 0$, $\gamma < 0$, $\beta > 0$, there are at most

three horizons for positive λ . They are cosmic horizon, black hole event horizon and black hole inner horizon, respectively, as shown in Fig. 1. For negative λ , there are at most two horizons. They are black hole event horizon and black hole inner horizon. In the next section, we shall study the thermodynamics of this spacetime.




FIG. 1: There are four cases for the scenario of horizons. The curves correspond to 1-horizon, 2-horizon, 3-horizon and 2-horizon from down to up, respectively. We fix $M = 1$, $Q = 1$, $\gamma = -1$ and choose $\beta = 0.1$, $\lambda = 0.5$; $\beta = 0.1$, $\lambda = 0.184$; $\beta = 0.2$, $\lambda = 0.14$ and $\beta = 0.277$, $\lambda = 0.12$, respectively.

IV. BLACK HOLE THERMODYNAMICS

In order to explore the black holes thermodynamics, we start from the calculation of its temperature. The black hole temperature is well defined in quite general setting due to its strict geometrical basis [32]. One can employ the so-called surface gravity which is defined as follows:

$$\kappa^2 = -\frac{1}{2}\nabla_\mu\chi_\nu\nabla^\mu\chi^\nu, \quad (22)$$

where χ_μ is a Killing vector field and it is null on the horizon. We can chose $\chi^\nu = \partial/\partial t$ because spacetime is the static. As a result, the black hole temperature takes the form

$$T = \frac{\kappa}{2\pi} = \frac{1}{4\pi}U'(r_+)$$

$$\begin{aligned}
&= \frac{M}{2\pi r_+^2} - \frac{\beta Q^2}{4\pi r_+^2 (r_+ + 2D)} - \frac{\beta Q^2}{4\pi r_+ (r_+ + 2D)^2} \\
&\quad - \frac{\lambda r_+}{6\pi} - \frac{\gamma \lambda Q^2}{12\pi M},
\end{aligned} \tag{23}$$

where r_+ represents the black hole event horizon which is determined by $U(r_+) = 0$. In Fig. 2(a) and Fig. 2(b) we plot the black hole temperature T as the function of black hole event horizon r_+ with running β and Q , respectively. Fig. 2(a) shows there are two phases of black holes with the same temperature, the so-called small and large black holes, respectively. When $\beta = 0.335$, there is uniquely one phase and its temperature is zero. With the increasing of β , the black holes make phases transition from 2-phase to 1-phase. Fig. 2(b) shows that there are three phases of black holes with the same temperature. They are large, middle and small black holes, respectively. With the decreasing of Q , the black holes make phases transition from 1-phase to 3-phase.




FIG. 2: Black hole temperature T as the function of black hole event horizon r_+ . (a) The curves correspond to $\beta = 0.2, 0.24, 0.28, 0.32, 0.335$, from up to down, respectively. We put $M = 1, Q = 1, \gamma = -1$. There are two phases of black holes with the same temperature, the so-called small (solid lines) and large black holes (dotted lines), respectively. When $\beta = 0.335$, there is uniquely one phase and its temperature is zero. (b) The curves correspond to $Q = 0.24, 0.22, 0.20, 0.18, 0.16, 0.14$, from down to up, respectively. We put $\lambda = -1.5, \beta = 1.4, \gamma = -0.1$. With the decreasing of Q , the black holes make phases transition from 1-phase to 3-phase which consists of small (solid line), middle (dash-dot line) and large (dotted line) black holes.

The entropy of black holes generally satisfies the area law which states that the entropy

is a quarter of the area of black hole event horizon [33]. The law applies to almost all kinds of black holes including Einstein-Maxwell-scalar black holes [34]. Therefore we have the entropy of the black hole

$$S = \frac{A}{4} = \pi r_+ (r_+ + 2D) , \quad (24)$$

According to definition of [35], the electric potential measured by the distant observer is

$$\Phi = \int_{r_+}^{+\infty} A'_0 dr = -A_0(r_+) . \quad (25)$$

The study on the thermodynamic phase structure of AdS black holes [36] tells us the cosmological constant acts as a thermodynamic pressure

$$P = -\frac{\lambda}{8\pi} . \quad (26)$$

The variation of thermal pressure requires the presence of a conjugate thermal volume in the first law of thermodynamics, which can lead to a variety of novel thermodynamic behaviour, for example, triple points [37], reentrant phase transitions [38], the emergence of polymer-like phase structure [39], the superfluid-like phase structure [40] and the Van der Waals transition [41].

We find the conjugate thermodynamic volume is

$$\mathfrak{V} \equiv \left(\frac{\partial M}{\partial P} \right)_{S,Q} = \frac{4\pi}{3} r (r + 2D) (r + D) . \quad (27)$$

Then the Smarr formula

$$M = 2TS - 2\mathfrak{V}P + Q\Phi , \quad (28)$$

is satisfied.

In the next, we shall make an examination whether the thermal quantities fulfill the requirement of the first law of thermodynamics. From the equation of horizon, we obtain

$$\lambda = \frac{3}{f_+} \left(1 - \frac{2M}{r_+} + \frac{\beta Q^2}{f_+} \right) , \quad (29)$$

where $f_+ = f(r = r_+)$. So in view of Eq. (26), the pressure P can be written as

$$P = -\frac{3}{8\pi f_+} \left(1 - \frac{2M}{r_+} + \frac{\beta Q^2}{f_+} \right) . \quad (30)$$

We treat the pressure P , the entropy S as the function of r_+ , M , Q . Then we have

$$dP = P_{,M}dM + P_{,Q}dQ + P_{,r_+}dr_+ , \quad (31)$$

$$dS = S_{,M}dM + S_{,Q}dQ + S_{,r_+}dr_+ . \quad (32)$$

By using the equations Eq. (23), Eq. (25) and Eq. (27), we find the first law of thermodynamics

$$dM = TdS + \Phi dQ + \mathfrak{V}dP , \quad (33)$$

is indeed satisfied.

It is pointed [36, 42] that once the thermodynamic pressure P and thermal volume \mathfrak{V} are introduced, the ADM mass M should be understood as enthalpy. Then the more convenient function to analyze thermodynamic behaviour of a system, especially in the case that there is some critical behaviour, is the Gibbs free energy which is defined in the following way

$$G = M - TS . \quad (34)$$

the Gibbs free energy G is understood to depend on M , Q , γ , β , r_+ .

In Fig. 3(a) and Fig. 3(b), we plot the $G(T)$ relations with running β and Q , respectively. With the increasing of β , the black holes make phases transition from 2-phase to 1-phase. It also tells us the Gibbs free energy of large black hole is decreasing with the increasing of Hawking temperature. On the other hand, the Gibbs free energy of small black hole first decreases and then increases with the increasing of Hawking temperature. As is known to us, the specific heat is $C_{Q,P} = -\partial^2 G / \partial T^2$. Therefore, the thermodynamically stable and unstable phases have the concave downward and upward $G(T)$ curves, respectively. Then we conclude that the large black holes are thermodynamically unstable while the small black holes are stable. Fig. 3(b) shows that with the decreasing of Q , the black holes make phases transition from 1-phase to 3-phase. In this case, we let Q be running. Then the Gibbs free energy of large, middle and small black holes all decreases with the increasing of Hawking temperature. But only large and small black holes are thermodynamically stable while the middle black holes are unstable.




FIG. 3: Black hole Gibbs free energy G as the function of temperature T . (a) The curves correspond to $\beta = 0.2, 0.24, 0.28, 0.32, 0.335$, from outer to inner, respectively. We put $M = 1, Q = 1, \gamma = -1$. There are two phases of black holes (dotted lines for small black hole and solid lines for large black holes) with the same temperature. When $\beta = 0.335$, the Gibbs free energy is $G = M = 1$. (b) The curves correspond to $Q = 0.22, 0.20, 0.18, 0.16, 0.14$, from up to down, respectively. We put $\lambda = -1.5, \beta = 1.4, \gamma = -0.1$. With the decreasing of Q , the black holes make phases transition from 1-phase to 3-phase.

V. SECOND SOLUTION

In this section, we construct the second solution in Einstein-Maxwell-scalar theory. We remember that the dilaton black hole in de Sitter universe for arbitrary coupling constant α is [10]

$$U = \left(1 - \frac{2m}{r}\right) \left(1 - \frac{Q^2}{mr}\right)^{\frac{1-\alpha^2}{1+\alpha^2}} - \frac{1}{3}\lambda f, \quad (35)$$

with

$$f = r^2 \left(1 - \frac{Q^2}{mr}\right)^{\frac{2\alpha^2}{1+\alpha^2}}. \quad (36)$$

The corresponding coupling function K and scalar potential V in the action are

$$K = e^{2\alpha\phi}, \quad (37)$$

and

$$V = \frac{2\lambda}{3(1+\alpha^2)^2} [\alpha^2 (3\alpha^2 - 1) e^{2\phi/\alpha} + (3 - \alpha^2) e^{-2\alpha\phi} + 8\alpha^2 e^{-\phi\alpha + \phi/\alpha}]. \quad (38)$$

Observing the λ term in Eq. (35), we find it is proportional to f . So one presume

$$U = \left(1 - \frac{2m}{r}\right) \left(1 + \frac{\gamma Q^2}{mr}\right)^{\frac{1-\alpha^2}{1+\alpha^2}} + \frac{\beta Q^2}{f} - \frac{1}{3}\lambda f, \quad (39)$$

$$f = r^2 \left(1 + \frac{\gamma Q^2}{mr}\right)^{\frac{2\alpha^2}{1+\alpha^2}}. \quad (40)$$

We notice that when the coupling constant $\alpha = 1$, the solution restores to the first solution in section III. Substituting Eqs. (39), (40) into the equations of motion (7,8,9,10), we obtain

$$\phi = -\frac{\alpha}{1+\alpha^2} \ln \left(1 + \frac{\gamma Q^2}{mr}\right), \quad (41)$$

$$A_0 = \frac{2}{1+\alpha^2} \left[\frac{\gamma Q}{r} - \frac{\beta Q}{2r} - \frac{\beta m Q \alpha^2}{2(mr + \gamma Q^2)} \right], \quad (42)$$

$$K = \frac{(\alpha^2 + 1) e^{2\alpha\phi}}{-2\gamma + \beta + \beta\alpha^2 e^{2\alpha\phi + 2\phi/\alpha}}, \quad (43)$$

with the scalar potential V of the same expression in Eq. (38). When $\alpha = 1$, they reduce to the counterparts in the first solution of section III. So equations from (38) to Eq. (43) constitute the second black hole solution in this study. We have replaced large M with small m in the expressions because it is M that represents the ADM mass. We shall show this point later.

The scalar charge D can be computed by

$$D = \frac{1}{4\pi} \lim_{S^\mu \rightarrow i^0} \oint \nabla_\mu \phi \, dS^\mu = \frac{\alpha\gamma Q^2}{m(1+\alpha^2)}. \quad (44)$$

Here the integration is taken over the surface of spacelike infinity, i^0 . It tells us the scalar charge is determined by m , electric charge Q , the coupling constant α and the dimensionless constant γ .

The electric charge of the black hole is shown to be

$$Q = \frac{1}{4\pi} \oint K(\phi) \nabla_\mu A_0 \, dS^\mu. \quad (45)$$

Here the integration is taken over a spacelike surface enclosing the origin. The electric charge is a conserved value and it does not depend on the choice of the surface.

We find the ADM mass M of the spacetime is

$$M = \frac{1}{16\pi} \lim_{S^\mu \rightarrow i^0} \oint g^{\alpha\beta} (g_{\alpha\mu,\beta} - g_{\alpha\beta,\mu}) \, dS^\mu = m + \frac{D}{2} \left(\alpha - \frac{1}{\alpha}\right). \quad (46)$$

where i^0 is the spacelike infinity. We see the ADM mass M is not equals to m provided that $\alpha \neq 1$.

The curvature singularity of this spacetime is at $r = -\gamma Q^2/m$. The situations of horizons is similar to the first solution. So we shall not present a detailed discussion. In the next section, we study the black hole thermodynamics.

VI. THERMODYNAMICS

Following the procedure in section IV, we figure out the thermodynamic quantities for the second solution, namely, the black hole temperature

$$T = \frac{\kappa}{2\pi} = \frac{1}{4\pi} U' (r_+) , \quad (47)$$

the entropy

$$S = \frac{A}{4} = \pi r_+^2 \left(1 + \frac{\gamma Q^2}{m r_+} \right)^{\frac{2\alpha^2}{1+\alpha^2}} , \quad (48)$$

the electric potential

$$\Phi = \int_{r_+}^{+\infty} A'_0 dr = -A_0 (r_+) , \quad (49)$$

the thermal pressure

$$P = -\frac{\lambda}{8\pi} , \quad (50)$$

and the thermal volume

$$\mathfrak{V} = \frac{4\pi}{3} \cdot \frac{(1 + \alpha^2) r + 2D}{r (1 + \alpha^2) (r + 2D)} f^2 . \quad (51)$$

With these expressions, the Smarr formula

$$M = 2TS - 2\mathfrak{V}P + Q\Phi , \quad (52)$$

is achieved. In the next, we check whether these thermodynamic quantities satisfy the first law of thermodynamics.

From the equation of horizon, we obtain

$$\lambda = \frac{3}{f_+} \left[\left(1 - \frac{2m}{r_+} \right) \left(1 + \frac{\gamma Q^2}{m r_+} \right)^{\frac{1-\alpha^2}{1+\alpha^2}} + \frac{\beta Q^2}{f_+} \right] , \quad (53)$$

where $f_+ = f (r = r_+)$. So the pressure P can be written as

$$P = -\frac{3}{8\pi f_+} \left[\left(1 - \frac{2m}{r_+} \right) \left(1 + \frac{\gamma Q^2}{m r_+} \right)^{\frac{1-\alpha^2}{1+\alpha^2}} + \frac{\beta Q^2}{f_+} \right] . \quad (54)$$

We take the pressure P , the entropy S as the function of r_+ , m and Q . Then we have

$$dP = P_{,m}dm + P_{,Q}dQ + P_{,r_+}dr_+ , \quad (55)$$

$$dS = S_{,m}dm + S_{,Q}dQ + S_{,r_+}dr_+ , \quad (56)$$

$$dM = M_{,m}dm + M_{,Q}dQ , \quad (57)$$

By using the equations Eq. (47), Eq. (49) and Eq. (51), we arrive at the first law of thermodynamics

$$dM = TdS + \Phi dQ + \mathfrak{A}dP . \quad (58)$$

In Fig. 4(a) and Fig. 5(a), we plot the $T - r_+$ relation and $G - T$ relation with $M = 1$, $Q = 1$, and $\gamma = -1$, $\beta = 0.1$. They show that with the increasing of α , the black holes make phase transition from 1-phase to 2-phase. In Fig. 4(b) and Fig. 5(b), we plot the $T - r_+$ relation and $G - T$ relation with $M = 1$, $\alpha = 5$, $\lambda = -1.5$, $\beta = 1.4$, $\gamma = -0.1$. They show that with the decreasing of Q , the black holes make phase transition from 1-phase to 3-phase. Same as the scenario of first black hole solution, the large black holes are thermodynamically unstable while the small black holes are stable in the case of two phases. For the structure of three phases, both the small and large black holes are all thermodynamically stable while middle black holes are unstable.

VII. CONCLUSION AND DISCUSSION

In conclusion, we construct two exact and electrically charged black holes in the Einstein-Maxwell-scalar theory. The two solutions are all asymptotically de Sitter or anti-de Sitter. They are the extensions of dilaton black holes in de Sitter or anti-de Sitter universe. The first solution is for the case of $\alpha = 1$ and the second solution is for arbitrary α . In this sense, the second solution is the extension of the first solution. An interesting fact is that the corresponding scalar potential $V(\phi)$ is the same as the dilaton ones. The only difference from the Einstein-Maxwell-dilaton theory is the presence of coupling function $K(\phi)$ in front of the Maxwell invariant. With the presence of $K(\phi)$, the well-known Reissner-Nordstrom-(anti) de Sitter solution is included in these solutions.

We calculate the ADM mass M , the electric charge Q and the scalar charge D of the black holes. The scalar charge D is not independent, but dependent on M and Q . This is the




FIG. 4: Black hole temperature T as the function of black hole event horizon r_+ . (a) The curves correspond to $\alpha = 0.74, 0.78, 0.84, 0.92, 1.0$, from down to up, respectively. We put $M = 1, Q = 1, \gamma = -1, \beta = 0.1$. With the increasing of α , the black holes make phase transition from 1-phase to 2-phase, the so-called small (solid lines) and large (dotted lines) black holes. (b) The curves correspond to $Q = 0.24, 0.22, 0.20, 0.18, 0.16, 0.14$, from down to up, respectively. We put $\alpha = 5, \lambda = -1.5, \beta = 1.4, \gamma = -0.1$. With the decreasing of Q , the black holes make phases transition from 1-phase to 3-phase (small, middle and large black holes).

same as the dilaton black holes. Then we compute the Hawking temperature, the entropy, the electric potential and the thermal volume. Afterwards, both the Smarr formula and the first law of thermodynamics are constructed. The evolution of Hawking temperature and Gibbs free energy shows that these black holes have two situations of phase structure, the scenario of two phases and three phases. In the scenario of two phases, we let the coupling constants β and α be running. In this case, the Gibbs free energy of large black hole is decreasing with the increasing of Hawking temperature. On the other hand, the Gibbs free energy of small black holes first decreases and then increases with the increasing of Hawking temperature. This shows the large black hole is thermodynamically unstable and the small black hole is stable. In the scenario of three phases, we let Q be running. In this case, the Gibbs free energy of large, middle and small black holes all decreases with the increasing of Hawking temperature. But only the large and small black holes are thermodynamically stable while the middle black holes are unstable.

Finally, since these black hole solutions are asymptotically anti-de Sitter, it is worth studying the corresponding ADS/CFT correspondence. On the other hand, the extension




FIG. 5: Black hole Gibbs free energy G as the function of temperature T . (a) The curves correspond to $\alpha = 0.74, 0.78, 0.84, 0.92, 1.0$, from inner to outer, respectively. We put $M = 1, Q = 1, \gamma = -1, \beta = 0.1$. There are two phases of black holes with the same temperature. (b) The curves correspond to $Q = 0.22, 0.20, 0.18, 0.16, 0.14$, from down to up, respectively. We put $\lambda = -1.5, \beta = 1.4, \gamma = -0.1, \alpha = 5$. With the increasing of Q , the black holes make phases transition from 1-phase to 3-phase.

of these static black holes to the rotating and higher dimensional cases are also worthy of study.

ACKNOWLEDGMENTS

This work is partially supported by China Program of International ST Cooperation 2016YFE0100300, the Strategic Priority Research Program “Multi-wavelength Gravitational Wave Universe” of the CAS, Grant No. XDB23040100, the Joint Research Fund in Astronomy (U1631118), and the NSFC under grants 11473044, 11633004 and the Project of CAS, QYZDJ-SSW-SLH017.

-
- [1] I. Z. Fisher, Zh. Eksp. Teor. Fiz. 18, 636 (1948) [Sov. Phys. JETP 18, 636 (1948)].
[2] M. B. Green, J.H. Schwarz and E. Witten, Superstring Theory, Cambridge University Press, 1987.

- [3] B. Harms and Y. Leblanc, Phys. Rev. D 46, 2334 (1992); C. F. E. Holzhey and F. Wilczek, Nucl. Phys. B380, 447(1992).
- [4] J. M. Maldacena, The Large N limit of superconformal field theories and supergravity, Int. J. Theor. Phys.38, 1113 (1999).
- [5] E. Witten, Adv. Theor. Math. Phys. 2, 253 (1998); E. Witten, Adv. Theor. Math. Phys. 2, 505 (1998).
- [6] G.W. Gibbons and K. Maeda, Nucl. Phys. B **298**, 741 (1988).
- [7] D. Garfinkle, G. Horowitz, and A. Strominger, Phys. Rev. D **43**, 3140 (1991).
- [8] D. Brill and G. Horowitz, Phys. Lett. B 262, 437 (1991); R. Gregory and J. Harvey, Phys. Rev. D 47, 2411 (1993); T. Koikawa and M. Yoshimura, Phys. Lett. B 189, 29 (1987); D. Boulware and S. Deser, Phys. Lett. B 175, 409 (1986); M. Rakhmanov, Phys. Rev. D 50, 5155 (1994).
- [9] K. C. K. Chan, J. H. Horn and R. B. Mann, Nucl. Phys. B447, 441 (1995) ; G. Clement and C. Leygnac, Phys.Rev. D70, 084018 (2004) ; Rong-Gen Cai and An Zhong Wang, Phys. Rev. D70, 084042 (2004); Rong-Gen Cai and Yuan Zhong Zhang, Phys. Rev. D64, 104015 (2001); Rong-Gen Cai, Jeong-Young Ji and Kwang-Sup Soh, Phys. Rev. D57, 6547 (1998); Rong-Gen Cai and Yuan-Zhong Zhang, Phys. Rev. D54, 4891(1996); G. Clement, D. Galtsov, C. Leygnac, Phys. Rev. D 67, 024012(2003); A. Sheykhi, M.H. Dehghani, N. Riazi, Phys. Rev. D 75, 044020(2007); A. Sheykhi, M.H. Dehghani, N. Riazi, J. Pakravan, Phys. Rev. D74, 084016 (2006); A. Sheykhi, Phys. Rev. D 76, 124025 (2007); A. Sheykhi, Phys. Lett. B 662, 7 (2008); M.H. Dehghani, J. Pakravan, S.H. Hendi, Phys. Rev. D 74, 104014(2006); M.H. Dehghani et al., J. Cosmol. Astropart. Phys. 02, 020 (2007); M. H. Dehghani, A. Sheykhi and S. H. Hendi, Phys. Lett. B659, 476 (2008) ; S.H. Hendi, J. Math. Phys. 49, 082501 (2008).
- [10] C. Gao and S. N. Zhang, Phys. Rev. D **70**, 124019 (2004).
- [11] C. Gao and S. N. Zhang, Phys. Lett.B 605 (2005) 185; C. Gao and S. N. Zhang, Phys. Lett. B 612, 127 (2005).
- [12] S. Hajkhalili, A. Sheykhi, Phys. Rev. D 99, 024028 (2019)
- [13] S. H. Hendi, A. Sheykhi, and M. H. Dehghani, Eur. Phys. J.C 70, 703 (2010).
- [14] S. Hajkhalili and A. Sheykhi, Int. J. Mod. Phys. D 27, 1850075 (2018).
- [15] A. Sheykhi, Phys. Rev. D 78, 064055 (2008) .
- [16] R. Yamazaki and D. Ida, Phys. Rev. D 64, 024009 (2001) .
- [17] T. Ghosh and S. SenGupta, Phys. Rev. D 76, 087504 (2007) ; A. Sheykhi and M. Allahverdizadeh, Phys. Rev. D 78, 064073 (2008) ; A. Sheykhi, Phys. Lett. B 672, 101 (2009) ; M. H. Dehghani and A. Bazrafshan, Int. J. Mod. Phys. D 19, 293 (2010) .
- [18] A. Sheykhi, M. H. Dehghani and S. H. Hendi, Thermodynamic instability of charged dilaton

- black holes in AdS spaces, *Phys. Rev. D* **81**, 084040(2010) .
- [19] A. Sheykhi, S. H. Hendi, F. Naeimipour, S. Panahiyan and B. Eslam Panah, Thermodynamic geometry of charged dilaton black holes in AdS spaces, *Can. J. Phys.* **94**, no.10, 1045 (2016).
- [20] S. J. Zhang and E. Abdalla, Holographic Thermalization in Charged Dilaton Antide Sitter Spacetime, *Nucl. Phys. B* **896**, 569 (2015) .
- [21] A. C. Li, H. Q. Shi and D. f. Zeng, *Phys. Rev. D* **97**, no.2, 026014 (2018); P. Wang, H. Wu and H. Yang, *Eur. Phys. J. C* **80**, 216 (2020), gr-qc:1910.07874; gr-qc:1904.12365; K. Liang, P. Wang, H. Wu and M. Yang, *Eur. Phys. J. C* **80**, 187 (2020) gr-qc: 1907.00799.
- [22] T. Kaluza, *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* 1921, 966 (1921); T. Kaluza, *Int. J. Mod. Phys. D* **27**(14), 1870001 (2018); O. Klein, *Z. Phys.* **37**, 895 (1926); 5. O. Klein, *Surv. High Energy Phys.* **5**, 241 (1986); T. Appelquist, A. Chodos, P.G.O. Freund, Reading, USA: Addison- Wesley (1987) 619 (Frontiers in physics, 65).
- [23] P. Van Nieuwenhuizen, *Phys. Rep.* **68**, 189 (1981)
- [24] J. Martin, J. Yokoyama, *JCAP* **0801**, 025 (2008); A. Maleknejad, M.M. Sheikh-Jabbari, J. Soda, *Phys. Rep.* **528**, 161. (2013). arXiv:1212.2921 [hep-th]
- [25] C. A. R. Herdeiro, E. Radu, N. Sanchis-Gual, and J. A. Font, *Phys. Rev. Lett.*, vol. 121, no. 10, p. 101102, 2018.
- [26] P. G. Fernandes, C. A. Herdeiro, A. M. Pombo, E. Radu, and N. Sanchis-Gual, *Classical and Quantum Gravity*, vol. 36, no. 13, p. 134002, 2019.
- [27] J. L. Blazquez-Salcedo, C. A. R. Herdeiro, J. Kunz, A. M. Pombo and E. Radu, *Phys. Lett. B* **806** (2020) 135493.
- [28] M. Cadoni, S. Mignemi and M. Serra, *Phys. Rev. D* **84**, 084046 (2011); M. Cadoni and E. Franzin, *Phys. Rev. D* **91**, 104011 (2015); Q. Wen, *Phys. Rev. D* **92**, 104002 (2015); M. Cadoni and E. Franzin, *Springer Proc. Phys.* **208**, 47, (2018), e-Print: 1510.02076.
- [29] R. Arnowitt, S. Deser, and C. W. Misner. The dynamics of general relativity. In L. Witten, editor, *Gravitation: An Introduction to Current Research*, pages 227-265. Wiley, 1962.
- [30] Charles W. Misner, Kip S. Thorne, and John A. Wheeler. *Gravitation*. W. H. Freeman, 1973.
- [31] Robert M. Wald. *General Relativity*. University of Chicago Press, 1984.
- [32] G. W. Gibbons, S. W. Hawking, *Phys. Rev. D* **15**, 2752 (1977).
- [33] J. D. Beckenstein, *Phys. Rev. D* **7**, 2333 (1973);
S. W. Hawking, *Nature*, (London) **248**, 30 (1974);
G. W. Gibbons and S. W. Hawking, *Phys. Rev. D* **15**, 2738 (1977).
- [34] C. J. Hunter, *Phys. Rev. D* **59**, 024009 (1999);
S. W. Hawking, C. J. Hunter and D. N. Page, *Phys. Rev. D* **59**, 044033 (1999);
R. B. Mann *Phys. Rev. D* **60**, 104047 (1999).

- [35] M. Cvetič and S. S. Gubser, *JHEP.* **04**, 024 (1999);
M. M. Caldarelli, G. Cognola and D. Klemm, *Class. Quantum Grav.* **17**, 399 (2000).
- [36] D. Kastor, S. Ray, and J. Traschen, *Class. Quantum Grav.* **26**, 195011 (2009);
D. Kubiznak and F. Simovic, *Class. Quantum Grav.* **33**, 245001 (2016).
- [37] N. Altamirano, D. Kubiznak, R. B. Mann, and Z. Sherkatghanad, *Class. Quantum Grav.* **31**, 042001 (2014).
- [38] N. Altamirano, D. Kubiznak, and R. B. Mann, *Phys. Rev. D* **88**, 101502 (2013).
- [39] A. M. Frassino, D. Kubiznak, R. B. Mann, and F. Simovic, *JHEP.* **2014**, 80 (2014);
B. P. Dolan, *Class. Quantum Grav.* **31**, 035022 (2014).
- [40] R. A. Hennigar, R. B. Mann, and E. Tjoa, *Phys. Rev. Lett.* **118**, 021301 (2017);
R. A. Hennigar, E. Tjoa, and R. B. Mann, *JHEP.* **2017**, 70 (2017);
H. Dykaar, R. A. Hennigar, and R. B. Mann, *JHEP.* **2017**, 45 (2017).
- [41] D. Kubiznak and R. B. Mann, *JHEP.* **2012**, 33 (2012).
- [42] S. Gunasekaran, R. B. Mann, D. Kubiznak, *JHEP.* **11**, 110 (2011).
D. Kubiznak, R. B. Mann, M. Teo, *Class. Quantum Grav.* **34**, 063001 (2017).