

# Crouzeix–Raviart and Raviart–Thomas finite-element error analysis on anisotropic meshes violating the maximum-angle condition

Hiroki Ishizaka · Kenta Kobayashi ·  
Takuya Tsuchiya

Received: date / Accepted: date

**Abstract** We investigate the piecewise linear nonconforming Crouzeix–Raviart and the lowest order Raviart–Thomas finite-element methods for the Poisson problem on three-dimensional anisotropic meshes. We first give error estimates of the Crouzeix–Raviart and the Raviart–Thomas finite-element approximate problems. We next present the equivalence between the Raviart–Thomas finite-element method and the enriched Crouzeix–Raviart finite-element method. We emphasise that we do not impose either shape-regular or maximum-angle condition during mesh partitioning. Numerical results confirm the results that we obtained.

**Keywords** Finite element · Raviart–Thomas · Crouzeix–Raviart ·  
Anisotropic meshes

## 1 Introduction

Let  $\Omega \subset \mathbb{R}^d$ ,  $d \in \{2, 3\}$ , be a bounded polyhedral domain. Furthermore, we assume that  $\Omega$  is convex if necessary. We consider the Poisson problem as follows. Find  $u : \Omega \rightarrow \mathbb{R}$  such that

$$-\Delta u = f \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega, \quad (1.1)$$

---

Hiroki Ishizaka  
Graduate School of Science and Engineering, Ehime University, Matsuyama, Japan  
E-mail: h.ishizaka005@gmail.com

Kenta Kobayashi  
Graduate School of Business Administration, Hitotsubashi University, Kunitachi, Japan  
E-mail: kenta.k@r.hit-u.ac.jp

Takuya Tsuchiya  
Graduate School of Science and Engineering, Ehime University, Matsuyama, Japan  
E-mail: tsuchiya@math.sci.ehime-u.ac.jp

where  $f \in L^2(\Omega)$  is a given function. This paper gives error estimates for the first-order Crouzeix–Raviart (CR) finite-element approximation on anisotropic meshes in three dimensions. Anisotropic meshes have different mesh sizes in different directions. The shape regularity assumption on triangulations  $\mathbb{T}_h$  is no longer valid on these meshes; see for example [2]. Furthermore, we do not impose the maximum-angle condition proposed in [4] during mesh partitioning. In many instances, the discussion also relates to two dimensions. We therefore discuss the problem here as uniformly valid in an arbitrary number of dimensions.

CR finite error estimates for the non-homogeneous Dirichlet Poisson problem are known. Let  $CR_{h0}^1$  be the CR finite-element space, to be defined in Section 2.3. Let  $u \in H_0^1(\Omega)$  and  $u_h^{CR} \in CR_{h0}^1$  be the exact and CR finite-element solutions, respectively. In [14, Corollary 2.2], adopting medius analysis, the estimate

$$|u - u_h^{CR}|_{H^1(\mathbb{T}_h)} \leq c_0 \left( \inf_{v_h \in CR_{h0}^1} |u - v_h|_{H^1(\mathbb{T}_h)} + Osc_1(f) \right), \quad (1.2)$$

is given, where  $|\cdot|_{H^1(\mathbb{T}_h)}$  denotes the broken (piecewise)  $H^1$ -semi norm defined in Section 2.2, and  $c_0$  a positive constant independent of  $h$ . Here, the oscillation  $Osc_1(f)$  is expressed as

$$Osc_1(f) := \left( \sum_{T \in \mathbb{T}_h} h_T^2 \left[ \inf_{\bar{f} \in \mathcal{P}^0(T)} \|f - \bar{f}\|_{L^2(T)}^2 \right] \right)^{1/2},$$

where  $\mathcal{P}^0(T)$  denotes the space of constant functions on  $T$ . Suppose that  $u \in H^2(\Omega)$  and oscillation  $Osc_1(f)$  vanishes. Let  $I_h u \in CR_{h0}^1$  be the nodal interpolation of  $u$  at the midpoints of the faces. Then, from the standard interpolation error estimate (see for example [11, Corollary 1.109]), we have

$$|u - u_h^{CR}|_{H^1(\mathbb{T}_h)} \leq c_0 |u - I_h u|_{H^1(\mathbb{T}_h)} \leq c_1 h |u|_{H^2(\Omega)},$$

where  $c_1$  represents a positive constant independent of  $h$  and  $u$  but depending on the parameter of the simplicial mesh; see for example [11, Definition 1.107]. This parameter is bounded if the simplicial mesh sequence is shape regular. However, the situation is different without the shape-regular condition. The aim of the present paper is to deduce an analogous error estimate on anisotropic finite-element meshes. Note that very flat elements might be included in the mesh sequence. In many papers reporting on such investigations, the maximum-angle condition instead of the shape-regular condition is imposed. However, the maximum-angle condition is not necessarily needed to obtain error estimates. Recently, in the two-dimensional instance, the CR finite-element analysis of the non-homogeneous Dirichlet–Poisson problem has been investigated under a more relaxed mesh condition, [19]. The present paper extends previous research to a three-dimensional setting.

However, it may not be easy to use the estimate (1.2) on anisotropic finite-element meshes. To overcome this difficulty, we use the interpolation error

estimates obtained in [16]. In that paper, the CR and Raviart–Thomas (RT) interpolation errors are bounded in terms of  $h$  and the new parameter  $H$ , see Corollary 2, 3.

The CR finite-element space is not in  $H_0^1(\Omega)$ . Hence, an error between the exact solution and the CR finite-element approximation solution with a  $H^1$ -broken seminorm is divided into two parts ([8, 11]). One is an approximation error that measures how well the exact solution is approximated by the CR finite-element functions, the other is a nonconformity error term. For the former, the CR interpolation error estimates (Corollary 2) are used. In the latter, the standard scaling argument is often used to obtain the error estimates. However, in this way, we are unable to derive the correct order on anisotropic meshes. To overcome this difficulty, we shall use the lowest-order RT interpolation error estimates on anisotropic meshes (Corollary 3). By this technique, we consequently have the error estimates in the  $H^1$ -broken seminorm (Theorem 6) and the  $L^2$  norm (Theorem 7) on anisotropic meshes.

Furthermore, we present an error estimate for the first-order RT finite-element approximation of the Poisson problem (1.1) based on the dual mixed formulation (Theorem 9). In the proof, we again use Corollary 3. We again emphasise that we do not impose either the shape-regular or the maximum-angle condition during mesh partitioning.

We next present the equivalence of the enriched piecewise linear CR finite-element method introduced by [15] and the first-order RT finite-element method. In two dimensions, the work [3] represents pioneering research. Marini [22] further found an expression relating RT and CR finite-element methods:

$$\bar{\sigma}_h^{RT}|_T = \nabla \bar{u}_h^{CR} - \frac{f_T^0}{2}(x - x_T) \quad \text{on } T, \quad (1.3)$$

where  $T$  denotes a mesh element,  $x_i$  ( $i = 1, 2, 3$ ) the vertices of triangle  $T$ ,  $x_T$  the barycentre of  $T$  such that  $x_T := \frac{1}{3}(x_1 + x_2 + x_3)$ , and  $\bar{\sigma}_h^{RT}$  and  $\bar{u}_h^{CR}$  respectively denote the RT and CR finite-element solutions with a given external piecewise-constant function  $f_T^0$ . It was recently proved [15] that the enriched piecewise-linear CR finite-element method is identical to the first-order RT finite-element method for both the Poisson and Stokes problems in any number of dimensions. In the present paper, we extend Marini's results to three dimensions (Lemma 10).

The remainder of the present paper is organised as follows. Section 2 introduces the weak form of the continuous problem (1.1), the finite-element meshes, and finite-element spaces. Furthermore, we propose a parameter  $H$ . Section 3 introduces discrete settings of the CR finite-element method for (1.1) and proposes error estimates. Section 4 proves error estimates for the first-order RT finite-element method based on the dual mixed formulation of the Poisson problem. Section 5 gives the equivalence of the RT and CR finite-element problems. Finally, Section 6 presents numerical results obtained using the Lagrange P1 element and the first-order CR element.

## 2 Preliminaries

### 2.1 Weak formulation

The variational formulation for the Poisson problem (1.1) is then as follows. Find  $u \in H_0^1(\Omega)$  such that

$$a_0(u, \varphi) = (f, \varphi) \quad \forall \varphi \in H_0^1(\Omega), \quad (2.1)$$

where  $a_0 : H^1(\Omega) \times H^1(\Omega) \rightarrow \mathbb{R}$  denotes a bilinear form defined by

$$a_0(u, \varphi) := (\nabla u, \nabla \varphi).$$

Here, we define  $H_0^1(\Omega)$  as the closure of  $C_0^\infty(\Omega)$  in the semi-norm  $|\cdot|_{H^1(\Omega)}$ . By the Lax–Milgram lemma, there exists a unique solution  $u \in H_0^1(\Omega)$  for any  $f \in L^2(\Omega)$  and it holds that

$$|u|_{H^1(\Omega)} \leq C_P(\Omega) \|f\|,$$

where  $C_P(\Omega)$  is the Poincaré constant depending on  $\Omega$ . Furthermore, if  $\Omega$  is convex, then  $u \in H^2(\Omega)$  and

$$|u|_{H^2(\Omega)} \leq \|\Delta u\|. \quad (2.2)$$

The proof can be found in, for example, [13, Theorem 3.1.1.2, Theorem 3.2.1.2].

### 2.2 Meshes, Mesh faces, Averages and Jumps

Let  $\mathbb{T}_h = \{T\}$  be a simplicial mesh of  $\overline{\Omega}$ , made up of closed  $d$ -simplices, such as

$$\overline{\Omega} = \bigcup_{T \in \mathbb{T}_h} T,$$

with  $h := \max_{T \in \mathbb{T}_h} h_T$ , where  $h_T := \text{diam}(T)$ . We assume that each face of any  $d$ -simplex  $T_1$  in  $\mathbb{T}_h$  is either a subset of the boundary  $\partial\Omega$  or a face of another  $d$ -simplex  $T_2$  in  $\mathbb{T}_h$ . That is,  $\mathbb{T}_h$  is a simplicial mesh of  $\overline{\Omega}$  without hanging nodes.

**Definition 1** For any  $T \in \mathbb{T}_h$ , we define the parameter  $H_T$  as

$$H_T := \frac{h_T^2}{|T|} \min_{1 \leq i \leq 3} |L_i| \quad \text{if } d = 2,$$

where  $L_i$  ( $i = 1, 2, 3$ ) denotes edges of the triangle  $T$ . Further, we define the parameter  $H_T$  as

$$H_T := \frac{h_T^2}{|T|} \min_{1 \leq i, j \leq 6, i \neq j} |L_i| |L_j| \quad \text{if } d = 3,$$

where  $L_i$  ( $i = 1, \dots, 6$ ) denotes edges of the tetrahedra  $T$ . Here,  $|T|$  denotes the measure of  $T$ . Furthermore, we set

$$H := H(h) := \max_{T \in \mathbb{T}_h} H_T.$$

We impose the following assumption.

**Assumption 1** *We assume that  $\{\mathbb{T}_h\}_{h>0}$  is a sequence of triangulations of  $\Omega$  such that*

$$\lim_{h \rightarrow 0} H(h) = 0.$$

**Remark 1** *The parameter  $H_T$  was introduced, and the interpolation errors are bounded (locally) in terms of  $h_T$  and  $H_T$  on anisotropic meshes without any geometric conditions in [16]. In two-dimensional case, the parameter  $H_T$  is equivalent to the circumradius of  $T$ . Hence, the maximum-angle condition or the semiregular condition holds if and only if there exists a constant  $\sigma_0$  such that  $H_T/h_T \leq \sigma_0$ . In three-dimensional case, it is conjectured that the maximum-angle condition holds if and only if the quantity  $H_T/h_T$  is bounded.*

We adopt the concepts of mesh faces, averages and jumps in the analysis of RT and CR finite element method. Let  $\mathcal{F}_h^i$  be the set of interior faces and  $\mathcal{F}_h^\partial$  the set of the faces on the boundary  $\partial\Omega$ . Let  $\mathcal{F}_h := \mathcal{F}_h^i \cup \mathcal{F}_h^\partial$ . For any  $F \in \mathcal{F}_h$ , we define the unit normal  $n_F$  to  $F$  as follows: (i) If  $F \in \mathcal{F}_h^i$  with  $F = T_1 \cap T_2$ ,  $T_1, T_2 \in \mathbb{T}_h$ , let  $n_1$  and  $n_2$  be the outward unit normals of  $T_1$  and  $T_2$ , respectively. Then,  $n_F$  is either of  $\{n_1, n_2\}$ ; (ii) If  $F \in \mathcal{F}_h^\partial$ ,  $n_F$  is the unit outward normal  $n$  to  $\partial\Omega$ .

Let  $k$  be a positive integer. We then define the broken (piecewise) Sobolev space as

$$H^k(\mathbb{T}_h) := \{\varphi \in L^2(\Omega); \varphi|_T \in H^k(T) \ \forall T \in \mathbb{T}_h\}$$

with the norm

$$|\varphi|_{H^1(\mathbb{T}_h)} := \left( \sum_{T \in \mathbb{T}_h} \|\nabla \varphi\|_{L^2(T)^d}^2 \right)^{1/2} \quad \varphi \in H^1(\mathbb{T}_h).$$

Let  $\varphi \in H^k(\mathbb{T}_h)$ . Suppose that  $F \in \mathcal{F}_h^i$  with  $F = T_1 \cap T_2$ ,  $T_1, T_2 \in \mathbb{T}_h$ . Set  $\varphi_1 := \varphi|_{T_1}$  and  $\varphi_2 := \varphi|_{T_2}$ . The jump and the average of  $\varphi$  across  $F$  is then defined as

$$[[\varphi]]_F := (\varphi_1 n_1 + \varphi_2 n_2) \cdot n_F, \quad \{\{\varphi\}\}_F := \frac{1}{2}(\varphi_1 + \varphi_2).$$

For a boundary face  $F \in \mathcal{F}_h^\partial$  with  $F = \partial T \cap \partial\Omega$ ,  $[[\varphi]]_F := \varphi|_T$  and  $\{\{\varphi\}\}_F := \varphi|_T$ . When  $v$  is an  $\mathbb{R}^d$ -valued function, we use the notation

$$[[v \cdot n]]_F := (v_1 - v_2) \cdot n_F, \quad \{\{v\}\}_F := \frac{1}{2}(v_1 + v_2)$$

for the jump of the normal component of  $v$ . For a boundary face  $F \in \mathcal{F}_h^\partial$  with  $F = \partial T \cap \partial\Omega$ ,  $[[v \cdot n]]_F := v|_T \cdot n$  and  $\{\{v\}\}_F := v|_T$ . Whenever no confusion can arise, we simply write  $[[\varphi]]$ ,  $\{\{\varphi\}\}$ ,  $[[v \cdot n]]$  and  $\{\{v\}\}$ , respectively.

Suppose that  $F \in \mathcal{F}_h^i$  with  $F = T_1 \cap T_2$ ,  $T_1, T_2 \in \mathbb{T}_h$ . For  $v \in H^1(\mathbb{T}_h)^d$  and  $\varphi \in H^1(\mathbb{T}_h)$ , it holds that

$$[(v\varphi) \cdot n]_F = \{\{v\}\}_F \cdot n_F [[\varphi]]_F + [[v \cdot n]]_F \{\{\varphi\}\}_F.$$

We here define a broken gradient operator as follows.

**Definition 2** For  $\varphi \in H^1(\mathbb{T}_h)$ , the broken gradient  $\nabla_h : H^1(\mathbb{T}_h) \rightarrow L^2(\Omega)^d$  is defined by

$$(\nabla_h \varphi)|_T := \nabla(\varphi|_T) \quad \forall T \in \mathbb{T}_h.$$

Note that  $H^1(\Omega) \subset H^1(\mathbb{T}_h)$  and the broken gradient coincides with the distributional gradient in  $H^1(\Omega)$ .

### 2.3 Finite Element Spaces and Interpolations Error Estimates

This section introduce the piecewise-constant, CR and RT finite element spaces.

Let  $T \in \mathbb{T}_h$ . For any  $k \in \mathbb{N}_0$ , let  $\mathcal{P}^k(T)$  be the space of polynomials with degree at most  $k$  in  $T$ .

**Theorem 1 (Poincaré inequality)** Let  $D \subset \mathbb{R}^d$  be a convex domain with diameter  $\text{diam}(D)$ . It then holds that, for  $\varphi \in H^1(D)$  with  $\int_D \varphi dx = 0$ ,

$$\|\varphi\|_{L^2(D)} \leq \frac{\text{diam}(D)}{\pi} |\varphi|_{H^1(D)}. \quad (2.3)$$

**Proof** The proof is found in [23, Theorem 3.2], also see [24].  $\square$

#### 2.3.1 Piecewise-constant finite element space

We define the standard piecewise constant space as

$$M_h^0 := \{q_h \in L^2(\Omega); q_h|_T \in \mathcal{P}^0(T) \quad \forall T \in \mathbb{T}_h\}.$$

The local interpolation  $\Pi_T^0$  from  $L^2(T)$  into the space  $\mathcal{P}^0(T)$  is defined by

$$\int_T (\Pi_T^0 q - q) dx = 0 \quad \forall q \in L^2(T).$$

Note that  $\Pi_T^0 q$  is the constant function equal to  $\frac{1}{|T|} \int_T q dx$ . We also define the global interpolation  $\Pi_h^0$  to the space  $M_h^0$  by

$$(\Pi_h^0 q)|_T = \Pi_T^0(q|_T) \quad \forall T \in \mathbb{T}_h, \quad \forall q \in L^2(\Omega).$$

The Poincaré inequality (2.3) directly yields the following error estimate of the local  $L^2$ -projection  $\Pi_T^0$ .

**Theorem 2** We have the error estimate of the local  $L^2$ -projection such that

$$\|\Pi_T^0 q - q\|_{L^2(T)} \leq \frac{h_T}{\pi} |q|_{H^1(T)} \quad \forall T \in \mathbb{T}_h, \quad \forall q \in H^1(T). \quad (2.4)$$

**Proof** For any  $q \in H^1(T)$ , we set  $w := \Pi_T^0 q - q$ . It then holds that

$$\int_T w dx = \int_T (\Pi_T^0 q - q) dx = \frac{1}{|T|} \int_T q dx |T| - \int_T q dx = 0.$$

Therefore, using the Poincaré inequality (2.3), we conclude (2.4).  $\square$

The global error estimate of the  $L^2$ -projection is obtained as follows.

**Corollary 1** *Let  $\{\mathbb{T}_h\}$  be a family of conformal meshes satisfying Assumption 1. It then holds that*

$$\|\Pi_h^0 q - q\|_{L^p(\Omega)} \leq \frac{h}{\pi} |q|_{H^1(\Omega)} \quad \forall q \in H^1(\Omega). \quad (2.5)$$

### 2.3.2 CR finite element space

We define the following CR finite element space as

$$CR_{h0}^1 := \left\{ \varphi_h \in L^2(\Omega); \varphi_h|_T \in \mathcal{P}^1(T) \quad \forall T \in \mathbb{T}_h, \int_F [[\varphi_h]]_F ds = 0 \quad \forall F \in \mathcal{F}_h \right\}.$$

Using the barycentric coordinates  $\lambda_i : \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $i = 1, \dots, d+1$ , we define the local basis functions as

$$\theta_i(x) := d \left( \frac{1}{d} - \lambda_i(x) \right), \quad 1 \leq i \leq d+1.$$

For  $i = 1, \dots, d+1$ , let  $F_i$  be the face of  $T$  and  $x_{F_i}$  the barycentre of the face  $F_i$ . We then define the local CR interpolation operator as

$$I_T^{CR} : H^1(T) \ni \varphi \mapsto I_T^{CR} \varphi := \sum_{i=1}^{d+1} \left( \frac{1}{|F_i|} \int_{F_i} \varphi ds \right) \theta_i \in \mathcal{P}^1.$$

Furthermore, it holds that

$$\frac{1}{|F_i|} \int_{F_i} (I_T^{CR} \varphi - \varphi) ds = 0, \quad i = 1, \dots, d+1, \quad \forall \varphi \in H^1(T).$$

We define the global CR interpolation  $I_h^{CR} : H_0^1(\Omega) \rightarrow CR_{h0}^1$  by

$$(I_h^{CR} \varphi)|_T = I_T^{CR}(\varphi|_T) \quad \forall T \in \mathbb{T}_h, \quad \forall \varphi \in H_0^1(\Omega).$$

We give the local CR interpolation error estimate.

**Theorem 3** *We have the following estimates such that for  $m \in \{0, 1\}$ ,*

$$|\varphi - I_T^{CR} \varphi|_{H^m(T)} \leq C_I^{CR,m} \left( \frac{H_T}{h_T} \right)^m h_T^{2-m} |\varphi|_{H^2(T)} \quad \forall T \in \mathbb{T}_h, \quad \forall \varphi \in H^2(T). \quad (2.6)$$

Here,  $C_I^{CR,m}$  is a positive constant independent of  $h_T$  and  $H_T$ .

**Proof** The proof is found in [16, Theorem 2].  $\square$

The global CR interpolation error estimates are obtained as follows.

**Corollary 2** *Let  $\{\mathbb{T}_h\}$  be a family of conformal meshes satisfying Assumption 1. Then, there exists constants  $C_G^{CR,0}, C_G^{CR,1} > 0$ , independent of  $H$  and  $h$ , such that*

$$\|\varphi - I_h^{CR} \varphi\|_{L^2(\Omega)} \leq C_G^{CR,0} h^2 |\varphi|_{H^2(\Omega)} \quad \forall \varphi \in H_0^1(\Omega) \cap H^2(\Omega), \quad (2.7)$$

$$|\varphi - I_h^{CR} \varphi|_{H^1(\mathbb{T}_h)} \leq C_G^{CR,1} H |\varphi|_{H^2(\Omega)} \quad \forall \varphi \in H_0^1(\Omega) \cap H^2(\Omega). \quad (2.8)$$

The inequality (2.6) with  $m = 1$  can be improved by replacing  $H_T$  with  $h_T$ . To this end, we use the Poincaré inequality (2.3).

**Theorem 4** *It then holds that*

$$|I_T^{CR} \varphi - \varphi|_{H^1(T)} \leq \frac{h_T}{\pi} |\varphi|_{H^2(T)} \quad \forall T \in \mathbb{T}_h, \quad \forall \varphi \in H^2(T). \quad (2.9)$$

**Proof** Let  $F_i, i = 1, \dots, d+1$  be the faces of the element  $T$ . We set  $\psi := I_T^{CR} \varphi - \varphi \in H^2(T)$ . From Green's formula and the property of the CR interpolation, we have

$$\int_T \frac{\partial \psi}{\partial x_j} dx = \int_{\partial T} \psi n_T^{(j)} ds = \sum_{i=1}^{d+1} n_T^{(j)} \int_{F_i} \psi ds = 0,$$

where  $n_T^{(j)}$  denotes the  $j$ th component of the outer unit normal vector  $n_T$ . From the Poincaré inequality (2.3), we have

$$\begin{aligned} |I_T^{CR} \varphi - \varphi|_{H^1(T)}^2 &= \sum_{j=1}^d \left\| \frac{\partial}{\partial x_j} (I_T^{CR} \varphi - \varphi) \right\|_{L^2(T)}^2 \\ &\leq \left( \frac{h_T}{\pi} \right)^2 \sum_{j=1}^d \left\| \frac{\partial}{\partial x_j} (I_T^{CR} \varphi - \varphi) \right\|_{H^1(T)}^2 \\ &= \left( \frac{h_T}{\pi} \right)^2 \sum_{j,k=1}^d \left\| \frac{\partial^2}{\partial x_j \partial x_k} (I_T^{CR} \varphi - \varphi) \right\|_{L^2(T)}^2 \\ &= \left( \frac{h_T}{\pi} \right)^2 |\varphi|_{H^2(T)}^2, \end{aligned}$$

which conclude (2.9).  $\square$

**Remark 2** *For  $i = 1, \dots, d+1$ , let  $x_{F_i}$  the barycentre of face  $F_i$ . If we choose the domain of the local CR interpolation operator as  $W^{\ell,p}(T) \subset C^0(T)$  with  $1 \leq p < \infty$  and  $d < \ell p$ , it is possible to define*

$$I_T^{CR,S} : W^{\ell,p}(T) \ni \varphi \mapsto I_T^{CR,S} \varphi := \sum_{i=1}^{d+1} \varphi(x_{F_i}) \theta_i \in \mathcal{P}^1.$$



However, the estimate [16, Theorem 2]

$$|I_T^{CR,S} \varphi - \varphi|_{H^1(T)} \leq C_I^{CR,1} H_T |\varphi|_{H^2(T)} \quad \forall \varphi \in H^2(T)$$

can not be improved by replacing  $H_T$  with  $h_T$ .

As a counter example, let us consider  $T$  with vertices  $x_1 := (0, 0, 0)^T$ ,  $x_2 := (h, 0, 0)^T$ ,  $x_3 := (\frac{h}{2}, h^\gamma, 0)^T$  and  $x_4 := (\frac{h}{2}, 0, \frac{h}{2})^T$ , where  $h := \frac{1}{N}$ ,  $N \in \mathbb{N}$  and  $\gamma \in \mathbb{R}$ ,  $1 < \gamma \leq 2$ . Let  $\varphi$  be a function such that

$$\varphi(x, y, z) := x^2 + y^2 + z^2.$$

If an exact solution  $\varphi$  is known, the error  $e_h := \varphi - \varphi_h$  and  $e_{h/2} := \varphi - \varphi_{h/2}$  are computed numerically for two mesh sizes  $h$  and  $h/2$ , where  $\varphi_h := I_T^{CR,S} \varphi$ . The convergence indicator  $r$  is defined by

$$r = \frac{1}{\log(2)} \log \left( \frac{\|e_h\|_X}{\|e_{h/2}\|_X} \right).$$

The parameter  $H_T$  is then  $H_T = \mathcal{O}(h^{2-\gamma})$ . We compute the convergence order with respect to the  $H_0^1$  norm defined by

$$Err_h^{CR,S}(H^1) := \frac{|\varphi - I_T^{CR,S} \varphi|_{H^1(T)}}{|\varphi|_{H^2(T)}},$$

for the case:  $\gamma = 1.5$  (Table 1).

**Table 1** Error of the local CR interpolation operator ( $\gamma = 1.5$ )

$N$	$h$	$H_T$	$Err_h^{CR,S}(H^1)$	$r$
128	7.8125e-03	3.8081e-01	2.8183e-03	
256	3.9062e-03	2.6723e-01	1.7641e-03	0.68
512	1.9531e-03	1.8823e-01	1.1587e-03	0.61
1024	9.7656e-04	1.3284e-01	7.8625e-04	0.60
2048	4.8828e-04	9.3842e-02	5.4390e-04	0.53
4096	2.4414e-04	6.6324e-02	3.8026e-04	0.52

### 2.3.3 RT finite element space

The lowest order RT finite element space is defined by

$$RT^0(T) := \{v; v(x) = p + xq, p \in \mathcal{P}^0(T)^d, q \in \mathcal{P}^0(T), x \in \mathbb{R}^d\}.$$

The functionals are defined by, for any  $v \in RT^0(T)$ ,

$$\chi_i(v) := \frac{1}{|F_i|} \int_{F_i} v \cdot n_i ds, \quad F_i \subset \partial T, \quad 1 \leq i \leq d+1,$$

where  $n_i$  denotes the outer unit normal vector of  $T$  along  $F_i$ . We set  $\Sigma := \{\chi_i\}_{i=1}^{d+1}$ . Note that  $\dim RT^0(T) = d + 1$ . The triple  $\{T, RT^0(T), \Sigma\}$  is then a finite element. We define the RT finite element space by

$$RT_h^0 := \{v_h \in L^2(\Omega)^d; v_h|_T \in RT^0(T), \forall T \in \mathbb{T}_h, [[v_h \cdot n]]_F = 0, \forall F \in \mathcal{F}_h^i\}.$$

Note that  $RT_h^0 \subset H(\operatorname{div}; \Omega) := \{v \in L^2(\Omega)^d; \operatorname{div} v \in L^2(\Omega)\}$ .

We next define the local RT interpolation as

$$I_T^{RT} : H^1(T)^d \rightarrow RT^0(T),$$

using

$$\int_{F_i} (v - I_T^{RT} v) \cdot n_i ds = 0, \quad F_i \subset \partial T, \quad i \in \{1, \dots, d+1\} \quad \forall v \in H^1(T)^d.$$

Further, we define the global RT interpolation  $I_h^{RT} : H^1(\Omega)^d \rightarrow RT_h^0$  by

$$(I_h^{RT} v)|_T = I_T^{RT}(v|_T) \quad \forall T \in \mathbb{T}_h, \quad \forall v \in H^1(\Omega)^d.$$

The local RT interpolation error estimate is as follows.

**Theorem 5** *We have the following estimates such that*

$$\|I_T^{RT} v - v\|_{L^2(T)^d} \leq C_I^{RT} H_T |v|_{H^1(T)^d} \quad \forall T \in \mathbb{T}_h, \quad \forall v \in H^1(T)^d. \quad (2.10)$$

Here,  $C_I^{RT}$  is a positive constant independent of  $H_T$ .

**Proof** The proof is found in [16, Theorem 3].  $\square$

The global RT interpolation error estimates are obtained as follows.

**Corollary 3** *Let  $\{\mathbb{T}_h\}$  be a family of conformal meshes satisfying Assumption 1. Then, there exists a constant  $C_G^{RT} > 0$ , independent of  $H$ , such that*

$$\|I_h^{RT} v - v\|_{L^2(\Omega)^d} \leq C_G^{RT} H |v|_{H^1(\Omega)^d} \quad \forall v \in H^1(\Omega)^d. \quad (2.11)$$

Between the RT interpolation  $I_h^{RT}$  and the  $L^2$ -projection  $\Pi_h^0$ , the following relation holds:

**Lemma 1** *For any  $v \in H^1(\Omega)^d$ , it holds that*

$$\operatorname{div}(I_h^{RT} v) = \Pi_h^0(\operatorname{div} v).$$

That is to say, the diagram

$$\begin{array}{ccc} H^1(\Omega)^d & \xrightarrow{\operatorname{div}} & L^2(\Omega) \\ I_h^{RT} \downarrow & & \downarrow \Pi_h^0 \\ RT_h^0 & \xrightarrow{\operatorname{div}} & M_h^0 \end{array}$$

commutes.

**Proof** The proof of this lemma is found in [7].  $\square$

The following relation plays an important role in the CR finite element analysis on anisotropic meshes.

**Lemma 2** *It holds that*

$$(v_h, \nabla_h \psi_h) + (\operatorname{div} v_h, \psi_h) = 0 \quad \forall v_h \in RT_h^0, \quad \forall \psi_h \in H_0^1(\Omega) + CR_{h0}^1. \quad (2.12)$$

**Proof** For any  $v_h \in RT_h^0$  and  $\psi_h \in H_0^1(\Omega) + CR_{h0}^1$ , using Green formula and the fact  $v_h \cdot n_F \in \mathcal{P}^0(F)$  for any  $F \in \mathcal{F}_h$ , we can derive

$$\begin{aligned} (v_h, \nabla_h \psi_h) + (\operatorname{div} v_h, \psi_h) &= \sum_{T \in \mathbb{T}_h} \int_{\partial T} (v_h \cdot n_T) \psi_h ds \\ &= \sum_{F \in \mathcal{F}_h} \int_F [[(v_h \psi_h) \cdot n_F]] ds \\ &= \sum_{F \in \mathcal{F}_h} \int_F ([v_h \cdot n_F] \{\{\psi_h\}\} + \{\{v_h\}\} \cdot n_F [[\psi_h]]) ds \\ &= 0. \end{aligned}$$

$\square$

## 2.4 Discrete Poincaré Inequality on Anisotropic Meshes

We propose the discrete Poincaré inequality on anisotropic meshes.

**Lemma 3 (Discrete Poincaré inequality on anisotropic meshes)** *Assume that  $\Omega$  is convex. If  $H \leq 1$ , there exists  $C(\Omega)$ , independent of  $h$ ,  $H$ , and the geometry of meshes, such that*

$$\|\varphi_h\| \leq C(\Omega) |\varphi_h|_{H^1(\mathbb{T}_h)} \quad \forall \varphi_h \in CR_{h0}^1. \quad (2.13)$$

**Proof** Let  $\varphi_h \in CR_{h0}^1$ . We consider the dual problem. Find  $z \in H^2(\Omega) \cap H_0^1(\Omega)$  such that

$$-\Delta z = \frac{\varphi_h}{\|\varphi_h\|} \quad \text{in } \Omega, \quad z = 0 \quad \text{on } \partial\Omega.$$

We then have a priori estimates:

$$|z|_{H^1(\Omega)} \leq C_P, \quad |z|_{H^2(\Omega)} \leq 1,$$

where  $C_P$  is the Poincaré constant. We use the duality argument to show the target inequality. That is to say, we have

$$\begin{aligned} \|\varphi_h\| &= \frac{1}{\|\varphi_h\|} (\varphi_h, \varphi_h) = (-\Delta z, \varphi_h) = (-\operatorname{div} \nabla z, \varphi_h) \\ &= (-\operatorname{div} \nabla z, \varphi_h - \Pi_h^0 \varphi_h) - (\nabla z - I_h^{RT}(\nabla z), \nabla_h \varphi_h) + (\nabla z, \nabla_h \varphi_h) \\ &\leq \|\Delta z\| \|\varphi_h - \Pi_h^0 \varphi_h\| + \|\nabla z - I_h^{RT}(\nabla z)\| |\varphi_h|_{H^1(\mathbb{T}_h)} + |z|_{H^1(\Omega)} |\varphi_h|_{H^1(\mathbb{T}_h)} \\ &\leq c(h + H|\nabla z|_{H^1(\Omega)} + C_P) |\varphi_h|_{H^1(\mathbb{T}_h)}, \end{aligned}$$

which leads to

$$\|\varphi_h\| \leq c(2 + C_p)|\varphi_h|_{H^1(\mathbb{T}_h)} \quad \text{if } H \leq 1.$$

We here used

$$\begin{aligned} - \int_{\Omega} \operatorname{div}(\nabla z) \varphi_h dx &= \int_{\Omega} (\Pi_h^0 \operatorname{div}(\nabla z) - \operatorname{div}(\nabla z)) \varphi_h dx - \int_{\Omega} (\Pi_h^0 \operatorname{div}(\nabla z)) \varphi_h dx \\ &= \int_{\Omega} (\Pi_h^0 \operatorname{div}(\nabla z) - \operatorname{div}(\nabla z)) (\varphi_h - \Pi_h^0 \varphi_h) dx \\ &\quad - \int_{\Omega} (\operatorname{div} I_h^{RT}(\nabla z)) \varphi_h dx \\ &= - \int_{\Omega} \operatorname{div}(\nabla z) (\varphi_h - \Pi_h^0 \varphi_h) dx \\ &\quad - \int_{\Omega} (\nabla z - I_h^{RT}(\nabla z)) \cdot \nabla_h \varphi_h dx + \int_{\Omega} \nabla z \cdot \nabla_h \varphi_h dx, \end{aligned}$$

where

$$\begin{aligned} \int_{\Omega} (\operatorname{div} I_h^{RT}(\nabla z)) \varphi_h dx &= \sum_{T \in \mathbb{T}_h} \int_{\partial T} n_T \cdot I_h^{RT}(\nabla z) \varphi_h ds - \int_{\Omega} I_h^{RT}(\nabla z) \cdot \nabla_h \varphi_h dx \\ &= \int_{\Omega} (\nabla z - I_h^{RT}(\nabla z)) \cdot \nabla_h \varphi_h dx - \int_{\Omega} \nabla z \cdot \nabla_h \varphi_h dx. \end{aligned}$$

□

### 3 CR Finite Element Approximation

#### 3.1 Finite Element Approximation

The CR finite element problem is to find  $u_h^{CR} \in CR_{h0}^1$  such that

$$a_{0h}(u_h^{CR}, \varphi_h) = (f, \varphi_h) \quad \forall \varphi_h \in CR_{h0}^1, \quad (3.1)$$

where  $a_{0h} : (CR_{h0}^1 + H_0^1(\Omega)) \times (CR_{h0}^1 + H_0^1(\Omega)) \rightarrow \mathbb{R}$  is defined by

$$a_{0h}(\psi_h, \varphi_h) := \sum_{T \in \mathbb{T}_h} \int_T \nabla \psi_h \cdot \nabla \varphi_h dx = (\nabla_h \psi_h, \nabla_h \varphi_h).$$

This problem is nonconforming because  $CR_{h0}^1 \not\subset H_0^1(\Omega)$ .

For the CR approximate solution  $u_h^{CR} \in CR_{h0}^1$  of (3.1), we have the a priori estimate, using (2.13),

$$|u_h^{CR}|_{H^1(\mathbb{T}_h)}^2 \leq \|f\| \|u_h^{CR}\| \leq C(\Omega) \|f\| |u_h^{CR}|_{H^1(\mathbb{T}_h)}.$$

By the Lax–Milgram lemma, there exists a unique solution  $u_h^{CR} \in CR_{h0}^1$  for any  $f \in L^2(\Omega)$ .

### 3.2 Classical Error Analysis

The starting point for error analysis is the Second Strang Lemma, e.g. see [11, Lemma 2.25],

$$|u - u_h^{CR}|_{H^1(\mathbb{T}_h)} \leq 2 \inf_{v_h \in CR_{h0}^1} |u - v_h|_{H^1(\mathbb{T}_h)} + \sup_{\varphi_h \in CR_{h0}^1} \frac{a_{0h}(u, \varphi_h) - (f, \varphi_h)}{|\varphi_h|_{H^1(\mathbb{T}_h)}}. \quad (3.2)$$

The first term of the inequality (3.2) is estimated as follows. Using the CR interpolation error estimate (2.8), we have, for any  $u \in H^2(\Omega)$ ,

$$\inf_{v_h \in CR_{h0}^1} |u - v_h|_{H^1(\mathbb{T}_h)} \leq |u - I_h^{CR}u|_{H^1(\mathbb{T}_h)} \leq cH|u|_{H^2(\Omega)}. \quad (3.3)$$

From the standard scaling argument, we have a consistency error inequality, e.g., see [11, Lemma 3.36].

**Lemma 4 (Asymptotic Consistency)** *Let  $u \in H_0^1(\Omega) \cap H^2(\Omega)$  be the solution of the homogeneous Dirichlet Poisson problem (1.1). It then holds that*

$$\frac{a_{0h}(u, \varphi_h) - (f, \varphi_h)}{|\varphi_h|_{H^1(\mathbb{T}_h)}} \leq c \left( \sum_{T \in \mathbb{T}_h} \frac{h_T^4}{(\min_{F \in \partial \mathbb{T}_h} \ell_F)^2} |u|_{H^2(T)}^2 \right)^{1/2} \quad \forall h, \quad \forall \varphi_h \in CR_{h0}^1, \quad (3.4)$$

where  $\partial \mathbb{T}_h$  denotes the set of all faces  $F$  of  $T \in \mathbb{T}_h$ . Here,  $\ell_F$  denotes the distance of the vertex of  $T$  opposite to  $F$  to the face.

**Proof** We follow [11, Lemma 3.36].

Let  $\varphi_h \in CR_{h0}^1$ . Because  $-\Delta u = f$ , we have

$$\begin{aligned} a_{0h}(u, \varphi_h) - (f, \varphi_h) &= \sum_{T \in \mathbb{T}_h} \int_T (\nabla u \cdot \nabla \varphi_h - f \varphi_h) dx \\ &= \sum_{T \in \mathbb{T}_h} \sum_{F \in \partial \mathbb{T}_h} \int_F (n_T \cdot \nabla) u \varphi_h ds. \end{aligned}$$

Because each face  $F$  of an element  $T$  located inside  $\Omega$  appears twice in the above sum, we have

$$a_{0h}(u, \varphi_h) - (f, \varphi_h) = \sum_{T \in \mathbb{T}_h} \sum_{F \in \partial \mathbb{T}_h} \int_F (n_T \cdot \nabla) u (\varphi_h - \overline{\varphi_h}) ds$$

with the mean value

$$\overline{\varphi_h} := \frac{1}{|F|} \int_F \varphi_h ds.$$

Furthermore, we get

$$a_{0h}(u, \varphi_h) - (f, \varphi_h) = \sum_{T \in \mathbb{T}_h} \sum_{F \in \partial \mathbb{T}_h} \int_F n_T \cdot (\nabla u - \overline{\nabla u}) (\varphi_h - \overline{\varphi_h}) ds$$

with the mean value

$$n_T \cdot \overline{\nabla u} := \frac{1}{|F|} \int_F (n_T \cdot \nabla) u ds.$$

The Cauchy–Schwarz inequality yields

$$a_{0h}(u, \varphi_h) - (f, \varphi_h) \leq \sum_{T \in \mathbb{T}_h} \sum_{F \in \partial \mathbb{T}_h} \|\nabla u - \overline{\nabla u}\|_{L^2(F)^d} \|\varphi_h - \overline{\varphi_h}\|_{L^2(F)}.$$

For  $F \in \partial \mathbb{T}_h$ , let  $\widehat{T} \subset \mathbb{R}^d$  be the reference simplex and let  $\Phi_T : \widehat{T} \rightarrow T$  be the corresponding affine transformation with Jacobian matrix  $A_T$ . Let  $\widehat{F} = \Phi_T^{-1}(F)$ . Using the standard scaling argument and the trace theorem on the reference element, we have

$$\|\varphi_h - \overline{\varphi_h}\|_{L^2(F)} \leq \left( \frac{|F|}{|\widehat{F}|} \right)^{1/2} \|\widehat{\varphi}_h - \overline{\widehat{\varphi}_h}\|_{L^2(\widehat{F})} \leq c \left( \frac{|F|}{|\widehat{F}|} \right)^{1/2} \|\widehat{\varphi}_h - \overline{\widehat{\varphi}_h}\|_{H^1(\widehat{T})}.$$

The Deny–Lions Lemma (see [11, Lemma B.67]) implies

$$\|\widehat{\varphi}_h - \overline{\widehat{\varphi}_h}\|_{H^1(\widehat{T})} \leq c |\widehat{\varphi}_h|_{H^1(\widehat{T})}.$$

Using the standard scaling argument again, we obtain

$$\begin{aligned} \|\varphi_h - \overline{\varphi_h}\|_{L^2(F)} &\leq c \left( \frac{|F|}{|\widehat{F}|} \right)^{1/2} |\widehat{\varphi}_h|_{H^1(\widehat{T})} \\ &\leq c \left( \frac{|F|}{|\widehat{F}|} \right)^{1/2} \|A_T\|_2 \left( \frac{|\widehat{T}|}{|T|} \right)^{1/2} |\varphi_h|_{H^1(T)} \\ &\leq c \left( \frac{|F|}{|T|} \right)^{1/2} h_T |\varphi_h|_{H^1(T)} = c \left( \frac{d}{\ell_F} \right)^{1/2} h_T |\varphi_h|_{H^1(T)}. \end{aligned}$$

Here,  $\|A_T\|_2$  denotes the matrix 2-norm as

$$\|A_T\|_2 := \sup_{0 \neq x \in \mathbb{R}^d} \frac{|A_T x|}{|x|},$$

where  $|x| := (\sum_{i=1}^d |x_i|^2)^{1/2}$  for  $x \in \mathbb{R}^d$ .

By analogous argument, we have

$$\|\nabla u - \overline{\nabla u}\|_{L^2(F)^d} \leq c \left( \frac{d}{\ell_F} \right)^{1/2} h_T |u|_{H^2(T)}.$$

We consequently get

$$\begin{aligned}
a_{0h}(u, \varphi_h) - (f, \varphi_h) &\leq c \sum_{T \in \mathbb{T}_h} \sum_{F \in \partial \mathbb{T}_h} \frac{h_T^2}{\ell_F} |u|_{H^2(T)} |\varphi_h|_{H^1(T)} \\
&\leq c \sum_{T \in \mathbb{T}_h} \frac{h_T^2}{\min_{F \in \partial \mathbb{T}_h} \ell_F} |u|_{H^2(T)} |\varphi_h|_{H^1(T)} \\
&\leq c \left( \sum_{T \in \mathbb{T}_h} \frac{h_T^4}{(\min_{F \in \partial \mathbb{T}_h} \ell_F)^2} |u|_{H^2(T)}^2 \sum_{T \in \mathbb{T}_h} |\varphi_h|_{H^1(T)}^2 \right)^{1/2},
\end{aligned}$$

which leads to (3.4).  $\square$

From (3.2), (3.3) and (3.4), we have

$$|u - u_h^{CR}|_{H^1(\mathbb{T}_h)} \leq cH|u|_{H^2(\Omega)} + c \left( \sum_{T \in \mathbb{T}_h} \frac{h_T^4}{(\min_{F \in \partial \mathbb{T}_h} \ell_F)^2} |u|_{H^2(T)}^2 \right)^{1/2}.$$

Since the order of the nonconforming term does not necessary becomes the order  $H$ , this inequality may be overestimated.

*Example: Let  $0 < h_T \leq 1$ . As examples, we consider two cases.*

(I) When we use meshes including the tetrahedra  $T$  with vertices  $(0, 0, 0)^T$ ,  $(h_T, 0, 0)^T$ ,  $(0, h_T, 0)^T$ , and  $(0, 0, h_T^\varepsilon)^T$ , we have

$$\frac{h_T^4}{(\min_{F \in \partial \mathbb{T}_h} \ell_F)^2} |u|_{H^2(T)}^2 \leq ch_T^{2(2-\varepsilon)} |u|_{H^2(T)}^2,$$

where  $1 < \varepsilon \leq 2$ . Since  $H = \mathcal{O}(h)$ , we get

$$|u - u_h^{CR}|_{H^1(\mathbb{T}_h)} \leq c(h + h^{2-\varepsilon}) |u|_{H^2(\Omega)}.$$

(II) When we use meshes including the tetrahedra  $T$  with vertices  $(0, 0, 0)^T$ ,  $(h_T, 0, 0)^T$ ,  $(0, h_T, 0)^T$ , and  $(h_T^\gamma, 0, h_T^\varepsilon)^T$ , we have

$$\frac{h_T^4}{(\min_{F \in \partial \mathbb{T}_h} \ell_F)^2} |u|_{H^2(T)}^2 \leq ch_T^{2(2-\varepsilon)} |u|_{H^2(T)}^2,$$

where  $1 < \gamma < \varepsilon \leq 1 + \gamma$  and  $1 < \varepsilon \leq 2$ . Since  $H = \mathcal{O}(h^{1+\gamma-\varepsilon})$ , we get

$$|u - u_h^{CR}|_{H^1(\mathbb{T}_h)} \leq c(h^{1+\gamma-\varepsilon} + h^{2-\varepsilon}) |u|_{H^2(\Omega)}.$$

### 3.3 Argument via the RT Interpolation Error

To overcome the difficulty, we use the relation (2.12) in Lemma 2, e.g., see also [1, 21].

**Lemma 5 (Asymptotic Consistency)** *We assume that  $\Omega$  is convex. Let  $\{\mathbb{T}_h\}$  be a family of conformal meshes satisfying Assumption 1. Let  $u \in H_0^1(\Omega) \cap H^2(\Omega)$  be the solution of the homogeneous Dirichlet Poisson problem (1.1). Then, there exists  $c$ , independent of  $H$ , such that*

$$\sup_{\varphi_h \in CR_{h0}^1} \frac{a_{0h}(u, \varphi_h) - (f, \varphi_h)}{|\varphi_h|_{H^1(\mathbb{T}_h)}} \leq cH\|f\|. \quad (3.5)$$

**Proof** Using (2.12), we have, for any  $w_h \in RT_h^0$ ,

$$\sup_{\varphi_h \in CR_{h0}^1} \frac{a_{0h}(u, \varphi_h) - (f, \varphi_h)}{|\varphi_h|_{H^1(\mathbb{T}_h)}} = \sup_{\varphi_h \in CR_{h0}^1} \frac{(\nabla u - w_h, \nabla_h \varphi_h) - (\operatorname{div} w_h + f, \varphi_h)}{|\varphi_h|_{H^1(\mathbb{T}_h)}}.$$

We set  $w_h := I_h^{RT} \nabla u$ . From Lemma 1, we get

$$\operatorname{div}(I_h^{RT} \nabla u) = \Pi_h^0 \operatorname{div}(\nabla u) = -\Pi_h^0 f.$$

Furthermore, we have, for any  $\varphi_h \in CR_{h0}^1$ ,

$$(-\Pi_h^0 f + f, \Pi_h^0 \varphi_h) = 0.$$

We thus obtain

$$\begin{aligned} & (\nabla u - I_h^{RT} \nabla u, \nabla_h \varphi_h) - (-\Pi_h^0 f + f, \varphi_h) \\ &= (\nabla u - I_h^{RT} \nabla u, \nabla_h \varphi_h) - (-\Pi_h^0 f + f, \varphi_h - \Pi_h^0 \varphi_h) \\ &\leq \|\nabla u - I_h^{RT} \nabla u\|_{L^2(\Omega)^d} |\varphi_h|_{H^1(\mathbb{T}_h)} + \|f - \Pi_h^0 f\| \|\varphi_h - \Pi_h^0 \varphi_h\| \\ &\leq cH|u|_{H^2(\Omega)} |\varphi_h|_{H^1(\mathbb{T}_h)} + ch\|f\| |\varphi_h|_{H^1(\mathbb{T}_h)}. \end{aligned}$$

□

We consequently obtain the error estimate of the CR finite element method on anisotropic meshes.

**Theorem 6** *We assume that  $\Omega$  is convex. Let  $\{\mathbb{T}_h\}$  be a family of conformal meshes satisfying Assumption 1. Let  $u \in H_0^1(\Omega) \cap H^2(\Omega)$  be the solution of the homogeneous Dirichlet Poisson problem (1.1) with data  $f \in L^2(\Omega)$ . Let  $u_h^{CR} \in CR_{h0}^1$  be the approximate solution of (3.1). Then, there exists  $c$ , independent of  $H$ , such that*

$$|u - u_h^{CR}|_{H^1(\mathbb{T}_h)} \leq cH\|f\|. \quad (3.6)$$



**Proof** Using (3.2), (2.8) and (3.5), we have

$$\begin{aligned} |u - u_h^{CR}|_{H^1(\mathbb{T}_h)} &\leq 2 \inf_{v_h \in CR_{h0}^1} |u - v_h|_{H^1(\mathbb{T}_h)} + \sup_{\varphi_h \in CR_{h0}^1} \frac{a_{0h}(u, \varphi_h) - (f, \varphi_h)}{|\varphi_h|_{H^1(\mathbb{T}_h)}} \\ &\leq 2|u - I_h^{CR}u|_{H^1(\mathbb{T}_h)} + cH\|f\| \leq cH\|f\|, \end{aligned}$$

which leads to the estimate (3.6).  $\square$

We next give the  $L^2$  error estimate of the CR finite element method on anisotropic meshes, see also [20, 21, 9].

**Theorem 7** *We assume that  $\Omega$  is convex. Let  $\{\mathbb{T}_h\}$  be a family of conformal meshes satisfying Assumption 1. Let  $u \in H_0^1(\Omega) \cap H^2(\Omega)$  be the solution of the homogeneous Dirichlet Poisson problem (1.1) with data  $f \in L^2(\Omega)$ . Let  $u_h^{CR} \in CR_{h0}^1$  be the approximate solution of (3.1). Then, there exists  $c$ , independent of  $H$ , such that*

$$\|u - u_h^{CR}\| \leq cH^2\|f\|. \quad (3.7)$$

**Proof** We set  $e_h := u - u_h^{CR}$ . Let  $z \in H^2(\Omega) \cap H_0^1(\Omega)$  satisfy

$$a_0(\varphi, z) = (\varphi, e_h) \quad \forall \varphi \in H_0^1(\Omega) \quad (3.8)$$

and  $z_h^{CR} \in CR_{h0}^1$  satisfy

$$a_{0h}(\varphi_h, z_h^{CR}) = (\varphi_h, e_h) \quad \forall \varphi_h \in CR_{h0}^1. \quad (3.9)$$

We then have

$$\begin{aligned} \|e_h\|^2 &= (e_h, e_h) = a_{0h}(u, z) - a_{0h}(u_h^{CR}, z_h^{CR}) \\ &= a_{0h}(u - u_h^{CR}, z - z_h^{CR}) + a_{0h}(u - u_h^{CR}, z_h^{CR}) + a_{0h}(u_h^{CR}, z - z_h^{CR}) \\ &= a_{0h}(u - u_h^{CR}, z - z_h^{CR}) \\ &\quad + a_{0h}(u - u_h^{CR}, z_h^{CR} - I_h^{CR}z) + a_{0h}(u - u_h^{CR}, I_h^{CR}z) \\ &\quad + a_{0h}(u_h^{CR} - I_h^{CR}u, z - z_h^{CR}) + a_{0h}(I_h^{CR}u, z - z_h^{CR}). \end{aligned} \quad (3.10)$$

Using Theorem 6, the first term on the right hand side of (3.10) can be estimated as

$$\begin{aligned} a_{0h}(u - u_h^{CR}, z - z_h^{CR}) &\leq |u - u_h^{CR}|_{H^1(\mathbb{T}_h)} |z - z_h^{CR}|_{H^1(\mathbb{T}_h)} \\ &\leq cH^2\|f\|\|e_h\|. \end{aligned} \quad (3.11)$$

For the second and fourth terms on the right hand side of (3.10), we have

$$\begin{aligned} &a_{0h}(u - u_h^{CR}, z_h^{CR} - I_h^{CR}z) \\ &= a_{0h}(u - u_h^{CR}, z_h^{CR} - z) + a_{0h}(u - u_h^{CR}, z - I_h^{CR}z) \\ &\leq |u - u_h^{CR}|_{H^1(\mathbb{T}_h)} (|z_h^{CR} - z|_{H^1(\mathbb{T}_h)} + |z - I_h^{CR}z|_{H^1(\mathbb{T}_h)}) \\ &\leq cH^2\|f\|\|e_h\|, \end{aligned} \quad (3.12)$$

and, analogously,

$$a_{0h}(u_h^{CR} - I_h^{CR}u, z - z_h^{CR}) \leq cH^2\|f\|\|e_h\|. \quad (3.13)$$

From (3.8), (3.9) and (2.12), we have

$$\begin{aligned} a_{0h}(u - u_h^{CR}, I_h^{CR}z) &= a_{0h}(u, I_h^{CR}z) - a_{0h}(u_h^{CR}, I_h^{CR}z) = (\nabla u, \nabla_h I_h^{CR}z) - (f, I_h^{CR}z) \\ &= (\nabla u, \nabla_h I_h^{CR}z - \nabla z) - (f, I_h^{CR}z - z) + (\nabla u, \nabla z) - (f, z) \\ &= (\nabla u - I_h^{RT}\nabla u, \nabla_h I_h^{CR}z - \nabla z) - (f + \operatorname{div}(I_h^{RT}\nabla u), I_h^{CR}z - z). \end{aligned}$$

From Lemma 1 and  $\operatorname{div}(I_h^{RT}\nabla u) = -\Pi_h^0 f$ , we have

$$\begin{aligned} a_{0h}(u - u_h^{CR}, I_h^{CR}z) &= (\nabla u - I_h^{RT}\nabla u, \nabla_h I_h^{CR}z - \nabla z) - (f - \Pi_h^0 f, I_h^{CR}z - z) \\ &\leq \|\nabla u - I_h^{RT}\nabla u\|_{L^2(\Omega)^d} \|I_h^{CR}z - z\|_{H^1(\mathbb{T}_h)} + \|f - \Pi_h^0 f\| \|I_h^{CR}z - z\| \\ &\leq cH^2\|f\|\|e_h\|. \end{aligned} \quad (3.14)$$

Analogously, from  $\operatorname{div}(I_h^{RT}\nabla z) = -\Pi_h^0 e_h$ , we have

$$\begin{aligned} a_{0h}(I_h^{CR}u, z - z_h^{CR}) &= (\nabla_h I_h^{CR}u - \nabla u, \nabla z - I_h^{RT}\nabla z) - (I_h^{CR}u - u, e_h + \operatorname{div}(I_h^{RT}\nabla z)) \\ &\leq |I_h^{CR}u - u|_{H^1(\mathbb{T}_h)} \|\nabla z - I_h^{RT}\nabla z\|_{L^2(\Omega)^d} + \|I_h^{CR}u - u\| \|e_h - \Pi_h^0 e_h\| \\ &\leq cH^2\|f\|\|e_h\|. \end{aligned} \quad (3.15)$$

Combining (3.10), (3.11), (3.12), (3.13), (3.14), and (3.15), we finally get

$$\|e_h\|^2 \leq cH^2\|f\|\|e_h\|,$$

which leads to the target estimate.  $\square$

## 4 RT Finite Element Error Estimates

### 4.1 Dual mixed formulation of the Poisson problem

The Poisson equation (1.1)  $-\Delta u = -\operatorname{div} \nabla u = f$  can be written as the following system. Find  $(\sigma, u) : \Omega \rightarrow \mathbb{R}^d \times \mathbb{R}$  such that

$$\sigma - \nabla u = 0 \quad \text{in } \Omega, \quad (4.1a)$$

$$\operatorname{div} \sigma = -f \quad \text{in } \Omega, \quad (4.1b)$$

$$u = 0 \quad \text{on } \partial\Omega. \quad (4.1c)$$

We consider the following dual mixed formulation: Find  $(\sigma, u) \in H(\operatorname{div}; \Omega) \times L^2(\Omega)$  such that

$$a(\sigma, v) + b(v, u) = 0 \quad \forall v \in H(\operatorname{div}; \Omega), \quad (4.2a)$$

$$b(\sigma, q) = -(f, q) \quad \forall q \in L^2(\Omega), \quad (4.2b)$$

where bilinear forms  $a : H(\operatorname{div}; \Omega) \times H(\operatorname{div}; \Omega) \rightarrow \mathbb{R}$  and  $b : H(\operatorname{div}; \Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$  are defined by

$$a(\sigma, v) := (\sigma, v), \quad b(v, q) := (\operatorname{div} v, q).$$

We set  $X_0 := \{v \in H(\operatorname{div}; \Omega); b(v, q) = 0 \quad \forall q \in L^2(\Omega)\}$ . Because there exists a constant  $c > 0$  such that

$$a(v, v) \geq c \|v\|_{H(\operatorname{div}; \Omega)}^2 \quad \forall v \in X_0$$

and the bilinear form  $b(\cdot, \cdot)$  satisfies the inf-sup condition

$$\inf_{0 \neq q \in L^2(\Omega)} \sup_{0 \neq v \in H(\operatorname{div}; \Omega)} \frac{b(v, q)}{\|v\|_{H(\operatorname{div}; \Omega)} \|q\|} \geq \beta_* > 0, \quad (4.3)$$

(4.2) is uniquely solvable; e.g., see [12, 6].

## 4.2 RT Approximate Problem

We consider the following RT approximate problem. Find  $(\sigma_h^{RT}, u_h^{RT}) \in RT_h^0 \times M_h^0$  such that

$$a(\sigma_h^{RT}, v_h) + b(v_h, u_h^{RT}) = 0, \quad \forall v_h \in RT_h^0, \quad (4.4a)$$

$$b(\sigma_h^{RT}, q_h) = -(f, q_h), \quad \forall q_h \in M_h^0. \quad (4.4b)$$

This setting is conforming because  $RT_h^0 \times M_h^0 \subset H(\operatorname{div}; \Omega) \times L^2(\Omega)$ . It is given later that the discrete inf-sup condition

$$\inf_{q_h \in M_h^0} \sup_{v_h \in RT_h^0} \frac{b(v_h, q_h)}{\|v_h\|_{H(\operatorname{div}; \Omega)} \|q_h\|} \geq c_* > 0$$

holds, where  $c_*$  is a constant independent of  $h$ .

## 4.3 Error Estimates of the RT Finite Element Approximation

This section gives error estimates of the mixed finite element approximation (4.4). We emphasise that we do not impose the shape regularity condition and the maximum-angle condition for the mesh partition. That is, we assume that  $\{\mathbb{T}_h\}$  is a family of conformal meshes satisfying Assumption 1.

**Lemma 6** *Let  $D \subset \mathbb{R}^d$  be a bounded domain. For any  $g \in L^2(D)$ , there exists  $v \in H^1(D)^d$  such that*

$$\operatorname{div} v = g \quad \text{in } D \quad (4.5)$$

and

$$\|v\|_{H^1(D)^d} \leq \|g\|_{L^2(D)}, \quad \|v\|_{L^2(\Omega)^d} \leq C_P(D) \|g\|_{L^2(D)}, \quad (4.6)$$

where  $C_P(D)$  is the Poincaré constant.

**Proof** The proof can be found in [5, Lemma 2.2].  $\square$

We next give the discrete inf-sup condition.

**Lemma 7 (Discrete inf-sup condition)** *If  $C_G^{RT} H \leq 1$ , there exists a constant  $c_*$ , depending only on the Poincaré constant, such that*

$$\inf_{q_h \in M_h^0} \sup_{v_h \in RT_h^0} \frac{b(v_h, q_h)}{\|v_h\|_{H(\operatorname{div}; \Omega)} \|q_h\|} \geq c_* > 0, \quad (4.7)$$

where  $C_G^{RT}$  is the constant appearing in Corollary 3.

**Proof** Let  $q_h \in M_h^0$ . From Lemma 6, there exists  $v \in H^1(\Omega)^d$  such that  $\operatorname{div} v = q_h$  in  $\Omega$ ,  $\|v\|_{H^1(\Omega)^d} \leq \|q_h\|$ , and  $\|v\|_{L^2(\Omega)^d} \leq C_P(\Omega) \|q_h\|$ .

By the Gauss theorem, we have

$$\sum_{T \in \mathbb{T}_h} \int_{\partial T} v \cdot n_T ds = \sum_{T \in \mathbb{T}_h} \int_T \operatorname{div} v dx = \int_{\Omega} q_h dx.$$

From the definition of the Raviart–Thomas interpolation, we conclude that

$$\begin{aligned} \int_{\Omega} \operatorname{div}(I_T^{RT} v) p_h dx &= \sum_{T \in \mathbb{T}_h} p_h \int_T \operatorname{div}(I_T^{RT} v) dx = \sum_{T \in \mathbb{T}_h} p_h \int_{\partial T} n_T \cdot (I_T^{RT} v) ds \\ &= \sum_{T \in \mathbb{T}_h} p_h \int_{\partial T} v \cdot n_T ds = \int_{\Omega} q_h p_h dx \quad \forall p_h \in M_h^0. \end{aligned}$$

Therefore, it follows that  $\operatorname{div}(I_h^{RT} v) = q_h$ .

From the definitions, we have

$$\begin{aligned} \|I_h^{RT} v\|_{H(\operatorname{div}; \Omega)}^2 &= \|I_h^{RT} v\|_{L^2(\Omega)^d}^2 + \|\operatorname{div}(I_h^{RT} v)\|^2 \\ &\leq 2\|I_h^{RT} v - v\|_{L^2(\Omega)^d}^2 + 2\|v\|_{L^2(\Omega)^d}^2 + \|q_h\|^2 \\ &\leq 2(C_G^{RT})^2 H^2 \|v\|_{H^1(\Omega)^d}^2 + 2C_P(\Omega)^2 \|q_h\|^2 + \|q_h\|^2 \\ &\leq (3 + 2C_P(\Omega)^2) \|q_h\|^2. \end{aligned}$$

We thus have

$$\sup_{v_h \in RT_h^0} \frac{b(v_h, q_h)}{\|v_h\|_{H(\operatorname{div}; \Omega)}} \geq \frac{b(I_h^{RT} v, q_h)}{\|I_h^{RT} v\|_{H(\operatorname{div}; \Omega)}} \geq \frac{1}{(3 + 2C_P(\Omega)^2)^{1/2}} \frac{(q_h, q_h)}{\|q_h\|},$$

and the proof of (4.7) is completed with  $c_* := (3 + 2C_P(\Omega)^2)^{-1/2}$ .  $\square$

From the discrete equations (4.4) and their continuous counterpart (4.2), we obtain the Galerkin orthogonality

$$a(\sigma - \sigma_h^{RT}, v_h) + b(v_h, u - u_h^{RT}) = 0 \quad \forall v_h \in RT_h^0, \quad (4.8a)$$

$$b(\sigma - \sigma_h^{RT}, q_h) = 0 \quad \forall q_h \in M_h^0. \quad (4.8b)$$

We then get the following Céa-lemma-type estimates with the help of (4.8) and the inf-sup condition (4.7).

**Theorem 8** *Let  $\sigma \in H^1(\Omega)^d$  and  $\sigma_h^{RT} \in RT_h^0$  be the solutions of (4.1) and (4.4), respectively. We then have*

$$\|\sigma - \sigma_h^{RT}\|_{L^2(\Omega)^d} \leq \|\sigma - I_h^{RT} \sigma\|_{L^2(\Omega)^d}. \quad (4.9)$$

Furthermore, let  $(\sigma, u) \in H^1(\Omega)^d \times L^2(\Omega)$  and  $(\sigma_h^{RT}, u_h^{RT}) \in RT_h^0 \times M_h^0$  be the solutions of (4.1) and (4.4), respectively. Then, if  $C_G^{RT} H \leq 1$ , it holds that

$$\|u - u_h^{RT}\| \leq \|u - \Pi_h^0 u\| + c_*^{-1} \|\sigma - \sigma_h^{RT}\|_{L^2(\Omega)^d}. \quad (4.10)$$

Here,  $C_G^{RT}$  and  $c_*$  are respectively the constants appearing in Corollary 3 and Lemma 7.

**Proof** The proof can be found in [5, Lemma 3.7, Lemma 3.9]. □

Using Theorem 8 and the interpolation error estimates of Corollary 1 and 3, we thus have the error estimates of the mixed finite element approximation (4.4) on anisotropic meshes violating the maximum-angle condition.

**Theorem 9** *let  $(\sigma, u) \in H^1(\Omega)^d \times H^1(\Omega)$  and  $(\sigma_h^{RT}, u_h^{RT}) \in RT_h^0 \times M_h^0$  be the solutions of (4.1) and (4.4), respectively. Then, there exists a constant  $c_1 > 0$ , independent of  $\sigma$ ,  $H$ , and the geometric properties of  $\mathbb{T}_h$ , such that*

$$\|\sigma - \sigma_h^{RT}\|_{L^2(\Omega)^d} \leq c_1 H |\sigma|_{H^1(\Omega)^d}. \quad (4.11)$$

Furthermore, if  $C_G^{RT} H \leq 1$ , there exists a constant  $c_2 > 0$ , depending on the discrete inf-sup condition but independent of  $\sigma$ ,  $u$ ,  $h$ ,  $H$ , and the geometric properties of  $\mathbb{T}_h$

$$\|u - u_h^{RT}\| \leq c_2 (h|u|_{H^1(\Omega)} + H|\sigma|_{H^1(\Omega)^d}). \quad (4.12)$$

Here,  $C_G^{RT}$  is the constant appearing in Corollary 3.




Fig. 1 Tetrahedron

## 5 Relationship between the RT and CR Finite Element Approximation

This section shows the relationship between the RT and CR problems. Find  $(\bar{\sigma}_h^{RT}, \bar{u}_h^{RT}) \in RT_h^0 \times M_h^0$  such that

$$a(\bar{\sigma}_h^{RT}, v_h) + b(v_h, \bar{u}_h^{RT}) = 0 \quad \forall v_h \in RT_h^0, \quad (5.1a)$$

$$b(\bar{\sigma}_h^{RT}, q_h) = -(\Pi_h^0 f, q_h) \quad \forall q_h \in M_h^0 \quad (5.1b)$$

and find  $\bar{u}_h^{CR} \in CR_{h0}^1$  such that

$$a_{0h}(\bar{u}_h^{CR}, \varphi_h) = (\Pi_h^0 f, \varphi_h) \quad \forall \varphi_h \in CR_{h0}^1. \quad (5.2)$$

Here, (5.2) is the CR approximation of the Poisson equation

$$-\Delta \bar{u} = \Pi_h^0 f \quad \text{in } \Omega, \quad \bar{u} = 0 \quad \text{on } \partial\Omega. \quad (5.3)$$

In the case of  $d = 2$ , it is well known that there exists a relationship between  $(\bar{\sigma}_h^{RT}, \bar{u}_h^{RT})$  and  $\bar{u}_h^{CR}$  introduced by Marini; for example, [22]. See also [20, 18, 21]. We here show the relation in the three dimensional case.

Let us consider a tetrahedron  $T \subset \mathbb{R}^3$  such as that in Figure 1. Let  $x_i$  ( $i = 1, 2, 3, 4$ ) be the vertices and  $m_{i,j}$  the midpoints of edges of the tetrahedron; that is,  $m_{i,j} := \frac{1}{2}(x_i + x_j)$ . Furthermore, for  $1 \leq i \leq 4$ , let  $F_i$  be the face of the tetrahedron opposite  $x_i$ . Then, by simple calculation, we find the equality

$$L := \sum_{i=1}^4 |x_i - x_T|^2 = |m_{1,4} - m_{2,3}|^2 + |m_{1,3} - m_{2,4}|^2 + |m_{1,2} - m_{3,4}|^2,$$

holds, where  $x_T$  is the barycentre of  $T$  such that  $x_T := \frac{1}{4} \sum_{i=1}^4 x_i$ .

We present a quadrature scheme over a simplex  $T \subset \mathbb{R}^3$  (e.g., [25, p.307]) that is easily conformed.

**Lemma 8** For any  $f \in \mathcal{C}^0(T)$ , the quadrature scheme

$$\int_T f(x)dx \sim -\frac{|T|}{20} \sum_{i=1}^4 f(x_i) + \frac{|T|}{5} \sum_{1 \leq i < j \leq 4} f(m_{i,j})$$

is exact for polynomials of degree less than or equal to 2;

$$\int_T f(x)dx + \frac{|T|}{20} \sum_{i=1}^4 f(x_i) - \frac{|T|}{5} \sum_{1 \leq i < j \leq 4} f(m_{i,j}) = 0 \quad \forall f \in \mathcal{P}^2(T). \quad (5.4)$$

Define the function  $\varphi_T$  by

$$\varphi_T(x) := \begin{cases} L - 12|x - x_T|^2, & \text{on } T, \\ 0, & \text{otherwise.} \end{cases} \quad (5.5)$$

We then have the following lemma.

**Lemma 9** It holds that

$$\frac{1}{|F_i|} \int_{F_i} \varphi_T(x)ds = 0, \quad i = 1, 2, 3, 4, \quad (5.6)$$

$$\frac{1}{|T|} \int_T \varphi_T(x)dx = \frac{2}{5}L, \quad (5.7)$$

$$\frac{1}{|T|} \int_T |\nabla \varphi_T(x)|^2 dx = \frac{144}{5}L. \quad (5.8)$$

**Proof** From second-order three-point numerical integration over  $F_1$ ,

$$\int_{F_1} f(x)ds = \frac{|F_1|}{3} (f(m_{2,3}) + f(m_{3,4}) + f(m_{2,4})) \quad \forall f \in \mathcal{P}^2(T),$$

we have

$$\begin{aligned} & \frac{1}{|F_1|} \int_{F_1} \varphi_T(x)ds \\ &= \frac{1}{3} (\varphi_T(m_{2,3}) + \varphi_T(m_{3,4}) + \varphi_T(m_{2,4})) \\ &= \frac{1}{3} (3L - 12(|m_{2,3} - x_T|^2 + |m_{3,4} - x_T|^2 + |m_{2,4} - x_T|^2)) \\ &= \frac{1}{3} \left( 3L - \frac{12}{4} (|m_{2,3} - m_{1,4}|^2 + |m_{3,4} - m_{1,2}|^2 + |m_{2,4} - m_{1,3}|^2) \right) = 0, \end{aligned}$$

which leads to (5.6).

Next, using (5.4), we have

$$\begin{aligned}
& \frac{1}{|T|} \int_T \varphi_T(x) dx \\
&= -\frac{1}{20} \sum_{i=1}^4 \varphi_T(x_i) + \frac{1}{5} \sum_{1 \leq i < j \leq 4} \varphi_T(m_{i,j}) \\
&= -\frac{1}{20} \left( 4L - 12 \sum_{i=1}^4 |x_i - x_T|^2 \right) + \frac{1}{5} \left( 6L - 12 \sum_{1 \leq i < j \leq 4} |m_{i,j} - x_T|^2 \right) \\
&= \frac{2}{5} L,
\end{aligned}$$

which leads to (5.7). We here used

$$\begin{aligned}
& \sum_{1 \leq i < j \leq 4} |m_{i,j} - x_T|^2 \\
&= |m_{1,2} - x_T|^2 + |m_{1,3} - x_T|^2 + |m_{1,4} - x_T|^2 \\
&\quad + |m_{2,3} - x_T|^2 + |m_{2,4} - x_T|^2 + |m_{3,4} - x_T|^2 \\
&= \frac{1}{4} (2|m_{1,2} - m_{3,4}|^2 + 2|m_{1,3} - m_{2,4}|^2 + 2|m_{1,4} - m_{2,3}|^2) = \frac{L}{2}.
\end{aligned}$$

We similarly obtain

$$\begin{aligned}
& \frac{1}{|T|} \int_T |\nabla \varphi_T(x)|^2 dx \\
&= \frac{24^2}{|T|} \int_T |x - x_T|^2 dx \\
&= -\frac{24^2}{20} \sum_{i=1}^4 |x_i - x_T|^2 + \frac{24^2}{5} \sum_{1 \leq i < j \leq 4} |m_{i,j} - x_T|^2 = \frac{144}{5} L,
\end{aligned}$$

which leads to (5.8).  $\square$

We set the bubble space  $B_h$  by

$$B_h := \{b_h \in L^2(\Omega); b_h|_T \in \text{span}\{\varphi_T\}, \forall T \in \mathbb{T}_h\}. \quad (5.9)$$

Then, for any  $\psi_h \in CR_{h0}^1$  and  $b_h \in B_h$ , because one writes  $b_h|_T = c_b \varphi_T$  for  $c_b \in \mathbb{R}$ , it holds that

$$\begin{aligned}
(\nabla_h \psi_h, \nabla_h b_h) &= \sum_{T \in \mathbb{T}_h} c_b \int_T \nabla \psi_h \cdot \nabla \varphi_T dx \\
&= \sum_{T \in \mathbb{T}_h} c_b \left\{ \sum_{F \subset \partial T} (n_F \cdot \nabla \psi_h) \int_F \varphi_T ds - \int_T \Delta \psi_h \varphi_T dx \right\} = 0.
\end{aligned}$$



We here used the facts that (5.6),  $n_F \cdot \nabla \psi_h$  is constant on  $F$ , and  $\Delta \psi_h = 0$  on  $T$ . That is to say, two finite element spaces  $CR_{h0}^1$  and  $B_h$  are orthogonal to each other.

Furthermore, we define the finite element space  $X_h^{bCR}$  by

$$X_h^{bCR} := CR_{h0}^1 + B_h = \{\psi_h + b_h; \psi_h \in CR_{h0}^1, b_h \in B_h\}. \quad (5.10)$$

We consider the following finite element problem. Find  $u_h^{bCR} \in X_h^{bCR}$  such that

$$a_{0h}(u_h^{bCR}, \varphi_h) = (\nabla_h u_h^{bCR}, \nabla_h \varphi_h) = (\Pi_h^0 f, \varphi_h) \quad \forall \varphi_h \in X_h^{bCR}. \quad (5.11)$$

The solution  $u_h^{bCR} \in X_h^{bCR}$  is then decomposed as  $u_h^{bCR} = \bar{u}_h^{CR} + b_h$  with  $\bar{u}_h^{CR} \in CR_{h0}^1$  and  $b_h \in B_h$ . Note that  $\bar{u}_h^{CR}$  and  $b_h$  respectively satisfy (5.2) and the equation

$$a_{0h}(b_h, c_h) = (\nabla_h b_h, \nabla_h c_h) = (\Pi_h^0 f, c_h) \quad \forall c_h \in B_h. \quad (5.12)$$

On each element  $T \in \mathbb{T}_h$ , (5.12) has the form

$$\gamma_T \int_T \nabla \varphi_T \cdot \nabla \varphi_T dx = \int_T \Pi_T^0 f \varphi_T dx, \quad \gamma_T \in \mathbb{R}.$$

From (5.7) and (5.8), we have

$$\gamma_T = \frac{1}{72} \Pi_T^0 f \quad \forall T \in \mathbb{T}_h. \quad (5.13)$$

**Theorem 10** *Let  $u_h^{bCR} \in X_h^{bCR}$  be the solution of (5.11) and  $(\bar{\sigma}_h^{RT}, \bar{u}_h^{RT}) \in RT_h^0 \times M_h^0$  the solution of (5.1). We then have  $\nabla_h u_h^{bCR} \in RT_h^0$  and*

$$\bar{\sigma}_h^{RT} = \nabla u_h^{bCR} \quad \forall T \in \mathbb{T}_h, \quad (5.14)$$

$$\bar{u}_h^{RT} = \Pi_T^0 u_h^{bCR} \quad \forall T \in \mathbb{T}_h. \quad (5.15)$$

**Proof** The proof can be found in [15].  $\square$

From Theorem 10, for  $d = 3$ , the following lemma holds.

**Lemma 10** *Let  $\bar{u}_h^{CR} \in CR_{h0}^1$  be the solution of (5.2) and  $(\bar{\sigma}_h^{RT}, \bar{u}_h^{RT}) \in RT_h^0 \times M_h^0$  be the solution of (5.1). We then have the relationships*

$$\bar{\sigma}_h^{RT}|_T = \nabla \bar{u}_h^{CR} - \frac{1}{3} \Pi_T^0 f(x - x_T) \quad \forall T \in \mathbb{T}_h, \quad (5.16)$$

$$\bar{u}_h^{RT}|_T = \Pi_T^0 \bar{u}_h^{CR} + \frac{1}{180} \Pi_T^0 f \sum_{i=1}^4 |x_i - x_T|^2 \quad \forall T \in \mathbb{T}_h. \quad (5.17)$$

Using relationship between the RT and CR finite element methods, we have the error estimate of the CR finite element approximation with the bubble function.

**Lemma 11** *We assume that  $\Omega$  is convex. Let  $\{\mathbb{T}_h\}$  be a family of conformal meshes satisfying Assumption 1. Let  $\bar{u} \in H_0^1(\Omega) \cap H^2(\Omega)$  be the solution of (5.3) and  $u_h^{bCR} \in X_h^{bCR}$  be the solution of the CR problem (5.11). There then exists a constant  $c > 0$  independent of  $\bar{u}$ ,  $h$ ,  $H$  and the geometric properties of  $\mathbb{T}_h$  such that*

$$|\bar{u} - u_h^{bCR}|_{H^1(\mathbb{T}_h)} \leq cH \|\Pi_h^0 f\|. \quad (5.18)$$

**Proof** Let  $(\bar{\sigma}_h^{RT}, \bar{u}_h^{RT}) \in RT_h^0 \times M_h^0$  be the solution of (5.1). From Theorem 10, it holds that  $\nabla_h u_h^{bCR} \in RT_h^0$  and  $\bar{\sigma}_h^{RT} = \nabla_h u_h^{bCR}$ . Setting  $\bar{\sigma} := \nabla \bar{u} \in H^1(\Omega)^d$ , we then have, using inequality (4.11), that

$$\begin{aligned} |\bar{u} - u_h^{bCR}|_{H^1(\mathbb{T}_h)} &= \left( \sum_{T \in \mathbb{T}_h} \|\bar{\sigma} - \bar{\sigma}_h^{RT}\|_{L^2(T)^d}^2 \right)^{1/2} \\ &\leq cH |\bar{\sigma}|_{H^1(\Omega)^d} = cH |\bar{u}|_{H^2(\Omega)} \leq cH \|\Pi_h^0 f\|. \end{aligned}$$

□

## 6 Numerical Results

This section presents results of numerical examples. Let  $\Omega := (0, 1)^3$ . Let  $u_h^L$  and  $u_h^{CR}$  be the  $\mathcal{P}^1$ -Lagrange and  $\mathcal{P}^1$ -CR finite element solutions, respectively, for the model problem

$$\begin{aligned} -\Delta u &= 2y(1-y)z(1-z) + 2x(1-x)z(1-z) + 2x(1-x)y(1-y) \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega, \end{aligned}$$

which is the exact solution  $u = x(1-x)y(1-y)z(1-z)$ .


Let  $M$  be the division number of each side of the bottom face and  $N$  the division number of the height of  $\Omega$  with  $N \sim M^\gamma$  (see Fig. 2). There are two elements as shown in Fig. 3.

If an exact solution  $u$  is known, the error  $e_h := u - u_h$  and  $e_{h/2} := u - u_{h/2}$  are computed numerically for two mesh sizes  $h$  and  $h/2$ . The convergence indicator  $r$  is defined by


$$r = \frac{1}{\log(2)} \log \left( \frac{\|e_h\|_X}{\|e_{h/2}\|_X} \right).$$

We set  $h := \frac{1}{M}$ . The parameter  $H$  is then  $H = \mathcal{O}(h^{2-\gamma})$ . We compute the convergence order with respect to  $H_0^1$  and  $L^2$  norms defined by

$$\begin{aligned} Err_h^L(H^1) &:= \frac{|u - u_h^L|_{H^1(\Omega)}}{\|\Delta u\|}, & Err_h^L(L^2) &:= \frac{\|u - u_h^L\|}{\|\Delta u\|}, \\ Err_h^{CR}(H^1) &:= \frac{|u - u_h^{CR}|_{H^1(\mathbb{T}_h)}}{\|\Delta u\|}, & Err_h^{CR}(L^2) &:= \frac{\|u - u_h^{CR}\|}{\|\Delta u\|}, \end{aligned}$$



**Fig. 2** Mesh:  $M = 8$ ,  $N = 22$



**Fig. 3** Elements

for three cases:  $\gamma = 1.5$ ,  $\gamma = 1.9$  and  $\gamma = 2.0$ . In order to compute the above norms, we use the five-order fifteen-point numerical integration introduced in [17]. The results are give in Table 2, Table 3 when  $\gamma = 1.5$ , Table 4, Table 5 when  $\gamma = 1.9$ , and Table 6, Table 7 when  $\gamma = 2.0$ . Further,  $N_p^L$  and  $N_p^{CR}$  denote respectively the degrees of freedom for the  $\mathcal{P}^1$ -Lagrange finite element and the  $\mathcal{P}^1$ -CR finite element.

**Table 2** Error of the  $\mathcal{P}^1$ -Lagrange finite element solution ( $\gamma = 1.5$ )

$M$	$N$	$h$	$H$	$N_p^L$	$Err_h^L(H^1)$	$r$	$Err_h^L(L^2)$	$r$
4	8	2.50e-01	5.00e-01	225	1.2043e-01		9.5321e-03	
8	22	1.25e-01	3.54e-01	1,863	7.0318e-02	0.78	3.1646e-03	1.59
16	64	6.25e-02	2.50e-01	18,785	4.4662e-02	0.65	1.2570e-03	1.33
32	182	3.13e-02	1.77e-01	199,287	2.9479e-02	0.60	5.4477e-04	1.21

Observing the numerical results, the convergence indicators  $r$  in each norms are respectively

$$\begin{aligned}
 |u - u_h^L|_{H^1(\Omega)} &= \mathcal{O}(H), & \|u - u_h^L\| &= \mathcal{O}(H^2), \\
 |u - u_h^{CR}|_{H^1(\mathbb{T}_h)} &= \mathcal{O}(h), & \|u - u_h^{CR}\| &= \mathcal{O}(h^2),
 \end{aligned}$$

**Table 3** Error of the  $\mathcal{P}^1$ -CR finite element solution ( $\gamma = 1.5$ )

$M$	$N$	$h$	$H$	$N_p^{CR}$	$Err_h^{CR}(H^1)$	$r$	$Err_h^{CR}(L^2)$	$r$
4	8	2.50e-01	5.00e-01	1,440	8.2569e-02		3.8242e-03	
8	22	1.25e-01	3.54e-01	14,912	4.0629e-02	1.02	8.8356e-04	2.11
16	64	6.25e-02	2.50e-01	168,448	2.0042e-02	1.02	2.0485e-04	2.11
32	182	3.13e-02	1.77e-01	1,889,024	9.9579e-03	1.01	4.8960e-05	2.07

**Table 4** Error of the  $\mathcal{P}^1$ -Lagrange finite element solution ( $\gamma = 1.9$ )

$M$	$N$	$h$	$H$	$N_p^L$	$Err_h^L(H^1)$	$r$	$Err_h^L(L^2)$	$r$
4	14	2.50e-01	8.71e-01	345	1.4873e-01		1.4032e-02	
8	52	1.25e-01	8.12e-01	4,293	1.2167e-01	0.29	9.3061e-03	0.59
16	194	6.25e-02	7.58e-01	56,355	1.0919e-01	0.16	7.4989e-03	0.31
32	724	3.13e-02	7.07e-01	789,525	1.0128e-01	0.11	6.4558e-03	0.22

**Table 5** Error of the  $\mathcal{P}^1$ -CR finite element solution ( $\gamma = 1.9$ )

$M$	$N$	$h$	$H$	$N_p^{CR}$	$Err_h^{CR}(H^1)$	$r$	$Err_h^{CR}(L^2)$	$r$
4	14	2.50e-01	8.71e-01	2,496	7.9756e-02		3.2993e-03	
8	52	1.25e-01	8.12e-01	35,072	3.9708e-02	1.01	7.7177e-04	2.10
16	194	6.25e-02	7.58e-01	509,568	1.9814e-02	1.00	1.8781e-04	2.04
32	724	3.13e-02	7.07e-01	7,508,480	9.9003e-03	1.00	4.6546e-05	2.01

**Table 6** Error of the  $\mathcal{P}^1$ -Lagrange finite element solution ( $\gamma = 2.0$ )

$M$	$N$	$h$	$H$	$N_p^L$	$Err_h^L(H^1)$	$r$	$Err_h^L(L^2)$	$r$
4	16	2.50e-01	1.00	425	1.5862e-01		1.5909e-02	
8	64	1.25e-01	1.00	5,265	1.4079e-01	0.17	1.2472e-02	0.35
16	256	6.25e-02	1.00	74,273	1.3597e-01	0.05	1.1646e-02	0.10
32	1,024	3.13e-02	1.00	1,116,225	1.3474e-01	0.01	1.1442e-02	0.03

where  $H = \mathcal{O}(h^{2-\gamma})$ . Meanwhile, the theoretical results are as follows:

$$\begin{aligned} |u - u_h^L|_{H^1(\Omega)} &= \mathcal{O}(H), \quad \|u - u_h^L\| = \mathcal{O}(H^2), \\ |u - u_h^{CR}|_{H^1(\mathbb{T}_h)} &= \mathcal{O}(H), \quad \|u - u_h^{CR}\| = \mathcal{O}(H^2), \end{aligned}$$

if  $\Omega$  is convex and  $u \in H^2(\Omega) \cap H_0^1(\Omega)$ . In this numerical examples, the CR finite element approximation is superior to the Lagrange finite element approximation on this anisotropic meshes. The theoretical explanation of this point is still open.

**Table 7** Error of the  $\mathcal{P}^1$ -CR finite element solution ( $\gamma = 2.0$ )

$M$	$N$	$h$	$H$	$N_p^{CR}$	$Err_h^{CR}(H^1)$	$r$	$Err_h^{CR}(L^2)$	$r$
4	16	2.50e-01	1.00	2,848	7.9473e-02		3.2264e-03	
8	64	1.25e-01	1.00	43,136	3.9647e-02	1.00	7.6153e-04	2.08
16	256	6.25e-02	1.00	672,256	1.9803e-02	1.00	1.8680e-04	2.03
32	1,024	3.13e-02	1.00	10,618,880	9.8984e-03	1.00	4.6458e-05	2.01

**Acknowledgements** This work was supported by JSPS KAKENHI Grant Number JP16H03950. We would like to thank the anonymous referee for the valuable comments.

## References

1. Acosta, G., Durán, R.G.: The maximum angle condition for mixed and nonconforming elements: Application to the Stokes equations, *SIAM J. Numer. Anal.* **37**, 18-36 (1999)
2. Apel, Th.: Anisotropic finite elements: Local estimates and applications. *Advances in Numerical Mathematics*. Teubner, Stuttgart, (1999)
3. Arnord, D.T., Brezzi, F.: Mixed and nonconforming finite element methods: implementation, postprocessing and error estimates. *RAIRO Modélisation mathématique et analyse numérique* **19**, 7-32 (1985)
4. Babuška, I., Aziz, A.K.: On the angle condition in the finite element method. *SIAM J. Numer. Anal.* **13**, 214-226 (1976)
5. Boffi, D., Brezzi, F., Demkowicz, L.F., Durán, R.G., Falk, R.S., Fortin, M.: Mixed Finite Elements, Compatibility Conditions, and Applications : Lectures Given at the C.I.M.E. Summer School, Italy, 2006. *Lecture Notes in Mathematics* **1939**, Springer, (2008)
6. Boffi, D., Brezzi, F., Fortin, M.: Mixed Finite Element Methods and Applications. Springer Verlag, New York (2013)
7. Braess, D.: Finite elements Theory, fast solvers, and application in solid mechanics. Cambridge, (2007)
8. Brenner, S. C. : Forty years of the Crouzeix–Raviart element, *Numer. Methods Partial Differential Equations* **31**, 367-396 (2015)
9. Brenner, S.C., Scott, L.R.: The Mathematical Theory of Finite Element Methods, Third Edition. Springer Verlag, New York (2008)
10. Ciarlet, P. G.: The Finite Element Method for Elliptic problems. SIAM, New York (2002)
11. Ern, A., Guermond, J.L.: Theory and Practice of Finite Elements. Springer Verlag, New York (2004)
12. Girault, V., Raviart, P.A.: Finite Element Methods for Navier-Stokes Equations. Springer-Verlag, (1986)
13. Grisvard, P.: Elliptic Problems in Nonsmooth Domains. SIAM, (2011)
14. Gudi, T.: A New Error Analysis for Discontinuous Finite Element Methods for Linear Elliptic Problems. *Math. Comp.* **79**, 2169-2189 (2010)
15. Hu, J., Ma, R.: The Enriched Crouzeix–Raviart Elements are Equivalent to the Raviart-Thomas Elements. *J. Sci. Comput.* **63**, 410-425 (2015)
16. Ishizaka, H., Kobayashi, K., Tsuchiya, T.: General theory of interpolation error estimates on anisotropic meshes. *Japan Journal of Industrial and Applied Mathematics*, <https://arxiv.org/abs/2002.09721> (2020)
17. Keast, P.: Moderate-degree tetrahedral quadrature formulas, *Computer Methods in Applied Mechanics and Engineering* **55**, Issue 3, 339-348 (1986)
18. Kikuchi, F., Saito, N.: Principle of Numerical Analysis. (in Japanese) Iwatani-Shoten, (2016)
19. Kobayashi, K., Tsuchiya, T.: Error analysis of Crouzeix-Raviart and Raviart-Thomas finite element methods. *Japan Journal of Industrial and Applied Mathematics* **35**, 1191-1211 (2018)
20. Liu, X., Kikuchi, F.: Estimation of error constants appearing in non-conforming linear triangular finite element, *Proceedings of APCOM'07-EPMESC XI* (2007)
21. Liu, X., Kikuchi, F.: Explicit estimation of error constants appearing in non-conforming linear triangular finite element method, *Applications of Mathematics* **63**, 381-397 (2018)
22. Marini, L.D.: An inexpensive method for the evaluation of the solution of the lowest order Raviart-Thomas mixed method. *SIAM J. Numer. Anal.* **22**, 493-496 (1985)
23. Mario, B.: A note on the Poincaré inequality for convex domains. *Z. Anal. ihre. Anwend.*, **22**, 751-756 (2003)
24. Payne, L.E., Weinberger, H.F.: An optimal Poincaré-inequality for convex domains, *Arch. Rational Mech. Anal.* **5**, 286-292 (1960)
25. Stroud, A.H.: Approximate Calculation of Multiple Integrals. Prentice-Hall, (1971)