

Conditional Independence Beyond Domain Separability

Discussion of Engelke and Hitz (2020)

Yuexia Zhang¹ and Linbo Wang^{†1,2}

¹Department of Computer and Mathematical Sciences, University of Toronto Scarborough

²Department of Statistical Sciences, University of Toronto

1 A general definition of conditional independence

We congratulate Engelke and Hitz on a thought-provoking paper on graphical models for extremes. A key contribution of the paper is the introduction of a novel definition of conditional independence for a multivariate Pareto distribution. Here, we outline a proposal for independence and conditional independence of general random variables whose support is a general set Ω in \mathbb{R}^d . Our proposal includes the authors' definition of conditional independence, and the analogous definition of independence as special cases. By making our proposal independent of the context of extreme value theory, we highlight the importance of the authors' contribution beyond this particular context.

Definition 1: Suppose that $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C)$ is a random vector with support Ω in \mathbb{R}^d , where A, B, C are disjoint sets whose union is $\{1, \dots, d\}$. Let $U \times V$ denote the Cartesian product of U and V .

- (a) We say \mathbf{Y}_A is *conditionally outer independent* of \mathbf{Y}_C given \mathbf{Y}_B if there exists a random vector $\mathbf{W} = (\mathbf{W}_A, \mathbf{W}_B, \mathbf{W}_C)$ with support $L_A \times L_B \times L_C$ in \mathbb{R}^d such that (i) $\Omega \subset L_A \times L_B \times L_C$; (ii) $(\mathbf{W} \mid \mathbf{W} \in \Omega) \stackrel{d}{=} \mathbf{Y}$; (iii) $\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid \mathbf{W}_B$. In this case, we write $\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \mid \mathbf{Y}_B$.

If $B = \emptyset$, we say \mathbf{Y}_A is *outer independent* of \mathbf{Y}_C , denoted as $\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C$.

- (b) We say \mathbf{Y}_A is *conditionally inner independent* of \mathbf{Y}_C given \mathbf{Y}_B if for any $S_A \times S_B \times S_C \subset \Omega$ such that S_k is a measurable subset of $\mathbb{R}^{\dim(Y_k)}$, $k \in \{A, B, C\}$ and $P(\mathbf{Y} \in S_A \times S_B \times S_C) > 0$, we have $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y}_B, \mathbf{Y} \in S_A \times S_B \times S_C)$. In this case, we write $\mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \mid \mathbf{Y}_B$.

If $B = \emptyset$, we say \mathbf{Y}_A is *inner independent* of \mathbf{Y}_C , denoted as $\mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C$.

Proposition 1: Suppose $\Omega = [0, \infty)^d \setminus [0, 1]^d$ as in EH's case. Then

$$\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \mid \mathbf{Y}_B \Leftrightarrow \mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \mid \mathbf{Y}_B \Leftrightarrow \mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C \mid \mathbf{Y}_B.$$

where $\perp\!\!\!\perp_e$ denotes the notion of conditional independence introduced by Engelke and Hitz (2020). In particular, if $B = \emptyset$, then

$$\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \Leftrightarrow \mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \Leftrightarrow \mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C.$$

Remark 1: We do not place any distributional assumptions on \mathbf{Y} in Proposition 1.

[†]E-mail address for correspondence: linbo.wang@utoronto.ca

Engelke and Hitz showed that if \mathbf{Y} is multivariate Pareto and admits a positive and continuous density, then $\mathbf{Y}_A \not\perp_e \mathbf{Y}_C$. This does not rule out the possibility of $\mathbf{Y}_A \perp_e \mathbf{Y}_C$ for general Pareto distributions. For example, consider two independent standard Pareto distributions X_1 and X_2 . Following eqn. (6) in Engelke and Hitz (2020), all the probability mass of \mathbf{Y} lies on $(1, \infty) \times \{0\}$ and $\{0\} \times (1, \infty)$ so that it does not admit a density with respect to Lebesgue measure. Nevertheless $Y_1 \perp_e Y_2$. This observation is generalized in Proposition 2.

Proposition 2: Suppose $\Omega = [([0, \infty)^{|A|} \setminus [0, 1]^{|A|}) \times \{0\}^{|B|} \times \{0\}^{|C|}] \cup [\{0\}^{|A|} \times ([0, \infty)^{|B|} \setminus [0, 1]^{|B|}) \times \{0\}^{|C|}] \cup [\{0\}^{|A|} \times \{0\}^{|B|} \times ([0, \infty)^{|C|} \setminus [0, 1]^{|C|})]$, where $|A|$ is the cardinality of set A . Then

$$\mathbf{Y}_A \perp_o \mathbf{Y}_C \mid \mathbf{Y}_B \Leftrightarrow \mathbf{Y}_A \perp_i \mathbf{Y}_C \mid \mathbf{Y}_B \Leftrightarrow \mathbf{Y}_A \perp_e \mathbf{Y}_C \mid \mathbf{Y}_B.$$

In particular, if $B = \emptyset$, then

$$\mathbf{Y}_A \perp_o \mathbf{Y}_C \Leftrightarrow \mathbf{Y}_A \perp_i \mathbf{Y}_C \Leftrightarrow \mathbf{Y}_A \perp_e \mathbf{Y}_C.$$

2 Proofs of propositions

2.1 Proof of Proposition 1 in the case where $B = \emptyset$

Proof. We need to show

$$\mathbf{Y}_A \perp_o \mathbf{Y}_C \Rightarrow \mathbf{Y}_A \perp_i \mathbf{Y}_C \Rightarrow \mathbf{Y}_A \perp_e \mathbf{Y}_C \Rightarrow \mathbf{Y}_A \perp_o \mathbf{Y}_C. \quad (1)$$

The first two claims are straightforward. We now prove the third claim in (1).

Let $H_k^1 \subset L_k^1 = [0, 1]^{|k|}$, $H_k^2 \subset L_k^2 = [0, \infty)^{|k|} \setminus [0, 1]^{|k|}$, $H_k = H_k^1 \cup H_k^2$, $L_k = L_k^1 \cup L_k^2$, $k \in \{A, C\}$. Obviously, $H_k^1 \cap H_k^2 = \emptyset$ and $L_k^1 \cap L_k^2 = \emptyset$. Besides, $\Omega = [0, \infty)^d \setminus [0, 1]^d \subset L_A \times L_C$.

We let

$$P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^1) = \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2)P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)}, \quad (2)$$

and

$$P(\mathbf{W}_A \in H_A^i, \mathbf{W}_C \in H_C^j) = \lambda P(\mathbf{Y}_A \in H_A^i, \mathbf{Y}_C \in H_C^j), \quad (i, j) \neq (1, 1),$$

where λ is a normalizing constant. Simple calculation yields

$$\lambda = \frac{1}{1 + \frac{P(\mathbf{Y}_A \in L_A^1, \mathbf{Y}_C \in L_C^2)P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_C \in L_C^1)}{P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_C \in L_C^2)}}.$$

It is easy to see that $(\mathbf{W} \mid \mathbf{W} \in \Omega) \stackrel{d}{=} \mathbf{Y}$. We then prove

$$\mathbf{W}_A \perp \mathbf{W}_C.$$

If $k \in A$ and $\{\mathbf{W}_A \in H_A\} \cap \{\mathbf{W}_A : W_k > 1\} = \emptyset$, then

$$P(\mathbf{W}_A \in H_A, \mathbf{W}_C \in H_C \mid W_k > 1) = P(\mathbf{W}_A \in H_A \mid W_k > 1)P(\mathbf{W}_C \in H_C \mid W_k > 1) = 0. \quad (3)$$

If $k \in A$ and $\{\mathbf{W}_A \in H_A\} \cap \{\mathbf{W}_A : W_k > 1\} \neq \emptyset$, then $\{\mathbf{W}_A \in H_A\} \cap \{\mathbf{W}_A : W_k > 1\} = \{\mathbf{W}_A \in H_A^2, W_k > 1\}$ and

$$\begin{aligned}
& P(\mathbf{W}_A \in H_A, \mathbf{W}_C \in H_C, W_k > 1) \\
&= \lambda P(\mathbf{Y}_A \in H_A, \mathbf{Y}_C \in H_C, Y_k > 1) \\
&= \lambda P(\mathbf{Y}_C \in H_C \mid \mathbf{Y}_A \in H_A, Y_k > 1) P(\mathbf{Y}_A \in H_A, Y_k > 1) \\
&= P(\mathbf{Y}_C \in H_C \mid Y_k > 1) P(\mathbf{W}_A \in H_A, W_k > 1) \quad (\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid Y_k > 1).
\end{aligned}$$

So we have

$$\begin{aligned}
P(\mathbf{W}_C \in H_C \mid \mathbf{W}_A \in H_A, W_k > 1) &= P(\mathbf{Y}_C \in H_C \mid Y_k > 1) \\
&= \frac{P(\mathbf{Y}_C \in H_C, Y_k > 1)}{P(Y_k > 1)} \\
&= \frac{P(\mathbf{W}_C \in H_C, W_k > 1)/\lambda}{P(W_k > 1)/\lambda} \\
&= \frac{P(\mathbf{W}_C \in H_C, W_k > 1)}{P(W_k > 1)} \\
&= P(\mathbf{W}_C \in H_C \mid W_k > 1).
\end{aligned} \tag{4}$$

Thus, from (3) and (4), we have for any $k \in A$, $\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid W_k > 1$. Similarly, we have for any $k \in C$, $\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid W_k > 1$. In summary,

$$\forall k \in \{1, \dots, d\} : \mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid W_k > 1. \tag{5}$$

We then show that for any $H_A^1, H_C^1, H_A^2, H_C^2$, we have

$$\frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1)} = \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^2)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^2)}. \tag{6}$$

To prove (6), note that

$$\begin{aligned}
& \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^2)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^2)} P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1) \\
&= \frac{P(\mathbf{W}_A \in H_A^1 \mid \mathbf{W}_C \in H_C^2)}{P(\mathbf{W}_A \in H_A^2 \mid \mathbf{W}_C \in H_C^2)} P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1) \\
&= \frac{P(\mathbf{W}_A \in H_A^1 \mid \mathbf{W}_C \in H_C^2, \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in H_A^2 \mid \mathbf{W}_C \in H_C^2, \mathbf{W}_C \in L_C^2)} P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1) \\
&= \frac{P(\mathbf{W}_A \in H_A^1 \mid \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in H_A^2 \mid \mathbf{W}_C \in L_C^2)} P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1) \quad \text{due to (5)} \\
&= \frac{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in L_C^2)} P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2) \\
&= \frac{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_A \in L_A^2, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)} P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2) \\
&= \frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)} P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2) \quad \text{due to (5)} \\
&= P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^1) \quad \text{due to (2)}.
\end{aligned}$$

We then show that

$$P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^2) = P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^1). \tag{7}$$

To prove (7), note that due to (6),

$$\frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1)} = \frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)}.$$

Then

$$\begin{aligned} \text{LHS of (7)} &= \frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in L_A^2)} \\ &= \frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2) + P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1)} \\ &= \frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2) + P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^1)} \\ &= \frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in L_A^1)} \\ &= P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^1) = \text{RHS of (7)}. \end{aligned}$$

Similarly we have

$$P(\mathbf{W}_C \in L_C^1 \mid \mathbf{W}_A \in L_A^2) = P(\mathbf{W}_C \in L_C^1 \mid \mathbf{W}_A \in L_A^1).$$

We then show that

$$P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^2) = P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^1).$$

To see this, due to (6),

$$\frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in H_C^1)} = \frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)}.$$

Furthermore, due to (7),

$$\frac{P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in L_A^1)}{P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in L_A^2)} = \frac{P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^1)}{P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^2)} = 1.$$

Due to (5) and (6), it follows directly that

$$\begin{aligned} P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^1) &= P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^2) \\ &= P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in H_A^2) = P(\mathbf{W}_C \in H_C^2). \end{aligned} \quad (8)$$

Similarly we have

$$P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in L_A^1) = P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in H_A^2) = P(\mathbf{W}_C \in H_C^1). \quad (9)$$

We still need to show

$$P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in H_A^1) = P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^1), \quad (10)$$

$$P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in H_A^1) = P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in L_A^1). \quad (11)$$

To show (10), note that due to (5) and (6),

$$\begin{aligned}
& P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in H_A^1) \\
&= \frac{P(\mathbf{W}_C \in H_C^2, \mathbf{W}_A \in H_A^1)}{P(\mathbf{W}_A \in H_A^1)} \\
&= \frac{\lambda P(\mathbf{Y}_C \in H_C^2, \mathbf{Y}_C \in L_C^2, \mathbf{Y}_A \in H_A^1)}{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^1) + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2)} \\
&= \frac{\lambda P(\mathbf{Y}_C \in H_C^2 \mid \mathbf{Y}_C \in L_C^2) P(\mathbf{Y}_A \in H_A^1, \mathbf{Y}_C \in L_C^2)}{\frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2) P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)} + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2)} \\
&= \frac{P(\mathbf{Y}_C \in H_C^2 \mid \mathbf{Y}_C \in L_C^2) P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2)}{\frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2) P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)} + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2)} \\
&= \frac{P(\mathbf{Y}_C \in H_C^2 \mid \mathbf{Y}_C \in L_C^2)}{\frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)} + 1}
\end{aligned}$$

does not depend on H_A^1 .

To show (11), note that due to (2) and (6),

$$\begin{aligned}
& P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in H_A^1) \\
&= \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^1) + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2)} \\
&= \frac{\frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2) P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)}}{\frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2) P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)} + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2)} \\
&= \frac{\frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)}}{\frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2)} + 1}
\end{aligned}$$

does not depend on H_A^1 .

From (8), (9), (10) and (11), we have

$$P(\mathbf{W}_C \in H_C \mid \mathbf{W}_A \in H_A) = P(\mathbf{W}_C \in H_C).$$

Hence $\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C$.

□

2.2 Proof of Proposition 1 for a general B

Proof. We need to show

$$\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \mid \mathbf{Y}_B \Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \mid \mathbf{Y}_B \Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C \mid \mathbf{Y}_B \Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \mid \mathbf{Y}_B. \quad (12)$$

The first two claims are straightforward. We now prove the third claim in (12).

Let $H_k^1 \subset L_k^1 = [0, 1]^{|k|}$, $H_k^2 \subset L_k^2 = [0, \infty)^{|k|} \setminus [0, 1]^{|k|}$, $H_k = H_k^1 \cup H_k^2$, $L_k = L_k^1 \cup L_k^2$, $k \in \{A, B, C\}$. Obviously, $H_k^1 \cap H_k^2 = \emptyset$ and $L_k^1 \cap L_k^2 = \emptyset$. Besides, $\Omega = [0, \infty)^d \setminus [0, 1]^d \subset L_A \times L_B \times L_C$.

We let

$$\begin{aligned} & P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in H_C^1) \\ &= \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)}, \end{aligned} \quad (13)$$

and for $(i, j, k) \neq (1, 1, 1)$,

$$P(\mathbf{W}_A \in H_A^i, \mathbf{W}_B \in H_B^j, \mathbf{W}_C \in H_C^k) = \lambda P(\mathbf{Y}_A \in H_A^i, \mathbf{Y}_B \in H_B^j, \mathbf{Y}_C \in H_C^k),$$

where λ is a normalizing constant. Simple calculation yields

$$\lambda = \frac{1}{1 + \frac{P(\mathbf{Y}_A \in L_A^1, \mathbf{Y}_B \in L_B^1, \mathbf{Y}_C \in L_C^2)P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_B \in L_B^1, \mathbf{Y}_C \in L_C^1)}{P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_B \in L_B^1, \mathbf{Y}_C \in L_C^2)}}.$$

It is easy to see that $(\mathbf{W} \mid \mathbf{W} \in \Omega) \stackrel{d}{=} \mathbf{Y}$. We then prove

$$\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid \mathbf{W}_B.$$

If $k \in A$ and $\{\mathbf{W}_A \in H_A\} \cap \{\mathbf{W}_A : W_k > 1\} = \emptyset$, then

$$\begin{aligned} P(\mathbf{W}_A \in H_A, \mathbf{W}_C \in H_C \mid \mathbf{W}_B \in H_B, W_k > 1) &= P(\mathbf{W}_A \in H_A \mid \mathbf{W}_B \in H_B, W_k > 1) \\ &\times P(\mathbf{W}_C \in H_C \mid \mathbf{W}_B \in H_B, W_k > 1) = 0. \end{aligned} \quad (14)$$

If $k \in A$ and $\{\mathbf{W}_A \in H_A\} \cap \{\mathbf{W}_A : W_k > 1\} \neq \emptyset$, then $\{\mathbf{W}_A \in H_A\} \cap \{\mathbf{W}_A : W_k > 1\} = \{\mathbf{W}_A \in H_A^2, W_k > 1\}$. Note that

$$\begin{aligned} & P(\mathbf{W}_A \in H_A, \mathbf{W}_B \in H_B, \mathbf{W}_C \in H_C, W_k > 1) \\ &= \lambda P(\mathbf{Y}_A \in H_A, \mathbf{Y}_B \in H_B, \mathbf{Y}_C \in H_C, Y_k > 1) \\ &= \lambda P(\mathbf{Y}_C \in H_C \mid \mathbf{Y}_A \in H_A, \mathbf{Y}_B \in H_B, Y_k > 1)P(\mathbf{Y}_A \in H_A, \mathbf{Y}_B \in H_B, Y_k > 1) \\ &= P(\mathbf{Y}_C \in H_C \mid \mathbf{Y}_B \in H_B, Y_k > 1)P(\mathbf{W}_A \in H_A, \mathbf{W}_B \in H_B, W_k > 1) \quad (\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid \mathbf{Y}_B, Y_k > 1). \end{aligned}$$

So we have

$$\begin{aligned} P(\mathbf{W}_C \in H_C \mid \mathbf{W}_A \in H_A, \mathbf{W}_B \in H_B, W_k > 1) &= P(\mathbf{Y}_C \in H_C \mid \mathbf{Y}_B \in H_B, Y_k > 1) \\ &= \frac{P(\mathbf{Y}_C \in H_C, \mathbf{Y}_B \in H_B, Y_k > 1)}{P(\mathbf{Y}_B \in H_B, Y_k > 1)} \\ &= \frac{P(\mathbf{W}_C \in H_C, \mathbf{W}_B \in H_B, W_k > 1)/\lambda}{P(\mathbf{W}_B \in H_B, W_k > 1)/\lambda} \quad (15) \\ &= \frac{P(\mathbf{W}_C \in H_C, \mathbf{W}_B \in H_B, W_k > 1)}{P(\mathbf{W}_B \in H_B, W_k > 1)} \\ &= P(\mathbf{W}_C \in H_C \mid \mathbf{W}_B \in H_B, W_k > 1). \end{aligned}$$

Thus, from (14) and (15), we have for any $k \in A$, $\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid \mathbf{W}_B, W_k > 1$. Similarly, we have for any $k \in C$, $\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid \mathbf{W}_B, W_k > 1$.

If $k \in B$ and $\{\mathbf{W}_B \in H_B\} \cap \{\mathbf{W}_B : W_k > 1\} = \emptyset$, then according to the definition in Shorack (2017, §7.4),

$$P(\mathbf{W}_C \in H_C \mid \mathbf{W}_A \in H_A, \mathbf{W}_B \in H_B, W_k > 1) = P(\mathbf{W}_C \in H_C). \quad (16)$$

If $k \in B$ and $\{\mathbf{W}_B \in H_B\} \cap \{\mathbf{W}_B : W_k > 1\} \neq \emptyset$, then $\{\mathbf{W}_B \in H_B\} \cap \{\mathbf{W}_B : W_k > 1\} = \{\mathbf{W}_B \in H_B^2, W_k > 1\}$. Besides, similar to the proof of (15), we have

$$P(\mathbf{W}_C \in H_C \mid \mathbf{W}_A \in H_A, \mathbf{W}_B \in H_B, W_k > 1) = P(\mathbf{W}_C \in H_C \mid \mathbf{W}_B \in H_B, W_k > 1). \quad (17)$$

Thus, from (16) and (17), we have for any $k \in B$, $\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid \mathbf{W}_B, W_k > 1$.

In summary,

$$\forall k \in \{1, \dots, d\} : \mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C \mid \mathbf{W}_B, W_k > 1. \quad (18)$$

From (18), obviously, we have

$$P(\mathbf{W}_C \in H_C \mid \mathbf{W}_A \in H_A, \mathbf{W}_B \in H_B^2) = P(\mathbf{W}_C \in H_C \mid \mathbf{W}_B \in H_B^2). \quad (19)$$

We then show that for any $H_A^1, H_B^1, H_C^1, H_A^2, H_C^2$, we have

$$\frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1)} = \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^2 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^2 \mid \mathbf{W}_B \in H_B^1)}. \quad (20)$$

To prove (20), note that

$$\begin{aligned} & \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^2 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^2 \mid \mathbf{W}_B \in H_B^1)} P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1) \\ &= \frac{P(\mathbf{W}_A \in H_A^1 \mid \mathbf{W}_C \in H_C^2, \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in H_A^2 \mid \mathbf{W}_C \in H_C^2, \mathbf{W}_B \in H_B^1)} P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1) \\ &= \frac{P(\mathbf{W}_A \in H_A^1 \mid \mathbf{W}_C \in L_C^2, \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in H_A^2 \mid \mathbf{W}_C \in L_C^2, \mathbf{W}_B \in H_B^1)} P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1) \quad \text{due to (18)} \\ &= \frac{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in H_A^2, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1)} P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1) \\ &= \frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1)} P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1) \quad \text{due to (18)} \\ &= P(\mathbf{W}_A \in H_A^1, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1) \quad \text{due to (13)}. \end{aligned}$$

We then show that

$$P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1) = P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1). \quad (21)$$

To prove (21), note that due to (20),

$$\frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^1 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1 \mid \mathbf{W}_B \in H_B^1)} = \frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1)}.$$

Then

$$\begin{aligned} \text{LHS of (21)} &= \frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2, \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1)} \\ &= \frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2, \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2, \mathbf{W}_B \in H_B^1) + P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^1, \mathbf{W}_B \in H_B^1)} \end{aligned}$$

$$\begin{aligned}
&= \frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1) + P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^1 \mid \mathbf{W}_B \in H_B^1)} \\
&= \frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^1 \mid \mathbf{W}_B \in H_B^1)} \\
&= P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1) = \text{RHS of (21)}.
\end{aligned}$$

Similarly we have

$$P(\mathbf{W}_C \in L_C^1 \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1) = P(\mathbf{W}_C \in L_C^1 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1).$$

We then show that

$$P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1) = P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1).$$

To see this, due to (20),

$$\frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1)} = \frac{P(\mathbf{W}_A \in L_A^1, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_C \in L_C^2 \mid \mathbf{W}_B \in H_B^1)}.$$

Furthermore, due to (21),

$$\frac{P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1)} = \frac{P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_C \in L_C^2 \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1)} = 1.$$

Due to (18) and (20), it follows directly that

$$\begin{aligned}
P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1) &= P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1) \\
&= P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in H_A^2, \mathbf{W}_B \in H_B^1) \\
&= P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_B \in H_B^1). \quad (22)
\end{aligned}$$

Similarly we have

$$\begin{aligned}
P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1) &= P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in H_A^2, \mathbf{W}_B \in H_B^1) = P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_B \in H_B^1). \quad (23)
\end{aligned}$$

We still need to show

$$P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1) = P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1), \quad (24)$$

$$P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1) = P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in L_A^1, \mathbf{W}_B \in H_B^1). \quad (25)$$

To show (24), note that due to (18) and (20),

$$\begin{aligned}
&P(\mathbf{W}_C \in H_C^2 \mid \mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1) \\
&= \frac{P(\mathbf{W}_C \in H_C^2, \mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1)}{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1)} \\
&= \frac{\lambda P(\mathbf{Y}_C \in H_C^2, \mathbf{Y}_C \in L_C^2, \mathbf{Y}_A \in H_A^1, \mathbf{Y}_B \in H_B^1)}{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^1) + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\lambda P(\mathbf{Y}_C \in H_C^2 \mid \mathbf{Y}_B \in H_B^1, \mathbf{Y}_C \in L_C^2) P(\mathbf{Y}_A \in H_A^1, \mathbf{Y}_B \in H_B^1, \mathbf{Y}_C \in L_C^2)}{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2) P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)} + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2) \\
&= \frac{P(\mathbf{Y}_C \in H_C^2 \mid \mathbf{Y}_B \in H_B^1, \mathbf{Y}_C \in L_C^2) P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)}{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2) P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)} + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2) \\
&= \frac{P(\mathbf{Y}_C \in H_C^2 \mid \mathbf{Y}_B \in H_B^1, \mathbf{Y}_C \in L_C^2)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)} + 1
\end{aligned}$$

does not depend on H_A^1 .

To show (25), note that due to (13) and (20),

$$\begin{aligned}
&P(\mathbf{W}_C \in H_C^1 \mid \mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1) \\
&= \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^1) + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)} \\
&= \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2) P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)} \\
&= \frac{P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2) P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)} + P(\mathbf{W}_A \in H_A^1, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2) \\
&= \frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in H_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)} \\
&= \frac{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^1)}{P(\mathbf{W}_A \in L_A^2, \mathbf{W}_B \in H_B^1, \mathbf{W}_C \in L_C^2)} + 1
\end{aligned}$$

does not depend on H_A^1 .

From (22), (23), (24) and (25), we have

$$P(\mathbf{W}_C \in H_C \mid \mathbf{W}_A \in H_A, \mathbf{W}_B \in H_B^1) = P(\mathbf{W}_C \in H_C \mid \mathbf{W}_B \in H_B^1).$$

Besides, because of (19), we can obtain that

$$P(\mathbf{W}_C \in H_C \mid \mathbf{W}_A \in H_A, \mathbf{W}_B \in H_B) = P(\mathbf{W}_C \in H_C \mid \mathbf{W}_B \in H_B).$$

We have hence completed the proof. \square

2.3 Proof of Proposition 2 in the case where $B = \emptyset$

Proof. Use the same notation as before, denote $[0, 1]^{|k|}$ as L_k^1 , $[0, \infty)^{|k|} \setminus [0, 1]^{|k|}$ as L_k^2 for $k \in \{A, C\}$. Besides, denote $\{0\}^{|k|}$ as $\mathbf{0}_k$.

We need to show

$$\text{the support of } \mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C) \text{ is } \Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2) \Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C, \quad (26)$$

$$\text{the support of } \mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C) \text{ is } \Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2) \Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C, \quad (27)$$

$$\text{the support of } \mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C) \text{ is } \Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2) \Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C. \quad (28)$$

Under (26), (27) and (28), we can obtain that if the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2)$, then

$$\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \Leftrightarrow \mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \Leftrightarrow \mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C.$$

First, we show (26). Since the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2)$, then $S_A \times S_C \subset (L_A^2 \times \mathbf{0}_C)$ or $S_A \times S_C \subset (\mathbf{0}_A \times L_C^2)$. When $S_A \times S_C \subset (L_A^2 \times \mathbf{0}_C)$, since $\mathbf{Y}_C = \mathbf{0}_C$ regardless of the value of \mathbf{Y}_A , then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y} \in S_A \times S_C)$. When $S_A \times S_C \subset (\mathbf{0}_A \times L_C^2)$, since $\mathbf{Y}_A = \mathbf{0}_A$ regardless of the value of \mathbf{Y}_C , then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y} \in S_A \times S_C)$. Thus, based on the definition of $\mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C$, we have $\mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C$.

Second, we show (27). Since the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2)$, then when $Y_k > 1$ for some $k \in A$, we have $\mathbf{Y} \in (L_A^2 \times \mathbf{0}_C)$. Since $\mathbf{Y}_C = \mathbf{0}_C$ regardless of the value of \mathbf{Y}_A , then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid Y_k > 1$. When $Y_k > 1$ for some $k \in C$, we have $\mathbf{Y} \in (\mathbf{0}_A \times L_C^2)$. Since $\mathbf{Y}_A = \mathbf{0}_A$ regardless of the value of \mathbf{Y}_C , then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid Y_k > 1$. Thus, based on the definition of $\mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C$, we have $\mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C$.

Third, we show (28). Since the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2)$, then

$$\begin{aligned} P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_C = \mathbf{0}_C \text{ or } \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \in L_C^2) \\ = P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_C = \mathbf{0}_C) + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \in L_C^2) = 1. \end{aligned}$$

Suppose $P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_C = \mathbf{0}_C) = \alpha$ ($0 < \alpha < 1$), then $P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \in L_C^2) = 1 - \alpha$. Now we introduce two new random vectors \mathbf{W}_A and \mathbf{W}_C , assume that they have the distributions as follows:

$$F_{\mathbf{W}_A}(\mathbf{w}_A) = \begin{cases} (1 - p_A)P(\mathbf{U}_A \leq \mathbf{w}_A), & \mathbf{w}_A \in L_A^1, \\ (1 - p_A) + p_AP(\mathbf{Y}_A \leq \mathbf{w}_A \mid \mathbf{Y}_A \in L_A^2, \mathbf{Y}_C = \mathbf{0}_C), & \mathbf{w}_A \in L_A^2, \end{cases}$$

and

$$F_{\mathbf{W}_C}(\mathbf{w}_C) = \begin{cases} (1 - p_C)P(\mathbf{U}_C \leq \mathbf{w}_C), & \mathbf{w}_C \in L_C^1, \\ (1 - p_C) + p_CP(\mathbf{Y}_C \leq \mathbf{w}_C \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \in L_C^2), & \mathbf{w}_C \in L_C^2, \end{cases}$$

where $p_A \in (0, 1)$, $p_C \in (0, 1)$, \mathbf{U}_A is a random vector which follows the uniform distribution in L_A^1 , and \mathbf{U}_C is a random vector which follows the uniform distribution in L_C^1 .

To make $\mathbf{W}_A \perp\!\!\!\perp \mathbf{W}_C$, let $F(\mathbf{w}_A, \mathbf{w}_C) = F_{\mathbf{W}_A}(\mathbf{w}_A)F_{\mathbf{W}_C}(\mathbf{w}_C)$. Thus, if $\mathbf{w}_A \in L_A^1, \mathbf{w}_C \in L_C^1$,

$$F(\mathbf{w}_A, \mathbf{w}_C) = (1 - p_A)(1 - p_C)P(\mathbf{U}_A \leq \mathbf{w}_A)P(\mathbf{U}_C \leq \mathbf{w}_C).$$

If $\mathbf{w}_A \in L_A^1, \mathbf{w}_C \in L_C^2$,

$$F(\mathbf{w}_A, \mathbf{w}_C) = (1 - p_A)P(\mathbf{U}_A \leq \mathbf{w}_A)\{(1 - p_C) + p_CP(\mathbf{Y}_C \leq \mathbf{w}_C \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \in L_C^2)\}.$$

If $\mathbf{w}_A \in L_A^2, \mathbf{w}_C \in L_C^1$,

$$F(\mathbf{w}_A, \mathbf{w}_C) = \{(1 - p_A) + p_AP(\mathbf{Y}_A \leq \mathbf{w}_A \mid \mathbf{Y}_A \in L_A^2, \mathbf{Y}_C = \mathbf{0}_C)\}(1 - p_C)P(\mathbf{U}_C \leq \mathbf{w}_C).$$

If $\mathbf{w}_A \in L_A^2, \mathbf{w}_C \in L_C^2$, then

$$\begin{aligned} F(\mathbf{w}_A, \mathbf{w}_C) = \{(1 - p_A) + p_AP(\mathbf{Y}_A \leq \mathbf{w}_A \mid \mathbf{Y}_A \in L_A^2, \mathbf{Y}_C = \mathbf{0}_C)\} \\ \times \{(1 - p_C) + p_CP(\mathbf{Y}_C \leq \mathbf{w}_C \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \in L_C^2)\}. \end{aligned}$$

If the density function of $F(\mathbf{w}_A, \mathbf{w}_C)$ exists, then

$$f(\mathbf{w}_A, \mathbf{w}_C) = \begin{cases} (1 - p_A)(1 - p_C), & \mathbf{w}_A \in L_A^1, \mathbf{w}_C \in L_C^1, \\ \frac{(1 - p_A)p_C}{1 - \alpha} \frac{\partial P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}, & \mathbf{w}_A \in L_A^1, \mathbf{w}_C \in L_C^2, \\ \frac{p_A(1 - p_C)}{\alpha} \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C)}{\partial \mathbf{w}_A}, & \mathbf{w}_A \in L_A^2, \mathbf{w}_C \in L_C^1, \\ \frac{p_AP_C}{\alpha(1 - \alpha)} \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C)}{\partial \mathbf{w}_A} \frac{\partial P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}, & \mathbf{w}_A \in L_A^2, \mathbf{w}_C \in L_C^2. \end{cases}$$

Since the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2)$, then $\partial P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \leq \mathbf{w}_C) / \partial \mathbf{w}_C \neq 0$ for $\mathbf{w}_C \in L_C^2$ and $\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C) / \partial \mathbf{w}_A \neq 0$ for $\mathbf{w}_A \in L_A^2$. Thus, from $f(\mathbf{w}_1, \mathbf{w}_2)$, we have the support of $\mathbf{W} = (\mathbf{W}_A, \mathbf{W}_C)$ is $[0, \infty)^{|A|+|C|}$, which contains Ω . Based on the definition of conditional density in Shorack (2017, §7.4), we have that the conditional density at $(\mathbf{w}_A, \mathbf{0}_C)$, $(\mathbf{w}_A \in L_A^2)$ and $(\mathbf{0}_A, \mathbf{w}_C)$, $(\mathbf{w}_C \in L_C^2)$ given that all the points are on the curve of Ω are

$$\begin{aligned} & \frac{f(\mathbf{w}_A, \mathbf{0}_C)}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{\frac{p_A(1-p_C)}{\alpha} \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C)}{\partial \mathbf{w}_A}}{\frac{p_A(1-p_C)}{\alpha} P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_C = \mathbf{0}_C) + \frac{(1-p_A)p_C}{1-\alpha} P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \in L_C^2)} \\ &= \frac{p_A(1-p_C) \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C)}{\partial \mathbf{w}_A}}{\alpha \{p_A(1-p_C) + (1-p_A)p_C\}}, \end{aligned}$$

and

$$\begin{aligned} & \frac{f(\mathbf{0}_A, \mathbf{w}_C)}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{\frac{(1-p_A)p_C}{1-\alpha} \frac{\partial P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}}{\frac{p_A(1-p_C)}{\alpha} P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_C = \mathbf{0}_C) + \frac{(1-p_A)p_C}{1-\alpha} P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \in L_C^2)} \\ &= \frac{(1-p_A)p_C \frac{\partial P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}}{(1-\alpha) \{p_A(1-p_C) + (1-p_A)p_C\}}. \end{aligned}$$

Then we can get the value of $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C$ or $\mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2)$. When $\mathbf{w}_A \in L_A^1, \mathbf{w}_C \in L_C^1$,

$$P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C \text{ or } \mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2) = 0 = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C).$$

When $\mathbf{w}_A \in L_A^1, \mathbf{w}_C \in L_C^2$,

$$\begin{aligned} & P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C \text{ or } \mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2) \\ &= \frac{\int_{[\mathbf{0}_C, \mathbf{w}_C] \setminus [0,1]^{|C|}} f(\mathbf{0}_A, \mathbf{t}) dt}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{(1-p_A)p_C P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha) \{p_A(1-p_C) + (1-p_A)p_C\}} \\ &= \frac{(1-p_A)p_C P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha) \{p_A(1-p_C) + (1-p_A)p_C\}}, \end{aligned}$$

where the last equation is due to the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_C)$ being $\Omega = (L_A^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_C^2)$. When $\mathbf{w}_A \in L_A^2, \mathbf{w}_C \in L_C^1$,

$$\begin{aligned} & P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C \text{ or } \mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2) \\ &= \frac{\int_{[\mathbf{0}_A, \mathbf{w}_A] \setminus [0,1]^{|A|}} f(\mathbf{t}, \mathbf{0}_C) dt}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{p_A(1-p_C) P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha \{p_A(1-p_C) + (1-p_A)p_C\}} \\ &= \frac{p_A(1-p_C) P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{\alpha \{p_A(1-p_C) + (1-p_A)p_C\}}. \end{aligned}$$

When $\mathbf{w}_A \in L_A^2, \mathbf{w}_C \in L_C^2$, due to $P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C) = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C) + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \leq \mathbf{w}_C)$,

$$\begin{aligned}
& P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C \text{ or } \mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2) \\
&= \frac{\int_{[\mathbf{0}_A, \mathbf{w}_A] \setminus [0,1]^{|A|}} f(\mathbf{t}, \mathbf{0}_C) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{w}_C) d\mathbf{w}_C} + \frac{\int_{[\mathbf{0}_C, \mathbf{w}_C] \setminus [0,1]^{|C|}} f(\mathbf{0}_A, \mathbf{t}) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{w}_C) d\mathbf{w}_C} \\
&= \frac{p_A(1-p_C)P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha\{p_A(1-p_C) + (1-p_A)p_C\}} + \frac{(1-p_A)p_C P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha)\{p_A(1-p_C) + (1-p_A)p_C\}} \\
&= \frac{\{(1-\alpha)p_A(1-p_C) - \alpha(1-p_A)p_C\}P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha(1-\alpha)\{p_A(1-p_C) + (1-p_A)p_C\}} + \frac{(1-p_A)p_C P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha)\{p_A(1-p_C) + (1-p_A)p_C\}}.
\end{aligned}$$

In summary,

$$\begin{aligned}
& P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C \text{ or } \mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2) = \\
& \begin{cases} 0, & \mathbf{w}_A \in L_A^1, \mathbf{w}_C \in L_C^1, \\ \frac{(1-p_A)p_C P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha)\{p_A(1-p_C) + (1-p_A)p_C\}}, & \mathbf{w}_A \in L_A^1, \mathbf{w}_C \in L_C^2, \\ \frac{p_A(1-p_C)P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{\alpha\{p_A(1-p_C) + (1-p_A)p_C\}}, & \mathbf{w}_A \in L_A^2, \mathbf{w}_C \in L_C^1, \\ \frac{\{(1-\alpha)p_A(1-p_C) - \alpha(1-p_A)p_C\}P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha(1-\alpha)\{p_A(1-p_C) + (1-p_A)p_C\}} + \frac{(1-p_A)p_C P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha)\{p_A(1-p_C) + (1-p_A)p_C\}}, & \mathbf{w}_A \in L_A^2, \mathbf{w}_C \in L_C^2. \end{cases} \quad (29)
\end{aligned}$$

From (29), $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C \text{ or } \mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2) = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C)$, if and only if

$$\begin{cases} \frac{(1-p_A)p_C}{(1-\alpha)\{p_A(1-p_C) + (1-p_A)p_C\}} = 1, \\ \frac{p_A(1-p_C)}{\alpha\{p_A(1-p_C) + (1-p_A)p_C\}} = 1, \\ \frac{\{(1-\alpha)p_A(1-p_C) - \alpha(1-p_A)p_C\}}{\alpha(1-\alpha)\{p_A(1-p_C) + (1-p_A)p_C\}} = 0, \end{cases}$$

or equivalently,

$$\alpha = \frac{p_A(1-p_C)}{p_A(1-p_C) + (1-p_A)p_C}. \quad (30)$$

We have hence shown that under condition (30), $(\mathbf{W} \mid \mathbf{W} \in \Omega) \stackrel{d}{=} \mathbf{Y}$ if the density function of $F(\mathbf{w}_A, \mathbf{w}_C)$ exists.

If the density function of $F(\mathbf{w}_A, \mathbf{w}_C)$ does not exist, from the definition of conditional probability in Shorack (2017, §7.4), we know that (29) still satisfies the condition for conditional probability. Thus, we can define $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C \text{ or } \mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2)$ as that in (29). Under condition (30), we also have $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_C = \mathbf{0}_C \text{ or } \mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_C \in L_C^2) = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_C \leq \mathbf{w}_C)$. Then $(\mathbf{W} \mid \mathbf{W} \in \Omega) \stackrel{d}{=} \mathbf{Y}$.

In summary, based on the definition of $\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C$, we have $\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C$. □

2.4 Proof of Proposition 2 for a general B

Proof. We need to show

$$\begin{aligned}
& \text{the support of } \mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C) \text{ is } \Omega = (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2) \\
& \Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \mid \mathbf{Y}_B, \quad (31)
\end{aligned}$$

the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$
 $\Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C \mid \mathbf{Y}_B$, (32)

the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$
 $\Rightarrow \mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \mid \mathbf{Y}_B$. (33)

Under (31), (32) and (33), we can have if the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$, then

$$\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \mid \mathbf{Y}_B \Leftrightarrow \mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \mid \mathbf{Y}_B \Leftrightarrow \mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C \mid \mathbf{Y}_B.$$

First, we show (31). Since the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$, then $S_A \times S_B \times S_C \subset (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C)$, $(\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C)$ or $(\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$. When $S_A \times S_B \times S_C \subset (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C)$, $\mathbf{Y}_B = \mathbf{0}_B$. Given $\mathbf{Y}_B = \mathbf{0}_B$, since $\mathbf{Y}_C = \mathbf{0}_C$ regardless of the value of \mathbf{Y}_A , then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y}_B, \mathbf{Y} \in S_A \times S_B \times S_C)$. When $S_A \times S_B \times S_C \subset (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C)$, $\mathbf{Y}_B \in L_B^2$. Given $\mathbf{Y}_B = \mathbf{y}_B$, since $\mathbf{Y}_A = \mathbf{0}_A$ and $\mathbf{Y}_C = \mathbf{0}_C$, then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y}_B, \mathbf{Y} \in S_A \times S_B \times S_C)$. When $S_A \times S_B \times S_C \subset (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$, $\mathbf{Y}_B = \mathbf{0}_B$. Given $\mathbf{Y}_B = \mathbf{0}_B$, since $\mathbf{Y}_A = \mathbf{0}_A$ regardless of the value of \mathbf{Y}_C , then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y}_B, \mathbf{Y} \in S_A \times S_B \times S_C)$. Thus, based on the definition of $\mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \mid \mathbf{Y}_B$, we have $\mathbf{Y}_A \perp\!\!\!\perp_i \mathbf{Y}_C \mid \mathbf{Y}_B$.

Second, we show (32). Since the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$, then when $Y_k > 1$ for some $k \in A$, we have $\mathbf{Y} \in (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C)$ and $\mathbf{Y}_B = \mathbf{0}_B$. Given $\mathbf{Y}_B = \mathbf{0}_B$ and $Y_k > 1$, since $\mathbf{Y}_C = \mathbf{0}_C$ regardless of the value of \mathbf{Y}_A , then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y}_B, Y_k > 1)$. When $Y_k > 1$ for some $k \in B$, we have $\mathbf{Y} \in (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C)$ and $\mathbf{Y}_B \in L_B^2$. Given $\mathbf{Y}_B = \mathbf{y}_B$ and $Y_k > 1$, since $\mathbf{Y}_A = \mathbf{0}_A$ and $\mathbf{Y}_C = \mathbf{0}_C$, then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y}_B, Y_k > 1)$. When $Y_k > 1$ for some $k \in C$, we have $\mathbf{Y} \in (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$ and $\mathbf{Y}_B = \mathbf{0}_B$. Given $\mathbf{Y}_B = \mathbf{0}_B$ and $Y_k > 1$, since $\mathbf{Y}_A = \mathbf{0}_A$ regardless of the value of \mathbf{Y}_C , then $\mathbf{Y}_A \perp\!\!\!\perp \mathbf{Y}_C \mid (\mathbf{Y}_B, Y_k > 1)$. Thus, based on the definition of $\mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C \mid \mathbf{Y}_B$, we have $\mathbf{Y}_A \perp\!\!\!\perp_e \mathbf{Y}_C \mid \mathbf{Y}_B$.

Third, we show (33). Since the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$, then

$$\begin{aligned} P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C \text{ or } \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \in L_B^2, \mathbf{Y}_C = \mathbf{0}_C \text{ or } \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \in L_C^2) \\ = P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C) + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \in L_B^2, \mathbf{Y}_C = \mathbf{0}_C) \\ + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \in L_C^2) = 1. \end{aligned}$$

Suppose that $P(\mathbf{Y}_A \in L_A^2, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C) = \alpha_1$ and $P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \in L_B^2, \mathbf{Y}_C = \mathbf{0}_C) = \alpha_2$, then $P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \in L_C^2) = 1 - \alpha_1 - \alpha_2$, where α_1 and α_2 satisfy the conditions of $\alpha_1 > 0, \alpha_2 > 0$ and $0 < \alpha_1 + \alpha_2 < 1$. Now we introduce three new random vectors $\mathbf{W}_A, \mathbf{W}_B$ and \mathbf{W}_C , assume that they have the distributions as follows:

$$F_{\mathbf{W}_A}(\mathbf{w}_A) = \begin{cases} (1 - p_A)P(\mathbf{U}_A \leq \mathbf{w}_A), & \mathbf{w}_A \in L_A^1, \\ (1 - p_A) + p_AP(\mathbf{Y}_A \leq \mathbf{w}_A \mid \mathbf{Y}_A \in L_A^2, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C), & \mathbf{w}_A \in L_A^2, \end{cases}$$

$$F_{\mathbf{W}_B}(\mathbf{w}_B) = \begin{cases} (1 - p_B)P(\mathbf{U}_B \leq \mathbf{w}_B), & \mathbf{w}_B \in L_B^1, \\ (1 - p_B) + p_BP(\mathbf{Y}_B \leq \mathbf{w}_B \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \in L_B^2, \mathbf{Y}_C = \mathbf{0}_C), & \mathbf{w}_B \in L_B^2, \end{cases}$$

and

$$F_{\mathbf{W}_C}(\mathbf{w}_C) = \begin{cases} (1 - p_C)P(\mathbf{U}_C \leq \mathbf{w}_C), & \mathbf{w}_C \in L_C^1, \\ (1 - p_C) + p_CP(\mathbf{Y}_C \leq \mathbf{w}_C \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \in L_C^2), & \mathbf{w}_C \in L_C^2, \end{cases}$$

where $p_A \in (0, 1)$, $p_B \in (0, 1)$, $p_C \in (0, 1)$, and $\mathbf{U}_A, \mathbf{U}_B, \mathbf{U}_C$ are random vectors which follow the uniform distributions in L_A^1, L_B^1 and L_C^1 , respectively.

To make $\mathbf{W}_A, \mathbf{W}_B, \mathbf{W}_C$ independent, let $F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = F_{\mathbf{W}_A}(\mathbf{w}_A)F_{\mathbf{W}_B}(\mathbf{w}_B)F_{\mathbf{W}_C}(\mathbf{w}_C)$. Furthermore, we have $\mathbf{W}_A \perp \mathbf{W}_C \mid \mathbf{W}_B$. Besides, if $\mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^1$,

$$F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = (1 - p_A)(1 - p_B)(1 - p_C)P(\mathbf{U}_A \leq \mathbf{w}_A)P(\mathbf{U}_B \leq \mathbf{w}_B)P(\mathbf{U}_C \leq \mathbf{w}_C).$$

If $\mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^2$,

$$F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = (1 - p_A)(1 - p_B)P(\mathbf{U}_A \leq \mathbf{w}_A)P(\mathbf{U}_B \leq \mathbf{w}_B) \\ \times \{(1 - p_C) + p_C P(\mathbf{Y}_C \leq \mathbf{w}_C \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \in L_C^2)\}.$$

If $\mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^1$,

$$F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = (1 - p_A)(1 - p_C)P(\mathbf{U}_A \leq \mathbf{w}_A)P(\mathbf{U}_C \leq \mathbf{w}_C) \\ \times \{(1 - p_B) + p_B P(\mathbf{Y}_B \leq \mathbf{w}_B \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \in L_B^2, \mathbf{Y}_C = \mathbf{0}_C)\}.$$

If $\mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^2$,

$$F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = (1 - p_A)P(\mathbf{U}_A \leq \mathbf{w}_A)\{(1 - p_B) + p_B P(\mathbf{Y}_B \leq \mathbf{w}_B \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \in L_B^2, \mathbf{Y}_C = \mathbf{0}_C)\} \\ \times \{(1 - p_C) + p_C P(\mathbf{Y}_C \leq \mathbf{w}_C \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \in L_C^2)\}.$$

If $\mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^1$,

$$F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = (1 - p_B)(1 - p_C)P(\mathbf{U}_B \leq \mathbf{w}_B)P(\mathbf{U}_C \leq \mathbf{w}_C) \\ \times \{(1 - p_A) + p_A P(\mathbf{Y}_A \leq \mathbf{w}_A \mid \mathbf{Y}_A \in L_A^2, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C)\}.$$

If $\mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^2$,

$$F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = (1 - p_B)P(\mathbf{U}_B \leq \mathbf{w}_B)\{(1 - p_A) + p_A P(\mathbf{Y}_A \leq \mathbf{w}_A \mid \mathbf{Y}_A \in L_A^2, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C)\} \\ \times \{(1 - p_C) + p_C P(\mathbf{Y}_C \leq \mathbf{w}_C \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \in L_C^2)\}.$$

If $\mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^1$,

$$F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = (1 - p_C)P(\mathbf{U}_C \leq \mathbf{w}_C)\{(1 - p_A) + p_A P(\mathbf{Y}_A \leq \mathbf{w}_A \mid \mathbf{Y}_A \in L_A^2, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C)\} \\ \times \{(1 - p_B) + p_B P(\mathbf{Y}_B \leq \mathbf{w}_B \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \in L_B^2, \mathbf{Y}_C = \mathbf{0}_C)\}.$$

If $\mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^2$,

$$F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = \{(1 - p_A) + p_A P(\mathbf{Y}_A \leq \mathbf{w}_A \mid \mathbf{Y}_A \in L_A^2, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C)\} \\ \times \{(1 - p_B) + p_B P(\mathbf{Y}_B \leq \mathbf{w}_B \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \in L_B^2, \mathbf{Y}_C = \mathbf{0}_C)\} \\ \times \{(1 - p_C) + p_C P(\mathbf{Y}_C \leq \mathbf{w}_C \mid \mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \in L_C^2)\}.$$

If the density function of $F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C)$ exists, then by calculation

$$f(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C) = \left\{ \begin{array}{ll} (1-p_A)(1-p_B)(1-p_C), & \mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^1, \\ \frac{(1-p_A)(1-p_B)p_C}{1-\alpha_1-\alpha_2} \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}, & \mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^2, \\ \frac{(1-p_A)p_B(1-p_C)}{\alpha_2} \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_B}, & \mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^1, \\ \frac{(1-p_A)p_B p_C}{\alpha_2(1-\alpha_1-\alpha_2)} \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_B} \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}, & \mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^2, \\ \frac{p_A(1-p_B)(1-p_C)}{\alpha_1} \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_A}, & \mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^1, \\ \frac{p_A(1-p_B)p_C}{\alpha_1(1-\alpha_1-\alpha_2)} \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_A} \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}, & \mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^2, \\ \frac{p_A p_B(1-p_C)}{\alpha_1 \alpha_2} \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_A} \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_B}, & \mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^1, \\ \frac{p_A p_B p_C}{\alpha_1 \alpha_2(1-\alpha_1-\alpha_2)} \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_A} \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_B} \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}, & \mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^2. \end{array} \right.$$

Since the support of $\mathbf{Y} = (\mathbf{Y}_A, \mathbf{Y}_B, \mathbf{Y}_C)$ is $\Omega = (L_A^2 \times \mathbf{0}_B \times \mathbf{0}_C) \cup (\mathbf{0}_A \times L_B^2 \times \mathbf{0}_C) \cup (\mathbf{0}_A \times \mathbf{0}_B \times L_C^2)$, then $\partial P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C) / \partial \mathbf{w}_C \neq 0$ for $\mathbf{w}_C \in L_C^2$, $\partial P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C = \mathbf{0}_C) / \partial \mathbf{w}_B \neq 0$ for $\mathbf{w}_B \in L_B^2$ and $\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C) / \partial \mathbf{w}_A \neq 0$ for $\mathbf{w}_A \in L_A^2$. Thus, from $f(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C)$, we can obtain that the support of $\mathbf{W} = (\mathbf{W}_A, \mathbf{W}_B, \mathbf{W}_C)$ is $[0, \infty)^{|A|+|B|+|C|}$, which contains Ω . Based on the definition of conditional density in Shorack (2017, §7.4), if $\mathbf{w}_A \in L_A^2$, $\mathbf{w}_B \in L_B^2$ and $\mathbf{w}_C \in L_C^2$, then we have that the conditional densities at $(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C)$, $(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C)$ and $(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C)$, given that all the points are on the curve of Ω are

$$\begin{aligned} & \frac{f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C)}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{p_A(1-p_B)(1-p_C) \frac{\partial P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_A}}{\alpha_1 \{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}, \\ & \frac{f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C)}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{(1-p_A)p_B(1-p_C) \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C=\mathbf{0}_C)}{\partial \mathbf{w}_B}}{\alpha_2 \{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}, \end{aligned}$$

and

$$\begin{aligned} & \frac{f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C)}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{(1-p_A)(1-p_B)p_C \frac{\partial P(\mathbf{Y}_A=\mathbf{0}_A, \mathbf{Y}_B=\mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\partial \mathbf{w}_C}}{(1-\alpha_1-\alpha_2) \{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}. \end{aligned}$$

Then we can get the value of $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W}_A \in L_A^2, \mathbf{W}_B = \mathbf{0}_B, \mathbf{W}_C = \mathbf{0}_C$ or $\mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_B \in L_B^2, \mathbf{W}_C = \mathbf{0}_C$ or $\mathbf{W}_A = \mathbf{0}_A, \mathbf{W}_B = \mathbf{0}_B, \mathbf{W}_C \in L_C^2) = P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega)$. When $\mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^1$,

$$P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) = 0 = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C).$$

When $\mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^2$,

$$\begin{aligned}
& P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) \\
&= \frac{\int_{[\mathbf{0}_C, \mathbf{w}_C] \setminus [0,1]^{|C|}} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{t}) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\
&= \frac{(1-p_A)(1-p_B)p_C P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha_1-\alpha_2)\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}} \\
&= \frac{(1-p_A)(1-p_B)p_C P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha_1-\alpha_2)\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}.
\end{aligned}$$

When $\mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^1$,

$$\begin{aligned}
& P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) \\
&= \frac{\int_{[\mathbf{0}_B, \mathbf{w}_B] \setminus [0,1]^{|B|}} f(\mathbf{0}_A, \mathbf{t}, \mathbf{0}_C) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\
&= \frac{(1-p_A)p_B(1-p_C) P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha_2\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}} \\
&= \frac{(1-p_A)p_B(1-p_C) P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\alpha_2\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}.
\end{aligned}$$

When $\mathbf{w}_A \in L_A^1, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^2$, due to $P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C) = P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C = \mathbf{0}_C) + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C)$,

$$\begin{aligned}
& P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) \\
&= \frac{\int_{[\mathbf{0}_B, \mathbf{w}_B] \setminus [0,1]^{|B|}} f(\mathbf{0}_A, \mathbf{t}, \mathbf{0}_C) d\mathbf{t} + \int_{[\mathbf{0}_C, \mathbf{w}_C] \setminus [0,1]^{|C|}} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{t}) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\
&= \frac{\{(1-p_A)p_B(1-p_C)(1-\alpha_1-\alpha_2) - (1-p_A)(1-p_B)p_C\alpha_2\} P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha_2(1-\alpha_1-\alpha_2)\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}} \\
&+ \frac{(1-p_A)(1-p_B)p_C P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha_1-\alpha_2)\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}.
\end{aligned}$$

When $\mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^1$,

$$\begin{aligned}
& P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) \\
&= \frac{\int_{[\mathbf{0}_A, \mathbf{w}_A] \setminus [0,1]^{|A|}} f(\mathbf{t}, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\
&= \frac{p_A(1-p_B)(1-p_C) P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha_1\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}} \\
&= \frac{p_A(1-p_B)(1-p_C) P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\alpha_1\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}.
\end{aligned}$$

When $\mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^1, \mathbf{w}_C \in L_C^2$, due to $P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C) = P(\mathbf{Y}_A \leq$

$$\mathbf{w}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C) + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C),$$

$$\begin{aligned} & P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) \\ &= \frac{\int_{[\mathbf{0}_A, \mathbf{w}_A] \setminus [0,1]^{|A|}} f(\mathbf{t}, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{t} + \int_{[\mathbf{0}_C, \mathbf{w}_C] \setminus [0,1]^{|C|}} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{t}) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{\{p_A(1-p_B)(1-p_C)(1-\alpha_1-\alpha_2) - (1-p_A)(1-p_B)p_C\alpha_1\}P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha_1(1-\alpha_1-\alpha_2)\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}} \\ &\quad + \frac{(1-p_A)(1-p_B)p_C P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{(1-\alpha_1-\alpha_2)\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}. \end{aligned}$$

When $\mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^1$, due to $P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C) = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C) + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C = \mathbf{0}_C)$,

$$\begin{aligned} & P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) \\ &= \frac{\int_{[\mathbf{0}_A, \mathbf{w}_A] \setminus [0,1]^{|A|}} f(\mathbf{t}, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{t} + \int_{[\mathbf{0}_B, \mathbf{w}_B] \setminus [0,1]^{|B|}} f(\mathbf{0}_A, \mathbf{t}, \mathbf{0}_C) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \frac{\{p_A(1-p_B)(1-p_C)\alpha_2 - (1-p_A)p_B(1-p_C)\alpha_1\}P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha_1\alpha_2\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}} \\ &\quad + \frac{(1-p_A)p_B(1-p_C)P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{\alpha_2\{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C\}}. \end{aligned}$$

When $\mathbf{w}_A \in L_A^2, \mathbf{w}_B \in L_B^2, \mathbf{w}_C \in L_C^2$, due to $P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C) = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C) + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C = \mathbf{0}_C) + P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C \leq \mathbf{w}_C)$,

$$\begin{aligned} & P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) \\ &= \frac{\int_{[\mathbf{0}_A, \mathbf{w}_A] \setminus [0,1]^{|A|}} f(\mathbf{t}, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{t} + \int_{[\mathbf{0}_B, \mathbf{w}_B] \setminus [0,1]^{|B|}} f(\mathbf{0}_A, \mathbf{t}, \mathbf{0}_C) d\mathbf{t} + \int_{[\mathbf{0}_C, \mathbf{w}_C] \setminus [0,1]^{|C|}} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{t}) d\mathbf{t}}{\int_{L_A^2} f(\mathbf{w}_A, \mathbf{0}_B, \mathbf{0}_C) d\mathbf{w}_A + \int_{L_B^2} f(\mathbf{0}_A, \mathbf{w}_B, \mathbf{0}_C) d\mathbf{w}_B + \int_{L_C^2} f(\mathbf{0}_A, \mathbf{0}_B, \mathbf{w}_C) d\mathbf{w}_C} \\ &= \left[\frac{\{p_A(1-p_B)(1-p_C)(1-\alpha_1-\alpha_2) - (1-p_A)(1-p_B)p_C\alpha_1\}P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B = \mathbf{0}_B, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha_1(1-\alpha_1-\alpha_2)} \right. \\ &\quad + \frac{\{(1-p_A)p_B(1-p_C)(1-\alpha_1-\alpha_2) - (1-p_A)(1-p_B)p_C\alpha_2\}P(\mathbf{Y}_A = \mathbf{0}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C = \mathbf{0}_C)}{\alpha_2(1-\alpha_1-\alpha_2)} \\ &\quad \left. + \frac{(1-p_A)(1-p_B)p_C P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)}{1-\alpha_1-\alpha_2} \right] \\ &\quad \times \frac{1}{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C}. \end{aligned}$$

By comparison, we can find that $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)$ if and only if

$$\begin{cases} \alpha_1 = \frac{p_A(1-p_B)(1-p_C)}{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C}, \\ \alpha_2 = \frac{(1-p_A)p_B(1-p_C)}{p_A(1-p_B)(1-p_C) + (1-p_A)p_B(1-p_C) + (1-p_A)(1-p_B)p_C}. \end{cases} \quad (34)$$

We have hence shown that under condition (34), $(\mathbf{W} \mid \mathbf{W} \in \Omega) \stackrel{d}{=} \mathbf{Y}$ if the density function of $F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C)$ exists.

If the density function of $F(\mathbf{w}_A, \mathbf{w}_B, \mathbf{w}_C)$ does not exist, from the definition of conditional probability in Shorack (2017, §7.4), we know that the above formulas for $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega)$ still satisfy the condition for conditional probability. Thus, we still can define $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega)$ in the same way as before. Under condition (34), we also have $P(\mathbf{W}_A \leq \mathbf{w}_A, \mathbf{W}_B \leq \mathbf{w}_B, \mathbf{W}_C \leq \mathbf{w}_C \mid \mathbf{W} \in \Omega) = P(\mathbf{Y}_A \leq \mathbf{w}_A, \mathbf{Y}_B \leq \mathbf{w}_B, \mathbf{Y}_C \leq \mathbf{w}_C)$. Then $(\mathbf{W} \mid \mathbf{W} \in \Omega) \stackrel{d}{=} \mathbf{Y}$.

In summary, according to the definition of $\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \mid \mathbf{Y}_B$, we have $\mathbf{Y}_A \perp\!\!\!\perp_o \mathbf{Y}_C \mid \mathbf{Y}_B$. □

References

- Engelke, S. and Hitz, A. S. (2020). Graphical models for extremes. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, page (forthcoming).
- Shorack, G. R. (2017). *Probability for Statisticians*. Springer.