

Frustrated Potts: Multiplicity Eliminates Chaos via Reentrance

Alpar Türkoğlu¹ and A. Nihat Berker^{2,3}

¹*Department of Electrical and Electronics Engineering,
Boğaziçi University, Bebek, İstanbul 34342, Turkey*

²*Faculty of Engineering and Natural Sciences, Kadir Has University, Cibali, İstanbul 34083, Turkey*

³*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

The frustrated q -state Potts model is solved exactly on a hierarchical lattice, yielding chaos under rescaling, namely the signature of a spin-glass phase, as previously seen for the Ising ($q = 2$) model. However, the ground-state entropy introduced by the ($q > 2$)-state antiferromagnetic Potts bond induces an escape from chaos as multiplicity q increases. The frustration versus multiplicity phase diagram has a reentrant (as a function of frustration) chaotic phase.

Frustration [1], meaning loops of equal-strength interactions that cannot all be simultaneously satisfied, diminishes ordered phases in the phase diagram and may drastically change the nature of certain ordered phases.[2] For example, the so-called Mattis phase [3], where spins are ordered in random directions but the system has no competing interactions between the spins and thus all interactions can be simultaneously satisfied, becomes a spin-glass phase with the introduction of the smallest amount of competing frustrated interactions [4], with residual entropy, unsaturated order at zero temperature, and the chaotic rescaling of the interactions, as measured by a positive Lyapunov exponent. The latter, chaos under scale change, is the signature of the spin-glass phase [5–13], as gauged quantitatively by the Lyapunov exponent. Chaotic interactions under scale change dictate chaotic correlation functions as a function of distance.[14]




FIG. 1. Top row: Construction of a hierarchical lattice, from Ref.[15]. The renormalization-group solution of a hierarchical lattice proceeds via renormalization group in the opposite direction of its construction. Middle row: The two units used in the construction of the frustrated hierarchical lattice.[5] On the left is the frustrated unit and on the right the repressed unit. In the former, p parallel bonds connect the internal spins s_3 and s_4 ($p = 4$ is shown here). The wiggly bonds are infinitely strong antiferromagnetic couplings. Bottom row: The assemblage of the units for the construction of the frustrated hierarchical lattice.

In fact, chaos under rescaling was seen in frustrated

systems, with no randomness, with the exact solution of hierarchical lattices [5–8]. When frustration is increased, a sequence of period doublings leads to the regime of spin-glass chaos [5]. The regime of period doublings that precedes spin-glass chaos also converts, under randomness, to chaotic bands [6]. Spin-glass chaos and its positive Lyapunov exponent was also calculated in the renormalization-group solution of cubic systems where frustration is introduced by quenched randomness.[4]




FIG. 2. Lower panel: The onset of chaos, by period doubling, under increased frustration for the $q = 3$ -state Potts model. For each p_b , the renormalization-group flows are in the vertical direction. The full lines represent attractive fixed points, limit cycles, and chaotic bands. The dashed lines represent the unstable fixed points, only some of which are shown. The $\tanh(K) = 0$ points are the stable fixed points which are the sinks of the disordered phase. The upper panel shows the calculated Lyapunov exponents. The upper inset shows the chaotic renormalization-group trajectory for $p_b = 28$.

Another important cause of ground-state degeneracy is in the ground-state participation of a multiplicity of spin states, as seen in antiferromagnetic ($q > 2$)-state Potts models [16–18]. In the latter systems, antiferromagnetic Potts interactions can all be satisfied, while local spins can take many different values, leading to a distinctive phase [16–18]. In the current work, we have studied the combination of both effects, chaos from frustration and degeneracy from multiplicity of states. We have exactly solved the q -state Potts models on the frustrated hierarchical model as in Ref.[5]. As seen below and in Fig. 1, hierarchical models are models obtained by self-embedding a graph infinitely many times and are exactly solvable by renormalization-group theory [15, 19, 20]. We find that the system escapes chaos through multiplicity and that chaos shows reentrant behavior.




FIG. 3. The $q = 6$ -state Potts model, under increased frustration, after the onset of chaos, leaves chaos through a set of reverse period doublings. The upper inset shows the chaotic renormalization-group trajectory for $p_b = 42$.

The model is constructed, by combining two units, embodying the different microscopic effects of competing interactions. The Hamiltonian of the system is

$$-\beta\mathcal{H} = K \sum_{\langle ij \rangle} \delta(s_i, s_j), \quad (1)$$

where $\beta = 1/k_B T$, at site i the Potts spin $s_i = 1, 2, \dots, q$ can be in q different states, the delta function $\delta(s_i, s_j) = 1(0)$ for $s_i = s_j (s_i \neq s_j)$, and the sum is over all interacting pairs of spins, represented by straight lines in Fig. 1. In unit C of Fig. 1, the correlations are repressed but not eliminated along the path of the unit, as $m_2 > m_1$ and




FIG. 4. The $q = 8$ -state Potts model, under increased frustration, through a set of period doublings followed by reverse period doublings, bypasses chaos.




FIG. 5. As the number q of Potts states increases, the system leaves chaos through reverse period doublings. Note the horizontal scale change in the figure at $q = 15$. For this calculation, $p_b = 40$. The upper inset shows the chaotic renormalization-group trajectory for $q = 5$.

the competing correlation on the longer unit is weaker. Unit B is frustrated: Both interaction loops across the unit are frustrated. The above are representative of the two qualitatively different effects of competing interactions in d -dimensional hypercubic lattices [5]. This competition occurs when, in two different paths connecting two points on the lattice, all interactions cannot be simultaneously satisfied.

The recursion relations of a hierarchical lattice are obtained by doing the partition function sum of the spins internal to the unit (in black in Fig. 1) while keeping the two external spins (in white in Fig. 1) fixed [15, 19, 20]. Thus, the two strands in unit C are disconnected with respect to this summation. If the two paths are of unequal length, the effective overall interaction along the longer path is weaker (as can be seen, e.g., from renormalization-group transformations on $d = 1$ chains), so that the overall interaction along the shorter path dominates but is weakened (repressed) by the parallel presence of the competing longer path. For this, it is important that one of the paths has an anti-aligning wiggly bond, to induce competition rather than cooperation. On the other hand, when the two paths are of equal length, a frustrated loop is generated and the effective (*i.e.*, renormalized) interactions along the two paths cancel out (are frustrated). Across unit B in Fig. 1 are two adjacent frustrated loops, each with one wiggly (anti-aligning) interaction, so that this unit is frustrated. Incidentally, this discussion shows that loops are essential to chaotic spin-glass behavior. Although the discussion here was given in context of d -dimensional hypercubic lattices, it is also applicable to scale-free networks of physical and social nature [21, 22].

By combining in parallel p_b and p_c units B and C , a family of models is created. In this work, we present results for $m_1 = 2, m_2 = 3, p = 4, p_c = 1$ and varying the number p_b of frustrated units. However, we have also explored a wide range of model parameters m_1, m_2, p, p_c and find that our chaos and reentrance results are generic.

Hierarchical models are solved exactly, by renormalization-group theory, proceeding in the reverse direction of the construction of the hierarchical model by decimating the interior spins of each level (black circles in Fig. 1), obtaining recursion relations $K' = K'(K)$ exactly.[15, 19, 20]. The onset of the chaotic bands, as the number p_b of frustrated units is increased, is shown in Fig. 2, for the number of Potts states $q = 3$. The chaotic band is an interaction interval in which, under the renormalization-group transformations, the points are visited in a chaotic sequence. It has been shown that chaotic interactions under renormalization-group dictate chaotic correlations between spins as a function of distance [14].

A measure of the strength of chaos are the calculated Lyapunov exponents [23, 24],

$$\lambda = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \ln \left| \frac{dK_{k+1}}{dK_k} \right|, \quad (2)$$




FIG. 6. The calculated frustration-multiplicity chaos phase diagram of the system, shows the (q, p_b) combinations for which chaos occurs. NC stands for non-chaos. The vertical scale changes at $p_b = 100$. Note the reentrance, as a function of frustration, of non-chaos. All the points of this phase diagram have been calculated.

where K_k is the interaction at step k of the renormalization-group trajectory. Thus, the Lyapunov exponent is calculated by logarithmically summing [23, 24] the derivative dK_{k+1}/dK_k of the interaction constant K_{k+1} after the $(k+1)$ th renormalization, with respect to the interaction constant K_k after the (k) th renormalization. The sum in Eq.(2) is to be taken within the asymptotic trajectory, so that we throw out the first 100 renormalization-group iterations to eliminate the transient points and subsequently use 600 iterations in the sum in Eq.(2), which assures convergence in the chaotic bands. It is seen that the Lyapunov exponent is non-positive outside chaos, barely touching zero at each period doubling.

A qualitative understanding of this chaos phenomenon is that strong interactions when frustrated generate weak interactions at the next length scale of the renormalization-group transformation. Thus, renormalization-group flows inside the ordered phase cannot go to a strong-coupling sink [25] as happens in non-frustrated systems. This mechanism and phenomenon occur in hierarchical as well as non-hierarchical lattices, as is seen in hypercubic lattices [4] and as was eventually accepted (subsequently to [5]) for spin-glass phases in general [5–13].

For $q = 6$ in Fig. 3, as frustration is increased, the system leaves chaos through a series of reverse period doublings (foldings). We thus have non-chaos reentrance [26] around the chaotic phase. For $q = 8$, shown in Fig. 4, doublings and foldings succeed, but chaos has disappeared. We can therefore look for reverse bifurcations as the number of Potts states q is increased for fixed frustration. This is seen in Fig. 5, where it is indeed seen that chaos disappears as the antiferromagnetic degeneracy of the Potts model is increased by increasing q .

The complete frustration versus multiplicity phase diagram has been calculated and is given in Fig. 6, clearly showing reentrance. Note the stability of the chaotic

phase around $q = 3$.

The disappearance of the spin-glass phase under increased frustration has been seen in $q = 2$ Ising models that are microscopically rewired, in $d = 2, 3$, by a renormalization-group study similar to our current work [4]. The full double reentrance as a function of frustration and q needs to be studied by other methods, such as Monte Carlo sampling [27]. The ousting of frustrated chaos by the multiplicity of local states could have a relevance to the mathematical Ising-type modeling of societal collective behavior [28]. Such studies typically assign Ising variables ($q = 2$) to individuals based on opinions and inclinations. Frustrated closed loops of interactions will

then lead to chaotic rescaling behavior, which implies large instabilities with respect to small changes in external parameters. Perhaps this is artificial, as individuals can be in more than two possible states, and chaos can be avoided as seen in this paper.

ACKNOWLEDGMENTS

Support by the Academy of Sciences of Turkey (TÜBA) is gratefully acknowledged.

-
- [1] G. Toulouse, Theory of frustration effect in spin glasses I., *Commun. Phys.* **2**, 115 (1977).
- [2] While frustration reduces the ordered phases in favor of the disordered phase, the spin-glass phase created by frustration replaces (as with the Mattis phase) or reduces (as with the ferromagnetic phase) conventionally ordered phases. See G. Gülpınar and A. N. Berker, Quenched-vacancy induced spin-glass order, *Phys. Rev. E* **79**, 021110 (2009).
- [3] D.C. Mattis, Solvable spin systems with random interactions, *Phys. Lett. A* **56**, 421 (1976).
- [4] E. Ilker and A. N. Berker, Overfrustrated and underfrustrated spin glasses in $d=3$ and 2 : Evolution of phase diagrams and chaos including spin-glass order in $d=2$, *Phys. Rev. E* **89**, 042139 (2014).
- [5] S. R. McKay, A. N. Berker, and S. Kirkpatrick, Spin-glass behavior in frustrated Ising models with chaotic renormalization-group trajectories, *Phys. Rev. Lett.* **48**, 767 (1982).
- [6] S. R. McKay, A. N. Berker, and S. Kirkpatrick, Amorphously packed, frustrated hierarchical models: Chaotic rescaling and spin-glass behavior, *J. Appl. Phys.* **53**, 7974 (1982).
- [7] A. N. Berker and S. R. McKay, Hierarchical models and chaotic spin glasses, *J. Stat. Phys.* **36**, 787 (1984).
- [8] E. J. Hartford and S. R. McKay, Ising spin-glass critical and multicritical fixed distributions from a renormalization-group calculation with quenched randomness, *J. Appl. Phys.* **70**, 6068 (1991).
- [9] Z. Zhu, A. J. Ochoa, S. Schnabel, F. Hamze, and H. G. Katzgraber, Best-case performance of quantum annealers on native spin-glass benchmarks: How chaos can affect success probabilities, *Phys. Rev. A* **93**, 012317 (2016).
- [10] W. Wang, J. Machta, and H. G. Katzgraber, Bond chaos in spin glasses revealed through thermal boundary conditions, *Phys. Rev. B* **93**, 224414 (2016).
- [11] L. A. Fernandez, E. Marinari, V. Martin-Mayor, G. Parisi, and D. Yllanes, Temperature chaos is a non-local effect, *J. Stat. Mech. - Theory and Experiment*, 123301 (2016).
- [12] A. Billoire, L. A. Fernandez, A. Maiorano, E. Marinari, V. Martin-Mayor, J. Moreno-Gordo, G. Parisi, F. Ricci-Tersenghi, J.J. Ruiz-Lorenzo, Dynamic variational study of chaos: Spin glasses in three dimensions, *J. Stat. Mech. - Theory and Experiment*, 033302 (2018).
- [13] R. Eldan, The Sherrington-Kirkpatrick spin glass exhibits chaos, arXiv:2004.14885 (2020)
- [14] N. Aral and A. N. Berker, Chaotic spin correlations in frustrated Ising hierarchical lattices, *Phys. Rev. B* **79**, 014434 (2009).
- [15] A. N. Berker and S. Ostlund, Renormalisation-group calculations of finite systems: Order parameter and specific heat for epitaxial ordering, *J. Phys. C* **12**, 4961 (1979).
- [16] A. N. Berker and L. P. Kadanoff, Ground-state entropy and algebraic order at low temperatures, *J. Phys. A* **13**, L259 (1980).
- [17] A. N. Berker and L. P. Kadanoff, corrigendum, *J. Phys. A* **13**, 3786 (1980).
- [18] H. Saleur, The antiferromagnetic Potts model in 2 dimensions: Berker-Kadanoff phase, antiferromagnetic transition, and the role of Beraha numbers, *Nuc. Phys. B* **360**, 219 (1991).
- [19] R. B. Griffiths and M. Kaufman, Spin systems on hierarchical lattices: Introduction and thermodynamic limit, *Phys. Rev. B* **26**, 5022R (1982).
- [20] M. Kaufman and R. B. Griffiths, Spin systems on hierarchical lattices: 2. Some examples of soluble models, *Phys. Rev. B* **30**, 244 (1984).
- [21] D. Eroğlu, J. S. W. Lamb, and T. Pereira, Synchronisation of chaos and its applications, *Contemp. Phys.* **58**, 207 (2017).
- [22] D. Eroğlu, M. Tanzi, S. van Strien, and T. Pereira, Revealing dynamics, communities, and criticality from data, *Phys. Rev. X* **10** 021047 (2020).
- [23] P. Collet and J.-P. Eckmann, *Iterated Maps on the Interval as Dynamical Systems* (Birkhäuser, Boston, 1980).
- [24] R. C. Hilborn, *Chaos and Nonlinear Dynamics*, 2nd ed. (Oxford University Press, New York, 2003).
- [25] A. N. Berker and M. Wortis, Blume-Emery-Griffiths-Potts model in two dimensions: Phase diagram and critical properties from a position-space renormalization-group, *Phys. Rev. B* **14**, 4946 (1976).
- [26] P. E. Cladis, New liquid-crystal phase diagram, *Phys. Rev. Lett.* **35**, 48 (1975).
- [27] N. G. Fytas, J. Zierenberg, P. E. Theodorakis, M. Weigel, W. Janke, and A. Malakis, Universality from disorder in the random-bond Blume-Capel model, *Phys. Rev. E* **97**, 040102 (2018).
- [28] G. Naldi, L. Pareschi, and G. Toscani, eds., *Mathematical Modeling of Collective Behavior in Socio-Economic and*

