

Modelling of daily reference evapotranspiration using deep neural network in different climates

Atilla Ozgur^a, Sevim Seda Yamac^b

^a*Department of Logistics and Mathematics, Jacobs University Bremen gGmbH ,Campus Ring 1, 28759 Bremen, Germany*

^b*Konya Food and Agriculture University, Faculty of Agriculture and Natural Sciences, Department of Plant Production and Technologies, Konya, Turkey*

Abstract

Precise and reliable estimation of reference evapotranspiration (ET_o) is an essential for the irrigation and water resources management. ET_o is difficult to predict due to its complex processes. This complexity can be solved using machine learning methods. This study investigates the performance of artificial neural network (ANN) and deep neural network (DNN) models for estimating daily ET_o . Previously proposed ANN and DNN methods have been realized, and their performances have been compared. Six input data including maximum air temperature (T_{max}), minimum air temperature (T_{min}), solar radiation (R_n), maximum relative humidity (RH_{max}), minimum relative humidity (RH_{min}) and wind speed (U_2) are used from 4 meteorological stations (Adana, Aksaray, Isparta and Nide) during 1999-2018 in Turkey. The results have shown that our proposed DNN models achieves satisfactory accuracy for daily ET_o estimation compared to previous ANN and DNN models. The best performance has been observed with the proposed model of DNN with SeLU activation function (P-DNN-SeLU) in Aksaray with coefficient of determination (R^2) of 0.9934, root mean square error (RMSE) of 0.2073 and mean absolute error (MAE) of 0.1590, respectively. Therefore, the P-DNN-SeLU model could be recommended for estimation of ET_o in other climate zones of the world.

Keywords: Penman Monteith equation, Artificial Neural Networks, Deep Learning, Machine Learning, Meteorological data, Deep Neural Networks

1. Introduction

Reference evapotranspiration (ET_o) is an essential hydrological component for the sustainable and efficient management of agricultural water resources and the optimum irrigation scheduling (Huang et al., 2019; Wu et al., 2019; Yama and Todorovic, 2020). Many direct and indirect methods have been recommended to estimate ET_o . The direct method of ET_o estimation could be accomplished by water budget measurement (e.g. weighting lysimeters) or water vapor transfer methods (e.g. eddy covariance and Bowen ratio) (Huang et al., 2019). Unfortunately, these methods are time consuming and costly. Additionally, they have some spatial and temporal limitations (Irmak et al., 2003; Dinpashoh, 2006; Peng et al., 2017). As an alternative to the direct methods, mathematical models based on meteorological data provided by weather stations can be used to estimate ET_o (Tao et al., 2018).

The FAO-56 Penman-Monteith (FAO-56 PM) equation is recommended by the Food and Agriculture Organization (FAO) of the United Nations as a reference model for ET_o estimate (Allen et al., 1998). The FAO-56 PM incorporates both the aerodynamics and thermodynamics aspects and has more accurate results compared to the other empirical methods (Fan et al., 2018; Wu et al., 2019). The FAO-56 PM model has been evaluated against various other methods under diverse areas, climates and time steps (daily, weekly and monthly). The results show that FAO-56 PM method has better performance than other empirical equations (Pereira et al., 2015; Lpez-Urrea et al., 2006). However, the FAO-56 PM requires numerous features for ET_o estimation, including the geological variables such as elevation and latitude, and meteorological variables such as maximum and minimum temperature, maximum and minimum relative humidity, wind speed and net solar radiation (Shiri et al., 2014; Feng et al., 2017; Peng et al., 2017). These requirements bring a major drawback to the application of the FAO-56 PM model. Due to the limited availability of meteorological data, mainly in developing countries, simplified empirical models with fewer requirements have been proposed (Valiantzas, 2013; Ahooghalandari et al., 2016), such as temperature based (Hargreaves and Samani, 1985), mass transfer based (Trabert, 2019) and radiation based models (Priestley and Taylor, 1972). These simplified methods are more accurate for monthly and weekly ET_o estimation while they are less accurate on a daily ET_o estimation (Torres et al., 2011).

The estimation of ET_o is considered as a complex and highly nonlinear dynamic process depending on quality of meteorological variables (Wu et al.,

2019). However, it is usually difficult to develop accurate empirical models considering all these nonlinear and complicated processes, especially when some important input parameters are lacking. Recently, machine learning methods have been widely used to estimate complex process of ET_o estimation because these methods do not require knowledge of internal variables to solve non-linear and multi-variable functions (Kisi, 2015; Yama et al., 2020). Thus, various machine learning methods have been suggested for estimation of ET_o . Among these, (1) artificial neural networks (ANN) (Antonopoulos and Antonopoulos, 2017; Ferreira et al., 2019), (2) support vector machines (SVM) (Fan et al., 2018; Ferreira et al., 2019), (3) tree based assemble methods (Kisi and Kilic, 2016; Fan et al., 2018), (4) boosting (Fan et al., 2019) can be mentioned.

Because of promising results and enormous potential for image processing and data analysis, the Deep Neural Network (DNN) methods have become increasingly popular in recent years (Kamilaris and Prenafeta-Bold, 2018). The DNN methods are actually improved versions of the ANN methods (LeCun et al., 2015). The ANN with single hidden layer is commonly called as multi-layer perceptrons (MLP) or feed forward neural networks, while the ANN with more than two hidden layers are called Deep Neural Networks. DNNs are interchangeably called as deep neural networks, deep learning methods, or deep neural nets. The DNN methods have been applied to different domains, such as speech recognition (Amodei et al., 2016), natural language processing (Young et al., 2018), and game playing (Guo et al., 2014). Likewise, the use of DNN methods recently increased in the area of hydrological (Wang et al., 2020; Lee et al., 2020; Bui et al., 2020) and agricultural (Golhani et al., 2018; Grinblat et al., 2016; Dyrmann et al., 2016) studies.

The current study makes following contributions: (1) Previously proposed methods are categorized in Table 1 and Table 2. (2) Previously proposed methods have been realized and their performances have been compared in current meteorological dataset (Table 5). (3) Different from previously proposed DNN method (Saggi and Jain, 2019), dropout layer has been used and its performance is measured. (4) Moreover, two activation function which are called rectifier linear units (ReLU) and scaled exponential linear units (SeLU) has been used in DNN method and compared with the other methods. (5) All methods (previous and new) are compared using 5-fold cross validation instead of using single train-test dataset split. In 5-fold cross validation, models were trained on dataset with 5 different splits.

2. Related works

In this section, previous studies were reviewed for estimation of daily ET_o value. Table 1 summarizes the previous ANN methods in the literature. Table 2 demonstrates the dataset information of the previous ANN methods.

Various ANN methods were proposed for estimation of ET_o value (Table 1). However, only two DNN methods were proposed in the literature (Saggi and Jain, 2019; Ferreira and da Cunha, 2020). The first DNN method used only 3 hidden layers with Rectifier Linear Units (ReLU) (Saggi and Jain, 2019). The second DNN method employed convolutional neural networks on hourly data (Ferreira and da Cunha, 2020). Since the currently proposed DNN methods use daily data, they are not comparable to the second DNN method.

Table 1: Previously proposed neural network models for ET_o estimation

	Study	Neural Architecture	Network	Activation Functions	Software
1	Landeras et al. (2008)	(4-6)(1-14)-1		NI	Statistica
2	Traore et al. (2010)	(3-5)-(1-20)-1		Sigmoid	NeuroSolution
3	Huo et al. (2012)	5-8-1, 3-4-5-1, 3-4-5-1	4-5-6-1,	Sigmoidal logistic	Matlab
4	Rahimikhoob (2014)	4-(1-10)-1		Logsigmoid	Weka
5	Shiri et al. (2014)	(4-5)-(1-14)-1		Sigmoid, linear	Matlab
6	Goci et al. (2015)	5-3-6-10-1		Continuous logsigmoid	Matlab
7	Kisi and Kilic (2016)	4-(3-9)-1,2-(3-10)-1		Sigmoid, linear	NI
8	Yassin et al. (2016)	(4-9)-(2-20)-1		Sigmoid, linear	Multiple Back-Propagation
9	Feng et al. (2016)	(2,3,5)-6-1		NI	Matlab
10	Antonopoulos and Antonopoulos (2017)	(2-4)-6-1, 2-4-1		Sigmoid	NI
11	Dou and Yang (2018)	4-11-1, 4-15-1		Sigmoid, linear	Matlab
12	Saggi and Jain (2019)	7-40-60-40-1		ReLU, softmax	H_2O

As can be seen in Table 1, all of the architectures had standard input layer, hidden layer(s) and output layer. For example, architecture of Landeras et al. (2008) was "(4-6)(1-14)-1". This means that they used 4 to 6 neurons in the input layer, 1 to 14 neurons in the single hidden layer and 1 neuron in the output layer. They performed empirical experiments and reported their best results among the tried number of neurons. Also, other studies performed similar empirical experiments (Landeras et al., 2008; Traore et al., 2010; Rahimikhoob, 2014; Shiri et al., 2014; Kisi and Kilic, 2016; Yassin

et al., 2016). As an another example, Goci et al. (2015) used architecture of "5-3-6-10-1". This means that they used 5 neurons in the input layer, 3-6-10 neurons in the 3-hidden layers and 1 neuron in the output layer. Since model of Goci et al. (2015) used more than 2 layers in the architecture, they could have chosen to call their method DNN but they did not. This could be the fact that they did not use any other DNN improvements in their experiments.

Activation functions that are used in literature are given in the third column (Table 1). All the previous studies used standard sigmoid and linear functions except for the previously applied DNN method (Saggi and Jain, 2019). In addition, last column shows the software that was used in studies. Most of the studies (5/12) used Matlab neural network toolbox. Unfortunately, some studies did not report the activation functions and software, making reproducibility of their studies harder if not impossible.

Table 2 gives information about the years, frequency and dataset split (train, validation and test) in the literature. It was important to show the split of datasets (train, validation and test) because this split affects the machine learning performances. For example, Traore et al. (2010) reported that they used the meteorological dataset between 2004 and 2005 years as cross validation to optimize ANN performance. However, this usage was for validation dataset. Similar usage were done also by Landaras et al. (2008); Yassin et al. (2016).

Among the reviewed studies, 10 of 12 studies used the daily data for estimation of ET_o , while the other 2 studies used the monthly data. Interestingly, no study evaluated their approaches using true cross validation in their experiments. According to the best knowledge of the authors, the present study was the first study that uses true cross validation in the literature of ET_o estimation.

Table 2: Dataset information of the previously proposed neural network methods for ET_o estimation

	Study	Years	Frequency	Train Validation Test
1	Landeras et al. (2008)	1999-2003	Daily	1999-2001 Train (75%) validation (25%), 2002-2003 Test
2	Traore et al. (2010)	1996-2006	Daily	1996-2003 Train, 2004-2005 Validation, 2006 Test
3	Huo et al. (2012)	1952-2001	Daily	1952-1986 Train, 1987-2001 Test
4	Rahimikhoob (2014)	1998-2007	Monthly	1998-2004 Train, 2005-2007 Test
5	Shiri et al. (2014)	2000-2008	Daily	2000-2005 Train, 2006-2008 Test
6	Goci et al. (2015)	1980-2010	Monthly	1980-1995 Train, 1996-2010 Test
7	Kisi and Kilic (2016)	1994-2009	Daily	1998-2001 Train, 2002-2005 Validation, 2003-2009 Test
8	Yassin et al. (2016)	1980-2010	Daily	Train (65%), Validation (35%) in 13 stations, Test using separate 6 stations
9	Feng et al. (2016)	1994-2013	Daily	65% Train, 35% Test
10	Antonopoulos and Antonopoulos (2017)	2009-2013	Daily	1 year Train, other 4 years Test
11	Dou and Yang (2018)	2001-2009 but 6 years of data used	Daily	4 years Train, 1 year Validation, 1 year Test
12	Saggi and Jain (2019)	1978-1999 and 2007-2016 H, 1970-1999 and 2007-2016 P	Daily	55% Train, 30% Validation, 15% Test

3. Materials and methods

In this study, two newly DNN models were proposed to estimate daily value of ET_o . These newly proposed models were called P-DNN-ReLU and P-DNN-SeLU that uses activation ReLU and SeLU functions. Previous DNN

and ANN models were reproduced and their results were also included in experiments. The dropout layer was also tried on all DNN models, though its effect on performance was not good.

In addition to current (P-DNN-ReLU and P-DNN-SeLU) and previous DNN model (Saggi and Jain, 2019), 11 previous ANN methods were also implemented (Landeras et al., 2008; Traore et al., 2010; Huo et al., 2012; Rahimikhoob, 2014; Shiri et al., 2014; Goci et al., 2015; Kisi and Kilic, 2016; Yassin et al., 2016; Feng et al., 2016; Antonopoulos and Antonopoulos, 2017; Dou and Yang, 2018). The ANN methods used neuron size between 1 to 30 in their hidden layers. Therefore, 30 different ANN models were trained for every station. For the DNN methods, the dropout layer was also used. Therefore, $(6 \times 6 \times 6) \times 3 = 648$ different DNN models were trained also for every station. In total, 678 models were trained in the present study. Finally, considering 4 stations and 5 cross validation, the experiments trained and tested $678 \times 5 \times 4 = 13560$ different models. All of these experimental results are available as a supplementary data. The flowchart of the modeling procedure is presented in Figure 5.

3.1. Background knowledge

3.1.1. Study area and dataset description

The daily data from 4 meteorological stations in Turkey were obtained from Turkish State Meteorological Service for the period of 1999-2018. Data features are maximum air temperature (T_{max}), minimum air temperature (T_{min}), solar radiation (R_n), maximum relative humidity (RH_{max}), minimum relative humidity (RH_{min}) and wind speed (U_2). Table 3 shows the statistical parameters of meteorological variables at Adana, Aksaray, Isparta and Nide sites. The map of the study area and the location of the 4 meteorological stations are shown in Figure 1.

According to the Kppen Geiger climate classification (Kottek et al., 2006), the climate of Adana and Isparta sites have a warm temperature with a dry summer, while the climate of Aksaray and Nide sites have a semi-arid with cold and snowy winters. In this way, the meteorological data collected from 4 different sites in Turkey was applied to two different climate types in this study. Table 3 sums up the geographical and meteorological information of the 4 stations in Turkey.




Figure 1: Geographical locations of the 4 meteorological stations in Turkey.

3.1.2. FAO Penman-Monteith equation

The FAO Penman-Monteith (FAO-56 PM) equation was proposed by Allen et al. (1998). It is used to predict daily ET_o ($mm\ day^{-1}$) and provided the reference data for the training and testing in the current study.

$$ET_o = \frac{0.408(R_n - G) + \gamma \frac{900}{T+273} U_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34 U_2)} \quad (1)$$

where ET_o is the reference evapotranspiration ($mm\ day^{-1}$). R_n is the net solar radiation ($MJ\ m^{-2}\ day^{-1}$), G is the soil heat flux density ($MJ\ m^{-2}\ day^{-1}$), T is the mean daily air temperature at 2m height ($^{\circ}C$), Δ is the slope of the saturated vapour pressure curve ($kPa\ ^{\circ}C^{-1}$), γ is the psychrometric constant ($0.066\ kPa\ ^{\circ}C^{-1}$), e_s and e_a are saturated and prevailing actual vapour pressure (kPa), respectively, and U_2 is the mean daily wind speed ($m\ s^{-1}$) measured at 2 m height.

The saturation vapor pressure e_s was estimated as:

$$e_s = \frac{e^o(T_{max}) + e^o(T_{min})}{2} \quad (2)$$

Table 3: Geographic information and statistical parameters of meteorological variables of the 4 stations (Adana, Aksaray, Isparta and Nide) in Turkey during 1999-2018

Station	Station Code	Longitude	Latitude	Altitude	Variable	Min	Max	Mean	Std ¹	Cv ²
Adana	1	35.34	37.00	23						
					T_{max}	5.30	44.40	25.74	7.68	0.30
					T_{min}	-3.20	29.80	14.85	7.28	0.49
					R_s	0.00	33.68	15.90	6.98	0.44
					R_{Hmax}	27.00	100.00	86.22	11.38	0.13
					R_{Hmin}	0.00	96.00	42.57	17.09	0.40
					U_2	0.30	6.00	1.62	0.70	0.43
					ET_0	0.51	12.73	4.45	2.20	0.50
Aksaray	2	34.00	38.37	970						
					T_{max}	-10.00	40.00	19.30	10.06	0.52
					T_{min}	-20.40	25.60	7.09	8.05	1.14
					R_s	0.69	32.43	16.98	7.82	0.46
					R_{Hmax}	20.00	100.00	71.40	16.46	0.23
					R_{Hmin}	0.00	98.00	37.69	17.01	0.45
					U_2	0.30	5.92	1.58	0.69	0.44
					ET_0	0.34	10.61	4.24	2.55	0.60
Isparta	3	30.57	37.78	997						
					T_{max}	-6.60	42.30	19.24	9.49	0.49
					T_{min}	-16.00	23.30	6.53	7.19	1.10
					R_s	0.00	32.52	15.30	7.99	0.52
					R_{Hmax}	14.00	100.00	81.44	12.94	0.16
					R_{Hmin}	0.00	99.00	40.93	16.87	0.41
					U_2	0.00	5.78	1.32	0.70	0.53
					ET_0	0.42	9.60	3.73	2.30	0.62
Nide	4	34.68	37.96	1211						
					T_{max}	-10.30	38.50	18.51	9.90	0.53
					T_{min}	-19.80	23.00	6.02	7.89	1.31
					R_s	0.68	35.10	18.75	8.38	0.45
					R_{Hmax}	24.00	104.00	75.55	15.13	0.20
					R_{Hmin}	2.00	96.00	37.44	17.49	0.47
					U_2	0.38	7.95	1.83	0.70	0.38
					ET_0	0.39	10.99	4.49	2.67	0.59

¹ Std: Standard deviation

² Cv: Covariance of variance

where $e^o(T)$ is the saturation vapour pressure (kPa) at the air temperature (T), and T_{max} and T_{min} are maximum and minimum daily air temperature ($^{\circ}C$), respectively. The saturation vapor pressure at the air temperature T was calculated as:

$$e^o(T) = 0.6108 e^{\left[\frac{17.27 T}{T+237.3}\right]} \quad (3)$$

The actual vapor pressure (e_a) was calculated as:

$$e_a = \frac{e^o(T_{min}) \frac{RH_{max}}{100} + e^o(T_{max}) \frac{RH_{min}}{100}}{2} \quad (4)$$

where RH_{max} is the maximum daily relative humidity, (RH_{min}) is the minimum daily relative humidity, $e^o(T_{min})$ is the saturation vapor pressure (kPa) at the minimum daily air temperature and $e^o(T_{max})$ is the saturation vapor pressure (kPa) at the maximum daily air temperature, respectively.

3.1.3. Artificial neural networks and deep neural networks

Artificial neural networks (ANN) are powerful machine learning methods that take their roots from biological neurons. The ANN uses artificial neurons modeled from biological neurons as fundamental building block. An artificial neuron has 3 characteristics. (1) inputs, (2) summation unit and (3) transfer (activation) function (Figure 2).




Figure 2: Visualization of the artificial neuron.

In the estimation of ET_o , inputs are features used in the FAO-56 PM equation. Inputs are multiplied with weights and added together in summation

unit. The summation value is sent to activation function and the output of this activation function is the output of the neuron. Diverse transfer functions are proposed in the neural network literature (Nwankpa et al., 2018). The most common activation functions are sigmoid, Gaussian and linear. Different transfer function are used for different purposes. For example, sigmoid function is used for binary classification whereas linear function is used for regression.

The ANN method is the umbrella term for machine learning methods that use neurons as building blocks. Nonetheless, most studies use terms of Neural Networks, Artificial Neural Networks (ANN), Multi Layer Perceptrons (MLP) and Feed Forward Neural Networks interchangeably. In the current study, artificial neural network term is preferred since this term is commonly used in related literature.




Figure 3: Visualization of Neural Network (3-4-5-1) architecture (Note: Figure is created with Weka (Hall et al., 2009)).

In the ANN architecture, multiple neurons are used in multiple layers. Most of the time, ANN architecture is identified with layer neuron counts such as 3 – 5 – 1. Here, it has 3 layers (one input, one hidden and one output layer) and there are 3 neurons in input layer, 5 neurons in one hidden layer and 1 neuron in output layer. According to number of features used, input layer have corresponding number of neurons. An example for an ANN architecture is depicted in Figure 3.

3.1.4. Deep neural networks

Deep neural networks (DNN) are advanced versions of ANN methods (Nielsen, 2015; LeCun et al., 2015). However, the differences between the classical ANN and DNN methods are not clearly defined in the literature, but following improvements are mostly related to DNN studies.

1. Using more than 2 hidden layers, so called deep layers.
2. Different neuron types, such as convolutional, pooling and dropout.
3. Introduction of new activation functions, such as rectified linear units (ReLU), softmax and scaled exponential linear units (SeLU) functions and many others.
4. Different training methods for back propagation that are more suited to using parallelization using multiple GPUs and CPUs.
5. Different initialization methods for neuron weights.

3.1.5. Cross validation

The k -fold cross validation ($k = 5$) method was used in this study. The data were divided into k parts and algorithms were trained using $k - 1$ parts. After that the trained model was tested on remaining 1 part. This procedure was repeated k times. As an example, in the 5 fold cross validation, dataset was divided into 5 parts. Using 4 parts, ANN and DNN models were trained, then tested on remaining 1 part. This procedure was repeated 5 times (Figure 4). Since experiments were repeated 5 times and performance metrics averaged, obtained performance metrics are more reliable than using only one train-test split.

Train	Train	Train	Train	Test
Train	Train	Train	Test	Train
Train	Train	Test	Train	Train
Train	Test	Train	Train	Train
Test	Train	Train	Train	Train

Figure 4: The visualization of 5-fold cross validation.

3.1.6. Performance evaluation of model parameters

The performance of the models were evaluated using the root mean square error (RMSE), mean absolute error (MAE), coefficient of determination (R^2) in the training and testing subsets.

The daily ET_o values, generated by the models (S_i), were transformed into daily errors, comparing them with observed values (O_i). \bar{O} is the mean value of the observed values and \bar{S} is the mean value of the observed values.

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (S_i - O_i)^2}{n}} \quad (5)$$

$$MAE = \left| \frac{\sum_{i=1}^n (S_i - O_i)}{n} \right| \quad (6)$$

$$R^2 = \frac{[\sum_{i=1}^n (O_i - \bar{O})(S_i - \bar{S})]^2}{\sum_{i=1}^n (O_i - \bar{O}) \sum_{i=1}^n (S_i - \bar{S})} \quad (7)$$

Smaller values of RMSE and MAE and higher values of R^2 indicates higher model performance.




Figure 5: The flowchart of the experimental procedures




Figure 6: Activation Functions used in Experiments.

3.2. Activation functions

As mentioned in section 3.1.4, differences between the DNN and ANN methods are the number of layers, activation functions, and neuron types. Figure 6 shows used activation functions in present study. Classical ANN methods mostly uses sigmoid function (Equation 8).

$$\text{sigmoid}(x) = \frac{1}{1 + e^{-x}} \quad (8)$$

ReLU function (Equation 9) was made popular by DNN methods (Glorot et al., 2011). Only the previously applied DNN method (Saggi and Jain, 2019) was used ReLU function in the hidden layers.

$$\text{ReLU}(x) = \max(0, x) \quad (9)$$

In addition to ReLU function, SeLU function (Equation 10) was also tried in the experiments. The SeLU function was introduced by Klambauer et al. (2017). Klambauer et al proposed the best constant values for (α, λ) as (1.67326324, 1.05070098), respectively.

$$\text{SeLU}(x) = \lambda \begin{cases} x & \text{if } x > 0 \\ \alpha e^x - \alpha & \text{if } x \leq 0 \end{cases} \quad (10)$$

3.3. Dropout layer

The dropout layer was first proposed by Srivastava et al. (2014). Dropout layer is one of the most popular solutions to over fitting problem in DNN methods. Dropout layer is used with single parameter, dropout rate, which changes between 0 and 1. Dropout rate controls chance of dropping out connections when neural network is training. For example, if dropout rate is 0.5, coming neural connections to dropout layer is dropped with 50% chance. An example can be seen in Figure 7 in which connections of crossed neurons are dropped out.




Figure 7: The architecture of dropout in neural networks. (Srivastava et al., 2014)

To be able to see if dropout layer is helpful for over fitting in the estimation of ET_o value, different dropout rates were tried between the hidden layers. Therefore, 3 different dropout layers were introduced in DNN models (Figure 8b and Figure 8d). In the experiment, dropout rate changed from 0 to 0.5 with a rate of 0.1 increase. In total, 6 dropout rates were tried [0, 0.1, 0.2, 0.3, 0.4, 0.5]. In the end, using 3 DNN models with 3 different layers and 6 dropout layers (3 DNN * 6 (Dropout Layer 2) * 6 (Dropout Layer 2) * 6 (Dropout Layer 3)), 648 models were applied in the experiments. According to our results, dropout layer is not useful for the estimation of ET_o values.

3.4. Proposed deep neural network models

In this study, 3 layer architecture (1-60-90-60-1) were proposed for DNN model. For this 3 layer architecture, both ReLU and SeLU activation functions

were used. Additionally, different dropout rates were also tried in the DNN model. Figure 8 shows the proposed DNN models.




Figure 8: Visualization of proposed DNN methods.

4. Results and discussion

4.1. Comparison of all the applied models

The best twenty performances metric of current and previous models for estimating daily ET_o at the 4 meteorological stations are presented in Table 4. As can be seen in the Table 4, R^2 , RMSE and MAE performance metrics ranged between 0.9913-0.9933, 0.1811-0.2471 $mm\ day^{-1}$, 0.1333-0.1874 $mm\ day^{-1}$, respectively. It was found that R^2 values of all the applied models were higher than 0.991, indicating a strong relationship between the ET_o values from the FAO-56 PM equation and those predicted by the applied models. The RMSE and MAE values were lower than 0.25 and 0.19 $mm\ day^{-1}$, which shows excellent performance for the estimation of daily ET_o . The proposed models (P-DNN-SeLU and P-DNN-ReLU) performed better than the previous models, with R^2 , RMSE and MAE ranging 0.9933-0.9932, 0.1811-0.2182 $mm\ day^{-1}$, 0.1333-0.1678 $mm\ day^{-1}$, respectively. These results confirmed that proposed DNN models are superior to previously applied ANN and DNN models for estimation of daily ET_o .

Table 4: The best twenty performance metric of all the currently and previously proposed models for estimation of ET_o at the 4 meteorological stations according to R^2

Order	model name	station name	R^2	RMSE	MAE
01	P-DNN-SeLU ¹	Aksaray	0.9934	0.2073	0.1591
02	P-DNN-ReLU	Nigde	0.9933	0.2182	0.1678
03	P-DNN-SeLU	Adana	0.9932	0.1812	0.1333
04	L-DNN-Saggi	Nigde	0.9930	0.2240	0.1740
05	L-DNN-Saggi	Aksaray	0.9928	0.2160	0.1654
06	P-DNN-ReLU	Aksaray	0.9928	0.2165	0.1651
07	L-DNN-Saggi	Aksaray	0.9926	0.2197	0.1683
08	L-ANN (60, 90, 60) ²	Nigde	0.9919	0.2400	0.1826
09	P-DNN-SeLU dropout 0 0.1 0	Aksaray	0.9919	0.2298	0.1788
10	L-ANN (9)	Aksaray	0.9918	0.2311	0.1786
11	L-ANN (19)	Nigde	0.9918	0.2421	0.1854
12	L-ANN (19)	Aksaray	0.9916	0.2334	0.1788
13	P-DNN-SeLU dropout 0 0.1 0	Adana	0.9916	0.2023	0.1483
14	L-ANN (12)	Nigde	0.9916	0.2453	0.1871
15	L-ANN (21)	Aksaray	0.9915	0.2348	0.1796
16	L-ANN (22)	Aksaray	0.9915	0.2353	0.1799
17	L-ANN (16)	Aksaray	0.9915	0.2354	0.1793
18	L-ANN (28)	Aksaray	0.9915	0.2355	0.1797
19	L-ANN (26)	Nigde	0.9914	0.2472	0.1875
20	L-ANN (60, 90, 60)	Aksaray	0.9913	0.2372	0.1804

¹ P- means currently proposed, L mean Literature.

² ANN(..) refers to Artificial Neural Network hidden layer neuron counts, for example (21) means 1 hidden layer with 21 neurons while (60,90,60) means 3 hidden layers with 60, 90 and 60 neurons.

However, the performance of the proposed activation SeLU decreased when dropout layer used (proposed activation SeLU dropout 0 0.1 0). This means that using dropout layers did not improve the modeling performances for the estimation of daily ET_o .

In general, it is observed that among the previously applied models, the Saggi and Jain model had the highest performance based on the performance metrics. Among the stations in the top twenty best performing models, Aksaray and Nigde stations located in the semi-arid region were the most appearing, while Adana and Isparta stations located in Mediterranean region

were the least appearing. This can be explained by the fact that the models performed better in semi arid region than the Mediterranean region.

4.2. Comparison of proposed and previously applied ANN and DNN models

Table 5: The highest performances of previously proposed methods for ET_o estimation according to R^2 score

	Study	ANN Hidden Layers	Best (R^2) Adana	Best (R^2) Aksaray	Best (R^2) Isparta	Best (R^2) Nigde
1	Landeras et al. (2008), Traore et al. (2010), Rahimikhoob (2014), Shiri et al. (2014), Yassin et al. (2016), Kisi and Kilic (2016), Antonopoulos and Antonopoulos (2017), Dou and Yang (2018)	(1-30)	0.9864	0.9918	0.9880	0.9918
2	Huo et al. (2012)	4-5, 5-6	0.9814	0.9883	0.9837	0.9885
3	Saggi and Jain (2019)	40-60-40	0.9905	0.9928	0.9896	0.9931
3	Proposed ReLU	60-90-60	0.9888	0.9928	0.9890	0.9934
4	Proposed SeLU	60-90-60	0.9925	0.9940	0.9887	0.9936

The comparison results of the previous models (ANN and DNN) and two current DNN models can be seen in Table 5. The table showed only the best performances among the experiments. According to Table 5, it was seen that R^2 values of all the previously applied and proposed models were higher than 0.98. DNN models compared to ANN models improved R^2 values in the 3-4 decimal points. If the complexity of the models and time to train are taken into account, these results may be considered as diminishing improvements according to requirements for ET_o .

4.3. Comparison of meteorological station performances

The best five performance metric of currently proposed and previous models for estimation of ET_o at the Adana, Aksaray, Isparta and Nide

Table 6: The best five performance metric of proposed and previously applied models for estimation of ET_o at the Adana, Aksaray, Isparta and Nide stations respectively according to R^2 score

Rank	model name	station name	R^2	RMSE	MAE
01	P-DNN-SeLU	Adana	0.9925	0.1912	0.1409
02	P-DNN-SeLU dropout 0.0 0.1 0.0	Adana	0.9916	0.2023	0.1483
03	L-DNN-Saggi	Adana	0.9905	0.2151	0.1520
04	P-DNN-ReLU	Adana	0.9888	0.2336	0.1664
05	ANN (60, 90, 60)	Adana	0.9865	0.2557	0.1860
01	P-DNN-SeLU	Aksaray	0.9940	0.1977	0.1503
02	L-DNN-Saggi	Aksaray	0.9928	0.2160	0.1654
03	P-DNN-ReLU	Aksaray	0.9928	0.2165	0.1652
04	P-DNN-SeLU dropout 0.0 0.1 0.0	Aksaray	0.9919	0.2298	0.1788
05	ANN (9)	Aksaray	0.9918	0.2311	0.1786
01	L-DNN-Saggi	Isparta	0.9896	0.2346	0.1779
02	P-DNN-ReLU	Isparta	0.9890	0.2405	0.1848
03	P-DNN-SeLU	Isparta	0.9887	0.2438	0.1864
04	ANN (16)	Isparta	0.9880	0.2513	0.1942
05	P-DNN-SeLU dropout 0.0 0.0 0.1	Isparta	0.9878	0.2538	0.1938
01	P-DNN-SeLU	Nigde	0.9936	0.2132	0.1644
02	P-DNN-ReLU	Nigde	0.9934	0.2168	0.1681
03	L-DNN-Saggi	Nigde	0.9931	0.2219	0.1708
04	ANN (60, 90, 60)	Nigde	0.9919	0.2400	0.1826
05	ANN (19)	Nigde	0.9918	0.2421	0.1854

stations are presented in Table 6. In general, Table 6 showed that the P-DNN-SeLU model had the highest performance in Aksaray, Nide and Adana stations. In that case, R^2 values of P-DNN-SeLU model are 0.9939, 0.9936 and 0.9924, RMSE values are 0.1977, 0.2131 and 0.1911, MAE values are 0.1502, 0.1643 and 0.1409. However, Saggi and Jain model had the highest performance in Isparta station with the value of R^2 to 0.9896, RMSE to 0.2346 and MAE to 0.1779, respectively. Obviously, the proposed activation SeLU model in Aksaray and Nide stations had slightly better prediction accuracy than the P-DNN-SeLU model in Adana and Isparta stations. It can be seen that the P-DNN-SeLU model showed more improvements in daily ET_o estimation in the semi arid region, compared with those in the Mediterranean regions.

The scatter plot of estimated ET_o values by P-DNN-SeLU, P-DNN-ReLU and Saggi and Jain models compared with the FAO-56 PM values at the Adana, Aksaray, Isparta and Nigde stations are presented in Figure 9. The figure showed that the plotted data points mostly correlated close towards the 1:1 line. However, the models in Isparta station yielded more scattered daily ET_o values compared to other three stations. The models were more close to those obtained with FAO-56 PM equation in Aksaray station. These results indicated that the models showed a much higher prediction accuracy of daily ET_o value in Aksaray station.




Figure 9: Scatter plots for 3 Deep Neural Networks.

5. Conclusions

This study assessed the application of 14 ANN methods (2 new DNN, 1 previous DNN and 11 previous ANN) for estimation of daily ET_o in the two different climate zones of Turkey. The models used 6 input meteorological data including T_{max} , T_{min} , R_n , RH_{max} , RH_{min} and U_2 from 4 weather stations (Adana, Aksaray, Isparta and Nide) during 1999-2018 in Turkey. The results demonstrated that all models of DNN and ANN achieved satisfactory accuracy for estimation of daily ET_o using available meteorological data. Especially, the DNN methods were highly effective in estimating ET_o value. It can be seen from the results that the performance of the models became more reliable when cross validation was used in the study. However, the result showed that dropout layer was not useful for ET_o estimation. In addition, using more powerful architecture did not improve the estimation of ET_o .

In general, among the meteorological stations, Aksaray station offered the best prediction accuracy, while Isparta station performed the least prediction accuracy in all the DNN and ANN models. The overall results indicated that the proposed model of P-DNN-SeLU made a significant improvement in accuracy among the other models. Therefore, P-DNN-SeLU model has a very high potential for estimation of daily ET_o in different climatic zones of Turkey, even possibly in other zones of the world. In short, P-DNN-SeLU model could be applied in the future studies to estimate ET_o under different climate conditions. Finally, further studies should be carried out to evaluate the applicability of P-DNN-SeLU model under limited input data, in places where meteorological variables are limited.

References

- Ahooghalandari, M., Khiadani, M., Jahromi, M. E., 2016. Developing equations for estimating reference evapotranspiration in australia. Water Resources Management.
- Allen, R., Pereira, L., Raes, D., Smith, M., 1998. Crop evapotranspiration guidelines for computing crop water requirements-fao irrigation and drainage paper 56. Tech. rep., Food and Agriculture Organization (FAO).
- Amodei, D., et al., 2016. Deep speech 2 : End-to-end speech recognition in english and mandarin. In: Proceedings of The 33rd International Conference on Machine Learning.

- Antonopoulos, V. Z., Antonopoulos, A. V., 2017. Daily reference evapotranspiration estimates by artificial neural networks technique and empirical equations using limited input climate variables. *Computers and Electronics in Agriculture* 132, 86 – 96.
URL <http://www.sciencedirect.com/science/article/pii/S0168169916305506>
- Bui, Q.-T., Nguyen, Q.-H., Nguyen, X. L., Pham, V. D., Nguyen, H. D., Pham, V.-M., 2020. Verification of novel integrations of swarm intelligence algorithms into deep learning neural network for flood susceptibility mapping. *Journal of Hydrology* 581, 124379.
URL <http://www.sciencedirect.com/science/article/pii/S002216941931114X>
- Dinpashoh, Y., 2006. Study of reference crop evapotranspiration in i.r. of iran. *Agricultural Water Management*.
- Dou, X., Yang, Y., 2018. Evapotranspiration estimation using four different machine learning approaches in different terrestrial ecosystems. *Computers and Electronics in Agriculture* 148, 95 – 106.
URL <http://www.sciencedirect.com/science/article/pii/S016816991731476X>
- Dyrmann, M., Karstoft, H., Midtiby, H. S., 2016. Plant species classification using deep convolutional neural network. *Biosystems Engineering* 151, 72 – 80.
URL <http://www.sciencedirect.com/science/article/pii/S1537511016301465>
- Fan, J., Ma, X., Wu, L., Zhang, F., Yu, X., Zeng, W., 2019. Light gradient boosting machine: An efficient soft computing model for estimating daily reference evapotranspiration with local and external meteorological data. *Agricultural Water Management* 225, 105758.
- Fan, J., Yue, W., Wu, L., Zhang, F., Cai, H., Wang, X., Lu, X., Xiang, Y., 2018. Evaluation of svm, elm and four tree-based ensemble models for predicting daily reference evapotranspiration using limited meteorological data in different climates of china. *Agricultural and Forest Meteorology* 263, 225 – 241.

- URL <http://www.sciencedirect.com/science/article/pii/S0168192318302855>
- Feng, Y., Cui, N., Gong, D., Zhang, Q., Zhao, L., 2017. Evaluation of random forests and generalized regression neural networks for daily reference evapotranspiration modelling. *Agricultural Water Management* 193, 163 – 173.
URL <http://www.sciencedirect.com/science/article/pii/S0378377417302597>
- Feng, Y., Cui, N., Zhao, L., Hu, X., Gong, D., 2016. Comparison of elm, gann, wnn and empirical models for estimating reference evapotranspiration in humid region of southwest china. *Journal of Hydrology* 536, 376 – 383.
URL <http://www.sciencedirect.com/science/article/pii/S0022169416300981>
- Ferreira, L. B., da Cunha, F. F., may 2020. New approach to estimate daily reference evapotranspiration based on hourly temperature and relative humidity using machine learning and deep learning. *Agricultural Water Management* 234, 106113.
- Ferreira, L. B., da Cunha, F. F., de Oliveira, R. A., Filho, E. I. F., 2019. Estimation of reference evapotranspiration in brazil with limited meteorological data using ann and svm a new approach. *Journal of Hydrology* 572, 556 – 570.
URL <http://www.sciencedirect.com/science/article/pii/S0022169419302689>
- Glorot, X., Bordes, A., Bengio, Y., 2011. Deep sparse rectifier neural networks. In: *AISTATS*.
- Goci, M., Motamedi, S., Shamshirband, S., Petkovi, D., Ch, S., Hashim, R., Arif, M., 2015. Soft computing approaches for forecasting reference evapotranspiration. *Computers and Electronics in Agriculture* 113, 164 – 173.
URL <http://www.sciencedirect.com/science/article/pii/S0168169915000526>
- Golhani, K., Balasundram, S. K., Vadamalai, G., Pradhan, B., 2018. A review of neural networks in plant disease detection using hyperspectral

- data. Information Processing in Agriculture 5 (3), 354 – 371.
 URL <http://www.sciencedirect.com/science/article/pii/S2214317317301774>
- Grinblat, G. L., Uzal, L. C., Larese, M. G., Granitto, P. M., 2016. Deep learning for plant identification using vein morphological patterns. Computers and Electronics in Agriculture 127, 418 – 424.
 URL <http://www.sciencedirect.com/science/article/pii/S0168169916304665>
- Guo, X., Singh, S., Lee, H., Lewis, R. L., Wang, X., 2014. Deep learning for real-time atari game play using offline monte-carlo tree search planning. In: Advances in Neural Information Processing Systems 27. Curran Associates, Inc.
 URL <http://papers.nips.cc/paper/5421-deep-learning-for-real-time-atari-game-play.pdf>
- Hall, M., Frank, E., Holmes, G., Pfahringer, B., Reutemann, P., Witten, I. H., 2009. The weka data mining software: An update. SIGKDD Explor. Newsl.
- Hargreaves, G., Samani, Z., 1985. Reference crop evapotranspiration from temperature. Applied Engineering in Agriculture.
- Huang, G., Wu, L., Ma, X., Zhang, W., Fan, J., Yu, X., Zeng, W., Zhou, H., 2019. Evaluation of catboost method for prediction of reference evapotranspiration in humid regions. Journal of Hydrology 574, 1029 – 1041.
 URL <http://www.sciencedirect.com/science/article/pii/S0022169419304251>
- Huo, Z., Feng, S., Kang, S., Dai, X., 2012. Artificial neural network models for reference evapotranspiration in an arid area of northwest china. Journal of Arid Environments 82, 81 – 90.
- Irmak, S., Irmak, A., Allen, R. G., Jones, J. W., 2003. Solar and net radiation-based equations to estimate reference evapotranspiration in humid climates. Journal of Irrigation and Drainage Engineering.
- Kamilaris, A., Prenafeta-Bold, F. X., 2018. Deep learning in agriculture: A survey. Computers and Electronics in Agriculture 147, 70 – 90.

- URL <http://www.sciencedirect.com/science/article/pii/S0168169917308803>
- Kisi, O., 2015. Pan evaporation modeling using least square support vector machine, multivariate adaptive regression splines and m5 model tree. *Journal of Hydrology*.
- Kisi, O., Kilic, Y., 2016. An investigation on generalization ability of artificial neural networks and m5 model tree in modeling reference evapotranspiration. *Theoretical and Applied Climatology* 126 (3), 413–425.
- Klambauer, G., Unterthiner, T., Mayr, A., Hochreiter, S., 2017. Self-normalizing neural networks. *CoRR*.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World map of the kppen-geiger climate classification updated. *Meteorologische Zeitschrift* 15, 259–263.
- Landeras, G., Ortiz-Barredo, A., Lpez, J. J., 2008. Comparison of artificial neural network models and empirical and semi-empirical equations for daily reference evapotranspiration estimation in the basque country (northern spain). *Agricultural Water Management*.
- LeCun, Y., Bengio, Y., Hinton, G., May 2015. Deep learning. *Nature* 521, 436.
URL <https://doi.org/10.1038/nature14539>
- Lee, T., Shin, J.-Y., Kim, J.-S., Singh, V. P., 2020. Stochastic simulation on reproducing long-term memory of hydroclimatological variables using deep learning model. *Journal of Hydrology* 582, 124540.
URL <http://www.sciencedirect.com/science/article/pii/S0022169419312752>
- Lpez-Urrea, R., de Santa Olalla, F. M., Fabeiro, C., Moratalla, A., 2006. Testing evapotranspiration equations using lysimeter observations in a semiarid climate. *Agricultural Water Management*.
- Nielsen, M., 2015. *Neural Networks and Deep Learning*. Determination press.
URL <http://neuralnetworksanddeeplearning.com/index.html>

- Nwankpa, C., Ijomah, W., Gachagan, A., Marshall, S., 2018. Activation functions: Comparison of trends in practice and research for deep learning. CoRR.
- Peng, L., Li, Y., Feng, H., 12 2017. The best alternative for estimating reference crop evapotranspiration in different sub-regions of mainland china. Scientific Reports 7.
- Pereira, L. S., Allen, R. G., Smith, M., Raes, D., 2015. Crop evapotranspiration estimation with fao56: Past and future. Agricultural Water Management.
- Priestley, C. H. B., Taylor, R. J., 1972. On the assessment of surface heat flux and evaporation using large-scale parameters. Monthly Weather Review 100 (2), 81–92.
- Rahimikhoob, A., 02 2014. Comparison between m5 model tree and neural networks for estimating reference evapotranspiration in an arid environment. Water Resources Management 28, 657–669.
- Saggi, M. K., Jain, S., 2019. Reference evapotranspiration estimation and modeling of the punjab northern india using deep learning. Computers and Electronics in Agriculture 156, 387 – 398.
URL <http://www.sciencedirect.com/science/article/pii/S0168169918308779>
- Shiri, J., Nazemi, A. H., Sadraddini, A. A., Landeras, G., Kisi, O., Fard, A. F., Marti, P., 2014. Comparison of heuristic and empirical approaches for estimating reference evapotranspiration from limited inputs in iran. Computers and Electronics in Agriculture.
- Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I., Salakhutdinov, R., 2014. Dropout: A simple way to prevent neural networks from overfitting. Journal of Machine Learning Research 15, 1929–1958.
URL <http://jmlr.org/papers/v15/srivastava14a.html>
- Tao, H., Diop, L., Bodian, A., Djaman, K., Ndiaye, P. M., Yaseen, Z. M., 2018. Reference evapotranspiration prediction using hybridized fuzzy model with firefly algorithm: Regional case study in burkina faso. Agricultural Water Management 208, 140 – 151.

- URL <http://www.sciencedirect.com/science/article/pii/S0378377418308151>
- Torres, A. F., Walker, W. R., McKee, M., 2011. Forecasting daily potential evapotranspiration using machine learning and limited climatic data. *Agricultural Water Management*.
- Trabert, W., 09 2019. Neue beobachtungen ber verdampfungsgeschwindigkeiten. *Meteorol. Z.* 13, 261–263.
- Traore, S., Wang, Y.-M., Kerh, T., 2010. Artificial neural network for modeling reference evapotranspiration complex process in sudano-sahelian zone. *Agricultural Water Management* 97 (5), 707 – 714.
- Valiantzas, J. D., 2013. Simplified forms for the standardized fao-56 penmanmonteith reference evapotranspiration using limited weather data. *Journal of Hydrology*.
- Wang, N., Zhang, D., Chang, H., Li, H., 2020. Deep learning of subsurface flow via theory-guided neural network. *Journal of Hydrology* 584, 124700.
URL <http://www.sciencedirect.com/science/article/pii/S0022169420301608>
- Wu, L., Zhou, H., Ma, X., Fan, J., Zhang, F., 2019. Daily reference evapotranspiration prediction based on hybridized extreme learning machine model with bio-inspired optimization algorithms: Application in contrasting climates of china. *Journal of Hydrology*, 123960.
URL <http://www.sciencedirect.com/science/article/pii/S0022169419306808>
- Yama, S. S., eker, C., Negi, H., 2020. Evaluation of machine learning methods to predict soil moisture constants with different combinations of soil input data for calcareous soils in a semi arid area. *Agricultural Water Management* 234, 106121.
URL <http://www.sciencedirect.com/science/article/pii/S037837741932356X>
- Yama, S. S., Todorovic, M., 2020. Estimation of daily potato crop evapotranspiration using three different machine learning algorithms and four scenarios of available meteorological data. *Agricultural Water Management*.

- Yassin, M. A., Alazba, A., Mattar, M. A., 2016. Artificial neural networks versus gene expression programming for estimating reference evapotranspiration in arid climate. *Agricultural Water Management*.
- Young, T., Hazarika, D., Poria, S., Cambria, E., 2018. Recent trends in deep learning based natural language processing [review article]. *IEEE Computational Intelligence Magazine*.