

# On critical elliptic problems with singular Trudinger-Moser nonlinearities\*

Shiqiu Fu and Kanishka Perera  
 Department of Mathematical Sciences  
 Florida Institute of Technology  
 Melbourne, FL 32901, USA  
*sfu2013@my.fit.edu & kperera@fit.edu*

## Abstract

We establish some existence results for a class of critical elliptic problems with singular exponential nonlinearities. We do not assume any global sign conditions on the nonlinearity, which makes our results new even in the nonsingular case.

## 1 Introduction

The purpose of this paper is to establish some existence results for the class of singular elliptic problems with exponential nonlinearities

$$\begin{cases} -\Delta u = h(u) \frac{e^{\alpha u^2}}{|x|^\gamma} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where  $\Omega$  is a bounded Lipschitz domain in  $\mathbb{R}^2$  containing the origin,  $\alpha > 0$ ,  $0 \leq \gamma < 2$ , and  $h$  is a continuous function such that

$$\lim_{|t| \rightarrow \infty} h(t) = 0 \quad (1.2)$$

and

$$0 < \beta := \liminf_{|t| \rightarrow \infty} th(t) < \infty. \quad (1.3)$$

---

\**MSC2010*: Primary 35B33, Secondary 35J20, 35J60, 35J75

*Key Words and Phrases*: Critical elliptic problems, singular exponential nonlinearities, existence

This problem is motivated by the following singular Trudinger-Moser embedding:

$$\int_{\Omega} \frac{e^{\alpha u^2}}{|x|^{\gamma}} dx < \infty \quad \forall u \in H_0^1(\Omega)$$

for all  $\alpha > 0$  and  $0 \leq \gamma < 2$ , and

$$\sup_{\|u\|_{H_0^1(\Omega)} \leq 1} \int_{\Omega} \frac{e^{\alpha u^2}}{|x|^{\gamma}} dx < \infty \quad (1.4)$$

if and only if

$$\frac{\alpha}{4\pi} + \frac{\gamma}{2} \leq 1$$

(see Adimurthi and Sandeep [2]). Problem (1.1) is critical with respect to this embedding and hence lacks compactness. The case  $\beta = \infty$  was considered in [2], so we will focus on the case  $0 < \beta < \infty$  here.

The nonsingular case  $\gamma = 0$  has been widely studied in the literature (see, e.g., Adimurthi [1], Adimurthi and Yadava [3], de Figueiredo et al. [4, 5], Marcos B. do Ó [8], de Figueiredo et al. [6, 7], Zhang et al. [10], Perera and Yang [9], and their references). However, in all these results it is assumed that  $h(t) > 0$  for all  $t > 0$  and  $h(t) < 0$  for all  $t < 0$ . We impose no such global sign conditions on the nonlinearity  $h$ , so our results here are new even in the nonsingular case.

The singular eigenvalue problem

$$\begin{cases} -\Delta u = \lambda \frac{u}{|x|^{\gamma}} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (1.5)$$

will play a major role in our results. The first eigenvalue of this eigenvalue problem is positive and is given by

$$\lambda_1(\gamma) = \inf_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{\int_{\Omega} |\nabla u|^2 dx}{\int_{\Omega} \frac{u^2}{|x|^{\gamma}} dx}. \quad (1.6)$$

Set

$$G(t) = \int_0^t h(s) e^{\alpha s^2} ds,$$

let  $d$  be the radius of the largest open ball centered at the origin that is contained in  $\Omega$ , and let

$$\kappa = \frac{(2 - \gamma)^2}{2d^{2-\gamma}}.$$

Our first result is the following theorem.

**Theorem 1.1.** Assume that  $\alpha > 0$  and  $0 \leq \gamma < 2$  satisfy  $\alpha/4\pi + \gamma/2 \leq 1$ ,  $h$  satisfies (1.2) and (1.3),  $G$  satisfies

$$G(t) \geq 0 \quad \text{for } t \geq 0, \quad (1.7)$$

$$G(t) \leq \frac{1}{2} (\lambda_1(\gamma) - \sigma_1) t^2 \quad \text{for } |t| \leq \delta \quad (1.8)$$

for some  $\sigma_1, \delta > 0$ , and

$$\beta > \frac{\kappa}{\alpha}. \quad (1.9)$$

Then problem (1.1) has a nontrivial solution.

This theorem is new even in the nonsingular case  $\gamma = 0$ . Indeed, the corresponding result for the nonsingular case is proved in de Figueiredo et al. [4, 5] and Marcos B. do Ó [8] only assuming that  $h(t) \geq 0$  for all  $t \geq 0$ . This implies our assumption (1.7), but clearly (1.7) is weaker. Moreover, we can further weaken the assumption (1.7) if we assume a larger lower bound on  $\beta$ . We have the following theorem.

**Theorem 1.2.** Assume that  $\alpha > 0$  and  $0 \leq \gamma < 2$  satisfy  $\alpha/4\pi + \gamma/2 \leq 1$ ,  $h$  satisfies (1.2) and (1.3),  $G$  satisfies

$$G(t) \geq -\frac{1}{2} \sigma_0 t^2 \quad \text{for } t \geq 0, \quad (1.10)$$

$$G(t) \leq \frac{1}{2} (\lambda_1(\gamma) - \sigma_1) t^2 \quad \text{for } |t| \leq \delta$$

for some  $\sigma_0, \sigma_1, \delta > 0$ , and

$$\beta > \begin{cases} \frac{2\kappa}{\alpha} \frac{e^{\sigma_0/\kappa}}{3 - e^{\sigma_0/\kappa}} & \text{if } \sigma_0 \leq \kappa \log 2 \\ \frac{2\kappa}{\alpha} e^{\sigma_0/\kappa} & \text{if } \sigma_0 > \kappa \log 2. \end{cases} \quad (1.11)$$

Then problem (1.1) has a nontrivial solution.

*Remark 1.3.* We note that Theorem 1.2 reduces to Theorem 1.1 when  $\sigma_0 = 0$ .

Now let  $(\lambda_k(\gamma))$  be the sequence of eigenvalues of the eigenvalue problem (1.5), repeated according to multiplicity (see Proposition 2.2 in the next section). Our last result is the following theorem.

**Theorem 1.4.** *Assume that  $\alpha > 0$  and  $0 \leq \gamma < 2$  satisfy  $\alpha/4\pi + \gamma/2 \leq 1$ ,  $h$  satisfies (1.2) and (1.3), and  $G$  satisfies*

$$G(t) \geq \frac{1}{2} (\lambda_{k-1}(\gamma) + \sigma_0) t^2 \quad \forall t, \quad (1.12)$$

$$G(t) \leq \frac{1}{2} (\lambda_k(\gamma) - \sigma_1) t^2 \quad \text{for } |t| \leq \delta \quad (1.13)$$

*for some  $k \geq 2$  and  $\sigma_0, \sigma_1, \delta > 0$ . Then there exists a constant  $c > 0$  depending on  $\Omega$ ,  $\alpha$ ,  $\gamma$ , and  $k$ , but not on  $\sigma_0$ ,  $\sigma_1$ , or  $\delta$ , such that if*

$$\beta > \frac{2\kappa}{\alpha} e^{c/\sigma_0}, \quad (1.14)$$

*then problem (1.1) has a nontrivial solution.*

This theorem is also new even in the nonsingular case  $\gamma = 0$ . The corresponding result for the nonsingular case is proved in de Figueiredo et al. [4, 5] only under the additional assumption that  $0 < 2G(t) \leq th(t) e^{\alpha t^2}$  for all  $t \in \mathbb{R} \setminus \{0\}$ .

Proofs of Theorems 1.1 and 1.2 will be given in Section 3 and the proof of Theorem 1.4 will be given in Section 4, after proving a suitable compactness property of the associated variational functional in the next section.

## 2 Preliminaries

Weak solutions of problem (1.1) coincide with critical points of the  $C^1$ -functional

$$E(u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - \int_{\Omega} \frac{G(u)}{|x|^\gamma} dx, \quad u \in H_0^1(\Omega).$$

We recall that a  $(PS)_c$  sequence of  $E$  is a sequence  $(u_j) \subset H_0^1(\Omega)$  such that  $E(u_j) \rightarrow c$  and  $E'(u_j) \rightarrow 0$ . Proofs of Theorems 1.1 and 1.2 will be based on the following compactness result.

**Proposition 2.1.** *Assume that  $\alpha > 0$  and  $0 \leq \gamma < 2$  satisfy  $\alpha/4\pi + \gamma/2 \leq 1$ , and  $h$  satisfies (1.2) and (1.3). Then for all  $c \neq 0$  with*

$$c < \frac{2\pi}{\alpha} \left(1 - \frac{\gamma}{2}\right),$$

*every  $(PS)_c$  sequence of  $E$  has a subsequence that converges weakly to a nontrivial solution of problem (1.1).*

*Proof.* Let  $(u_j) \subset H_0^1(\Omega)$  be a  $(PS)_c$  sequence of  $E$ . Then

$$E(u_j) = \frac{1}{2} \|u_j\|^2 - \int_{\Omega} \frac{G(u_j)}{|x|^\gamma} dx = c + o(1) \quad (2.1)$$

and

$$E'(u_j) u_j = \|u_j\|^2 - \int_{\Omega} u_j h(u_j) \frac{e^{\alpha u_j^2}}{|x|^\gamma} dx = o(\|u_j\|). \quad (2.2)$$

First we show that  $(u_j)$  is bounded in  $H_0^1(\Omega)$ . Multiplying (2.1) by 4 and subtracting (2.2) gives

$$\|u_j\|^2 + \int_{\Omega} \left( u_j h(u_j) e^{\alpha u_j^2} - 4G(u_j) \right) \frac{dx}{|x|^\gamma} = 4c + o(\|u_j\| + 1),$$

so it suffices to show that  $th(t) e^{\alpha t^2} - 4G(t)$  is bounded from below. Let  $0 < \varepsilon \leq \beta/5$ . By (1.2) and (1.3), for some constant  $C_\varepsilon > 0$ ,

$$|G(t)| \leq \varepsilon e^{\alpha t^2} + C_\varepsilon \quad (2.3)$$

and

$$th(t) e^{\alpha t^2} \geq (\beta - \varepsilon) e^{\alpha t^2} - C_\varepsilon \quad (2.4)$$

for all  $t$ , and the desired conclusion follows.

Since  $(u_j)$  is bounded in  $H_0^1(\Omega)$ , a renamed subsequence converges to some  $u$  weakly in  $H_0^1(\Omega)$ , strongly in  $L^p(\Omega)$  for all  $p \in [1, \infty)$ , and a.e. in  $\Omega$ . We have

$$E'(u_j) v = \int_{\Omega} \nabla u_j \cdot \nabla v dx - \int_{\Omega} v h(u_j) \frac{e^{\alpha u_j^2}}{|x|^\gamma} dx \rightarrow 0 \quad (2.5)$$

for all  $v \in H_0^1(\Omega)$ . By (1.2), given any  $\varepsilon > 0$ , there exists a constant  $C_\varepsilon > 0$  such that

$$|h(t) e^{\alpha t^2}| \leq \varepsilon e^{\alpha t^2} + C_\varepsilon \quad \forall t. \quad (2.6)$$

By (2.2),

$$\sup_j \int_{\Omega} u_j h(u_j) \frac{e^{\alpha u_j^2}}{|x|^\gamma} dx < \infty,$$

which together with (2.4) gives

$$\sup_j \int_{\Omega} \frac{e^{\alpha u_j^2}}{|x|^\gamma} dx < \infty. \quad (2.7)$$

For  $v \in C_0^\infty(\Omega)$ , it follows from (2.6) and (2.7) that the sequence  $(v h(u_j) e^{\alpha u_j^2}/|x|^\gamma)$  is uniformly integrable and hence

$$\int_{\Omega} v h(u_j) \frac{e^{\alpha u_j^2}}{|x|^\gamma} dx \rightarrow \int_{\Omega} v h(u) \frac{e^{\alpha u^2}}{|x|^\gamma} dx$$

by Vitali's convergence theorem, so it follows from (2.5) that

$$\int_{\Omega} \nabla u \cdot \nabla v dx - \int_{\Omega} v h(u) \frac{e^{\alpha u^2}}{|x|^\gamma} dx = 0.$$

Then this holds for all  $v \in H_0^1(\Omega)$  by density, so the weak limit  $u$  is a solution of problem (1.1).

Suppose that  $u = 0$ . Then

$$\int_{\Omega} \frac{G(u_j)}{|x|^\gamma} dx \rightarrow 0$$

since (2.3) and (2.7) imply that the sequence  $(G(u_j)/|x|^\gamma)$  is uniformly integrable, so (2.1) gives  $c \geq 0$  and

$$\|u_j\| \rightarrow (2c)^{1/2}. \quad (2.8)$$

Let  $2c < \nu < 4\pi(1 - \gamma/2)/\alpha$ . Then  $\|u_j\| \leq \nu^{1/2}$  for all  $j \geq j_0$  for some  $j_0$ . Let  $q = 4\pi(1 - \gamma/2)/\alpha\nu > 1$  and let  $1/(1 - 1/q) < r < 2/\gamma(1 - 1/q)$ . By the Hölder inequality,

$$\left| \int_{\Omega} u_j h(u_j) \frac{e^{\alpha u_j^2}}{|x|^\gamma} dx \right| \leq \left( \int_{\Omega} |u_j h(u_j)|^p dx \right)^{1/p} \left( \int_{\Omega} \frac{e^{q\alpha u_j^2}}{|x|^\gamma} dx \right)^{1/q} \left( \int_{\Omega} \frac{dx}{|x|^{\gamma r(1-1/q)}} \right)^{1/r},$$

where  $1/p + 1/q + 1/r = 1$ . The first integral on the right-hand side converges to zero since  $th(t)$  is bounded and  $u = 0$ , the second integral is bounded for  $j \geq j_0$  by (1.4) since  $q\alpha u_j^2 = 4\pi(1 - \gamma/2)\tilde{u}_j^2$ , where  $\tilde{u}_j = u_j/\nu^{1/2}$  satisfies  $\|\tilde{u}_j\| \leq 1$ , and the last integral is finite since  $\gamma r(1 - 1/q) < 2$ , so

$$\int_{\Omega} u_j h(u_j) \frac{e^{\alpha u_j^2}}{|x|^\gamma} dx \rightarrow 0.$$

Then  $u_j \rightarrow 0$  by (2.2) and hence  $c = 0$  by (2.8), contrary to assumption. So  $u$  is nontrivial.  $\square$

We close this preliminary section with a basic result for the singular eigenvalue problem (1.5). Set  $\omega(x) = |x|^{-\gamma}$  and let  $L^2(\Omega, \omega)$  be the weighted Lebesgue space with the norm

$$\|u\|_{2, \omega} = \left( \int_{\Omega} \omega(x) |u(x)|^2 dx \right)^{1/2}. \quad (2.9)$$

**Proposition 2.2.** *The eigenvalues of the eigenvalue problem (1.5) are positive, have finite multiplicities, and form a nondecreasing sequence  $\lambda_k(\gamma) \rightarrow \infty$ . The space  $L^2(\Omega, \omega)$  has an orthonormal basis consisting of eigenfunctions that are also orthogonal in  $H_0^1(\Omega)$ . Moreover, the eigenfunctions belong to  $C^\alpha(\overline{\Omega})$  for some  $\alpha \in (0, 1)$ .*

*Proof.* Since  $H_0^1(\Omega)$  is compactly embedded in  $L^p(\Omega)$  for  $1 \leq p < \infty$  by the Sobolev embedding theorem and  $L^p(\Omega) \hookrightarrow L^2(\Omega, \omega)$  for  $p > 4/(2 - \gamma)$  by the Hölder inequality,  $H_0^1(\Omega)$  is compactly embedded in  $L^2(\Omega, \omega)$ . The eigenvalue problem (1.5) can be written as

$$Su = \lambda^{-1} u,$$

where  $S : L^2(\Omega, \omega) \rightarrow H_0^1(\Omega)$ ,  $f \mapsto u$  is the solution operator for the singular boundary value problem

$$\begin{cases} -\Delta u = \frac{f(x)}{|x|^\gamma} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Since  $S : L^2(\Omega, \omega) \rightarrow L^2(\Omega, \omega)$  is a compact symmetric operator, the first part of the proposition follows from the spectral theorem.

Let  $u$  be an eigenfunction, let  $1 < q < 2/\gamma$ , and let  $1 < s < 2/\gamma q$ . By the Hölder inequality,

$$\int_{\Omega} \left| \frac{u}{|x|^\gamma} \right|^q dx \leq \left( \int_{\Omega} |u|^{qr} dx \right)^{1/r} \left( \int_{\Omega} \frac{dx}{|x|^{\gamma qs}} \right)^{1/s},$$

where  $1/r + 1/s = 1$ . The first integral on the right-hand side is finite by the Sobolev embedding, and so is the second integral since  $\gamma qs < 2$ , so  $u/|x|^\gamma \in L^q(\Omega)$ . By the Calderon-Zygmund inequality, then  $u \in W^{2,q}(\Omega)$ , which is embedded in  $C^\alpha(\overline{\Omega})$  for  $\alpha = 2 - 2/q$  when  $1 < q < 2$ .  $\square$

### 3 Proofs of Theorems 1.1 and 1.2

In this section we prove Theorems 1.1 and 1.2 by showing that the functional  $E$  has the mountain pass geometry with the mountain pass level  $c \in (0, 2\pi(1 - \gamma/2)/\alpha)$  and applying Proposition 2.1.

**Lemma 3.1.** *If (1.8) holds, then there exists a  $\rho > 0$  such that*

$$\inf_{\|u\|=\rho} E(u) > 0.$$

*Proof.* Since (1.2) implies that  $h$  is bounded, there exists a constant  $C_\delta > 0$  such that

$$|G(t)| \leq C_\delta |t|^3 e^{\alpha t^2} \quad \text{for } |t| > \delta,$$

which together with (1.8) gives

$$\int_{\Omega} \frac{G(u)}{|x|^\gamma} dx \leq \frac{1}{2} (\lambda_1(\gamma) - \sigma_1) \int_{\Omega} \frac{u^2}{|x|^\gamma} dx + C_\delta \int_{\Omega} |u|^3 \frac{e^{\alpha u^2}}{|x|^\gamma} dx. \quad (3.1)$$

By (1.6),

$$\int_{\Omega} \frac{u^2}{|x|^\gamma} dx \leq \frac{\rho^2}{\lambda_1(\gamma)}, \quad (3.2)$$

where  $\rho = \|u\|$ . Let  $2 < r < 4/\gamma$ . By the Hölder inequality,

$$\int_{\Omega} |u|^3 \frac{e^{\alpha u^2}}{|x|^\gamma} dx \leq \left( \int_{\Omega} |u|^{3p} dx \right)^{1/p} \left( \int_{\Omega} \frac{e^{2\alpha u^2}}{|x|^\gamma} dx \right)^{1/2} \left( \int_{\Omega} \frac{dx}{|x|^{\gamma r/2}} \right)^{1/r}, \quad (3.3)$$

where  $1/p + 1/r = 1/2$ . The first integral on the right-hand side is bounded by  $C\rho^{3p}$  for some constant  $C > 0$  by the Sobolev embedding. Since  $2\alpha u^2 = 2\alpha\rho^2 \tilde{u}^2$ , where  $\tilde{u} = u/\rho$  satisfies  $\|\tilde{u}\| = 1$ , the second integral is bounded when  $\rho^2 \leq 2\pi(1 - \gamma/2)/\alpha$  by (1.4). The last integral is finite since  $\gamma r < 4$ . So combining (3.1)–(3.3) gives

$$\int_{\Omega} \frac{G(u)}{|x|^\gamma} dx \leq \frac{1}{2} \left( 1 - \frac{\sigma_1}{\lambda_1(\gamma)} \right) \rho^2 + O(\rho^3) \quad \text{as } \rho \rightarrow 0.$$

Then

$$E(u) \geq \frac{1}{2} \frac{\sigma_1}{\lambda_1(\gamma)} \rho^2 + O(\rho^3),$$

and the desired conclusion follows from this for sufficiently small  $\rho > 0$ .  $\square$

We have  $B_d(0) \subset \Omega$ . For  $j \geq 2$ , let

$$\omega_j(x) = \frac{1}{\sqrt{2\pi}} \begin{cases} \sqrt{\log j} & \text{if } |x| \leq d/j \\ \frac{\log(d/|x|)}{\sqrt{\log j}} & \text{if } d/j < |x| < d \\ 0 & \text{otherwise.} \end{cases} \quad (3.4)$$

It is easily seen that  $\omega_j \in H_0^1(\Omega)$  with  $\|\omega_j\| = 1$ . Moreover,

$$\int_{\Omega} \frac{\omega_j}{|x|^\gamma} dx = \frac{d^{2-\gamma}}{(2-\gamma)^2} \sqrt{\frac{2\pi}{\log j}} \left( 1 - \frac{1}{j^{2-\gamma}} \right) \quad (3.5)$$



and

$$\int_{\Omega} \frac{\omega_j^2}{|x|^\gamma} dx = \frac{2d^{2-\gamma}}{(2-\gamma)^3 \log j} \left[ 1 - \frac{(2-\gamma) \log j + 1}{j^{2-\gamma}} \right]. \quad (3.6)$$

**Lemma 3.2.** *Assume that  $h$  satisfies (1.2) and (1.3).*

- (i) *For all  $j \geq 2$ ,  $E(t\omega_j) \rightarrow -\infty$  as  $t \rightarrow \infty$ .*
- (ii) *If (1.8) holds, then there exists a  $j_0 \geq 2$  such that*

$$\sup_{t \geq 0} E(t\omega_{j_0}) < \frac{2\pi}{\alpha} \left( 1 - \frac{\gamma}{2} \right) \quad (3.7)$$

*in each of the following cases:*

- (a) *(1.7) and (1.9) hold,*
- (b) *(1.10) and (1.11) hold.*

*Proof.* Fix  $\varepsilon > 0$ . By (1.3),  $\exists M_\varepsilon > 0$  such that

$$th(t) e^{\alpha t^2} > (\beta - \varepsilon) e^{\alpha t^2} \quad \text{for } |t| > M_\varepsilon. \quad (3.8)$$

Since  $e^{\alpha t^2} > \alpha^2 t^4 / 2$  for all  $t$ , then there exists a constant  $C_\varepsilon > 0$  such that

$$th(t) e^{\alpha t^2} \geq \frac{1}{2} (\beta - \varepsilon) \alpha^2 t^4 - C_\varepsilon |t| \quad (3.9)$$

and

$$G(t) \geq \frac{1}{8} (\beta - \varepsilon) \alpha^2 t^4 - C_\varepsilon |t| \quad (3.10)$$

for all  $t$ . Since  $\|\omega_j\| = 1$  and  $\omega_j \geq 0$ , then

$$E(t\omega_j) \leq \frac{t^2}{2} - \frac{1}{8} (\beta - \varepsilon) \alpha^2 t^4 \int_{\Omega} \frac{\omega_j^4}{|x|^\gamma} dx + C_\varepsilon t \int_{\Omega} \frac{\omega_j}{|x|^\gamma} dx,$$

and (i) follows.

Set

$$H_j(t) = E(t\omega_j) = \frac{t^2}{2} - \int_{\Omega} \frac{G(t\omega_j)}{|x|^\gamma} dx, \quad t \geq 0.$$

If (ii) is false, then it follows from Lemma 3.1 and (i) that for all  $j \geq 2$ ,  $\exists t_j > 0$  such that

$$H_j(t_j) = \frac{t_j^2}{2} - \int_{\Omega} \frac{G(t_j\omega_j)}{|x|^\gamma} dx = \sup_{t \geq 0} H_j(t) \geq \frac{2\pi}{\alpha} \left( 1 - \frac{\gamma}{2} \right), \quad (3.11)$$

$$H'_j(t_j) = t_j - \int_{\Omega} \omega_j h(t_j\omega_j) \frac{e^{\alpha t_j^2 \omega_j^2}}{|x|^\gamma} dx = 0. \quad (3.12)$$

Since  $G(t) \geq -C_\varepsilon t$  for all  $t \geq 0$  by (3.10), (3.11) gives

$$t_j^2 \geq t_0^2 - 2\delta_j t_j, \quad (3.13)$$

where

$$t_0 = \sqrt{\frac{4\pi}{\alpha} \left(1 - \frac{\gamma}{2}\right)}$$

and

$$\delta_j = C_\varepsilon \int_{\Omega} \frac{\omega_j}{|x|^\gamma} dx \rightarrow 0 \quad \text{as } j \rightarrow \infty$$

by (3.5). First we will show that  $t_j \rightarrow t_0$ .

By (3.13),  $t_j \geq \sqrt{t_0^2 + \delta_j^2} - \delta_j$  and hence

$$\liminf_{j \rightarrow \infty} t_j \geq t_0. \quad (3.14)$$

Write (3.12) as

$$t_j^2 = \int_{\{t_j \omega_j > M_\varepsilon\}} t_j \omega_j h(t_j \omega_j) \frac{e^{\alpha t_j^2 \omega_j^2}}{|x|^\gamma} dx + \int_{\{t_j \omega_j \leq M_\varepsilon\}} t_j \omega_j h(t_j \omega_j) \frac{e^{\alpha t_j^2 \omega_j^2}}{|x|^\gamma} dx =: I_1 + I_2. \quad (3.15)$$

Set  $r_j = d e^{-M_\varepsilon \sqrt{2\pi \log j}/t_j}$ . Since  $\liminf t_j > 0$ , for all sufficiently large  $j$ ,  $d/j < r_j < d$  and  $t_j \omega_j(x) > M_\varepsilon$  if and only if  $|x| < r_j$ . So (3.8) gives

$$\begin{aligned} I_1 &\geq (\beta - \varepsilon) \int_{\{|x| < r_j\}} \frac{e^{\alpha t_j^2 \omega_j^2}}{|x|^\gamma} dx = (\underline{\beta} - \varepsilon) \left( \int_{\{|x| \leq d/j\}} \frac{e^{\alpha t_j^2 \omega_j^2}}{|x|^\gamma} dx \right. \\ &\quad \left. + \int_{\{d/j < |x| < r_j\}} \frac{e^{\alpha t_j^2 \omega_j^2}}{|x|^\gamma} dx \right) =: (\beta - \varepsilon) (I_3 + I_4). \end{aligned} \quad (3.16)$$

We have

$$I_3 = e^{\alpha t_j^2 \log j / 2\pi} \int_{\{|x| \leq d/j\}} \frac{dx}{|x|^\gamma} = \frac{2\pi}{2 - \gamma} \left(\frac{d}{j}\right)^{2-\gamma} j^{\alpha t_j^2 / 2\pi} = \frac{2\pi d^{2-\gamma}}{2 - \gamma} j^{\alpha(t_j^2 - t_0^2)/2\pi}. \quad (3.17)$$

Since  $th(t) e^{\alpha t^2} \geq -C_\varepsilon t$  for all  $t \geq 0$  by (3.9),

$$I_2 \geq -C_\varepsilon t_j \int_{\{t_j \omega_j \leq M_\varepsilon\}} \frac{\omega_j}{|x|^\gamma} dx \geq -\delta_j t_j. \quad (3.18)$$

Combining (3.15)–(3.18) and noting that  $I_4 \geq 0$  gives

$$t_j^2 \geq (\beta - \varepsilon) \frac{2\pi d^{2-\gamma}}{2-\gamma} j^{\alpha(t_j^2 - t_0^2)/2\pi} - \delta_j t_j.$$

It follows from this that

$$\limsup_{j \rightarrow \infty} t_j \leq t_0,$$

which together with (3.14) shows that  $t_j \rightarrow t_0$ .

Next we estimate  $I_4$ . We have

$$\begin{aligned} I_4 &= \int_{\{d/j < |x| < r_j\}} \frac{e^{\alpha t_j^2 [\log(d/|x|)]^2 / 2\pi \log j}}{|x|^\gamma} dx \\ &= 2\pi \left( \int_{d/j}^d e^{\alpha t_j^2 [\log(d/r)]^2 / 2\pi \log j} r^{1-\gamma} dr - \int_{r_j}^d e^{\alpha t_j^2 [\log(d/r)]^2 / 2\pi \log j} r^{1-\gamma} dr \right) \\ &= 2\pi d^{2-\gamma} \left( \log j \int_0^1 e^{-(2-\gamma)t[1-(t_j/t_0)^2 t]} \log j dt - \int_{s_j}^1 s^{1-\gamma} e^{\alpha t_j^2 (\log s)^2 / 2\pi \log j} ds \right), \end{aligned} \tag{3.19}$$

where  $t = \log(d/r)/\log j$ ,  $s = r/d$ , and  $s_j = r_j/d = e^{-M_\varepsilon \sqrt{2\pi \log j}/t_j} \rightarrow 0$ . For  $s_j < s < 1$ ,  $\alpha t_j^2 (\log s)^2 / 2\pi \log j$  is bounded by  $\alpha M_\varepsilon^2$  and goes to zero as  $j \rightarrow \infty$ , so the last integral converges to

$$\int_0^1 s^{1-\gamma} ds = \frac{1}{2-\gamma}.$$

So combining (3.15)–(3.19) and letting  $j \rightarrow \infty$  gives

$$t_0^2 \geq (\beta - \varepsilon) \frac{2\pi d^{2-\gamma}}{2-\gamma} (L_1 + L_2 - 1),$$

where

$$L_1 = \liminf_{j \rightarrow \infty} j^{\alpha(t_j^2 - t_0^2)/2\pi},$$

$$L_2 = \liminf_{j \rightarrow \infty} \int_0^1 n e^{-nt[1-(t_j/t_0)^2 t]} dt,$$

and  $n = (2-\gamma) \log j \rightarrow \infty$ . Letting  $\varepsilon \rightarrow 0$  in this inequality gives

$$\beta \leq \frac{2\kappa}{\alpha(L_1 + L_2 - 1)}. \tag{3.20}$$

We will show that this leads to a contradiction if (a) or (b) holds.

(a) By (1.7),  $G(t_j \omega_j) \geq 0$  and hence (3.11) gives  $t_j \geq t_0$ , so  $L_1 \geq 1$  and

$$L_2 \geq \lim_{n \rightarrow \infty} \int_0^1 n e^{-nt(1-t)} dt = 2$$

(see de Figueiredo et al. [4, 5]). Then (3.20) gives  $\beta \leq \kappa/\alpha$ , contradicting (1.9).

(b) By (1.10),  $G(t_j \omega_j) \geq -\sigma_0 t_j^2 \omega_j^2/2$  and hence (3.11) and (3.6) give

$$t_j^2 - t_0^2 \geq -\sigma_0 t_j^2 \int_{\Omega} \frac{\omega_j^2}{|x|^\gamma} dx \geq -\frac{\sigma_0 t_j^2}{\kappa(2-\gamma) \log j} = -\frac{\sigma_0 t_j^2}{\kappa n},$$

so

$$t_j^2 - t_0^2 \geq -\frac{\sigma_0 t_0^2}{\kappa(2-\gamma) \log j + \sigma_0} \geq -\frac{2\pi\sigma_0}{\alpha\kappa \log j}$$

and

$$\left(\frac{t_j}{t_0}\right)^2 \geq \frac{\kappa n}{\kappa n + \sigma_0} \geq 1 - \frac{\sigma_0}{\kappa n}.$$

So

$$L_1 = \liminf_{j \rightarrow \infty} e^{\alpha(t_j^2 - t_0^2) \log j / 2\pi} \geq e^{-\sigma_0/\kappa}$$

and

$$L_2 \geq \liminf_{n \rightarrow \infty} \int_0^1 n e^{-nt(1-t) - \sigma_0 t^2/\kappa} dt \geq e^{-\sigma_0/\kappa} \left( \lim_{n \rightarrow \infty} \int_0^1 n e^{-nt(1-t)} dt \right) = 2e^{-\sigma_0/\kappa}.$$

On the other hand,

$$L_2 \geq \lim_{n \rightarrow \infty} \int_0^1 n e^{-nt} dt = 1.$$

Then (3.20) gives

$$\beta \leq \frac{2\kappa}{\alpha(e^{-\sigma_0/\kappa} + \max\{2e^{-\sigma_0/\kappa}, 1\} - 1)},$$

contradicting (1.11). □

We can now conclude the proofs of Theorems 1.1 and 1.2. Let  $j_0$  be as in Lemma 3.2 (ii). By Lemma 3.2 (i),  $\exists R > \rho$  such that  $E(R\omega_{j_0}) \leq 0$ , where  $\rho$  is as in Lemma 3.1. Let

$$\Gamma = \{\gamma \in C([0, 1], H_0^1(\Omega)) : \gamma(0) = 0, \gamma(1) = R\omega_{j_0}\}$$

be the class of paths joining the origin to  $R\omega_{j_0}$ , and set

$$c := \inf_{\gamma \in \Gamma} \max_{u \in \gamma([0,1])} E(u).$$

By Lemma 3.1,  $c > 0$ . Since the path  $\gamma_0(t) = tR\omega_{j_0}$ ,  $t \in [0, 1]$  is in  $\Gamma$ ,

$$c \leq \max_{u \in \gamma_0([0,1])} E(u) \leq \sup_{t \geq 0} E(t\omega_{j_0}) < \frac{2\pi}{\alpha} \left(1 - \frac{\gamma}{2}\right)$$

by (3.7). If there are no  $(PS)_c$  sequences of  $E$ , then  $E$  satisfies the  $(PS)_c$  condition vacuously and hence has a critical point  $u$  at the level  $c$  by the mountain pass theorem. Then  $u$  is a solution of problem (1.1) and  $u$  is nontrivial since  $c > 0$ . So we may assume that  $E$  has a  $(PS)_c$  sequence. Then this sequence has a subsequence that converges weakly to a nontrivial solution of problem (1.1) by Proposition 2.1.

## 4 Proof of Theorem 1.4

In this section we prove Theorem 1.4 by showing that the functional  $E$  has the linking geometry with the minimax level  $c \in (0, 2\pi(1 - \gamma/2)/\alpha)$  and applying Proposition 2.1. We have the direct sum decomposition

$$H_0^1(\Omega) = V \oplus W, \quad u = v + w,$$

where  $V$  is the subspace spanned by the eigenfunctions of  $\lambda_1(\gamma), \dots, \lambda_{k-1}(\gamma)$  and  $W$  is the closure of the subspace spanned by the eigenfunctions of  $\lambda_k(\gamma), \lambda_{k+1}(\gamma), \dots$ . It follows from Proposition 2.2 that

$$\int_{\Omega} |\nabla v|^2 dx \leq \lambda_{k-1}(\gamma) \int_{\Omega} \frac{v^2}{|x|^\gamma} dx \quad \forall v \in V \quad (4.1)$$

and

$$\int_{\Omega} |\nabla w|^2 dx \geq \lambda_k(\gamma) \int_{\Omega} \frac{w^2}{|x|^\gamma} dx \quad \forall w \in W. \quad (4.2)$$

**Lemma 4.1.** *If (1.13) holds, then there exists a  $\rho > 0$  such that*

$$\inf_{\substack{w \in W \\ \|w\| = \rho}} E(w) > 0.$$

*Proof.* As in the proof of Lemma 3.1 in the last section, (1.13) gives

$$\int_{\Omega} \frac{G(w)}{|x|^\gamma} dx \leq \frac{1}{2} (\lambda_k(\gamma) - \sigma_2) \int_{\Omega} \frac{w^2}{|x|^\gamma} dx + C_\delta \int_{\Omega} |w|^3 \frac{e^{\alpha w^2}}{|x|^\gamma} dx$$

for some constant  $C_\delta > 0$  and the last integral is  $O(\rho^3)$  as  $\rho = \|w\| \rightarrow 0$ . Since

$$\int_{\Omega} \frac{w^2}{|x|^\gamma} dx \leq \frac{\rho^2}{\lambda_k(\gamma)}$$

by (4.2), then

$$E(w) \geq \frac{1}{2} \frac{\sigma_2}{\lambda_k(\gamma)} \rho^2 + O(\rho^3) \quad \text{as } \rho \rightarrow 0,$$

and the desired conclusion follows for sufficiently small  $\rho > 0$ .  $\square$

Let  $\omega_j$  be as in (3.4), and set

$$Q_{j,R} = \{u = v + t\omega_j : v \in V, t \geq 0, \|u\| \leq R\}$$

for  $R > 0$ .

**Lemma 4.2.** *Assume that  $h$  satisfies (1.2) and (1.3).*

(i) *If (1.12) holds, then for all  $j \geq 2$ , there exists a  $R_j > 0$  such that*

$$\sup_{u \in \partial Q_{j,R}} E(u) = 0 \quad \forall R \geq R_j.$$

(ii) *If (1.12) and (1.13) hold, then there exists a constant  $c > 0$  depending only on  $\Omega$ ,  $\alpha$ ,  $\gamma$ , and  $k$  such that whenever*

$$\beta > \frac{2\kappa}{\alpha} e^{c/\sigma_0},$$

*there exists a  $j_0 \geq 2$  such that*

$$\sup_{v \in V, t \geq 0} E(v + t\omega_{j_0}) < \frac{2\pi}{\alpha} \left(1 - \frac{\gamma}{2}\right). \quad (4.3)$$

*Proof.* As in the proof of Lemma 3.2, for each  $\varepsilon > 0$ , there exist constants  $M_\varepsilon, C_\varepsilon > 0$  such that

$$th(t) e^{\alpha t^2} > (\beta - \varepsilon) e^{\alpha t^2} \quad \text{for } |t| > M_\varepsilon \quad (4.4)$$

and

$$th(t) e^{\alpha t^2} \geq \frac{1}{2} (\beta - \varepsilon) \alpha^2 t^4 - C_\varepsilon |t|, \quad (4.5)$$

$$G(t) \geq \frac{1}{8} (\beta - \varepsilon) \alpha^2 t^4 - C_\varepsilon |t|$$

for all  $t$ . Then for all  $u \neq 0$ ,

$$E(tu) \leq \frac{t^2}{2} \int_{\Omega} |\nabla u|^2 dx - \frac{1}{8} (\beta - \varepsilon) \alpha^2 t^4 \int_{\Omega} \frac{u^4}{|x|^\gamma} dx + C_\varepsilon t \int_{\Omega} \frac{|u|}{|x|^\gamma} dx \rightarrow -\infty$$

as  $t \rightarrow \infty$ . On the other hand, by (1.12) and (4.1),

$$E(v) \leq \frac{1}{2} \left( \int_{\Omega} |\nabla v|^2 dx - \lambda_{k-1}(\gamma) \int_{\Omega} \frac{v^2}{|x|^\gamma} dx \right) \leq 0 \quad \forall v \in V.$$

Since  $Q_{j,R}$  lies in a finite dimensional subspace, (i) follows.

We will show that if (1.12) and (1.13) hold, but (4.3) does not hold for any  $j_0 \geq 2$ , then there exists a constant  $c > 0$  depending only on  $\Omega$ ,  $\alpha$ ,  $\gamma$ , and  $k$  such that

$$\beta \leq \frac{2\kappa}{\alpha} e^{c/\sigma_0}. \quad (4.6)$$

We have

$$\sup_{v \in V, t \geq 0} E(v + t\omega_j) \geq \frac{2\pi}{\alpha} \left(1 - \frac{\gamma}{2}\right) \quad \forall j \geq 2.$$

It follows from Lemma 4.1 and (i) that the above supremum is attained at some point  $u_j = v_j + t_j \omega_j$ ,  $v_j \in V$ ,  $t_j > 0$  such that  $E'(u_j) = 0$  on  $\{v + t\omega_j : v \in V, t \geq 0\}$ . Then

$$E(u_j) = \frac{1}{2} \|u_j\|^2 - \int_{\Omega} \frac{G(u_j)}{|x|^\gamma} dx \geq \frac{2\pi}{\alpha} \left(1 - \frac{\gamma}{2}\right), \quad (4.7)$$

$$E'(u_j) u_j = \|u_j\|^2 - \int_{\Omega} u_j h(u_j) \frac{e^{\alpha u_j^2}}{|x|^\gamma} dx = 0. \quad (4.8)$$

Since  $\|\omega_j\| = 1$  and  $G(t) \geq 0$  for all  $t$  by (1.12), (4.7) gives

$$\|v_j\| + t_j \geq t_0,$$

where

$$t_0 = \sqrt{\frac{4\pi}{\alpha} \left(1 - \frac{\gamma}{2}\right)}.$$

First we will show that  $t_j \rightarrow t_0$  and  $v_j \rightarrow 0$  as  $j \rightarrow \infty$ .

Combining (4.7) with (1.12) and (4.1) gives

$$t_j^2 + 2t_j \int_{\Omega} \nabla v_j \cdot \nabla \omega_j dx \geq t_0^2 + 2(\lambda_{k-1}(\gamma) + \sigma_0) t_j \int_{\Omega} \frac{v_j \omega_j}{|x|^\gamma} dx + \sigma_0 \int_{\Omega} \frac{v_j^2}{|x|^\gamma} dx. \quad (4.9)$$

Since  $\nabla \omega_j = 0$  outside  $\{d/j < |x| < d\}$  and  $\omega_j$  is harmonic in  $\{d/j < |x| < d\}$ ,

$$\begin{aligned} \left| \int_{\Omega} \nabla v_j \cdot \nabla \omega_j dx \right| &= \left| \int_{\partial\{d/j < |x| < d\}} v_j \frac{\partial \omega_j}{\partial n} ds \right| \leq |v_j|_{\infty} \int_{\partial\{d/j < |x| < d\}} |\nabla \omega_j| ds \\ &= 2 |v_j|_{\infty} \sqrt{\frac{2\pi}{\log j}}, \end{aligned} \quad (4.10)$$

and

$$\left| \int_{\Omega} \frac{v_j \omega_j}{|x|^{\gamma}} dx \right| \leq |v_j|_{\infty} \int_{\Omega} \frac{\omega_j}{|x|^{\gamma}} dx \leq \frac{d^{2-\gamma}}{(2-\gamma)^2} |v_j|_{\infty} \sqrt{\frac{2\pi}{\log j}} \quad (4.11)$$

by (3.5), where  $|\cdot|_{\infty}$  denotes the  $L^{\infty}(\Omega)$  norm. Combining (4.9)–(4.11) and noting that  $\lambda_{k-1}(\gamma) + \sigma_0 < \lambda_k(\gamma)$  by (1.12) and (1.13) gives

$$\sigma_0 |v_j|_{2,\omega}^2 \leq t_j^2 - t_0^2 + 2 \left( 2 + \lambda_k(\gamma) \frac{d^{2-\gamma}}{(2-\gamma)^2} \right) t_j |v_j|_{\infty} \sqrt{\frac{2\pi}{\log j}}, \quad (4.12)$$

where  $|\cdot|_{2,\omega}$  is defined in (2.9). Since  $V$  is finite dimensional and  $\sigma_0 > 0$ , it follows from this that  $\|v_j\| = O(t_j)$  and then

$$\liminf_{j \rightarrow \infty} t_j \geq t_0. \quad (4.13)$$

Next combining (4.8) with (4.4) and (4.5) gives

$$\begin{aligned} \|u_j\|^2 &= \int_{\{|u_j| > M_{\varepsilon}\}} u_j h(u_j) \frac{e^{\alpha u_j^2}}{|x|^{\gamma}} dx + \int_{\{|u_j| \leq M_{\varepsilon}\}} u_j h(u_j) \frac{e^{\alpha u_j^2}}{|x|^{\gamma}} dx \\ &\geq (\beta - \varepsilon) \int_{\{|u_j| > M_{\varepsilon}\}} \frac{e^{\alpha u_j^2}}{|x|^{\gamma}} dx - C_{\varepsilon} \int_{\{|u_j| \leq M_{\varepsilon}\}} \frac{|u_j|}{|x|^{\gamma}} dx. \end{aligned} \quad (4.14)$$

For  $|x| \leq d/j$ ,

$$|u_j| \geq t_j \omega_j - |v_j| \geq t_j \sqrt{\frac{\log j}{2\pi}} - |v_j|_{\infty},$$

and the last expression is greater than  $M_{\varepsilon}$  for all sufficiently large  $j$  since  $|v_j|_{\infty} = O(t_j)$  and  $\liminf t_j > 0$ , so

$$\begin{aligned} \int_{\{|u_j| > M_{\varepsilon}\}} \frac{e^{\alpha u_j^2}}{|x|^{\gamma}} dx &\geq e^{\alpha (t_j \sqrt{\log j / 2\pi} - |v_j|_{\infty})^2} \int_{\{|x| \leq d/j\}} \frac{dx}{|x|^{\gamma}} \\ &= \frac{2\pi}{2-\gamma} \left( \frac{d}{j} \right)^{2-\gamma} j^{\alpha (t_j - |v_j|_{\infty} \sqrt{2\pi / \log j})^2 / 2\pi} = \frac{2\pi d^{2-\gamma}}{2-\gamma} j^{\alpha [(t_j - |v_j|_{\infty} \sqrt{2\pi / \log j})^2 - t_0^2] / 2\pi} \end{aligned}$$



for large  $j$ . So (4.14) gives

$$(\beta - \varepsilon) j^\alpha [(t_j - |v_j|_\infty \sqrt{2\pi/\log j})^2 - t_0^2]/2\pi \leq \frac{2 - \gamma}{2\pi d^{2-\gamma}} \left[ (\|v_j\| + t_j)^2 + C_\varepsilon \int_\Omega \frac{|v_j|}{|x|^\gamma} dx + \delta_j t_j \right], \quad (4.15)$$

where

$$\delta_j = C_\varepsilon \int_\Omega \frac{\omega_j}{|x|^\gamma} dx \rightarrow 0 \quad \text{as } j \rightarrow \infty$$

by (3.5). Since  $\|v_j\| = O(t_j)$ , it follows from this that

$$\limsup_{j \rightarrow \infty} t_j \leq t_0,$$

which together with (4.13) shows that  $t_j \rightarrow t_0$ . Then (4.12) implies that  $v_j \rightarrow 0$ .

Now the right-hand side of (4.15) goes to  $2\kappa/\alpha$  as  $j \rightarrow \infty$ . If  $\beta \leq 2\kappa/\alpha$ , then we may take any  $c > 0$ , so suppose that  $\underline{\beta} > 2\kappa/\alpha$ . Then for  $\varepsilon < \underline{\beta} - 2\kappa/\alpha$  and all sufficiently large  $j$ , (4.15) gives  $j^\alpha [(t_j - |v_j|_\infty \sqrt{2\pi/\log j})^2 - t_0^2]/2\pi \leq 1$  and hence

$$t_j \leq t_0 + |v_j|_\infty \sqrt{\frac{2\pi}{\log j}}.$$

Combining this with (4.12) gives

$$\|v_j\| \leq \frac{c_1}{\sigma_0 \sqrt{\log j}}, \quad t_j^2 - t_0^2 \geq -\frac{c_2}{\sigma_0 \log j}$$

for some constants  $c_1, c_2 > 0$  depending only on  $\Omega, \alpha, \gamma$ , and  $k$ . Then  $\alpha [(t_j - |v_j|_\infty \sqrt{2\pi/\log j})^2 - t_0^2]/2\pi \geq -c/\sigma_0 \log j$  for some constant  $c > 0$  depending only on  $\Omega, \alpha, \gamma$ , and  $k$ , so (4.15) gives

$$(\beta - \varepsilon) e^{-c/\sigma_0} \leq \frac{2\kappa}{\alpha},$$

and letting  $\varepsilon \rightarrow 0$  gives (4.6).  $\square$

We can now conclude the proof of Theorem 1.4. Let  $c > 0$  be as in Lemma 4.2 (ii), let  $\beta$  satisfy (1.14), and let  $j_0$  also be as in Lemma 4.2 (ii). By Lemma 4.2  $\exists R > \rho$  such that

$$\sup_{u \in \partial Q_{j_0, R}} E(u) = 0,$$

where  $\rho$  is as in Lemma 4.1. Let

$$\Gamma = \left\{ \gamma \in C(Q_{j_0, R}, H_0^1(\Omega)) : \gamma|_{\partial Q_{j_0, R}} = id \right\},$$

and set

$$c := \inf_{\gamma \in \Gamma} \max_{u \in \gamma(Q_{j_0, R})} E(u).$$

By Lemma 4.1,  $c > 0$ . Since the identity mapping is in  $\Gamma$ ,

$$c \leq \max_{u \in Q_{j_0, R}} E(u) \leq \sup_{v \in V, t \geq 0} E(v + t\omega_{j_0}) < \frac{2\pi}{\alpha} \left(1 - \frac{\gamma}{2}\right)$$

by (4.3). If there are no  $(PS)_c$  sequences of  $E$ , then  $E$  satisfies the  $(PS)_c$  condition vacuously and hence has a critical point  $u$  at the level  $c$  by the linking theorem. Then  $u$  is a solution of problem (1.1) and  $u$  is nontrivial since  $c > 0$ . So we may assume that  $E$  has a  $(PS)_c$  sequence. Then this sequence has a subsequence that converges weakly to a nontrivial solution of problem (1.1) by Proposition 2.1.

## References

- [1] Adimurthi. Existence of positive solutions of the semilinear Dirichlet problem with critical growth for the  $n$ -Laplacian. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 17(3):393–413, 1990.
- [2] Adimurthi and K. Sandeep. A singular Moser-Trudinger embedding and its applications. *NoDEA Nonlinear Differential Equations Appl.*, 13(5-6):585–603, 2007.
- [3] Adimurthi and S. L. Yadava. Bifurcation results for semilinear elliptic problems with critical exponent in  $\mathbf{R}^2$ . *Nonlinear Anal.*, 14(7):607–612, 1990.
- [4] D. G. de Figueiredo, O. H. Miyagaki, and B. Ruf. Elliptic equations in  $\mathbf{R}^2$  with nonlinearities in the critical growth range. *Calc. Var. Partial Differential Equations*, 3(2):139–153, 1995.
- [5] D. G. de Figueiredo, O. H. Miyagaki, and B. Ruf. Corrigendum: “Elliptic equations in  $\mathbf{R}^2$  with nonlinearities in the critical growth range”. *Calc. Var. Partial Differential Equations*, 4(2):203, 1996.
- [6] Djairo G. de Figueiredo, João Marcos do Ó, and Bernhard Ruf. On an inequality by N. Trudinger and J. Moser and related elliptic equations. *Comm. Pure Appl. Math.*, 55(2):135–152, 2002.
- [7] Djairo G. de Figueiredo, João Marcos do Ó, and Bernhard Ruf. Elliptic equations and systems with critical Trudinger-Moser nonlinearities. *Discrete Contin. Dyn. Syst.*, 30(2):455–476, 2011.

- [8] João Marcos B. do Ó. Semilinear Dirichlet problems for the  $N$ -Laplacian in  $\mathbb{R}^N$  with nonlinearities in the critical growth range. *Differential Integral Equations*, 9(5):967–979, 1996.
- [9] Yang Yang and Kanishka Perera.  $N$ -Laplacian problems with critical Trudinger-Moser nonlinearities. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 16(4):1123–1138, 2016.
- [10] Zhitao Zhang, Marta Calanchi, and Bernhard Ruf. Elliptic equations in  $\mathbb{R}^2$  with one-sided exponential growth. *Commun. Contemp. Math.*, 6(6):947–971, 2004.