

BOCHNER-RIESZ OPERATORS IN GRAND LEBESGUE SPACES

Maria Rosaria FORMICA ¹, Eugeny OSTROVSKY ² and Leonid SIROTA ³

¹ Università degli Studi di Napoli “Parthenope”, via Generale Parisi 13,
Palazzo Pacanowsky, 80132, Napoli, Italy.

e-mail: mara.formica@uniparthenope.it

^{2,3} Bar-Ilan University, Department of Mathematics and Statistics,
52900, Ramat Gan, Israel.

e-mail: eugostrovsky@list.ru

e-mail: sirota3@bezeqint.net

Abstract

We provide the conditions for the boundedness of the Bochner-Riesz operator acting between two different Grand Lebesgue Spaces. Moreover we obtain a lower estimate for the constant appearing in the Lebesgue-Riesz norm estimation of the Bochner-Riesz operator and we investigate the convergence of the Bochner-Riesz approximation in Lebesgue-Riesz spaces.

Keywords: Bochner-Riesz operator, Lebesgue-Riesz spaces, Grand Lebesgue spaces, Fourier transform, convolution, Beckner-Brascamp-Lieb inequality, Gaussian density, dominated convergence theorem, Bessel’s and Gamma functions, modulus of continuity.

2010 Mathematics Subject Classification: 46E30, 44A35, 42B15.

1 Introduction

The well-known (pseudo-differential) Bochner-Riesz linear operator $B_R^\alpha[f]$, acting on measurable functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ($n > 1$), is defined as follows

$$B_R^\alpha[f](t) \stackrel{def}{=} \int_{\mathbb{R}^n} e^{-i(t,y)} \left(1 - \frac{|y|^2}{R^2} \right)_+^\alpha \tilde{f}(y) dy, \quad (1.1)$$

where

$$R = \text{const} > 0, \quad \alpha = \text{const}, \quad z_+ = \max(z, 0), \quad z \in \mathbb{R}, \quad (1.2)$$

(x, y) stands for the scalar (inner) product of two vectors $x, y \in \mathbb{R}^n$, $|y|^2 := (y, y)$ and \tilde{f} denotes, as ordinary, the Fourier transform

$$\tilde{f}(y) = \int_{\mathbb{R}^n} e^{i(x,y)} f(x) dx. \quad (1.3)$$

The applications of these operators are described in particular in functional analysis (see [8], [21]) and in statistics of random processes and field (see [5], [37]).

The classical Lebesgue-Riesz norm $\|f\|_p$, $p \in [1, \infty]$, for the function f , is denoted by

$$\|f\|_p := \left[\int_{\mathbb{R}^n} |f(x)|^p dx \right]^{1/p}, \quad 1 \leq p < \infty,$$

$$\|f\|_\infty := \operatorname{ess\,sup}_{x \in \mathbb{R}^n} |f(x)|, \quad p = \infty$$

and the corresponding Banach space is, as usually,

$$L_p = L_p(\mathbb{R}^n) = \{ f : \|f\|_p < \infty \}.$$

Denote, for an arbitrary linear or quasi linear operator U , acting from L_p to L_q , $p, q \geq 1$, its norm

$$\|U\|_{p,q} = \|U\|_{p \rightarrow q} := \sup_{0 \neq f \in L_p} \left[\frac{\|Uf\|_q}{\|f\|_p} \right].$$

Of course, the operator U is bounded as the operator acting from the space L_p into the space L_q iff $\|U\|_{p,q} < \infty$.

More generally, for the operator U acting from some Banach space F equipped with the norm $\|\cdot\|_F$ into another (in the general case) Banach space D , having the norm $\|\cdot\|_D$, denote as usually

$$\|U\|_{F \rightarrow D} := \sup_{0 \neq f \in F} \left[\frac{\|Uf\|_D}{\|f\|_F} \right].$$

There exists a huge numbers of works devoted to the p, q estimates for Bochner-Riesz operators B_R^α , as a rule for the case $q = p$, see e.g. [21, chapter 5], [7, 8, 11, 26, 31, 38, 39], etc.

Our aim, in this paper, is to extend some results contained in the above mentioned works concerning upper estimates for the norm of the Bochner-Riesz operator, in the case of different Lebesgue-Riesz spaces (Section 2) and to extend these estimates to the so-called Grand Lebesgue Spaces (GLS), see Section 3.

We deduce also a non trivial quantitative lower estimate for the coefficient in the Lebesgue-Riesz norm estimation for the Bochner-Riesz operator (Section 4) and we study the convergence of the Bochner-Riesz approximation in Lebesgue-Riesz spaces (Section 5).

2 Norm estimates for the Bochner-Riesz operator acting between different Lebesgue-Riesz spaces

We clarify slightly the known results about the Lebesgue-Riesz p, r norms for the operators B_R^α defined in (1.1), see e.g. [8, 21, 10, 26, 39], ecc.

It is important to observe that the Bochner-Riesz operator B_R^α may be rewritten as a convolution, namely

$$B_R^\alpha[f](t) = f * K_\lambda^R(t) = \int_{\mathbb{R}^n} f(t-s) K_\lambda^R(s) ds, \quad (2.1)$$

where $R = \text{const} > 0$,

$$K_\lambda^R(z) = C(\alpha) R^n \frac{J_\lambda(R|z|)}{|Rz|^\lambda}, \quad \lambda = \alpha + \frac{n}{2},$$

$$C(\alpha) := \frac{1}{\Gamma(\alpha+1)}, \quad \alpha > -1,$$

$\Gamma(\cdot)$ is the Gamma-function, $J_\lambda(\cdot)$ is the Bessel function of order λ and $|z| = \sqrt{(z, z)}$.

Notice that, under our restriction on α ,

$$\int_{\mathbb{R}^n} K_\lambda^R(t) dt = 1. \quad (2.2)$$

Evidently,

$$K_\lambda^R(z) = R^n K_\lambda(Rz), \quad (2.3)$$

where in turn

$$K_\lambda(z) \stackrel{\text{def}}{=} K_\lambda^1(z) = C(\alpha) \frac{J_\lambda(|z|)}{|z|^\lambda}, \quad (2.4)$$

see, e.g., [8, pp. 171-172].

Note that, from the identity (2.3), it follows

$$\|K_\lambda^R(z)\|_q = R^{n-n/q} \|K_\lambda\|_q, \quad q \geq 1, \quad (2.5)$$

as long as $R > 0$.

Assume here and in the sequel

$$-1 < \alpha \leq \alpha_0, \quad \alpha_0 := \frac{n-1}{2}, \quad (2.6)$$

and introduce the value

$$q_0 := \frac{n}{(n+1)/2 + \alpha}, \quad (2.7)$$

so that

$$q_0 \geq 1 \iff \frac{-(n+1)}{2} < \alpha \leq \alpha_0,$$

where the inequality on the left hand side is true due to the restriction $\alpha > -1$.

Moreover, using the well-known results about the behavior of the Bessel functions (see, e.g., [8, p. 172]),

$$K_\lambda(z) \asymp |z|^{-(\frac{n+1}{2} + \alpha)}, \quad \text{as } |z| \rightarrow \infty,$$

it is easy to estimate

$$\|K_\lambda\|_q \asymp (q - q_0)^{-1/q}, \quad q > q_0, \quad (2.8)$$

and $\|K_\lambda\|_q = \infty$ otherwise (see also, e.g., [21, pp. 339-341]).

Correspondingly,

$$\|K_\lambda^R\|_q \asymp R^{n-n/q} (q - q_0)^{-1/q}, \quad q > q_0. \quad (2.9)$$

Furthermore, we recall the Beckner-Brascamp-Lieb-Young inequality for the convolution (see [3, 4])

$$\|f * g\|_r \leq \left(\frac{C_p C_q}{C_r} \right)^d \|f\|_p \|g\|_q, \quad 1 + 1/r = 1/p + 1/q, \quad (2.10)$$

where $p, q, r \geq 1$ and

$$C_m = \left(\frac{m^{1/m}}{(m')^{1/m'}} \right)^{\frac{1}{2}}, \quad m' = m/(m-1).$$

Note that

$$\frac{C_p C_q}{C_r} \leq 1.$$

Recently in [20] has been given a generalization of the convolution inequality in the context of the Grand Lebesgue Spaces (see Section 3 for the definition), built on a unimodular locally compact topological group.

The estimate (2.10) is essentially non-improbable. Indeed, the equality in (2.10) is attained iff both functions f, g are proportional to Gaussian densities, namely there exists positive constants c_1, c_2, c_3, c_4 such that

$$f(x) = c_1 \exp(-c_2 \|x\|^2), \quad g(x) = c_3 \exp(-c_4 \|x\|^2), \quad x \in \mathbb{R}^n, \quad (2.11)$$

where $\|\cdot\|$ is the euclidean norm. Then the convolution $f * g$ is also Gaussian and

$$\|f * g\|_r = \left(\frac{C_p C_q}{C_r} \right)^d \|f\|_p \|g\|_q, \quad 1/p + 1/q = 1 + 1/r, \quad p, q, r > 1. \quad (2.12)$$

The following result about the boundedness of the Bochner-Riesz operator, acting between different Lebesgue-Riesz spaces, holds.

Theorem 2.1. *Let $p > 1$ and $f \in L_p(\mathbb{R}^n)$. Let α be a constant such that*

$$-1 < \alpha \leq \alpha_0 := \frac{n-1}{2} \quad (2.13)$$

and

$$q_0 = \frac{n}{(n+1)/2 + \alpha}. \quad (2.14)$$

Let $q, r \geq 1$ such that $1 + 1/r = 1/q + 1/p$ and assume $q > q_0$, $r > r_0$, $p \leq p_0$, where

$$r_0 = r_0(\alpha, n; p) := \frac{pq_0}{p + q_0 - pq_0}, \quad (2.15)$$

$$p_0 = q'_0 = (1 - 1/q_0)^{-1} \quad (2.16)$$

Define, for $R = \text{const} > 0$,

$$W(\alpha, n, R; p, r) \stackrel{\text{def}}{=} R^{n(1/p-1/r)} \cdot [q(p, r) - q_0(\alpha, n)]^{1/p-1-1/r}. \quad (2.17)$$

Then the Bochner-Riesz operator satisfies

$$\|B_R^\alpha[f]\|_r \leq C(\alpha, R) \cdot W(\alpha, n, R; p, r) \cdot \|f\|_p. \quad (2.18)$$

Proof.

If $f \in L_p$ for some value $p > 1$, then by (2.10) it follows

$$\|B_R^\alpha[f]\|_r \leq \|K_\lambda^R\|_q \|f\|_p, \quad (2.19)$$

where $1 + 1/r = 1/q + 1/p$ and $p, q, r \geq 1, q > q_0$. On the other words,

$$q = q(p, r) \stackrel{\text{def}}{=} \frac{pr}{pr + p - r}, \quad q > q_0, \quad r > p.$$

Moreover it is easy to verify that r_0 , defined in (2.15), is such that $r_0 > p$. Under our restrictions $W(\alpha, n, R; p, r)$ is finite and positive.

So we conclude that estimate (2.18) holds. □

3 Main result. Boundedness of Bochner-Riesz operators in Grand Lebesgue Spaces (GLS)

We recall here, for reader convenience, some known definitions and facts from the theory of Grand Lebesgue Spaces (GLS).

Definition 3.1. Let $\psi = \psi(p)$, $p \in (a, b)$, $a, b = \text{const}$, $1 \leq a < b \leq \infty$, be a positive measurable numerical valued function, such that

$$\inf_{p \in (a, b)} \psi(p) > 0. \quad (3.1)$$

The (Banach) Grand Lebesgue Space $G\psi = G\psi(a, b)$ consists of all the real (or complex) numerical valued measurable functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$ having finite norm, defined by

$$\|f\|_{G\psi} := \sup_{p \in (a, b)} \left[\frac{\|f\|_p}{\psi(p)} \right]. \quad (3.2)$$

We agree to write $G\psi$ in the case when $a = 1$ and $b = \infty$.

The function ψ is named *generating function* for the space $G\psi$ and we denote by $\{\psi(\cdot)\}$ the set of all such functions.

For instance

$$\psi(p) := p^{1/m}, \quad p \in [1, \infty), \quad m > 0, \quad (3.3)$$

or

$$\psi(p) := (p - a)^{-\alpha} (b - p)^{-\beta}, \quad p \in (a, b), \quad 1 \leq a < b < \infty, \quad \alpha, \beta \geq 0, \quad (3.4)$$

are generating functions.

If

$$\psi(p) = 1, \quad p = r; \quad \psi(p) = +\infty, \quad p \neq r, \quad r \in [1, \infty),$$

where $C/\infty := 0$, $C \in \mathbb{R}$ (extremal case), then the corresponding $G\psi$ space coincides with the classical Lebesgue-Riesz space $L_r = L_r(\mathbb{R}^d)$.

The Grand Lebesgue Spaces and several generalizations of them have been widely investigated, mainly in the case of GLS on sets of finite measure, (see, e.g., [9, 23, 24, 12, 30, 32, 6, 18], etc). They play an important role in the theory of Partial Differential Equations (PDEs) (see, e.g., [22, 13, 15, 2], etc.), in interpolation theory (see, e.g., [14, 17]), in the theory of Probability ([34, 36, 19]), in Statistics [32, chapter 5], in theory of random fields [27], [35], in Functional Analysis [32], [33], [35] and so one.

These spaces are rearrangement invariant (r.i.) Banach functional (complete) spaces; their fundamental functions have been considered in [35]. They do not coincide, in the general case, with the classical Banach rearrangement functional spaces: Orlicz, Lorentz, Marcinkiewicz etc., see [30, 33]. The belonging of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ to some $G\psi$ space is closely related with its tail function behavior as $t \rightarrow 0+$ as well as when $t \rightarrow \infty$, see [27, 29].

The Grand Lebesgue Spaces can be considered not only on the Euclidean space \mathbb{R}^n equipped with the Lebesgue measure, but also on an arbitrary measurable space with sigma-finite non-trivial measure.

In the following Theorem we investigate the boundedness of the Bochner-Riesz operator acting from some Grand Lebesgue Space $G\psi = G\psi(a, b)$ into another one $G\nu$. We will consider the same restrictions and quantities defined in Theorem 2.1.

Theorem 3.1. *Let $1 \leq a < b \leq \infty$ and $f \in G\psi(a, b)$. Let α be a constant such that*

$$-1 < \alpha \leq \alpha_0 := \frac{n-1}{2}, \quad n > 1.$$

Let $p > 1$, $q, r \geq 1$, p_0, q_0, r_0 and $W(\alpha, n, R; p, r)$ defined as in Theorem 2.1.

Denote

$$s = s(\alpha, n; r) := \min \left\{ b, \frac{rq_0}{rq_0 + q_0 - r} \right\},$$

and

$$d := r_0(\alpha, n; b) = \frac{bq_0}{q_0 + b - bq_0}.$$

For $r \in (d, \infty)$ let $\nu(r)$ be the following generating function

$$\nu(r) = \nu[\psi](r) = \nu[\psi](\alpha, n, R; r) := \inf_{p \in (a, s)} [W(\alpha, n, R; p, r) \psi(p)]. \quad (3.5)$$

Then

$$\|B_R^\alpha[f]\|_{G\nu} \leq C(\alpha, n, R) \|f\|_{G\psi}. \quad (3.6)$$

Proof.

The proof is simple and alike as the one in [36]. First we observe that $\nu(r)$ is finite. One can assume, without loss of generality, $\|f\|_{G\psi} = 1$, then $\|f\|_p \leq \psi(p)$, $p \in (a, b)$. Applying the inequality (2.18) we have

$$\|B_R^\alpha[f]\|_r \leq C(\alpha, n, R) \cdot W(\alpha, n, R; p, r) \cdot \psi(p). \quad (3.7)$$

Taking the minimum over p subject to our limitations, we get

$$\begin{aligned} \|B_R^\alpha[f]\|_r &\leq C(\alpha, n, R) \cdot \inf_{p \in (a, s)} [W(\alpha, n, R; p, r) \cdot \psi(p)] \\ &= C(\alpha, n, R) \nu(r) = C(\alpha, n, R) \nu(r) \|f\|_{G\psi}, \end{aligned} \quad (3.8)$$

which is quite equivalent to our claim in (3.6). □

4 Lower bound for the coefficient in the Lebesgue-Riesz norm estimate for the Bochner-Riesz operator.

Let $p, r > 1$, $n > 1$ and let us introduce the following variable

$$Q_n(p, r) \stackrel{\text{def}}{=} \sup_{\alpha > 0} \sup_{R > 0} W(\alpha, n, R; p, r),$$

where $W(\alpha, n, R; p, r)$ is defined in Section 2. Our target in this Section is a *lower* bound for the above variable.

Theorem 4.1. *Let $p, r > 1$, $n > 1$ and*

$$\Theta(n, p) := (2\pi)^{n(1-p)/2p} p^{-n/2p}.$$

Then

$$Q_n(p, r) \geq \Theta\left(n, \frac{pr}{pr + p - r}\right), \quad r > p. \quad (4.1)$$

Remark 4.1. The *possible* case when $Q_n(p, r) = +\infty$ can not be excluded.

Proof.

We will apply equality (2.12), in which we choose the ordinary Gaussian density

$$f_0(x) := (2\pi)^{-n/2} \exp(-\|x\|^2/2), \quad x \in \mathbb{R}^n,$$

and take $\alpha = R^2/2$. Obviously

$$Q_n(p, r) \geq \lim_{R \rightarrow \infty} W(R^2/2, n, R; p, r).$$

We have

$$B_R^{R^2}[f](t) = \int_{\mathbb{R}^n} e^{-i(t,y)} \left(1 - \frac{\|y\|^2}{R^2}\right)^{R^2/2} I(\|y\| < R) \tilde{f}(y) dy,$$

where $I(A)$ denotes the indicator function of the (measurable) set A , $A \subset \mathbb{R}^n$.

Therefore, as $R \rightarrow \infty$,

$$B_R^{R^2}[f](t) \rightarrow \int_{\mathbb{R}^n} e^{-i(t,y)} \exp\{-\|y\|^2/2\} dy \tilde{f}(y) dy = [\sqrt{2\pi}]^n [f * f_0](t), \quad (4.2)$$

by virtue of dominated convergence theorem.

If we take $f = f_0$, then in (4.2) the convolution of two Gaussian densities appears. It remains to apply the relation (2.12); we omit some simple calculations. □

5 Convergence of Bochner-Riesz operators.

We investigate here the convergence, as $R \rightarrow \infty$, of the family of Bochner-Riesz approximations $B_R^\alpha[f]$ to the source function f in the Lebesgue-Riesz norm $L_p(\mathbb{R}^n)$, $p \in (1, \infty)$, in addition to the similar results in [21, 11, 26], etc.

For any function $f \in L_p(\mathbb{R}^n)$, its modulus of L_p continuity is defined alike as in approximation theory [1, chapter V] :

$$\omega_p[f](\delta) \stackrel{\text{def}}{=} \sup_{h: |h| \leq \delta} \|T_h[f] - f\|_p, \quad \delta \geq 0,$$

where $T_h[f]$ denotes the shift operator

$$T_h[f](t) = f(t - h), \quad t, h \in \mathbb{R}^n.$$

Obviously, $\omega_p[f](\delta) \leq 2\|f\|_p$ and

$$\lim_{\delta \rightarrow 0+} \omega_p[f](\delta) = 0, \quad f \in L_p. \quad (5.1)$$

Theorem 5.1. *Let $p \in (1, \infty)$, $\alpha \leq \alpha_0 = (n-1)/2$ and $f \in L_p(\mathbb{R}^n)$. Then*

$$\lim_{R \rightarrow \infty} \|B_R^\alpha[f] - f\|_p = 0. \quad (5.2)$$

Proof.

The difference $\Delta_R[f](t) = B_R^\alpha[f] - f$ has the form

$$\Delta_R[f](t) = \int_{\mathbb{R}^n} \left\{ f\left(t - \frac{v}{R}\right) - f(t) \right\} K_\lambda(|v|) dv.$$

We apply now the triangle inequality for the L_p norm in the integral form

$$\begin{aligned} \|\Delta_R[f]\|_p &\leq \int_{\mathbb{R}^n} \|T_{|v|/R}[f] - f\|_p |K_\lambda(|v|)| dv \\ &\leq \int_{\mathbb{R}^n} \omega_p[f](|v|/R) |K_\lambda(|v|)| dv. \end{aligned}$$

Note that, under the above conditions,

$$\int_{\mathbb{R}^n} |K_\lambda(|v|)| dv < \infty,$$

so that (5.2) follows again from the dominated convergence theorem. \square

Remark 5.1. As a slight consequence, under the above conditions, if $f \in L_p(\mathbb{R}^n)$, then

$$\|B_R^\alpha[f]\|_p \leq \|f\|_p + \int_{\mathbb{R}^n} \omega_p[f](|v|/R) |K_\lambda(|v|)| dv$$

and, consequently,

$$\forall f \in L_p(\mathbb{R}^n) \Rightarrow \sup_{R \geq 1} \|B_R^\alpha[f]\|_p \leq 3\|f\|_p.$$

The case $p = \infty$ requires a separate consideration. Introduce the Banach space $C_0(\mathbb{R}^n)$ as a collection of all bounded and *uniformly continuous* functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$, equipped with the ordinary norm,

$$\|f\|_\infty \stackrel{def}{=} \sup_{t \in \mathbb{R}^n} |f(t)|.$$

As above

$$\omega_\infty[f](\delta) \stackrel{def}{=} \sup_{h: |h| \leq \delta} \|T_h[f] - f\|_\infty, \quad \delta \geq 0.$$

Evidently, $\omega_\infty[f](\delta) \leq 2\|f\|_\infty$ and

$$\lim_{\delta \rightarrow 0+} \omega_\infty[f](\delta) = 0, \quad f \in L_p. \quad (5.3)$$

The assertion of Theorem 5.1 under the same conditions remains true in the case $p = \infty$.

Theorem 5.2. *Under the same conditions of Theorem 5.1, for any function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ from the space $C_0(\mathbb{R}^n)$, its Bochner-Riesz approximation B_R^α converges uniformly to the source function f , that is*

$$\lim_{R \rightarrow \infty} \|B_R^\alpha[f] - f\|_\infty = 0. \quad (5.4)$$

Proof. The proof is the same as in Theorem 5.1 and may be omitted. \square

Remark 5.2. *As before, if $f \in C_0(\mathbb{R}^n)$, then*

$$\|B_R^\alpha[f]\|_\infty \leq \|f\|_\infty + \int_{\mathbb{R}^n} \omega_\infty[f](|v|/R) |K_\lambda(|v|)| \, dv$$

and moreover

$$\forall f \in C_0(\mathbb{R}^n) \Rightarrow \sup_{R \geq 1} \|B_R^\alpha[f]\|_\infty \leq 3\|f\|_\infty.$$

6 Concluding remarks.

A. In our opinion, the method described in this paper may be essentially generalized on more operators of convolutions type, linear or not. See some preliminary results [30].

B. It is interesting to generalize the estimates obtained in the previous Sections to the so-called *maximal* operators associated with the Bochner-Riesz one considered here, in the spirit of the works [16, 25], and so one:

$$\|\sup_{R \geq 1} B_R^\alpha[f]\|_p \leq M(\alpha, n; p) \cdot \|f\|_p, \quad p \geq p_0,$$

or

$$\|\sup_{R \geq 1} B_R^\alpha[f]\|_r \leq M(\alpha, n; p, r) \cdot \|f\|_p, \quad p \geq p_0, \quad r = r(p),$$

in order to obtain the GLS estimate for the Bochner-Riesz *maximal* operator of the form

$$\|\sup_{R \geq 1} B_R^\alpha[f]\|_{G\zeta} \leq L(\alpha, n; \zeta, \psi) \cdot \|f\|_{G\psi}$$

for some generating functions ψ, ζ .

Acknowledgement. The first author has been partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) and by Università degli Studi di Napoli Parthenope through the project “sostegno alla Ricerca individuale”.

References

- [1] N.I. Achiezer. *Theory of approximation*, F. Ungar, (1956), (Translated from Russian).
- [2] I. Ahmed, A. Fiorenza, M.R. Formica, A. Gogatishvili and J.M. Rakotoson. *Some new results related to Lorentz G -Gamma spaces and interpolation*. J. Math. Anal. Appl., 483, **2**, 123623 (2020).
- [3] W. Beckner, *Inequalities in Fourier analysis*, Ann. of Math. **102** (1975), 159–182.
- [4] H.J. Brascamp and E.H. Lieb, *Best constants in Young's inequality, its converse, and its generalization to more than three functions*, Advances in Math. **20** (2) (1976), 151–173.
- [5] V.V. Buldygin and E.V. Ilarionov, *A problem in the statistics of random fields*. (Russian) Probabilistic infinite-dimensional analysis, pp. 6–14, 123, Akad. Nauk Ukrain. SSR, Inst. Mat., Kiev, 1981.
- [6] C. Capone, M.R. Formica and R. Giova, *Grand Lebesgue spaces with respect to measurable functions*, Nonlinear Anal. **85** (2013), 125–131.
- [7] A. Córdoba, *A note on Bochner-Riesz operators*, Duke Math. J. **46** (1979), no. 3, 505–511.
- [8] J. Duoandikoetxea, *Fourier analysis*, Graduate Studies in Math., vol. 29. American Mathematical Society, Providence, RI, 2001. Translated and revised from the 1995 Spanish original by David Cruz-Uribe.
- [9] S. V. Ermakov, and E. I. Ostrovsky, *Continuity Conditions, Exponential Estimates, and the Central Limit Theorem for Random Fields*, Moscow, VINITI, 1986 (in Russian).
- [10] C. Fefferman, *Inequalities for strongly singular convolution operators*, Acta Math. **124** (1970), 9–36.
- [11] C. Fefferman, *A note on spherical summation multipliers*, Israel J. Math., **15**, (1973), 44–52.
- [12] A. Fiorenza, *Duality and reflexivity in grand Lebesgue spaces*, Collect. Math. **51** (2) (2000), 131–148.
- [13] A. Fiorenza, M.R. Formica and A. Gogatishvili, *On grand and small Lebesgue and Sobolev spaces and some applications to PDE's*. Differ. Equ. Appl., **10** (1), (2018), 21–46.
- [14] A. Fiorenza, M.R. Formica, A. Gogatishvili, T. Kopaliani and J.M. Rakotoson, *Characterization of interpolation between grand, small or classical Lebesgue spaces*, Nonlinear Anal. **177** (2018), 422–453.
- [15] A. Fiorenza, M.R. Formica and J.M. Rakotoson, *Pointwise estimates for GT -functions and applications*, Differential Integral Equations **30** (11-12) (2017), 809–824.
- [16] A. Fiorenza, B. Gupta and P. Jain, *The maximal theorem for weighted grand Lebesgue spaces*, Studia Math. **188**, no.2, (2008), 123–133.
- [17] A. Fiorenza and G.E. Karadzhov, *Grand and small Lebesgue spaces and their analogs*, Z. Anal. Anwend. **23** (4) (2004), 657–681.
- [18] M.R. Formica and R. Giova, *Boyd indices in generalized grand Lebesgue spaces and applications*, Mediterr. J. Math., **12** (3) (2015), 987–995.
- [19] M.R. Formica, Y.V. Kozachenko, E. Ostrovsky, L. Sirota, *Exponential tail estimates in the law of ordinary logarithm (LOL) for triangular arrays of random variables*. Lith. Math. J. (2020), DOI 10.1007/s10986-020-09481-x.

- [20] M.R. Formica, E. Ostrovsky, L. Sirota, *Grand Lebesgue Spaces are really Banach algebras relative to the convolution on unimodular locally compact groups equipped with Haar measure*, to appear on Math. Nachr.
- [21] L. Grafakos, *Modern Fourier analysis*, Third edition. Graduate Texts in Mathematics, 250. Springer, New York, 2014.
- [22] L. Greco, T. Iwaniec and C. Sbordone, *Inverting the p -harmonic operator*, Manuscripta Math. **92**, no. 2, (1997), 249–258.
- [23] T. Iwaniec and C. Sbordone, *On the integrability of the Jacobian under minimal hypotheses*, Arch. Rational Mech. Anal. **119**, no. 2, (1992), 129–143.
- [24] T. Iwaniec, P. Koskela and J. Onninen, *Mappings of finite distortion: monotonicity and continuity*, Invent. Math. **144**, no. 3, (2001), 507–531.
- [25] P. Jain and S. Kumari, *On grand Lorentz spaces and the maximal operator*. Georgian Math. J. **19**, no. 2, (2012), 235–246.
- [26] D.N. Karasev and V.A. Nogin, *$L_p - L_q$ estimates for the Bochner-Riesz operator of complex order*, Z. Anal. Anwendungen **21**, no. 4, (2002), 915–929.
- [27] Yu.V. Kozachenko and E.I. Ostrovsky, *Banach Spaces of random variables of sub-Gaussian type*, Teor. Veroyatn. Mat. Stat., Kiev, **32** (134), (1985), 42–53, (in Russian). English transl.: Theory Probab. Math. Stat., **32**, (1986), 45–56.
- [28] Yu.V. Kozachenko, Yu.Yu. Mlavets, and N.V. Yurchenko, *Weak convergence of stochastic processes from spaces $F_\psi(\Omega)$* , Statistics, Optimization & Information Computing, **6** (2), (2018), 266–277.
- [29] Yu.V. Kozachenko, E. Ostrovsky and L. Sirota, *Relations between exponential tails, moments and moment generating functions for random variables and vectors*.
arXiv:1701.01901v1 [math.FA] 8 Jan 2017
- [30] E. Liflyand, E. Ostrovsky and L. Sirota, *Structural properties of bilateral grand Lebesgue spaces*, Turkish J. Math. **34**, no. 2, (2010), 207–219.
- [31] S. Lu and D. Yan, *Bochner-Riesz means on Euclidean spaces*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2013, 376 pp.
- [32] E. Ostrovsky, *Exponential Orlicz's spaces: new norms and applications*.
arXiv:FA/0406534, v.1, (25.06.2004.)
- [33] E. Ostrovsky and L. Sirota, *Some new rearrangement invariant spaces: theory and applications*.
arXiv:math.FA/0605732 v1, 29, (May 2006).
- [34] E. Ostrovsky and L. Sirota, *Fourier Transforms in Exponential Rearrangement Invariant Spaces*.
arXiv:math.FA/040639, v1, (20.6.2004.)
- [35] E. Ostrovsky and L. Sirota, *Moment Banach spaces: Theory and applications*, HIAT Journal of Science and Engineering C, 4(1–2), (2007), 233–262.
- [36] E. Ostrovsky and L. Sirota, *Boundedness of operators in bilateral Grand Lebesgue Spaces, with exact and weakly exact constant calculation*. arXiv:1104.2963v1 [math.FA] 15 Apr 2011.
- [37] E. Ostrovsky, S.Yu Tsykunova, *Asymptotic properties of the distribution of the maximum of a Gaussian nonstationary process that arises in covariance statistics*, (Russian) Teor. Veroyatnost. i Primenen. **39** (1994), no. 3, 641–649; translation in Theory Probab. Appl. **39** (1994), no. 3, 527–534 (1995).
- [38] E.M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, no. 30 Princeton University Press, Princeton, N.J. 1970.
- [39] E.M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. With the assistance of Timothy S. Murphy. Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.