

# Change-point tests for the tail parameter of Long Memory Stochastic Volatility time series

Annika Betken\*

Faculty of Mathematics, Ruhr-Universität Bochum  
and

Davide Giraudo\*

Faculty of Mathematics, Ruhr-Universität Bochum  
and

Rafał Kulik

Department of Mathematics and Statistics, University of Ottawa

January 28, 2022

## Abstract

We consider a change-point test based on the Hill estimator to test for structural changes in the tail index of Long Memory Stochastic Volatility time series. In order to determine the asymptotic distribution of the corresponding test statistic, we prove a uniform reduction principle for the tail empirical process in a two-parameter Skorohod space. It is shown that such a process displays a dichotomous behavior according to an interplay between the Hurst parameter, i.e., a parameter characterizing the dependence in the data, and the tail index. Our theoretical results are accompanied by simulation studies and the analysis of financial time series with regard to structural changes in the tail index.

*Keywords:* stochastic volatility; long-range dependence; change-point tests; tail empirical process; heavy tails; chaining

---

\*Research supported by the German National Academic Foundation and Collaborative Research Center SFB 823 *Statistical modelling of nonlinear dynamic processes*

# 1 Introduction and motivation

The tail behavior of the marginal distribution of time series is of major relevance for statistics in applied sciences such as econometrics and hydrology, where heavy-tailed data occurs frequently. More precisely, time series from finance such as the log returns of exchange rates and stock market indices display heavy tails; see [Mandelbrot \(1963\)](#). Furthermore, drastic events like the financial crisis in 2008 substantiate the importance of studying time series models that underlie financial data. Against this background, the identification of changes in the tail behavior of data-generating stochastic processes, that result in an increase or decrease in the probability of extreme events, is of utmost interest. In particular, the analysis of the tail behavior of financial data may pave the way for a corresponding adjustment of risk management for capital investments and, therefore, prevent from huge capital losses. Indeed, there is empirical evidence that the tail behavior of financial time series may change over time: [Quintos et al. \(2001\)](#) identify changes in the tail of Asian stock market indices, [Galbraith and Zernov \(2004\)](#) find evidence for changes in the tail behavior of returns on U.S. equities, and [Werner and Upper \(2004\)](#) detect structural breaks in high-frequency data of Bund future returns.

## 1.1 Tail index estimation and change-point problem

Let  $X_j$ ,  $j \in \mathbb{N}$ , be a stationary time series whose marginal tail distribution function  $\bar{F}$  is regularly varying with index  $-\alpha$ ,  $\alpha > 0$ , i.e.,  $\mathbb{P}(X > x) = x^{-\alpha}L(x)$ , where  $L$  is slowly varying at infinity. Since the tail behavior of  $X_j$ ,  $j \in \mathbb{N}$ , is primarily determined by the value of the tail index  $\alpha$ , identifying a change in the tail of data-generating processes corresponds to testing for a change-point in this parameter.

In particular, this means that, given a set of observations  $X_1, \dots, X_n$  with  $\mathbb{P}(X_j > x) = x^{-\alpha_j}L(x)$ ,  $j = 1, \dots, n$ , we aim at deciding on the testing problem  $(H, A)$  with

$$H : \alpha_1 = \dots = \alpha_n$$

and

$$A : \alpha_1 = \dots = \alpha_k \neq \alpha_{k+1} = \dots = \alpha_n \\ \text{for some } k \in \{1, \dots, n-1\}.$$

Test statistics that are designed for identifying structural changes in the tail index are naturally derived from an estimation of the tail index  $\alpha$ . For some general results on tail index estimation see [Drees \(1998a\)](#) and [Drees \(1998b\)](#). In this article, we focus on two estimators that are motivated by the fact that for a random variable  $X$  with tail index  $\alpha$

$$\lim_{u \rightarrow \infty} \mathbb{E} \left[ \log \left( \frac{X}{u} \right) \mid X > u \right] = \lim_{u \rightarrow \infty} \frac{\mathbb{E} \left[ \log \left( \frac{X}{u} \right) \mathbf{1} \{X > u\} \right]}{\mathbb{P}(X > u)} = \frac{1}{\alpha} =: \gamma.$$

When we are given a set of observations  $X_1, \dots, X_n$ , an approximation of the unknown distribution of  $X$  by its empirical analogue gives the following estimator for the tail index:

$$\hat{\gamma} := \frac{1}{\sum_{j=1}^n \mathbf{1} \{X_j > u_n\}} \sum_{j=1}^n \log \left( \frac{X_j}{u_n} \right) \mathbf{1} \{X_j > u_n\}, \quad (1)$$

where  $u_n$ ,  $n \in \mathbb{N}$ , is a sequence with  $u_n \rightarrow \infty$  and  $n\bar{F}(u_n) \rightarrow \infty$ . Replacing the deterministic levels  $u_n$  in the formula for  $\hat{\gamma}$  by  $X_{n:n-k_n}$  for some  $k_n$ ,  $1 \leq k_n \leq n-1$ , where  $X_{n:n} \geq X_{n:n-1} \geq \dots \geq X_{n:1}$  are the order statistics of the sample  $X_1, \dots, X_n$ , yields the Hill estimator

$$\hat{\gamma}_{\text{Hill}} = \frac{1}{k_n} \sum_{i=1}^{k_n} \log \left( \frac{X_{n:n-i+1}}{X_{n:n-k_n}} \right).$$

As the most popular estimator for the tail index, established in [Hill \(1975\)](#), the Hill estimator has been widely studied in the literature. Its limiting distribution was obtained under various model assumptions, including linear processes ([Resnick and Stărică \(1997\)](#)),  $\beta$ -mixing processes ([Drees \(2000\)](#)), and Long Memory Stochastic Volatility models ([Kulik and Soulier \(2011\)](#)). The first article that establishes a theory for change-point tests that are based on the Hill estimator seems to be [Quintos et al. \(2001\)](#). While [Quintos et al. \(2001\)](#) consider independent, identically distributed observations, ARCH- and GARCH-type processes, [Kim and Lee \(2011\)](#) and [Kim and Lee \(2012\)](#) extend their results to  $\beta$ -mixing processes and residual-based change-point tests for AR( $p$ ) processes with heavy-tailed innovations. In contrast, we study change-point tests for the tail index of Long Memory Stochastic Volatility time series based on the two estimators  $\hat{\gamma}$  and  $\hat{\gamma}_{\text{Hill}}$ . In fact, our results are the first to consider the change-point problem for stochastic volatility models and time series with long-range dependence.

To motivate the design of test statistics for deciding on the change-point problem  $(H, A)$ , we temporarily assume that the change-point location is known, i.e., for a given  $k \in \{1, \dots, n-1\}$  we consider the testing problem  $(H, A_k)$  with

$$A_k : \alpha_1 = \dots = \alpha_k \neq \alpha_{k+1} = \dots = \alpha_n.$$

For this testing problem, change-point tests have been considered in [Phillips et al. \(1990\)](#) and [Koedijk et al. \(1990\)](#). In order to decide on  $(H, A_k)$ , we compare an estimation  $\hat{\gamma}_k$  of the tail index based on the observations  $X_1, \dots, X_k$  to an estimation  $\hat{\gamma}_n$  of the tail index based on the whole sample  $X_1, \dots, X_n$ . This idea leads to studying the following test statistic

$$\Gamma_{k,n} = \frac{k}{n} \left| \frac{\hat{\gamma}_k}{\hat{\gamma}_n} - 1 \right|.$$

Under the assumption that the change-point location is unknown under the alternative, it seems natural to consider the statistic  $\Gamma_{k,n}$  for every potential change-point location  $k$  and to decide in favor of the alternative hypothesis  $A$  if the maximum of its values exceeds a predefined threshold. As a result, a change-point test for the testing problem  $(H, A)$  that rests upon the estimator  $\hat{\gamma}$  defined by (1) bases test decisions on the values of the statistic

$$\Gamma_n := \sup_{t \in [t_0, 1]} t \left| \frac{\hat{\gamma}_{[nt]}}{\hat{\gamma}_n} - 1 \right| \quad (2)$$

with  $t_0 \in (0, 1)$  and with the sequential version of  $\hat{\gamma}$  defined by

$$\hat{\gamma}_{[nt]} := \frac{1}{\sum_{j=1}^{[nt]} \mathbf{1}\{X_j > u_n\}} \sum_{j=1}^{[nt]} \log \left( \frac{X_j}{u_n} \right) \mathbf{1}\{X_j > u_n\}. \quad (3)$$

Likewise, a test statistic based on the Hill estimator is given by

$$\tilde{\Gamma}_n := \sup_{t \in [t_0, 1]} t \left| \frac{\hat{\gamma}_{\text{Hill}}(t)}{\hat{\gamma}_{\text{Hill}}(1)} - 1 \right|$$

with the sequential version of  $\hat{\gamma}_{\text{Hill}}$  defined by

$$\hat{\gamma}_{\text{Hill}}(t) := \frac{1}{[k_n t]} \sum_{i=1}^{[k_n t]} \log \left( \frac{X_{[nt]:[nt]-i+1}}{X_{[nt]:[nt]-k_{[nt]}}} \right).$$

In this context, the most comprehensive theory for change-point tests is presented in [Hoga \(2017\)](#). The author considers a number of test statistics based on different tail index estimators and derives their asymptotic distributions under the assumption of  $\beta$ -mixing data generating processes.

In the following, we derive the asymptotic distribution of both estimators, i.e.,  $\hat{\gamma}_{[nt]}$  and  $\hat{\gamma}_{\text{Hill}}(t)$ , and the corresponding tests statistics, i.e.,  $\Gamma_n$  and  $\tilde{\Gamma}_n$ , under the hypothesis of stationary time series data. For this purpose, we first prove a limit theorem for the tail empirical process of Long Memory Stochastic Volatility time series in two parameters. This limit theorem does not necessarily relate to the change-point context. It

can therefore be considered of independent interest and, thus, as the main theoretical result of our work. Our theoretical results are accompanied by simulation studies. As an empirical application of our tests, we consider Standard & Poor's 500 daily closing index covering the period from January 2008 to December 2008, the year of the financial crisis. We identify a change in the data at exactly one day after Lehman Brothers filed for bankruptcy protection, an event which is thought to have played a major role in the unfolding of the crisis in 2007 - 2008.

## 1.2 Tail empirical process

In order to derive the limit distribution of the tail estimators  $\hat{\gamma}_{[nt]}$  and  $\hat{\gamma}_{\text{Hill}}(t)$ , parametrized in  $t$ , and the corresponding test statistics  $\Gamma_n$  and  $\tilde{\Gamma}_n$ , it is crucial to note that

$$\hat{\gamma}_{[nt]} = \frac{1}{\sum_{j=1}^{[nt]} \mathbf{1}\{X_j > u_n\}} \sum_{j=1}^{[nt]} \log\left(\frac{X_j}{u_n}\right) \mathbf{1}\{X_j > u_n\} = \frac{1}{\tilde{T}_n(1, t)} \int_1^\infty s^{-1} \tilde{T}_n(s, t) ds, \quad (4)$$

where

$$\tilde{T}_n(s, t) = \frac{1}{n\bar{F}(u_n)} \sum_{j=1}^{[nt]} \mathbf{1}\{X_j > u_n s\}.$$

As a result, asymptotics of the considered statistics can be derived from a limit theorem for the two-parameter tail empirical process

$$e_n(s, t) := \left\{ \tilde{T}_n(s, t) - T(s, t) \right\}, \quad s \in [1, \infty], \quad t \in [0, 1], \quad (5)$$

where  $T(s, t)$  does not correspond to the mean of  $\tilde{T}_n(s, t)$ , but rather to the limit of that mean, i.e., to

$$T(s, t) := ts^{-\alpha}. \quad (6)$$

Among others, the tail empirical process in one parameter, i.e.,  $e_n(s, 1)$ ,  $s \in [1, \infty]$ , has previously been studied in [Mason \(1988\)](#), [Einmahl \(1990\)](#), and [Einmahl \(1992\)](#) for independent, identically distributed observations, in [Rootzén \(2009\)](#) for absolutely regular processes, and in [Kulik and Soulier \(2011\)](#) for Long Memory Stochastic Volatility time series. For the latter, the convergence of the two-parameter tail empirical process will be discussed in Section [2.2](#).

### 1.3 Long Memory Stochastic Volatility model

A phenomenon that is often encountered in the context of financial time series corresponds to the observation that the observations seem to be uncorrelated, whereas their absolute values or higher moments tend to be highly correlated. Another characteristic of financial time series is the occurrence of heavy tails. In particular, the distribution of the considered data often exhibits tails that are heavier than those of a normal distribution. The previously described features of financial data can be covered by stochastic volatility models.

#### Stochastic volatility model

The Long Memory Stochastic Volatility model that is taken as a basis of the theoretical results established in this article can be considered as a generalization of stochastic volatility models considered, for example, in [Taylor \(1986\)](#). Initially, this model had been introduced by [Breidt et al. \(1998\)](#) and, independently, by [Harvey \(2002\)](#). An overview of stochastic volatility models with long-range dependence and their basic properties is given in [Deo et al. \(2006\)](#) and in [Hurvich and Soulier \(2009\)](#).

Stochastic volatility time series  $X_j$ ,  $j \in \mathbb{N}$ , are typically defined via

$$X_j = Z_j \varepsilon_j \quad \text{with} \quad Z_j = \exp\left(\frac{1}{2} Y_j\right), \quad (7)$$

where  $\varepsilon_j$ ,  $j \in \mathbb{N}$ , is a sequence of independent, identically distributed random variables with mean 0, and  $Y_j$ ,  $j \in \mathbb{N}$ , is a Gaussian process, independent of  $\varepsilon_j$ ,  $j \in \mathbb{N}$ .

While these models are often restricted to modeling a relatively fast decay of dependence in  $Y_j$ ,  $j \in \mathbb{N}$ , the so-called Long Memory Stochastic Volatility model allows for long-range dependence. In what follows, we will specify a corresponding dependence structure for  $Y_j$ ,  $j \in \mathbb{N}$ .

#### Subordinated Gaussian processes

The rate of decay of the autocovariance function is crucial to the definition of long-range dependence in time series.

**Definition 1.1.** *A (second-order) stationary, real-valued time series  $Y_j$ ,  $j \in \mathbb{Z}$ , is called*

long-range dependent if its autocovariance function  $\gamma$  satisfies

$$\gamma_Y(k) := \text{Cov}(Y_1, Y_{k+1}) \sim k^{-D} L_\gamma(k), \quad \text{as } k \rightarrow \infty,$$

with  $D \in (0, 1)$  for some slowly varying function  $L_\gamma$ . We refer to  $D$  as long-range dependence (LRD) parameter; see [Pipiras and Taqqu \(2017\)](#), p. 17.

We will focus our considerations on long-range dependent subordinated Gaussian time series.

**Definition 1.2.** Let  $Y_j$ ,  $j \in \mathbb{N}$ , be a Gaussian process. A process  $Z_j$ ,  $j \in \mathbb{N}$ , satisfying  $Z_j = G(Y_j)$  for some measurable function  $G: \mathbb{R} \rightarrow \mathbb{R}$  is called subordinated Gaussian process.

**Remark 1.3.** For any particular distribution function  $F$ , an appropriate choice of the transformation  $G$  in Definition 1.2 yields a subordinated Gaussian process with marginal distribution  $F$ . Moreover, there exist algorithms for generating Gaussian processes that, after suitable transformation, yield subordinated Gaussian processes with marginal distribution  $F$  and a predefined covariance structure; see [Pipiras and Taqqu \(2017\)](#), Section 5.8.4. To that effect, subordinated Gaussian processes constitute a comprehensive model for long-range dependent time series.

The subordinated random variables  $Z_j = G(Y_j)$ ,  $j \in \mathbb{N}$ , can be considered as elements of the Hilbert space  $L^2 := L^2(\mathbb{R}, \varphi(x)dx)$ , i.e., the space of all measurable, real-valued functions which are square-integrable with respect to the measure  $\varphi(x)dx$  associated with the standard normal density function  $\varphi$ , equipped with the inner product

$$\langle G_1, G_2 \rangle_{L^2} := \int_{-\infty}^{\infty} G_1(x)G_2(x)\varphi(x)dx = \mathbb{E}[G_1(Y)G_2(Y)],$$

where  $G_1, G_2 \in L^2(\mathbb{R}, \varphi(x)dx)$  and  $Y$  denotes a standard normally distributed random variable. In order to characterize the dependence structure of subordinated Gaussian processes, we consider their expansion in Hermite polynomials.

**Definition 1.4.** For  $n \geq 0$ , the Hermite polynomial of order  $n$  is defined by

$$H_n(x) = (-1)^n e^{\frac{1}{2}x^2} \frac{d^n}{dx^n} e^{-\frac{1}{2}x^2}, \quad x \in \mathbb{R}.$$

The sequence of Hermite polynomials constitutes an orthogonal basis of  $L^2$ . In particular, it holds that

$$\langle H_n, H_m \rangle_{L^2} = \begin{cases} n! & \text{if } n = m, \\ 0 & \text{if } n \neq m. \end{cases}$$

As a result, every  $G \in L^2(\mathbb{R}, \varphi(x)dx)$  has an expansion in Hermite polynomials, i.e., for  $G \in L^2(\mathbb{R}, \varphi(x)dx)$  and  $Y$  standard normally distributed, we have

$$G(Y) \stackrel{L^2}{=} \sum_{r=0}^{\infty} \frac{J_r(G)}{r!} H_r(Y), \text{ i.e., } \lim_{n \rightarrow \infty} \left\| G(Y) - \sum_{r=0}^n \frac{J_r(G)}{r!} H_r(Y) \right\|_{L^2} = 0, \quad (8)$$

where  $\|\cdot\|_{L^2}$  denotes the norm induced by the inner product  $\langle \cdot, \cdot \rangle_{L^2}$ , and the so-called *Hermite coefficients*  $J_r(G)$ ,  $r \geq 1$ , are given by

$$J_r(G) := \langle G, H_r \rangle_{L^2} = \mathbb{E} G(Y) H_r(Y), \quad r \geq 1.$$

Given the Hermite expansion (8), it is possible to characterize the dependence structure of subordinated Gaussian time series  $G(Y_j)$ ,  $j \in \mathbb{N}$ . Under the assumption that the Gaussian sequence  $Y_j$ ,  $j \in \mathbb{N}$ , is stationary and that  $G$  is a one-to-one function, the behavior of the autocorrelations of the transformed process is completely determined by the dependence structure of the underlying process. However, this is not the case in general. In fact, it holds that

$$\text{Cov}(G(Y_1), G(Y_{k+1})) = \sum_{r=1}^{\infty} \frac{J_r^2(G)}{r!} (\gamma_Y(k))^r, \quad (9)$$

where  $\gamma_Y(k)$  denotes the autocovariance function of  $Y_n$ ,  $n \in \mathbb{N}$ ; see [Pipiras and Taqqu \(2017\)](#).

Under the assumption that, as  $k$  tends to  $\infty$ ,  $\gamma_Y(k)$  converges to 0 with a certain rate, the asymptotically dominating term in the series (9) is the summand corresponding to the smallest integer  $r$  for which the Hermite coefficient  $J_r(G)$  is non-zero. This index, which decisively depends on  $G$ , is called Hermite rank.

**Definition 1.5.** *Let  $G \in L^2(\mathbb{R}, \varphi(x)dx)$ ,  $\mathbb{E}[G(Y)] = 0$  for standard normally distributed  $Y$  and let  $J_r(G)$ ,  $r \geq 1$ , be the Hermite coefficients in the Hermite expansion of  $G$ . The smallest index  $k \geq 1$  for which  $J_k(G) \neq 0$  is called the Hermite rank of  $G$ , i.e.,*

$$r := \min \{k \geq 1 : J_k(G) \neq 0\}.$$



It follows from (9) that subordination of long-range dependent Gaussian time series potentially generates time series whose autocovariances decay faster than the autocovariances of the underlying Gaussian process. In some cases, the subordinated time series is long-range dependent as well, in other cases subordination may even yield short-range dependence. Given that  $\text{Cov}(Y_1, Y_{k+1}) \sim k^{-D}L(k)$ , as  $k \rightarrow \infty$ , and given that  $G \in L^2(\mathbb{R}, \varphi(x)dx)$  is a function with Hermite rank  $r$ , we have

$$\text{Cov}(G(Y_1), G(Y_{k+1})) \sim J_r^2(G)r!k^{-Dr}L_\gamma^r(k), \quad \text{as } k \rightarrow \infty.$$

It immediately follows that subordinated Gaussian time series  $G(Y_j)$ ,  $j \in \mathbb{N}$ , are long-range dependent with LRD parameter  $D_G := Dr$  and slowly-varying function  $L_G(k) = J_r^2(G)r!L_\gamma^r(k)$  whenever  $Dr < 1$ .

Given the previous definitions, we specify model assumptions that are taken as a basis for the results in the following sections.

**Definition 1.6.** *Let the data generating process  $X_j$ ,  $j \in \mathbb{N}$ , satisfy*

$$X_j = Z_j \varepsilon_j, \quad j \in \mathbb{N},$$

where  $\varepsilon_j$ ,  $j \in \mathbb{N}$ , is a sequence of independent, identically distributed random variables with mean 0, and  $Z_j$ ,  $j \in \mathbb{N}$ , is a long-range dependent subordinated Gaussian process with  $Z_j = \sigma(Y_j)$ ,  $j \in \mathbb{N}$ , for some stationary, long-range dependent Gaussian process  $Y_j$ ,  $j \in \mathbb{N}$ , with LRD parameter  $D$  and a non-negative measurable function  $\sigma$  (not equal to 0). More precisely, assume that  $Y_j$ ,  $j \in \mathbb{N}$ , admits a linear representation with respect to an independent, standard normally distributed sequence  $\eta_k$ ,  $k \in \mathbb{Z}$ , i.e.,

$$Y_j = \sum_{k=1}^{\infty} c_k \eta_{j-k}, \quad j \in \mathbb{N},$$

with  $\sum_{k=1}^{\infty} c_k^2 = 1$ . Furthermore, suppose that  $(\varepsilon_j, \eta_j)$ ,  $j \in \mathbb{Z}$ , is a sequence of independent, identically distributed random vectors. A sequence of random variables  $X_j$ ,  $j \in \mathbb{N}$ , which satisfies the previous assumption is called a Long Memory Stochastic Volatility (LMSV) time series.

**Remark 1.7.** The model assumptions generalize the preceding concepts of stochastic volatility models with long-range dependence by allowing for general subordinated Gaussian sequences  $Z_j$ ,  $j \in \mathbb{N}$ , and dependence between  $Y_j$ ,  $j \in \mathbb{N}$ , and  $\varepsilon_j$ ,  $j \in \mathbb{N}$ . Instead

of claiming mutual independence of  $Y_j$ ,  $j \in \mathbb{N}$ , and  $\varepsilon_j$ ,  $j \in \mathbb{N}$ , the sequence of random vectors  $(\eta_j, \varepsilon_j)$  is assumed to be independent. In particular, this implies that for a fixed index  $j$ , the random variables  $Y_j$  and  $\varepsilon_j$  are independent, while  $Y_j$  may depend on  $\varepsilon_i$ ,  $i < j$ . In many cases, an LMSV model incorporating this dependence structure is referred to as *LMSV with leverage*, as it allows for so-called *leverage effects* in financial time series. Not taking account of leverage, Definition 1.6 corresponds to the LMSV model considered in Kulik and Soulier (2011), while a similar model with leverage is considered in Bilayi-Biakana et al. (2019).

It can be shown that random variables  $X_j$ ,  $j \in \mathbb{N}$ , satisfying Definition 1.6 are uncorrelated, while their squares inherit the dependence structure from the subordinated Gaussian sequence  $Z_j^2$ ,  $j \in \mathbb{N}$ . Moreover,  $X_j$ ,  $j \in \mathbb{N}$ , inherits the tail behavior from the sequence  $\varepsilon_j$ ,  $j \in \mathbb{N}$ , if the marginal distribution of the random variables  $\varepsilon_j$ ,  $j \in \mathbb{N}$ , has a regularly varying right tail, i.e.,  $\bar{F}_\varepsilon(x) := \mathbb{P}(\varepsilon_1 > x) = x^{-\alpha}L(x)$  for some  $\alpha > 0$  and a slowly varying function  $L$ , and if  $\mathbb{E}[\sigma^{\alpha+\delta}(Y_1)] < \infty$  for some  $\delta > 0$ . More precisely, under these assumptions the following asymptotic equivalence holds:

$$\mathbb{P}(X_1 > x) \sim \mathbb{E}[\sigma^\alpha(Y_1)] \mathbb{P}(\varepsilon_1 > x), \text{ as } x \rightarrow \infty.$$

This result is known as Breiman’s Lemma; see Breiman (1965). On this account, it follows that Definition 1.6 is suited for modeling the previously described characteristic features of financial time series. In all following sections, we will therefore assume that the data-generating process  $X_j$ ,  $j \in \mathbb{N}$ , corresponds to a LMSV time series specified by Definition 1.6.

## 1.4 Organisation of the paper

Equipped with the introductory remarks and definitions, we are in a position to discuss the structure of the paper. In Section 2 we state the technical assumptions that are needed for our theoretical results. These are followed by the main theorem on convergence of the two-parameter tail empirical process (Theorem 2.6). Convergence of estimators of the tail index (Corollaries 2.7 and 2.8) and the test statistics (Corollaries 2.10 and 2.11) are immediate consequences. Simulation studies are presented in Section 3, while real-data analysis can be found in Section 4. All the proofs are included in Section 5. In order to establish convergence of the two-parameter tail empirical process, we decompose it

into a martingale and a long-range dependent part. The latter is dealt with in Section 5.1.2. For the former, we establish finite dimensional convergence (Section 5.1.3) using classical tools from martingale theory, while tightness of the two-parameter martingale part is handled by chaining. This is a theoretical novelty in the present context since the methods used in related papers are not suitable (the method used in Kulik and Soulier (2011) cannot be applied to models with leverage, while the approach in Bilayi-Biakana et al. (2019) is not well-suited for two-parameter processes).

## 2 Main results

### 2.1 Assumptions

In this section, we establish the assumptions guaranteeing convergence of the two-parameter empirical process for LMSV time series. Initially, we specify the LMSV model yielding the main assumptions for the theory:

**Assumption 2.1** (Main Assumptions). *Let  $X_j = Z_j \varepsilon_j$ ,  $j \in \mathbb{N}$ , satisfy Definition 1.6 with  $Z_j = \sigma(Y_j)$ ,  $j \in \mathbb{N}$ , for some stationary, long-range dependent Gaussian process  $Y_j$ ,  $j \in \mathbb{N}$ , with autocovariance function  $\gamma_Y(k) := \text{Cov}(Y_1, Y_{k+1}) \sim k^{-D} L_\gamma(k)$ , as  $k \rightarrow \infty$ , and some independent, identically distributed sequence  $\varepsilon_j$ ,  $j \in \mathbb{N}$ , with regularly varying right tail, i.e.,  $\bar{F}_\varepsilon(x) := \mathbb{P}(\varepsilon_1 > x) = x^{-\alpha} L(x)$  for some  $\alpha > 0$  and a slowly varying function  $L$ . Moreover, let  $r$  denote the Hermite rank of  $\Psi(y) := \sigma^\alpha(y)$  and assume that  $r < 1/D$ .*

In the following, we list some technical conditions that characterize the behavior of the slowly varying function  $L$  and the moments of  $\sigma(Y_1)$ . For this, we introduce another condition on the distribution function  $F_\varepsilon$ .

**Definition 2.2** (Second order regular variation). *Let  $\bar{F}_\varepsilon(x) = x^{-\alpha} L(x)$  for some  $\alpha > 0$  and some slowly varying function  $L$  that is represented by*

$$L(x) = c \exp \left( \int_1^x \frac{\eta(u)}{u} du \right)$$

*for some constant  $c$  and a measurable function  $\eta$ . Furthermore, we assume that there exists a bounded, decreasing function  $\eta^*$  on  $[0, \infty)$ , regularly varying at infinity with pa-*

parameter  $\rho \geq 0$ , i.e.,  $\eta^*(x) = x^{-\rho} L_{\eta^*}(x)$ , such that

$$|\eta(s)| \leq C\eta^*(s),$$

for some constant  $C$  and for all  $s \geq 0$ . We say that  $\bar{F}_\varepsilon$  is second order regularly varying with tail index  $\alpha$  and rate function  $\eta^*$  and we write  $\bar{F}_\varepsilon \in 2RV(\alpha, \eta^*)$ .

Second-order regular variation allows to control the difference between  $\bar{F}_\varepsilon$  and the function  $u \mapsto u^{-\alpha}$ ; see Lemma 6.1 and 6.2 in the Appendix. Moreover, the specific form of  $L$  guarantees continuity of  $\bar{F}_\varepsilon$ .

**Assumption 2.3** (Technical Assumptions). *Suppose the main assumptions hold. Additionally, we assume that*

(TA.1)  $\bar{F}_\varepsilon \in 2RV(\alpha, \eta^*)$  and  $\eta$  is regularly varying with index  $\rho$ ;

(TA.2)  $\eta^*(u_n) = o\left(\frac{d_{n,r}}{n} + \frac{1}{\sqrt{n\bar{F}(u_n)}}\right)$ , where  $d_{n,r}$  is defined by

$$d_{n,r}^2 = \text{Var}\left(\sum_{j=1}^n H_r(Y_j)\right) \sim c_r n^{2-rD} L_\gamma^r(n), \quad c_r = \frac{2r!}{(1-Dr)(2-Dr)}; \quad (10)$$

(TA.3)  $\mathbb{E}[\sigma^{\alpha+\max\{\rho,\alpha\}+\varepsilon}(Y_1)] < \infty$  for some  $\varepsilon > 0$ ;

(TA.4)  $\mathbb{E}[(\sigma(Y_1))^{-1}] < \infty$ .

**Remark 2.4.** Assumption (TA.2) handles the bias which is created by centering the tail empirical process not by its mean, but rather by the limit of that mean.

**Example 2.5.** The most commonly used second order assumption is that

$$L(x) = c \exp\left(\int_1^x \frac{\eta(u)}{u} du\right)$$

with  $\eta(s) = s^{-\alpha\beta}$  for some  $\beta > 0$ . It then holds that  $\bar{F}_\varepsilon(s) = C(s^{-\alpha} + \mathcal{O}(s^{-(\alpha(\beta+1))}))$ , for  $s \rightarrow \infty$ , for some constant  $c > 0$ . Furthermore, we have

$$\sup_{s \geq s_0} \left| \frac{\bar{F}_\varepsilon(u_n s)}{\bar{F}_\varepsilon(u_n)} - s^{-\alpha} \right| = \mathcal{O}(u_n^{-\alpha\beta}).$$

In this case, (TA.2) can be replaced by the assumption  $u_n^{-\alpha\beta} = o\left(\frac{d_{n,r}}{n} + \frac{1}{\sqrt{n\bar{F}(u_n)}}\right)$ .

## 2.2 Convergence of the tail empirical process

Recall that the tail empirical in two parameters is defined by

$$e_n(s, t) := \frac{1}{n\bar{F}(u_n)} \sum_{j=1}^{\lfloor nt \rfloor} \mathbf{1}\{X_j > u_n s\} - ts^{-\alpha}, \quad s \in [1, \infty], \quad t \in [0, 1].$$

The following theorem establishes a characterization of its limit.

**Theorem 2.6.** *Let  $X_j$ ,  $j \in \mathbb{N}$ , be a stationary time series with marginal tail distribution function  $\bar{F}$ . Moreover, assume that Assumptions 2.1 and 2.3 hold.*

1. If  $\frac{n}{d_{n,r}} = o\left(\sqrt{n\bar{F}(u_n)}\right)$ ,

$$\frac{n}{d_{n,r}} e_n(s, t) \Rightarrow \frac{s^{-\alpha}}{\mathbb{E}[\sigma^\alpha(Y_1)]} \frac{J_r(\Psi)}{r!} Z_{r,H}(t), \quad (11)$$

where  $\Psi(y) = \sigma^\alpha(y)$ ,  $r$  is the Hermite rank of  $\Psi$ ,  $Z_{r,H}$  is an  $r$ -th order Hermite process,  $H = 1 - \frac{rD}{2}$ , and  $d_{n,r}^2$  is defined in (10).

2. If  $\sqrt{n\bar{F}(u_n)} = o\left(\frac{n}{d_{n,r}}\right)$ ,

$$\sqrt{n\bar{F}(u_n)} e_n(s, t) \Rightarrow B(s^{-\alpha}, t), \quad (12)$$

where  $B$  denotes a standard Brownian sheet.

The convergence holds in a two-parameter Skorohod space, i.e.,  $\Rightarrow$  denotes weak convergence in  $D([1, \infty] \times [0, 1])$ .

The dichotomy of the limiting process is explained by the decomposition of the tail empirical process into the sum of a martingale and a partial sum of long-range dependent random variables, which can be viewed as a special case of Doob's decomposition; see Section 5.1.1. If  $\frac{n}{d_{n,r}} = o\left(\sqrt{n\bar{F}(u_n)}\right)$ , the martingale part in the decomposition becomes negligible, such that the limiting process arises from the convergence of the long-range dependent part. If  $\sqrt{n\bar{F}(u_n)} = o\left(\frac{n}{d_{n,r}}\right)$ , the long-range dependent part in the decomposition becomes negligible, such that the limiting process arises from the convergence of the martingale part. The same decomposition has already been employed in Kulik and Soulier (2011), Betken and Kulik (2019), and Bilayi-Biakana et al. (2019).

## 2.3 Convergence of the tail estimators

Recall that the considered tail index estimators are defined by

$$\widehat{\gamma}_{[nt]} := \frac{1}{\sum_{j=1}^{[nt]} \mathbf{1}\{X_j > u_n\}} \sum_{j=1}^{[nt]} \log \left( \frac{X_j}{u_n} \right) \mathbf{1}\{X_j > u_n\}$$

and

$$\widehat{\gamma}_{\text{Hill}}(t) := \frac{1}{[k_n t]} \sum_{i=1}^{[k_n t]} \log \left( \frac{X_{[nt]:[nt]-i+1}}{X_{[nt]:[nt]-k_{[nt]}}} \right).$$

Based on 2.6 the limiting distributions of  $\widehat{\gamma}_{[nt]}$  and  $\widehat{\gamma}_{\text{Hill}}(t)$  can be established in  $D[t_0, 1]$  for any  $t_0 \in (0, 1)$ .

**Corollary 2.7.** *Let  $X_j$ ,  $j \in \mathbb{N}$ , be a stationary time series with marginal tail distribution function  $\bar{F}$ . Moreover, assume that Assumptions 2.1 and 2.3 hold.*

1. If  $\frac{n}{d_{n,r}} = o\left(\sqrt{n\bar{F}(u_n)}\right)$ , then

$$\frac{n}{d_{n,r}} t (\widehat{\gamma}_{[nt]} - \gamma) \Rightarrow 0$$

in  $D[t_0, 1]$  for all  $t_0 \in (0, 1)$ .

2. If  $\sqrt{n\bar{F}(u_n)} = o\left(\frac{n}{d_{n,r}}\right)$ , then

$$\sqrt{n\bar{F}(u_n)} t (\widehat{\gamma}_{[nt]} - \gamma) \Rightarrow \int_1^\infty s^{-1} B(s^{-\alpha}, t) ds - \alpha^{-1} B(1, t) \quad (13)$$

in  $D[t_0, 1]$  for all  $t_0 \in (0, 1)$ .

**Corollary 2.8.** *Let  $X_j$ ,  $j \in \mathbb{N}$ , be a stationary time series with marginal tail distribution function  $\bar{F}$ . Moreover, assume that Assumptions 2.1 and 2.3 hold.*

1. If  $\frac{n}{d_{n,r}} = o\left(\sqrt{n\bar{F}(u_n)}\right)$ , then

$$\frac{n}{d_{n,r}} t (\widehat{\gamma}_{\text{Hill}}(t) - \gamma) \Rightarrow 0$$

in  $D[t_0, 1]$  for all  $t_0 \in (0, 1)$ .

2. If  $\sqrt{n\bar{F}(u_n)} = o\left(\frac{n}{d_{n,r}}\right)$ , then

$$\sqrt{n\bar{F}(u_n)} t (\widehat{\gamma}_{\text{Hill}}(t) - \gamma) \Rightarrow \int_1^\infty s^{-1} B(s^{-\alpha}, t) ds - \alpha^{-1} B(1, t) \quad (14)$$

in  $D[t_0, 1]$  for all  $t_0 \in (0, 1)$ .

**Remark 2.9.** 1. Following [Kulik and Soulier \(2011\)](#), we conjecture that the proper scaling in the first case is  $a_n = \sqrt{n\bar{F}(u_n)}$ , as well, yielding the same limit as in the second case. However, within the scope of this article, we will not consider the corresponding argument in detail.

2. The limit in (13) and (14) corresponds to  $\gamma B(t)$ ,  $t \in [0, 1]$ , where  $B$  is a standard Brownian motion.

## 2.4 Asymptotic distribution of the test statistics

Recall that the considered test statistics for the change-point problem  $(H, A)$  are defined by

$$\Gamma_n := \sup_{t \in [t_0, 1]} t \left| \frac{\widehat{\gamma}_{[nt]}}{\widehat{\gamma}_n} - 1 \right|$$

and

$$\widetilde{\Gamma}_n := \sup_{t \in [t_0, 1]} t \left| \frac{\widehat{\gamma}_{\text{Hill}}(t)}{\widehat{\gamma}_{\text{Hill}}(1)} - 1 \right|.$$

Using the convergence obtained in Corollaries 2.7 and 2.8, we derive the asymptotic distribution of the test statistics.

**Corollary 2.10.** *Let  $X_j$ ,  $j \in \mathbb{N}$ , be a stationary time series with marginal tail distribution function  $\bar{F}$ . Moreover, assume that Assumptions 2.1 and 2.3 hold. If  $\sqrt{n\bar{F}(u_n)} = o\left(\frac{n}{d_{n,r}}\right)$ , then, for all  $t_0 \in (0, 1)$ ,*

$$\sqrt{n\bar{F}(u_n)} \sup_{t \in [t_0, 1]} t \left| \frac{\widehat{\gamma}_{[nt]}}{\widehat{\gamma}_n} - 1 \right| \Rightarrow \sup_{t \in [t_0, 1]} |B(t) - tB(1)|,$$

where  $B(t)$ ,  $t \in [0, 1]$ , denotes a standard Brownian motion.

**Corollary 2.11.** *Let  $X_j$ ,  $j \in \mathbb{N}$ , be a stationary time series with marginal tail distribution function  $\bar{F}$ . Moreover, assume that Assumptions 2.1 and 2.3 hold. If  $\sqrt{n\bar{F}(u_n)} = o\left(\frac{n}{d_{n,r}}\right)$ , then, for all  $t_0 \in (0, 1)$ ,*

$$\sqrt{n\bar{F}(u_n)} \sup_{t \in [t_0, 1]} t \left| \frac{\widehat{\gamma}_{\text{Hill}}(t)}{\widehat{\gamma}_{\text{Hill}}(1)} - 1 \right| \Rightarrow \sup_{t \in [t_0, 1]} |B(t) - tB(1)|,$$

where  $B(t)$ ,  $t \in [0, 1]$ , denotes a standard Brownian motion.

### 3 Simulations

For all simulations, the following specifications are made:

$$X_j = \sigma(Y_j)\varepsilon_j, \quad j \geq 1, \quad (15)$$

where

- $\varepsilon_j, j \geq 1$ , is an independent, identically distributed sequence of Pareto distributed random variables generated by the function `rgpd` (`fExtremes` package in R);
- $Y_j, j \geq 1$ , is a fractional Gaussian noise sequence generated by the function `simFGNO` (`longmemo` package in R) with Hurst parameter  $H$ ;
- $\sigma(y) = \exp(y)$ .

Under the alternative, we insert a change of height  $h$  at location  $k = \lfloor n\tau \rfloor$  by simulating independent, identically Pareto distributed observations  $\varepsilon_j, j \geq 1$ , with  $\varepsilon_j, j = 1, \dots, k$ , having tail index  $\alpha_1 = \dots = \alpha_k = \alpha$  and  $\varepsilon_j, j = k + 1, \dots, n$ , having tail index  $\alpha_{k+1} = \dots = \alpha_n = \alpha + h$ .

We base test decisions on the statistic  $\tilde{\Gamma}_n := \max_{1 \leq k \leq n-1} \Gamma_{k,n}$ , where

$$\tilde{\Gamma}_{k,n} = \frac{k}{n} \left| \frac{\hat{\gamma}_{\text{Hill}}\left(\frac{k}{n}\right)}{\hat{\gamma}_{\text{Hill}}(1)} - 1 \right| \quad \text{with} \quad \hat{\gamma}_{\text{Hill}}(t) = \frac{1}{\lfloor k_n t \rfloor} \sum_{i=1}^{\lfloor k_n t \rfloor} \log \left( \frac{X_{\lfloor nt \rfloor : \lfloor nt \rfloor - i + 1}}{X_{\lfloor nt \rfloor : \lfloor nt \rfloor - \lfloor k_n t \rfloor}} \right), \quad (16)$$

and we choose a significance level of 5%.

For the computation of the test statistic, the choice of  $k_n$ , i.e., the number of largest observations that contribute to the estimation of the tail index, is considered a delicate issue. In fact, it has been shown in [Hall \(1982\)](#) that the optimal choice of  $k_n$  depends on the tail behavior of the data-generating process. Due to this circularity, [DuMouchel \(1983\)](#) suggests to chose  $k_n$  proportional to the sample size. As noted in [Quintos et al. \(2001\)](#), a corresponding choice of  $k_n$  has been shown to perform well in simulations and is widely used by practitioners. Hence, we choose  $k_n = \lfloor np \rfloor$ , i.e.,  $p$  defines the proportion of the data that the estimation of the tail index is based on.

The power of the testing procedures is analyzed by considering different choices for the height of the level shift, denoted by  $h$ , and the location of the change-point, denoted by  $\tau$ . In the tables, the columns that are superscribed by  $h = 0$  correspond to the frequency of a type 1 error, i.e., the rejection rate under the hypothesis.



Considering all simulation results, the first thing to note is that these concur with the expected behavior of change-point tests: An increasing sample size goes along with an improvement of the finite sample performance, i.e., the empirical size approaches the level of significance and the empirical power increases, the empirical power of the testing procedures increases when the height of the level shift increases, and the empirical power is higher for breakpoints located in the middle of the sample than for change-point locations that lie close to the boundary of the testing region.

Both Hurst parameter and tail index, seem to have a significant effect on the rejection rates of the change-point test. An increase in dependence, i.e., an increase of the Hurst parameter  $H$ , leads to an increase in the number of rejections. On the one hand, this leads to an increase of power, on the other hand, it results in a larger deviation of the empirical size from the significance level. An increase of tail thickness, i.e., a decrease of the tail parameter  $\alpha$ , however, results in an improvement of the test's performance in that the empirical power increases while the empirical size draws closer to the level of significance. Moreover, the empirical power of the test is higher for changes to heavier tails, i.e., the test tends to detect changes with a negative change-point height  $h$  better.

Technically speaking, the particular case of a change with height  $h = -1$  from  $\alpha = 1$  to  $\alpha = 0$  does not fall under our model assumptions. For a change after a proportion  $\tau = 0.5$  of the data, the empirical power is extremely low in this case. However, for an early change, i.e., for  $\tau = 0.25$ , the empirical power is comparatively high.

		$\alpha = 2.5$					$\alpha = 2$					$\alpha = 1$					
$p$	$n$	$h = 0$	$h = 0.5$	$h = 1$	$h = -0.5$	$h = -1$	$h = 0$	$h = 0.5$	$h = 1$	$h = -0.5$	$h = -1$	$h = 0$	$h = 0.5$	$h = 1$	$h = -0.5$	$h = -1$	
$H = 0.6$	0.1	300	0.088	0.088	0.086	0.109	0.192	0.097	0.086	0.086	0.145	0.418	0.100	0.110	0.128	0.641	0.056
		500	0.078	0.069	0.065	0.105	0.249	0.078	0.083	0.069	0.148	0.602	0.092	0.129	0.141	0.842	0.053
		1000	0.071	0.063	0.059	0.106	0.391	0.062	0.072	0.073	0.195	0.853	0.077	0.188	0.222	0.979	0.054
	0.2	300	0.071	0.065	0.058	0.078	0.176	0.072	0.073	0.071	0.112	0.485	0.076	0.125	0.178	0.816	0.095
		500	0.049	0.059	0.059	0.076	0.227	0.060	0.061	0.071	0.123	0.687	0.067	0.187	0.249	0.951	0.133
		1000	0.044	0.050	0.055	0.086	0.387	0.047	0.053	0.075	0.185	0.911	0.062	0.308	0.428	0.999	0.235
$H = 0.7$	0.1	300	0.112	0.103	0.096	0.137	0.217	0.114	0.100	0.102	0.159	0.443	0.106	0.130	0.131	0.642	0.055
		500	0.093	0.086	0.087	0.123	0.262	0.096	0.082	0.091	0.166	0.626	0.095	0.131	0.148	0.835	0.052
		1000	0.084	0.069	0.070	0.118	0.385	0.077	0.077	0.086	0.213	0.866	0.084	0.199	0.239	0.979	0.062
	0.2	300	0.087	0.083	0.083	0.105	0.196	0.080	0.085	0.090	0.131	0.502	0.084	0.134	0.184	0.824	0.092
		500	0.075	0.080	0.071	0.099	0.256	0.073	0.073	0.081	0.147	0.702	0.075	0.196	0.263	0.959	0.122
		1000	0.068	0.063	0.067	0.109	0.408	0.068	0.077	0.090	0.211	0.919	0.066	0.317	0.458	0.999	0.235
$H = 0.8$	0.1	300	0.140	0.125	0.124	0.149	0.248	0.130	0.135	0.124	0.186	0.477	0.116	0.132	0.148	0.637	0.053
		500	0.131	0.122	0.113	0.156	0.313	0.108	0.119	0.112	0.197	0.662	0.101	0.146	0.171	0.842	0.053
		1000	0.108	0.109	0.108	0.167	0.446	0.107	0.113	0.115	0.254	0.881	0.090	0.217	0.264	0.984	0.058
	0.2	300	0.118	0.110	0.116	0.135	0.250	0.107	0.112	0.118	0.174	0.560	0.092	0.171	0.217	0.837	0.078
		500	0.122	0.111	0.103	0.157	0.319	0.101	0.109	0.118	0.192	0.743	0.079	0.209	0.299	0.957	0.123
		1000	0.098	0.107	0.113	0.165	0.491	0.100	0.117	0.142	0.269	0.935	0.076	0.360	0.493	0.999	0.215
$H = 0.9$	0.1	300	0.175	0.164	0.165	0.192	0.308	0.166	0.151	0.162	0.215	0.530	0.120	0.152	0.167	0.650	0.059
		500	0.167	0.165	0.164	0.201	0.395	0.152	0.151	0.159	0.244	0.715	0.105	0.166	0.198	0.834	0.053
		1000	0.175	0.171	0.176	0.247	0.554	0.156	0.166	0.185	0.322	0.924	0.104	0.239	0.289	0.982	0.056
	0.2	300	0.169	0.158	0.168	0.194	0.341	0.140	0.162	0.161	0.215	0.646	0.102	0.200	0.268	0.846	0.063
		500	0.177	0.183	0.171	0.213	0.458	0.153	0.158	0.180	0.275	0.821	0.101	0.262	0.373	0.964	0.079
		1000	0.207	0.203	0.215	0.281	0.625	0.175	0.192	0.230	0.372	0.966	0.100	0.414	0.557	0.999	0.154

Table 1: Rejection rates of the change-point test based on the statistic  $\Gamma_n$ ,  $k_n = \lfloor np \rfloor$ , for LMSV time series (Pareto distributed  $\varepsilon_j$ ,  $j \geq 1$ ) of length  $n$  with Hurst parameter  $H$ , tail index  $\alpha$  and a shift in the mean of height  $h$  after a proportion  $\tau = 0.5$ . The calculations are based on 5,000 simulation runs.

		$\alpha = 2.5$					$\alpha = 2$					$\alpha = 1$					
$p$	$n$	$h = 0$	$h = 0.5$	$h = 1$	$h = -0.5$	$h = -1$	$h = 0$	$h = 0.5$	$h = 1$	$h = -0.5$	$h = -1$	$h = 0$	$h = 0.5$	$h = 1$	$h = -0.5$	$h = -1$	
$H = 0.6$	0.1	300	0.088	0.086	0.085	0.104	0.127	0.097	0.099	0.092	0.105	0.145	0.100	0.148	0.198	0.183	0.069
		500	0.078	0.071	0.071	0.083	0.129	0.078	0.072	0.075	0.105	0.203	0.092	0.155	0.238	0.254	0.078
		1000	0.071	0.058	0.060	0.076	0.151	0.062	0.061	0.075	0.106	0.373	0.077	0.216	0.376	0.594	0.137
	0.2	300	0.071	0.069	0.068	0.075	0.099	0.072	0.074	0.073	0.089	0.156	0.076	0.149	0.230	0.272	0.658
		500	0.049	0.052	0.059	0.063	0.120	0.060	0.062	0.070	0.084	0.262	0.067	0.185	0.328	0.532	0.891
		1000	0.044	0.050	0.052	0.056	0.160	0.047	0.055	0.072	0.096	0.521	0.062	0.295	0.550	0.912	0.997
$H = 0.7$	0.1	300	0.112	0.100	0.110	0.124	0.139	0.114	0.102	0.104	0.125	0.168	0.106	0.146	0.191	0.176	0.061
		500	0.093	0.091	0.092	0.100	0.145	0.096	0.096	0.091	0.110	0.206	0.095	0.178	0.245	0.251	0.082
		1000	0.084	0.074	0.075	0.092	0.176	0.077	0.076	0.091	0.116	0.393	0.084	0.236	0.388	0.591	0.122
	0.2	300	0.0868	0.081	0.073	0.087	0.113	0.080	0.085	0.097	0.100	0.185	0.084	0.148	0.246	0.297	0.653
		500	0.075	0.071	0.076	0.080	0.122	0.073	0.077	0.082	0.104	0.285	0.075	0.206	0.343	0.532	0.879
		1000	0.068	0.068	0.073	0.084	0.187	0.068	0.066	0.088	0.114	0.571	0.066	0.305	0.567	0.922	0.994
$H = 0.8$	0.1	300	0.140	0.135	0.132	0.136	0.152	0.130	0.141	0.126	0.134	0.186	0.116	0.164	0.211	0.182	0.062
		500	0.131	0.122	0.126	0.130	0.166	0.108	0.123	0.137	0.135	0.216	0.101	0.185	0.283	0.251	0.073
		1000	0.108	0.117	0.115	0.127	0.220	0.107	0.108	0.128	0.145	0.434	0.090	0.266	0.420	0.599	0.113
	0.2	300	0.118	0.119	0.121	0.123	0.149	0.107	0.119	0.109	0.125	0.203	0.092	0.177	0.261	0.300	0.619
		500	0.122	0.107	0.117	0.123	0.173	0.101	0.111	0.121	0.135	0.326	0.079	0.227	0.370	0.540	0.851
		1000	0.098	0.113	0.113	0.137	0.259	0.100	0.111	0.129	0.157	0.625	0.076	0.332	0.588	0.922	0.994
$H = 0.9$	0.1	300	0.175	0.181	0.187	0.169	0.173	0.166	0.165	0.177	0.168	0.192	0.120	0.193	0.268	0.176	0.054
		500	0.167	0.181	0.180	0.167	0.204	0.152	0.164	0.175	0.160	0.252	0.105	0.212	0.324	0.266	0.056
		1000	0.175	0.180	0.192	0.195	0.289	0.156	0.170	0.200	0.202	0.509	0.104	0.295	0.492	0.602	0.080
	0.2	300	0.169	0.171	0.174	0.166	0.184	0.140	0.155	0.179	0.163	0.261	0.102	0.206	0.349	0.304	0.501
		500	0.177	0.175	0.197	0.183	0.252	0.153	0.164	0.197	0.182	0.414	0.101	0.259	0.455	0.580	0.759
		1000	0.207	0.215	0.229	0.236	0.377	0.175	0.200	0.222	0.243	0.755	0.100	0.412	0.684	0.941	0.966

Table 2: Rejection rates of the change-point test based on the statistic  $\Gamma_n$ ,  $k_n = \lfloor np \rfloor$ , for LMSV time series (Pareto distributed  $\varepsilon_j$ ,  $j \geq 1$ ) of length  $n$  with Hurst parameter  $H$ , tail index  $\alpha$  and a shift in the mean of height  $h$  after a proportion  $\tau = 0.25$ . The calculations are based on 5,000 simulation runs.

## 4 Data

The analysis of financial time series, such as stock market prices, usually focuses on log returns instead of the observed data itself. As an example, we consider the log returns of the daily closing indices of Standard & Poor's 500 (S&P 500, in short) defined by

$$L_t := \log R_t, \quad R_t := \frac{P_t}{P_{t-1}},$$

where  $P_t$  denotes the value of the index on day  $t$ , in the period from January 2008 to December 2008; see Figure 1.




Figure 1: Daily closing index of Standard & Poor's 500 and its log returns from January 2008 to December 2008. The data has been obtained from Google Finance.

Comparing the plots of the sample autocorrelation function of the log returns and the sample autocorrelation function of their absolute values in Figure 2, we observe a phenomenon that is often encountered in the context of financial data: the log returns of the index appear to be uncorrelated, whereas the absolute log returns tend to be highly correlated.

Moreover, the plot in Figure 1 shows that the considered time series exhibits *volatility clustering*, meaning that large price changes, i.e., log returns with relatively large absolute values, tend to cluster. This indicates that observations are not independent across time, although the absence of linear autocorrelation suggests that the dependence is nonlinear; see [Cont \(2005\)](#).




Figure 2: Sample autocorrelation of the log returns and the absolute log returns of Standard & Poor's 500 daily closing index from January 2005 to December 2010. The two dashed horizontal lines mark the bounds for the 95% confidence interval of the autocovariances under the assumption of data generated by white noise.

Another characteristic of financial time series is the occurrence of heavy tails. In particular, probability distributions of log returns often exhibit tails which are heavier than those of a normal distribution. For the S&P 500 data, this property is highlighted by the Q-Q plot in Figure 3.

All of the previously described features of financial data can be covered by the LMSV model considered in our paper.

In view of the fact that the LMSV model captures properties of the log returns of Standard & Poor's 500 daily closing index, we are interested in analyzing the data with respect to a change in the tail index.

As in our simulations, we base the test decision on the statistic defined in (16). We choose  $k_n = \lfloor np \rfloor$ , i.e.,  $p$  defines the proportion of the data that the estimation of the tail index is based on. Choosing  $p = 0.1$ , the value of the test statistic corresponds to  $\tilde{\Gamma}_n = 1.467503$ . The 95%-quantile of the limit distribution  $\sup_{t \in [0,1]} |B(t) - tB(1)|$  equals 1.3463348. Choosing the critical value for the hypothesis test correspondingly, the value of  $\Gamma_n$  therefore indicates a change-point in the tail index at a level of significance of 5%.

A natural estimate for the change-point location is given by that point in time  $k$ ,




Figure 3: Q-Q plot for the log returns of Standard & Poor's 500 daily closing index from January 2005 to December 2010.

where  $\Gamma_{k,n}$  attains its maximum. For the considered data, this point in time corresponds to September 16, 2008, i.e., one day after September 15, 2008, the day Lehman Brothers filed for bankruptcy protection; see Figure 4.




Figure 4: Log returns of the daily closing index of Standard & Poor's 500 from January 2008 to December 2008. The red dashed line indicates the estimated change-point location.

## 5 Proofs

### 5.1 Proof of Theorem 2.6

#### 5.1.1 Decomposition of the tail empirical process

Recall that

$$e_n(s, t) = \left\{ \tilde{T}_n(s, t) - T(s, t) \right\},$$

where

$$\tilde{T}_n(s, t) := \frac{1}{n\bar{F}(u_n)} \sum_{j=1}^{\lfloor nt \rfloor} \mathbf{1}\{X_j > u_n s\} \quad \text{and} \quad T(s, t) = ts^{-\alpha}.$$

To prove Theorem 2.6, we consider the following decomposition:

$$e_n(s, t) = \left\{ \tilde{T}_n(s, t) - T_n(s, t) \right\} + \left\{ T_n(s, t) - T(s, t) \right\},$$

where

$$T_n(s, t) := \mathbb{E} \left[ \tilde{T}_n(s, t) \right] = \frac{\lfloor nt \rfloor}{n} \frac{\bar{F}(u_n s)}{\bar{F}(u_n)}.$$

Obviously, it holds that

$$\lim_{n \rightarrow \infty} T_n(s, t) = T(s, t)$$

for  $s > 0$  and  $t \in [0, 1]$ . In particular, the convergence holds uniformly on compact subsets of  $(0, \infty) \times [0, 1]$ . Moreover, for any  $s_0 > 0$  it holds that

$$\sup_{s \geq s_0, t \in [0, 1]} |T_n(s, t) - T(s, t)| \leq \sup_{s \geq s_0} \frac{\bar{F}(u_n s)}{\bar{F}(u_n)} \sup_{t \in [0, 1]} \left| \frac{\lfloor nt \rfloor}{n} - t \right| + \sup_{s \geq s_0} \left| \frac{\bar{F}(u_n s)}{\bar{F}(u_n)} - s^{-\alpha} \right|.$$

Note that

$$\sup_{t \in [0, 1]} \left| \frac{\lfloor nt \rfloor}{n} - t \right| = o \left( \frac{d_{n,r}}{n} + \frac{1}{\sqrt{n\bar{F}(u_n)}} \right).$$

Due to Proposition 2.8 in Kulik and Soulier (2011) and (TA.2) we have

$$\sup_{s \geq s_0} \left| \frac{\bar{F}(u_n s)}{\bar{F}(u_n)} - s^{-\alpha} \right| = o \left( \frac{d_{n,r}}{n} + \frac{1}{\sqrt{n\bar{F}(u_n)}} \right),$$

which implies

$$\sup_{s \geq s_0, t \in [0, 1]} |T_n(s, t) - T(s, t)| = o \left( \frac{d_{n,r}}{n} + \frac{1}{\sqrt{n\bar{F}(u_n)}} \right).$$

Since

$$\lim_{n \rightarrow \infty} \frac{\bar{F}_\varepsilon(u_n)}{\bar{F}(u_n)} = \frac{1}{\mathbb{E}[\sigma^\alpha(Y_1)]}$$

by (TA.3) and Breiman's Lemma, it therefore suffices to study weak convergence of the process

$$\tilde{e}_n(s, t) = \frac{1}{n\bar{F}_\varepsilon(u_n)} \sum_{j=1}^{\lfloor nt \rfloor} (\mathbf{1}\{X_j > u_n s\} - \bar{F}_\varepsilon(u_n s)).$$

For this, we consider the following decomposition:

$$\tilde{\varepsilon}_n(s, t) =: M_n(s, t) + R_n(s, t), \quad (17)$$

where

$$M_n(s, t) := \frac{1}{n\bar{F}_\varepsilon(u_n)} \sum_{j=1}^{\lfloor nt \rfloor} (\mathbf{1}\{X_j > u_n s\} - \mathbb{E}[\mathbf{1}\{X_j > u_n s\} | \mathcal{F}_{j-1}]),$$

$$R_n(s, t) := \frac{1}{n\bar{F}_\varepsilon(u_n)} \sum_{j=1}^{\lfloor nt \rfloor} (\mathbb{E}[\mathbf{1}\{X_j > u_n s\} | \mathcal{F}_{j-1}] - \bar{F}(u_n s)),$$

and

$$\mathcal{F}_j := \sigma(\varepsilon_k, \eta_k, k \in \mathbb{Z}, k \leq j). \quad (18)$$

We call  $M_n$  the *martingale part*, while we refer to  $R_n$  as the *long-range dependent part*.

### 5.1.2 The long-range dependent part

**Proposition 5.1** (Weak convergence of  $R_n(s, t)$ ). *Under the assumptions of Theorem 2.6, the following holds:*

$$\frac{n}{d_{n,r}} R_n(s, t) \Rightarrow s^{-\alpha} \frac{1}{r!} J_r(\Psi) Z_{r,H}(t),$$

where  $\Rightarrow$  denotes weak convergence in  $D([1, \infty] \times [0, 1])$ .

*Proof.* Note that

$$\mathbb{E}[\mathbf{1}\{X_j > u_n s\} | \mathcal{F}_{j-1}] = \bar{F}_\varepsilon\left(\frac{u_n s}{\sigma(Y_j)}\right) \quad (19)$$

and

$$\mathbb{E}\left[\bar{F}_\varepsilon\left(\frac{u_n s}{\sigma(Y_j)}\right)\right] = \mathbb{E}[\mathbb{E}[\mathbf{1}\{X_j > u_n s\} | \mathcal{F}_{j-1}]] = \mathbb{E}[\mathbf{1}\{X_j > u_n s\}] = \bar{F}(u_n s).$$

As a result, we can rewrite  $R_n(s, t)$  as follows:

$$R_n(s, t) = \frac{1}{n\bar{F}_\varepsilon(u_n)} \sum_{j=1}^{\lfloor nt \rfloor} \left( \bar{F}_\varepsilon\left(\frac{u_n s}{\sigma(Y_j)}\right) - \mathbb{E}\left[\bar{F}_\varepsilon\left(\frac{u_n s}{\sigma(Y_j)}\right)\right] \right) = \frac{1}{n} \sum_{j=1}^{\lfloor nt \rfloor} (\Psi_n(Y_j, s) - \mathbb{E}[\Psi_n(Y_j, s)]),$$

where

$$\Psi_n(y, s) = \frac{\bar{F}_\varepsilon\left(\frac{u_n s}{\sigma(y)}\right)}{\bar{F}_\varepsilon(u_n)}. \quad (20)$$



Due to regular variation of  $\bar{F}_\varepsilon$ , we have

$$\Psi_n(y, s) = \frac{\bar{F}_\varepsilon\left(\frac{u_n s}{\sigma(y)}\right)}{\bar{F}_\varepsilon(u_n)} \sim \left(\frac{s}{\sigma(y)}\right)^{-\alpha} = s^{-\alpha} \Psi(y). \quad (21)$$

Furthermore, it holds that

$$\begin{aligned} R_n(s, t) &= \frac{1}{n} \sum_{j=1}^{\lfloor nt \rfloor} (\Psi_n(Y_j, s) - \mathbb{E}[\Psi_n(Y_j, s)]) \\ &= \frac{1}{n} \sum_{j=1}^{\lfloor nt \rfloor} (\Psi_s(Y_j) - \mathbb{E}[\Psi_s(Y_j)]) + \frac{1}{n} \sum_{j=1}^{\lfloor nt \rfloor} (\Psi_n(Y_j, s) - \Psi_s(Y_j)) \\ &\quad + \frac{1}{n} \sum_{j=1}^{\lfloor nt \rfloor} (\mathbb{E}[\Psi_s(Y_j)] - \mathbb{E}[\Psi_n(Y_j, s)]), \end{aligned} \quad (22)$$

where  $\Psi_s(y) := s^{-\alpha} \Psi(y)$ .

As  $\mathbb{E}[\sigma^{2\alpha}(Y_1)] < \infty$  by (TA.3), the functional non-central limit theorem of Taqqu (1979) yields

$$\frac{n}{d_{n,r}} \frac{1}{n} \sum_{j=1}^{\lfloor nt \rfloor} (\Psi_s(Y_j) - \mathbb{E}[\Psi_s(Y_j)]) = s^{-\alpha} \frac{n}{d_{n,r}} \frac{1}{n} \sum_{j=1}^{\lfloor nt \rfloor} (\Psi(Y_j) - \mathbb{E}[\Psi(Y_j)]) \Rightarrow s^{-\alpha} \frac{1}{r!} J_r(\Psi) Z_{r,H}(t).$$

In the following, we will see that the first and the second summand in (22) are negligible. For this, it suffices to show that

$$\lim_{n \rightarrow \infty} \frac{\lfloor nt \rfloor}{d_{n,r}} \mathbb{E} [|\Psi_n(Y_1, s) - s^{-\alpha} \Psi(Y_1)|] = 0.$$

Note that

$$\mathbb{E} [|\Psi_n(Y_1, s) - s^{-\alpha} \Psi(Y_1)|] = \int_{\mathbb{R}} \left| \frac{\bar{F}_\varepsilon\left(\frac{u_n s}{\sigma(y)}\right)}{\bar{F}_\varepsilon(u_n)} - \left(\frac{s}{\sigma(y)}\right)^{-\alpha} \right| \varphi(y) dy.$$

Due to second order regular variation of  $\bar{F}_\varepsilon$ , Lemma 6.1 implies that for any  $\varepsilon > 0$

$$\left| \frac{\bar{F}_\varepsilon\left(\frac{u_n s}{\sigma(y)}\right)}{\bar{F}_\varepsilon(u_n)} - \left(\frac{s}{\sigma(y)}\right)^{-\alpha} \right| \leq C \eta^*(u_n) \left(\frac{s}{\sigma(y)}\right)^{-\rho-\alpha} \left(\max\left\{\frac{s}{\sigma(y)}, \frac{\sigma(y)}{s}\right\}\right)^\varepsilon.$$

Thus, it follows that

$$\sup_{s \geq s_0} \mathbb{E} [|\Psi_n(Y_1, s) - s^{-\alpha} \Psi(Y_1)|] \leq C \eta^*(u_n) s_0^{-\alpha-\rho-\varepsilon} \int_{\mathbb{R}} \sigma^{\alpha+\rho+\varepsilon}(y) \varphi(y) dy.$$

By (TA.2) and (TA.3), i.e., since  $\eta^*(u_n) = o(d_{n,r}/n)$  and  $\mathbb{E}[\sigma^{\alpha+\rho+\varepsilon}(Y_1)] < \infty$ , it holds that

$$\sup_{s \geq s_0} \mathbb{E} [|\Psi_n(Y_1, s) - s^{-\alpha} \Psi(Y_1)|] = o\left(\frac{d_{n,r}}{n}\right).$$

This completes the proof of Proposition 5.1.  $\square$

### 5.1.3 The martingale part

The goal of this section is to prove the following proposition:

**Proposition 5.2.** *Under the assumptions of Theorem 2.6, for any  $R > 1$ , the sequence  $\sqrt{n\bar{F}_\varepsilon(u_n)}M_n(s, t)$ ,  $n \geq 1$ , converges in distribution to  $\sqrt{\mathbb{E}[\sigma^\alpha(Y_1)]}B(s^{-\alpha}, t)$  in  $D([1, R] \times [0, 1])$ , where  $B$  denotes a standard Brownian sheet.*

First, we establish convergence of the finite dimensional distributions. Then, we proceed with a proof of tightness. For the latter, we use chaining arguments; see Section 5.1.3).

#### The martingale part: Convergence of the finite dimensional distributions

We have to prove that for all positive integers  $d_1$  and  $d_2$ , all  $1 \leq s_{d_1} < \dots < s_1$  with  $s_i \in \mathbb{R}$  and all  $0 \leq t_1 < \dots < t_{d_2} \leq 1$ , the vector with entries  $\sqrt{n\bar{F}_\varepsilon(u_n)}M_n(s_i, t_j)$ ,  $1 \leq i \leq d_1$ ,  $1 \leq j \leq d_2$ , converges in distribution to  $\sqrt{\mathbb{E}[\sigma^\alpha(Y_1)]}B(s_i^{-\alpha}, t_j)$ ,  $1 \leq i \leq d_1$ ,  $1 \leq j \leq d_2$ . For this, it suffices to consider the case  $d_1 = d_2$ . Indeed, if  $d_1 < d_2$ , we include  $s_i$ ,  $d_1 + 1 \leq i \leq d_2$ , in a decreasing order between  $s_{d_1}$  and  $s_{d_1-1}$ , i.e., such that  $1 \leq s_{d_1} < s_{d_2} < \dots < s_{d_1+1} < s_{d_1-1} < \dots < s_1$ , and if  $d_2 < d_1$ , we include  $t_j$ ,  $d_2 + 1 \leq j \leq d_1$  between  $t_{d_2-1}$  and  $t_{d_2}$  in an increasing order, i.e.,  $t_{d_2-1} < t_{d_2+1} < \dots < t_{d_1} < t_{d_2} \leq 1$ . Letting  $d = \max\{d_1, d_2\}$ , the convergence in distribution of the vector with entries  $\sqrt{n\bar{F}_\varepsilon(u_n)}M_n(s_i, t_j)$ ,  $1 \leq i \leq d$ ,  $1 \leq j \leq d$ , to  $\sqrt{\mathbb{E}[\sigma^\alpha(Y_1)]}B(s_i^{-\alpha}, t_j)$ ,  $1 \leq i \leq d$ ,  $1 \leq j \leq d$ , implies convergence of  $\sqrt{n\bar{F}_\varepsilon(u_n)}M_n(s_i, t_j)$ ,  $1 \leq i \leq d_1$ ,  $1 \leq j \leq d_2$ , to  $\sqrt{\mathbb{E}[\sigma^\alpha(Y_1)]}B(s_i^{-\alpha}, t_j)$ ,  $1 \leq i \leq d_1$ ,  $1 \leq j \leq d_2$ .

Let  $d \geq 1$  be an integer and let  $s_1, \dots, s_d$  and  $t_1, \dots, t_d$  be real numbers such that  $1 \leq s_d < \dots < s_1 \leq R$  and  $0 =: t_0 \leq t_1 < \dots < t_d \leq 1$ . Define  $t$  intervals  $I_{n,1} := (s_1 u_n, \infty)$ , and for  $2 \leq i \leq d$ , let  $I_{n,i} := (s_i u_n, s_{i-1} u_n]$ . Moreover, define random variables

$$Z_{i,j}^n := \frac{1}{\sqrt{n\bar{F}_\varepsilon(u_n)}} \sum_{k=[nt_{j-1}]+1}^{\lfloor nt_j \rfloor} (\mathbf{1}\{X_k \in I_{n,i}\} - \mathbb{E}[\mathbf{1}\{X_k \in I_{n,i}\} | \mathcal{F}_{k-1}])$$

for  $1 \leq i, j \leq d$ .

**Lemma 5.3.** *The sequence of random vectors  $(Z_{i,j}^n)_{1 \leq i, j \leq d}$ ,  $n \geq 1$ , converges in distribution to  $(Z_{i,j})_{1 \leq i, j \leq d}$ , where the random variables  $Z_{i,j}$ ,  $1 \leq i, j \leq d$ , are independent, and for all  $i, j \in \{1, \dots, d\}$ , the random variable  $Z_{i,j}$  is Gaussian, centered, and has variance  $\mathbb{E}[\sigma^\alpha(Y_1)](t_j - t_{j-1})(s_i^{-\alpha} - s_{i-1}^{-\alpha})$ , where  $s_0^{-\alpha} = 0$ .*

This lemma directly provides convergence of the finite dimensional distributions, after having applied the continuous mapping theorem to the function

$$(x_{i,j})_{i,j=1}^d \mapsto \left( \sum_{i'=1}^i \sum_{j'=1}^j x_{i',j'} \right)_{i,j=1}^d .$$

*Proof of Lemma 5.3.* By the Cramér-Wold device, we have to prove that for each collection of real numbers  $a_{i,j}$ ,  $1 \leq i, j \leq d$ , the sequence  $\sum_{i,j=1}^d a_{i,j} Z_{i,j}^n$ ,  $n \geq 1$ , converges to a normal distribution with mean zero and variance

$$\sigma^2 := \mathbb{E}[\sigma^\alpha(Y_1)] \sum_{i,j=1}^d (t_j - t_{j-1}) (s_i^{-\alpha} - s_{i-1}^{-\alpha}) a_{i,j}^2. \quad (23)$$

We will prove this by applying the following central limit theorem for martingale difference arrays. For this, recall that  $\Delta_{n,k}$ ,  $n \geq 1$ ,  $1 \leq k \leq n$ , is a martingale difference array with respect to the filtration  $\mathcal{F}_{n,k}$ ,  $n \geq 1$ ,  $0 \leq k \leq n$ , if for all  $n$ ,  $\mathcal{F}_{n,k} \subset \mathcal{F}_{n,k+1}$ ,  $1 \leq k \leq n-1$ ,  $\Delta_{n,k}$  is  $\mathcal{F}_{n,k}$ -measurable and  $\mathbb{E}[\Delta_{n,k} | \mathcal{F}_{n,k-1}] = 0$ .

**Theorem 5.4** (Theorem VIII. 1 in [Pollard \(1984\)](#)). *Let  $(\Delta_{n,k})_{n \geq 1, 1 \leq k \leq n}$  be a martingale difference array with respect to the filtration  $(\mathcal{F}_{n,k})_{n \geq 1, 0 \leq k \leq n}$ , such that  $\Delta_{n,k}$  is square integrable for all  $n$  and  $k$ . Moreover, assume that*

1. *for each positive  $\epsilon$ ,*

$$\sum_{k=1}^n \mathbb{E}[\Delta_{n,k}^2 \mathbf{1}\{|\Delta_{n,k}| > \epsilon\} | \mathcal{F}_{n,k-1}] \rightarrow 0 \text{ in probability}; \quad (24)$$

2.

$$\sum_{k=1}^n \mathbb{E}[\Delta_{n,k}^2 | \mathcal{F}_{n,k-1}] \rightarrow \sigma^2 \text{ in probability}. \quad (25)$$

*Then, the sequence  $\sum_{k=1}^n \Delta_{n,k}$ ,  $n \geq 1$ , converges in distribution to a normal law with mean zero and variance  $\sigma^2$  defined in (23).*

We express  $\tilde{Z}_n := \sum_{i,j=1}^d a_{i,j} Z_{i,j}^n$  as a sum of martingale differences. For  $1 \leq i \leq d$ , define

$$D_{n,i,k} := \frac{1}{\sqrt{nF_\epsilon(u_n)}} (\mathbf{1}\{X_k \in I_{n,i}\} - \mathbb{E}[\mathbf{1}\{X_k \in I_{n,i}\} | \mathcal{F}_{k-1}]).$$

If for some  $j \in \{1, \dots, d\}$  the integer  $k$  satisfies  $\lfloor nt_j \rfloor + 1 \leq k \leq \lfloor nt_{j+1} \rfloor$ , then we define

$$\Delta_{n,k} := \sum_{i=1}^d a_{i,j} D_{n,i,k},$$

and if  $\lfloor nt_d \rfloor + 1 \leq k \leq n$ , we define  $\Delta_{n,k} := 0$ .

In this way,  $\tilde{Z}_n = \sum_{k=1}^n \Delta_{n,k}$  and defining  $\mathcal{F}_{n,j}$  as the  $\sigma$ -algebra  $\mathcal{F}_j$  given by (18), the array  $\Delta_{n,k}$ ,  $n \geq 1$ ,  $1 \leq k \leq n$ , is a square integrable martingale difference array with respect to the filtration  $\mathcal{F}_{n,k}$ ,  $n \geq 1$ ,  $0 \leq k \leq n$ . Let us check (24). Observe that for all  $1 \leq k \leq n$  and  $1 \leq i \leq d$ ,  $|D_{n,i,k}| \leq 2 (n\bar{F}_\varepsilon(u_n))^{-1/2}$ . Hence, for all  $1 \leq k \leq n$ ,

$$|\Delta_{n,k}| \leq 2 \sum_{i,j=1}^d |a_{i,j}| (n\bar{F}_\varepsilon(u_n))^{-1/2}.$$

Consequently, for a fixed  $\varepsilon$ , when  $n$  is such that  $2 \sum_{i,j=1}^d |a_{i,j}| (n\bar{F}_\varepsilon(u_n))^{-1/2} < \varepsilon$ , the indicator  $\mathbf{1}\{|\Delta_{n,k}| > \varepsilon\}$  vanishes and hence (24) holds.

Let us check (25). It suffices to prove that for all  $j \in \{1, \dots, d\}$

$$\mathbb{E} \left[ \left| \sum_{k=\lfloor nt_{j-1} \rfloor + 1}^{\lfloor nt_j \rfloor} \mathbb{E} [\Delta_{n,k}^2 | \mathcal{F}_{k-1}] - \mathbb{E} [\sigma^\alpha(Y_1)] (t_j - t_{j-1}) \sum_{i=1}^d a_{i,j}^2 (s_i^{-\alpha} - s_{i-1}^{-\alpha}) \right| \right] \rightarrow 0. \quad (26)$$

To this aim, we decompose  $\mathbb{E} [\Delta_{n,k}^2 | \mathcal{F}_{k-1}]$ :

$$\mathbb{E} [\Delta_{n,k}^2 | \mathcal{F}_{k-1}] = \sum_{i=1}^d a_{i,j}^2 \mathbb{E} [D_{n,i,k}^2 | \mathcal{F}_{k-1}] + 2 \sum_{1 \leq i_1 < i_2 \leq d} a_{i_1,j} a_{i_2,j} \mathbb{E} [D_{n,i_1,k} D_{n,i_2,k} | \mathcal{F}_{k-1}].$$

Observe that using (19)

$$\begin{aligned} \mathbb{E} [D_{n,i,k}^2 | \mathcal{F}_{k-1}] &= \frac{1}{n\bar{F}_\varepsilon(u_n)} \left( \bar{F}_\varepsilon \left( \frac{u_n s_i}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i-1}}{\sigma(Y_k)} \right) \right) \\ &\quad - \frac{1}{n\bar{F}_\varepsilon(u_n)} \left( \bar{F}_\varepsilon \left( \frac{u_n s_i}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i-1}}{\sigma(Y_k)} \right) \right)^2, \end{aligned}$$

where we set  $\bar{F}_\varepsilon(s_0 x) = 0$  for all  $x$ . Moreover, it holds that

$$\begin{aligned} &\mathbb{E} [D_{n,i_1,k} D_{n,i_2,k} | \mathcal{F}_{k-1}] \\ &= \frac{1}{n\bar{F}_\varepsilon(u_n)} \left( \bar{F}_\varepsilon \left( \frac{u_n s_{i_1}}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i_1-1}}{\sigma(Y_k)} \right) \right) \left( \bar{F}_\varepsilon \left( \frac{u_n s_{i_2}}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i_2-1}}{\sigma(Y_k)} \right) \right). \end{aligned}$$

Hence, the expression on the left hand side of (26) is bounded by

$$\mathbb{E} [|R_{n,1}|] + \mathbb{E} [|R_{n,2}|] + \mathbb{E} [|R_{n,3}|],$$

where

$$\begin{aligned} R_{n,1} &:= \sum_{i=1}^d a_{i,j}^2 \frac{1}{n\bar{F}_\varepsilon(u_n)} \sum_{k=\lfloor nt_{j-1} \rfloor + 1}^{\lfloor nt_j \rfloor} \left( \bar{F}_\varepsilon \left( \frac{u_n s_i}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i-1}}{\sigma(Y_k)} \right) \right) \\ &\quad - \mathbb{E} [\sigma^\alpha(Y_1)] (t_j - t_{j-1}) \sum_{i=1}^d a_{i,j}^2 (s_i^{-\alpha} - s_{i-1}^{-\alpha}), \end{aligned}$$

$$R_{n,2} := \sum_{i=1}^d a_{i,j}^2 \frac{1}{n\bar{F}_\varepsilon(u_n)} \sum_{k=\lfloor nt_{j-1} \rfloor + 1}^{\lfloor nt_j \rfloor} \left( \bar{F}_\varepsilon \left( \frac{u_n s_i}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i-1}}{\sigma(Y_k)} \right) \right)^2,$$

$$R_{n,3} := \frac{1}{n\bar{F}_\varepsilon(u_n)} \sum_{k=\lfloor nt_{j-1} \rfloor + 1}^{\lfloor nt_j \rfloor} \sum_{1 \leq i_1 < i_2 \leq d} a_{i_1,j} a_{i_2,j} \\ \times \left( \bar{F}_\varepsilon \left( \frac{u_n s_{i_1}}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i_1-1}}{\sigma(Y_k)} \right) \right) \left( \bar{F}_\varepsilon \left( \frac{u_n s_{i_2}}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i_2-1}}{\sigma(Y_k)} \right) \right).$$

Here we omit the dependence on  $j \in \{1, \dots, d\}$  in  $R_{n,1}$  and  $R_{n,2}$  in order to ease notations.

We start with  $R_{n,1}$ . Using stationarity, it suffices to prove that for all  $i \in \{1, \dots, d\}$ ,

$\mathbb{E}[|R_{n,1,i}|] \rightarrow 0$ , where

$$R_{n,1,i} := \frac{1}{n\bar{F}_\varepsilon(u_n)} \sum_{k=1}^{\lfloor nt_j \rfloor - \lfloor nt_{j-1} \rfloor} \left( \bar{F}_\varepsilon \left( \frac{u_n s_i}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i-1}}{\sigma(Y_k)} \right) \right) \\ - \mathbb{E}[\sigma^\alpha(Y_1)] (t_j - t_{j-1}) (s_i^{-\alpha} - s_{i-1}^{-\alpha}).$$

Applying Lemma 6.2 with  $t = u_n$ ,  $a = s_i/\sigma(Y_k)$ ,  $b = s_{i-1}/\sigma(Y_k)$  and  $\epsilon = 1$ , we get

$$|R_{n,1,i}| \leq \left| \frac{1}{n} \sum_{k=1}^{\lfloor nt_j \rfloor - \lfloor nt_{j-1} \rfloor} \left( \frac{\sigma^\alpha(Y_k)}{s_i^\alpha} - \frac{\sigma^\alpha(Y_k)}{s_{i-1}^\alpha} \right) - (t_j - t_{j-1}) (s_i^{-\alpha} - s_{i-1}^{-\alpha}) \mathbb{E}[\sigma^\alpha(Y_1)] \right| \\ + C\eta^*(u_n) \frac{1}{n} \sum_{k=1}^{\lfloor nt_j \rfloor - \lfloor nt_{j-1} \rfloor} \left( \max \left\{ \frac{\sigma(Y_k)}{s_i}, 1 \right\} \right)^{\alpha+\rho+1} \frac{s_i - s_{i-1}}{\sigma(Y_k)}.$$

By (TA.3) and (TA.4) we have

$$\mathbb{E} \left[ \frac{(\max \{ \sigma(Y_0), 1 \})^{\alpha+\rho+1}}{\sigma(Y_0)} \right] < \infty.$$

Using the ergodic theorem for the first term, the fact that  $\eta^*(u_n) \rightarrow 0$  allows us to conclude that  $\mathbb{E}[|R_{n,1}|] \rightarrow 0$  as  $n$  goes to infinity.

The treatment of  $R_{n,2}$  and  $R_{n,3}$  is the same: we take expectations and bound one of the factors  $\bar{F}_\varepsilon \left( \frac{u_n s_i}{\sigma(Y_k)} \right) - \bar{F}_\varepsilon \left( \frac{u_n s_{i-1}}{\sigma(Y_k)} \right)$  using (43) in Lemma 6.2 when  $i = 1$  and (45) in Lemma 6.2 when  $2 \leq i \leq d$ . We then conclude by the dominated convergence theorem.

This finishes the proof of Lemma 5.3.  $\square$

### The martingale part: Tightness

**Lemma 5.5.** *Under the assumptions of Theorem 2.6, for any  $R > 1$ , the sequence*

$$\sqrt{n\bar{F}_\varepsilon(u_n)} M_n(s, t), \quad n \geq 1,$$

*is tight in  $D([1, R] \times [0, 1])$ .*

*Proof.* Define

$$m_n(s, t) := \sqrt{n\bar{F}_\varepsilon(u_n)} M_n(s, t).$$

In order to prove tightness of  $m_n(s, t)$ , we validate the following tightness criterion: for all  $\epsilon > 0$

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \mathbb{P} \left( \sup_{\substack{|s_2 - s_1| < \delta \\ 1 \leq s_1, s_2 \leq R}} \sup_{\substack{|t_2 - t_1| < \delta \\ 0 \leq t_1, t_2 \leq 1}} |m_n(s_2, t_2) - m_n(s_1, t_1)| > \epsilon \right) = 0.$$

Writing

$$m_n(s_2, t_2) - m_n(s_1, t_1) = m_n(s_2, t_2) - m_n(s_1, t_2) + m_n(s_1, t_2) - m_n(s_1, t_1),$$

it suffices to show

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \mathbb{P} \left( \sup_{\substack{|s_2 - s_1| < \delta \\ 1 \leq s_1, s_2 \leq R}} \sup_{t \in [0, 1]} |m_n(s_2, t) - m_n(s_1, t)| > \epsilon \right) = 0 \quad (27)$$

and

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \mathbb{P} \left( \sup_{1 \leq s \leq R} \sup_{\substack{|t_2 - t_1| < \delta \\ 0 \leq t_1, t_2 \leq 1}} |m_n(s, t_2) - m_n(s, t_1)| > \epsilon \right) = 0. \quad (28)$$

#### 5.1.4 Proof of (27).

In order to prove (27), we apply a chaining technique.

For this, we define the intervals

$$I_{1,k} := [1 + 2k\delta, 1 + 2(k+1)\delta] \quad \text{and} \quad I_{2,k} := [1 + (2k+1)\delta, 1 + (2(k+1)+1)\delta]$$

for  $k = 0, \dots, L_\delta := \lfloor \frac{R-1}{2\delta} \rfloor$ . Then, the expression inside  $\mathbb{P}$  in (27) is bounded by

$$\max_{0 \leq k \leq L_\delta} \sup_{s_1, s_2 \in I_{1,k}} \sup_{t \in [0, 1]} |m_n(s_2, t) - m_n(s_1, t)| + \max_{0 \leq k \leq L_\delta} \sup_{s_1, s_2 \in I_{2,k}} \sup_{t \in [0, 1]} |m_n(s_2, t) - m_n(s_1, t)|. \quad (29)$$

In the following, we consider the first summand only, since for the second summand analogous considerations hold, i.e., it remains to show that

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \mathbb{P} \left( \max_{0 \leq k \leq L_\delta} \sup_{s_1, s_2 \in I_{1,k}} \sup_{t \in [0, 1]} |m_n(s_2, t) - m_n(s_1, t)| > \epsilon \right) = 0.$$

For this, it suffices to show that

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{\delta} \max_{0 \leq k \leq L_\delta} \mathbb{P} \left( \sup_{s_1, s_2 \in I_{1,k}} \sup_{t \in [0,1]} |m_n(s_2, t) - m_n(s_1, t)| > \epsilon \right) = 0.$$

We write  $I_{1,k} = [a_k, a_{k+1}]$ , i.e.,  $a_k := 1 + 2k\delta$  and  $a_{k+1} := 1 + 2(k+1)\delta$ . Note that

$$\sup_{s_1, s_2 \in I_{1,k}} \sup_{t \in [0,1]} |m_n(s_2, t) - m_n(s_1, t)| \leq 2 \sup_{x \in [0, 2\delta]} \sup_{t \in [0,1]} |m_n(a_k, t) - m_n(a_k + x, t)|.$$

Define refining partitions  $x_i(k)$  for  $k = 0, \dots, K_n$  with  $K_n \rightarrow \infty$ , for  $n \rightarrow \infty$ , by

$$x_i(k) := \sup \left\{ x \in [1, R] \mid \frac{\bar{F}_\epsilon(u_n x)}{\bar{F}_\epsilon(u_n)} \geq \frac{i}{2^k} \delta \right\}, \quad i = 0, \dots, \lfloor 2^k \delta^{-1} \rfloor,$$

and choose  $i_k(x)$  such that

$$a_k + x \in (x_{i_k(x)+1}(k), x_{i_k(x)}(k)].$$

From the definition of  $m_n$  we obtain

$$|m_n(y, t) - m_n(x, t)| = \left| \frac{1}{\sqrt{n\bar{F}_\epsilon(u_n)}} \sum_{j=1}^{\lfloor nt \rfloor} (\mathbf{1}\{u_n x < X_j \leq u_n y\} - \mathbb{E}[\mathbf{1}\{u_n x < X_j \leq u_n y\} | \mathcal{F}_{j-1}]) \right|.$$

Then, with  $m_n(x) := \sup_{t \in [0,1]} |m_n(x, t)|$  and  $\bar{m}_n(x, y) := \sup_{t \in [0,1]} |m_n(y, t) - m_n(x, t)|$ , it follows that

$$\begin{aligned} & \sup_{t \in [0,1]} |m_n(a_k, t) - m_n(a_k + x, t)| \\ & \leq \bar{m}_n(a_k, x_{i_{K_n}(x)+1}(K_n)) + \sum_{k=1}^{K_n} \bar{m}_n(x_{i_k(x)+1}(k), x_{i_{k-1}(x)+1}(k-1)) + \bar{m}_n(x_{i_0(x)+1}(0), a_k + x). \end{aligned}$$

As a result, we have

$$\begin{aligned} & \mathbb{P} \left( \sup_{x \in [0, 2\delta]} \sup_{t \in [0,1]} |m_n(a_k, t) - m_n(a_k + x, t)| > \epsilon \right) \leq \mathbb{P} \left( \sup_{x \in [0, 2\delta]} \bar{m}_n(a_k, x_{i_{K_n}(x)+1}(K_n)) > \frac{\epsilon}{4} \right) \\ & + \sum_{k=1}^{K_n} \mathbb{P} \left( \sup_{x \in [0, 2\delta]} \bar{m}_n(x_{i_k(x)+1}(k), x_{i_{k-1}(x)+1}(k-1)) > \frac{\epsilon}{(k+3)^2} \right) \\ & + \mathbb{P} \left( \sup_{x \in [0, 2\delta]} \bar{m}_n(x_{i_0(x)+1}(0), a_k + x) > \epsilon - \sum_{k=0}^{\infty} \frac{\epsilon}{(k+3)^2} - \frac{\epsilon}{4} \right). \end{aligned} \quad (30)$$

Since

$$\sum_{k=0}^{\infty} \frac{\epsilon}{(k+3)^2} \leq \frac{\epsilon}{2},$$

it follows that

$$\mathbb{P} \left( \sup_{x \in [0, 2\delta]} \bar{m}_n(x_{i_0(x)+1}(0), a_k + x) > \epsilon - \sum_{k=0}^{\infty} \frac{\epsilon}{(k+3)^2} - \frac{\epsilon}{4} \right) \leq \mathbb{P} \left( \sup_{x \in [0, 2\delta]} \bar{m}_n(x_{i_0(x)+1}(0), a_k + x) > \frac{\epsilon}{4} \right).$$

Additionally, we conclude that

$$\begin{aligned} & \sum_{k=1}^{K_n} \mathbb{P} \left( \sup_{x \in [0, 2\delta]} \bar{m}_n(x_{i_k(x)+1}(k), x_{i_{k-1}(x)+1}(k-1)) > \frac{\epsilon}{(k+3)^2} \right) \\ & \leq \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \mathbb{P} \left( \bar{m}_n(x_{i+1}(k), x_i(k)) > \frac{\epsilon}{(k+3)^2} \right). \end{aligned}$$

For further estimation, we use Freedman's inequality: if  $(d_j, \mathcal{F}_j)$ ,  $j \geq 1$ , is a martingale difference sequence such that  $\sup_j |d_j| \leq c$ , where  $c$  is a positive constant, then for all  $x, y > 0$

$$\begin{aligned} \mathbb{P} \left( \left\{ \max_{1 \leq \ell \leq n} \left| \sum_{j=1}^{\ell} d_j \right| > x \right\} \cap \left\{ \sum_{j=1}^n \mathbb{E} [d_j^2 \mid \mathcal{F}_{j-1}] \leq y \right\} \right) & \leq 2 \exp \left( -\frac{x^2}{2(y + \frac{2}{3}cx)} \right) \\ & \leq 2 \max \left\{ \exp \left( -\frac{x^2}{4y} \right), \exp \left( -\frac{3}{8}x/c \right) \right\}; \end{aligned}$$

see Theorem 1.6 in [Freedman \(1975\)](#).

For this purpose, we define

$$d_{j,i,k,n} := \mathbf{1} \{u_n x_{i+1}(k) < X_j \leq u_n x_i(k)\} - \mathbb{E} [\mathbf{1} \{u_n x_{i+1}(k) < X_j \leq u_n x_i(k)\} \mid \mathcal{F}_{j-1}].$$

Then, it follows that

$$\bar{m}_n(x_{i+1}(k), x_i(k)) = \max_{1 \leq \ell \leq n} \left| \frac{1}{\sqrt{n\bar{F}_\epsilon(u_n)}} \sum_{j=1}^{\ell} d_{j,i,k,n} \right|.$$

Furthermore, for  $B > 0$ , which is chosen later, define

$$y_{n,k,i} := n\bar{F}_\epsilon(u_n) B \left( \frac{\bar{F}_\epsilon(u_n x_{i+1}(k))}{\bar{F}_\epsilon(u_n)} - \frac{\bar{F}_\epsilon(u_n x_i(k))}{\bar{F}_\epsilon(u_n)} \right)$$

and

$$z_{n,k} := \frac{\epsilon \sqrt{n\bar{F}_\epsilon(u_n)}}{(k+3)^2}.$$

Since  $\bar{F}_\epsilon$  is continuous and since we can assume without loss of generality that  $\bar{F}_\epsilon$  is ultimately strictly decreasing, it holds that

$$\frac{\bar{F}_\epsilon(u_n x_{i+1}(k))}{\bar{F}_\epsilon(u_n)} - \frac{\bar{F}_\epsilon(u_n x_i(k))}{\bar{F}_\epsilon(u_n)} = \frac{\delta}{2^k},$$



and, consequently,

$$y_{n,k,i} = n\bar{F}_\varepsilon(u_n)B\frac{\delta}{2^k}.$$

For an estimation by Freedman's inequality, we have to specify  $K_n$ . Choosing  $K_n := \lceil \log_2(\delta a_n C) \rceil$  for some constant  $C$ , where  $a_n$ ,  $n \geq 1$ , is a sequence with  $a_n \rightarrow \infty$  and  $a_n = o\left(\frac{n}{d_{n,r}} + \sqrt{n\bar{F}_\varepsilon(u_n)}\right)$ , it follows (with  $c = 2$ ) that

$$\frac{z_{n,k}^2}{4y_{n,k,i}} = \frac{\epsilon^2 2^k}{4B(k+3)^4 \delta} \leq \frac{\epsilon^2 C \sqrt{n\bar{F}_\varepsilon(u_n)}}{4B(k+3)^4} \leq \frac{3\epsilon \sqrt{n\bar{F}_\varepsilon(u_n)}}{8 \cdot 2(k+3)^2} = \frac{3}{8} \frac{z_{n,k}}{c}$$

for a corresponding choice of  $C$ . Therefore, we have

$$\max \left\{ \exp\left(-\frac{z_{n,k}^2}{4y_{n,k,i}}\right), \exp\left(-\frac{3}{8} z_{n,k}/c\right) \right\} = \exp\left(-\frac{z_{n,k}^2}{4y_{n,k,i}}\right).$$

As a result, an application of Freedman's inequality yields

$$\begin{aligned} & \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \mathbb{P} \left( \bar{m}_n(x_{i+1}(k), x_i(k)) > \frac{\epsilon}{(k+3)^2}, \sum_{j=1}^n \mathbb{E} [d_{j,i,k,n}^2 | \mathcal{F}_{j-1}] \leq y_{n,k,i} \right) \\ & \leq \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \mathbb{P} \left( \max_{1 \leq \ell \leq n} \left| \sum_{j=1}^{\ell} d_{j,i,k,n} \right| > \frac{\epsilon \sqrt{n\bar{F}_\varepsilon(u_n)}}{(k+3)^2}, \sum_{j=1}^n \mathbb{E} [d_{j,i,k,n}^2 | \mathcal{F}_{j-1}] \leq y_{n,k,i} \right) \\ & \leq \sum_{k=1}^{K_n} (\lfloor 2^k \delta^{-1} \rfloor + 1) \exp\left(-\frac{\epsilon^2}{4(k+3)^4} \frac{2^k}{B\delta}\right). \end{aligned}$$

Noting that there exists a constant  $D > 0$  such that for all  $k$

$$(\lfloor 2^k \delta^{-1} \rfloor + 1) \exp\left(-\frac{\epsilon^2}{4(k+3)^4 \delta} \frac{2^k}{B}\right) \leq \lfloor 2^{\frac{k}{2}} \delta^{-1} \rfloor \exp\left(-D \frac{\epsilon^2}{4} \frac{2^{\frac{k}{2}}}{B\delta}\right),$$

elementary calculations yield

$$\begin{aligned} & \sum_{k=1}^{K_n} (\lfloor 2^k \delta^{-1} \rfloor + 1) \exp\left(-\frac{\epsilon^2}{4(k+3)^4} \frac{2^k}{B\delta}\right) \\ & \leq \frac{4B}{\epsilon^2 \log(2)} \left( \exp\left(-D \frac{\epsilon^2}{4B\delta}\right) - \exp\left(-D \frac{\epsilon^2}{4B\delta} 2^{\frac{K_n}{2}}\right) \right). \end{aligned}$$

It follows that

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{\delta} \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \mathbb{P} \left( \max_{1 \leq \ell \leq n} \left| \sum_{j=1}^{\ell} d_{j,i,k,n} \right| > z_{n,k}, \sum_{j=1}^n \mathbb{E} [d_{j,i,k,n}^2 | \mathcal{F}_{j-1}] \leq y_{n,k,i} \right) = 0.$$

Therefore, in order to finish the proof of (27), it remains to show that

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{\delta} \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \mathbb{P} \left( \sum_{j=1}^n \mathbb{E} [d_{j,i,k,n}^2 | \mathcal{F}_{j-1}] > y_{n,k,i} \right) = 0.$$

Since for an event  $A$  and a  $\sigma$ -algebra  $\mathcal{F}$

$$\mathbb{E}[(\mathbf{1}(A) - \mathbb{E}[\mathbf{1}(A) | \mathcal{F}])^2 | \mathcal{F}] = \mathbb{E}[\mathbf{1}(A) | \mathcal{F}] - (\mathbb{E}[\mathbf{1}(A) | \mathcal{F}])^2 \leq \mathbb{E}[\mathbf{1}(A) | \mathcal{F}],$$

it holds that

$$\mathbb{E}[d_{j,i,k,n}^2 | \mathcal{F}_{j-1}] \leq \bar{F}_\varepsilon \left( \frac{u_n x_{i+1}(k)}{\sigma(Y_j)} \right) - \bar{F}_\varepsilon \left( \frac{u_n x_i(k)}{\sigma(Y_j)} \right).$$

Therefore, we arrive at

$$\mathbb{P} \left( \sum_{j=1}^n \mathbb{E}[d_{j,i,k,n}^2 | \mathcal{F}_{j-1}] > y_{n,k,i} \right) \leq \mathbb{P} \left( \frac{1}{y_{n,k,i}} \sum_{j=1}^n \left( \bar{F}_\varepsilon \left( \frac{u_n x_{i+1}(k)}{\sigma(Y_j)} \right) - \bar{F}_\varepsilon \left( \frac{u_n x_i(k)}{\sigma(Y_j)} \right) \right) > 1 \right).$$

Note that

$$\frac{1}{y_{n,k,i}} \sum_{j=1}^n \left( \bar{F}_\varepsilon \left( \frac{u_n x_{i+1}(k)}{\sigma(Y_j)} \right) - \bar{F}_\varepsilon \left( \frac{u_n x_i(k)}{\sigma(Y_j)} \right) \right) = A_1(n, k, i) + A_2(n, k, i) + A_3(n, k, i), \quad (31)$$

where

$$\begin{aligned} A_1(n, k, i) &:= \frac{\bar{F}_\varepsilon(u_n)}{y_{n,k,i}} \sum_{j=1}^n \left\{ \frac{\bar{F}_\varepsilon \left( \frac{u_n x_{i+1}(k)}{\sigma(Y_j)} \right)}{\bar{F}_\varepsilon(u_n)} - \frac{\bar{F}_\varepsilon \left( \frac{u_n x_i(k)}{\sigma(Y_j)} \right)}{\bar{F}_\varepsilon(u_n)} - \sigma^\alpha(Y_j) (x_{i+1}(k)^{-\alpha} - x_i(k)^{-\alpha}) \right\}, \\ A_2(n, k, i) &:= \frac{\bar{F}_\varepsilon(u_n)}{y_{n,k,i}} \sum_{j=1}^n \sigma^\alpha(Y_j) \left\{ (x_{i+1}(k)^{-\alpha} - x_i(k)^{-\alpha}) - \left( \frac{\bar{F}_\varepsilon(u_n x_{i+1}(k))}{\bar{F}_\varepsilon(u_n)} - \frac{\bar{F}_\varepsilon(u_n x_i(k))}{\bar{F}_\varepsilon(u_n)} \right) \right\}, \\ A_3(n, k, i) &:= \frac{\bar{F}_\varepsilon(u_n)}{y_{n,k,i}} \sum_{j=1}^n \sigma^\alpha(Y_j) \left( \frac{\bar{F}_\varepsilon(u_n x_{i+1}(k))}{\bar{F}_\varepsilon(u_n)} - \frac{\bar{F}_\varepsilon(u_n x_i(k))}{\bar{F}_\varepsilon(u_n)} \right), \end{aligned}$$

so that

$$\begin{aligned} &\mathbb{P} \left( \frac{1}{y_{n,k,i}} \sum_{j=1}^n \left[ \bar{F}_\varepsilon \left( \frac{u_n x_{i+1}(k)}{\sigma(Y_j)} \right) - \bar{F}_\varepsilon \left( \frac{u_n x_i(k)}{\sigma(Y_j)} \right) \right] > 1 \right) \\ &\leq \mathbb{P} \left( |A_1(n, k, i)| > \frac{1}{3} \right) + \mathbb{P} \left( |A_2(n, k, i)| > \frac{1}{3} \right) + \mathbb{P} \left( |A_3(n, k, i)| > \frac{1}{3} \right). \quad (32) \end{aligned}$$

According to Lemma 6.2 in the appendix it holds that

$$|A_2(n, k, i)| \leq \frac{2^k \bar{F}_\varepsilon(u_n)}{n \bar{F}_\varepsilon(u_n) B \delta} \mathcal{O}(\eta^*(u_n)) (x_{i+1}(k) - x_i(k)) \sum_{j=1}^n \sigma^\alpha(Y_j).$$

Therefore, given Assumption (TA.2), it follows that

$$\begin{aligned} &\lim_{n \rightarrow \infty} \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \frac{1}{\delta} \mathbb{P} \left( |A_2(n, k, i)| > \frac{1}{3} \right) \\ &\leq \lim_{n \rightarrow \infty} \mathcal{O}(\eta^*(u_n)) \frac{1}{\delta} \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \frac{2^k}{B \delta} (x_{i+1}(k) - x_i(k)) \mathbb{E}[\sigma^\alpha(Y_1)] \\ &= \lim_{n \rightarrow \infty} \mathcal{O}(\eta^*(u_n)) \mathcal{O} \left( \frac{1}{\delta} \sum_{k=1}^{K_n} \frac{2^k}{B \delta} \right) = \lim_{n \rightarrow \infty} \mathcal{O}(\eta^*(u_n)) \mathcal{O} \left( \frac{1}{\delta} \frac{2^{K_n}}{B \delta} \right) = 0. \end{aligned}$$

For the first summand in (31), it follows by Lemma 6.2 in the appendix that

$$\begin{aligned}
|A_1(n, k, i)| &\leq \frac{\bar{F}_\varepsilon(u_n)}{y_{n,k,i}} \sum_{j=1}^n C\eta^*(u_n) \left( \frac{x_{i+1}(k)}{\sigma(Y_j)} - \frac{x_i(k)}{\sigma(Y_j)} \right) \left( \min \left\{ \frac{x_{i+1}(k)}{\sigma(Y_j)}, 1 \right\} \right)^{-\alpha-\rho-\epsilon} \\
&\leq \frac{\bar{F}_\varepsilon(u_n)}{n\bar{F}_\varepsilon(u_n)B\frac{\delta}{2^k}} C\eta^*(u_n)(x_{i+1}(k) - x_i(k)) \sum_{j=1}^n \sigma^{-1}(Y_j) (\max \{ \sigma(Y_j), 1 \})^{\alpha+\rho+\epsilon} \\
&= \frac{2^k}{B\delta} C\eta^*(u_n)(x_{i+1}(k) - x_i(k)) \frac{1}{n} \sum_{j=1}^n \sigma^{-1}(Y_j) (\max \{ \sigma(Y_j), 1 \})^{\alpha+\rho+\epsilon}.
\end{aligned}$$

According to Assumptions (TA.3) and (TA.4) the expectation of the summands on the right-hand side is finite and hence, for each  $\delta > 0$ ,

$$\begin{aligned}
&\lim_{n \rightarrow \infty} \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \frac{1}{\delta} \mathbb{P} \left( |A_1(n, k, i)| > \frac{1}{3} \right) \\
&\leq \lim_{n \rightarrow \infty} \frac{1}{\delta} \sum_{k=1}^{K_n} \sum_{i=0}^{\lfloor 2^k \delta^{-1} \rfloor} \frac{2^k}{nB\delta} \mathcal{O}(\eta^*(u_n))(x_{i+1}(k) - x_i(k)) \sum_{j=1}^n \mathbb{E} [\sigma^{-1}(Y_j) (\max \{ \sigma(Y_j), 1 \})^{\alpha+\rho+\epsilon}] \\
&= \lim_{n \rightarrow \infty} \mathcal{O}(\eta^*(u_n)) \mathcal{O} \left( \frac{1}{\delta} \sum_{k=1}^{K_n} \frac{2^k}{B\delta} \right) = \lim_{n \rightarrow \infty} \mathcal{O}(\eta^*(u_n)) \mathcal{O} \left( \frac{1}{\delta} \frac{2^{K_n}}{B\delta} \right) \\
&= \lim_{n \rightarrow \infty} \mathcal{O}(\eta^*(u_n)) \mathcal{O} \left( \frac{a_n}{B\delta} \right) = 0.
\end{aligned}$$

Finally, we consider the last summand in (31):

$$A_3(n, k, i) = \frac{1}{nB} \sum_{j=1}^n \sigma^\alpha(Y_j).$$

Obviously,  $A_3(n, k, i)$  depends neither on  $k$  nor on  $i$ . Due to the non-central limit theorem in Taqqu (1979), it holds that

$$\frac{1}{nB} \sum_{j=1}^n (\sigma^\alpha(Y_j) - \mathbb{E}[\sigma^\alpha(Y_1)]) = \mathcal{O}_P \left( \frac{d_{n,r}}{n} \right).$$

Choosing  $B > 6\mathbb{E}[\sigma^\alpha(Y_1)]$ , it follows that

$$\mathbb{P} \left( \frac{1}{nB} \sum_{j=1}^n \sigma^\alpha(Y_j) > \frac{1}{3} \right) \leq \mathbb{P} \left( \frac{1}{nB} \sum_{j=1}^n (\sigma^\alpha(Y_j) - \mathbb{E}[\sigma^\alpha(Y_1)]) > \frac{1}{6} \right) = \mathcal{O} \left( \frac{d_{n,r}}{n} \right).$$

Hence, we have

$$\begin{aligned}
&\lim_{n \rightarrow \infty} \sum_{k=1}^{K_n} \sum_{i=0}^{\frac{2^k}{\delta}-1} \frac{1}{\delta} \mathbb{P} \left( |A_3(n, k, i)| > \frac{1}{3} \right) \\
&= \lim_{n \rightarrow \infty} \mathbb{P} \left( \frac{1}{nB} \sum_{j=1}^n \sigma^\alpha(Y_j) > \frac{1}{3} \right) \sum_{k=1}^{K_n} \sum_{i=0}^{\frac{2^k}{\delta}-1} \frac{1}{\delta} \\
&= \lim_{n \rightarrow \infty} \mathcal{O} \left( \frac{d_{n,r}}{n} \right) \sum_{k=1}^{K_n} \frac{2^k}{\delta^2} = \lim_{n \rightarrow \infty} \mathcal{O} \left( \frac{d_{n,r}}{n} \frac{2^{K_n}}{\delta^2} \right) = 0.
\end{aligned}$$

All in all, the previous considerations show that the second summand in (30) converges to 0. Since the other terms in (30) can be treated analogously, this finishes the proof of (27).

### 5.1.5 Proof of (28).

Initially, note that

$$\begin{aligned} & \sup_{\substack{|t_2-t_1|<\delta \\ 0\leq t_1, t_2\leq 1}} |m_n(s, t_2) - m_n(s, t_1)| \\ &= \sup_{\substack{|t_2-t_1|<\delta \\ 0\leq t_1, t_2\leq 1}} \left| \frac{1}{\sqrt{n\bar{F}_\varepsilon(u_n)}} \sum_{j=\lfloor nt_1 \rfloor + 1}^{\lfloor nt_2 \rfloor} (\mathbf{1}\{X_j > u_n s\} - \mathbb{E}[\mathbf{1}\{X_j > u_n s\} | \mathcal{F}_{j-1}]) \right|. \end{aligned}$$

As before, we apply a chaining technique in order to prove (28). For this, we define the intervals

$$I_{1,k} := [2k\delta, 2(k+1)\delta] \quad \text{and} \quad I_{2,k} := [(2k+1)\delta, (2(k+1)+1)\delta].$$

for  $k = 0, \dots, L_\delta := \lfloor \frac{1}{2\delta} \rfloor$ .

Then, similarly to (29), it holds that

$$\begin{aligned} & \sup_{1\leq s\leq R} \sup_{\substack{|t_2-t_1|<\delta \\ 0\leq t_1, t_2\leq 1}} |m_n(s, t_2) - m_n(s, t_1)| \\ & \leq \sup_{1\leq s\leq R} \max_{0\leq k\leq L_\delta} \sup_{t_1, t_2 \in I_{1,k}} |m_n(s, t_2) - m_n(s, t_1)| + \sup_{1\leq s\leq R} \max_{0\leq k\leq L_\delta} \sup_{t_1, t_2 \in I_{2,k}} |m_n(s, t_2) - m_n(s, t_1)|. \end{aligned}$$

Again, we restrict our considerations to the first summand and we note that it suffices to show that

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{\delta} \mathbb{P} \left( \sup_{1\leq s\leq R} \sup_{t_2, t_1 \in I_{1,k}} |m_n(s, t_2) - m_n(s, t_1)| > \epsilon \right) = 0.$$

Since, due to stationarity of the data-generating process,

$$\sup_{1\leq s\leq R} \sup_{t_2, t_1 \in I_{1,k}} |m_n(s, t_2) - m_n(s, t_1)| \stackrel{D}{=} \sup_{1\leq s\leq R} \sup_{t_2, t_1 \in I_{1,0}} |m_n(s, t_2) - m_n(s, t_1)|,$$

verification of (28) follows by the same argument as verification (27).  $\square$

## 5.2 Proof of Corollaries 2.7, 2.8, 2.10, and 2.11

*Proof of Corollary 2.7.* An argument from Kulik and Soulier (2011) is repeated and hence many technicalities are omitted. Note that the arguments below are model-free and only use the conclusion of Theorem 2.6.

It holds that

$$\widehat{\gamma}_{\lfloor nt \rfloor} = \frac{A_n(t)}{B_n(t)},$$

where

$$A_n(t) := \frac{1}{n\overline{F}(u_n)} \sum_{j=1}^{\lfloor nt \rfloor} \log \left( \frac{X_j}{u_n} \right) \mathbf{1}\{X_j > u_n\} \quad \text{and} \quad B_n(t) := \frac{1}{n\overline{F}(u_n)} \sum_{j=1}^{\lfloor nt \rfloor} \mathbf{1}\{X_j > u_n\}.$$

By subtracting and adding  $t\alpha^{-1}/B_n(t)$ , the following equality holds:

$$ta_n (\widehat{\gamma}_{\lfloor nt \rfloor} - \alpha^{-1}) = \frac{a_n t}{B_n(t)} (A_n(t) - t\alpha^{-1}) + \frac{a_n t}{\alpha} \left( \frac{t}{B_n(t)} - 1 \right),$$

where  $a_n = n/d_{n,r}$  if  $n/d_{n,r} = o\left(\sqrt{n\overline{F}(u_n)}\right)$  and  $a_n = \sqrt{n\overline{F}(u_n)}$  if  $\sqrt{n\overline{F}(u_n)} = o(n/d_{n,r})$ .

We note that, compared to [Kulik and Soulier \(2011\)](#), the last term appears additionally due to the fact that we consider the two-parameter processes.

Rewriting  $A_n(t) - t\alpha^{-1}$  as an integral and replacing  $\widetilde{T}_n - T$  by  $e_n$ , we have

$$t (\widehat{\gamma}_{\lfloor nt \rfloor} - \alpha^{-1}) = \frac{a_n t}{B_n(t)} \int_1^\infty s^{-1} e_n(s, t) ds - \frac{a_n t}{\alpha B_n(t)} e_n(1, t). \quad (33)$$

We will show weak convergence of the sequence

$$Y_n^{(R)}(t) := \frac{a_n t}{B_n(t)} \int_1^R s^{-1} e_n(s, t) ds - \frac{a_n t}{\alpha B_n(t)} e_n(1, t), \quad n \geq 1, \quad (34)$$

in  $D[t_0, 1]$  by an application of the continuous mapping theorem. For this, we have to initially show that terms of the form  $t/B_n(t)$  are negligible. Noting that  $B_n(t) = \widetilde{T}_n(1, t)$ , it follows by [Theorem 2.6](#) that

$$\sup_{t_0 \leq t \leq 1} \left| t \frac{B_n(1)}{B_n(t)} - 1 \right| \xrightarrow{P} 0. \quad (35)$$

Indeed, [Theorem 2.6](#) implies that  $\sup_{t_0 \leq t \leq 1} |B_n(t) - t| \rightarrow 0$  in probability and

$$\begin{aligned} \sup_{t_0 \leq t \leq 1} \left| t \frac{B_n(1)}{B_n(t)} - 1 \right| &\leq \frac{1}{B_n(t_0)} \sup_{t_0 \leq t \leq 1} |tB_n(1) - B_n(t)| \\ &\leq \frac{1}{B_n(t_0)} \sup_{t_0 \leq t \leq 1} |t(B_n(1) - 1) + t - B_n(t)| \leq \frac{2}{B_n(t_0)} \sup_{t_0 \leq t \leq 1} |B_n(t) - t| \rightarrow 0. \end{aligned}$$

As a consequence, rewriting  $Y_n^{(R)}$  as

$$\begin{aligned} Y_n^{(R)}(t) &= a_n \int_1^R s^{-1} e_n(s, t) ds - \alpha^{-1} a_n e_n(1, t) \\ &\quad + \left( \frac{t}{B_n(t)} - 1 \right) \int_1^R a_n s^{-1} e_n(s, t) ds - \left( \frac{t}{B_n(t)} - 1 \right) \alpha^{-1} a_n e_n(1, t) \quad (36) \end{aligned}$$

and combining Theorem 2.6 with (35) shows that the limit of the sequence  $Y_n^{(R)}, n \geq 1$  corresponds to the limit of

$$Z_n^{(R)} := a_n \int_1^R s^{-1} e_n(s, t) ds - \alpha^{-1} a_n e_n(1, t), \quad n \geq 1. \quad (37)$$

By Theorem 2.6 and the continuous mapping theorem, we conclude that  $Z_n^{(R)}, n \geq 1$ , converges in distribution to  $\int_1^R \xi(s, t) ds - \alpha^{-1} \xi(1, t)$ , where  $\xi(s, t)$  is the limiting process in (11) or (12), respectively. Using the same arguments as in Kulik and Soulier (2011), it can be shown that the convergence of  $Y_n^{(R)}, n \geq 1$ , can be easily extended to convergence of

$$\frac{a_n t}{B_n(t)} \int_1^\infty s^{-1} e_n(s, t) ds - \frac{a_n t}{\alpha B_n(t)} e_n(1, t), \quad n \geq 1. \quad (38)$$

Hence, we conclude that

$$t a_n (\widehat{\gamma}_{[nt]} - \alpha^{-1}) \Rightarrow \int_1^\infty \xi(s, t) ds - \alpha^{-1} \xi(1, t)$$

in  $D[t_0, 1]$ .

If  $\frac{n}{d_{n,r}} = o\left(\sqrt{n\overline{F}(u_n)}\right)$ , separation of the variables  $s$  and  $t$  shows that the limit vanishes. This finishes the proof of Corollary 2.7.  $\square$

*Proof of Corollary 2.8.* Define

$$\widehat{T}_n(s, t) := \frac{1}{[k_n t]} \sum_{j=1}^{[nt]} \mathbf{1}\{X_j > s X_{[nt]:[nt]-[k_n t]}\}.$$

Then, it holds that

$$\int_1^\infty s^{-1} \widehat{T}_n(s, t) ds = \frac{1}{[k_n t]} \sum_{j=1}^{[nt]} \int_1^\infty s^{-1} \mathbf{1}\{X_j > s X_{[nt]:[nt]-[k_n t]}\} ds = \widehat{\gamma}_{\text{Hill}}(t).$$

According to Skorokhod's representation theorem (Theorem 2.3.4 in Shorack and Wellner (1986)) and Theorem 2.6, we may assume without loss of generality that

$$\sup_{s \in [1, \infty], t \in [0, 1]} \left| a_n \left( \frac{1}{n\overline{F}(u_n)} \sum_{j=1}^{[nt]} \mathbf{1}\{X_j > u_n s\} - s^{-\alpha} t \right) - \xi(s, t) \right| \rightarrow 0 \text{ almost surely,}$$

where  $a_n = n/d_{n,r}$  if  $n/d_{n,r} = o\left(\sqrt{n\overline{F}(u_n)}\right)$ ,  $a_n = \sqrt{n\overline{F}(u_n)}$  if  $\sqrt{n\overline{F}(u_n)} = o(n/d_{n,r})$ , and  $\xi$  denotes the corresponding limiting process in Theorem 2.6. In order to apply

Vervaat's Lemma as stated by Lemma 5 in [Einmahl et al. \(2010\)](#), we rephrase the above convergence as:

$$\sup_{s \in [0,1], t \in [t_0,1]} \left| a_n (\Gamma_{n,t}(s) - s) - \frac{1}{t} \xi(s^{-\frac{1}{\alpha}}, t) \right| \longrightarrow 0 \text{ almost surely} \quad (39)$$

for any  $t_0 > 0$  and with

$$\Gamma_{n,t}(s) := \frac{1}{\bar{F}(u_n)} \frac{1}{[nt]} \sum_{j=1}^{[nt]} \mathbf{1} \left\{ X_j > u_n s^{-\frac{1}{\alpha}} \right\}.$$

Choosing  $k_n = n\bar{F}(u_n)$ , it follows that

$$\Gamma_{n,t}^-(s) = (u_n^{-1} X_{[nt]:[nt]-s[k_n t]})^{-\alpha}.$$

As a result, Vervaat's Lemma yields

$$\sup_{s \in [0,1], t \in [t_0,1]} \left| a_n \left( (u_n^{-1} X_{[nt]:[nt]-s[k_n t]})^{-\alpha} - s \right) + \frac{1}{t} \xi(s^{-\frac{1}{\alpha}}, t) \right| \longrightarrow 0 \text{ almost surely.} \quad (40)$$

Setting  $s_n = s u_n^{-1} X_{[nt]:[nt]-[k_n t]}$ , we arrive at

$$\begin{aligned} & \sup_{s \in [1,\infty], t \in [t_0,1]} \left| a_n \left( \frac{1}{n\bar{F}(u_n)} \frac{1}{t} \sum_{j=1}^{[nt]} \mathbf{1} \{ X_j > s X_{[nt]:[nt]-[k_n t]} \} - s^{-\alpha} \right) - \frac{1}{t} (\xi(s, t) - s^{-\alpha} \xi(1, t)) \right| \\ & \leq \sup_{s \in [1,\infty], t \in [t_0,1]} \left| a_n \left( \frac{1}{n\bar{F}(u_n)} \frac{1}{t} \sum_{j=1}^{[nt]} \mathbf{1} \{ X_j > u_n s \} - s^{-\alpha} \right) - \frac{1}{t} \xi(s, t) \right| \\ & \quad + \sup_{s \in [1,\infty], t \in [t_0,1]} \left| a_n (s_n^{-\alpha} - s^{-\alpha}) + s^{-\alpha} \frac{1}{t} \xi(1, t) \right| + \sup_{s \in [1,\infty], t \in [t_0,1]} \left| \frac{1}{t} (\xi(s_n, t) - \xi(s, t)) \right|. \end{aligned}$$

The first two summands on the right-hand side converge to 0 almost surely according to (39) and (40). Moreover, we have  $s_n^{-\alpha} = s^{-\alpha} (1 + o(1))$  a.s. uniformly in  $t$  and  $s$  such that it follows by a continuity argument that the third summand converges to 0 almost surely, as well.

Since  $k_n = n\bar{F}(u_n)$ , we have

$$\begin{aligned} a_n \left( \widehat{T}_n(s, t) - T(s, 1) \right) &= a_n \left( \frac{1}{[k_n t]} \sum_{j=1}^{[nt]} \mathbf{1} \{ X_j > s X_{[nt]:[nt]-[k_n t]} \} - s^{-\alpha} \right) \\ &= a_n \left( \frac{1}{n\bar{F}(u_n)} \frac{1}{t} \sum_{j=1}^{[nt]} \mathbf{1} \{ X_j > s X_{[nt]:[nt]-[k_n t]} \} - s^{-\alpha} \right) + o_P(1) \\ &\Rightarrow \frac{1}{t} (\xi(s, t) - s^{-\alpha} \xi(1, t)). \end{aligned}$$

Similar to the proof of Corollary 2.7, it follows that

$$a_n t (\widehat{\gamma}_{\text{Hill}}(t) - \gamma) = t \int_1^\infty s^{-1} a_n \left( \widehat{T}_n(s, t) - T(s, 1) \right) ds \Rightarrow \int_1^\infty \xi(s, t) ds - \alpha^{-1} \xi(1, t)$$

in  $D[t_0, 1]$ . □

*Proof of Corollaries 2.10 and 2.11.* Since, due to Corollaries 2.7 and 2.8,  $\widehat{\gamma}_n$  and  $\widehat{\gamma}_{\text{Hill}}(1)$  converge to  $\gamma$  in probability, Corollaries 2.10 and 2.11 follow from Slutsky's theorem and an application of the continuous mapping theorem to the processes

$$\begin{aligned} a_n t (\widehat{\gamma}_{[nt]} - \gamma) &\Rightarrow \int_1^\infty s^{-1} \xi(s, t) ds - \alpha^{-1} \xi(1, t), \quad t \in [t_0, 1], \\ a_n t (\widehat{\gamma}_{\text{Hill}}(t) - \gamma) &\Rightarrow \int_1^\infty s^{-1} \xi(s, t) ds - \alpha^{-1} \xi(1, t), \quad t \in [t_0, 1], \end{aligned}$$

and the function

$$f \mapsto \sup_{t \in [t_0, 1]} |f(t) - t f(1)|.$$

□

## 6 Appendix: Second-order regular variation

The following lemmata originate in Kulik and Soulier (2011) and are essential to the proof of Theorem 2.6.

**Lemma 6.1** (Lemma 4.1 in Kulik and Soulier (2011)). *Assume that  $\overline{F}_\varepsilon \in 2RV(\alpha, \eta^*)$ . For positive  $\varepsilon$  there exists a constant  $C$  such that*

$$\forall t \geq 1, \forall z > 0, \quad \left| \frac{\overline{F}_\varepsilon(z t)}{\overline{F}_\varepsilon(t)} - z^{-\alpha} \right| \leq C \eta^*(t) z^{-\alpha-\rho} (\max\{z, z^{-1}\})^\varepsilon. \quad (41)$$

This implies that

$$\sup_{s \geq s_0} \left| \frac{\overline{F}_\varepsilon(u_n s)}{\overline{F}_\varepsilon(u_n)} - s^{-\alpha} \right| = O(\eta^*(u_n)). \quad (42)$$

Using boundedness of  $\eta^*$ , we get the following simplified version of inequality (41):

$$\forall t \geq 1, \forall z > 0, \quad \frac{\overline{F}_\varepsilon(z t)}{\overline{F}_\varepsilon(t)} \leq z^{-\alpha} + C_\varepsilon z^{-\alpha-\rho} (\max\{z, z^{-1}\})^\varepsilon. \quad (43)$$

We also need the following bound on the increments of  $\overline{F}_\varepsilon$ .



**Lemma 6.2** (Lemma 4.2 in [Kulik and Soulier \(2011\)](#)). Assume that  $\bar{F}_\varepsilon \in 2RV(\alpha, \eta^*)$ . For positive  $\varepsilon$  there exists a constant  $C$  such that for all  $t \geq 1$  and  $b > a > 0$

$$\left| \frac{\bar{F}_\varepsilon(at) - \bar{F}_\varepsilon(bt)}{\bar{F}_\varepsilon(t)} - (a^{-\alpha} - b^{-\alpha}) \right| \leq C \eta^*(t) (\min\{a, 1\})^{-\alpha-\rho-\varepsilon} (b-a). \quad (44)$$

Using again boundedness of  $\eta^*$ , we get the following simplified version of inequality (44):

$$\frac{\bar{F}_\varepsilon(at) - \bar{F}_\varepsilon(bt)}{\bar{F}_\varepsilon(t)} \leq a^{-\alpha} - b^{-\alpha} + C_\varepsilon (\min\{a, 1\})^{-\alpha-\rho-\varepsilon} (b-a). \quad (45)$$

## References

- Betken, A. and Kulik, R. (2019). Testing for change in long-memory stochastic volatility time series. *Journal of Time Series Analysis*, 40(5):707 – 738.
- Bilayi-Biakana, C., Ivanoff, G., and Kulik, R. (2019). The tail empirical process for long memory stochastic volatility models with leverage. *Electronic Journal of Statistics*, 13(2):3453 – 3484.
- Breidt, F. J., Crato, N., and de Lima, P. (1998). The detection and estimation of long memory in stochastic volatility. *Journal of Econometrics*, 83(1 – 2):325 – 348.
- Breiman, L. (1965). On some limit theorems similar to the arc-sin law. *Teoriya Veroyatnostei i ee Primeneniya*, 10:351 – 360.
- Cont, R. (2005). Long range dependence in financial markets. In *Fractals in engineering*, pages 159 – 179. Springer.
- Deo, R., Hsieh, M., Hurvich, C. M., and Soulier, P. (2006). Long memory in nonlinear processes. In *Dependence in probability and statistics*, volume 187 of *Lecture Notes in Statistics*, pages 221 – 244. Springer, New York.
- Drees, H. (1998a). A general class of estimators of the extreme value index. *Journal of Statistical Planning and Inference*, 66(1):95 – 112.
- Drees, H. (1998b). On smooth statistical tail functionals. *Scandinavian Journal of Statistics*, 25:187 – 210.

- Drees, H. (2000). Weighted approximations of tail processes for  $\beta$ -mixing random variables. *Annals of Applied Probability*, 10(4):1274 – 1301.
- DuMouchel, W. H. (1983). Estimating the stable index  $\alpha$  in order to measure tail thickness: a critique. *Annals of Statistics*, 11(4):1019 – 1031.
- Einmahl, J. H. J. (1990). The empirical distribution function as a tail estimator. *Statistica Neerlandica*, 44(2):79 – 82.
- Einmahl, J. H. J. (1992). Limit theorems for tail processes with application to intermediate quantile estimation. *Journal of Statistical Planning and Inference*, 32(1):137 – 145.
- Einmahl, J. H. J., Gantner, M., and Sawitzki, G. (2010). Asymptotics of the shorth plot. *Journal of Statistical Planning and Inference*, 140(11):3003 – 3012.
- Freedman, D. A. (1975). On tail probabilities for martingales. *The Annals of Probability*, pages 100 – 118.
- Galbraith, J. W. and Zernov, S. (2004). Circuit breakers and the tail index of equity returns. *Journal of Financial Econometrics*, 2(1):109 – 129.
- Hall, P. (1982). On some simple estimates of an exponent of regular variation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 44(1):37 – 42.
- Harvey, A. C. (2002). Long memory in stochastic volatility. In *Forecasting Volatility in the Financial Markets*, pages 307 – 320. Butterworth-Heinemann Finance.
- Hill, B. M. (1975). A simple general approach to inference about the tail of a distribution. *Annals of Statistics*, 3(5):1163 – 1174.
- Hoga, Y. (2017). Change point tests for the tail index of  $\beta$ -mixing random variables. *Econometric Theory*, 33(4):915 – 954.
- Hurvich, C. M. and Soulier, P. (2009). *Stochastic Volatility Models with Long Memory*, pages 345 – 354. Springer.
- Kim, M. and Lee, S. (2011). Change point test for tail index for dependent data. *Metrika*, 74(3):297 – 311.

- Kim, M. and Lee, S. (2012). Change point test of tail index for autoregressive processes. *Journal of the Korean Statistical Society*, 41(3):305 – 312.
- Koedijk, K. G., Schafgans, M. M., and De Vries, C. G. (1990). The tail index of exchange rate returns. *Journal of International Economics*, 29(1-2):93 – 108.
- Kulik, R. and Soulier, P. (2011). The tail empirical process for long memory stochastic volatility sequences. *Stochastic Processes and their Applications*, 121(1):109 – 134.
- Mandelbrot, B. B. (1963). The Variation of Certain Speculative Prices. *Business*, 36:394 – 419.
- Mason, D. M. (1988). A strong invariance theorem for the tail empirical process. In *Annales de l'IHP Probabilités et statistiques*, volume 24, pages 491 – 506.
- Phillips, P. C., Loretan, M., et al. (1990). Testing covariance stationarity under moment condition failure with an application to common stock returns. Technical report, Cowles Foundation for Research in Economics, Yale University.
- Pipiras, V. and Taqqu, M. S. (2017). *Long-Range Dependence and Self-Similarity*, volume 45. Cambridge University Press.
- Pollard, D. (1984). *Convergence of stochastic processes*. Springer Series in Statistics. Springer-Verlag, New York.
- Quintos, C., Fan, Z., and Phillips, P. (2001). Structural change tests in tail behaviour and the Asian crisis. *The Review of Economic Studies*, 68(3):633 – 663.
- Resnick, S. I. and Stărică, C. (1997). Asymptotic behavior of hill's estimator for autoregressive data. *Communications in Statistics. Stochastic Models*, 13(4):703 – 721. Heavy tails and highly volatile phenomena.
- Rootzén, H. (2009). Weak convergence of the tail empirical process for dependent sequences. *Stochastic Processes and their Applications*, 119(2):468 – 490.
- Shorack, G. R. and Wellner, J. A. (1986). *Empirical processes with applications to statistics*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York.

Taqqu, M. S. (1979). Zeitschrift für wahrscheinlichkeitstheorie und verwandte gebiete. *Zeitschrift*, 50(1):53 – 83.

Taylor, S. J. (1986). *Modelling Financial Time Series*. Wiley, New York.

Werner, T. and Upper, C. (2004). Time variation in the tail behavior of Bund future returns. *Journal of Futures Markets: Futures, Options, and Other Derivative Products*, 24(4):387 – 398.