
ASYMPTOTIC OPTIMALITY OF CONDITIONED STOCHASTIC GRADIENT DESCENT

Rémi Leluc
 LTCI, Télécom Paris
 Institut polytechnique de Paris
 91120 Palaiseau, France
 remi.leluc@gmail.com

Francois Portier
 LTCI, Télécom Paris
 Institut polytechnique de Paris
 91120 Palaiseau, France
 francois.portier@gmail.com

January 18, 2022

ABSTRACT

In this paper, we investigate a general class of stochastic gradient descent (SGD) algorithms, called conditioned SGD, based on a preconditioning of the gradient direction. Under some mild assumptions, namely the L -smoothness of the non-convex objective function and some weak growth condition on the noise, we establish the almost sure convergence and the asymptotic normality for a broad class of conditioning matrices. In particular, when the conditioning matrix is an estimate of the inverse Hessian at the optimal point, the algorithm is proved to be asymptotically optimal. The benefits of this approach are validated on simulated and real datasets.

1 Introduction

Consider the following type of unconstrained optimization problem: $\min_{\theta \in \mathbb{R}^d} \{F(\theta) = \mathbb{E}_{\xi} [f(\theta, \xi)]\}$, where f is a loss function and ξ is a random variable. This key methodological problem, known under the name of *stochastic programming* (Shapiro et al., 2014), includes many flagship machine learning applications such as *empirical risk minimization* (Bottou et al., 2018), *adaptive importance sampling* (Delyon and Portier, 2018) and *reinforcement learning* (Sutton and Barto, 2018). When F is differentiable, a common approach is to rely on first-order methods. However, in many scenarios and particularly in large-scale learning, the gradient of F may be hard to evaluate or even intractable. Instead, a random unbiased estimate of the gradient is available at a cheap computing cost and the state-of-the-art algorithm, *stochastic gradient descent* (SGD), just moves along this estimate at each iteration. It is an iterative algorithm, simple and computationally fast, but its convergence towards the optimum is generally slow. *Conditioned* SGD, which consists in multiplying the gradient estimate by some conditioning matrix at each iteration, can lead to better results in practice as shown in several recent studies (Amari, 1998; Kakade, 2002; Byrd et al., 2011; Duchi et al., 2011; Gower et al., 2020).

Conditioned SGD generalizes *standard SGD* by adding a conditioning step to refine the descent direction. Starting from $\theta_0 \in \mathbb{R}^d$, the algorithm of interest is defined by the following iteration

$$\theta_{k+1} = \theta_k - \alpha_{k+1} C_k g(\theta_k, \xi_{k+1}), \quad k \geq 0,$$

where $g(\theta_k, \xi_{k+1})$ is some unbiased gradient valued in \mathbb{R}^d , $C_k \in \mathbb{R}^{d \times d}$ is called *conditioning matrix* and $(\alpha_k)_{k \geq 1}$ is some learning rate sequence that should decrease throughout the algorithm.

Interestingly, the optimal choice according to the asymptotic variance is the inverse of the Hessian matrix at optimal point, i.e., $C_k = \nabla^2 F(\theta^*)^{-1}$; see (Benveniste et al., 2012, Chapter 3) or Section 4 in this paper. When using such a conditioning matrix, the rate of convergence remains the same and only

the asymptotic variance can be reduced; e.g., [Agarwal et al. \(2009\)](#). An important question, which is still open to the best of our knowledge, is whether the optimal variance can be achieved by such an algorithm for non-convex objective F . We show that the answer is positive under mild conditions on the matrix C_k .

Related work. Early seminal works around standard SGD (when C_k is the identity matrix) are part of the *stochastic approximation* literature initiated by [Robbins and Monro \(1951\)](#) and [Kiefer et al. \(1952\)](#); the almost sure convergence is studied in [Robbins and Siegmund \(1971\)](#) and [Bertsekas and Tsitsiklis \(2000\)](#); rates of convergence are investigated in [Kushner and Huang \(1979\)](#) and [Pelletier \(1998a\)](#); non-asymptotic bounds are given in [Moulines and Bach \(2011\)](#). The asymptotic normality is usually obtained using two different approaches: a diffusion-based method is employed in [Pelletier \(1998b\)](#) and [Benaïm \(1999\)](#) whereas martingale tools are used in [Sacks \(1958\)](#) and [Kushner and Clark \(1978\)](#). We refer to [Nevelson and Khas'minskiĭ \(1976\)](#); [Delyon \(1996\)](#); [Benveniste et al. \(2012\)](#); [Duflo \(2013\)](#) for general textbooks on *stochastic approximation*.

The aforementioned results do not apply directly to conditioned SGD because of the presence of the matrix sequence $(C_k)_{k \geq 0}$ involving an additional source of randomness in the algorithm. Seminal papers dealing with the weak convergence of conditioned SGD are [Venter \(1967\)](#) and [Fabian \(1968\)](#). Within a restrictive framework (univariate case $d = 1$ and strong assumptions on the function F), their results are encouraging because the limiting variance of the procedure is shown to be smaller than the limiting variance of standard SGD. Venter's and Fabian's results have then been extended to more general situations ([Fabian, 1973](#); [Nevelson and Khas'minskiĭ, 1976](#); [Wei, 1987](#)). In [Wei \(1987\)](#), the framework is still restrictive not only because the random errors are assumed to be independent and identically distributed but also because the objective F must satisfy their assumption (4.10) which hardly extends to objectives other than quadratic. More recently, [Bercu et al. \(2020\)](#) have obtained the asymptotic normality as well as the efficiency of certain conditioned SGD estimates in the particular case of *logistic regression*. The previous approach has been generalized not long ago in [Boyer and Godichon-Baggioni \(2020\)](#) where the use of the Woodbury matrix identity is promoted to compute the Hessian inverse. Several theoretical results, including the weak convergence of conditioned SGD, are obtained for convex objective functions. An alternative to *conditioning*, called *averaging*, developed by [Polyak \(1990\)](#) and [Polyak and Juditsky \(1992\)](#), allows to recover the same asymptotic variance as *conditioned SGD*. When dealing with convex objectives, the theory behind this averaging technique is a well-studied topic ([Moulines and Bach, 2011](#); [Gadat and Panloup, 2017](#); [Dieuleveut et al., 2020](#)). However, it is inevitably associated with a large bias caused by poor initialization. Furthermore, *conditioned SGD* methods proved to be the current state-of-the-art for training machine learning models ([Zhang, 2004](#); [LeCun et al., 2012](#)) and are implemented in widely used programming tools ([Pedregosa et al., 2011](#); [Abadi et al., 2016](#)).

Contributions. There are two key results established in this paper. A first result provides conditions on the sequence $(C_k)_{k \geq 0}$ for the almost sure convergence of the conditioned SGD. A second result (our main contribution) deals with the asymptotic normality of the rescaled sequence of iterates. The working conditions are relatively weak as the function F is only required to be L -smooth, which is classical in general non-convex learning and the variance noise of the gradient might be unbounded using a growth condition related to *expected smoothness* ([Gower et al., 2019](#)). Interestingly, our asymptotic normality result consists of the following continuity property: whenever the matrix C_k converges to C , the algorithm behaves in the same way as an oracle version in which C would be used instead of C_k . We stress that contrary to [Boyer and Godichon-Baggioni \(2020\)](#), no convexity assumption is needed on the objective function and no rate of convergence is required on the sequence $(C_k)_{k \geq 0}$. This is important because, in most cases, deriving a convergence rate on $(C_k)_{k \geq 0}$ requires a specific convergence rate on the iterates $(\theta_k)_{k \geq 0}$ and this last point cannot be directly inferred from the almost sure convergence. Finally, for illustrative purposes, we provide a feasible algorithm that achieve minimum variance and that is considered in a numerical study.

To obtain these results, instead of approximating the rescaled sequence of iterates by a continuous diffusion (as for instance in [Pelletier \(1998b\)](#)), we rely on a discrete-time approach where the recursion scheme is directly analyzed (as for instance in [Delyon \(1996\)](#)). More precisely, the sequence of iterates is studied with the help of an auxiliary linear algorithm whose limiting distribution can be deduced from

the central limit theorem for martingale increments (Hall and Heyde, 1980). The limiting variance is derived from a discrete time matrix-valued dynamical system algorithm. It corresponds to the solution of a Lyapunov equation involving the matrix C . It allows a special choice for C which guarantees an optimal variance. Finally, in order to examine the remaining part, a particular recursion is identified. By studying it on a particular event, we show that this remaining part is negligible. The point of view adopted in this paper regarding the theoretical analysis is the one of *weak convergence*. It is an appropriate tool to deal with efficiency issues because algorithms can be compared easily through their asymptotic variances.

Outline. In Section 2, the mathematical framework is introduced. In Section 3, an asymptotic study of standard SGD is given. Section 4 is dedicated to conditioned SGD and contains our main results. Finally, numerical experiments are performed in Section 5 for illustrative purposes.

2 Mathematical background

In this Section, the mathematical background of stochastic gradient descent (SGD) methods is presented and motivated with the help of some examples.

Consider the problem of finding the minimizer $\theta^* \in \mathbb{R}^d$ of a function $F : \mathbb{R}^d \rightarrow \mathbb{R}$, that is,

$$\theta^* = \arg \min_{\theta \in \mathbb{R}^d} F(\theta).$$

In many scenarios, the gradient of F cannot be fully computed and only a stochastic unbiased version of it is available. The SGD algorithm moves the iterate along this direction. To increase the efficiency, the random generators used to derive the unbiased gradients might evolve during the algorithm, *e.g.*, using the past iterations. To analyse such algorithms, we consider the following probabilistic setting.

Definition 1. A stochastic algorithm is a sequence $(\theta_k)_{k \geq 0}$ of random variables defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and valued in \mathbb{R}^d . Define $(\mathcal{F}_k)_{k \geq 0}$ as the natural σ -field associated to the stochastic algorithm $(\theta_k)_{k \geq 0}$, *i.e.*, $\mathcal{F}_k = \sigma(\theta_0, \theta_1, \dots, \theta_k)$, $k \geq 0$. A policy is a sequence of random probability measures $(P_k)_{k \geq 0}$, each defined on a measurable space (S, \mathcal{S}) that are adapted to \mathcal{F}_k .

Given a policy $(P_k)_{k \geq 0}$ and a learning rates sequence $(\alpha_k)_{k \geq 1}$ of positive numbers, the SGD algorithm (Robbins and Monro, 1951) is defined by the update rule

$$\theta_{k+1} = \theta_k - \alpha_{k+1} g(\theta_k, \xi_{k+1}) \quad \text{where} \quad \xi_{k+1} \sim P_k, \quad (1)$$

with $g : \mathbb{R}^d \times S \rightarrow \mathbb{R}^d$. Hence the policy $(P_k)_{k \geq 0}$ is used at each iteration to produce random gradients through the function g . Those gradients are assumed to be unbiased.

Assumption 1 (Unbiased gradient). *The gradient generator $g : \mathbb{R}^d \times S \rightarrow \mathbb{R}^d$ is such that for all $\theta \in \mathbb{R}^d$, $g(\theta, \cdot)$ is measurable, and*

$$\forall k \geq 0, \quad \mathbb{E}[g(\theta_k, \xi_{k+1}) | \mathcal{F}_k] = \nabla F(\theta_k).$$

In this paper, we consider the *decreasing step-size* scheme where the learning rates sequence should go to 0 according to the following Robbins-Monro condition (Robbins and Monro, 1951). This condition is meant to eventually anneal the noise but not too fast so that the iterates $(\theta_k)_{k \geq 0}$ can reach interesting places in a finite time.

Assumption 2 (Robbins-Monro). *The sequence $(\alpha_k)_{k \geq 1}$ is positive decreasing to 0 and satisfies the Robbins-Monro condition: $\sum_{k \geq 1} \alpha_k = +\infty$ and $\sum_{k \geq 1} \alpha_k^2 < +\infty$.*

We emphasize three important examples covered by the developed approach. In each case, we provide explicit ways to generate the stochastic gradient.

Example 1. (*Empirical Risk Minimization*) Given some observed data $z_1, \dots, z_n \in \mathbb{R}^p$ and a differentiable loss function $\ell : \mathbb{R}^d \times \mathbb{R}^p \rightarrow \mathbb{R}$, the objective function F approximates the true risk

$\mathbb{E}_z[\ell(\theta, z)]$ using its empirical counterpart

$$F(\theta) = \frac{1}{n} \sum_{i=1}^n \ell(\theta, z_i).$$

Classically, the gradient estimates at θ_k are given by the policy

$$g(\theta_k, \xi_{k+1}) = \nabla_{\theta} \ell(\theta_k, \xi_{k+1}) \quad \text{where} \quad \xi_{k+1} \sim \sum_{i=1}^n \delta_{z_i} / n.$$

Another one, more subtle, referred to as mini-batching (Gower et al., 2019), consists in generating uniformly a set of n_k samples (z_1, \dots, z_{n_k}) and computing the gradient as the average $n_k^{-1} \sum_{j=1}^{n_k} \nabla_{\theta} \ell(\theta_k, z_j)$. Note that interestingly, we allow changes of the minibatch size throughout the algorithm. Our framework also includes adaptive non-uniform sampling (Papa et al., 2015) and survey sampling (Cl  men  on et al., 2019), which use $P_k = \sum_{i=1}^n w_i^{(k)} \delta_{z_i}$ with \mathcal{F}_k -adapted weights satisfying $\sum_{i=1}^n w_i^{(k)} = 1$ for each $k \geq 0$.

Example 2. (*Adaptive importance sampling*) Given a target function f , which might result from the likelihood of some data, and a parametric family of sampler $\{q_{\theta} : \theta \in \Theta\}$, the objective function is

$$F(\theta) = - \int \log(q_{\theta}(y)) f(y) dy.$$

Other losses can be considered and we refer to (Delyon and Portier, 2018) for some details and further references about adaptive importance sampling. A common choice in practice for the policy is given by

$$g(\theta_k, \xi_{k+1}) = -\nabla_{\theta} \log(q_{\theta_k}(\xi_{k+1})) \frac{f(\xi_{k+1})}{q_{\theta_k}(\xi_{k+1})} \quad \text{where} \quad \xi_{k+1} \sim q_{\theta_k}.$$

Example 3. (*Policy-gradient methods*) In reinforcement learning (Sutton and Barto, 2018), the goal of the agent is to find the best action-selection policy to maximize the expected reward. Policy-gradient methods (Baxter and Bartlett, 2001; Williams, 1992) use a parameterized policy $\{\pi_{\theta} : \theta \in \Theta\}$ to optimize an expected reward function F given by

$$F(\theta) = \mathbb{E}_{\xi \sim \pi_{\theta}} [\mathcal{R}(\xi)],$$

where ξ is a trajectory including nature states and selected actions. Using the policy gradient theorem, one has $\nabla F(\theta) = \mathbb{E}_{\xi \sim \pi_{\theta}} [\mathcal{R}(\xi) \nabla_{\theta} \log \pi_{\theta}(\xi)]$, leading to the algorithm REINFORCE (Williams, 1992) given by

$$g(\theta_k, \xi_{k+1}) = \mathcal{R}(\xi_{k+1}) \nabla_{\theta} \log \pi_{\theta_k}(\xi_{k+1}) \quad \text{where} \quad \xi_{k+1} \sim \pi_{\theta_k}.$$

3 The asymptotics of stochastic gradient descent

In this Section, we gradually introduce the working assumptions along with some discussions. We provide an asymptotic analysis of SGD including the almost sure convergence, the asymptotic normality and a discussion of several references. Even if the considered approach brings little novelty compared to existing work, the results of the Section serve essentially as a starting point for further comparison with conditioned SGD.

To perform the asymptotic analysis of stochastic gradient methods, one needs to make assumptions on the objective function (Assumptions 3 and 4) and on the stochastic perturbation (Assumption 5). Although stochastic gradient descent is often associated with convex optimization, the analysis presented here is only based on the following standard notion of smoothness for F .

Assumption 3 (L-smooth). *The objective function $F : \mathbb{R}^d \rightarrow \mathbb{R}$ is continuously differentiable and the gradient function $\nabla F : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is Lipschitz continuous with Lipschitz constant $L > 0$, i.e.*

$$\forall \theta, \eta \in \mathbb{R}^d, \quad \|\nabla F(\theta) - \nabla F(\eta)\|_2 \leq L \|\theta - \eta\|_2.$$

The following identifiability condition shall be useful to recover a unique argument of the minimum.

Assumption 4 (Identifiability). F is coercive, i.e., $\lim_{\|\theta\|_2 \rightarrow +\infty} F(\theta) = +\infty$ and the equation $\nabla F(\theta) = 0$ has a unique solution θ^* .

Note that the two previous assumptions together imply that θ^* is the unique minimizer of F . With some additional work, the previous condition could be refined assuming instead that the level sets of stationary points $\{\theta, \nabla F(\theta) = 0\} \cap \{\theta, F(\theta) = y\}$ are locally finite for every $y \in \mathbb{R}^d$ (Gadat et al., 2018).

To handle the stochastic noise associated to the gradient estimates, we consider a relatively weak growth condition, related to the notion of *expected smoothness* as introduced in Gower et al. (2019) (see also Gazagnadou et al. (2019); Gower et al. (2020)). In particular, we extend the condition of Gower et al. (2019) to our general context in which the sampling distributions are allowed to change along the algorithm.

Assumption 5 (Growth condition). With probability 1, there exist $0 \leq \mathcal{L}, \sigma^2 < \infty$ such that

$$\forall \theta \in \mathbb{R}^d, \forall k \in \mathbb{N}, \quad \mathbb{E} [\|g(\theta, \xi_{k+1})\|_2^2 | \mathcal{F}_k] \leq 2\mathcal{L} (F(\theta) - F(\theta^*)) + \sigma^2.$$

This almost-sure bound on the stochastic noise $\mathbb{E} [\|g(\theta, \xi_k)\|_2^2 | \mathcal{F}_{k-1}]$ is the key to prove the almost sure convergence of the algorithm.

Note that Assumption 5, often referred to as a *growth condition*, is mild since it allows the noise to be large when the iterate is far away from the optimal point. In that aspect, it contrasts with uniform bounds of the form $\mathbb{E} [\|g(\theta_k, \xi_{k+1})\|_2^2 | \mathcal{F}_k] \leq \sigma^2$ for some deterministic $\sigma^2 > 0$ (see Nemirovski et al. (2009); Nemirovski and Yudin (1983); Shalev-Shwartz et al. (2011)). Observe that such uniform bound is recovered by taking $\mathcal{L} = 0$ in Assumption 5 but cannot hold when the objective function F is strongly convex (Nguyen et al., 2018). Besides, fast convergence rates have been derived in Schmidt and Roux (2013) under the *strong-growth condition*: $\mathbb{E} [\|g(\theta, \xi_{k+1})\|_2^2 | \mathcal{F}_k] \leq M \|\nabla F(\theta)\|_2^2$ for some $M > 0$. Similarly to our growth condition, Bertsekas and Tsitsiklis (2000) and Bottou et al. (2018) performed an analysis under the condition $\mathbb{E} [\|g(\theta, \xi_{k+1})\|_2^2 | \mathcal{F}_k] \leq M \|\nabla F(\theta)\|_2^2 + \sigma^2$ for $M, \sigma^2 > 0$. Under Assumptions 3 and 4, we have $\|\nabla F(\theta)\|_2^2 \leq 2\mathcal{L} (F(\theta) - F(\theta^*))$ (Gower et al., 2019, Proposition A.1) so our growth condition is less restrictive. If F satisfies the Polyak-Lojasiewicz condition (Karimi et al., 2016), then our growth condition becomes a bit stronger. Another weak growth condition has been used for a non-asymptotic study in Moulines and Bach (2011).

The weak growth condition on the stochastic noise is general and can be achieved in practice with the following Lemma, whose proof can be found in the appendix.

Lemma 1. Suppose that for all $k \geq 0, \theta \in \mathbb{R}^d, F(\theta) = \mathbb{E} [f(\theta, \xi_{k+1}) | \mathcal{F}_k]$ with $\xi_{k+1} \sim P_k$. Assume that, with probability 1, the function $\theta \mapsto f(\theta, \xi_{k+1})$ is L -smooth and there exists $m \in \mathbb{R}$ such that for all $\theta \in \mathbb{R}^d, f(\theta, \xi_{k+1}) \geq m$. Then the growth condition of Assumption 5 is satisfied.

The following proposition reveals that all these assumptions are sufficient to ensure the almost sure convergence of the algorithm.

Theorem 1 (Almost sure convergence). Suppose that Assumptions 1 to 5 are fulfilled. Then the sequence of iterates $(\theta_k)_{k \geq 0}$ obtained by the SGD rule (1) converges almost surely towards the minimizer $\theta_k \rightarrow \theta^*$.

Compared to (Nguyen et al., 2018, Theorem 1 and 3), our conditions are not concerned with the whole sequence of learning rates but only with their limiting behavior. The proof of Theorem 1 follows from an application of Theorem 3 - one of our main results given in Section 4 - which covers the use of a conditioning matrix.

Since the almost sure convergence of stochastic gradient descent has been verified, we can now consider the convergence rate at which the sequence $(\theta_k - \theta^*)$ remains bounded. Under a few additional assumptions, the SGD iterates are asymptotically normal with convergence rate $\sqrt{\alpha_k}$. First let us be more specific about the choice of the learning rates sequence.

Assumption 6 (Learning rates). The sequence of step-size is equal to $\alpha_k = \alpha k^{-\beta}$ with $\beta \in (1/2, 1]$.

There is also a need for stability around the minimizer which translates into an assumption on the Hessian matrix at optimal point. Such a matrix is assumed to be well-defined. Let $\mathcal{S}_d^{++}(\mathbb{R})$ denote the space of real symmetric positive definite matrices.

Assumption 7 (Hessian). *The Hessian matrix at optimal point is positive definite, i.e., $H = \nabla^2 F(\theta^*) \in \mathcal{S}_d^{++}(\mathbb{R})$ and the mapping $\theta \mapsto \nabla^2 F(\theta)$ is continuous at θ^* .*

In addition, to identify the limiting covariance of the asymptotic regime, the conditional covariance matrices of $g(\theta_k, \xi_{k+1})$ are assumed to converge. To this aim, define for all $k \geq 0$,

$$\begin{aligned} w_{k+1} &= \nabla F(\theta_k) - g(\theta_k, \xi_{k+1}) \\ \Gamma_k &= \mathbb{E} [w_{k+1} w_{k+1}^T | \mathcal{F}_k]. \end{aligned}$$

Assumption 8 (Covariance matrix). *There exists $\Gamma \in \mathcal{S}_d^{++}(\mathbb{R})$ such that almost surely $\Gamma_k \xrightarrow[k \rightarrow +\infty]{} \Gamma$.*

Finally, in order to derive a central limit theorem for the iterates of the algorithm, there is an extra need for stability which is synonymous with a uniform bound on the noise around the minimizer.

Assumption 9 (Lyapunov bound). *There exist $\delta, \varepsilon > 0$ such that almost surely*

$$\sup_{k \geq 0} \mathbb{E} [\|w_{k+1}\|_2^{2+\delta} | \mathcal{F}_k] \mathbf{1}_{\{\|\theta_k - \theta^*\| \leq \varepsilon\}} < \infty.$$

The following asymptotic normality result can be either derived from (Pelletier, 1998b, Theorem 1) or as a direct corollary of our main result, Theorem 4, given in Section 4.

Theorem 2 (Weak convergence). *Suppose that Assumptions 1 to 9 are fulfilled. Assume that $(H - \zeta I)$ is positive definite where $\zeta = \mathbf{1}_{\{\beta=1\}}/2\alpha$. Let $(\theta_k)_{k \geq 0}$ be obtained by the SGD rule (1) then*

$$\frac{1}{\sqrt{\alpha_k}} (\theta_k - \theta^*) \rightsquigarrow \mathcal{N}(0, \Sigma), \quad \text{as } k \rightarrow \infty,$$

where the covariance matrix Σ satisfies the following Lyapunov equation

$$(H - \zeta I_d) \Sigma + \Sigma (H - \zeta I_d)^T = \Gamma.$$

Several remarks are to be explored. First, since Γ and $(H - \zeta I)$ are positive definite matrices, there exists a unique solution Σ to the Lyapunov equation $(H - \zeta I_d) \Sigma + \Sigma (H - \zeta I_d)^T = \Gamma$ given by $\Sigma = \int_0^{+\infty} \exp[-t(H - \zeta I_d)] \Gamma \exp[-t(H - \zeta I_d)^T] dt$. Second, the previous result can be expressed as $k^{\beta/2} (\theta_k - \theta^*) \rightsquigarrow \mathcal{N}(0, \alpha \Sigma)$. Hence, the fastest rate of convergence is obtained when $\beta = 1$ for which we recover the classical $1/\sqrt{k}$ -rate of a Monte-Carlo estimate. In this case, the coefficient α should be chosen large enough to ensure the convergence through the condition $H - I_d/(2\alpha) \succ 0$, but also such that the covariance matrix $\alpha \Sigma$ is small. The choice of α is discussed in the next Section and should be replaced with a matrix gain.

4 Conditioned stochastic gradient descent

4.1 Minimum variance

To motivate the use of conditioning matrices in SGD, we raise the question of variance optimality when α_k decreases as $1/k$, so the rate of convergence in Theorem 2 is optimal, and the scalar gain α is replaced by a conditioning matrix $C \in \mathcal{S}_d^{++}(\mathbb{R})$. That is, we consider the iteration scheme, for $k \geq 1$,

$$\theta_{k+1} = \theta_k - \left(\frac{C}{k+1} \right) g(\theta_k, \xi_{k+1}). \quad (2)$$

As a corollary of Theorem 4 (given below) or inferring from the results in Pelletier (1998b), we can derive the following result. Define \mathcal{C}_H as the set of real symmetric positive definite matrices $C \in \mathcal{S}_d^{++}(\mathbb{R})$ such that all the eigenvalues of the matrix $CH - (I/2)$ are positive.

Proposition 1. *Suppose that Assumptions 1, 3, 4, 5, 7, 8 and 9 are fulfilled. Let $(\theta_k)_{k \geq 0}$ be obtained by (2) with $C \in \mathcal{C}_H$. Then we have*

$$\sqrt{k}(\theta_k - \theta^*) \rightsquigarrow \mathcal{N}(0, \Sigma_C),$$

where Σ_C satisfies:

$$(CH - I_d/2)\Sigma_C + \Sigma_C(CH - I_d/2)^T = C\Gamma C^T. \quad (3)$$

The best conditioning matrix C that could be chosen regarding the asymptotic variance is specified in the next proposition whose proof is given in the appendix.

Proposition 2 (Optimal choice). *The choice $C^* = H^{-1}$ is optimal in the sense that $\Sigma_{C^*} \preceq \Sigma_C$ for all $C \in \mathcal{C}_H$. Moreover, we have $\Sigma_{C^*} = H^{-1}\Gamma H^{-1}$.*

In deterministic gradient descent, it is well-known that the rate of convergence is improved when the gradient is multiplied by the inverse of the Hessian matrix, referred to as the Newton algorithm, whose convergence rate is quadratic, instead of linear for gradient descent. Due to Proposition 2 where we have seen that the smallest limiting variance is nonzero, a faster rate of convergence cannot be expected with SGD. However, an improvement in the limiting variance is still possible.

Other approaches to improve standard SGD include adaptive regularization (Duchi et al., 2011), variance reduction (Defazio et al., 2014; Johnson and Zhang, 2013), dual coordinate ascent (Shalev-Shwartz and Zhang, 2013), and averaging (Polyak, 1990; Polyak and Juditsky, 1992; Delyon and Juditsky, 1992). Note that Polyak averaging has the same limit as the optimal one for conditioned SGD but the method is known to suffer from a difficult initialization and large bias (see Figures in Section 5).

4.2 General results

We now introduce the general framework of conditioned SGD as an extension of the standard SGD presented in Section 2. Conditioned SGD is defined by the following update rule, for $k \geq 0$,

$$\theta_{k+1} = \theta_k - \alpha_{k+1} C_k g(\theta_k, \xi_{k+1}), \quad (4)$$

where the matrix $C_k \in \mathcal{S}_d^{++}(\mathbb{R})$, the *conditioning matrix*, is a \mathcal{F}_k -measurable real symmetric positive definite matrix that refines the conditioning of the gradient estimate. The success of the proposed approach relies on the following condition which ensures a suitable control on the eigenvalues of the conditioning matrices.

Assumption 10 (Eigenvalues and learning rates). *Let $(\beta_k)_{k \geq 1}$ and $(\gamma_k)_{k \geq 1}$ be such that*

$$\forall k \geq 1, \quad \beta_k I_d \preceq C_{k-1} \preceq \gamma_k I_d.$$

The sequences $(\alpha_k)_{k \geq 1}, (\beta_k)_{k \geq 1}, (\gamma_k)_{k \geq 1}$ are positive and satisfy the following conditions $\sum_{k \geq 1} \alpha_k \beta_k = +\infty$ a.s. and $\sum_{k \geq 1} (\alpha_k \gamma_k)^2 < +\infty$ a.s.

Our first main result, whose proof is in the appendix, provides conditions for the almost sure convergence of conditioned SGD.

Theorem 3 (Almost sure convergence). *Suppose that Assumptions 1, 3, 4, 5 and 10 are fulfilled. Then the sequence of iterates $(\theta_k)_{k \geq 0}$ obtained by the conditioned SGD (4) converges almost surely towards the minimizer $\theta_k \rightarrow \theta^*$.*

Note that compared to the conditions of Theorem 1, only the standard Robbins-Monro condition, Assumption 2, has been replaced by Assumption 10. To recover the setting of classical SGD, one can simply choose $C_k = I_d$, in which case, Assumption 10 meets Assumption 2. This makes Theorem 1 a particular case of our result.

Following the same path as in the SGD Section, it is now interesting to search for an appropriate rescaled process to obtain some convergence rate and asymptotic normality results. In fact the only additional assumption we need, compared to SGD, is the convergence of the sequence $(C_k)_{k \geq 0}$ towards a symmetric positive definite matrix.

Assumption 11 (Convergence of the conditioning matrix). *There exists $C \in \mathcal{S}_d^{++}(\mathbb{R})$ such that almost surely $C_k \xrightarrow{k \rightarrow +\infty} C$.*

Theorem 4 (Weak convergence). *Suppose that Assumptions 1 and 3 to 11 are fulfilled. Assume that all the eigenvalues of $(CH - \zeta I)$ are positive where $\zeta = \mathbb{1}_{\{\beta=1\}}/2\alpha$. Let $(\theta_k)_{k \geq 0}$ be obtained by conditioned SGD (4) then*

$$\frac{1}{\sqrt{\alpha_k}}(\theta_k - \theta^*) \rightsquigarrow \mathcal{N}(0, \Sigma_C), \quad \text{as } k \rightarrow \infty,$$

where Σ_C satisfies:

$$(CH - \zeta I_d) \Sigma_C + \Sigma_C (CH - \zeta I_d)^T = C \Gamma C^T.$$

Sketch of the proof. In a similar spirit as in Delyon (1996), the proof is based on the approximation $\nabla F(\theta_k) = \nabla F(\theta^*) + H(\theta_k - \theta^*) + o(\theta_k - \theta^*) \simeq H(\theta_k - \theta^*)$ and relies on the introduction of a linear stochastic algorithm. Avoiding some technicalities related to the introduction of some event, we introduce the matrix $K = CH$ along with the iteration

$$\tilde{\Delta}_{k+1} = \tilde{\Delta}_k - \alpha_{k+1} K \tilde{\Delta}_k + \alpha_{k+1} C_k w_{k+1}, \quad k \geq 1,$$

and prove that the difference $(\theta_k - \theta^*) - \tilde{\Delta}_k$ is negligible. The analysis of $\tilde{\Delta}_k$ is carried out with martingale tools.

Comparison with previous works. Theorem 4 stated above is comparable to Theorem 1 given in Pelletier (1998b). The framework presented in Pelletier (1998b) is more general as it focuses on general stochastic algorithms but their results are conditioned by the event $\{\theta_k \rightarrow \theta^*\}$. Conversely, our paper guarantees the almost sure convergence in the context of stochastic gradient descent. Moreover, our result on the weak convergence can not be recovered from the one of Pelletier (1998b) due to their Assumption A1.2. Indeed, this assumption would require that the sequence $(C_k)_{k \geq 0}$ converges towards C faster than $\sqrt{\alpha_k}$ which is either hardly met in practice or difficult to check. In a more restrictive setting of convex objective, another way to obtain the weak convergence of $(\theta_k)_{k \geq 0}$ is to rely on the results of Boyer and Godichon-Baggioni (2020). However, once again, a convergence rate on the matrix sequence $(C_k)_{k \geq 0}$ is needed in their work. This implies the derivation of an additional result on the almost sure convergence rate of the iterates. In the subsequent Section, we show that our conditions on the matrices C_k are easily satisfied in common situations.

4.3 Asymptotic optimality for conditioned SGD.

As emphasized by Assumption 10, the conditioning matrices C_k are not allowed to have too large eigenvalues. This could result in a certain instability in practice caused by large jumps towards wrong directions. In addition, as indicated by Assumption 11, the sequence $(C_k)_{k \geq 0}$ needs to converge to some matrix C . Following the result presented in Proposition 2, this matrix C should be close to the inverse of the Hessian $H = \nabla^2 F(\theta^*)$. We present a way to satisfy those assumptions.

Mixture regularization. To ensure a control on the largest eigenvalue of the conditioning matrix C_k , we rely on a weighted mixture involving Φ_k , an estimate that will be specified, and the identity matrix I_d . More precisely, let $(\gamma_k)_{k \geq 1}$ be a positive sequence and define

$$\forall k \in \mathbb{N}, \quad C_k = (\Phi_k + \gamma_{k+1}^{-1} I_d)^{-1}. \quad (5)$$

Averaging past estimates. It remains to define the matrix Φ_k . In practice, similarly to exact gradients, one may not have access to values of the Hessian matrix but only stochastic versions of it (see Section 5 for more details). As a consequence, we consider the following framework which involves random Hessian matrices. As for gradients, a policy $(P'_k)_{k \geq 0}$ is used at each iteration to produce random Hessians through a generator function H which satisfies the following property.

Assumption 12 (Unbiased and bounded Hessians). *The Hessian generator $H : \mathbb{R}^d \times S \rightarrow \mathbb{R}^{d \times d}$ is uniformly bounded around the minimizer and is such that for all $\theta \in \mathbb{R}^d$, $H(\theta, \cdot)$ is measurable with*

$$\forall k \geq 0, \quad \mathbb{E} [H(\theta_k, \xi'_{k+1}) | \mathcal{F}_k] = \nabla^2 F(\theta_k).$$

One simple and reasonable approach is to take Φ_k equal to $H(\theta_k, \xi'_{k+1})$, where only the latest point is used. However, in view of the almost sure convergence $\theta_k \rightarrow \theta^*$, it is also possible to average all past Hessian estimates to get a more robust global estimate. A weighted sum of past Hessian estimates may be used as a compromise between the two approaches and is defined as

$$\Phi_k = \sum_{j=0}^k \nu_{j,k} H(\theta_j, \xi'_{j+1}) \quad \text{with} \quad \sum_{j=0}^k \nu_{j,k} = 1. \quad (6)$$

Proposition 3. *Suppose that Assumptions 1, 3, 4, 5 and 12 are fulfilled. Let $(\theta_k)_{k \geq 0}$ be the sequence of iterates obtained by conditioned SGD (4) and consider Φ_k defined in Equation (6). If $\sup_{0 \leq j \leq k} \nu_{j,k} = O(1/k)$, then we have $\Phi_k \rightarrow H = \nabla^2 F(\theta^*)$ almost surely.*

A natural choice is to take equal weights $\nu_{j,k} = (k+1)^{-1}$. However, since the past iterates that are close to the current point are more likely to bring more relevant information through their Hessian estimates, we may use adaptive weights of the form $\nu_{j,k} \propto \exp(-\eta \|\theta_j - \theta_k\|_1)$ with a parameter $\eta \geq 0$ that recovers equal weights with $\eta = 0$. These two weights sequences satisfy the assumption of Proposition 3. They will be considered in the numerical illustration of the next section. Based on Theorem 4 and Proposition 3, we have the following result.

Theorem 5 (Asymptotic optimality of the iterates). *Suppose that Assumptions 1, 3, 4, 5, 7, 8, 9 are fulfilled. Let $(\theta_k)_{k \geq 0}$ be the sequence of iterates obtained by conditioned SGD (4) with $\alpha_k = 1/k$, C_k given by (5) and Φ_k defined by (6). If $\sup_{0 \leq j \leq k} \nu_{j,k} = O(1/k)$ and $\sum_{k \geq 1} (\gamma_k/k)^2 < \infty$, we have*

$$\sqrt{k}(\theta_k - \theta^*) \rightsquigarrow \mathcal{N}(0, H^{-1} \Gamma H^{-1}).$$

The previous Theorem shows the success of the proposed approach as the asymptotic variance obtained is the optimal one. It provides the user a practical choice for the sequence of rate, $\alpha_k = 1/k$ and also removes the assumption that $2\alpha H \succ I_d$ which is usually needed in SGD (see Proposition 2). Another remarkable result, which directly follows from the previous theorem, is now stated as a corollary.

Corollary 1 (Asymptotic optimality of the excess risk). *Let $(Z_1, \dots, Z_d) \sim \mathcal{N}(0, I_d)$ and $(\lambda_k)_{k=1, \dots, d}$ be the eigenvalues of the matrix $H^{-1/2} \Gamma H^{-1/2}$. Under the assumptions of Theorem 5, we have the convergence in distribution*

$$k(F(\theta_k) - F(\theta^*)) \rightsquigarrow \sum_{k=1}^d \lambda_k Z_k^2.$$

The above statement provides insights about the convergence speed. It claims that the convergence rate of $F(\theta_k)$ towards the optimum $F(\theta^*)$, in $1/k$, is faster than the convergence rate of the iterates, in $1/\sqrt{k}$. Another important feature, which is a consequence of Proposition 2, is that the eigenvalues $(\lambda_k)_{k=1, \dots, d}$ that appear in the limiting distribution are the smallest ones among all the other possible version of conditioned SGD (defined by the conditioning matrix C).

5 Numerical illustration

This Section is dedicated to numerical experiments on both simulated and real data for illustrative purposes of *conditioned* stochastic gradient descent methods. Consider the empirical risk minimization framework applied to Generalized Linear Models. Given a data matrix $X = (x_{i,j}) \in \mathbb{R}^{n \times d}$ with labels $y \in \mathbb{R}^n$ and a regularization parameter $\lambda > 0$, we are interested in solving the following optimization problem

$$\min_{\theta \in \mathbb{R}^d} \left\{ F(\theta) = \frac{1}{n} \sum_{i=1}^n f_i(\theta) \right\}, \quad f_i(\theta) = \mathcal{L}(x_i^T \theta, y_i) + \lambda \Omega(\theta),$$

where $\mathcal{L} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is smooth loss function and $\Omega : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is a smooth convex regularizer chosen as Tikhonov regularization $\Omega(\theta) = \frac{1}{2} \|\theta\|_2^2$. The gradient and Hessian of each component f_i are

given for all $i = 1, \dots, n$ by

$$\nabla f_i(\theta) = \mathcal{L}'(x_i^T \theta, y_i) x_i + \lambda \theta, \quad \nabla^2 f_i(\theta) = \mathcal{L}''(x_i^T \theta, y_i) x_i x_i^T + \lambda I_d,$$

where $\mathcal{L}'(\cdot, \cdot)$ and $\mathcal{L}''(\cdot, \cdot)$ are the first and second derivative of $\mathcal{L}(\cdot, \cdot)$ with respect to the first argument. As stated in Example 1 of Section 2, stochastic versions of both the gradient and the Hessian of the objective function F are computed using only a small batch $B \subset \{1, \dots, n\}$ of data ($|B| \ll n$):

$$\nabla_B F(\theta) = \frac{1}{|B|} \sum_{i \in B} \nabla f_i(\theta), \quad \nabla_B^2 F(\theta) = \frac{1}{|B|} \sum_{i \in B} \nabla^2 f_i(\theta).$$

Note that these random generators meet Assumptions 1 and 12 as they produce unbiased estimates of the gradient and the Hessian matrix respectively. We consider two different losses, namely least-squares and logistic. These losses are respectively associated to the Ridge regression problem with $y \in \mathbb{R}^n$ and the binary classification task with $y \in \{-1, +1\}^n$. The regularization parameter is set to the classical value $\lambda = 1/n$. Denote by $\sigma(z) = 1/(1 + \exp(-z))$ the sigmoid function, we have the following closed-form equations

(Ridge regression)	(Logistic regression)
$\mathcal{L}(x_i^T \theta, y_i) = (y_i - x_i^T \theta)^2 / 2$	$\mathcal{L}(x_i^T \theta, y_i) = \log(1 + \exp(-y_i x_i^T \theta))$
$\mathcal{L}'(x_i^T \theta, y_i) = x_i^T \theta - y_i$	$\mathcal{L}'(x_i^T \theta, y_i) = \sigma(x_i^T \theta) - y_i$
$\mathcal{L}''(x_i^T \theta, y_i) = 1$	$\mathcal{L}''(x_i^T \theta, y_i) = \sigma(x_i^T \theta)(1 - \sigma(x_i^T \theta))$

Starting from the null vector $\theta_0 = (0, \dots, 0) \in \mathbb{R}^d$, we use the update rule of Algorithm 1 with the optimal learning rate of the form $\alpha_k = \alpha/(k + k_0)$ (Bottou et al., 2018) and set $\gamma_k = \gamma\sqrt{k}$ in the experiments where α, k_0 and γ are tuned using a grid search. The means and standard deviations of the optimality ratio $k \mapsto [F(\theta_k) - F(\theta^*)]/[F(\theta_0) - F(\theta^*)]$, obtained over 30 independent runs, are presented in Figures below. For ease of reproducibility, the code is available online¹.

Algorithm 1 *Conditioned*-SGD with Hessian averaging

Require: $\theta_0 \in \mathbb{R}^d$, N (iterations), $(\alpha_k)_{k \geq 0}$, $(\gamma_k)_{k \geq 0}$

1. **for** $k = 0, \dots, N - 1$ **do**
 2. Compute gradient and hessian estimates $g_k = g(\theta_k, \xi_{k+1})$, $H_k = H(\theta_k, \xi'_{k+1})$
 3. Compute average matrix Φ_k as given by (6)
 4. Compute conditioning matrix C_k as given by (5)
 5. Update $\theta_{k+1} = \theta_k - \alpha_{k+1} C_k g_k$
 6. **end for**
 7. Return final point θ_N
-

We compare the performance of classical stochastic gradient descent (sgd) and its conditioned variant (csgd) where the matrix Φ_k is an averaging of past Hessian estimates as given in Equation (6). We shall compare equal weights $\nu_{j,k} = (k+1)^{-1}$ and adaptive weights $\nu_{j,k} \propto \exp(-\eta \|\theta_j - \theta_k\|_1)$ with $\eta > 0$ to give more importance to Hessian estimates associated to iterates which are closed to the current point. The methods in competition are: standard stochastic gradient descent (*sgd*), its Polyak-averaging variant (*sgd_avg*) and the conditioned stochastic gradient descent methods with equal and adaptive weights, respectively the curves associated to *csgd*($\eta = 0$) and *csgd*($\eta > 0$).

5.1 Simulated data

First, we focus on simulated data with $n = 5,000$ samples in dimensions $d \in \{20, 100\}$. Stochastic gradient methods are known to greatly benefit from mini-batch instead of picking a single random sample when computing the gradient estimate. We use a batch-size equal to $|B| = 4$ (*resp.* $|B| = 32$) in dimension $d = 20$ (*resp.* $d = 100$). In Figure 1, we can see that *conditioned* sgd outperforms standard sgd for both Ridge and Logistic regression.

¹<https://github.com/RemiLELUC/ConditionedSGD>




Figure 1: Optimality ratio $k \mapsto [F(\theta_k) - F(\theta^*)]/[F(\theta_0) - F(\theta^*)]$ for simulated data.

5.2 Real-world data

We now turn our attention to real-world datasets and consider again the Ridge and Logistic regression. The different datasets and their attributes are described as follows:

- *Boston Housing dataset* (Harrison Jr and Rubinfeld, 1978): This dataset contains information collected by the U.S Census Service concerning housing in the area of Boston Mass. It contains $n = 506$ samples in dimension $d = 14$.
- *Diabetes dataset* (Dua and Graff, 2017): Ten baseline variables, age, sex, body mass index, average blood pressure, and six blood serum measurements were obtained for each of $n = 442$ diabetes patients, as well as the response of interest, a quantitative measure of disease progression one year after baseline.
- *Mushroom dataset* (Chang and Lin, 2011): This data set includes descriptions of hypothetical samples corresponding to 23 species of gilled mushrooms in the Agaricus and Lepiota Family. Each species is identified as definitely edible, definitely poisonous, or of unknown edibility and not recommended. This latter class was combined with the poisonous one leading to a total of $n = 8,124$ samples in dimension $d = 112$.
- *Phishing dataset* (Chang and Lin, 2011): These data consist of a collection of legitimate, as well as phishing website instances. Each website is represented by the set of features that denote whether the website is legitimate or not. It contains $n = 11,055$ samples in dimension $d = 68$.

Figures 2(a) and 2(b) are dedicated to Ridge regression on the Boston and Diabetes datasets. In these small dimension settings, the conditioned sgd methods outperform both classical sgd and the Polyak-averaging variant. Similarly, concerning the classification task, Figures 2(c) and 2(d) reveal the good performance of conditioned sgd methods in a high-dimensional setting.




Figure 2: Optimality ratio $k \mapsto [F(\theta_k) - F(\theta^*)]/[F(\theta_0) - F(\theta^*)]$ for real data.

6 Conclusion and Discussion

We derived an asymptotic theory for *conditioned* stochastic gradient descent in a general non-convex setting. The use of appropriate *conditioning* matrices with the help of Hessian estimates has been the key to achieve asymptotic optimality in the sense of minimal variance.

While our study focuses on the weak convergence of the iterates (an appropriate tool to deal with efficiency issues because algorithms can be compared easily through their asymptotic variances), it would be interesting to complement our asymptotic results with concentration inequalities and non-asymptotic bounds. This may be done at the expense of additional assumptions, *e.g.*, strong convexity of the objective function combined with bounded gradients.

From a practical point of view, the approach proposed in Section 4.3 is not computationally optimal as it requires $O(d^3)$ operations due to matrix inversion. However, conditioned sgd methods do not actually require the full computation of a matrix inversion but rely on matrix-vector products which may be performed in $O(d^2)$ operations. More precisely, designing an efficient *conditioned* SGD algorithm involves a careful trade-off between the low-memory storage of the scaling matrix representation C_k and the quality of its approximation of the inverse Hessian $\nabla^2 F(\theta^*)^{-1}$. Two classical choices are diagonal approximations (Becker et al., 1988; Duchi et al., 2011) and low rank approximations, the most famous ones being Broyden-Fletcher-Goldfarb-Shanno (BFGS) algorithm (Broyden, 1970; Fletcher, 1970; Goldfarb, 1970; Shanno, 1970) and its limited memory variant L-BFGS (Liu and Nocedal, 1989). Those algorithms approximately invert Hessian matrices in $O(d)$ operations. More recently, this technique was extended to the online learning framework (Schraudolph et al., 2007) and a purely stochastic setting (Moritz et al., 2016). Though beyond the scope of the present paper, the weak convergence of these methods might be investigated with the help of Theorem 4.

Appendix

In Appendix [A](#) we present results on the expected smoothness condition and its links with our growth condition. Appendix [B](#) deals with the almost sure convergence (Theorem [3](#)) and Appendix [C](#) is the proof of the central limit theorem (Theorem [4](#)). Appendix [D](#) presents additional propositions, namely the optimality of the inverse of the Hessian matrix and the almost sure convergence of the matrix Φ_k built in Equation [\(6\)](#). Appendix [E](#) gathers Robbins-Siegmund theorem and technical Lemmas that are useful for the analysis.

A Auxiliary results on expected smoothness

The following Lemma gives sufficient conditions to meet the weak growth condition on the stochastic noise as stated in Assumption [5](#).

Lemma 2. *Suppose that for all $k \geq 1, \theta \in \mathbb{R}^d, F(\theta) = \mathbb{E}[f(\theta, \xi_k) | \mathcal{F}_{k-1}]$ with $\xi_k \sim P_{k-1}$. Assume that for all $\xi_k \sim P_{k-1}$, the function $\theta \mapsto f(\theta, \xi_k)$ is L -smooth almost surely and there exists $m \in \mathbb{R}$ such that for all $x \in \mathbb{R}^d, f(x, \xi_k) \geq m$. Then a gradient estimate is given by $g(\theta, \xi) = \nabla f(\theta, \xi)$ and the growth condition of Assumption [5](#) is satisfied with $\sigma^2 = 2L(F(\theta^*) - m)$ and*

$$\forall \theta \in \mathbb{R}^d, \forall k \in \mathbb{N}, \quad \mathbb{E}[\|g(\theta, \xi_k)\|_2^2 | \mathcal{F}_{k-1}] \leq 2L(F(\theta) - F(\theta^*)) + \sigma^2.$$

Proof. For all $\xi_k \sim P_{k-1}$, Lipschitz continuity of the gradient $\theta \mapsto \nabla f(\theta, \xi_k)$ implies (see [Nesterov \(2013\)](#))

$$f(y, \xi_k) \leq f(\theta, \xi_k) + \langle \nabla f(\theta, \xi_k), y - \theta \rangle + (L/2)\|y - \theta\|_2^2.$$

Plug $y = \theta - (1/L)\nabla f(\theta, \xi_k)$ and use the lower bound $f(y, \xi_k) \geq m$ to obtain

$$\frac{1}{2L}\|\nabla f(\theta, \xi_k)\|_2^2 \leq f(\theta, \xi_k) - f(y, \xi_k) \leq f(\theta, \xi_k) - m,$$

which gives,

$$\|g(\theta, \xi_k)\|_2^2 \leq 2L(f(\theta, \xi_k) - f(\theta^*, \xi_k)) + 2L(f(\theta^*, \xi_k) - m)$$

and conclude by taking the conditional expectation with respect to \mathcal{F}_{k-1} . □

The next Lemma links our weak growth condition with the notion of expected smoothness as introduced in [Gower et al. \(2019\)](#). In particular, this notion can be extended to our general context where the sampling distribution can evolve through the stochastic algorithm.

Lemma 3. (*Expected smoothness*) *Assume that with probability one,*

$$\sup_{k \geq 1} \sup_{x \neq x^*} \frac{\mathbb{E}[\|g(\theta, \xi_k) - g(\theta^*, \xi_k)\|_2^2 | \mathcal{F}_{k-1}]}{F(\theta) - F(\theta^*)} < \infty \quad \text{and} \quad \sup_{k \geq 1} \mathbb{E}[\|g(\theta^*, \xi_k)\|_2^2 | \mathcal{F}_{k-1}] < \infty.$$

Then there exist $0 \leq \mathcal{L}, \sigma^2 < \infty$ such that

$$\forall \theta \in \mathbb{R}^d, \forall k \in \mathbb{N}, \quad \mathbb{E}[\|g(\theta, \xi_k)\|_2^2 | \mathcal{F}_{k-1}] \leq 2\mathcal{L}(F(\theta) - F(\theta^*)) + 2\sigma^2.$$

Proof. For all $\theta \in \mathbb{R}^d$ and all $k \in \mathbb{N}$, we have

$$\begin{aligned} \|g(\theta, \xi_k)\|_2^2 &= \|g(\theta, \xi_k) - g(\theta^*, \xi_k) + g(\theta^*, \xi_k)\|_2^2 \\ &\leq 2\|g(\theta, \xi_k) - g(\theta^*, \xi_k)\|_2^2 + 2\|g(\theta^*, \xi_k)\|_2^2. \end{aligned}$$

Using the expected smoothness, with probability one, there exists $0 \leq \mathcal{L} < \infty$ such that

$$\mathbb{E}[\|g(\theta, \xi_k) - g(\theta^*, \xi_k)\|_2^2 | \mathcal{F}_{k-1}] \leq \mathcal{L}(F(\theta) - F(\theta^*)).$$

Since the noise at optimal point is almost surely finite there exists $0 \leq \sigma^2 < \infty$ such that

$$\mathbb{E}[\|g(\theta^*, \xi_k)\|_2^2 | \mathcal{F}_{k-1}] \leq \sigma^2,$$

which allows to conclude by taking the conditional expectation. □

B Proof of the almost sure convergence (Theorem 3)

The idea behind the proof of the almost sure convergence is to apply the Robbins-Siegmund Theorem (Theorem 7) which can be found in Section E.

For any matrix $A \in \mathbb{R}^{d \times d}$, we denote by $\|A\| = \max_{\|u\|_2=1} \|Au\|_2$ the operator norm associated to the Euclidian norm and by $\rho(A)$ the spectral radius of A , *i.e.*, $\rho(A) = \max\{|\lambda_1|, \dots, |\lambda_n|\}$ where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of A . We also introduce $\lambda_{\min}(A) = \min\{|\lambda_1|, \dots, |\lambda_n|\}$. Note that when A is symmetric $\|A\| = \rho(A)$ and recall that the spectral radius is a (submultiplicative) norm on the real linear space of symmetric matrices.

Since $\theta \mapsto F(\theta)$ is L -smooth, we have the quadratic bound (see Nesterov (2013))

$$\forall \theta, \eta \in \mathbb{R}^d \quad F(\eta) \leq F(\theta) + \langle \nabla F(\theta), \eta - \theta \rangle + \frac{L}{2} \|\eta - \theta\|_2^2.$$

Using the update rule $\theta_{k+1} = \theta_k - \alpha_{k+1} C_k g(\theta_k, \xi_{k+1})$, we get

$$\begin{aligned} F(\theta_{k+1}) &\leq F(\theta_k) + \langle \nabla F(\theta_k), \theta_{k+1} - \theta_k \rangle + \frac{L}{2} \|\theta_{k+1} - \theta_k\|_2^2 \\ &= F(\theta_k) - \alpha_{k+1} \langle \nabla F(\theta_k), C_k g(\theta_k, \xi_{k+1}) \rangle + \frac{L}{2} \alpha_{k+1}^2 \|C_k g(\theta_k, \xi_{k+1})\|_2^2. \end{aligned}$$

The last term can be upper bounded using the matrix norm and Assumption 10 as

$$\|C_k g(\theta_k, \xi_{k+1})\|_2^2 \leq \|C_k\|^2 \|g(\theta_k, \xi_{k+1})\|_2^2 \leq \gamma_{k+1}^2 \|g(\theta_k, \xi_{k+1})\|_2^2,$$

and we have the inequality

$$F(\theta_{k+1}) \leq F(\theta_k) - \alpha_{k+1} \langle \nabla F(\theta_k), C_k g(\theta_k, \xi_{k+1}) \rangle + \frac{L}{2} (\alpha_{k+1} \gamma_{k+1})^2 \|g(\theta_k, \xi_{k+1})\|_2^2.$$

Introduce $\mu_k = \alpha_k \gamma_k$ and $\eta_k = \alpha_k \beta_k$, we have $\sum_{k \geq 1} \eta_k = +\infty$ and $\sum_{k \geq 1} \mu_k^2 < +\infty$ a.s. in virtue of Assumption 10. The random variables $F(\theta_k), C_k$ are \mathcal{F}_k -measurable and the gradient estimate is unbiased with respect to \mathcal{F}_k . Taking the conditional expectation denoted by \mathbb{E}_k leads to

$$\begin{aligned} \mathbb{E}_k [F(\theta_{k+1})] - F(\theta_k) &\leq -\alpha_{k+1} \langle \nabla F(\theta_k), \mathbb{E}_k [C_k g(\theta_k, \xi_{k+1})] \rangle + \frac{L}{2} \mu_{k+1}^2 \mathbb{E}_k [\|g(\theta_k, \xi_{k+1})\|_2^2] \\ &= -\alpha_{k+1} \nabla F(\theta_k)^T C_k \nabla F(\theta_k) + \frac{L}{2} \mu_{k+1}^2 \mathbb{E}_k [\|g(\theta_k, \xi_{k+1})\|_2^2]. \end{aligned}$$

On the one hand for the first term, using Assumption 10 ,

$$\nabla F(\theta_k)^T C_k \nabla F(\theta_k) \geq \lambda_{\min}(C_k) \|\nabla F(\theta_k)\|_2^2 \geq \beta_{k+1} \|\nabla F(\theta_k)\|_2^2.$$

On the other hand, using Assumption 5, there exist $0 \leq \mathcal{L}, \sigma^2 < \infty$ such that almost surely

$$\forall k \in \mathbb{N}, \quad \mathbb{E}_k [\|g(\theta_k, \xi_{k+1})\|_2^2] \leq 2\mathcal{L} (F(\theta_k) - F(\theta^*)) + \sigma^2.$$

Inject these bounds in the previous inequality and subtract $F(\theta^*)$ on both sides to have

$$\mathbb{E}_k [F(\theta_{k+1}) - F(\theta^*)] \leq (1 + L\mathcal{L}\mu_{k+1}^2) [F(\theta_k) - F(\theta^*)] - \eta_{k+1} \|\nabla F(\theta_k)\|_2^2 + (L/2)\mu_{k+1}^2 \sigma^2.$$

Introduce $V_k = F(\theta_k) - F(\theta^*)$, $W_k = \eta_{k+1} \|\nabla F(\theta_k)\|_2^2$, $a_k = L\mathcal{L}\mu_{k+1}^2$ and $b_k = (L/2)\mu_{k+1}^2 \sigma^2$. These four random sequences are non-negative \mathcal{F}_k -measurable sequences with $\sum_k a_k < \infty$ and $\sum_k b_k < \infty$ almost surely. Moreover we have

$$\forall k \in \mathbb{N}, \quad \mathbb{E} [V_{k+1} | \mathcal{F}_k] \leq (1 + a_k) V_k - W_k + b_k.$$

We can apply Robbins-Siegmund Theorem 7 to have

$$(a) \sum_{k \geq 0} W_k < \infty \text{ a.s.} \quad (b) V_k \xrightarrow{\text{a.s.}} V_\infty, \mathbb{E} [V_\infty] < \infty. \quad (c) \sup_{k \geq 0} \mathbb{E} [V_k] < \infty.$$

Therefore we have almost surely that $(F(\theta_k))$ converges to a finite value $F_\infty \in L^1$ and $\sum_{k \geq 0} \eta_{k+1} \|\nabla F(\theta_k)\|_2^2 < +\infty$. There exists an event $\Omega_0 \subset \Omega$ such that, $\mathbb{P}(\Omega_0) = 1$ and for every $\omega \in \Omega_0$, $\limsup_k F(\theta_k(\omega)) < \infty$ and the series $\sum_k \eta_{k+1} \|\nabla F(\theta_k(\omega))\|_2^2$ converges. Since $\lim_{\|\theta\| \rightarrow \infty} F(\theta) = \infty$, we deduce that for every $\omega \in \Omega_0$, the sequence $(\theta_k(\omega))_{k \geq 0}$ is bounded in a finite dimensional euclidian space. Therefore the limit set $\chi_\infty(\omega)$ (set of accumulation points) of the sequence $(\theta_k(\omega))$ is non-empty. The convergence of the series $\sum_k \eta_{k+1} \|\nabla F(\theta_k(\omega))\|_2^2 < \infty$ along with the condition $\sum_k \eta_{k+1} = +\infty$ only imply that

$$\liminf_{k \rightarrow \infty} \|\nabla F(\theta_k(\omega))\|_2^2 = 0, \quad \mathbb{P} - a.s.$$

Hence, since $\theta \mapsto \nabla F(\theta)$ is continuous, there exists a limit point $\theta_\infty(\omega) \in \chi_\infty(\omega)$ such that $\|\nabla F(\theta_\infty(\omega))\|_2^2 = 0$, i.e., $\nabla F(\theta_\infty(\omega)) = 0$. Because the set of solutions $\{x \in \mathbb{R}^d, \nabla F(x) = 0\}$ is reduced to the singleton $\{x^*\}$, we have $\theta_\infty(\omega) = \theta^*$. Since $(F(\theta_k(\omega)))$ converges, it implies that $\lim_k F(\theta_k(\omega)) = F(\theta^*)$ and for every limit point $\theta \in \chi_\infty(\omega)$, we have $F(\theta) = F(\theta^*)$. Since the set $\{\theta \in \mathbb{R}^d, F(\theta) = F(\theta^*)\}$ is equal to $\{\theta^*\}$, the limit set $\chi_\infty(\omega)$ is also reduced to $\{\theta^*\}$.

C Proof of the weak convergence (Theorem 4)

Structure of the proof.

In virtue of Assumption 9, there exist $\delta, \varepsilon > 0$ such that almost surely

$$\sup_{k \geq 0} \mathbb{E}[\|w_{k+1}\|_2^{2+\delta} | \mathcal{F}_k] \mathbf{1}_{\{\|\theta_k - \theta^*\|_2 \leq \varepsilon\}} < \infty. \quad (7)$$

An important event in the following is

$$\mathcal{A}_k = \{\|\theta_k - \theta^*\|_2 \leq \varepsilon, \|C_k\| < 2\|C\|, \|\Gamma_k\| \leq 2\|\Gamma\|\}.$$

By assumption, this event has probability going to 1.

Introduce the difference

$$\Delta_k = \theta_k - \theta^*,$$

and remark that Δ_k is subjected to the iteration:

$$\begin{aligned} \Delta_0 &= \theta_0 - \theta^*, \\ \Delta_{k+1} &= \Delta_k - \alpha_{k+1} C_k \nabla F(\theta_k) + \alpha_{k+1} C_k w_{k+1}, \quad k \geq 0, \end{aligned}$$

with $w_{k+1} = \nabla F(\theta_k) - g(\theta_k, \xi_{k+1})$. We have by Assumption 11 that $C_k \rightarrow C$ almost surely and we can define $K = \lim_{k \rightarrow \infty} C_k H = CH$. The proof relies on the introduction of an auxiliary stochastic algorithm which follows the iteration:

$$\begin{aligned} \tilde{\Delta}_0 &= x_0 - x^* \\ \tilde{\Delta}_{k+1} &= \tilde{\Delta}_k - \alpha_{k+1} K \tilde{\Delta}_k + \alpha_{k+1} C_k w_{k+1} \mathbf{1}_{\mathcal{A}_k}, \quad k \geq 0 \end{aligned}$$

The previous algorithm is a linear approximation of the algorithm that defines Δ_k in the sense that $\nabla F(\theta_k) = \nabla F(\theta^*) + H(\theta_k - \theta^*) + o(\theta_k - \theta^*) \simeq H(\theta_k - \theta^*)$ has been linearly expanded around θ^* . Writing

$$\Delta_k = \tilde{\Delta}_k + (\Delta_k - \tilde{\Delta}_k),$$

and invoking the Slutsky lemma, the proof will be complete as soon as we obtain that

$$\alpha_k^{-1/2} \tilde{\Delta}_k \rightsquigarrow \mathcal{N}(0, \Sigma), \quad (8)$$

$$(\Delta_k - \tilde{\Delta}_k) = o_{\mathbb{P}}(\alpha_k^{1/2}). \quad (9)$$

Denote by \sqrt{H} the positive square root of the real symmetric positive definite matrix H and consider the transformation $\Theta_k = \sqrt{H}\tilde{\Delta}_k$ which satisfies

$$\begin{aligned}\Theta_0 &= \sqrt{H}\tilde{\Delta}_0 \\ \Theta_{k+1} &= \Theta_k - \alpha_{k+1}\tilde{K}\Theta_k + \alpha_{k+1}\sqrt{H}C_k w_{k+1}\mathbb{1}_{\mathcal{A}_k}, \quad k \geq 1,\end{aligned}$$

where $\tilde{K} = \sqrt{H}C\sqrt{H} \in \mathcal{S}_d^{++}(\mathbb{R})$ is a real symmetric positive definite matrix. The sequence $(\Theta_k)_{k \geq 0}$ is easier to study than $\tilde{\Delta}_k$ because contrary to \tilde{K} , the matrix $K = CH$ is not symmetric in general unless C and H commute. In view of Assumption 7 and 11, the eigenvalues of \tilde{K} are real and positive. Denote by λ_m (*resp.* λ_M) the smallest (*resp.* the largest) eigenvalue of \tilde{K} , *i.e.*,

$$\lambda_m = \lambda_{\min}(\tilde{K}), \quad \lambda_M = \lambda_{\max}(\tilde{K}).$$

Because CH is similar to \tilde{K} , they share the same eigenvalues. Since by assumption, the eigenvalues of $(CH - \zeta I_d)$ are positive, we have $2\alpha\lambda_m > \mathbb{1}_{\{\beta=1\}}$. For all $k \geq 1$, introduce the real symmetric matrix $A_k = I - \alpha_k\tilde{K}$. Observe that all these matrices commute, *i.e.*, for any $i, j \geq 0$, we have $A_i A_j = A_j A_i$. For any $k, n \geq 0$, denote the matrices product

$$\begin{cases} \Pi_{n,k} &= A_n \dots A_{k+1} \text{ if } k < n \\ \Pi_{n,k} &= I_d \text{ if } k \geq n, \Pi_n = \Pi_{n,0} \end{cases}$$

Since the matrices A_k commute, we have $\Pi_{n,k}^T = \Pi_{n,k}$ is also real symmetric.

Step 1. Proof of Equation (8).

The random process $(\Theta_k)_{k \geq 0}$ follows the recursion equation

$$\Theta_k = A_k \Theta_{k-1} + \alpha_k \sqrt{H} C_{k-1} w_k \mathbb{1}_{\mathcal{A}_{k-1}}.$$

We have by induction

$$\Theta_n = \Pi_n \Theta_0 + \sum_{k=1}^n \alpha_k \Pi_{n,k} \sqrt{H} C_{k-1} w_k \mathbb{1}_{\mathcal{A}_{k-1}},$$

and the rescaled process is equal to

$$\frac{\Theta_n}{\sqrt{\alpha_n}} = \underbrace{\frac{\Pi_n}{\sqrt{\alpha_n}} \Theta_0}_{\text{initial error } Y_n} + \underbrace{\sum_{k=1}^n \frac{\alpha_k}{\sqrt{\alpha_n}} \Pi_{n,k} \sqrt{H} C_{k-1} w_k \mathbb{1}_{\mathcal{A}_{k-1}}}_{\text{sampling error } M_n}.$$

Bound on the initial error.

Define $\tau_n = \sum_{k=1}^n \alpha_k$ the partial sum of the learning rates. Since Π_n is symmetric, we have $\rho(\Pi_n \Theta_0) \leq \rho(\Pi_n) \|\Theta_0\|_2$. In view of Lemma 7, since $\alpha_k \rightarrow 0$, there exists $j \geq 1$ such that

$$\rho(\Pi_n) \leq \rho(\Pi_j) \exp(-\lambda_m(\tau_n - \tau_j)).$$

Therefore, the initial error is bounded by

$$\rho(Y_n) \leq \rho(\Pi_j) \exp(\lambda_m \tau_j) \|\Theta_0\|_2 \exp(d_n) \quad \text{with} \quad d_n = -\lambda_m \tau_n - \log(\sqrt{\alpha_n}).$$

Using Lemma 8, we can treat the two cases $\beta < 1$ and $\beta = 1$. On the one hand, if $\beta < 1$ then we always have $d_n \rightarrow -\infty$. On the other hand, if $\beta = 1$, we have $d_n \sim (\frac{1}{2} - \alpha\lambda_m) \log(n)$ and the condition $2\alpha\lambda_m - 1 > 0$ ensures $d_n \rightarrow -\infty$. In both cases we get $\exp(d_n) \rightarrow 0$ and the initial error vanishes to 0.

Weak convergence of the sampling error.

Consider the random process

$$M_n = \alpha_n^{-1/2} \sum_{k=1}^n \alpha_k \Pi_{n,k} \sqrt{H} C_{k-1} w_k \mathbb{1}_{\mathcal{A}_{k-1}}.$$

Note that θ_k , \mathcal{A}_k and C_k are \mathcal{F}_k -measurable. As a consequence, M_n is a sum of martingale increments and we may rely on the following central limit theorem for martingale arrays.

Theorem 6. (*Hall and Heyde, 1980, Corollary 3.1*) Let $(W_{n,i})_{1 \leq i \leq n, n \geq 1}$ be a triangular array of random vectors such that

$$\mathbb{E}[W_{n,i} \mid \mathcal{F}_{i-1}] = 0, \quad \text{for all } 1 \leq i \leq n, \quad (10)$$

$$\sum_{i=1}^n \mathbb{E}[W_{n,i} W_{n,i}^T \mid \mathcal{F}_{i-1}] \rightarrow V^* \geq 0, \quad \text{in probability,} \quad (11)$$

$$\sum_{i=1}^n \mathbb{E}[\|W_{n,i}\|^2 \mathbf{1}_{\{\|W_{n,i}\| > \varepsilon\}} \mid \mathcal{F}_{i-1}] \rightarrow 0, \quad \text{in probability,} \quad (12)$$

then, $\sum_{i=1}^n W_{n,i} \rightsquigarrow \mathcal{N}(0, V^*)$, as $n \rightarrow \infty$.

We start by verifying (11). Let $D_k = \sqrt{H} C_{k-1} \Gamma_{k-1} C_{k-1}^T \sqrt{H} \mathbf{1}_{\mathcal{A}_{k-1}} \in \mathcal{S}_d(\mathbb{R})$. The quadratic variation of M_n is given by

$$\Sigma_n = \alpha_n^{-1} \sum_{k=1}^n \alpha_k^2 \Pi_{n,k} D_k \Pi_{n,k}^T.$$

First we can check that Σ_n is bounded. Using the triangle inequality and since the operator norm is submultiplicative, we have

$$\|\Sigma_n\| \leq \alpha_n^{-1} \sum_{k=1}^n \alpha_k^2 \|\Pi_{n,k} D_k \Pi_{n,k}^T\| \leq \alpha_n^{-1} \sum_{k=1}^n \alpha_k^2 \|D_k\| \|\Pi_{n,k}\|^2 = \alpha_n^{-1} \sum_{k=1}^n \alpha_k^2 \|D_k\| \rho(\Pi_{n,k})^2,$$

where we use in the last equality that $\Pi_{n,k}$ is real symmetric so $\|\Pi_{n,k}\| = \rho(\Pi_{n,k})$. On the event \mathcal{A}_{k-1} , the matrices C_{k-1} and Γ_{k-1} are bounded as $\|C_{k-1}\| \leq 2\|C\|$ and $\|\Gamma_{k-1}\| \leq 2\|\Gamma\|$ leading to the following bound for the matrix D_k ,

$$\begin{aligned} \|D_k\| &= \|\sqrt{H} C_{k-1} \Gamma_{k-1} C_{k-1}^T \sqrt{H} \mathbf{1}_{\mathcal{A}_{k-1}}\| \\ &\leq \|H\| \|\Gamma_{k-1}\| \|C_{k-1}\|^2 \mathbf{1}_{\mathcal{A}_{k-1}} \\ &\leq 8\|H\| \|\Gamma\| \|C\|^2 = U_D. \end{aligned}$$

It follows that

$$\|\Sigma_n\| \leq U_D \alpha_n^{-1} \sum_{k=1}^n \alpha_k^2 \rho(\Pi_{n,k})^2.$$

In view of Lemma 7, we shall split the summation from $k = 1, \dots, j$ and $k = j+1, \dots, n$ as

$$\begin{aligned} \alpha_n^{-1} \sum_{k=1}^n \alpha_k^2 \rho(\Pi_{n,k})^2 &= \alpha_n^{-1} \sum_{k=1}^j \alpha_k^2 \rho(\Pi_{n,k})^2 + \alpha_n^{-1} \sum_{k=j+1}^n \alpha_k^2 \rho(\Pi_{n,k})^2 \\ &\leq \underbrace{\alpha_n^{-1} \sum_{k=1}^j \alpha_k^2 \rho(\Pi_{n,k})^2}_{a_n} + \underbrace{\alpha_n^{-1} \sum_{k=j+1}^n \alpha_k^2 \prod_{i=k+1}^n (1 - \lambda_m \alpha_i)}_{b_n}. \end{aligned}$$

For the first term a_n , we have for all $k = 1, \dots, j$

$$\rho(\Pi_{n,k}) \leq \rho(\Pi_{n,j}) \leq \prod_{i=j+1}^n (1 - \lambda_m \alpha_i) \leq \exp(-\lambda_m (\tau_n - \tau_j)),$$

which implies since (α_k) is decreasing with $\alpha_1 = \alpha$ that

$$\sum_{k=1}^j \alpha_k^2 \rho(\Pi_{n,k})^2 \leq \alpha \tau_j \exp(-2\lambda_m (\tau_n - \tau_j)).$$

Therefore, similarly to the initial error term, we get

$$a_n \leq \alpha \tau_j \exp(2\lambda_m \tau_j) \exp(d_n) \quad \text{with} \quad d_n = -2\lambda_m \tau_n - \log(\alpha_n),$$

and the condition $2\alpha\lambda_m - 1 > 0$ ensures $d_n \rightarrow -\infty$ so that a_n goes to 0 and is almost surely bounded by U_a .

For the second term b_n , we can apply Lemma 5 and need to distinguish between the two cases:

- ($\beta = 1$) If $\alpha_n = \alpha/n$, since $2\alpha\lambda_m > 1$, we can apply Lemma 5 ($p = 1, m = 2, \lambda = \lambda_m \alpha, x_j = 0, \varepsilon_k = \alpha^2$) and obtain

$$b_n \leq \frac{\alpha^2}{2\alpha\lambda_m - 1} = U_b.$$

- ($\beta < 1$) If $\alpha_n = \alpha/n^\beta$, we deduce the same as before because $\lambda_m > 0$.

Finally in both cases, we get

$$\|\Sigma_n\| \leq U_D (U_a + U_b). \quad (13)$$

We now derive the limit of Σ_n . We shall use a recursion equation to recover a stochastic approximation scheme. Note that

$$\alpha_n \Sigma_n = \sum_{k=1}^n \alpha_k^2 \Pi_{n,k} D_k \Pi_{n,k}^T \quad (14)$$

$$= \alpha_n^2 D_n + A_n \left(\sum_{k=1}^{n-1} \alpha_k^2 \Pi_{n-1,k} D_k \Pi_{n-1,k}^T \right) A_n^T, \quad (15)$$

and recognize

$$\alpha_n \Sigma_n = \alpha_n^2 D_n + \alpha_{n-1} A_n \Sigma_{n-1} A_n^T.$$

Replacing the symmetric matrix $A_n = I - \alpha_n \tilde{K}$, we get (because Σ_n is bounded almost surely)

$$\begin{aligned} \alpha_n \Sigma_n &= \alpha_n^2 D_n + \alpha_{n-1} (I - \alpha_n \tilde{K}) \Sigma_{n-1} (I - \alpha_n \tilde{K}) \\ &= \alpha_n^2 D_n + \alpha_{n-1} \left[\Sigma_{n-1} - \alpha_n \Sigma_{n-1} \tilde{K} - \alpha_n \tilde{K} \Sigma_{n-1} + O(\alpha_n^2) \right]. \end{aligned}$$

Divide by α_n to obtain

$$\Sigma_n = \alpha_n D_n + \frac{\alpha_{n-1}}{\alpha_n} \left[\Sigma_{n-1} - \alpha_n (\tilde{K} \Sigma_{n-1} + \Sigma_{n-1} \tilde{K}) + O(\alpha_n^2) \right],$$

and we recognize a stochastic approximation scheme

$$\Sigma_n = \Sigma_{n-1} - \alpha_n \left[\tilde{K} \Sigma_{n-1} + \Sigma_{n-1} \tilde{K} - D_n \right] + \frac{\alpha_{n-1} - \alpha_n}{\alpha_n} \Sigma_{n-1} + O(\alpha_{n-1} \alpha_n + |\alpha_{n-1} - \alpha_n|)$$

Recall that when $\beta < 1$ we have

$$\frac{1}{\alpha_n} - \frac{1}{\alpha_{n-1}} \rightarrow 0, \text{ i.e., } \frac{\alpha_{n-1} - \alpha_n}{\alpha_n} = o(\alpha_n).$$

- ($\beta = 1$) If $\alpha_n = \alpha/n$ we get

$$\begin{aligned} \Sigma_n &= \Sigma_{n-1} - \frac{\alpha}{n} \left[\tilde{K} \Sigma_{n-1} + \Sigma_{n-1} \tilde{K} - \frac{1}{\alpha} \Sigma_{n-1} - D_n \right] + O(n^{-2}) \\ \Sigma_n &= \Sigma_{n-1} - \frac{\alpha}{n} \left[\left(\tilde{K} - \frac{I}{2\alpha} \right) \Sigma_{n-1} + \Sigma_{n-1} \left(\tilde{K} - \frac{I}{2\alpha} \right) - D_n \right] + O(n^{-2}). \end{aligned}$$

• ($\beta < 1$) If $\alpha_n = \alpha/n^\beta$ we get

$$\Sigma_n = \Sigma_{n-1} - \alpha_n \left[\tilde{K} \Sigma_{n-1} + \Sigma_{n-1} \tilde{K} - D_n \right] + o(\alpha_n).$$

Recall that $\zeta = \mathbb{1}_{\{\beta=1\}}/(2\alpha)$ and define $\tilde{K}_\zeta = \tilde{K} - \zeta I$, so that in both cases, the recursion equation becomes

$$\Sigma_n = \Sigma_{n-1} - \alpha_n \left[\tilde{K}_\zeta \Sigma_{n-1} + \Sigma_{n-1} \tilde{K}_\zeta^T - D_n \right] + o(\alpha_n).$$

We can vectorize this equation. The vectorization of an $m \times n$ matrix $A = (a_{i,j})$, denoted $\text{vec}(A)$, is the $mn \times 1$ column vector obtained by stacking the columns of the matrix A on top of one another:

$$\text{vec}(A) = [a_{1,1}, \dots, a_{m,1}, a_{1,2}, \dots, a_{m,2}, \dots, a_{1,n}, \dots, a_{m,n}]^T.$$

Applying this operator to our stochastic approximation scheme gives

$$\text{vec}(\Sigma_n) = \text{vec}(\Sigma_{n-1}) - \alpha_n \left[\text{vec} \left(\tilde{K}_\zeta \Sigma_{n-1} + \Sigma_{n-1} \tilde{K}_\zeta^T \right) - \text{vec}(D_n) \right] + o(\alpha_n).$$

Denote by \otimes the Kronecker product, we have the following property

$$\text{vec} \left(K_\zeta \Sigma_{n-1} + \Sigma_{n-1} K_\zeta^T \right) = (I_d \otimes K_\zeta + K_\zeta^T \otimes I_d) \text{vec}(\Sigma_{n-1}).$$

Define D as the almost sure limit of D_n , i.e.

$$D = \lim_{n \rightarrow \infty} D_n = \sqrt{H} C \Gamma C \sqrt{H}.$$

Introduce $v_n = \text{vec}(\Sigma_n)$ and $Q = \left(I_d \otimes \tilde{K}_\zeta + \tilde{K}_\zeta^T \otimes I_d \right)$. We have almost surely

$$\begin{aligned} v_n &= v_{n-1} - \alpha_n (Q v_{n-1} - \text{vec}(D)) + \alpha_n \text{vec}(D_n - D) + o(\alpha_n) \\ &= v_{n-1} - \alpha_n (Q v_{n-1} - \text{vec}(D)) + \varepsilon_n \alpha_n \end{aligned}$$

where $\varepsilon_n \rightarrow 0$ almost surely. This is a stochastic approximation scheme with the affine function $h(v) = Qv - \text{vec}(D)$ for $v \in \mathbb{R}^{d^2}$. Let v^* be the solution of $h(v) = 0$ which is well defined since $Q = \left(I_d \otimes \tilde{K}_\zeta + \tilde{K}_\zeta^T \otimes I_d \right)$ is invertible. Indeed, the eigenvalues of Q are $\mu_i + \mu_j$, $1 \leq i, j \leq d$, where the μ_i , $i = 1, \dots, d$ are the eigenvalues of \tilde{K}_ζ . Equivalently, the eigenvalues of Q are of the form $(\lambda_i - \zeta) + (\lambda_j - \zeta)$ where the λ_i , $i = 1, \dots, d$ are the eigenvalues of \tilde{K} . Because $\lambda_m > \zeta$, we have that $Q \succ 0$. As a consequence

$$\begin{aligned} (v_n - v^*) &= (v_{n-1} - v^*) - \alpha_n (h(v_{n-1}) - h(v^*)) + \varepsilon_n \alpha_n \\ &= (v_{n-1} - v^*) - \alpha_n Q (v_{n-1} - v^*) + \varepsilon_n \alpha_n \\ &= B_n (v_{n-1} - v^*) + \varepsilon_n \alpha_n, \end{aligned}$$

with $B_n = (I_{d^2} - \alpha_n Q)$. By induction, we obtain

$$(v_n - v^*) = (B_n \dots B_1) (v_0 - v^*) + \sum_{k=1}^n \alpha_k (B_n \dots B_{k+1}) \varepsilon_k,$$

Define $\lambda_Q = \lambda_{\min}(Q) > 0$ and remark that

$$\|B_n \dots B_{k+1}\| \leq \prod_{j=k+1}^n \|B_j\| = \prod_{j=k+1}^n (1 - \alpha_j \lambda_Q).$$

It follows that

$$\begin{aligned} \|v_n - v^*\|_2 &\leq \|B_n \dots B_1\| \|v_0 - v^*\|_2 + \sum_{k=1}^n \alpha_k \|B_n \dots B_{k+1}\| \|\varepsilon_k\|_2 \\ &\leq \prod_{j=1}^n (1 - \alpha_j \lambda_Q) \|v_0 - v^*\|_2 + \sum_{k=1}^n \alpha_k \prod_{j=k+1}^n (1 - \alpha_j \lambda_Q) \|\varepsilon_k\|_2 \end{aligned}$$

Applying Lemma 5 we obtain that the right-hand side term goes to 0. The left-hand side term goes to 0 under the effect of the product by definition of $(\alpha_k)_{k \geq 1}$. We therefore conclude that $v_n \rightarrow v^*$ almost surely. From easy manipulation involving $\text{vec}(\cdot)$ and \otimes , this is equivalent to $\Sigma_n \rightarrow \Sigma$, where Σ is the solution of the Lyapunov equation

$$(\tilde{K} - \zeta I)\Sigma + \Sigma(\tilde{K} - \zeta I) = D.$$

Now we turn our attention to (12). We need to show that almost surely,

$$\alpha_n^{-1} \sum_{k=1}^n \alpha_k^2 \mathbb{E}[\|\Pi_{n,k} \sqrt{H} C_{k-1} w_k\|_2^2 \mathbb{1}_{\{\alpha_k \|\Pi_{n,k} \sqrt{H} C_{k-1} w_k\|_2 > \varepsilon \alpha_n^{1/2}\}} \mid \mathcal{F}_{k-1}] \mathbb{1}_{\mathcal{A}_{k-1}} \rightarrow 0.$$

We have

$$\begin{aligned} & \mathbb{E}[\alpha_n^{-1} \alpha_k^2 \|\Pi_{n,k} \sqrt{H} C_{k-1} w_k\|_2^2 \mathbb{1}_{\{\alpha_k \|\Pi_{n,k} \sqrt{H} C_{k-1} w_k\|_2 > \varepsilon \alpha_n^{1/2}\}} \mid \mathcal{F}_{k-1}] \\ & \leq \varepsilon^{-\delta} \mathbb{E}[(\alpha_n^{-1/2} \alpha_k \|\Pi_{n,k} \sqrt{H} C_{k-1} w_k\|_2)^{2+\delta} \mid \mathcal{F}_{k-1}] \\ & \leq \varepsilon^{-\delta} (\alpha_n^{-1/2} \alpha_k \|\Pi_{n,k} \sqrt{H} C_{k-1}\|_2)^{2+\delta} \mathbb{E}[\|w_k\|_2^{2+\delta} \mid \mathcal{F}_{k-1}]. \end{aligned}$$

Let $U(\omega) = \sup_{k \geq 1} \mathbb{E}[\|w_k\|_2^{2+\delta} \mid \mathcal{F}_{k-1}] \mathbb{1}_{\mathcal{A}_{k-1}}$ which is almost surely finite by Assumption 9. We get

$$\begin{aligned} & \mathbb{E}[\alpha_n^{-1} \alpha_k^2 \|\Pi_{n,k} \sqrt{H} C_{k-1} w_k\|_2^2 \mathbb{1}_{\{\alpha_k \|\Pi_{n,k} \sqrt{H} C_{k-1} w_k\|_2 > \varepsilon \alpha_n^{1/2}\}} \mid \mathcal{F}_{k-1}] \mathbb{1}_{\mathcal{A}_{k-1}} \\ & \leq \varepsilon^{-\delta} \left(2 \|\sqrt{H}\| \|C\|\right)^{2+\delta} U(\omega) (\alpha_n^{-1/2} \alpha_k \rho(\Pi_{n,k}))^{2+\delta} \end{aligned}$$

Hence by showing that

$$\sum_{k=1}^n (\alpha_n^{-1/2} \alpha_k \rho(\Pi_{n,k}))^{2+\delta} \rightarrow 0,$$

we will obtain (12). The previous convergence can be deduced from Lemma 5 with $p = 1 + \delta/2$, $m = 2 + \delta$, $\epsilon_k = \alpha_k^{\delta/2}$, checking that $(2 + \delta)\alpha \lambda_m > 1 + \delta/2$.

Step 2. Proof of Equation (9).

A preliminary step to the derivation of Equation (9) is to obtain that $\tilde{\Delta}_k \rightarrow 0$ almost surely. For any θ and η in \mathbb{R}^d , we have

$$\|\theta\|^2 = \|\eta\|^2 + 2\eta^T(\theta - \eta) + \|\theta - \eta\|^2$$

implying that for all $k \geq 0$

$$\mathbb{E}[\|\Theta_{k+1}\|^2 \mid \mathcal{F}_k] = \|\tilde{\Theta}_k\|^2 - 2\alpha_{k+1} \Theta_k^T \tilde{K} \Theta_k + \alpha_{k+1}^2 \mathbb{E}[\|\tilde{K} \Theta_k - C_k w_{k+1} \mathbb{1}_{\mathcal{A}_k}\|^2 \mid \mathcal{F}_k].$$

Since (w_k) is a martingale increment and using matrix norms we get

$$\begin{aligned} \mathbb{E}[\|\tilde{K} \Theta_k - C_k w_{k+1} \mathbb{1}_{\mathcal{A}_k}\|^2 \mid \mathcal{F}_k] &= \mathbb{E}[\|\tilde{K} \Theta_k\|^2 \mid \mathcal{F}_k] + \mathbb{E}[\|C_k w_{k+1} \mathbb{1}_{\mathcal{A}_k}\|^2 \mid \mathcal{F}_k] \\ &\leq \lambda_M^2 \|\Theta_k\|^2 + \gamma_{k+1}^2 \mathbb{E}[\|w_{k+1} \mathbb{1}_{\mathcal{A}_k}\|^2 \mid \mathcal{F}_k]. \end{aligned}$$

Injecting this bound in the previous equality yields

$$\mathbb{E}[\|\Theta_{k+1}\|^2 \mid \mathcal{F}_k] \leq \|\Theta_k\|^2 (1 + \alpha_{k+1}^2 \lambda_M^2) - 2\alpha_{k+1} \Theta_k^T \tilde{K} \Theta_k + (\alpha_{k+1} \gamma_{k+1})^2 \mathbb{E}[\|w_{k+1}\|^2 \mid \mathcal{F}_k] \mathbb{1}_{\mathcal{A}_k}.$$

Since, using (7),

$$\sum_{k \geq 0} (\alpha_{k+1} \gamma_{k+1})^2 \mathbb{E}[\|w_{k+1}\|^2 \mid \mathcal{F}_k] \mathbb{1}_{\mathcal{A}_k} \leq \left(\sup_{k \geq 0} \mathbb{E}[\|w_{k+1}\|^2 \mid \mathcal{F}_k] \mathbb{1}_{\mathcal{A}_k} \right) \left(\sum_{k \geq 0} (\alpha_{k+1} \gamma_{k+1})^2 \right) < \infty,$$

we are in position to apply the Robbins-Siegmund Theorem 7 and we obtain the almost sure convergence of $\sum_k \alpha_{k+1} \Theta_k^T \tilde{K} \Theta_k$ and $\|\Theta_k\|_2^2 \rightarrow V_\infty$. Because \tilde{K} is positive definite, it gives that, with probability 1, $\sum_{k \geq 0} \alpha_{k+1} \|\Theta_k\|^2 < +\infty$, from which, we deduce $\liminf_k \|\Theta_k\|^2 = 0$. Therefore one can extract a subsequence Θ_k such that $\|\Theta_k\|^2 \rightarrow 0$. Using the above second condition yields $V_\infty = 0$ and we conclude that $\tilde{\Delta}_k = H^{-1/2} \Theta_k \rightarrow 0$.

Define the difference

$$E_k = \Delta_k - \tilde{\Delta}_k.$$

Since $\theta \mapsto \nabla^2 F(\theta)$ is continuous at θ^* , we can apply a coordinate-wise mean value theorem. Indeed, for any $\theta \in \mathbb{R}^d$, we have $\nabla F(\theta) = (\partial_1 F(\theta), \dots, \partial_d F(\theta))$ where for all $j = 1, \dots, d$, the partial derivatives functions $\partial_j F : \mathbb{R}^d \rightarrow \mathbb{R}$ are Lipschitz continuous. Denote by $\nabla(\partial_j F) : \mathbb{R}^d \rightarrow \mathbb{R}^d$ the gradient of the partial derivative $\partial_j F$, i.e., $\nabla(\partial_j F)(\theta) = (\partial_{1,j}^2 F(\theta), \dots, \partial_{d,j}^2 F(\theta))$. For any $\theta, \eta \in B(\theta^*, \varepsilon)$, there exists $\xi_j \in \mathbb{R}^d$ such that

$$\partial_j F(\theta) - \partial_j F(\eta) = \nabla(\partial_j F)(\xi_j)(\theta - \eta).$$

We construct a Hessian matrix by rows $H(\xi) = H(\xi_1, \dots, \xi_d)$ where the j -th row is equal to $\nabla(\partial_j F)(\xi_j) = (\partial_{1,j}^2 F(\xi_j), \dots, \partial_{d,j}^2 F(\xi_j))$

$$H(\xi) = \begin{bmatrix} \partial_{1,1}^2 F(\xi_1) & \dots & \partial_{1,d}^2 F(\xi_1) \\ \vdots & \ddots & \vdots \\ \partial_{d,1}^2 F(\xi_d) & \dots & \partial_{d,d}^2 F(\xi_d) \end{bmatrix}$$

and we can write

$$\nabla F(\theta) - \nabla F(\eta) = H(\xi)(\theta - \eta).$$

There exists $\xi_k = (\xi_k^{(1)}, \dots, \xi_k^{(d)})$ with $\xi_k^{(j)} \in [\theta^* + E_k, \theta_k]$ and $\xi'_k = (\xi_k'^{(1)}, \dots, \xi_k'^{(d)})$ with $\xi_k'^{(j)} \in [\theta^* + E_k, \theta^*]$ such that

$$\nabla F(\theta^* + E_k) - \nabla F(\theta_k) = -H(\xi_k) \tilde{\Delta}_k \quad (16)$$

$$\nabla F(\theta^* + E_k) = H(\xi'_k) E_k. \quad (17)$$

Let $\eta > 0$ such that $2\alpha\lambda_m(1 - 3\eta) > 1$. This choice will come clear at the end of the reasoning. On the one hand, we have $C_k \rightarrow C$. On the other hand, using Lemma 6, the spectrum of $C_k H$ is real and positive. Hence, we have the convergence of the eigenvalues of $C_k H$ towards the eigenvalues of $K = CH$. This follows from the definition of eigenvalues as roots of the characteristic polynomial and the fact that the roots of any polynomial $P \in \mathbb{C}[X]$ are continuous functions of the coefficients (Zedek, 1965). Consequently, there exists $n_1(\omega)$ such that for all $k \geq n_1(\omega)$,

$$(1 - \eta)\lambda_m \leq \lambda_{\min}(C_k H) \leq \lambda_{\max}(C_k H) \leq (1 + \eta)\lambda_M. \quad (18)$$

We can define $n_2(\omega)$ such that for all $k \geq n_2(\omega)$

$$\mathcal{A}_k \text{ is realized.} \quad (19)$$

Since $\|\sqrt{H^{-1}} H(\xi'_k) \sqrt{H^{-1}} - I_d\| \rightarrow 0$ as $k \rightarrow \infty$, there is $n_3(\omega)$ and $n_4(\omega)$ such that for all $k \geq n_3(\omega)$

$$\|\sqrt{H^{-1}} H(\xi'_k) \sqrt{H^{-1}} - I_d\| \leq \frac{\eta}{1 + \eta} \frac{\lambda_m}{\lambda_M}, \quad (20)$$

and for all $k \geq n_4(\omega)$,

$$\|\sqrt{H^{-1}} H(\xi'_k) \sqrt{H^{-1}}\| \leq 1. \quad (21)$$

Since $\alpha_k \rightarrow 0$, there is n_5 such that for all $k \geq n_5$

$$\alpha_{k+1} \leq \frac{2\eta\lambda_m}{(1+\eta)^2\lambda_M^2}. \quad (22)$$

To use the previous local properties, define $n_0(\omega) = n_1(\omega) \vee n_2(\omega) \vee n_3(\omega) \vee n_4(\omega) \vee n_5$ and introduce the set \mathcal{E}_j along with its complement \mathcal{E}_j^c , defined by

$$\mathcal{E}_j = \{\omega : j \geq n_0(\omega)\}.$$

Let $\delta > 0$ and take $j \geq 1$ large enough such that $\mathbb{P}(\mathcal{E}_j^c) \leq \delta$. Invoking the Markov inequality, we have for all $\gamma > 0$

$$\begin{aligned} \mathbb{P}(\alpha_k^{-1/2}\|E_k\| > \gamma) &= \mathbb{P}(\alpha_k^{-1/2}\|E_k\| > \gamma, \mathcal{E}_j) + \mathbb{P}(\alpha_k^{-1/2}\|E_k\| > \gamma, \mathcal{E}_j^c) \\ &\leq \mathbb{P}(\alpha_k^{-1/2}\|E_k\| > \gamma, \mathcal{E}_j) + \delta \\ &\leq \alpha_k^{-1/2}\gamma^{-1}\mathbb{E}[\|E_k\|\mathbb{1}_{\mathcal{E}_j}] + \delta \end{aligned}$$

Because δ is arbitrary, we only need to show that for any value of $j \geq 1$,

$$e_k := \mathbb{E}[\|E_k\|\mathbb{1}_{\mathcal{E}_j}] = o(\alpha_k^{1/2}).$$

To prove this fact, we shall recognize a stochastic algorithm for the sequence e_k .

Let $k \geq j$ and assume further that \mathcal{E}_j is realized. We have, because of (19),

$$E_{k+1} = \Delta_k - \tilde{\Delta}_k - \alpha_{k+1}C_k\nabla F(\theta_k) + \alpha_{k+1}K\tilde{\Delta}_k.$$

Introducing $\tilde{E}_k = \sqrt{H}E_k$, we find

$$\tilde{E}_{k+1} = \tilde{E}_k - \alpha_{k+1}\sqrt{H}C_k\nabla F(\theta_k) + \alpha_{k+1}\sqrt{H}K\tilde{\Delta}_k,$$

and using (16), it comes that

$$\begin{aligned} \tilde{E}_{k+1} &= \tilde{E}_k - \alpha_{k+1}\sqrt{H}C_k\nabla F(\theta^* + E_k) - \alpha_{k+1}\sqrt{H}C_kH(\xi_k)\tilde{\Delta}_k + \alpha_{k+1}\sqrt{H}K\tilde{\Delta}_k \\ &= \tilde{E}_k - \alpha_{k+1}\sqrt{H}C_k\nabla F(\theta^* + E_k) + \alpha_{k+1}\sqrt{H}(K - C_kH(\xi_k))\tilde{\Delta}_k. \end{aligned}$$

Using Minkowski inequality, we have

$$\|\tilde{E}_{k+1}\| \leq \|\tilde{E}_k - \alpha_{k+1}\sqrt{H}C_k\nabla F(\theta^* + E_k)\| + \|\alpha_{k+1}\sqrt{H}(K - C_kH(\xi_k))\tilde{\Delta}_k\|.$$

We shall now focus on the first term. Still on the set \mathcal{E}_j , we have

$$\begin{aligned} &\|\tilde{E}_k - \alpha_{k+1}\sqrt{H}C_k\nabla F(\theta^* + E_k)\|^2 \\ &= \|\tilde{E}_k\|^2 - 2\alpha_{k+1}\langle \tilde{E}_k, \sqrt{H}C_k\nabla F(\theta^* + E_k) \rangle + \alpha_{k+1}^2\|\sqrt{H}C_k\nabla F(\theta^* + E_k)\|^2 \end{aligned} \quad (23)$$

We have on the one hand using (17)

$$\begin{aligned} \langle \tilde{E}_k, \sqrt{H}C_k\nabla F(\theta^* + E_k) \rangle &= \langle \tilde{E}_k, \sqrt{H}C_kH(\xi'_k)E_k \rangle \\ &= \langle \tilde{E}_k, \sqrt{H}C_kHE_k \rangle + \langle \tilde{E}_k, \sqrt{H}C_k(H(\xi'_k) - H)E_k \rangle \end{aligned}$$

Due to (18), the first term satisfies

$$\begin{aligned} \langle \tilde{E}_k, \sqrt{H}C_kHE_k \rangle &= \langle \tilde{E}_k, \sqrt{H}C_k\sqrt{H}\tilde{E}_k \rangle \\ &\geq \lambda_{\min}(C_kH)\|\tilde{E}_k\|^2 \\ &\geq (1-\eta)\lambda_m\|\tilde{E}_k\|^2 \end{aligned}$$

The second term satisfies

$$\begin{aligned} \langle \tilde{E}_k, \sqrt{H}C_k(H(\xi'_k) - H)E_k \rangle &= \langle \tilde{E}_k, \sqrt{H}C_k\sqrt{H}(\sqrt{H^{-1}}H(\xi'_k)\sqrt{H^{-1}} - I_d)\tilde{E}_k \rangle \\ &\geq -\left| \langle \tilde{E}_k, \sqrt{H}C_k\sqrt{H}(\sqrt{H^{-1}}H(\xi'_k)\sqrt{H^{-1}} - I_d)\tilde{E}_k \rangle \right| \end{aligned}$$

Using Cauchy-Schwarz inequality, the submultiplicativity of the norm, (18) and (20), we have

$$\begin{aligned} & \left| \langle \tilde{E}_k, \sqrt{H}C_k\sqrt{H}(\sqrt{H^{-1}}H(\xi'_k)\sqrt{H^{-1}} - I_d)\tilde{E}_k \rangle \right| \\ & \leq \|\sqrt{H}C_k\sqrt{H}\| \|\sqrt{H^{-1}}H(\xi'_k)\sqrt{H^{-1}} - I_d\| \|\tilde{E}_k\|^2 \\ & \leq \eta\lambda_m \|\tilde{E}_k\|^2. \end{aligned}$$

Finally, it follows that

$$\langle \tilde{E}_k, \sqrt{H}C_k\nabla F(\theta^* + E_k) \rangle \geq (1 - 2\eta)\lambda_m \|\tilde{E}_k\|^2 \quad (24)$$

On the other hand using (17), (18) and (21),

$$\begin{aligned} \|\sqrt{H}C_k\nabla F(\theta^* + E_k)\|^2 &= \|\sqrt{H}C_kH(\xi'_k)E_k\|^2 \\ &= \|\sqrt{H}C_k\sqrt{H}(\sqrt{H^{-1}}H(\xi'_k)\sqrt{H^{-1}})\tilde{E}_k\|^2 \\ &\leq \lambda_{\max}(C_kH)^2 \|\sqrt{H^{-1}}H(\xi'_k)\sqrt{H^{-1}}\|^2 \|\tilde{E}_k\|^2 \\ &\leq (1 + \eta)^2 \lambda_M^2 \|\tilde{E}_k\|^2 \end{aligned} \quad (25)$$

Putting together (23), (24), (25) and using (22) gives that, on \mathcal{E}_j ,

$$\begin{aligned} & \|\tilde{E}_k - \alpha_{k+1}\sqrt{H}C_k\nabla F(\theta^* + E_k)\|^2 \\ & \leq \|\tilde{E}_k\|^2 (1 - 2\alpha_{k+1}(1 - 2\eta)\lambda_m + \alpha_{k+1}^2(1 + \eta)^2 \lambda_M^2) \\ & \leq \|\tilde{E}_k\|^2 (1 - 2\alpha_{k+1}(1 - 3\eta)\lambda_m). \end{aligned}$$

By the Minkowski inequality and the fact that $(1 - x)^{1/2} \leq 1 - x/2$, on \mathcal{E}_j , it holds

$$\begin{aligned} \|\tilde{E}_{k+1}\| &\leq \|\tilde{E}_k\| (1 - 2\alpha_{k+1}(1 - 3\eta)\lambda_m)^{1/2} + \alpha_{k+1} \|\sqrt{H}(K - C_kH(\xi_k))\tilde{\Delta}_k\| \\ &\leq \|\tilde{E}_k\| (1 - \alpha_{k+1}(1 - 3\eta)\lambda_m) + \alpha_{k+1} \|\sqrt{H}\| \|(K - C_kH(\xi_k))\tilde{\Delta}_k\| \end{aligned}$$

Hence, we have shown that for any $k \geq j$,

$$\|\tilde{E}_k\| \mathbf{1}_{\mathcal{E}_j} \leq \|\tilde{E}_k\| \mathbf{1}_{\mathcal{E}_j} (1 - \alpha_{k+1}(1 - 3\eta)\lambda_m) + \alpha_{k+1} \|\sqrt{H}\| \|(K - C_kH(\xi_k))\mathbf{1}_{\mathcal{E}_j}\tilde{\Delta}_k\|.$$

It follows that, for any $k \geq j$,

$$e_{k+1} \leq e_k (1 - \alpha_{k+1}(1 - 3\eta)\lambda_m) + \alpha_{k+1} \|\sqrt{H}\| \mathbb{E}[\|U_k\tilde{\Delta}_k\|],$$

with $U_k = (K - C_kH(\xi_k))\mathbf{1}_{\mathcal{E}_j}$. Because with probability 1, $\|U_k\|$ is bounded, we can apply the Lebesgue dominated convergence theorem to obtain that $\varepsilon_k = \mathbb{E}[\|U_k\|^2] \rightarrow 0$. From the Cauchy-Schwarz inequality, we get

$$\mathbb{E}[\|U_k\tilde{\Delta}_k\|] \leq \sqrt{\varepsilon_k} \sqrt{\mathbb{E}[\|\tilde{\Delta}_k\|_2^2]}.$$

On the other hand, we have already shown in (13) that $\rho(\Sigma_k) = \|\Sigma_k\| \leq U_D(U_a + U_b)$. Since $\tilde{\Delta}_k = \sqrt{H^{-1}}\Theta_k = \sqrt{H^{-1}}\sqrt{\alpha_k}(Y_k + M_k)$, we have

$$\mathbb{E}[\|\tilde{\Delta}_k\|_2^2] \leq 2(\alpha_k/\lambda_m)(\|Y_k\|_2^2 + \mathbb{E}[\|M_k\|_2^2]),$$

where the last term is the leading term and satisfies

$$\mathbb{E}[\|M_k\|_2^2] = \mathbb{E}[\text{Tr}(\Sigma_k)] \leq d\mathbb{E}[\rho(\Sigma_k)].$$

Therefore, we have

$$\mathbb{E}[\|\tilde{\Delta}_k\|_2^2] \leq \alpha_k A$$

for some $A > 0$. Consequently, for all $k \geq j$,

$$e_{k+1} \leq e_k (1 - \alpha_{k+1}(1 - 3\eta)\lambda_m) + \alpha_{k+1}^{3/2} A' \|\sqrt{H}\| \varepsilon_k^{1/2}.$$

The condition $2\alpha\lambda_m(1 - 3\eta) > 1$ ensures that we can apply Lemma 5 with $(m\lambda > p)$, $m = 1$, $p = 1/2$, $\lambda = \alpha(1 - 3\eta)\lambda_m$. we finally get

$$\limsup_k (e_k/\alpha_k^{1/2}) = 0.$$

As a consequence, $e_k = o(\sqrt{\alpha_k})$, which concludes the proof.

Since $\alpha_k^{-1/2}\sqrt{H}\tilde{\Delta}_k \rightarrow \mathcal{N}(0, \Sigma)$, we have $\alpha_k^{-1/2}\tilde{\Delta}_k \rightarrow \mathcal{N}(0, \tilde{\Sigma})$ where $\tilde{\Sigma} = \sqrt{H^{-1}}\Sigma\sqrt{H^{-1}}$. Recall that Σ satisfies the Lyapunov equation

$$(\sqrt{HC}\sqrt{H} - \zeta I_d)\Sigma + \Sigma(\sqrt{HC}\sqrt{H} - \zeta I_d) = \sqrt{H}C\Gamma C\sqrt{H}.$$

Multiplying on the left and right sides by $\sqrt{H^{-1}}$, we get

$$C\sqrt{H}\Sigma\sqrt{H^{-1}} - \zeta\sqrt{H^{-1}}\Sigma\sqrt{H^{-1}} + \sqrt{H^{-1}}\Sigma\sqrt{HC} - \zeta\sqrt{H^{-1}}\Sigma\sqrt{H^{-1}} = C\Gamma C,$$

where we recognize the following Lyapunov equation

$$(CH - \zeta I_d)\tilde{\Sigma} + \tilde{\Sigma}(CH - \zeta I_d)^T = C\Gamma C.$$

D Additional propositions

Proposition 4. *The choice $C^* = H^{-1}$ is optimal in the sense that $\Sigma_{C^*} \preceq \Sigma_C$, $\forall C \in \mathcal{C}_H$. Moreover, we have $\Sigma_{C^*} = H^{-1}\Gamma H^{-1}$.*

Proof. Define $\Delta_C = \Sigma_C - H^{-1}\Gamma H^{-1}$ and check that Δ_C satisfies

$$(CH - I_d/2)\Delta_C + \Delta_C(CH - I_d/2)^T = (C - H^{-1})\Gamma(C - H^{-1}).$$

Because Γ is symmetric positive semi-definite, we have using Lemma 10 that the term on the right side is symmetric positive semi-definite. Therefore, in view of Proposition 6, we get that Δ_C is symmetric positive semi-definite $\Delta_C \succeq 0$ which implies $\Sigma_C \succeq H^{-1}\Gamma H^{-1}$ for all $C \in \mathcal{C}_H$. The equality is reached for $C^* = H^{-1}$ with $\Delta_C = 0$, $\Sigma_{C^*} = H^{-1}\Gamma H^{-1}$. \square

Proposition 5. *Suppose that Assumptions 1, 3, 4, 5 and 12 are fulfilled. Let $(\theta_k)_{k \geq 0}$ be the sequence of the iterates obtained by conditioned SGD (4) and consider Φ_k defined in Equation (6). If $\sup_{0 \leq j \leq k} \nu_{j,k} = O(1/k)$ then we have $\Phi_k \rightarrow H = \nabla^2 F(\theta^*)$ almost surely.*

Proof. We use the decomposition

$$\Phi_k - H = \sum_{j=0}^k \nu_{j,k} (\nabla^2 F(\theta_j) - H) + \sum_{j=0}^k \nu_{j,k} (H(\theta_j, \xi'_{j+1}) - \nabla^2 F(\theta_j)).$$

The continuity of $\nabla^2 F$ at θ^* and the fact that $\theta_j \rightarrow \theta^*$ a.s. imply that $\|\nabla^2 F(\theta_j) - H\| \rightarrow 0$ a.s. Since $\sup_{0 \leq j \leq k} \nu_{j,k} = O(1/k)$, there exists $a > 0$ such that

$$\left\| \sum_{j=0}^k \nu_{j,k} (\nabla^2 F(\theta_j) - H) \right\| \leq \frac{a}{k+1} \sum_{j=0}^k \|\nabla^2 F(\theta_j) - H\|,$$

which goes to 0 in virtue of Cesaro's Lemma, therefore $\lim_{k \rightarrow \infty} \sum_{j=0}^k \nu_{j,k} (\nabla^2 F(\theta_j) - H) = 0$. The second term is a sum of martingale increments and shall be treated with Freedman inequality and Borel-Cantelli Lemma. Introduce the martingale increments

$$\forall 0 \leq j \leq k, \quad X_{j+1,k} = \nu_{j,k} (H(\theta_j, \xi'_{j+1}) - \nabla^2 F(\theta_j)).$$

For a fixed k , we have $X_{j+1,k} = \left(x_{j+1}^{(i,l)}\right)_{1 \leq i,l \leq d}$ where we remove the index k for the sake of clarity. Because the Hessian generator is unbiased, we have for all coordinates

$$\mathbb{E} \left[x_{j+1}^{(i,l)} | \mathcal{F}_j \right] = 0 \quad \text{for all } 0 \leq j \leq k.$$

By definition of the Hessian generator and using that $(\nabla^2 F(\theta_j))$ is bounded, we get that $\|H(\theta_j, \xi'_{j+1}) - \nabla^2 F(\theta_j)\| = O(1)$ for all $j \geq 0$. For any $b > 0$, consider the following event

$$\Omega_b = \left\{ \sup_{k \geq 0} \max_{j=0, \dots, k} (k+1) \left| x_{j+1}^{(i,l)} \right| \leq b \right\},$$

and note that since $\nu_{j,k} = O(1/k)$ we have $\mathbb{P}(\Omega_b) \rightarrow 1$ as $b \rightarrow \infty$. On this event, the martingale increments and the variance term are bounded as

$$\max_{j=0, \dots, k} \left| x_{j+1}^{(i,l)} \right| \leq b(k+1)^{-1}, \quad \sum_{j=0}^k \mathbb{E} \left[\left(x_{j+1}^{(i,l)} \right)^2 | \mathcal{F}_j \right] \leq b^2(k+1)^{-1}.$$

Using Freedman inequality (Theorem 8), we have for all coordinates $i, l = 1, \dots, d$,

$$\mathbb{P} \left(\left| \sum_{j=0}^k x_{j+1}^{(i,l)} \right| > \varepsilon, \Omega_b \right) \leq 2 \exp \left(-\frac{\varepsilon^2(k+1)}{2b(b+\varepsilon)} \right).$$

The last term is the general term of a convergent series. Apply Borel-Cantelli Lemma (Lemma 9) to finally get almost surely on Ω_b that $\lim_{k \rightarrow \infty} \sum_{j=0}^k x_{j+1}^{(i,l)} = 0$. Since $b > 0$ is arbitrary and $\mathbb{P}(\Omega_b) \rightarrow 1$ when $b \rightarrow \infty$, we have almost surely $\lim_{k \rightarrow \infty} \sum_{j=0}^k x_{j+1}^{(i,l)} = 0$. This is true for all the coordinates of the martingale increments and therefore

$$\lim_{k \rightarrow \infty} \sum_{j=0}^k \nu_{j,k} \left(H(\theta_j, \xi'_{j+1}) - \nabla^2 F(\theta_j) \right) = 0 \text{ a.s.}$$

□

E Auxiliary results

E.1 Robbins-Siegmund Theorem

Theorem 7. (*Robbins and Siegmund (1971)*) Consider a filtration $(\mathcal{F}_n)_{n \geq 0}$ and four sequences of random variables $(V_n)_{n \geq 0}$, $(W_n)_{n \geq 0}$, $(a_n)_{n \geq 0}$ and $(b_n)_{n \geq 0}$ that are adapted and non-negative. Assume that almost surely $\sum_k a_k < \infty$ and $\sum_k b_k < \infty$. Assume moreover that $\mathbb{E}[V_0] < \infty$ and

$$\forall n \in \mathbb{N}, \quad \mathbb{E}[V_{n+1} | \mathcal{F}_n] \leq (1 + a_n)V_n - W_n + b_n,$$

Then it holds

$$(a) \sum_k W_k < \infty \text{ a.s.} \quad (b) V_n \xrightarrow{\text{a.s.}} V_\infty, \mathbb{E}[V_\infty] < \infty. \quad (c) \sup_{n \geq 0} \mathbb{E}[V_n] < \infty.$$

E.2 Auxiliary lemmas

Lemma 4. Let $(u_n)_{n \geq 1}$, $(v_n)_{n \geq 1}$ and $(\alpha_n)_{n \geq 1}$ be non-negative sequences such that $\alpha_n \rightarrow 0$ and $\sum_n \alpha_n = +\infty$. Assume that there exists a real number $m \geq 1$ and $j \geq 1$ such that for all $n \geq j$,

$$u_n \leq (1 - \alpha_n)^m u_{n-1} + \alpha_n v_n,$$

then it holds that

$$\limsup_{n \rightarrow +\infty} u_n \leq \limsup_{n \rightarrow +\infty} v_n.$$

Proof. Denote $x_+ = \max(x, 0)$. One has $(x + y)_+ \leq x_+ + y_+$. Set $\varepsilon > 0$ and $v = \limsup_n v_n + \varepsilon$. Then there exists an integer $N \geq 1$ such that $(1 - \alpha_n)^m \leq (1 - \alpha_n)$ and $v_n < v$, i.e., $(v_n - v)_+ = 0$ for $n \geq N$. We have for large enough $n \geq N \vee j$,

$$u_n - v \leq (1 - \alpha_n)(u_{n-1} - v) + \alpha_n(v_n - v),$$

and taking the positive part gives

$$(u_n - v)_+ \leq (1 - \alpha_n)(u_{n-1} - v)_+ + \alpha_n(v_n - v)_+ = (1 - \alpha_n)(u_{n-1} - v)_+.$$

Since $\sum_n \alpha_n = +\infty$, this inequality implies that $(u_n - v)_+$ tends to zero, but this is true for all $\varepsilon > 0$ so v is arbitrarily close to $\limsup_n v_n$ and the result follows. \square

Lemma 5. Let $(\alpha_n)_{n \geq 1}$ be a non-negative sequence converging to zero, and λ, m and p three real numbers with $\lambda > 0, m \geq 1, p \geq 0$. Consider two non-negative sequences $(x_n), (\varepsilon_n)$ and an integer $j \geq 1$ such that

$$\begin{aligned} \forall n \geq j, \quad x_n &= (1 - \lambda\alpha_n)^m x_{n-1} + \alpha_n^{p+1} \varepsilon_n, \\ \text{i.e.,} \quad x_n &= \prod_{i=j}^n (1 - \lambda\alpha_i)^m x_{j-1} + \sum_{k=j}^n \alpha_k^{p+1} \left(\prod_{i=k+1}^n (1 - \lambda\alpha_i)^m \right) \varepsilon_k. \end{aligned}$$

The following holds

- if $\alpha_n = n^{-\beta}, \beta \in (1/2, 1)$, then for any p

$$\limsup_{n \rightarrow +\infty} \frac{x_n}{\alpha_n^p} \leq \frac{1}{m\lambda} \limsup_{n \rightarrow +\infty} \varepsilon_n.$$

- if $\alpha_n = 1/n$, then for any $p < m\lambda$

$$\limsup_{n \rightarrow +\infty} \frac{x_n}{\alpha_n^p} \leq \frac{1}{m\lambda - p} \limsup_{n \rightarrow +\infty} \varepsilon_n.$$

In particular, when $\varepsilon_n \rightarrow 0$ with $j = 1$ and $x_0 = 0$,

$$\begin{aligned} \lim_{n \rightarrow +\infty} \sum_{k=1}^n \alpha_k \prod_{i=k+1}^n (1 - \lambda\alpha_i)^m \varepsilon_k &= 0, \\ (m\lambda > 1) \lim_{n \rightarrow +\infty} \frac{1}{\alpha_n} \sum_{k=1}^n \alpha_k^2 \prod_{i=k+1}^n (1 - \lambda\alpha_i)^m \varepsilon_k &= 0. \end{aligned}$$

Before proving this result, note that if we consider $\alpha_n = \alpha/n^\beta$ then we can write

$$x_n = (1 - \lambda\alpha_n)^m x_{n-1} + \alpha_n^{p+1} \varepsilon_n = (1 - (\lambda\alpha)n^{-\beta})^m x_{n-1} + (n^{-\beta})^{p+1} (\alpha^{p+1} \varepsilon_n)$$

and apply the result with $\tilde{\lambda} = \lambda\alpha$ and $\tilde{\varepsilon}_n = \alpha^{p+1} \varepsilon_n$.

Proof. We apply Lemma 4 to the sequence $u_n = \frac{x_n}{\alpha_n^p}$. We have for all $n \geq j$,

$$\begin{aligned} u_n &= \frac{1}{\alpha_n^p} \left((1 - \lambda\alpha_n)^m x_{n-1} + \alpha_n^{p+1} \varepsilon_n \right) \\ &= \left(\frac{\alpha_{n-1}}{\alpha_n} \right)^p (1 - \lambda\alpha_n)^m u_{n-1} + \alpha_n \varepsilon_n \\ &= \exp \left(p \log \left(\frac{\alpha_{n-1}}{\alpha_n} \right) + m \log(1 - \lambda\alpha_n) \right) u_{n-1} + \alpha_n \varepsilon_n. \end{aligned}$$

Define

$$\lambda_n = \frac{1}{\alpha_n} \left(1 - \exp \left(p \log \left(\frac{\alpha_{n-1}}{\alpha_n} \right) + m \log(1 - \lambda\alpha_n) \right) \right),$$

so we get the recursion equation

$$\forall n \geq j, \quad u_n = (1 - \lambda_n \alpha_n) u_{n-1} + \lambda_n \alpha_n \frac{\varepsilon_n}{\lambda_n}.$$

• if $\alpha_n = n^{-\beta}$, $\beta \in (1/2, 1)$ then $1/\alpha_n - 1/\alpha_{n-1} \rightarrow 0$ and the ratio α_{n-1}/α_n tends to 1 with

$$\log\left(\frac{\alpha_{n-1}}{\alpha_n}\right) = \left(\frac{\alpha_{n-1}}{\alpha_n} - 1\right)(1 + o(1)) = \alpha_{n-1} \left(\frac{1}{\alpha_n} - \frac{1}{\alpha_{n-1}}\right)(1 + o(1)) = o(\alpha_n).$$

Besides, $m \log(1 - \lambda \alpha_n) = -m \lambda \alpha_n + o(\alpha_n)$ when $n \rightarrow +\infty$ and we get

$$\lambda_n = \frac{1}{\alpha_n} [1 - \exp(-m \lambda \alpha_n + o(\alpha_n))],$$

which implies that λ_n converges to $m\lambda$. We conclude with Lemma 4.

• if $\alpha_n = 1/n$ then the ratio α_{n-1}/α_n tends to 1 with

$$\log\left(\frac{\alpha_{n-1}}{\alpha_n}\right) = \log\left(1 + \frac{1}{n-1}\right) = \alpha_n + o(\alpha_n).$$

We still have $m \log(1 - \lambda \alpha_n) = -m \lambda \alpha_n + o(\alpha_n)$ when $n \rightarrow +\infty$ and therefore

$$\lambda_n = \frac{1}{\alpha_n} [1 - \exp((p - m\lambda)\alpha_n + o(\alpha_n))],$$

which implies λ_n converges to $(m\lambda - p)$ and we conclude in the same way. \square

Lemma 6. Let $A, B \in \mathcal{S}_d^{++}(\mathbb{R})$ be two real symmetric positive definite matrices then the eigenvalues of AB are real and positive with

$$Sp(AB) \subset [\lambda_{\min}(A)\lambda_{\min}(B); \lambda_{\max}(A)\lambda_{\max}(B)].$$

Proof. Denote by \sqrt{B} the unique positive square root of B . The matrix AB is similar to the real symmetric positive definite matrix $\sqrt{B}A\sqrt{B}$. Therefore its eigenvalues are real and positive. Since $A \mapsto \lambda_{\max}(A)$ is a sub-multiplicative matrix norm on $\mathcal{S}_d^{++}(\mathbb{R})$, $\lambda_{\max}(AB) \leq \lambda_{\max}(A)\lambda_{\max}(B)$ which gives $\lambda_{\max}((AB)^{-1}) \leq \lambda_{\max}(A^{-1})\lambda_{\max}(B^{-1})$, i.e., $\lambda_{\min}(AB)^{-1} \leq \lambda_{\min}(A)^{-1}\lambda_{\min}(B)^{-1}$, and finally $\lambda_{\min}(A)\lambda_{\min}(B) \leq \lambda_{\min}(AB)$. \square

Lemma 7. Let $S \in \mathcal{S}_d^{++}(\mathbb{R})$ be a real symmetric positive definite matrix. Let $(\alpha_k)_{k \geq 1}$ be a positive decreasing sequence converging to 0 such that $\sum_k \alpha_k = +\infty$. Denote by λ_m the smallest eigenvalue of S . It holds that there exists $j \geq 1$ such that for any $k > j$, all the eigenvalues of the real symmetric matrix $A_k = I - \alpha_k S$ are positive and we have

$$\begin{aligned} \rho(\Pi_n) &= \rho(A_n \dots A_1) \xrightarrow{n \rightarrow +\infty} 0, \\ \forall k > j, \quad \rho(\Pi_{n,k}) &= \rho(A_n \dots A_{k+1}) \leq \prod_{i=k+1}^n (1 - \alpha_i \lambda_m). \end{aligned}$$

Proof. For any $k \in \mathbb{N}$, the eigenvalues of the real symmetric matrix $A_k = I - \alpha_k S$ are given by $Sp(A_k) = \{(1 - \alpha_k \lambda), \lambda \in Sp(S)\}$. Since $\alpha_k \rightarrow 0$, there exists $j \geq 1$ such that $\alpha_k \lambda_m < 1$ for all $k > j$. Therefore for any $k > j$, we have $Sp(A_k) \subset \mathbb{R}_+^*$ and the largest eigenvalue is $\rho(A_k) = 1 - \alpha_k \lambda_m$. Since ρ is a sub-multiplicative norm for real symmetric matrices, we get $\rho(\Pi_n) \leq \prod_{k=1}^n \rho(A_k) = \prod_{k=1}^j \rho(A_k) \prod_{k=j+1}^n \rho(A_k)$. The second product can be upper bounded with the convexity of exponential,

$$\prod_{k=j+1}^n \rho(A_k) = \prod_{k=j+1}^n (1 - \alpha_k \lambda_m) \leq \prod_{k=j+1}^n \exp(-\alpha_k \lambda_m) = \exp(-\lambda_m(\tau_n - \tau_j)) \xrightarrow{n \rightarrow +\infty} 0.$$

Similarly we have for all $k > j$, $\rho(\Pi_{n,k}) \leq \prod_{i=k+1}^n \rho(A_i) \leq \prod_{i=k+1}^n (1 - \alpha_i \lambda_m)$. \square

Lemma 8. Let $\alpha_n = \alpha n^{-\beta}$ with $\beta \in (1/2, 1]$ then it holds

$$(\beta < 1) \sum_{k=1}^n \alpha_k \sim \frac{n\alpha_n}{1-\beta} = \frac{\alpha}{1-\beta} n^{1-\beta}, \quad (\beta = 1) \sum_{k=1}^n \alpha_k \sim \alpha \log(n).$$

Proof. By series-integral comprison, $\int_1^{n+1} t^{-\beta} dt \leq \sum_{k=1}^n k^{-\beta} \leq 1 + \int_1^n t^{-\beta} dt$. \square

Lemma 9. (*Borel, 1909*) Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $(A_k) \in \mathcal{F}^{\mathbb{N}}$.

$$\text{If } \sum_k \mathbb{P}(A_k) < \infty \text{ then } \mathbb{P}\left(\limsup_{k \rightarrow \infty} A_k\right) = 0.$$

In particular, for a sequence of random variables (X_k) and limit random variable X , define for $\varepsilon > 0$ the event $A_k(\varepsilon) = \{\omega : \|X_k(\omega) - X(\omega)\| > \varepsilon\}$. If $\sum_k \mathbb{P}(A_k(\varepsilon)) < \infty$ then $X_k \rightarrow X$ a.s.

Theorem 8. (*Delyon and Portier, 2021, Theorem 17*)(*Freedman inequality*) Let $(X_j)_{1 \leq j \leq n}$ be random variables such that $\mathbb{E}[X_j | \mathcal{F}_{j-1}] = 0$ for all $1 \leq j \leq n$ then, for all $t \geq 0$ and $v, m > 0$,

$$\mathbb{P}\left(\left|\sum_{j=1}^n X_j\right| \geq t, \max_{j=1, \dots, n} |X_j| \leq m, \sum_{j=1}^n \mathbb{E}[X_j^2 | \mathcal{F}_{j-1}] \leq v\right) \leq 2 \exp\left(-\frac{t^2/2}{v + tm/3}\right).$$

Lemma 10. Let $A \in \mathbb{R}^{n \times n}$ be a symmetric positive semi-definite matrix. Then for any $B \in \mathbb{R}^{m \times n}$, the matrix $BAB^T \in \mathbb{R}^{m \times m}$ is symmetric positive semi-definite.

Proof. First note that $(BAB^T)^T = (B^T)^T A^T B^T = BAB^T$ because A is symmetric. Then for any vector $x \in \mathbb{R}^m$, we have $x^T (BAB^T)x = (B^T x)^T A (B^T x) \geq 0$ since A is positive semi-definite. \square

Proposition 6. (*Khalil, 2002, Theorem 4.6*) Let H be a positive definite matrix and Γ a symmetric positive definite matrix of same dimension. Then there exists a symmetric positive definite matrix Σ , unique solution of the Lyapunov equation

$$H\Sigma + \Sigma H^T = \Gamma,$$

which is given by

$$\Sigma = \int_0^{+\infty} e^{-tH} \Gamma e^{-tH^T} dt.$$

The results remains true if the matrix Γ is only symmetric positive semi-definite: in that case the matrix Σ is also symmetric positive semi-definite and is the solution of the Lyapunov equation.

References

- M. Abadi, P. Barham, J. Chen, Z. Chen, A. Davis, J. Dean, M. Devin, S. Ghemawat, G. Irving, M. Isard, et al. Tensorflow: A system for large-scale machine learning. In *12th {USENIX} symposium on operating systems design and implementation ({OSDI} 16)*, pages 265–283, 2016.
- A. Agarwal, M. J. Wainwright, P. L. Bartlett, and P. K. Ravikumar. Information-theoretic lower bounds on the oracle complexity of convex optimization. In *Advances in Neural Information Processing Systems*, pages 1–9, 2009.
- S.-I. Amari. Natural gradient works efficiently in learning. *Neural computation*, 10(2):251–276, 1998.
- J. Baxter and P. L. Bartlett. Infinite-horizon policy-gradient estimation. *Journal of Artificial Intelligence Research*, 15:319–350, 2001.
- S. Becker, Y. Le Cun, et al. Improving the convergence of back-propagation learning with second order methods. In *Proceedings of the 1988 connectionist models summer school*, pages 29–37, 1988.
- M. Benaïm. Dynamics of stochastic approximation algorithms. In *Seminaire de probabilités XXXIII*, pages 1–68. Springer, 1999.
- A. Benveniste, M. Métivier, and P. Priouret. *Adaptive algorithms and stochastic approximations*, volume 22. Springer Science & Business Media, 2012.
- B. Bercu, A. Godichon, and B. Portier. An efficient stochastic newton algorithm for parameter estimation in logistic regressions. *SIAM Journal on Control and Optimization*, 58(1):348–367, 2020.
- D. P. Bertsekas and J. N. Tsitsiklis. Gradient convergence in gradient methods with errors. *SIAM Journal on Optimization*, 10(3):627–642, 2000.
- M. É. Borel. Les probabilités dénombrables et leurs applications arithmétiques. *Rendiconti del Circolo Matematico di Palermo (1884-1940)*, 27(1):247–271, 1909.
- L. Bottou, F. E. Curtis, and J. Nocedal. Optimization methods for large-scale machine learning. *Siam Review*, 60(2):223–311, 2018.
- C. Boyer and A. Godichon-Baggioni. On the asymptotic rate of convergence of stochastic newton algorithms and their weighted averaged versions. *arXiv preprint arXiv:2011.09706*, 2020.
- C. G. Broyden. The convergence of a class of double-rank minimization algorithms 1. general considerations. *IMA Journal of Applied Mathematics*, 6(1):76–90, 1970.
- R. H. Byrd, G. M. Chin, W. Neveitt, and J. Nocedal. On the use of stochastic hessian information in optimization methods for machine learning. *SIAM Journal on Optimization*, 21(3):977–995, 2011.
- C.-C. Chang and C.-J. Lin. Libsvm: a library for support vector machines. *ACM transactions on intelligent systems and technology (TIST)*, 2(3):1–27, 2011.
- S. Cléménçon, P. Bertail, E. Chautru, and G. Papa. Optimal survey schemes for stochastic gradient descent with applications to m-estimation. *ESAIM: Probability and Statistics*, 23:310–337, 2019.
- A. Defazio, F. Bach, and S. Lacoste-Julien. Saga: A fast incremental gradient method with support for non-strongly convex composite objectives. In *Advances in neural information processing systems*, pages 1646–1654, 2014.
- B. Delyon. General results on the convergence of stochastic algorithms. *IEEE Transactions on Automatic Control*, 41(9):1245–1255, 1996.
- B. Delyon and A. Juditsky. Stochastic optimization with averaging of trajectories. *Stochastics: An International Journal of Probability and Stochastic Processes*, 39(2-3):107–118, 1992.
- B. Delyon and F. Portier. Asymptotic optimality of adaptive importance sampling. In *Proceedings of the 32nd International Conference on Neural Information Processing Systems*, pages 3138–3148. Curran Associates Inc., 2018.
- B. Delyon and F. Portier. Safe adaptive importance sampling: A mixture approach. *The Annals of Statistics*, 49(2):885–917, 2021.
- A. Dieuleveut, A. Durmus, and F. Bach. Bridging the gap between constant step size stochastic gradient descent and markov chains. *Annals of Statistics*, 48(3):1348–1382, 2020.

- D. Dua and C. Graff. UCI machine learning repository, 2017. URL <http://archive.ics.uci.edu/ml>.
- J. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of machine learning research*, 12(Jul):2121–2159, 2011.
- M. Duflo. *Random iterative models*, volume 34. Springer Science & Business Media, 2013.
- V. Fabian. On asymptotic normality in stochastic approximation. *The Annals of Mathematical Statistics*, 39(4):1327–1332, 1968.
- V. Fabian. Asymptotically efficient stochastic approximation; the rm case. *The Annals of Statistics*, 1(3):486–495, 1973.
- R. Fletcher. A new approach to variable metric algorithms. *The computer journal*, 13(3):317–322, 1970.
- S. Gadat and F. Panloup. Optimal non-asymptotic bound of the ruppert-polyak averaging without strong convexity. *arXiv preprint arXiv:1709.03342*, 2017.
- S. Gadat, F. Panloup, S. Saadane, et al. Stochastic heavy ball. *Electronic Journal of Statistics*, 12(1):461–529, 2018.
- N. Gazagnadou, R. M. Gower, and J. Salmon. Optimal mini-batch and step sizes for saga. *arXiv preprint arXiv:1902.00071*, 2019.
- D. Goldfarb. A family of variable-metric methods derived by variational means. *Mathematics of computation*, 24(109):23–26, 1970.
- R. M. Gower, N. Loizou, X. Qian, A. Sailanbayev, E. Shulgin, and P. Richtárik. Sgd: General analysis and improved rates. In *International Conference on Machine Learning*, pages 5200–5209. PMLR, 2019.
- R. M. Gower, P. Richtárik, and F. Bach. Stochastic quasi-gradient methods: Variance reduction via jacobian sketching. *Mathematical Programming*, pages 1–58, 2020.
- P. Hall and C. Heyde. *Martingale Limit Theory and Its Application*. Probability and mathematical statistics. Academic Press, 1980. ISBN 9781483240244. URL <https://books.google.fr/books?id=wdLajgEACAAJ>.
- D. Harrison Jr and D. L. Rubinfeld. Hedonic housing prices and the demand for clean air. *Journal of environmental economics and management*, 5(1):81–102, 1978.
- R. Johnson and T. Zhang. Accelerating stochastic gradient descent using predictive variance reduction. In *Advances in neural information processing systems*, pages 315–323, 2013.
- S. M. Kakade. A natural policy gradient. In *Advances in neural information processing systems*, pages 1531–1538, 2002.
- H. Karimi, J. Nutini, and M. Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, pages 795–811. Springer, 2016.
- H. K. Khalil. *Nonlinear systems; 3rd ed.* Prentice-Hall, Upper Saddle River, NJ, 2002. URL <https://cds.cern.ch/record/1173048>.
- J. Kiefer, J. Wolfowitz, et al. Stochastic estimation of the maximum of a regression function. *The Annals of Mathematical Statistics*, 23(3):462–466, 1952.
- H. J. Kushner and D. S. Clark. Stochastic approximation methods for constrained and unconstrained systems. 1978.
- H. J. Kushner and H. Huang. Rates of convergence for stochastic approximation type algorithms. *SIAM Journal on Control and Optimization*, 17(5):607–617, 1979.
- Y. A. LeCun, L. Bottou, G. B. Orr, and K.-R. Müller. Efficient backprop. In *Neural networks: Tricks of the trade*, pages 9–48. Springer, 2012.
- D. C. Liu and J. Nocedal. On the limited memory bfgs method for large scale optimization. *Mathematical programming*, 45(1):503–528, 1989.

- P. Moritz, R. Nishihara, and M. Jordan. A linearly-convergent stochastic l-bfgs algorithm. In *Artificial Intelligence and Statistics*, pages 249–258. PMLR, 2016.
- E. Moulines and F. R. Bach. Non-asymptotic analysis of stochastic approximation algorithms for machine learning. In *Advances in Neural Information Processing Systems*, pages 451–459, 2011.
- A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on optimization*, 19(4):1574–1609, 2009.
- A. S. Nemirovski and D. B. Yudin. Problem complexity and method efficiency in optimization. 1983.
- Y. Nesterov. *Introductory lectures on convex optimization: A basic course*, volume 87. Springer Science & Business Media, 2013.
- M. B. Nevelson and R. Z. Khas'minskiĭ. *Stochastic approximation and recursive estimation*, volume 47. American Mathematical Soc., 1976.
- L. M. Nguyen, P. H. Nguyen, M. van Dijk, P. Richtárik, K. Scheinberg, and M. Takáč. Sgd and hogwild! convergence without the bounded gradients assumption. *arXiv preprint arXiv:1802.03801*, 2018.
- G. Papa, P. Bianchi, and S. Cléménçon. Adaptive sampling for incremental optimization using stochastic gradient descent. In *International Conference on Algorithmic Learning Theory*, pages 317–331. Springer, 2015.
- F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, et al. Scikit-learn: Machine learning in python. *the Journal of machine Learning research*, 12:2825–2830, 2011.
- M. Pelletier. On the almost sure asymptotic behaviour of stochastic algorithms. *Stochastic processes and their applications*, 78(2):217–244, 1998a.
- M. Pelletier. Weak convergence rates for stochastic approximation with application to multiple targets and simulated annealing. *Annals of Applied Probability*, pages 10–44, 1998b.
- B. T. Polyak. A new method of stochastic approximation type. *Avtomatika i telemekhanika*, (7):98–107, 1990.
- B. T. Polyak and A. B. Juditsky. Acceleration of stochastic approximation by averaging. *SIAM Journal on Control and Optimization*, 30(4):838–855, 1992.
- H. Robbins and S. Monro. A stochastic approximation method. *The annals of mathematical statistics*, pages 400–407, 1951.
- H. Robbins and D. Siegmund. A convergence theorem for non negative almost supermartingales and some applications. In *Optimizing methods in statistics*, pages 233–257. Elsevier, 1971.
- J. Sacks. Asymptotic distribution of stochastic approximation procedures. *The Annals of Mathematical Statistics*, 29(2):373–405, 1958.
- M. Schmidt and N. L. Roux. Fast convergence of stochastic gradient descent under a strong growth condition. *arXiv preprint arXiv:1308.6370*, 2013.
- N. N. Schraudolph, J. Yu, and S. Günter. A stochastic quasi-newton method for online convex optimization. In *Artificial intelligence and statistics*, pages 436–443. PMLR, 2007.
- S. Shalev-Shwartz and T. Zhang. Stochastic dual coordinate ascent methods for regularized loss minimization. *Journal of Machine Learning Research*, 14(Feb):567–599, 2013.
- S. Shalev-Shwartz, Y. Singer, N. Srebro, and A. Cotter. Pegasos: Primal estimated sub-gradient solver for svm. *Mathematical programming*, 127(1):3–30, 2011.
- D. F. Shanno. Conditioning of quasi-newton methods for function minimization. *Mathematics of computation*, 24(111):647–656, 1970.
- A. Shapiro, D. Dentcheva, and A. Ruszczyński. *Lectures on stochastic programming: modeling and theory*. SIAM, 2014.
- R. S. Sutton and A. G. Barto. *Reinforcement learning: An introduction*. MIT press, 2018.
- J. H. Venter. An extension of the robbins-monro procedure. *The Annals of Mathematical Statistics*, 38(1):181–190, 1967.

- C. Wei. Multivariate adaptive stochastic approximation. *The Annals of Statistics*, 15(3):1115–1130, 1987.
- R. J. Williams. Simple statistical gradient-following algorithms for connectionist reinforcement learning. *Machine learning*, 8(3-4):229–256, 1992.
- M. Zedek. Continuity and location of zeros of linear combinations of polynomials. *Proceedings of the American Mathematical Society*, 16(1):78–84, 1965.
- T. Zhang. Solving large scale linear prediction problems using stochastic gradient descent algorithms. In *Proceedings of the twenty-first international conference on Machine learning*, page 116, 2004.