

On Hausdorff operators on homogeneous spaces of locally compact groups

A. R. Mirotin

amirotin@yandex.ru

Abstract. Hausdorff operators on the real line and multidimensional Euclidean spaces originated from some classical summation methods. Now it is an active research area. Hausdorff operators on general groups were defined and studied by the author since 2019. The purpose of this paper is to define and study Hausdorff operators on Lebesgue and real Hardy spaces over homogeneous spaces of locally compact groups. We introduce in particular an atomic Hardy space over homogeneous spaces of locally compact groups and obtain conditions for boundedness of Hausdorff operators on such spaces. Several corollaries are considered and unsolved problems are formulated.

Key words. Hausdorff operator, locally compact group, homogeneous space, atomic Hardy space, Lebesgue space, bounded operator.

1 Introduction and preliminaries

Hausdorff operators were introduced by Hardy [1, Chapter XI] on the segment, and by C. Georgakis [2] and independently by E. Liflyand and F. Moricz [3] on the whole real line. Their multidimensional generalizations were considered later by Liflyand and Lerner [4]. Now it is an active research area. It is enough to note that the Google search by request "Hausdorff operator" gives more than 1 200 000 results. See also survey articles [5], [6] for historical remarks and the state of the art up to 2014.

Hausdorff operators on general groups were defined and studied by the author in [7] and [8]. The purpose of this paper is to define and study Hausdorff operators on Lebesgue and real Hardy spaces over homogeneous spaces of locally compact groups.

In what follows G stands for a locally compact group with left Haar measure ν . We denote by $\text{Aut}(G)$ the space of all topological automorphisms of G endowed with its natural topology (see, e.g. [9]), $\mathcal{L}(Y)$ denotes the space of linear bounded operators on a normed space Y .

In [7] the next definition was proposed.

Definition [7]. Let (T, m) be a measure space, G a topological group, $A : T \rightarrow \text{Aut}(G)$ a measurable map, and Ψ a locally integrable function on

T . We define the *Hausdorff operator* with the kernel Ψ over the group G by the formula

$$\mathcal{H}f(x) = \int_T \Psi(t)f(A(t)(x))dm(t).$$

By [7, Lemma 1] this operator is bounded in $L^p(G)$ ($1 \leq p \leq \infty$) provided $\Psi(t)(\text{mod}A(t))^{-1/p} \in L^1(T, m)$ and

$$\|\mathcal{H}\|_{\mathcal{L}(L^p(G))} \leq \int_T |\Psi(t)|(\text{mod}A(t))^{-1/p} dm(t).$$

Moreover, in [7], [8] conditions for boundedness of Hausdorff operators on the real Hardy space $H^1(G)$ over metrizable locally compact group G with doubling condition were obtained.

The aim of this work is to define Hausdorff operators on homogeneous spaces of locally compact groups and to prove analogs of aforementioned results for this situation.

Let K be a compact subgroup of G with normalized Haar measure β . Consider the quotient space G/K of left cosets $\dot{x} := xK = \pi_K(x)$ ($x \in G$) where $\pi_K : G \rightarrow G/K$ stands for a natural projection. We shall assume that the measure ν is normalized in such a way that (generalized) Weil's formula

$$\int_G g(x)dx = \int_{G/K} \left(\int_K g(xk)dk \right) d\lambda(\dot{x}) \quad (1)$$

holds for all $g \in L^1(G)$, where λ denotes some left- G invariant measure on G/K (see, e. g., [10, Chapter VII, §2, No. 5, Theorem 2] and especially remark c) after this theorem).¹ Here and below we write dx instead of $d\nu(x)$ and dk instead of $d\beta(k)$. We shall write also $d\dot{x}$ instead of $d\lambda(\dot{x})$.

The function $g : G \rightarrow \mathbb{C}$ is called *right- K -invariant* if $g(xk) = g(x)$ for all $x \in G$, $k \in K$. For such a function we put $\dot{g}(\dot{x}) := g(x)$. This definition is correct and for $g \in L^1(G)$ formula (1) implies that

$$\int_G g(x)dx = \int_{G/K} \dot{g}(\dot{x})d\dot{x} \quad (2)$$

(recall that $\int_K dk = 1$).

The map $g \mapsto \dot{g}$ is a bijection between the set of all right- K -invariant functions on G (all right- K -invariant functions from $L^1(G)$) and the set of all functions on G/K (respectively functions from $L^1(G/K, \lambda)$).

¹ G -left invariance of λ means that $\lambda(xE) = \lambda(E)$ for every Borel subset E of G/K and for every $x \in G$. This measure is unique up to constant multiplier.

Let an automorphism $A \in \text{Aut}(G)$ maps K onto itself. Since

$$A(\dot{x}) := A(xK) = \{A(x)A(k) : k \in K\} = A(x)K = \pi_K(A(x))$$

we get a homeomorphism $\dot{A} : G/K \rightarrow G/K$, $\dot{A}(\dot{x}) := \pi_K(A(x))$. Then for every right- K -invariant function g on G we have $\dot{g}(\dot{A}(\dot{x})) = g(A(x))$.

From now on we put

$$\text{Aut}_K(G) := \{\dot{A} : A \in \text{Aut}(G), A(K) = K\}.$$

Definition 1. Let (Ω, μ) be a measure space, $(\dot{A}(u))_{u \in \Omega} \subset \text{Aut}_K(G)$ a family of homeomorphisms of G/K , and $\Phi \in L^1_{loc}(\Omega, \mu)$. For a function f on G/K we define a *Hausdorff operator* on G/K as follows

$$(\mathcal{H}_{\Phi, \dot{A}} f)(\dot{x}) := \int_{\Omega} \Phi(u) f(\dot{A}(u)(\dot{x})) d\mu(u).$$

As was mentioned by Hardy in the case $\Omega = [0, 1]$ [1, Theorem 217] his Hausdorff operator possesses some regularity property. A Hausdorff operator in the sense of Definition 1 also enjoys this property as the next proposition shows.

Proposition 1. *Suppose that the conditions of Definition 1 are fulfilled and the group G is σ -compact. In order that the transformation $\mathcal{H}_{\Phi, \dot{A}}$ should be regular, i.e. that $f \in C(G/K)$, $f(\dot{x}) \rightarrow l$ when $\dot{x} \rightarrow \infty$ should imply $\mathcal{H}_{\Phi, \dot{A}} f(\dot{x}) \rightarrow l$, it is necessary and sufficient that $\int_{\Omega} \Phi(u) d\mu(u) = 1$.*

Proof. If $f(x) = 1$ then $\mathcal{H}_{\Phi, \dot{A}} f(\dot{x}) = \int_{\Omega} \Phi(u) d\mu(u)$. Thus, $\int_{\Omega} \Phi(u) d\mu(u) = 1$ is a necessary condition.

To prove the sufficiency, first note that since $\dot{A}(u)$ has continuous inverse, $f(\dot{x}) \rightarrow l$ when $\dot{x} \rightarrow \infty$ implies $f(\dot{A}(u)\dot{x}) \rightarrow l$ when $\dot{x} \rightarrow \infty$. Indeed, $f(\dot{x}) \rightarrow l$ when $\dot{x} \rightarrow \infty$ means that for every $\varepsilon > 0$ there is such compact $C_{\varepsilon} \subset G/K$ that $|f(\dot{x}) - l| < \varepsilon$ for $x \in G \setminus C_{\varepsilon}$. Now if $\dot{x} \in (G/K) \setminus \dot{A}(u)^{-1}(C_{\varepsilon})$ we get $\dot{A}(u)\dot{x} \in (G/K) \setminus C_{\varepsilon}$ and therefore $|f(\dot{A}(u)\dot{x}) - l| < \varepsilon$.

But if, in addition, $f \in C(G/K)$ the function f is bounded on G/K and therefore $\mathcal{H}_{\Phi, \dot{A}} f(\dot{x}) \rightarrow l$ by the Lebesgue Theorem (one can apply the Lebesgue Theorem, since G/K is σ -compact).

Thus, Proposition 1 shows that Hausdorff operators in a sense of Definition 1 gives us a family (for various Φ , $\dot{A}(u)$, and Ω) of generalized limits at infinity for functions on G/K .

2 Hausdorff operators on $L^p(G/K)$

In the following we put $L^p(G/K) := L^p(G/K, \lambda)$ ($p \in [1, \infty]$). Formula (2) implies that $\|g\|_{L^p(G)} = \|\dot{g}\|_{L^p(G/K)}$. For every right- K -invariant function $g \in L^p(G)$.

In this section we give conditions of boundedness of Hausdorff operators on $L^p(G/K)$.

Let (Ω, μ) and $(\dot{A}(u))_{u \in \Omega}$ be as in Definition 1. For a function Φ on Ω let

$$\|\Phi\|_{p,A} := \int_{\Omega} |\Phi(u)| (\text{mod} A(u))^{-1/p} d\mu(u).$$

Theorem 1. *Suppose that the conditions of Definition 1 are fulfilled, $p \in [1, \infty]$, and $\|\Phi\|_{p,A} < \infty$. Then $\mathcal{H}_{\Phi, \dot{A}}$ is bounded in $L^p(G/K)$ and*

$$\|\mathcal{H}_{\Phi, \dot{A}}\|_{\mathcal{L}(L^p(G/K))} \leq \|\Phi\|_{p,A}.$$

Proof. Let $1 < p < \infty$. Every function $f \in L^p(G/K)$ has the form $f = \dot{g}$ for a unique right K -invariant function $g \in L^p(G)$. Using Minkowski's integral inequality we have that

$$\begin{aligned} \|\mathcal{H}_{\Phi, \dot{A}} f\|_{L^p(G/K)} &= \left(\int_{G/K} \left| \int_{\Omega} \Phi(u) f(\dot{A}(u)(\dot{x})) d\mu(u) \right|^p d\dot{x} \right)^{1/p} \\ &\leq \int_{\Omega} |\Phi(u)| \left(\int_{G/K} |f(\dot{A}(u)(\dot{x}))|^p d\dot{x} \right)^{1/p} d\mu(u). \end{aligned}$$

Since the function $x \mapsto g(A(u)(x))$ is right K -invariant, as well, formula (2) yields

$$\int_{G/K} |f(\dot{A}(u)(\dot{x}))|^p d\dot{x} = \int_G |g(A(u)(x))|^p dx.$$

On the other hand, by [10, VII.1.4, formula (31)] we have

$$\int_G |g(A(u)(x))|^p dx = (\text{mod} A(u))^{-1} \int_G |g(x)|^p dx.$$

Thus,

$$\begin{aligned} \|\mathcal{H}_{\Phi, \dot{A}} f\|_{L^p(G/K)} &\leq \int_{\Omega} |\Phi(u)| (\text{mod} A(u))^{-1/p} d\mu(u) \left(\int_G |g(x)|^p dx \right)^{1/p} \\ &= \|\Phi\|_{p,A} \left(\int_{G/K} |f(\dot{x})|^p d\dot{x} \right)^{1/p} = \|\Phi\|_{p,A} \|f\|_{L^p(G/K)}. \end{aligned}$$

For $p = 1$ the statement of theorem 1 follows from Fubini theorem and for $p = \infty$ it is obvious.

The following simple example is intended to illustrate preceding constructions.

Example 1. Let G be the multiplicative group $\mathbb{C}^\times$ of the complex field $\mathbb{C}$ and $K := \{z \in \mathbb{C}^\times : |z| = 1\}$ the circle group. Then G/K can be identified with $(0, \infty)$ via the map $zK = \dot{z} \mapsto r$ where $r = |z|$. In other words, we use $(0, \infty)$ as a model of G/K , the positive number r representing the circle of radius r . Automorphisms of G have the form $A(re^{i\alpha}) = r^u e^{i\alpha}$ or $A(re^{i\alpha}) = r^u e^{-i\alpha}$ ($u \in \mathbb{R}$). Thus $\text{Aut}(G)/K = \text{Aut}(G)$. It follows that $\dot{A}(\dot{z}) = r^u$ ($u \in \mathbb{R}$). So the general form of a Hausdorff operator on $(0, \infty)$ looks as follows ($f : (0, \infty) \rightarrow \mathbb{C}$, $r > 0$)

$$\mathcal{H}_{\Phi, \dot{A}} f(r) = \int_{\mathbb{R}} \Phi(u) f(r^u) d\mu(u)$$

(we take $\Omega = \mathbb{R}$, and μ any positive measure on $\mathbb{R}$). Since G is commutative, $\text{mod} A = 1$ for all $A \in \text{Aut}(G)$. So Theorem 1 implies that $\mathcal{H}_{\Phi, \dot{A}}$ is bounded on $L^p(0, \infty)$ if $\Phi \in L^1(\mu)$ and in this case $\|\mathcal{H}_{\Phi, \dot{A}}\|_{\mathcal{L}(L^p)} \leq \|\Phi\|_{L^1(\mu)}$.

If we take in Theorem 1 the space $\mathbb{Z}_+$ (endowed with counting measure) as Ω , we arrive at the following

Corollary 1. *Let $\Phi(j)$ be a sequence of complex numbers. Consider a discrete Hausdorff operator over G/K* ²

$$\mathcal{H}_{\Phi, \dot{A}} f(x) := \sum_{j=0}^{\infty} \Phi(j) f(\dot{A}(j)(x)).$$

Then

$$\|\mathcal{H}_{\Phi, \dot{A}}\|_{\mathcal{L}(L^p(G/K))} \leq \sum_{j=0}^{\infty} |\Phi(j)| (\text{mod} A(j))^{-1/p}.$$

Putting

$$\Phi = \frac{\chi_{\{\text{mod} A(u) \geq 1\}}}{\text{mod} A}$$

in Definition 1 (χ_E denotes the indicator function of the set E) we get a *Césaro operator* over G/K (cf. [7])

$$\mathcal{C}_{A, \mu} f(x) := \int_{\{\text{mod} A(u) \geq 1\}} \frac{f(\dot{A}(u)(x))}{\text{mod} A(u)} d\mu(u).$$

Corollary 2. *For a Césaro operator over G/K the following estimate holds*

$$\|\mathcal{C}_{A, \mu}\|_{\mathcal{L}(L^p(G/K))} \leq \int_{\{\text{mod} A(u) \geq 1\}} \frac{d\mu(u)}{(\text{mod} A(u))^{1+1/p}}.$$

² Discrete Hausdorff operators were introduced in [11], [12].

3 Hausdorff operators on Hardy space $H^1(G/K)$

The goal of this section is to introduce the atomic Hardy space $H^1(G/K)$ and to obtain conditions for boundedness of Hausdorff operators on this space.

In the rest of the paper as in [8] we assume in addition that G is metrizable via a left invariant metric ρ and the following *doubling condition* in a sense of [13] holds:

there exists a constant C such that

$$\nu(B(x, 2r)) \leq C\nu(B(x, r))$$

for each $x \in G$ and $r > 0$.

Here $B(x, r)$ denotes the ball of radius r around x . The *doubling constant* is the smallest constant $C \geq 1$ for which the last inequality holds. We denote this constant by C_ν . Then for each $x \in G, k \geq 1$ and $r > 0$

$$\nu(B(x, kr)) \leq C_\nu k^s \nu(B(x, r)), \quad (3)$$

where $s = \log_2 C_\nu$ (see, e.g., [14, p. 76]). The number s takes the role of a "dimension" for a doubling metric measure space G .

To introduce the space $H^1(G/K)$ first recall that a function a on G is an $((1, \infty)$ -)atom if

- (i) the support of a is contained in a ball $B(x, r)$;
- (ii) $\|a\|_\infty \leq \frac{1}{\nu(B(x, r))}$;
- (iii) $\int_G a(x) dx = 0$.

In case $\nu(G) < \infty$ we shall assume $\nu(G) = 1$. Then the constant function having value 1 is also considered to be an atom.

Definition 2. We define the Hardy space $H^1(G/K)$ as a space of such functions $f = \dot{g}$ on G/K that g admits an atomic decomposition of the form

$$g = \sum_{j=1}^{\infty} \alpha_j a_j, \quad (4)$$

where a_j are right- K -invariant atoms and $\sum_{j=1}^{\infty} |\alpha_j| < \infty$. In this case,

$$\|f\|_{H^1(G/K)} := \inf \sum_{j=1}^{\infty} |\alpha_j|,$$

and infimum is taken over all decompositions above of g .

Thus a function $f = \dot{g}$ from $H^1(G/K)$ admits an atomic decomposition $f = \sum_{j=1}^{\infty} \alpha_j \dot{a}_j$ such that $\sum_{j=1}^{\infty} |\alpha_j| < \infty$, and $\|f\|_{H^1(G/K)} = \|g\|_{H^1(G)}$.

Remark 1. Real Hardy spaces over compact connected (not necessary quasi-metric) Abelian groups were defined in [15]. The case of semisimple Lie groups was considered earlier in [16].

Proposition 2. *The space $H^1(G/K)$ is Banach. If for some $h \in H^1(G)$ the inequality $\int_K h(k)dk \neq 0$ holds the space $H^1(G/K)$ is nontrivial.*

Proof. We prove that right- K -invariant atoms exist. To this end for an atom a on G let's consider the function

$$a'(x) := c \int_K a(xk)dk.$$

Then a' is right- K -invariant and satisfies (i) for every constant $c > 0$. Indeed, if a is supported in a ball $B = B(e, r_B)$, then $a'(x) = 0$ for $x \notin BK$. Since

$$\rho(e, xk) \leq \rho(e, x) + \rho(x, xk) = \rho(e, x) + \rho(e, k) < r_B + \text{dist}(e, K) =: r'_B$$

for $x \in B$, the set BK is contained in a ball $B' = B(e, r'_B)$. Thus, a' is supported in B' . From now on we assume that $c = \nu(B)/\nu(B')$. Then (ii) holds for a' because $\|a'\|_\infty \leq c\|a\|_\infty \leq c/\nu(B) = 1/\nu(B')$ for such c . The property (iii) for a' follows from

$$\int_G a'(x)dx = c \int_G \int_K a(xk)dkdx = c \int_K \Delta_G(k) \int_G a(x)dxdk = 0.$$

Finally, we shall show that $H^1(G/K)$ is complete. First note that since $\|a\|_{L^1(G)} \leq 1$ for each atom a , we have $\|g\|_{L^1(G)} \leq \|\dot{g}\|_{H^1(G/K)}$ for each right- K -invariant function $g \in H^1(G)$. Then for a function $f = \dot{g}$ we have $\|f\|_{L^1(G/K)} \leq \|f\|_{H^1(G/K)}$.

Let a sequence $f_j \in H^1(G/K)$ be such that $\sum_j \|f_j\|_{H^1(G/K)} < \infty$. It is enough to prove that the series $\sum_j f_j$ converges in $H^1(G/K)$. The sequence S_n of partial sums of this series is a Cauchy sequence in $L^1(G/K)$ because for $m < n$

$$\|S_n - S_m\|_{L^1(G/K)} \leq \sum_{j=m+1}^n \|f_j\|_{L^1(G/K)} \leq \sum_{j=m+1}^n \|f_j\|_{H^1(G/K)}.$$

So the series $\sum_j f_j$ converges in $L^1(G/K)$ to some function f . On the other hand, each f_j has an atomic decomposition $f_j = \sum_i \alpha_{ij} \dot{a}_{ji}$ such that $\sum_i |\alpha_{ij}| < 2\|f_j\|_{H^1(G/K)}$. Then f has an atomic decomposition

$$f = \sum_j \sum_i \alpha_{ij} \dot{a}_{ji},$$

and

$$\|f\|_{H^1(G/K)} \leq \sum_j \sum_i |\alpha_{ij}| < 2 \sum_j \|f_j\|_{H^1(G/K)} < \infty.$$

Thus $f \in H^1(G/K)$. Moreover,

$$\left\| f - \sum_{j=1}^n f_j \right\|_{H^1(G/K)} \leq \sum_{j=n+1}^{\infty} \|f_j\|_{H^1(G/K)} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

This completes the proof.

In the proof of Theorem 2 the next lemmas play an important role.

Lemma 1. [7] *Let (Ω, q, μ) be σ -compact quasi-metric space with quasi-metric q and positive Radon measure μ , (X, m) be a measure space and $\mathcal{F}(X)$ be some Banach space of m -measurable functions on X . Assume that the convergence of a sequence strongly in $\mathcal{F}(X)$ yields the convergence of some subsequence to the same function for m -almost all $x \in X$. Let $F(u, x)$ be a function such that $F(u, \cdot) \in \mathcal{F}(X)$ for μ -almost all $u \in \Omega$ and the map $u \mapsto F(u, \cdot) : \Omega \rightarrow \mathcal{F}(X)$ is Bochner integrable with respect to μ . Then for m -almost all $x \in X$*

$$\left((B) \int_{\Omega} F(u, \cdot) d\mu(u) \right) (x) = \int_{\Omega} F(u, x) d\mu(u).$$

Lemma 2. [8] *Every automorphism $A \in \text{Aut}(G)$ is Lipschitz. Moreover, one can choose the Lipschitz constant to be*

$$L_A = \kappa_{\rho} \bmod A,$$

where the constant κ_{ρ} depends on the metric ρ only.

Now we are in position to prove the following

Theorem 2. *Under assumptions of Definition 1 let (Ω, q, μ) be σ -compact quasi-metric space with quasi-metric q and positive Radon measure μ and $\Phi \in L^1(k^s \mu)$ where $k(u) := \kappa_{\rho} / \bmod(A(u))$. Then the operator $\mathcal{H}_{\Phi, A}$ is bounded on the space $H^1(G/K)$ and*

$$\|\mathcal{H}_{\Phi, A}\|_{\mathcal{L}(H^1(G/K))} \leq C_{\nu} \|\Phi\|_{L^1(k^s \mu)}.$$

Proof. We proceed as in the proof of main theorem in [7], [8]. If we set $X = G/K$ and $m = \lambda$ the pair (X, m) satisfies the conditions of Lemma 1 with $H^1(G/K)$ in place of $\mathcal{F}(X)$. Indeed, let $f_n = \dot{g}_n \in H^1(G/K)$, $f = \dot{g} \in H^1(G/K)$, and $\|f_n - f\|_{H^1(G/K)} \rightarrow 0$ ($n \rightarrow \infty$). Since

$$\|f_n - f\|_{L^1(G/K)} = \int_{G/K} |\pi_K(g_n - g)| d\lambda$$

$$= \int_G |g_n(x) - g(x)| dx \leq \|g_n - g\|_{H^1(G)} = \|f_n - f\|_{H^1(G/K)} \rightarrow 0,$$

there is a subsequence of f_n that converges to f λ -a.e.

Then Definition 2 and lemma 1 imply for $f \in H^1(G/K)$ that

$$\mathcal{H}_{\Phi, \dot{A}} f = \int_{\Omega} \Phi(u) f \circ \dot{A}(u) d\mu(u)$$

(the Bochner integral).

Therefore (below $f = \dot{g}$)

$$\begin{aligned} \|\mathcal{H}_{\Phi, \dot{A}} f\|_{H^1(G/K)} &\leq \int_{\Omega} |\Phi(u)| \|f \circ \dot{A}(u)\|_{H^1(G/K)} d\mu(u) \\ &= \int_{\Omega} |\Phi(u)| \|g \circ A(u)\|_{H^1(G)} d\mu(u). \end{aligned} \quad (5)$$

If g has representation (4) then

$$g \circ A(u) = \sum_{j=1}^{\infty} \alpha_j a_j \circ A(u). \quad (6)$$

We claim that $b_{j,u} := C_{\nu}^{-1} k(u)^{-s} a_j \circ A(u)$ is an atom, too. Indeed, Lemma 2 implies that

$$A(u)^{-1} B(x, r) \subseteq B(x', k(u)r),$$

where $x' = A(u)^{-1}(x)$. If a_j is supported in $B(x_j, r_j)$ then $b_{j,u}$ is supported in $B(x'_j, k(u)r_j)$. So the condition (i) holds for $b_{j,u}$.

Next, by (3) we have

$$\nu(B(x'_j, k(u)r_j)) \leq C_{\nu} k(u)^s \nu(B(x_j, r_j)).$$

This implies that

$$\|a_j \circ A(u)\|_{\infty} = \|a_j\|_{\infty} \leq \frac{1}{\nu(B(x_j, r_j))} \leq C_{\nu} k(u)^s \frac{1}{\nu(B(x'_j, k(u)r_j))}.$$

So, the condition (ii) is also fulfilled for $b_{j,u}$.

The validity of (iii) follows from [10, VII.1.4, formula (31)].

Finally, since formula (6) can be rewritten as

$$g \circ A(u) = \sum_{j=1}^{\infty} (C_{\nu} k(u)^s \alpha_j) b_{j,u},$$

we get

$$\|g \circ A(u)\|_{H^1(G)} \leq C_\nu k(u)^s \sum_{j=1}^{\infty} |\alpha_j|,$$

which in turn implies that

$$\|g \circ A(u)\|_{H^1(G)} \leq C_\nu k(u)^s \|g\|_{H^1(G)} = C_\nu k(u)^s \|f\|_{H^1(G/K)}.$$

Thus, the statement of the theorem follows from formula (5).

Corollary 3. *Let the assumptions of Theorem 2 holds. Then we have for a discrete Hausdorff operator over G/K (see Corollary 1)*

$$\|\mathcal{H}_{\Phi, A}\|_{\mathcal{L}(H^1(G/K))} \leq C_\nu \kappa_\rho^s \sum_{j=0}^{\infty} \frac{|\Phi(j)|}{(\text{mod} A(j))^s}.$$

Corollary 4. *Let the assumptions of Theorem 2 holds. Then we have for a Cesaro operator over G/K (see Corollary 2)*

$$\|\mathcal{C}_{A, \mu}\|_{\mathcal{L}(H^1(G/K))} \leq C_\nu \kappa_\rho^s \int_{\{\text{mod} A(u) \geq 1\}} \frac{d\mu(u)}{(\text{mod} A(u))^{1+s}}.$$

(Recall, that for this operator

$$\Phi = \frac{\chi_{\{\text{mod} A(u) \geq 1\}}}{\text{mod} A}.)$$

Concluding remarks

It would be of interest to apply Theorems 1 and 2 to classical homogeneous spaces such as Euclidean plane $\mathbb{R}^2 = M(2)/O(2)$, sphere $S^2 = O(3)/O(2)$, non-Euclidean plane $\mathbb{H} = SU(1, 1)/SO(2)$ [17, §4], to other Riemannian symmetric spaces etc. Would also be of interest to generalize Theorems 1 and 2 to the case when the group K is noncompact.

This is a preprint of the paper [18].

References

- [1] Hardy GH. *Divergent Series*. Oxford : Clarendon Press; 1949. 396 p.
- [2] Georgakis C. The Hausdorff mean of a Fourier–Stieltjes transform. *Proc. Amer. Math. Soc.* 1992; 116: 465–471. DOI: <https://doi.org/10.1090/S0002-9939-1992-1096210-9>.

- [3] Liflyand E., Móricz F. The Hausdorff operator is bounded on the real Hardy space $H^1(\mathbb{R})$. *Proc. Amer. Math. Soc.* 2000; 128: 1391–1396. DOI: <https://doi.org/10.1090/S0002-9939-99-05159-X>.
- [4] Lerner A., Liflyand E. Multidimensional Hausdorff operators on the real Hardy space. *J. Austr. Math. Soc.* 2007; 83: 79–86. doi:10.1017/S1446788700036399.
- [5] Liflyand E. Hausdorff operators on Hardy spaces. *Eurasian Math. J.* 2013; 4(4): 101 – 141.
- [6] Chen J., Fan D., Wang S. Hausdorff operators on Euclidean space (a survey article). *Appl. Math. J. Chinese Univ. Ser. B.* 2013; 28(4): 548–564. DOI:<https://doi.org/10.1007/s11766-013-3228-1>.
- [7] Mirotin AR. Boundedness of Hausdorff operators on Hardy spaces H^1 over locally compact groups. *J. Math. Anal. Appl.* 2019; 473: 519 – 533. DOI: 10.1016/j.jmaa.2018.12.065.
- [8] Mirotin AR. Addendum to "Boundedness of Hausdorff operators on Hardy spaces H^1 over locally compact groups". *J. Math. Anal. Appl.* 2019; 479(1): 872 – 874. DOI:<https://doi.org/10.1016/j.jmaa.2018.12.065>.
- [9] Hewitt E. , Ross K. *Abstract Harmonic Analysis. Vol. 1.* Berlin-Gettingen-Heidelberg: Springer-Verlag; 1963. 519 p.
- [10] Bourbaki N. *Elements de mathematique. 2nd edn. Integration, Chaps. I-IV.* Paris: Hermann; 1965.
- [11] Mirotin AR. The structure of normal Hausdorff operators on Lebesgue spaces. *Forum Math.* 2020; 32: 111–119. DOI: <https://doi.org/10.1515/forum-2019-0097>.
- [12] Mirotin AR. On the Structure of Normal Hausdorff Operators on Lebesgue Spaces. *Functional Analysis and Its Applications.* 2019; 53: 261–269. DOI: <https://doi.org/10.4213/faa3645>
- [13] Coifman RR, Weiss G. Extensions of Hardy spaces and their use in analysis, *Bull. Amer. Math. Soc.* 1977; 83(4): 569 – 645. DOI: <https://doi.org/10.1090/S0002-9904-1977-14325-5>.
- [14] Heinonen J. , Koskela P. , Shanmugalingam N. , Tyson JT. *Sobolev Spaces on Metric Measure Spaces. An Approach Based on Upper*

- Gradients*. Cambridge: Cambridge University Press; 2015. 434 p. DOI:<https://doi.org/10.1017/CBO9781316135914>.
- [15] Mirotin AR. On the general form of linear functionals on the Hardy spaces H^1 over compact Abelian groups and some of its applications. *Indag. Math.* 2017; 28: 451 – 462. DOI:<https://doi.org/10.1016/j.indag.2016.11.023>.
 - [16] Kawazoe T. Atomic Hardy spaces on semisimple Lie groups. *Japan. J. Math.* 1985; 11 (2): 293 - 343. DOI: <https://doi.org/10.4099/math1924.11.293>.
 - [17] Helgason S. *Topics in harmonic analysis on homogeneous spaces*. Boston - Basel - Stuttgart: Birkhauser; 1981. 142 p. DOI: <https://doi.org/10.1002/zamm.19820620921>.
 - [18] Mirotin AR. Hausdorff operators on homogeneous spaces of locally compact groups. *Journal of Belorussian State University. Mathematics. Informatics*. 2020; No 2, 28 – 35.