

Detecting Abrupt Changes in High-Dimensional Self-Exciting Poisson Processes

Daren Wang¹, Yi Yu², and Rebecca Willett¹

¹Department of Statistics, University of Chicago

²Department of Statistics, University of Warwick

Abstract

High-dimensional self-exciting point processes have been widely used in many application areas to model discrete event data in which past and current events affect the likelihood of future events. In this paper, we are concerned with detecting abrupt changes of the coefficient matrices in discrete-time high-dimensional self-exciting Poisson processes, which have yet to be studied in the existing literature due to both theoretical and computational challenges rooted in the non-stationary and high-dimensional nature of the underlying process. We propose a penalized dynamic programming approach which is supported by a theoretical rate analysis and numerical evidence.

Keywords: Self-exciting Poisson process; High-dimensional statistics; Piecewise stationarity; Penalized dynamic programming.

1 Introduction

Self-exciting point processes (SEPPs) are useful in modelling many types of discrete event data in which past and current event help determine the likelihood of future events. Such data are ubiquitous in application areas including crime science (e.g. [Egesdal et al., 2010](#)), national security (e.g. [Lewis et al., 2012](#)), finance (e.g. [Chavez-Demoulin and McGill, 2012](#)) and neuroscience (e.g. [Linderman et al., 2016](#)), to name but a few.

SEPPs were, arguably, first rigorously studied in a mathematical framework by [Hawkes \(1971\)](#), where the eponymous Hawkes process was proposed. Since the debut of the Hawkes process, there have been tremendous efforts poured into different aspects of understanding and utilizing the univariate Hawkes process; see [Laub et al. \(2015\)](#) and [Reinhart \(2019\)](#) for comprehensive and contemporary reviews. More recently, due to the availability of richer datasets and computational resources, attention has shifted to multivariate and even high-dimensional SEPPs, where different coordinates might correspond to different geographic locations, different neurons in a biological neural network, people in a social network, etc. See, for instance, [Hall et al. \(2016\)](#), [Mark et al. \(2018\)](#), [Chavez-Demoulin and McGill \(2012\)](#) and [Ertekin et al. \(2015\)](#).

In these high-dimensional settings, understanding how events in one coordinate influence the likelihood of events in another coordinate provides valuable insight into the underlying process. We call the collection of these influences between pairs of coordinates a “network”, and this paper describes novel methods for detecting abrupt changes in this network with theoretical performance

bounds that characterize the accuracy of the change point estimation and how strong the signals must be to ensure reliable estimation.

While change point detection has a long and rich history, we are unaware of any preexisting change point methodology that can be used to detect changes in SEPPs in high dimensions. Some recent high-dimensional change point detection work is briefly discussed as follows. Wang et al. (2018) and Padilla et al. (2019) studied the change point detection in Bernoulli networks and dynamic random dot product graphs, respectively. Cho and Fryzlewicz (2015), Cho (2016), Matteson and James (2014), Wang and Samworth (2018), Dette and Gösmann (2018) and others investigated high-dimensional mean change problems. Wang et al. (2017), Aue et al. (2009) and others were concerned with high/multi-dimensional covariance structure changes. Safikhani and Shojaie (2017) and Wang et al. (2019) exploited the high-dimensional vector autoregressive models and provided change point detection results thereof. Li et al. (2017) focused on a low-dimensional Hawkes process setting in which the processes may be characterized by a small number of parameters.

The lack of results on the change point analysis on high-dimensional SEPPs can be related to its nonlinearities inherent to the model. Note that, SEPPs can be viewed as a nonlinear autoregressive process, and even detecting changes for linear vector autoregressive processes is an active area of investigation (Wang et al., 2019). The nonlinearities associated with SEPPs further complicate the change point detection problem.

This paper describes a computationally- and statistically-efficient methodology for detecting changes in the network underlying SEPPs. At the heart of our method lies a penalized dynamic programming algorithm that estimates the times at which each change occurs when the underlying network is sparse, i.e. when the number of network edges is small relative to the number of pairs of network nodes. In this paper, we also apply our method to neuron spike train data sets to help pinpoint the times at which the functional networks might change due to the changes of the state of consciousness.

1.1 Problem formulation

The detailed model considered in this paper is introduced as follows.

Model 1. Let $\{X(t)\}_{t=1}^T \subset \mathbb{Z}^M$ be a discrete-time Poisson process. For each $t \in \{1, \dots, T\}$, let $\mathcal{X}(t) = (X(1), \dots, X(t)) \in \mathbb{R}^{M \times t}$ consist all the history up to time t . For each $t \in \{1, \dots, T-1\}$ and $m \in \{1, \dots, M\}$, suppose that given $\mathcal{X}(t)$, all coordinates $\{X_m(t+1)\}$ are conditionally independent and the conditional distribution of $X_m(t+1)$ is a Poisson distribution, i.e.

$$X_m(t+1)|\mathcal{X}(t) \sim \text{POISSON}(\exp\{\lambda_m(t)\}), \quad (1)$$

where

$$\lambda_m(t) = v + A_m^*(t)g_t\{\mathcal{X}(t)\}, \quad (2)$$

the matrix $A^*(t) \in \mathbb{R}^{M \times M}$ is the coefficient matrix at time point t , $A_m^*(t)$ is the m -th row of $A^*(t)$ and $g_t(\cdot) : \mathbb{R}^{M \times t} \rightarrow \mathbb{R}^M$ is a M -dimensional vector-valued function.

Suppose that there exists an integer $K \geq 0$ and time points $\{\eta_k\}_{k=0}^{K+1}$, called change points, satisfying $1 = \eta_0 < \eta_1 < \dots < \eta_K \leq T < \eta_{K+1} = T+1$ and

$$A^*(t) \neq A^*(t-1) \quad \text{if and only if} \quad t \in \{\eta_k\}_{k=1}^K.$$

Let the minimal spacing and the minimal jump size be defined as

$$\Delta = \min_{k=1, \dots, K+1} (\eta_k - \eta_{k-1}) \quad \text{and} \quad \kappa = \min_{k=1, \dots, K+1} \|A^*(\eta_k) - A^*(\eta_{k-1})\|_F,$$

respectively, where $\|\cdot\|_F$ denotes the Frobenius norm of a matrix.

It is worth mentioning that $\{X(t)\}_{t=1}^T$ defined in Model 1 is an SEPP, where each $X_m(t)$ is conditionally distributed as a Poisson random variable. We therefore refer to (1) a self-exciting Poisson processes. In the sequel, when there is no ambiguity, we will also refer self-exciting Poisson processes as SEPPs.

In fact, Model 1 is a generalization of a stationary SEPP process, which assumes that the coefficient matrices $A^*(t) = A^*(1)$, $t \in \{1, \dots, T\}$. Stationary SEPP models have been well-studied in the existing literature, including Hall et al. (2018) and Mark et al. (2018), where it has been shown that the coefficient matrix of the point process can be estimated by an ℓ_1 -penalized likelihood estimator.

Given $\{X(t)\}_{t=1}^T$ satisfying Model 1, our main task is to estimate $\{\eta_k\}_{k=1}^K$ accurately. To be specific, we seek estimators $\{\hat{\eta}_k\}_{k=1}^{\hat{K}}$ such that as the sample size $T \rightarrow \infty$, with probability tending to 1, it holds that

$$\hat{K} = K \quad \text{and} \quad \frac{\epsilon}{\Delta} = \frac{\max_{k=1, \dots, K} |\hat{\eta}_k - \eta_k|}{\Delta} \rightarrow 0. \quad (3)$$

For the change point estimators satisfying (3), we call them *consistent* change point estimators. We will also call ϵ the *localization error*.

To the best of our knowledge, we are the first to study the high-dimensional SEPPs with change points. In addition to the mathematical introduction of the model, we investigate the consistency of the abrupt change point location estimators, under minimal conditions. The proposed penalized dynamic programming approach in Section 2 is computationally efficient and tailored for this novel setting.

Notation

For any integer pair $(t_1, t_2) \in \mathbb{Z}^2$, let $[t_1, t_2]$ denote the integer interval $[t_1, t_2] \cap \mathbb{Z}$. Same notation applies to open intervals. For any matrix $A \in \mathbb{R}^{M \times M}$, let A_m denote the m th row of A and $A_{m, m'}$ denote the (m, m') th entry of A . With some abuse of notation, for any vector v and any matrix M , let $\|v\|_2$, $\|v\|_1$, $\|M\|_F$ and $\|M\|_1$ be the ℓ_2 - and ℓ_1 -norms of v , the Frobenius norm of M and the ℓ_1 -norm of $\text{vec}(M)$, respectively, where $\text{vec}(M)$ is the vectorized version of M by stacking all the columns of M . For any $v(t) : [1, T] \rightarrow \mathbb{R}^m$, let

$$\|Dv\|_0 = \sum_{t=2}^T I\{v(t-1) \neq v(t)\},$$

where $I\{\cdot\} \in \{0, 1\}$ is the indicator function. For any set $S \subset \{(m, m') : m, m' = 1, \dots, M\}$, let $A_S \in \mathbb{R}^{M \times M}$ satisfy

$$(A_S)_{m, m'} = \begin{cases} A_{m, m'}, & (m, m') \in S, \\ 0, & \text{otherwise.} \end{cases}$$

Given any $A(t) : [1, T] \rightarrow \mathbb{R}^{M \times M}$ and any $J \subset [1, T]$, if $A(\cdot)$ is unchanged in I , then we denote $A(I) = A(t)$, $t \in I$.

2 The Penalized Dynamic Programming Algorithm

To detect the change points in Model 1, we propose the penalized dynamic programming (PDP) algorithm, which is stated in (7) with necessary notation in (4), (5) and (6). The PDP consists of two layers: estimation of the coefficient matrices $A^*(t)$, $t \in [1, T]$, and estimation of the change points.

For the coefficient matrix estimation, we let $\widehat{A}(I)$ be the penalized log-likelihood estimator of the coefficient matrix over an integer interval $I \subset [1, T]$, i.e.

$$\widehat{A}(I) = \underset{A \in \mathcal{C}}{\operatorname{argmin}} H(A, I), \quad (4)$$

where $H(A, I)$ and \mathcal{C} are the penalized log-likelihood function and the constrained domain of the coefficient matrices, respectively. To be specific, with a pre-specified tuning parameter $\lambda > 0$ and $I = [s, e]$, let

$$H(A, I) = \sum_{t=s}^{e-1} \sum_{m=1}^M \left(\exp[v + A_m g_t\{\mathcal{X}(t)\}] - X_m(t+1)[v + A_m g_t\{\mathcal{X}(t)\}] + \lambda |I|^{1/2} \|A\|_1 \right) \quad (5)$$

and

$$\mathcal{C} = \left\{ A \in \mathbb{R}^{M \times M} : \max_{m=1, \dots, M} \|A_m\|_1 \leq 1 \right\}. \quad (6)$$

The loss function $H(\cdot, \cdot)$ is a penalized logarithmic conditional likelihood function, recalling that $X_m(t+1)$ given $\mathcal{X}(t)$ follows a Poisson distribution with intensity $\exp[v + A_m g_t\{\mathcal{X}(t)\}]$. The penalty term $\lambda |I|^{1/2}$ in (5) is introduced in a way such that the tuning parameter λ is independent of the interval length. The term $|I|^{1/2}$ reflects the order of the standard error of the sum of $|I|$ marginal log-likelihood functions. We elaborate on this scaling factor and its derivation in Lemma 8 and its proof.

The constraint on \mathcal{C} is to ensure that the SEPP process as vector-valued time series is stable (see e.g. Lütkepohl, 2005). As for stationary SEPP estimation, Mark et al. (2018) proposed a constraint similar to (6).

Given the above framework, we can now consider estimating change points by setting

$$\widehat{\mathcal{P}} = \underset{\mathcal{P}}{\operatorname{argmin}} \left\{ \sum_{I \in \mathcal{P}} H(\widehat{A}(I), I) + \gamma |\mathcal{P}| \right\}, \quad (7)$$

where $\gamma > 0$ is a tuning parameter, the minimization is over all possible interval partitions of $[1, T]$ and \mathcal{P} denotes one such partition. To be specific, an interval partition has the form $\mathcal{P} = \{I_k, k = 1, \dots, K_{\mathcal{P}}\}$ and satisfies $I_{k'} \cap I_k = \emptyset$ and $\bigcup_{k=1}^{K_{\mathcal{P}}} I_k = [1, T]$. Once $\widehat{\mathcal{P}}$ is at hand, we let $\widehat{K} = |\widehat{\mathcal{P}}| - 1 \geq 0$, $\eta_{\widehat{K}+1} = T + 1$ and

$$\widehat{\mathcal{P}} = \left\{ \{1, \dots, \widehat{\eta}_1 - 1\}, \{\widehat{\eta}_k, \dots, \widehat{\eta}_{k+1} - 1\}_{k=1}^{\widehat{K}} \right\}.$$

We call $\{\widehat{\eta}_k\}_{k=1}^{\widehat{K}}$ the change point estimators induced by $\widehat{\mathcal{P}}$.

The optimization problem in (7) is known as the minimal partition problem on a linear chain graph and can be solved using dynamic programming (e.g. Friedrich et al., 2008) with the worst case computational cost of order $O\{T^2 \text{Cost}(T)\}$, where $\text{Cost}(t)$ denotes, in our case, the computational cost of computing $\hat{A}(I)$ in the interval I with $|I| = t$. We remark that there has been a line of attack on the computational aspect of optimizing the minimal partition problem, including Killick et al. (2012) and Maidstone et al. (2017), among others.

For completeness, we summarize the PDP procedure in Algorithm 1 below. The quantities and functions involved there are defined in (4), (5) and (6).

Algorithm 1 Penalized dynamic programming. $\text{PDP}(\{X(t)\}_{t=1}^n, \lambda, \gamma)$

INPUT: Data $\{X(t)\}_{t=1}^T$, tuning parameters $\lambda, \gamma > 0$.

$(\mathcal{B}, s, t, \text{FLAG}) \leftarrow (\emptyset, 0, 2, 0)$

while $s < T - 1$ **do**

$s \leftarrow s + 1$

while $t < n$ and $\text{FLAG} = 0$ **do**

$t \leftarrow t + 1$

if $\min_{l=s+1, \dots, t-1} \{H(\hat{A}([s, l]), [s, l]) + H(\hat{A}([l+1, t]), [l+1, t]) + \gamma < H(\hat{A}([s, t]), [s, t])\}$ **then**

$$s \leftarrow \min_{l=s+1, \dots, t-1} \operatorname{argmin} \{H(\hat{A}([s, l]), [s, l]) + H(\hat{A}([l+1, t]), [l+1, t]) + \gamma < H(\hat{A}([s, t]), [s, t])\}$$

$$[l+1, t] + \gamma < H(\hat{A}([s, t]), [s, t])\}$$

$\mathcal{B} \leftarrow \mathcal{B} \cup \{s\}$

$\text{FLAG} \leftarrow 1$

end if

end while

end while

OUTPUT: The set of estimated change points \mathcal{B} .

2.1 The localization rate of the Penalized Dynamic Programming estimators

In order to establish the consistency of the change point estimators resulting from the PDP procedure detailed in Algorithm 1, we first impose Assumption 1.

Assumption 1. Let $\{X(t)\}_{t=1}^T \subset \mathbb{Z}^M$ be a discrete-time SEPP generated according to Model 1 and satisfying the following.

A1. There exists a subset $S \subset \{(m, m') : m, m' = 1, \dots, M\}$ such that for all $t \in [1, T]$, $A_{m, m'}^*(t) = 0$, if $(m, m') \notin S$. Let $d = |S|$.

A2. It holds that

$$\max_{t=1, \dots, T} \max_{m=1, \dots, M} \|A_m^*(t)\|_1 \leq 1.$$

A3. For any $\xi > 0$, there exist absolute constants $C_{\Delta, 1}, C_{\Delta, 2} > 0$ such that

$$\Delta \geq C_{\Delta, 1} T \quad \text{and} \quad \Delta \geq C_{\Delta, 2} \log^{2+\xi}(TM) d^2 \max\{\kappa^{-2}, \kappa^{-4}\}.$$

A4. *There exist absolute constants $p \in \mathbb{Z}^+$ and $\omega > 0$ such that for any t , the matrix*

$$\mathbb{E}[g_t\{\mathcal{X}(t)\}g_t\{\mathcal{X}(t)\}^\top | \mathcal{X}(t-p)] - \omega I_M$$

is positive definite, where $I_M \in \mathbb{R}^{M \times M}$ is an identity matrix. In addition, v and $\|g_t(\cdot)\|_\infty$, for all t , are uniformly upper bounded by an absolute constant $C_g > 0$.

Model 1 and Assumption 1 completely characterize the problem with model parameters M (the dimensionality of the time series), d (the sparsity parameter indicating an upper bound of the number of nonzero entries in all the coefficient matrices), Δ (the minimal spacing between change points), and κ (the minimal jump size), along with the sample size T . The consistency we are to establish is based on allowing M and d to diverge and κ to vanish as the sample size T diverges unbounded.

The number of parameters at each time point is of order M^2 , which is allowed to well exceed the sample size. A sparsity constraint therefore comes into force in Assumption **A1**, which is a standard assumption in the high-dimensional statistics literature. Note that the set S is the union of all (m, m') pairs with a nonzero entry in any coefficient matrix. Assumption **A2** echoes the imposition of the constraint domain \mathcal{C} (6) in the optimization (4), to ensure the stationarity of the SEPP. In fact, the constant one in the upper bound can be relaxed to any absolute constant and is set to be one in this paper for identification issue. To be specific, what goes into the model is the product of $A_m(t)$ and $g_t\{\mathcal{X}(t)\}$, and the latter is assumed to be upper bounded in sup-norm in Assumption **A4**.

Assumption **A3** can be regarded as a signal-to-noise assumption. It is required that the minimal spacing Δ is at least of a constant fraction of the total sample size, which implies the number of change points is of order $O(1)$. This might appear to be strong compared to other change point detection literature, however, the problem we are facing here is challenging due to the nonlinearity of the SEPP model. In fact, Assumption **A3** is a mild condition and covers some challenging scenarios. For instance, Assumption **A3** holds if $M \asymp \exp(T^{1/2})$, $d \asymp T^{1/4}$ and $\kappa \asymp \log(T)$. The quantity ξ can be set arbitrarily small and it ensures the consistency of the estimator which will be explained after Theorem 1.

Assumption **A4** can be interpreted as the restricted eigenvalue condition for SEPP processes. We refer readers to Section 4 of Mark et al. (2018) for a number of common self-excited point process models satisfying Assumption **A4**.

In what follows, we show the consistency of PDP in Theorem 1.

Theorem 1. *Let $\{X(t)\}_{t=1}^T \subset \mathbb{Z}^M$ be an SEPPs generated from Model 1 and satisfying Assumption 1. Let $\{\hat{\eta}_k\}_{k=1}^{\hat{K}}$ be the change point estimators from the PDP algorithm detailed in Algorithm 1 with tuning parameters*

$$\lambda = C_\lambda \log(TM) \quad \text{and} \quad \gamma = C_\gamma \log^2(TM) d (1 + d\kappa^{-2}), \quad (8)$$

where $C_\lambda, C_\gamma > 0$ are absolute constants, depending only on $p, \omega, C_{\Delta,1}, C_{\Delta,2}$ and C_g . We have that

$$\mathbb{P} \left\{ \hat{K} = K \quad \text{and} \quad \max_{k=1, \dots, K} |\hat{\eta}_k - \eta_k| \leq C_\epsilon d^2 \log^2(TM) \max\{\kappa^{-2}, \kappa^{-4}\} \right\} \geq 1 - 2(TM)^{-1},$$

where $C_\epsilon > 0$ is an absolute constant only depending on $p, \omega, C_{\Delta,1}, C_{\Delta,2}$ and C_g .

The proof of Theorem 1 is deferred to Appendix A, where it can be seen that the order of the estimation error is of the form

$$\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}.$$

Due to the signal-to-noise ratio condition in Assumption A3, we have that

$$\frac{\max_{k=1,\dots,K} |\hat{\eta}_k - \eta_k|}{\Delta} \lesssim \frac{d^2 \log^2(TM) \max\{\kappa^{-2}, \kappa^{-4}\}}{\Delta} \lesssim \frac{d^2 \log^2(TM) \max\{\kappa^{-2}, \kappa^{-4}\}}{d^2 \log^{2+\xi}(TM) \max\{\kappa^{-2}, \kappa^{-4}\}} \rightarrow 0,$$

as $T \rightarrow \infty$. This explains the role of the quantity ξ in Assumption A3 and shows the consistency of the PDP algorithm. In fact, if we let $d = 1$ and assume $\kappa > 1$, then the localization error we derived here coincides with the optimal localization error in the univariate mean change point detection problem (e.g. Wang et al., 2020).

Two tuning parameters are involved, where λ is used in the optimization (5) to recover the sparsity in estimating high-dimensional coefficient matrices, and γ is involved in optimizing (7) to penalize the over-partitioning. The order of λ required in (8) is a logarithmic quantity in T and M , which is resulted from a union bound argument applied to a sub-exponential concentration bound. The requirement on γ is essentially that $\gamma \asymp \lambda^2 (d + d^2 \kappa^{-2})$, which can be intuitively explained as an upper bound on the difference between $H(\hat{A}(I_1), I_1) + H(\hat{A}(I_2), I_2)$ and $H(\hat{A}(I_1 \cup I_2), I_1 \cup I_2)$, where I_1 and I_2 are two relatively long, non-overlapping and adjacent intervals, and there is no true change point near the shared endpoint of I_1 and I_2 . In this case, one would not wish to partition $I_1 \cup I_2$ into I_1 and I_2 . If we only focus on the log-likelihood functions, over-estimating will result in that

$$H(\hat{A}(I_1), I_1) + H(\hat{A}(I_2), I_2) < H(\hat{A}(I_1 \cup I_2), I_1 \cup I_2).$$

The penalty we impose through γ will therefore avoid this over-partitioning.

2.2 Comparisons with related work

In a broad sense, as we have mentioned, there have been numerous existing papers on different aspects of SEPPs. In fact, another related area is the analysis of piecewise-stationary time series models, where we also see a vast volume of existing papers. The two most related papers are Mark et al. (2018), which is concerned with a stationary, high-dimensional SEPP process, and Wang et al. (2019), which studies a piecewise-stationary high-dimensional linear process.

Mark et al. (2018) studied a stationary version of Model 1 with $K = 0$. The penalized estimator of the coefficient matrix developed there is almost identical to the ones summoned in our problem in (4). There are a few fundamental differences between this paper and Mark et al. (2018). (1) Due to the piecewise-stationarity assumed in Model 1, when estimating the coefficient matrices in (4) and (5), it is possible that there exists a true change point in the interval of interest and the estimator we seek is an estimator of a mixture of different true coefficient matrices. (2) We provide a more refined analysis as an improved version of Mark et al. (2018), for instance, the optimization constrain domain \mathcal{C} defined in (6) is a cleaner version of its counterpart in Mark et al. (2018); a subspace compatibility condition is required in Mark et al. (2018) to control the ratio of different norms of the coefficient matrix, and this assumption is shown to be redundant in our new analysis.

The other closest-related work is Wang et al. (2019), where the change point localizing problem in the piecewise-stationary vector autoregressive models is investigated and a penalized dynamic programming approach was deployed there. The main differences between this paper and Wang

et al. (2019) comes from the underlying model. The vector autoregressive model is a linear model in the sense that given $\mathcal{X}(t)$ the history data up till time point t , the conditional expectation of $X(t+1)$ is a linear combination of the columns of $\mathcal{X}(t)$, which is not the case here. The self-exciting point process is a nonlinear model, and as we have mentioned, the logarithm of the conditional intensity is a linear function of the history. Another key difference is that Wang et al. (2019) are concerned with sub-Gaussian innovation sequences, while the counting processes we study here determine the heavy-tail properties of the data.

3 Numerical Experiments

In this section, we further examine the performances of the PDP algorithm by numerical experiments, with simulated data analyzed in Section 3.1 and a real data set in Section 3.2.

3.1 Simulated data analysis

We generate data according to Model 1 and Assumption 1. In particular, we adopt the setting in Mark et al. (2018) and assume that the design function $g_t(\cdot)$ is defined to be

$$g_t\{\mathcal{X}(t)\} = (\min\{\mathcal{X}_1(t), C_g\}, \dots, \min\{\mathcal{X}_M(t), C_g\})^\top \in \mathbb{R}^M, \quad (9)$$

where $C_g > 0$ is a constant, $\mathcal{X}(t)$ is an $M \times t$ matrix and $\mathcal{X}_m(t)$ denotes the m th row of $\mathcal{X}(t)$, $m \in \{1, \dots, M\}$. For the two tuning parameters λ and γ defined in (5) and (7), respectively, with the theoretical guidance in Theorem 1, we fix $\lambda = 90 \log(TM)$ and $\gamma = \log^2(M)/2$ in all experiments in this section.

Since the piecewise-stationary SEPP model is first introduced here, we do not have direct competitors. For illustration purpose, however, we compare our PDP algorithm with the SBS-MVTS algorithm (Cho and Fryzlewicz, 2015) and E-Divisive procedure (Matteson and James, 2014), both of which are designed to detect abrupt change points in multivariate time series, but neither of which is designed specifically for the scenarios we are studying here. Having said this, there are the reasons we choose these two competitors. The SBS-MVTS can identify covariance changes in the high-dimensional autoregressive time series and the E-Divisive procedure can estimate of both the number and locations of change points under mild assumptions on the first or second moments of the underlying distributions. Since Poisson random variables have the same means and variances, these two competitors may be able to detect the changes in Poisson processes with piecewise-constant parameters. In all the simulated experiments, the tuning parameters for SBS-MVTS algorithm and E-Divisive procedure are selected according to the information-type criteria and permutation tests in the R (R Core Team, 2017) packages wbs (Baranowski and Fryzlewicz, 2019) and ecp (Nicholas A. James and Matteson, 2019), respectively.

Let $\{\hat{\eta}_k\}_{k=1}^{\hat{K}}$ and $\{\eta_k\}_{k=1}^K$ be a collection of change point estimates and a collection of true change points, respectively. We evaluate the estimators' performances by the absolute error $|K - \hat{K}|$ and their Hausdorff distance. The Hausdorff distance between two sets \mathcal{A} and \mathcal{B} is defined as

$$\mathcal{D}(\mathcal{A}, \mathcal{B}) = \max\{d(\mathcal{A}|\mathcal{B}), d(\mathcal{B}|\mathcal{A})\}, \quad (10)$$

where

$$d(\mathcal{A}|\mathcal{B}) = \max_{a \in \mathcal{A}} \min_{b \in \mathcal{B}} |a - b|.$$

In the sequel, we consider three settings. Recall that T is the total number of time points, M is the dimensionality of the time series and C_g is the threshold used in the design function $g_t(\cdot)$, which is specified in (9). Every setting is repeated 100 times. Additional setting details are listed below.

- (a) One change point and varying jump size. Fix $T = 300$, $M = 30$, $C_g = 6$ and the intercept $v = 1/2$, which is defined in (2). Let

$$A^*(t) = \begin{cases} (\rho v_1, \rho v_2, 0_{M \times (M-2)}) \in \mathbb{R}^{M \times M}, & t \in [1, 150], \\ (\rho v_2, \rho v_1, 0_{M \times (M-2)}) \in \mathbb{R}^{M \times M}, & t \in [151, 300], \end{cases}$$

where $v_1 \in \mathbb{R}^M$ with odd coordinates being 1 and even coordinates being -1 , $v_2 = -v_1$, $0_{M \times (M-2)} \in \mathbb{R}^{M \times (M-2)}$ is an all zero matrix and $\rho \in \{0.15, 0.20, 0.25, 0.30, 0.35\}$.

- (b) Two change points and varying minimal spacing. Let $T \in \{180, 240, 300, 360, 420\}$, $M = 40$, $C_g = 8$ and the intercept $v = 1/4$. Let the coefficient matrices satisfy $(A^*(t))_{ij} = 0$, $|i-j| > 1$, $t \in [1, T]$,

$$(A^*(t))_{ij} = \begin{cases} \begin{cases} 0.15 & t \in [1, T/3] \cup (2T/3, T], \\ -0.15 & t \in (T/3, 2T/3], \end{cases} & i = j, \\ \begin{cases} -0.15 & t \in [1, T/3], \\ 0.15 & t \in (T/3, T], \end{cases} & i - j = -1, \\ \begin{cases} 0.15 & t \in [1, 2T/3], \\ -0.15 & t \in (2T/3, T], \end{cases} & i - j = 1. \end{cases}$$

- (c) Two change points and varying dimension. Let $T = 450$, $C_g = 4$, $v = 1/5$ and $M \in \{15, 20, 25, 30, 35\}$. Let

$$A(t) = \begin{cases} (v_1, v_2, v_3, 0_{M \times (M-3)}), & t \in [1, 150], \\ (v_2, v_3, v_3, 0_{M \times (M-3)}), & t \in [151, 300], \\ (v_3, v_2, v_1, 0_{M \times (M-3)}), & t \in [301, 450], \end{cases}$$

where $v_1, v_2, v_3 \in \mathbb{R}^M$ are

$$\begin{aligned} v_1 &= (-0.075, 0.15, 0.3, -0.3, 0, \dots, 0)^\top, \\ v_2 &= (\underbrace{0, \dots, 0}_4, 0.375, -0.225, -0.075, 1.5, 0.225, 0, \dots, 0)^\top, \\ v_3 &= (\underbrace{0, \dots, 0}_8, -0.15, -0.075, 0.45, -0.225, 0, \dots, 0)^\top. \end{aligned}$$

We collect the simulation results in Table 1, each cell containing the mean and standard errors of 100 repetitions. The Hausdorff distances are visualized in Figure 1 to improve readability. These three settings have ranged over various situations. It is clearly that PDP outperforms both competitors in all settings on both metrics.

		Setting (a)				
Method	Metric	$\rho = 0.15$	$\rho = 0.20$	$\rho = 0.25$	$\rho = 0.30$	$\rho = 0.35$
PDP	\mathcal{D}	3.1(9.8)	1.1(1.0)	0.7(0.5)	0.6(0.5)	0.6(0.5)
SBS-MVTS		282.6(69.1)	226.5(119.9)	114.7(130.8)	47.3(52.9)	9.3(21.3)
E-Divisive		151.0(0.0)	151.0(0.0)	151.0(0.0)	151.0(0.0)	151.0(0.0)
PDP	$ \widehat{K} - K $	0.0(0.0)	0.0(0.0)	0.0(0.0)	0.0(0.0)	0.0(0.0)
SBS-MVTS		0.9(0.2)	0.7(0.4)	0.4(0.5)	0.5(0.5)	0.1(0.3)
E-Divisive		300.0(0.0)	300.0(0.1)	300.0(0.5)	296.4(16.2)	287.2(31.4)
		Setting (b)				
		$T = 180$	$T = 240$	$T = 300$	$T = 360$	$T = 420$
PDP	\mathcal{D}	11.5(6.2)	3.7(4.6)	2.5(4.6)	2.8(4.3)	1.2(3.6)
SBS-MVTS		177.0(21.1)	233.3(38.1)	270.1(85.5)	243.8 (156.1)	263.5(185.2)
E-Divisive		61.0(0.0)	81.0(0.0)	101.0(0.0)	121.0(0.0)	141.0(0.0)
PDP	$ \widehat{K} - K $	0.0(0.0)	0.0(0.0)	0.0(0.0)	0.0(0.0)	0.0(0.0)
SBS-MVTS		2.0(0.2)	1.9(0.3)	1.9(0.4)	1.6(0.7)	1.6(0.6)
E-Divisive		178.9(0.3)	238.9(0.3)	298.9(0.3)	358.8(0.4)	418.8(0.4)
		Setting (c)				
		$M = 15$	$M = 20$	$M = 25$	$M = 30$	$M = 35$
PDP	\mathcal{D}	3.3(5.0)	3.6(5.5)	3.2(4.5)	5.0(12.4)	6.1(13.2)
SBS-MVTS		401.4(112.8)	378.2(129.9)	411.3(101.7)	377.7(134.1)	375.4(134.5)
E-Divisive		151.0(0.0)	151.0(0.0)	151.0(0.0)	151.0(0.0)	151.0(0.0)
PDP	$ \widehat{K} - K $	0.0(0.0)	0.0(0.0)	0.0(0.0)	0.0(0.0)	0.0(0.0)
SBS-MVTS		1.8(0.4)	1.7(0.5)	1.9(0.3)	1.8(0.4)	1.8(0.4)
E-Divisive		448.6(0.5)	449.0(0.3)	449.0(0.1)	449.0(0.0)	449.0(0.0)

Table 1: Simulation results. Each cell is in the form of mean(standard error). For the metrics, \mathcal{D} denotes the Hausdorff distance defined in (10) and $|\widehat{K} - K|$ denotes the absolute errors in estimating the numbers of the change points. PDP uniformly outperforms the other two methods across a range of parameter values, including ρ (reflecting the jump size), T (the number of samples), and M (the dimension of the time series).

3.2 Real data example

We consider the neuron spike train data set previously analyzed in [Watson et al. \(2016b\)](#). The three chosen data sets are from [Watson et al. \(2016a\)](#) and each consists of wake-sleep episodes of multi-neuron spike train recording sessions of one laboratory animal. Each wake-sleep episode includes at least 7 minutes of wake time followed by at least 20 minutes of sleep time. Note that the wake and sleep periods were recorded so the true change point in each dataset is the end of the wake period. For each data set, we first compute the Firing Rate (FR) of each neuron using a 5-second discretization time window and then apply Algorithm 1 with $\lambda = 800$ and $\gamma = \log^2(M)/2$, the same as in Section 3.1. For comparison, we also apply the SBS-MVTS algorithm ([Cho and Fryzlewicz, 2015](#)) and E-Divisive procedure ([Matteson and James, 2014](#)).

These three subsets are on subjects 20140528_565um, BWRat17_121912 and BWRat19_032413. The numbers of neurons, i.e. the dimensions of the time series M , are 24, 33 and 41, respectively.




Figure 1: A visualization of the mean Hausdorff distance metric \mathcal{D} (10) in Table 1. The methods concerned are: Algorithm 1, PDP; SBS, SBS-MVTS; ECP, E-Divisive. In each panel, the y-axis represents the mean Hausdorff distance across 100 repetitions and the x-axis represents the varying parameter in each setting. PDP uniformly outperforms the other two methods across a range of parameter values, including ρ (reflecting the jump size), T (the number of samples), and M (the dimension of the time series).

The total numbers of 5-second time intervals, i.e. the total number of time points T considered in Model 1, are 3750, 2995 and 3920, respectively. The true change points are at point 788, 1184 and 2001, respectively.

The results are summarized in Table 2 and are depicted in Figure 2. As we can see from the table and the figure, our PDP algorithm consistently outperforms the other two algorithms in these real data examples.

Subject	Metric	PDP	SBS-MVTS	E-Divisive
20140528_565um	\mathcal{D}	38	382	2966
	$ \hat{K} - K $	0	0	740
BWRat17_121912	\mathcal{D}	84	140	1816
	$ \hat{K} - K $	0	0	595
BWRat19_032413	\mathcal{D}	1	99	1996
	$ \hat{K} - K $	0	0	773

Table 2: The results of three algorithms on multi-neuron spike train data sets. For the metrics, \mathcal{D} denotes the Hausdorff distance defined in (10) and $|\hat{K} - K|$ denotes the absolute errors in estimating the numbers of the change points. PDP uniformly outperforms the other two methods.




Figure 2: The true change points and the estimators provided by PDP and SBS-MVTS in the multi-neuron spike train data sets. Each panel corresponds to a subject. The y-axis represents the sum of the FRs across all neurons and the x-axis represents the ordered time intervals. The estimators of the E-Divisive procedure are not included because the corresponding \hat{K} 's are too large. PDP uniformly outperforms the other two methods. See Table 2 for detailed information.

4 Discussions

In this paper, we studied piecewise-stationary discrete-time high-dimensional self-exciting Poisson processes, which, or at least the theoretical properties of which were not studied in the literature. The number of stationary segments in the whole time series is assumed to be an unknown constant. All the other model parameters are allowed to be functions of the sample size T . We proposed a computationally-efficient and theoretically-guaranteed algorithm.

In the numerical experiments, we fix tuning parameters. One future research direction is to investigate data-driven methods for tuning parameter selection. Possible methods include variants of stationary bootstrap (Politis and Romano, 1994) or variants of information criteria (e.g. Chen and Chen, 2012).

Another future research direction is to extend the techniques we derived in this paper to other popular time series models. For instance, one key feature of the SEPPs we are concerned in this paper is the varying variance structure and heavy tail behaviours. These share similarities with the GARCH models, which are widely used in finance.

References

- Alexander Aue, Siegfried Hörmann, Lajos Horváth, and Matthew Reimherr. Break detection in the covariance structure of multivariate time series models. *The Annals of Statistics*, 37(6B):4046–4087, 2009.
- Kazuoki Azuma. Weighted sums of certain dependent random variables. *Tohoku Mathematical Journal, Second Series*, 19(3):357–367, 1967.
- Rafal Baranowski and Piotr Fryzlewicz. *wbs: Wild Binary Segmentation for Multiple Change-Point Detection*, 2019. URL <https://cran.r-project.org/web/packages/wbs/index.html>. R package version 1.4.
- Valérie Chavez-Demoulin and JA McGill. High-frequency financial data modeling using Hawkes processes. *Journal of Banking & Finance*, 36(12):3415–3426, 2012.
- Jiahua Chen and Zehua Chen. Extended bic for small-n-large-p sparse glm. *Statistica Sinica*, pages 555–574, 2012.
- Haeran Cho. Change-point detection in panel data via double cusum statistic. *Electronic Journal of Statistics*, 10(2):2000–2038, 2016.
- Haeran Cho and Piotr Fryzlewicz. Multiple-change-point detection for high dimensional time series via sparsified binary segmentation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 77(2):475–507, 2015.
- Holger Dette and Josua Gösmann. Relevant change points in high dimensional time series. *Electronic Journal of Statistics*, 12(2):2578–2636, 2018.
- Mike Egesdal, Chris Fathauer, Kym Louie, Jeremy Neuman, George Mohler, and Erik Lewis. Statistical and stochastic modeling of gang rivalries in Los Angeles. *SIAM Undergraduate Research Online*, 3:72–94, 2010.

- Şeyda Ertekin, Cynthia Rudin, Tyler H McCormick, et al. Reactive point processes: A new approach to predicting power failures in underground electrical systems. *The Annals of Applied Statistics*, 9(1):122–144, 2015.
- Felix Friedrich, Angela Kempe, Volkmar Liebscher, and Gerhard Winkler. Complexity penalized estimation: fast computation. *Journal of Computational and Graphical Statistics*, 17(1):201–224, 2008.
- Eric C Hall, Garvesh Raskutti, and Rebecca Willett. Inference of high-dimensional autoregressive generalized linear models. *arXiv preprint arXiv:1605.02693*, 2016.
- Eric C Hall, Garvesh Raskutti, and Rebecca M Willett. Learning high-dimensional generalized linear autoregressive models. *IEEE Transactions on Information Theory*, 65(4):2401–2422, 2018.
- Alan G Hawkes. Spectra of some self-exciting and mutually exciting point processes. *Biometrika*, 58(1):83–90, 1971.
- Rebecca Killick, Paul Fearnhead, and Idris A Eckley. Optimal detection of changepoints with a linear computational cost. *Journal of the American Statistical Association*, 107(500):1590–1598, 2012.
- Patrick J Laub, Thomas Taimre, and Philip K Pollett. Hawkes processes. *arXiv preprint arXiv:1507.02822*, 2015.
- Erik Lewis, George Mohler, P Jeffrey Brantingham, and Andrea L Bertozzi. Self-exciting point process models of civilian deaths in Iraq. *Security Journal*, 25(3):244–264, 2012.
- Shuang Li, Yao Xie, Mehrdad Farajtabar, Apurv Verma, and Le Song. Detecting changes in dynamic events over networks. *IEEE Transactions on Signal and Information Processing over Networks*, 3(2):346–359, 2017.
- Scott Linderman, Ryan P Adams, and Jonathan W Pillow. Bayesian latent structure discovery from multi-neuron recordings. In *Advances in neural information processing systems*, pages 2002–2010, 2016.
- Helmut Lütkepohl. *New introduction to multiple time series analysis*. Springer Science & Business Media, 2005.
- Robert Maidstone, Toby Hocking, Guillem Rigai, and Paul Fearnhead. On optimal multiple changepoint algorithms for large data. *Statistics and Computing*, 27(2):519–533, 2017.
- Benjamin Mark, Garvesh Raskutti, and Rebecca Willett. Network estimation from point process data. *IEEE Transactions on Information Theory*, 65(5):2953–2975, 2018.
- David S Matteson and Nicholas A James. A nonparametric approach for multiple change point analysis of multivariate data. *Journal of the American Statistical Association*, 109(505):334–345, 2014.
- Wenyu Zhang Nicholas A. James and David S. Matteson. *ecp: Non-Parametric Multiple Change-Point Analysis of Multivariate Data*, 2019. URL <https://cran.r-project.org/web/packages/ecp/index.html>. R package version 3.1.2.

- Oscar Hernan Madrid Padilla, Yi Yu, and Carey E Priebe. Change point localization in dependent dynamic nonparametric random dot product graphs. *arXiv preprint arXiv:1911.07494*, 2019.
- Dimitris N Politis and Joseph P Romano. The stationary bootstrap. *Journal of the American Statistical association*, 89(428):1303–1313, 1994.
- David Pollard. Empirical processes: theory and applications. In *NSF-CBMS regional conference series in probability and statistics*, pages i–86. JSTOR, 1990.
- R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2017. URL <https://www.R-project.org/>.
- Alex Reinhart. Self-exciting point processes, 2019. URL <https://www.refsmmat.com/notebooks/self-exciting-point-processes.html>.
- Abolfazl Safikhani and Ali Shojaie. Joint structural break detection and parameter estimation in high-dimensional non-stationary var models. *arXiv preprint arXiv:1711.07357*, 2017.
- Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.
- Daren Wang, Yi Yu, and Alessandro Rinaldo. Optimal covariance change point localization in high dimension. *arXiv preprint arXiv:1712.09912*, 2017.
- Daren Wang, Yi Yu, and Alessandro Rinaldo. Optimal change point detection and localization in sparse dynamic networks. *arXiv preprint arXiv:1809.09602*, 2018.
- Daren Wang, Yi Yu, Alessandro Rinaldo, and Rebecca Willett. Localizing changes in high-dimensional vector autoregressive processes. *arXiv preprint arXiv:1909.06359*, 2019.
- Daren Wang, Yi Yu, Alessandro Rinaldo, et al. Univariate mean change point detection: Penalization, cusum and optimality. *Electronic Journal of Statistics*, 14(1):1917–1961, 2020.
- Tengyao Wang and Richard J Samworth. High dimensional change point estimation via sparse projection. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(1): 57–83, 2018.
- Brendon O Watson, Daniel Levenstein, J Palmer Greene, Jennifer N Gelinias, and György Buzsáki. Multi-unit spiking activity recorded from rat frontal cortex (brain regions mpfc, ofc, acc, and m2) during wake-sleep episode wherein at least 7 minutes of wake are followed by 20 minutes of sleep, 2016a. URL <http://dx.doi.org/10.6080/K02N506Q>.
- Brendon O Watson, Daniel Levenstein, J Palmer Greene, Jennifer N Gelinias, and György Buzsáki. Network homeostasis and state dynamics of neocortical sleep. *Neuron*, 90(4):839–852, 2016b.

A Proof of Theorem 1

In all the appendices, we do not distinguish the notation of every single absolute constant. For notational simplicity, in the appendices, we drop the subscript of the function $g_t(\cdot)$.

Theorem 1 is an immediate consequence of Proposition 1 and 2.

Proposition 1. *Let $\widehat{\mathcal{P}}$ be the defined in (7). Under all the assumptions in Theorem 1, with probability at least $1 - (TM)^{-1}$, the following hold uniformly for any $I = (s, e) \in \widehat{\mathcal{P}}$.*

a. *If I contains only one change point η , then there exists an absolute constant $C > 0$ such that*

$$\min\{e - \eta, \eta - s\} \leq C \left(\frac{p\lambda^2 d}{\kappa^2} + \frac{p\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2} \right).$$

b. *If I contains exactly two change points η_k and η_{k+1} , then there exists an absolute constant $C > 0$ such that*

$$\max\{e - \eta_{k+1}, \eta_k - s\} \leq C \left(\frac{p\lambda^2 d}{\kappa^2} + \frac{p\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2} \right).$$

c. *If $|\widehat{\mathcal{P}}| > 1$, then let I and J be two consecutive intervals in $\widehat{\mathcal{P}}$. The interval $I \cup J$ contains at least one change point.*

d. *The interval I does not contain more than two change points.*

Proposition 2. *Let $\widehat{\mathcal{P}}$ be the defined in (7). Assume that $K \leq |\widehat{\mathcal{P}}| \leq 3K$. Under all the assumptions in Theorem 1, with probability at least $1 - (TM)^{-1}$, it holds that $|\widehat{\mathcal{P}}| = K$.*

B Proof of Proposition 1

Proof of Proposition 1. Proposition 1 is an immediate consequence of Lemmas 2, 3, 4 and 5. For illustration, we only prove the claim a.

Let $I_1 = (s, \eta]$ and $I_2 = (\eta, e]$. Note that

$$H(\widehat{A}(I), I) \leq H(\widehat{A}(I_1), I_1) + H(\widehat{A}(I_2), I_2) + \gamma \leq H(A^*(I_1), I_1) + H(A^*(I_2), I_2) + \gamma \quad (11)$$

where the first inequality follows from the fact that $\widehat{\mathcal{P}}$ is a minimizer defined in (7) and the second inequality follows from the definitions of $\widehat{A}(I_1)$ and $\widehat{A}(I_2)$ as in (4). Since there are at most T^2 integer intervals in $[1, T]$, a union bound argument leads to that with probability at least $1 - (TM)^{-2}$, Lemma 2 holds uniformly for all integer intervals in $[1, T]$. The claim a. therefore holds. \square

The following lemma is based on the strong convexity of the log-likelihood function.

Lemma 1. *For $A \in \mathcal{C}$, where \mathcal{C} is defined in (6), under Assumption 1, for any $m \in [1, M]$ and $t \in [1, T - 1]$, it holds that*

$$\begin{aligned} \exp[v + A_m g\{\mathcal{X}(t)\}] - \exp[v + A_m^*(t)g\{\mathcal{X}(t)\}] - X_m(t+1)\Delta_m(t)g\{\mathcal{X}(t)\} \\ \geq c[\Delta_m(t)g\{\mathcal{X}(t)\}]^2 + \epsilon_m(t)\Delta_m(t)g\{\mathcal{X}(t)\}, \end{aligned}$$

where $\Delta(t) = A - A^*(t)$ and $\epsilon(t) = X(t+1) - \exp[v + A_m^*(t)g\{\mathcal{X}(t)\}]$.

Proof. For any $C > 0$ and $a, b \in [-C, C]$, due to the strong concavity of $\exp(\cdot)$, it holds that

$$\exp(a) \geq \exp(b) + \exp(a)(b - a) + c(b - a)^2,$$

where c is constant depending on C . Since for any $A \in \mathcal{C}$ and $m \in [1, M]$, there exists a constant C depending on \mathcal{C} and C_g satisfying $|v + A_m g\{\mathcal{X}(t)\}| \leq C$, it holds that

$$\begin{aligned} & \exp[v + A_m g\{\mathcal{X}(t)\}] - \exp[v + A_m^*(t)g\{\mathcal{X}(t)\}] \\ & \geq -\exp[v + A_m^*(t)g\{\mathcal{X}(t)\}]\Delta_m(t)g\{\mathcal{X}(t)\} + c[\Delta_m(t)g\{\mathcal{X}(t)\}]^2, \end{aligned}$$

which leads to the claim. \square

Lemma 2. For $I = (s, e) \subset (0, T + 1)$, assume that I contains only one change point η . Denote $I_1 = (s, \eta]$ and $I_2 = (\eta, e]$. Assume that $\|A^*(I_1) - A^*(I_2)\|_F = \kappa > 0$. If

$$H\{\widehat{A}(I), I\} \leq H\{A^*(I_1), I_1\} + H\{A^*(I_2), I_2\} + \gamma, \quad (12)$$

then with probability at least $1 - 4(TM)^{-4}$, there exists an absolute constant $C > 0$ such that

$$\min\{|I_1|, |I_2|\} \leq C_\epsilon \left(\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2} \right).$$

Proof. Without loss of generality, assume $|I_1| \geq |I_2|$. Equation (12) implies that there exists a constant $c > 0$ such that

$$c \sum_{t \in I} \sum_{m=1}^M [\Delta_m(t)g\{\mathcal{X}(t)\}]^2 \leq - \sum_{t \in I} \sum_{m=1}^M \epsilon_m(t)\Delta_m(t)g\{\mathcal{X}(t)\} \quad (13)$$

$$- \lambda \sqrt{|I|} \|\widehat{A}\|_1 + \lambda \left\{ \sqrt{|I_1|} \|A^*(I_1)\|_1 + \sqrt{|I_2|} \|A^*(I_2)\|_1 \right\} + \gamma, \quad (14)$$

where $\epsilon_m(t) = X_m(t+1) - \exp[v + A_m^*(t)g\{\mathcal{X}(t)\}]$ and $\Delta_m(t) = \widehat{A}_m - A_m^*(t)$.

Step 1. Note that

$$\begin{aligned} & \sum_{t \in I} \sum_{m=1}^M \epsilon_m(t)\Delta_m(t)g\{\mathcal{X}(t)\} = \sum_{t \in I} \sum_{m, m'=1}^M \epsilon_m(t)\Delta_{m, m'}(t)g_{m'}\{\mathcal{X}(t)\} \\ & = \sum_{t \in I} \sum_{(m, m') \in S} \epsilon_m(t)\Delta_{m, m'}(t)g_{m'}\{\mathcal{X}(t)\} + \sum_{t \in I} \sum_{(m, m') \in S^c} \epsilon_m(t)\widehat{A}_{m, m'}g_{m'}\{\mathcal{X}(t)\}. \end{aligned}$$

It follows from Lemma 9 that with probability at least $1 - (TM)^{-4}$, there exists an absolute constant $C > 0$ such that

$$\begin{aligned} & \sum_{t \in I} \sum_{(m, m') \in S} \epsilon_m(t)\Delta_{m, m'}(t)g_{m'}\{\mathcal{X}(t)\} \\ & \leq \sup_{(m, m') \in S} \left| \frac{\sum_{t \in I} \epsilon_m(t)\Delta_{m, m'}(t)g_{m'}\{\mathcal{X}(t)\}}{\sqrt{\sum_{t \in I} \Delta_{m, m'}^2(t)}} \right| \sum_{(m, m') \in S} \sqrt{\sum_{t \in I} \Delta_{m, m'}^2(t)} \end{aligned}$$

$$\leq C \log(TM) \sqrt{d \sum_{t \in I} \|\Delta(t)\|_{\mathbb{F}}^2}.$$

It follows from Lemma 10 that with probability at least $1 - (TM)^{-4}$, there exists an absolute constant $C > 0$ such that

$$\begin{aligned} \sum_{t \in I} \sum_{(m, m') \in S^c} \epsilon_m(t) \widehat{A}_{m, m'} g_{m'}\{\mathcal{X}(t)\} &= \sum_{(m, m') \in S^c} \widehat{A}_{m, m'} \sum_{t \in I} \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\} \\ &\leq C \log^{1/2}(TM) \sqrt{|I|} \|\widehat{A}_{S^c}\|_1. \end{aligned}$$

Step 2. For (14), since both $A^*(I_1)$ and $A^*(I_2)$ are supported on S , it follows that

$$\begin{aligned} & -\lambda \sqrt{|I|} \|\widehat{A}\|_1 + \lambda \left(\sqrt{|I_1|} \|A^*(I_1)\|_1 + \sqrt{|I_2|} \|A^*(I_2)\|_1 \right) \\ &= -\lambda \sqrt{|I|} (\|\widehat{A}_S\|_1 + \|\widehat{A}_{S^c}\|_1) + \lambda \left(\sqrt{|I_1|} \|A_S^*(I_1)\|_1 + \sqrt{|I_2|} \|A_S^*(I_2)\|_1 \right) \\ &\leq -\lambda \sqrt{|I|} \|\widehat{A}_{S^c}\|_1 + \lambda \sqrt{|I_1|} \|A_S^*(I_1) - \widehat{A}_S\|_1 + \lambda \sqrt{|I_2|} \|A_S^*(I_2)\|_1 \\ &\leq -\lambda \sqrt{|I|} \|\widehat{A}_{S^c}\|_1 + \lambda \sqrt{|I_1| d} \|A_S^*(I_1) - \widehat{A}_S\|_{\mathbb{F}} + \lambda \sqrt{|I_2|} d \\ &\leq -\lambda \sqrt{|I|} \|\widehat{A}_{S^c}\|_1 + \lambda \sqrt{d \sum_{t \in I} \|\Delta(t)\|_{\mathbb{F}}^2} + \lambda \sqrt{|I_2|} d. \end{aligned}$$

Step 3. Applying Steps 1 and 2 to (13) and (14), respectively, leads to

$$c \sum_{t \in I} \sum_{m=1, \dots, M} [\Delta_m(t) g\{\mathcal{X}(t)\}]^2 \leq 3\lambda \sqrt{d \sum_{t \in I} \|\Delta(t)\|_{\mathbb{F}}^2} - 2\lambda/3 \sqrt{|I|} \|\widehat{A}_{S^c}\|_1 + \lambda \sqrt{|I_2|} d + \gamma, \quad (15)$$

with $\lambda \geq C_\lambda \log(TM)$. For any $q \in \mathbb{Z}^+ \cap [0, p-1]$, let

$$\mathcal{T}_q = \{pb + q : b \in \mathbb{Z}^+, 1 \leq pb + q \leq T\}.$$

Then

$$\begin{aligned} c \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} [\Delta_m(t) g\{\mathcal{X}(t)\}]^2 &= \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \Delta_m(t) \mathbb{E}[g\{\mathcal{X}(t)\} g\{\mathcal{X}(t)\}^\top | \mathcal{X}(t-p)] \Delta_m^\top(t) \\ &\quad - \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \Delta_m(t) y(t) \Delta_m^\top(t), \end{aligned} \quad (16)$$

where $y(t) = \mathbb{E}[g\{\mathcal{X}(t)\} g\{\mathcal{X}(t)\}^\top | \mathcal{X}(t-p)] - g\{\mathcal{X}(t)\} g\{\mathcal{X}(t)\}^\top \in \mathbb{R}^{M \times M}$. By Assumption A4, (16) implies that

$$\begin{aligned} & c \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} [\Delta_m(t) g\{\mathcal{X}(t)\}]^2 \\ &\geq \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_{\mathbb{F}}^2 \left\{ \omega - \frac{\sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \Delta_m(t) y(t) \Delta_m^\top(t)}{\sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_{\mathbb{F}}^2} \right\}. \end{aligned} \quad (17)$$

Now consider

$$Q_1 = \left\{ q : \omega \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_{\mathbb{F}}^2 \geq 4 \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \Delta_m(t) y(t) \Delta_m^\top(t) \right\}$$

and

$$Q_2 = \left\{ q : \omega \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_{\mathbb{F}}^2 < 4 \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \Delta_m(t) y(t) \Delta_m^\top(t) \right\}.$$

Step 4. For $q \in Q_1$, it follows from (17) and the definition of Q_1 that

$$\sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} [\Delta_m(t) g\{\mathcal{X}(t)\}]^2 \geq \omega/2 \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_{\mathbb{F}}^2. \quad (18)$$

Since $\widehat{A}, A^*(I_1) \in \mathcal{C}$, one has for $t \in I_1$,

$$\begin{aligned} & \max_{m=1, \dots, M} \sum_{m'=1, \dots, M} |\Delta_{m, m'}(t)| \\ & \leq \max_{m=1, \dots, M} \sum_{m'=1, \dots, M} |\widehat{A}_{m, m'}(t)| + \max_{m=1, \dots, M} \sum_{m'=1, \dots, M} |A_{m, m'}^*(I_1)| \leq 2, \end{aligned}$$

due to Assumption **A2**. Since $A^*(I_2) \in \mathcal{C}$, for $t \in I_2$,

$$\max_{m=1, \dots, M} \sum_{m'=1, \dots, M} |\Delta_{m, m'}(t)| \leq 2.$$

Consequently,

$$\max_{m=1, \dots, M} \sum_{m'=1, \dots, M} \sup_{t \in I} |\Delta_{m, m'}(t)| \leq 2. \quad (19)$$

Step 5. For $q \in Q_2$, observe that

$$\begin{aligned} & \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \Delta_m(t) y(t) \Delta_m^\top(t) = \sum_{t \in I \cap \mathcal{T}_q} \sum_{m, m', m''=1, \dots, M} \Delta_{m, m'}(t) y_{m', m''}(t) \Delta_{m, m''}(t) \\ & = \sum_{t \in I \cap \mathcal{T}_q} \sum_{(m, m') \in S} \sum_{m''=1, \dots, M} \Delta_{m, m'}(t) y_{m', m''}(t) \Delta_{m, m''}(t) \end{aligned} \quad (20)$$

$$+ \sum_{t \in I \cap \mathcal{T}_q} \sum_{(m, m') \in S^c} \sum_{m''=1, \dots, M} \widehat{A}_{m, m'} y_{m', m''}(t) \Delta_{m, m''}(t). \quad (21)$$

For Equation (20), due to Lemma 11, it holds with probability at least $1 - (TM)^{-4}$ that

$$\sum_{t \in I \cap \mathcal{T}_q} \sum_{(m, m') \in S} \sum_{m''=1, \dots, M} \Delta_{m, m'}(t) y_{m', m''}(t) \Delta_{m, m''}(t)$$

$$\begin{aligned}
&= \sum_{(m,m') \in S, m''=1, \dots, M} \left\{ \frac{\sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}(t) y_{m',m''}(t) \Delta_{m,m''}(t)}{\sqrt{\sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t) \Delta_{m,m''}^2(t)}} \right\} \sqrt{\sum_{t \in I \cap \mathcal{T}+q} \Delta_{m,m'}^2(t) \Delta_{m,m''}^2(t)} \\
&\leq \max_{m,m',m''=1, \dots, M} \left| \frac{\sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}(t) y_{m',m''}(t) \Delta_{m,m''}(t)}{\sqrt{\sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t) \Delta_{m,m''}^2(t)}} \right| \sum_{(m,m') \in S, m''=1, \dots, M} \sqrt{\sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t) \Delta_{m,m''}^2(t)} \\
&\leq C \log(TM) \sum_{(m,m') \in S, m''=1, \dots, M} \sqrt{\sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t) \Delta_{m,m''}^2(t)} \\
&\leq C \log(TM) \sum_{(m,m') \in S} \sqrt{\sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t)} \max_{m=1, \dots, M} \sum_{m''=1, \dots, M} \sup_{t \in I} |\Delta_{m,m''}(t)| \\
&\leq 2C \log(TM) \sqrt{d} \sqrt{\sum_{(m,m') \in S} \sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t)} \\
&\leq 32C \omega^{-1} \log^2(TM) d + \frac{\omega}{16} \sum_{(m,m') \in S} \sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t),
\end{aligned}$$

where the second inequality follows from Lemma 11 and the fourth inequality follows from (19). For (21), similarly, we have that with probability at least $1 - (TM)^{-4}$,

$$\begin{aligned}
&\sum_{t \in I \cap \mathcal{T}_q} \sum_{(m,m') \in S^c} \sum_{m''=1, \dots, M} \hat{A}_{m,m'} y_{m',m''}(t) \Delta_{m,m''}(t) \\
&= \sum_{m''=1, \dots, M} \sum_{(m,m') \in S^c} \hat{A}_{m,m'} \sum_{t \in I \cap \mathcal{T}_q} y_{m',m''}(t) \Delta_{m,m''}(t) \\
&\leq \sum_{m''=1, \dots, M} \sum_{(m,m') \in S^c} |\hat{A}_{m,m'}| \left| \sum_{t \in I \cap \mathcal{T}_q} \frac{y_{m',m''}(t) \Delta_{m,m''}(t)}{\sqrt{\sum_{t \in I \cap \mathcal{T}+q} \Delta_{m,m''}^2(t)}} \sqrt{\sum_{t \in I \cap \mathcal{T}+q} \Delta_{m,m''}^2(t)} \right| \\
&\leq C \sum_{m''=1, \dots, M} \sum_{(m,m') \in S^c} |\hat{A}_{m,m'}| \log(TM) \sqrt{|I|} \max_{t \in I} |\Delta_{m,m''}(t)| \leq C \|\hat{A}_{S^c}\|_1 \sqrt{|I|} \log(TM)
\end{aligned}$$

where the second inequality follows from Lemma 11 and the last inequality follows from (19). Combining the above calculations with (20) and (21), we have that

$$\begin{aligned}
&\sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \Delta_m(t) y(t) \Delta_m^\top(t) \\
&\leq 32C \omega^{-1} \log^2(TM) d + \frac{\omega}{16} \sum_{(m,m') \in S} \sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t) + C \|\hat{A}_{S^c}\|_1 \sqrt{|I|} \log(TM) \quad (22)
\end{aligned}$$

Since $q \in Q_2$, it holds that

$$\begin{aligned}
&\omega \sum_{q \in Q_2} \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_{\mathbb{F}}^2 \\
&\leq 4 \sum_{q \in Q_2} \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \Delta_m(t) y(t) \Delta_m^\top(t)
\end{aligned}$$

$$\leq Cp\omega^{-1} \log^2(TM)d + \frac{\omega}{4} \sum_{q \in Q_2} \sum_{(m,m') \in S} \sum_{t \in I \cap \mathcal{T}_q} \Delta_{m,m'}^2(t) + \lambda/6 \|\widehat{A}_{S^c}\|_1 \sqrt{|I|}, \quad (23)$$

where the first inequality follows from the definition of Q_2 , and the last inequality follows from (22) and that $\lambda \geq Cp \log(TM)$. Equation (23) directly leads to that

$$\omega \sum_{q \in Q_2} \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_F^2 \leq 2Cp\omega^{-1} \log^2(TM)d + \lambda/3 \|\widehat{A}_{S^c}\|_1 \sqrt{|I|}. \quad (24)$$

Therefore

$$\begin{aligned} & \omega \sum_{t \in I} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_F^2 \\ = & \omega \sum_{q \in Q_1} \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_F^2 + \omega \sum_{q \in Q_2} \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_F^2 \\ \leq & 2 \sum_{q \in Q_1} \sum_{t \in I \cap \mathcal{T}_q} \sum_{m=1, \dots, M} [\Delta_m(t)g\{\mathcal{X}(t)\}]^2 + 2Cp\omega^{-1} \log^2(TM)d + \lambda/3 \|\widehat{A}_{S^c}\|_1 \sqrt{|I|} \\ \leq & \sum_{t \in I} \sum_{m=1, \dots, M} [\Delta_m(t)g\{\mathcal{X}(t)\}]^2 + 2Cp\omega^{-1} \log^2(TM)d + \lambda/3 \|\widehat{A}_{S^c}\|_1 \sqrt{|I|} \\ \leq & 6/c\lambda \sqrt{d \sum_{t \in I} \|\Delta(t)\|_F^2 - 4\lambda/(3c) \sqrt{|I|} \|\widehat{A}_{S^c}\|_1 + 2\lambda/c \sqrt{|I_2|}d + \gamma} \\ & + 2Cp\omega^{-1} \log^2(TM)d + \lambda/3 \|\widehat{A}_{S^c}\|_1 \sqrt{|I|}, \end{aligned}$$

where the first inequality follows from (18) and (24) and the last inequality follows from (15). The above display together directly yields that

$$\omega \sum_{t \in I} \|\Delta(t)\|_F^2 + (\lambda/3) \|\widehat{A}_{S^c}\|_1 \sqrt{|I|} \leq \frac{pd}{\omega} \lambda^2 + \lambda \sqrt{|I_2|}d + \gamma. \quad (25)$$

Step 6. Observe that

$$\begin{aligned} \sum_{t \in I} \|\Delta(t)\|_F^2 & \geq \inf_{B \in \mathbb{R}^{M \times M}} \sum_{t \in I} \|B - A^*(t)\|_F^2 = \inf_{B \in \mathbb{R}^{M \times M}} \{|I_1| \|B - A^*(I_1)\|_F^2 + |I_2| \|B - A^*(I_2)\|_F^2\} \\ & = \frac{|I_1| |I_2|}{|I_1| + |I_2|} \kappa^2 \geq \frac{|I_2|}{2} \kappa^2, \end{aligned}$$

where $|I_1| \geq |I_2|$ is used in the last inequality. Due to (25), it holds that

$$\frac{1}{2} |I_2| \omega \kappa^2 \leq \frac{pd}{\omega} \lambda^2 + \lambda \sqrt{|I_2|}d + \gamma \leq \frac{pd\lambda^2}{\omega} + \left(\frac{8\lambda^2 d^2}{\omega \kappa^2} + \frac{1}{4} |I_2| \omega \kappa^2 \right) + \gamma,$$

where the last inequality follows from Hölder's inequality. We thus have that with probability at least $1 - 4(TM)^{-4}$,

$$|I_2| \leq C_\epsilon \left(\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2} \right).$$

□

Lemma 3. *Under all the assumptions in Theorem 1, let $I = (s, e] \subset [1, T]$ be any interval containing exactly two change points η_{r+1} and η_{r+2} , $I_1 = (s, \eta_{r+1}]$, $I_2 = (\eta_{r+1}, \eta_{r+2}]$ and $I_3 = (\eta_{r+2}, e]$. Let $\kappa_j = \|A^*(I_j) - A^*(I_{j+1})\|_F$ for $j = 1, 2$ and $\kappa = \min\{\kappa_1, \kappa_2\}$. If*

$$H(\widehat{A}(I), I) \leq \sum_{j=1}^3 H(A^*(I_j), I_j) + 2\gamma, \quad (26)$$

then with probability at least $1 - 4(TM)^{-4}$ it holds that

$$\max\{|I_1|, |I_3|\} \leq C_\epsilon \left(\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2} \right).$$

Proof. Without loss of generality, assume that $|I_1| \geq |I_3|$. Then (26) implies that

$$\begin{aligned} c \sum_{t \in I} \sum_{m=1, \dots, M} [\Delta_m(t) g\{\mathcal{X}(t)\}]^2 &\leq \sum_{t \in I} \sum_{m=1, \dots, M} \epsilon_m(t) \Delta_m(t) g\{\mathcal{X}(t)\} \\ &\quad - \lambda \sqrt{|I|} \|\widehat{A}(I)\|_1 + \lambda \sum_{j=1}^3 \left(\sqrt{|I_j|} \|A^*(I_j)\|_1 \right) + 2\gamma, \end{aligned} \quad (27)$$

where $\epsilon_m(t) = \exp[v + A_m(t)g\{\mathcal{X}(t)\}] - X_m(t+1)$ and $\Delta(t) = \widehat{A}(I) - A^*(I_j)$ for $t \in I_j$, $j = 1, 2, 3$.

There are two possible scenarios: (1) $\min\{|I_1|, |I_2|\} \geq |I_3|$ and (2) $|I_3| > |I_2|$. We remark that (2) is simpler than (1), so in the sequel we will assume (1). Let \widetilde{I} and \widetilde{J} be the shorter and longer one between I_1 and I_2 , respectively.

Step 1. For the case Observe that

$$\begin{aligned} & - \lambda \sqrt{|I|} \|\widehat{A}(I)\|_1 + \lambda \sum_{j=1}^3 \left(\sqrt{|I_j|} \|A^*(I_j)\|_1 \right) \\ & \leq - \lambda \sqrt{|I|} \|\widehat{A}(I)\|_1 + \lambda \sqrt{|\widetilde{J}|} \|A^*(\widetilde{J})\|_1 + 2\lambda \sqrt{|\widetilde{I}|} d \\ & \leq - \lambda \sqrt{|I|} \|\widehat{A}_{Sc}(I)\|_1 + \lambda \sqrt{|\widetilde{J}|} \|\widehat{A}_S(I) - A^*(\widetilde{J})\|_1 + 2\lambda \sqrt{|\widetilde{I}|} d \\ & \leq - \lambda \sqrt{|I|} \|\widehat{A}_{Sc}(I)\|_1 + \lambda \sqrt{d \sum_{t \in I} \|\Delta(t)\|_F^2} + 2\lambda \sqrt{|\widetilde{I}|} d, \end{aligned}$$

where the first inequality follows from the assumption that $|I_1| \geq |I_3|$ and the fact that $\|A^*(I_j)\|_1 \leq d$ for all $j = 1, 2, 3$.

Step 2. Following from similar arguments as those in the **Step 1** in the proof of Lemma 2, one has

$$\begin{aligned} & \sum_{t \in I} \epsilon_m(t) \Delta_{m,m'}(t) g_{m'}\{\mathcal{X}(t)\} \\ & \leq C \log(TM) \sqrt{d \sum_{t \in I} \|\Delta(t)\|_F^2} + C \log^{1/2}(TM) \sqrt{|I|} \|\widehat{A}_{Sc}(I)\|_1 \end{aligned}$$

$$\leq \lambda/6 \sqrt{d \sum_{t \in I} \|\Delta(t)\|_{\mathbb{F}}^2} + \lambda/6 \|\widehat{A}_{S^c}(I)\|_1 \sqrt{|I|},$$

where $\lambda \geq C_\lambda \log^{3/2}(TM)$ is used in the last inequality.

Step 3. Following from similar arguments as those in the **Step 5** in the proof of Lemma 2, one has

$$\omega \sum_{t \in I} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_{\mathbb{F}}^2 \leq 2 \sum_{t \in I} \sum_{m=1, \dots, M} [\Delta_m(t) g\{\mathcal{X}(t)\}]^2 + 2C\omega^{-1} p \log^2(TM) d + \lambda/3 \sqrt{|I|} \|\widehat{A}_{S^c}(I)\|_1$$

Step 4. Combing all the previous steps gives

$$\omega \sum_{t \in I} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_{\mathbb{F}}^2 + \lambda/3 \sqrt{|I|} \|\widehat{A}(I)\|_1 \leq 4C\omega^{-1} p \lambda^2 d + 2\lambda d \sqrt{|I|} + 2\gamma. \quad (28)$$

Observe that

$$\begin{aligned} \sum_{t \in I} \|\Delta(t)\|_{\mathbb{F}}^2 &\geq \sum_{t \in I_1 \cup I_2} \|\Delta(t)\|_{\mathbb{F}}^2 \geq \inf_{B \in \mathbb{R}^{M \times M}} \sum_{t \in I_1 \cup I_2} \|B - A^*(t)\|_{\mathbb{F}}^2 \\ &= \inf_{B \in \mathbb{R}^{M \times M}} \{|I_1| \|B - A^*(I_1)\|_{\mathbb{F}}^2 + |I_2| \|B - A^*(I_2)\|_{\mathbb{F}}^2\} = \frac{|I_1| |I_2|}{|I_1| + |I_2|} \kappa^2 \geq \frac{\min\{|I_1|, |I_2|\}}{2} \kappa^2. \end{aligned}$$

If $|I_2| \leq |I_1|$, then it follows that

$$\frac{|I_2| \kappa^2 \omega}{2} \leq 4C\omega^{-1} p \lambda^2 d + 2\lambda d \sqrt{|I_2|} + 2\gamma \leq 4C\omega^{-1} p \lambda^2 d + \frac{\Delta \kappa^2 \omega}{4} + \frac{8\lambda^2 d^2}{\kappa^2 \omega} + 2\gamma.$$

This leads to that

$$|I_2| \lesssim \frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2},$$

which contradicts with Assumption **A3**. Then there exists an absolute constant such that

$$|I_1| \leq C_\epsilon \left(\frac{\lambda^2 d}{\kappa^2} + \frac{p \lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2} \right).$$

Since by assumption, $|I_3| \leq |I_1|$, the desired results follows. \square

Lemma 4. Under all the assumptions in Theorem 1, let $I = (s, e] \subset [1, T]$ be any interval which contains no change point. Let I_1 and I_2 be two intervals such that $I_1 \sqcup I_2 = I$. Then with probability at least $1 - (TM)^{-4}$,

$$H(\widehat{A}(I_1), I_1) + H(\widehat{A}(I_2), I_2) + \gamma \geq H(A^*(I), I).$$

Proof. We prove by contradiction, assuming that

$$H(\widehat{A}(I_1), I_1) + H(\widehat{A}(I_2), I_2) + \gamma < H(A^*(I), I).$$

Denote

$$\Delta(t) = \begin{cases} \widehat{A}(I_1) - A^*(I), & t \in I_1; \\ \widehat{A}(I_2) - A^*(I), & t \in I_2. \end{cases}$$

Standard calculations give

$$c \sum_{t \in I} \sum_{m=1, \dots, M} [\Delta_m(t) g\{\mathcal{X}(t)\}]^2 + \gamma \leq \sum_{t \in I} \sum_{m=1, \dots, M} \epsilon_m(t) \Delta_m(t) g\{\mathcal{X}(t)\} \quad (29)$$

$$+ \lambda \sqrt{|I|} \|A^*(I)\|_1 - \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}(I_2)\|_1 \right\}, \quad (30)$$

where $\epsilon_m(t) = \exp[v + A_m(t) g\{\mathcal{X}(t)\}] - X_m(t + 1)$.

Step 1. For (30), we have that

$$\begin{aligned} & \lambda \sqrt{|I|} \|A^*(I)\|_1 - \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}(I_2)\|_1 \right\} \\ & \leq \lambda (\sqrt{|I_1|} + \sqrt{|I_2|}) \|A^*(I)\|_1 - \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}_S(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}_S(I_2)\|_1 \right\} \\ & \quad - \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}_{S^c}(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}_{S^c}(I_2)\|_1 \right\} \\ & \leq \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}_S(I_1) - A^*(I)\|_1 + \sqrt{|I_2|} \|\widehat{A}_S(I_2) - A^*(I)\|_1 \right\} \\ & \quad - \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}_{S^c}(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}_{S^c}(I_2)\|_1 \right\} \\ & \leq \lambda \sqrt{2|I_1| \|\widehat{A}_S(I_1) - A^*(I)\|_1^2 + 2|I_2| \|\widehat{A}_S(I_2) - A^*(I)\|_1^2} \\ & \quad - \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}_{S^c}(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}_{S^c}(I_2)\|_1 \right\} \\ & \leq \lambda \sqrt{2|I_1| d \|\widehat{A}_S(I_1) - A^*(I)\|_F^2 + 2|I_2| d \|\widehat{A}_S(I_2) - A^*(I)\|_F^2} \\ & \quad - \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}_{S^c}(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}_{S^c}(I_2)\|_1 \right\} \\ & = \sqrt{2\lambda d \sum_{t \in I} \|\Delta(t)\|_F^2} - \lambda \left\{ \sqrt{|I_1|} \|\widehat{A}_{S^c}(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}_{S^c}(I_2)\|_1 \right\}. \end{aligned}$$

Step 2. Using similar calculations as **Step 1** in the proof of Lemma 2, one has that with probability at least $1 - (TM)^4$,

$$\begin{aligned} & \sum_{t \in I} \epsilon_m(t) \Delta_{m,m'}(t) g_{m'}\{\mathcal{X}(t)\} \\ & \leq C \log(TM) \sqrt{d \sum_{t \in I} \|\Delta(t)\|_F^2} + C \log^{1/2}(TM) \left\{ \|\widehat{A}_{S^c}(I_1)\|_1 \sqrt{|I_1|} + \|\widehat{A}_{S^c}(I_2)\|_1 \sqrt{|I_2|} \right\} \\ & \leq \lambda/6 \sqrt{d \sum_{t \in I} \|\Delta(t)\|_F^2} + \lambda/6 \left\{ \|\widehat{A}_{S^c}(I_1)\|_1 \sqrt{|I_1|} + \|\widehat{A}_{S^c}(I_2)\|_1 \sqrt{|I_2|} \right\}, \end{aligned}$$

where $\lambda \geq C_\lambda \log(TM)$ is used in the last inequality.

Step 3. Using similar calculations as **Step 5** in the proof of Lemma 2, one has that with probability at least $1 - (TM)^4$,

$$\omega \sum_{t \in I} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_F^2 \leq \omega/c \sum_{t \in I} \sum_{m=1, \dots, M} [\Delta_m(t) g\{\mathcal{X}(t)\}]^2$$

$$+ 2C\omega^{-1}p \log^2(TM)d + \lambda/3 \left\{ \|\widehat{A}_{S^c}(I_1)\|_1 \sqrt{|I_1|} + \|\widehat{A}_{S^c}(I_2)\|_1 \sqrt{|I_2|} \right\}.$$

Step 4. Combing all the previous steps gives

$$\omega \sum_{t \in I} \sum_{m=1, \dots, M} \|\Delta_m(t)\|_F^2 + \gamma + \lambda/3 \left\{ \sqrt{|I_1|} \|\widehat{A}(I_1)\|_1 + \sqrt{|I_2|} \|\widehat{A}(I_2)\|_1 \right\} \leq 4C\omega^{-1}p \log^2(TM)d,$$

which directly implies that $\gamma \leq 2C_\epsilon p \lambda^2 d$. This leads to a contradiction with (8) and completes the proof. \square

Lemma 5. *Under all the assumptions in Theorem 1, let $I = (s, e] \subset [1, T]$ satisfying $I \cap \{\eta_k\}_{k=1}^K = \{\eta_{r+1}, \dots, \eta_{r+J}\}$, with $J \geq 3$. Denote $I_1 = (s, \eta_{r+1}]$, $I_j = (\eta_{r+j-1}, \eta_{r+j}]$ for all $j = 2, \dots, J$ and $I_{J+1} = (\eta_{r+J}, e]$. Denote $\kappa_j = \|A^*(I_j) - A^*(I_{j+1})\|_F$ for $j = 1, \dots, J$ and $\kappa = \min_{j=1, \dots, J} \kappa_j$. It holds that probability at least $1 - (TM)^{-4}$,*

$$H(\widehat{A}(I), I) > \sum_{j=1}^{J+1} H(A^*(I_j), I_j) + J\gamma. \quad (31)$$

Proof. We prove by contradiction, assuming that

$$H(\widehat{A}(I), I) \leq \sum_{j=1}^{J+1} H(A^*(I_j), I_j) + J\gamma.$$

Note that by assumption **A3** $\min\{|I_2|, |I_3|\} \geq cT$. Without loss of generality, assume that $|I_2| \geq |I_3|$. Equation (31) implies that

$$\sum_{t \in I} \sum_{1 \leq m \leq M} (\Delta_m(t)g(\mathcal{X}(t)))^2 \leq \sum_{t \in I} \sum_{1 \leq m \leq M} \epsilon_m(t) \Delta_m(t)^\top g(\mathcal{X}(t)) \quad (32)$$

$$- \lambda \sqrt{|I|} \|\widehat{A}(I)\|_1 + \lambda \sum_{j=1}^{J+1} \left(\sqrt{|I_j|} \|A^*(I_j)\|_1 \right) + \gamma \quad (33)$$

where $\epsilon_m(t) = \exp(v + A_m(t)g(\mathcal{X}(t))) - X_m(t+1)$ and $\Delta(t) = \widehat{A}(I) - A^*(I_j)$ for $t \in I_j \subset I$.

Step 1. Observe that by assumption **A3** there exists $c > 0$ such that $|I_3| \geq cT \geq c|I_j|$ for any $1 \leq j \leq J$. Therefore

$$\begin{aligned} & - \lambda \sqrt{|I|} \|\widehat{A}(I)\|_1 + \lambda \sum_{j=1}^{J+1} \left(\sqrt{|I_j|} \|A^*(I_j)\|_1 \right) \\ & \leq - \lambda \sqrt{|I|} \|\widehat{A}(I)\|_1 + \lambda \sqrt{|I_2|} \|A^*(I_2)\|_1 + 2 \sum_{j \neq 2} \lambda \sqrt{|I_j|} |S| \\ & \leq - \lambda \sqrt{|I|} \|\widehat{A}_{S^c}(I)\|_1 + \lambda \sqrt{|I_2|} \|\widehat{A}_S(I) - A^*(I_2)\|_1 + c^{-1/2} \lambda \sqrt{|I_3|} |S| \\ & \leq - \lambda \sqrt{|I|} \|\widehat{A}_{S^c}(I)\|_1 + \lambda \sqrt{|S|} \sum_{t \in I} \|\Delta(t)\|_F^2 + c^{-1/2} \lambda \sqrt{|I_3|} |S| \end{aligned}$$

where the first inequality follows from the fact that $\|A^*(I_j)\|_1 \leq |S|$ for all $1 \leq j \leq J$ and the second inequality follows from the observation that $|I_3| \geq c|I_j|$.

Step 2. Using similar calculations as **Step 1** in the proof of Lemma 2, one has

$$\begin{aligned} & \sum_{t \in I} \epsilon_m(t) \Delta_{m,m'}(t) g_{m'}(\mathcal{X}(t)) \\ & \leq C \log^{3/2}(TM) \sqrt{|S| \sum_{t \in I} \|\Delta(t)\|_{\mathbb{F}}^2} + C \log^{3/2}(TM) \sqrt{|I|} \|\widehat{A}_{S^c}(I)\|_1 \\ & \leq (\lambda/6) \sqrt{|S| \sum_{t \in I} \|\Delta(t)\|_{\mathbb{F}}^2} + (\lambda/6) \|\widehat{A}_{S^c}(I)\|_1 \sqrt{|I|} \end{aligned}$$

where $\lambda \geq C_\lambda \log^{3/2}(TM)$ is used in the last inequality.

Step 3. Using similar calculations as **Step 6** in the proof of Lemma 2, one has

$$\omega \sum_{t \in I} \sum_{1 \leq m \leq M} \|\Delta_m(t)\|_{\mathbb{F}}^2 \leq \sum_{t \in I} \sum_{1 \leq m \leq M} (\Delta_m(t)^\top g(\mathcal{X}(t)))^2 + 2C\omega^{-1}p \log^2(TM) |S| + (\lambda/3) \sqrt{|I|} \|\widehat{A}_{S^c}(I)\|_1$$

Step 4. Combing all the previous steps gives

$$\omega \sum_{t \in I} \sum_{1 \leq m \leq M} \|\Delta_m(t)\|_{\mathbb{F}}^2 + (\lambda/3) \sqrt{|I|} \|\widehat{A}(I)\|_1 \leq 4C\omega^{-1}p \lambda^2 |S| + 2\lambda |S| \sqrt{|I_3|} + J\gamma. \quad (34)$$

Observe that

$$\begin{aligned} \sum_{t \in I} \|\Delta(t)\|_{\mathbb{F}}^2 & \geq \sum_{t \in I_2 \cup I_3} \|\Delta(t)\|_{\mathbb{F}}^2 \geq \inf_B \sum_{t \in I_2 \cup I_3} \|B - A^*(t)\|_{\mathbb{F}}^2 = \inf_B |I_2| \|B - A^*(I_2)\|_{\mathbb{F}}^2 + |I_3| \|B - A^*(I_3)\|_{\mathbb{F}}^2 \\ & = \frac{|I_2||I_3|}{|I_2| + |I_3|} \kappa^2 \geq \frac{\min\{|I_2|, |I_3|\}}{2} \kappa^2 \geq \frac{c|I_3|}{2} \kappa^2, \end{aligned}$$

where $|I_2| \geq |I_3| \geq c|I_2|$ is used in the last inequality. So (34) implies that

$$\frac{c}{2} |I_3| \omega \kappa^2 \leq \frac{Cp|S|}{\omega} \lambda^2 + \lambda \sqrt{|I_3|} |S| + J\gamma \leq \frac{Cp|S|}{\omega} \lambda^2 + \left(\frac{8\lambda^2 |S|^2}{c\omega \kappa^2} + \frac{c}{4} |I_3| \omega \kappa^2 \right) + J\gamma.$$

Note that $J \leq K \leq c^{-1}$ and therefore this implies

$$|I_3| \leq C_\omega \left(\frac{p\lambda^2 |S|}{\kappa^2} + \frac{p\lambda^2 |S|^2}{\kappa^4} + \frac{\gamma}{\kappa^2} \right).$$

Since by assumption, $|I_3| \leq |I_2|$. This leads to the contradiction with Assumption **A3** and completes the proof. \square

C Proof of Proposition 2

Proof of Proposition 2. For a collection of generic strictly increasing time points $\{\eta'_j\}_{j=0}^{J+1}$, where $\eta'_0 = 1$ and $\eta'_{J+1} = T + 1$, denote $I_j = [\eta'_{j-1}, \eta'_j]$ and

$$\mathcal{L}(\{\eta'_j\}_{j=0}^{J+1}) = \sum_{j=1}^{J+1} H(\widehat{A}(I_j), I_j).$$

In addition assume that $\{\eta_k\}_{k=1}^K \subset \{\eta'_j\}_{j=1}^J$ so that $A^*(t)$ is unchanged in each of the interval I_j . Let

$$\mathcal{L}^*(\{\eta'_j\}_{j=0}^{J+1}) = \sum_{j=1}^{J+1} H(A^*(I_j), I_j).$$

Let $\{\widehat{\eta}_k\}_{k=1}^{\widehat{K}}$ denote the change points induced by $\widehat{\mathcal{P}}$. If one can show that

$$\begin{aligned} & \mathcal{L}^*(\eta_1, \dots, \eta_K) + K\gamma \\ & \geq \mathcal{L}(\eta_1, \dots, \eta_K) + K\gamma \end{aligned} \tag{35}$$

$$\geq \mathcal{L}(\widehat{\eta}_1, \dots, \widehat{\eta}_{\widehat{K}}) + \widehat{K}\gamma \tag{36}$$

$$\geq \mathcal{L}^*(\mathbf{Sort}(\widehat{\eta}_1, \dots, \widehat{\eta}_{\widehat{K}}, \eta_1, \dots, \eta_K)) + \widehat{K}\gamma - CdK\lambda^2 - CK\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}} \tag{37}$$

and that

$$\mathcal{L}^*(\eta_1, \dots, \eta_K) \leq \mathcal{L}^*(\mathbf{Sort}(\widehat{\eta}_1, \dots, \widehat{\eta}_{\widehat{K}}, \eta_1, \dots, \eta_K)), \tag{38}$$

then it must hold that $|\widehat{\mathcal{P}}| = K$. To see this, if $\widehat{K} \geq K + 1$, then

$$\gamma \leq (\widehat{K} - K)\gamma \leq CdK\lambda^2 + CK\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}}, \tag{39}$$

which is a contradiction to (8) if C_γ is sufficiently large.

Since $\widehat{A}(I) = \operatorname{argmin}_{A \in \mathcal{C}} H(A, I)$, we have that

$$H(\widehat{A}(I), I) \leq H(A^*(I), I).$$

Then (35) is a direct consequence of the above display.

Moreover, (36) is a direct consequence of (7).

To show (37), consider any $I = (s, e] \in \widehat{\mathcal{P}}$. By Proposition 1, with probability at least $1 - C(TM)^4$, I contains at most two change points. Therefore the three cases in Lemma 7 directly lead to

$$\mathcal{L}(\widehat{\eta}_1, \dots, \widehat{\eta}_{\widehat{K}}) \geq \mathcal{L}^*(\mathbf{Sort}(\widehat{\eta}_1, \dots, \widehat{\eta}_{\widehat{K}}, \eta_1, \dots, \eta_K)) - CKd\lambda^2 - CK\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}}.$$

Lastly, suppose I is any generic interval in $[1, T]$ containing no change points and that $I_1 \sqcup I_2 = I$. For any $a, b > 0$, we have that $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$. This inequality directly implies that

$$H(A^*(I), I) \leq H(A^*(I_1), I_1) + H(A^*(I_2), I_2). \tag{40}$$

Note that (38) is a straight forward consequence of Equation (40). This completes the proof. \square

Lemma 6 (Standard GLM inequality). *Under all the assumptions in Theorem 1, suppose I is any generic interval such that $|I| \geq Cp|S| \log^2(TM)$ for sufficiently large C and that I contains no change points. Let*

$$\widehat{A}(I) = \arg \min_{A \in \mathcal{C}} H(A, I),$$

where $H(A, I)$ is defined in (5). Then

$$\|\widehat{A}(I) - A^*(I)\|_{\mathbb{F}}^2 \leq \frac{Cd \log^2(TM)}{|I|} \quad \text{and} \quad \|\widehat{A}(I) - A^*(I)\|_1 \leq \frac{Cd \log(TM)}{\sqrt{|I|}}.$$

Lemma 6 is an immediate consequence of Lemma 1 and results in Appendix D, based on standard Lasso arguments (e.g. Wang et al., 2019).

Corollary 1. *Under all the assumptions in Theorem 1 and notation in Lemma 6, it holds that with probability at least $1 - C(TM)^{-4}$,*

$$H(\widehat{A}(I), I) \geq H(A^*(I), I) + C \log^2(TM)d.$$

Proof. Denote that $\Delta = \widehat{A}(I) - A^*(I)$ and that $\epsilon_m = \exp[v + A_m^*(I)g\{\mathcal{X}(t)\}] - X_m(t+1)$. Observe that

$$\begin{aligned} & \sum_{t \in I} \sum_{m=1, \dots, M} \exp[v + \widehat{A}_m(I)g\{\mathcal{X}(t)\}] - \exp[v + A_m^*(I)g\{\mathcal{X}(t)\}] - X_m(t+1) \Delta_m g\{\mathcal{X}(t)\} \\ & \geq \sum_{t \in I} \sum_{m=1, \dots, M} \epsilon_m \Delta_m g\{\mathcal{X}(t)\} + c_\sigma \left[\{\widehat{A}_m(I) - A_m^*(I)\} g\{\mathcal{X}(t)\} \right]^2 \\ & \geq \sum_{t \in I} \sum_{m=1, \dots, M} \epsilon_m \Delta_m g\{\mathcal{X}(t)\} \geq -C\sqrt{|I|} \log(TM) \sum_{m=1, \dots, M} \|\Delta_m\|_1 \geq -C \log^2(TM)d, \end{aligned}$$

where the third inequality follows from Lemma 10 and the last inequality follows from Lemma 6. \square

Lemma 7. *Under all the conditions in Theorem 1, suppose $I = (s, e]$ is any interval in $[1, T]$ containing at most two change points.*

Case 1. *If I contains no change points, then with probability at least $1 - (TM)^{-2}$,*

$$H(A^*(I), I) \leq H(\widehat{A}(I), I) + Cd\lambda^2 + C\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}}.$$

Case 2. *If I contains exactly one change point η_k , letting $J_1 = (s, \eta_k]$ and $J_2 = (\eta_k, e]$, then with probability at least $1 - (TM)^{-2}$,*

$$H(A^*(J_1), J_1) + H(A^*(J_2), J_2) \leq H(\widehat{A}(I), I) + Cd\lambda^2 + C\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}}.$$

Case 3. *If I contains exactly two change points η_k and η_{k+1} , letting $J_1 = (s, \eta_k]$, $J_2 = (\eta_k, \eta_{k+1}]$ and $J_3 = (\eta_{k+1}, e]$, then with probability at least $1 - (TM)^{-2}$,*

$$\sum_{j=1}^3 H(A^*(J_j), J_j) \leq H(\widehat{A}(I), I) + Cd\lambda^2 + C\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}}.$$

Proof. We only prove **Case 3**, as the other two cases are easier and similar. Since $|J_2| \geq \Delta \geq Cd^2 \log^2(TM)$, Observe that by definition of H ,

$$H(A^*(J_2), J_2) \leq H(\widehat{A}(J_2), J_2) + Cd\lambda^2 \leq H(\widehat{A}(I), J_2) + Cd\lambda^2, \quad (41)$$

where the first inequality follows from Corollary 1 and the second inequality follows from the definition of $\widehat{A}(J_2)$.

In addition, observe that with probability at least $1 - C(TM)^{-4}$,

$$\begin{aligned} & H(A^*(J_1), J_1) - \sum_{t \in J_1} \sum_{m=1, \dots, M} \exp[v + \widehat{A}_m(I)g\{\mathcal{X}(t)\}] - \lambda\sqrt{|J_1|}\|A^*(J_1)\|_1 \\ &= \sum_{t \in J_1} \sum_{m=1, \dots, M} \exp[v + A_m^*(J_1)g\{\mathcal{X}(t)\}] - \exp[v + \widehat{A}_m(I)g\{\mathcal{X}(t)\}] + \lambda\sqrt{|J_1|}\|A^*(J_1)\|_1 \\ &\leq \sum_{t \in J_1} \sum_{m=1, \dots, M} \{\widehat{A}_m(I) - A_m^*(J_1)\}g\{\mathcal{X}(t)\}\epsilon_m(t) - c \left[\{\widehat{A}_m(I) - A_m^*(J_1)\}g\{\mathcal{X}(t)\} \right]^2 + \lambda d\sqrt{|J_1|} \\ &\leq \|\widehat{A}(I) - A^*(J_1)\|_1 \max_{m, m'=1, \dots, M} \left| \sum_{t \in J_1} g_{m'}\{\mathcal{X}(t)\}\epsilon_m(t) \right| + \lambda d\sqrt{|J_1|} \\ &\leq C\lambda d\sqrt{|J_1|} + \lambda d\sqrt{|J_1|} \\ &\leq C\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}}, \end{aligned} \quad (42)$$

where the first inequality follows from the same argument in Lemma 1, the third inequality follows from Lemma 10 and the last inequality follows from Proposition 1 b.

In addition, it holds that

$$H(A^*(J_3), J_3) - \sum_{t \in J_3} \sum_{m=1}^M \exp[v + \widehat{A}_m(I)g\{\mathcal{X}(t)\}] \leq C\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}}. \quad (43)$$

Therefore

$$\begin{aligned} & \sum_{j=1}^3 H(A^*(J_j), J_j) \\ &\leq H(\widehat{A}(I), J_2) + \sum_{t \in J_1 \cup J_3} \sum_{m=1}^M \exp[v + \widehat{A}_m(I)g\{\mathcal{X}(t)\}] + C\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}} + Cd\lambda^2 \\ &\leq H(\widehat{A}(I), I) + C\lambda^2 d \sqrt{\frac{\lambda^2 d}{\kappa^2} + \frac{\lambda^2 d^2}{\kappa^4} + \frac{\gamma}{\kappa^2}} + Cd\lambda^2. \end{aligned}$$

where the first inequality follows from Equation (41), Equation (42) and Equation (43), and the second inequality follows from the observation that

$$\lambda\sqrt{|J_2|}\|\widehat{A}(I)\| \leq \lambda\sqrt{|I|}\|\widehat{A}(I)\|.$$

□

D Deviation Bounds

Lemma 8. *Under all the assumptions in Theorem 1, for any $t \in [1, T - 1]$, let $\epsilon(t) = X(t + 1) - \exp[v + A_m^*(t)g\{\mathcal{X}(t)\}]$. For any deterministic $v \in \mathbb{R}^T$ and any integer interval $I \subset [1, T - 1]$, it holds that for any $\delta > 0$,*

$$\max_{m, m'=1, \dots, M} \mathbb{P} \left[\sum_{t \in I} v_t \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\} \geq \delta \right] \leq 2 \exp \left(-\frac{C\delta^2}{\sum_{t \in I} v_t^2} \right).$$

Proof. For any $m, m' \in [1, M]$ and $t \in [1, T - 1]$, let $Y_t = \sum_{i=1}^t v_i \epsilon_m(i) g_{m'}\{\mathcal{X}(i)\}$. Due to Model 1, we have that $\{Y_t\}$ is a martingale sequence with respect to the filtration $\{\mathcal{F}_t\}$, $\mathcal{F}_t = \sigma\{X_1, \dots, X_t\}$. In addition, with $Y_0 = 0$, for any $t \in [1, T - 1]$,

$$\mathbb{E}(|Y_t - Y_{t-1}|) = \mathbb{E}(|v_t \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\}|) \leq C|v_t|.$$

The final result follows from Azuma's inequality (Azuma, 1967). \square

Lemma 9. *Under all the assumptions in Theorem 1, let $I \subset [1, T]$ be an integer interval and $R \in \mathbb{Z}_+ \cap [1, T]$. Denote the event*

$$\mathcal{A}_R(I) = \left\{ \max_{m, m'=1, \dots, M} \sup_{v: \|Dv\|_0=R, \|v\|_2=1} \sum_{t \in I} v_t \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\} \geq \sqrt{C \log^2(TM)R} \right\},$$

for some sufficiently large constant C . It holds that $\mathbb{P}\{\mathcal{A}_R(I)\} \leq (TM)^{-4R}$.

Proof. In this proof, we use C to refer an absolute constant, which is not necessarily the same throughout the proof.

For any $R \in [1, T] \cap \mathbb{Z}_+$, since $\|Dv\|_0 = R$, v has exactly R change points and there are at most $\binom{T}{R}$ possible choices of the locations of change points. Given the collection of change points $\{\eta_k\}_{k=1}^R \subset [1, T]$, denote by $\mathcal{S}(\{\eta_k\}_{k=1}^R)$ the linear subspace of all piecewise-constant functions with all change points at $\{\eta_k\}_{k=1}^R$. Let $\mathcal{N}_{1/4}(\{\eta_k\}_{k=1}^R)$ be a $1/4$ -net of $\mathcal{S}(\{\eta_k\}_{k=1}^R) \cap \mathcal{S}(0, 1)$, where $\mathcal{S}(0, 1)$ is the unit sphere in \mathbb{R}^T . Since $\mathcal{S}(\{\eta_k\}_{k=1}^R)$ is an affine subspace with dimension $R+1$, $\mathcal{N}_{1/4}(\{\eta_k\}_{k=1}^R)$ can be chosen such that $|\mathcal{N}_{1/4}(\{\eta_k\}_{k=1}^R)| \leq 12^{R+1}$, see e.g. Lemma 4.1 in Pollard (1990) and Lemma 4.2.8 in Vershynin (2018).

Then we have for any fixed $m, m' \in [1, M]$ and any set of $\{\eta_k\}_{k=1}^R \subset [1, T]$, it holds that

$$\begin{aligned} & \mathbb{P} \left[\sup_{v \in \mathcal{S}(\{\eta_k\}_{k=1}^R), \|v\|_2=1} \sum_{t \in I} v_t \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\} \geq \sqrt{C \log^2(TM)R} \right] \\ & \leq \mathbb{P} \left[\sup_{v \in \mathcal{N}_{1/4}(\{\eta_k\}_{k=1}^R)} \sum_{t \in I} v_t \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\} + \max_{t \in I} |4^{-1} \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\}| \geq \sqrt{C \log^2(TM)R} \right] \\ & \leq \mathbb{P} \left[\sup_{v \in \mathcal{N}_{1/4}(\{\eta_k\}_{k=1}^R)} \sum_{t \in I} v_t \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\} \geq \sqrt{C \log^2(TM)R} \right] \\ & \quad \times \mathbb{P} \left[\max_{t \in I} |4^{-1} \epsilon_m(t) g_{m'}\{\mathcal{X}(t)\}| < C \log(TM) \right] \end{aligned}$$

$$\begin{aligned}
& + \mathbb{P} \left[\max_{t \in I} |4^{-1} \epsilon_m(t) g_{m'} \{\mathcal{X}(t)\}| \geq C \log(TM) \right] \\
\leq & 12^{R+1} \sup_{v \in \mathcal{N}_{1/4}(\{\eta_k\}_{k=1}^R)} \mathbb{P} \left[\sum_{t \in I} v_t \epsilon_m(t) g_{m'} \{\mathcal{X}(t)\} \geq \sqrt{C \log^2(TM) R} \right] \\
& + \mathbb{P} \left[\max_{t \in I} |4^{-1} \epsilon_m(t) g_{m'} \{\mathcal{X}(t)\}| \geq C \log(TM) \right] \\
\leq & 12^{R+1} \times 2 \exp \left\{ -\frac{C \log^2(TM) R}{\sum_{t \in I} v_t^2} \right\} + 2T \exp \left(-\frac{4C \log(TM)}{C_g} \right) \\
\leq & C \exp \{-C \log(TM) R + R \log(12)\},
\end{aligned}$$

where the fourth inequality follows from Lemma 8 and the sub-Exponential property of Poisson random variables.

Therefore for any fixed integer interval I and positive constant $R \leq T$, it holds that

$$\mathbb{P}\{\mathcal{A}_R(I)\} \leq M^2 \binom{T}{R} C \exp\{-C \log(TM) R\} \leq (TM)^{-4R},$$

with sufficiently large absolute constants. \square

Lemma 10. *Let $I \subset [1, T]$ be an integer interval. Denote the event*

$$\mathcal{B}(I) = \left\{ \max_{m, m'=1, \dots, M} \sum_{t \in I} \epsilon_m(t) g_{m'} \{\mathcal{X}(t)\} \geq \sqrt{C|I| \log(TM)} \right\},$$

for some sufficiently large constant C . It holds that $\mathbb{P}\{\mathcal{B}(I)\} \leq (TM)^{-4}$.

Proof. It follows from Lemma 8 that, with sufficiently large constants,

$$\begin{aligned}
& \mathbb{P} \left\{ \max_{m, m'=1, \dots, M} \sum_{t \in I} \epsilon_m(t) g_{m'} \{\mathcal{X}(t)\} \geq \sqrt{C|I| \log(TM)} \right\} \\
\leq & M^2 \mathbb{P} \left\{ \sum_{t \in I} \epsilon_1(t) g_2 \{\mathcal{X}(t)\} \geq \sqrt{C|I| \log(TM)} \right\} \leq 2M^2 \exp \left(-\frac{C|I| \log(TM)}{2C_g^2 L^2 \sum_{t \in I} |I|} \right) \leq (TM)^{-4}.
\end{aligned}$$

\square

For any $q = 0, \dots, p-1$, denote

$$\mathcal{T}_q = \{pb + q : b \in \mathbb{Z}^+, pb + q = 1, \dots, T\}.$$

Lemma 11. *Under all the assumptions in Theorem 1, let $y(t) = \mathbb{E}[g\{\mathcal{X}(t)\}g\{\mathcal{X}(t)\}^\top | \mathcal{X}(t-p)] - g\{\mathcal{X}(t)\}g\{\mathcal{X}(t)\}^\top \in \mathbb{R}^{M \times M}$, $R \in \mathbb{Z} \cap [1, T]$, $I \subset [1, T]$ be an integer interval and*

$$\mathcal{C}_R(I) = \left\{ \max_{m, m'=1, \dots, M} \sup_{v: \|Dv\|_0=R, \|v\|_2=1} \sum_{t \in I \cap \mathcal{T}_q} v_t y_{m, m'}(t) \geq \sqrt{C \log(TM)} \right\}.$$

It holds that $\mathbb{P}\{\mathcal{C}_R(I)\} \leq (TM)^{-4}$, with sufficiently large $C > 0$.

Proof. Let v be any fixed vector in \mathbb{R}^T . It holds that

$$\mathbb{E}\{v_t y(t) | \mathcal{F}_{t-p}\} = 0 \in \mathbb{R}^{M \times M}.$$

As a result, $\{Z_t\}_{t \in \mathcal{T}_q}$ is a martingale sequence with respect to the filtration $\{\mathcal{F}_t\}_{t \in \mathcal{T}_q}$, where $Z_t = \sum_{s \in \mathcal{T}_q, s \leq t} v_s y_{m,m'}(s)$. In addition, $\mathbb{E}|Z_t - Z_{t-p}| = \mathbb{E}|v_t y_{m,m'}| \leq 2C_g |v_t|$. By Azuma's inequality (Azuma, 1967), for any $m, m' \in [1, M]$,

$$\mathbb{P} \left\{ \sum_{t \in I} v_t y_{m,m'}(t) \geq \sqrt{C \log(TM)} \right\} \leq \exp \left\{ -\frac{C \log(TM)}{\sum_{t \in I} v_t^2} \right\}.$$

The desired result follows from similar covering arguments as in Lemma 9. □