

A CLASS OF n -TUPLES WITH FINITE ASCENT AND PERTURBATIONS

Boulbeba ABDELMOUMEN, Hafedh DAMMAK and Sonia YENGUI

Département de Mathématiques, Faculté des sciences de Sfax,
Route de Soukra, Km 3.5, BP 1171, 3000, Sfax, Tunisie.

e-mail: boulbeba.abbelmoumen@ipeis.rnu.tn
e-mail: sonia.yengui@ipeis.rnu.tn
e-mail: dammak-hafedh@yahoo.fr

Abstract. In this paper, we are interested to study the stability of the ascent of a mutually commuting n -tuple T submitted to a perturbation by an n -tuple S which commutes mutually with T . This study lead us to generalize some well known results for operators and introduce a class of n -tuples in which we obtain the stability of the ascent of T .

Keywords: MSC (2010) Perturbation theory, Fredholm n -tuple, invariant subspaces.

Mathematics Subject Classifications: 47A55, 47A53, 47A15.

1 Introduction

The problem of stability of the semi-Fredholm operators under additive perturbations has been around for many years (see for instance [1, 2, 3, 4, 5, 6]). It is obvious to ask if these results are extended for some class of n -tuples.

The ultimate aim of this paper is to study the stability of the ascent of an n -tuple $T = (T_1, \dots, T_n)$ of mutually commuting operators, when T is submitted to a perturbation by an n -tuple $S = (S_1, \dots, S_n)$ which commutes mutually with T (see Theorem 4.2). In order to make the problem precise, we introduce a new class of n -tuples, denoted by $\mathcal{K}_n^e(X)$, which is equal, for $n = 1$, to the set of essentially Kato operators. In [9, Proposition 2.6], T. West proves that if T is upper semi-Fredholm operator, then

$$a(T) < \infty \iff \overline{N^\infty(T)} \cap R^\infty(T) = \{0\}. \quad (1.1)$$

In the present paper, we prove that the result (1.1) remains true for an n -tuple which belong to a class denoted by $\mathcal{C}_n(X)$ (see Corollary 2.1). In Theorem 3.1, we prove that, for all $T \in \mathcal{K}_n^e(X)$, there exists $\varepsilon > 0$ such that, for all S be in $\mathcal{B}_n(X) \cap B(O, \varepsilon)$ and mutually commuting with T , we have

$$R^\infty(T + S) \cap \overline{N^\infty(T + S)} \subset R^\infty(T) \cap \overline{N^\infty(T)}.$$

This result is well known for the case of essentially Kato operators (see [7, section 21, Theorem 14]). Finally, in Theorem 4.2, we study the stability of the ascent of a mutually commuting

n -tuple T subjected to a perturbation by an n -tuple S which commutes mutually with T . More precisely, we prove the following:

$$T \in \mathcal{G}_{n\alpha}(X), \ a(T) < +\infty \text{ and } S \in \mathcal{P}(\mathcal{G}_{n\alpha}(X)), \ \alpha > 0 \implies a(T + S) < +\infty.$$

This stability result, generalize the following well known Rakočević theorem:

For T, K be two bounded linear operators such that $TK = KT$. We have :

$$T \in \Phi_+(X), \ a(T) < \infty \text{ and } K \in \mathcal{P}(\Phi_+(X)) \implies a(T + K) < \infty.$$

Let us introduce some notations. For X and Y be two Banach spaces we denote by $\mathcal{B}(X, Y)$ the set of all bounded linear operators from X to Y . We write for short $\mathcal{B}(X) = \mathcal{B}(X, X)$. By X^* we denote the dual of X . Let M be a subset of a Banach space X . The annihilator of M is the closed subspace of X defined by

$$M^\perp := \{f \in X^*; \ f(x) = 0, \ \forall x \in M\},$$

while the pre-annihilator of a subset W of X^* is the closed subspace of X defined by

$${}^\perp W := \{x \in X; \ f(x) = 0, \ \forall f \in W\}.$$

In the following lemma, we recall some well known results.

Lemma 1.1 [7] *Let X be a Banach space.*

(i) *If M is a subspace of X , then ${}^\perp(M^\perp) = \overline{M}$.*

(ii) *If $\{M_\alpha\}_\alpha$ is any family of subsets of X , then $(\bigcup_\alpha M_\alpha)^\perp = \bigcap_\alpha M_\alpha^\perp$.*

If $\{L_\alpha\}_\alpha$ is any family of subsets of X^ , then ${}^\perp(\bigcup_\alpha L_\alpha) = \bigcap_\alpha {}^\perp L_\alpha$.*

(iii) *Let $n \in \mathbb{N} \setminus \{0, 1\}$ and M_1, \dots, M_n be closed subspaces of X .*

If, $\forall j \in \{1, \dots, (n-1)\}$, $\sum_{i=1}^j M_i$ is closed, then $(\sum_{i=1}^n M_i)^\perp = \bigcap_{i=1}^n M_i^\perp$. ◇

For an operator $T \in \mathcal{B}(X)$ define T^* the dual operator of T . We write $N(T)$ for the null space and $R(T)$ for the range of T . The generalized range and generalized kernel of T are defined respectively by $R^\infty(T) = \bigcap_{k=0}^{+\infty} R(T^k)$ and $N^\infty(T) = \bigcup_{k=0}^{+\infty} N(T^k)$. T is said to be essentially Kato if $R(T)$ is closed and there exists a finite-dimensional subspace $F \subset X$ such that

$$N^\infty(T) \subset R^\infty(T) + F.$$

The set of upper semi-Fredholm operators is defined by

$$\Phi_+(X) = \{T \in \mathcal{B}(X); \ \dim N(T) < \infty \text{ and } R(T) \text{ is closed in } X\}.$$

The perturbation class associated with $\Phi_+(X)$ is defined by

$$\mathcal{P}(\Phi_+(X)) := \{T \in \mathcal{B}(X); T + S \in \Phi_+(X), \forall S \in \Phi_+(X)\}.$$

Let $T = (T_1, \dots, T_n)$ be n -tuple. T is said to be of mutually commuting operators on a Banach space X if $\forall (i, j) \in \{1, \dots, n\}^2$, $T_i T_j = T_j T_i$. We denote $\mathcal{B}_n(X)$ the set of n -tuples of mutually commuting operators on a Banach space X . T is said to be of mutually commuting with an n -tuple $S = (S_1, \dots, S_n)$ if $\forall (i, j) \in \{1, \dots, n\}^2$, $T_i S_j = S_j T_i$. Define $T^* = (T_1^*, \dots, T_n^*)$ the dual of T . We use the standard multiindex notation. We denote for all $k \in \mathbb{N}$:

$$T^k = (T_1^k, \dots, T_n^k), N(T^k) = \bigcap_{i=1}^n N(T_i^k) \text{ and } R(T^k) = \sum_{i=1}^n R(T_i^k).$$

The generalized range and generalized kernel of the n -tuple T are defined respectively by $R^\infty(T) = \bigcap_{k=0}^{+\infty} R(T^k)$ and $N^\infty(T) = \bigcup_{k=0}^{+\infty} N(T^k)$. The n -tuple T is said to be essentially Kato if $R(T)$ is closed and there exists a finite-dimensional subspace $F \subset N^\infty(T)$ such that

$$N^\infty(T) \subset R^\infty(T) + F.$$

We write $\delta_T : X \rightarrow X^n$ the bounded operator defined by $\delta_T(x) = (T_1 x, \dots, T_n x)$ ($x \in X$). The set of upper semi-Fredholm and lower semi-Fredholm n -tuples of commuting operators on X are defined respectively by

$$\begin{aligned} \Phi_+^n(X) &= \{T = (T_1, \dots, T_n) \in \mathcal{B}_n(X); \dim N(T) < \infty \text{ and } R(\delta_T) \text{ is closed}\} \\ \Phi_-^n(X) &= \{T = (T_1, \dots, T_n) \in \mathcal{B}_n(X); \operatorname{codim} R(T) < \infty\}. \end{aligned}$$

For T be in $\mathcal{B}_n(X)$, the ascent of T is defined by

$$a(T) = \min\{k; N(T^k) = N(T^{k+1})\}.$$

If no such k exists, then we set $a(T) = \infty$.

The plan of this paper is as follows. In Section 2, we exhibit a large class of n -tuples in which the formula in [9, Proposition 2.6] remains valid. In Section 3, we give quantitative stability result for generalized range and generalized kernel of n tuple in the class $\mathcal{K}_n^e(X)$. The main result of this section is Theorem 3.1. In Section 4, we study the stability of the ascent of an n -tuple (see Theorem 4.2).

2 Class $\mathcal{C}_n(X)$ of n -tuples

The ultimate of this section is to extend Proposition 2.6 in [9] to a class of n -tuples of commuting operators.

Proposition 2.1 *Let $T = (T_1, \dots, T_n)$ be in $\mathcal{B}_n(X)$. We assume that the following conditions hold:*

(H₁) $\dim N(T)$ is finite.

(H₂) There exists $m \in \mathbb{N}^$ such that, $\forall k \geq m, \forall (i, j) \in \{1, \dots, n\}^2, T_i^k(N(T_i^{k+1})) \subset N(T_j)$.*

Then

$$\overline{N^\infty(T)} \cap R^\infty(T) = \{0\} \text{ implies that } T \text{ has a finite ascent.} \quad \diamond$$

Proof. For $n \in \mathbb{N}$ we denote by $V_n := N(T) \cap R(T^n)$. Since $(V_n)_n$ is a decreasing serie of vector subspaces of finite dimensions, then there exists $m \in \mathbb{N}$ such that $V_k = V_m, \forall k \geq m$. Let $x \in N(T^{m+1})$. Using hypothesis (H₂), we obtain $T_i^m(x) \in N(T_j), \forall (i, j) \in \{1, \dots, n\}^2$ and then

$$T_i^m(x) \in \bigcap_{j=1}^n N(T_j) = N(T). \text{ Hence,}$$

$$\forall i \in \{1, \dots, n\}, T_i^m(x) \in N(T) \cap R(T^m) = V_m \subset \overline{N^\infty(T)} \cap R^\infty(T) = \{0\}.$$

Thus, $x \in N(T^m)$. Finally, $N(T^m) = N(T^{m+1})$ and therefore $a(T)$ is finite. Q.E.D.

Examples 2.1 *For $i \in \mathbb{N}^*$, we consider the unilateral backward weighted shift operator T_i defined on $l^r(\mathbb{N}, \mathbb{C}), r \geq 1$, by:*

$$T_i((x_n)_n) = (w_n x_n)_n \text{ with } w_n = 0 \iff n = i - 1.$$

For all $i \in \{1, \dots, n\}$, we have

$$N(T_i^{k+1}) = \{0\}^i \times \mathbb{C} \times \{0\}^{\mathbb{N}}.$$

Let $(x_n)_{n \in \mathbb{N}} \in N(T_i^{k+1})$, then $(x_n)_{n \in \mathbb{N}} = (0, \dots, 0, x_i, 0, \dots)$. Thus,

$$T_i^k(N(T_i^{k+1})) = \{0\} \subset N(T_j), \forall j \in \{1, \dots, n\}.$$

Moreover, we have $N^\infty(T) = \{0\}$. Hence, according to Proposition 2.1, $a(T)$ is finite.

Remark 2.1 *Let $T = (T_1, \dots, T_n) \in \mathcal{B}_n(X)$. We suppose that, $\forall i \in \{1, \dots, n\}, a(T_i)$ is finite. We denote by $p = \max\{a(T_i), 1 \leq i \leq n\}$. Then we obtain*

$$N(T^{p+1}) = \bigcap_{i=1}^n N(T_i^{p+1}) = \bigcap_{i=1}^n N(T_i^p) = N(T^p)$$

and we conclude that $a(T)$ is finite. The converse is false, indeed, if we consider $T = (T_1, T_2)$ such that T_1 is invertible and $a(T_2)$ is infinite, then we obtain,

$$\forall k \in \mathbb{N}, N(T^k) = N(T_1^k) \cap N(T_2^k) = \{0\} = N(T^{k+1}).$$

Thus, $a(T)$ is finite.

In the following result, we study the converse of Proposition 2.1.

Proposition 2.2 *Let $T = (T_1, \dots, T_n)$ be in $\mathcal{B}_n(X)$. We assume that the following conditions hold.*

(H₃) *There exists $A = (A_1, \dots, A_n) \in \mathcal{B}_n(X)$ such that A is mutually commuting with T satisfying $N(A) = \{0\}$.*

(H₄) *There exists $m \in \mathbb{N}$ such that, $\forall p \geq m$, $\delta_A(N(T^p) \cap R(T^p)) \subset R(\delta_{T^p})$, where*

$$\begin{aligned} \delta_A : X &\longrightarrow X^n \\ x &\longmapsto (A_1x, \dots, A_nx) \end{aligned}$$

Then

$$T \text{ has finite ascent implies that } \overline{N^\infty(T)} \cap R^\infty(T) = \{0\}.$$

◇

Proof. Let $p \geq \sup(a(T), m)$ and $V_p := N(T^p) \cap R(T^p)$. Let $x \in V_p$, then, by hypothesis (H₄), $\delta_A(x) \in R(\delta_{T^p})$. Then there exists $z \in X$ such that $A_j(x) = T_j^p(z)$, $\forall j \in \{1, \dots, n\}$. Since, $\forall i \in \{1, \dots, n\}$, $A_i T_i = T_i A_i$, then we get $A_j T_j^p(x) = T_j^{2p}(z)$, $\forall j \in \{1, \dots, n\}$. Moreover, we have $x \in N(T^p)$, then $z \in N(T^{2p}) = N(T^p)$. Hence,

$$0 = T_j^p(z) = A_j(x), \quad \forall j \in \{1, \dots, n\}.$$

Thus, $x \in N(A) = \{0\}$ and therefore $V_p = \{0\}$. On the other hand, the fact that $R^\infty(T) \subset R(T^p)$ and $\overline{N^\infty(T)} \subset N(T^p)$, we obtain

$$\overline{N^\infty(T)} \cap R^\infty(T) \subset V_p = \{0\}.$$

Finally, $\overline{N^\infty(T)} \cap R^\infty(T) = \{0\}$.

Q.E.D.

Definition 2.1 *Let $T = (T_1, \dots, T_n)$ be in $\mathcal{B}_n(X)$. T is said to be in $\mathcal{C}_n(X)$ if it satisfies the following hypotheses (\mathcal{H}):*

$$(\mathcal{H}) \left\{ \begin{array}{l} (i) \ T \in \Phi_+^n(X); \\ (ii) \ \text{There exists } k_0 \in \mathbb{N} \text{ such that, } \forall k \geq k_0, \forall (i, j) \in \{1, \dots, n\}^2, i \neq j, \\ \qquad \qquad \qquad T_i^k(N(T_i^{k+1})) \subset N(T_j); \\ (iii) \ \text{There exists } m \in \mathbb{N} \text{ such that, } \forall p \geq m, \text{ there exists } A = (A_1, \dots, A_n) \in \mathcal{B}_n(X) \\ \qquad \qquad \qquad \text{mutually commuting with } T \text{ satisfying:} \\ \qquad \qquad \qquad N(A) = \{0\} \text{ and } \delta_A(N(T^p) \cap R(T^p)) \subset R(\delta_{T^p}). \end{array} \right.$$

Remark 2.2 Notice that for $n = 1$, $T \in \mathcal{C}_1(X)$, if and only if, $T \in \Phi_+(X)$. Indeed, if $T \in \mathcal{C}_1(X)$ then $T \in \Phi_+(X)$. Conversely, if $T \in \Phi_+(X)$, then the hypothesis (\mathcal{H}) is satisfied if we take $A = id_X$.

The following example proves that $\mathcal{C}_n(X)$ is not empty.

Examples 2.2 Let $T = (T_1, \dots, T_n)$ be in $\mathcal{B}_n(X)$ satisfying the following hypotheses:

- (i) T_1 is invertible in $\mathcal{L}(X)$.
- (ii) For all $i \in \{2, \dots, n\}$, $T_i \in \Phi_+(X)$ and $a(T_i) < +\infty$.

Then $T \in \mathcal{C}_n(X)$. Indeed, let $i \in \{2, \dots, n\}$, since $a(T_i) < +\infty$, then there exists $m \in \mathbb{N}$ such that, $\forall k \geq m$, $N(T_i^k) = N(T_i^m)$. For $k \geq m$, we have

$$T_i^k(N(T_i^{k+1})) = T_i^k(N(T_i^m)) = T_i^{k-m}(T_i^m(N(T_i^m))) = \{0\}.$$

Thus, $(\mathcal{H})(ii)$ is satisfied. Moreover, since T_1 is invertible, then, $\forall p \geq 1$, $N(T^p) = \{0\}$ and therefore $(\mathcal{H})(iii)$ is satisfied.

As a consequence of Propositions 2.1 and 2.2, we get the following result which extends Proposition 2.6 in [9].

Corollary 2.1 Let $T = (T_1, \dots, T_n)$ be in $\mathcal{C}_n(X)$. We have

$$T \text{ has finite ascent} \iff \overline{N^\infty}(T) \cap R^\infty(T) = \{0\}.$$

◇

3 Class $\mathcal{K}_n^e(X)$ of n -tuples

Definition 3.1 Let $\alpha > 0$ and $T = (T_1, \dots, T_n) \in \mathcal{B}_n(X)$. T is said to be essentially kato-stable if there exists $\varepsilon > \alpha$ such that, for all $S = (S_1, \dots, S_n) \in \mathcal{B}_n(X) \cap B(O, \varepsilon)$ and mutually commuting with T , the following assertions holds:

- (i) $T + S$ is essentially kato i.e. there exists a finite dimensional subspace F_S such that

$$N^\infty(T + S) \subset R^\infty(T + S) + F_S.$$

$$(ii) \sum_{i=1}^n (T_i^* + S_i^*) R^\infty(T^*) = R^\infty(T^*).$$

$$(iii) \prod_{i=1}^n (T_i + S_i) \left(R^\infty(T + S) \cap N^\infty(T + S) \right) = R^\infty(T + S) \cap N^\infty(T + S).$$

$$(iv) \forall k \in \mathbb{N}, \forall i \in \{1, \dots, n\}, R(T_i^* + S_i^*)^k \text{ and } R(T_i^{*k}) \text{ are closed in } X^*.$$

$$(v) \forall k \in \mathbb{N}, \forall p \in \{1, \dots, n\}, R((T + S)^k), \sum_{i=1}^p N(T_i^k) \text{ and } \sum_{i=1}^p N((T_i + S_i)^k) \text{ are closed in } X.$$

We denote by $\mathcal{K}_{n\alpha}^e(X)$ the set of all essentially kato-stable n -tuples and we consider

$$\mathcal{K}_n^e(X) := \bigcup_{\alpha>0} \mathcal{K}_{n\alpha}^e(X).$$

Hence, we have $T \in \mathcal{K}^e(X)$, if and only if, there exists $\varepsilon > 0$ such that, $\forall S = (S_1, \dots, S_n) \in \mathcal{B}_n(X) \cap B(O, \varepsilon)$, the assertions (i) - (v) of the previous definition are satisfied.

Remark 3.1 For $n = 1$, we have $T \in \mathcal{K}_1^e(X)$ is equivalent to T is essentially Kato operator. Indeed, it is clear that if $T \in \mathcal{K}_1^e(X)$, then T is essentially Kato. Conversely, if T is essentially Kato, then according [7, Theorems 5, 7 and 8 section 21], $T \in \mathcal{K}_1^e(X)$, when $\alpha < \lim_{n \rightarrow +\infty} (\gamma(T^n))^{\frac{1}{n}}$.

The following proposition proves that $\mathcal{K}_n^e(X)$ is a non-empty set.

Proposition 3.1 Let $T = (T_1, \dots, T_n)$ be in $\mathcal{B}_n(X)$ such that T_1 is invertible in $\mathcal{B}(X)$ and, $\forall i \in \{2, \dots, n\}$, $T_i \in \phi_+(X)$. Then

$$T \in \mathcal{K}_n^e(X).$$

◇

Proof.

Since T_1 is invertible, then there exists ε_1 such that, $\forall S_1 \in B(O, \varepsilon_1)$, $T_1 + S_1$ is invertible in $\mathcal{B}(X)$. Let $S = (S_1, \dots, S_n)$ be such that $\|S\| = \sum_{i=1}^n \|S_i\| < \varepsilon_1$. Then $\|S_1\| < \varepsilon_1$. Hence,

$$N^\infty(T + S) \subset N^\infty(T_1 + S_1) \subset \{0\}$$

and therefore the assertions (i) and (iii) of Definition 3.1 are satisfied. Moreover, since T_1^* is invertible, then $R^\infty(T^*) = X^*$ and $(T_i^* + S_i^*)(R^\infty(T^*)) = X^*$. So the assertion (ii) of Definition 3.1 is satisfied. Finally, since, $\forall i \in \{1, \dots, n\}$, $T_i \in \phi_+(X)$, then there exists $\varepsilon \leq \varepsilon_1$ such that,

for all $S_i \in B(O, \varepsilon)$, $(T_i + S_i) \in \phi_+(X)$. So, for all $k \in \mathbb{N}$ and all $j \in \{1, \dots, n\}$, $\sum_{i=1}^j N(T_i + S_i)^k$

and $R(T^* + S^*)^k = X^*$ are subsets of finite dimensional. Thus, (iv) and (v) of Definition 3.1 are satisfied. Q.E.D.

In the following Proposition and according to Definition 3.1, we deduce some useful properties in the set $\mathcal{K}_{n\alpha}^e(X)$.

Proposition 3.2 (i) If $0 < \alpha \leq \beta$, then $\mathcal{K}_{n\beta}^e(X) \subset \mathcal{K}_{n\alpha}^e(X)$.

(ii) For $\alpha > 0$ and $\beta > 0$, we have $\mathcal{K}_{n\alpha}^e(X) \cap \mathcal{K}_{n\beta}^e(X) = \mathcal{K}_{n\max(\alpha, \beta)}^e(X)$.

(iii) If T is in $\mathcal{K}_{n\alpha}^e(X)$, then $\forall \lambda \in \mathbb{C}^*$ such that $|\lambda| > 1$, $\lambda T \in \mathcal{K}_{n\alpha}^e(X)$.

◇

Proof. (i) Let $T \in \mathcal{K}_{n\beta}^e(X)$, then there exists $\varepsilon > \beta$ and hence $\varepsilon > \alpha$ such that, $\forall S \in \mathcal{B}_n(X) \cap B(O, \varepsilon)$ and mutually commuting with T , the assertions (i) - (v) of Definition 3.1 hold.

(ii) If $\varepsilon > \alpha$ and $\varepsilon > \beta$, then $\varepsilon > \max(\alpha, \beta)$.

(iii) Let $\varepsilon > 0$ such that $T \in \mathcal{K}_{n\alpha}^e(X)$. Then for $|\lambda| > 1$, we have $\frac{1}{\lambda}S \in \mathcal{B}_n(X) \cap \mathcal{B}(O, \varepsilon)$ and hence (i) – (v) of Definition 3.1 are satisfied. Thus, $\lambda T \in \mathcal{K}_{n\alpha}^e(X)$. Q.E.D.

The main result of this section is the following.

Theorem 3.1 *Let $T = (T_1, \dots, T_n)$ be in $\mathcal{B}_n(X)$. Suppose that $T \in \mathcal{K}_n^e(X)$, then there exists $\varepsilon > 0$ such that, for all $S \in \mathcal{B}_n(X) \cap \mathcal{B}(O, \varepsilon)$ and mutually commuting with T , we have*

$$R^\infty(T + S) \cap \overline{N^\infty(T + S)} \subset R^\infty(T) \cap \overline{N^\infty(T)}. \quad \diamond$$

To prove this Theorem we shall need two Lemmas.

Lemma 3.1 *Let $T = (T_1, \dots, T_n)$ and $S = (S_1, \dots, S_n)$ be in $\mathcal{B}_n(X)$ mutually commuting with T . We assume that the following conditions hold:*

$$(A_1) \sum_{i=1}^n (T_i^* + S_i^*) R^\infty(T^*) = R^\infty(T^*).$$

$$(A_2) \sum_{i=1}^p N(T_i^k) \text{ and } \sum_{i=1}^p N((T_i + S_i)^k) \text{ are closed in } X, \forall (p, k) \in \{1, \dots, n\} \times \mathbb{N}.$$

$$(A_3) R(T_i^{*k}) \text{ and } R((T_i^* + S_i^*)^k) \text{ are closed in } X^*, \forall (i, k) \in \{1, \dots, n\} \times \mathbb{N}.$$

Then

$$\overline{N^\infty(T + S)} \subset \overline{N^\infty(T)}. \quad \diamond$$

Proof.

- We claim that

$$\overline{N^\infty(T + S)} = {}^\perp \left(R^\infty(T^* + S^*) \right). \quad (3.1)$$

By Lemma 1.1, we obtain:

$$\begin{aligned} \overline{N^\infty(T + S)} &= {}^\perp \left[\left(\bigcup_{k=0}^{+\infty} \left(\bigcap_{i=1}^n N((T_i + S_i)^k) \right) \right)^\perp \right] = {}^\perp \left[\bigcap_{k=0}^{+\infty} \left(\bigcap_{i=1}^n N((T_i + S_i)^k) \right)^\perp \right] \\ &= {}^\perp \left[\bigcap_{k=0}^{+\infty} \left(\bigcap_{i=1}^n {}^\perp R((T_i^* + S_i^*)^k) \right)^\perp \right]. \end{aligned}$$

The fact that $R((T_i^* + S_i^*)^k)$ is closed $\forall (i, k) \in \{1, \dots, n\} \times \mathbb{N}$ and by Lemma 1.1 we infer that:

$$\begin{aligned} \overline{N^\infty(T + S)} &= {}^\perp \left[\bigcap_{k=0}^{+\infty} \left(\sum_{i=1}^n {}^\perp (R((T_i^* + S_i^*)^k))^\perp \right) \right] = {}^\perp \left[\bigcap_{k=0}^{+\infty} \left(\sum_{i=1}^n \overline{R((T_i^* + S_i^*)^k)} \right) \right] \\ &= {}^\perp \left[\bigcap_{k=0}^{+\infty} \left(\sum_{i=1}^n R((T_i^* + S_i^*)^k) \right) \right]. \end{aligned}$$

Hence, $\overline{N^\infty(T+S)} = {}^\perp(R^\infty(T^*+S^*))$ and our claim is proved.

- We claim that

$$R^\infty(T^*) \subset R^\infty(T^*+S^*). \quad (3.2)$$

Let $x \in R^\infty(T^*)$. Using (A_1) , there exist $x_1, \dots, x_n \in R^\infty(T^*)$ such that

$$x = \sum_{i=1}^n (T_i^* + S_i^*)(x_i).$$

By induction on $p \in \mathbb{N}^*$, we will have

$$x = \sum_{k_1+\dots+k_n=np} (T_1^* + S_1^*)^{k_1} \dots (T_n^* + S_n^*)^{k_n} (z_{k_1, \dots, k_n}),$$

where $z_{k_1, \dots, k_n} \in R^\infty(T^*)$. Since $k_1 + \dots + k_n = np$, then there exists $i \in \{1, \dots, n\}$ such that $k_i \geq p$ and hence

$$(T_1^* + S_1^*)^{k_1} \dots (T_n^* + S_n^*)^{k_n} (z_{k_1, \dots, k_n}) \in (T_i^* + S_i^*)^{k_i} (R^\infty(T^*)) \subset R((T_i^* + S_i^*)^p).$$

Thus, $x \in \sum_{i=1}^n R((T_i^* + S_i^*)^p)$. Hence,

$$x \in \bigcap_{p=0}^{+\infty} \left(\sum_{i=1}^n R((T_i^* + S_i^*)^p) \right) = R^\infty(T^* + S^*),$$

which proves our claim.

Now, using Equation (3.2), we obtain ${}^\perp R^\infty(T^* + S^*) \subset {}^\perp R^\infty(T^*)$. Then Equation (3.1) implies that

$$\overline{N^\infty(T+S)} \subset \overline{N^\infty(T)}. \quad \text{Q.E.D.}$$

Lemma 3.2 *Let $T = (T_1, \dots, T_n)$ be in $\mathcal{B}_n(X)$. We assume that, there exists $\varepsilon > 0$ such that, for all $S = (S_1, \dots, S_n) \in \mathcal{B}_n(X) \cap B(O, \varepsilon)$ and mutually commuting with T , the following conditions hold:*

(A_4) $T + S$ is essentially Kato and $R((T + S)^k)$ is closed in X , $\forall k \in \mathbb{N}$.

(A_5) $g_{T,S}(R^\infty(T + S) \cap N^\infty(T + S)) = R^\infty(T + S) \cap N^\infty(T + S)$, where $g_{T,S}$ is the operator defined by:

$$g_{T,S} = \prod_{i=1}^n (T_i + S_i).$$

Then

$$R^\infty(T + S) \cap \overline{N^\infty(T + S)} \subset R^\infty(T). \quad \diamond$$

Proof. Since $T + S$ is essentially Kato, then there exists a finite dimensional subspace F such that $F \cap R^\infty(T + S) = \{0\}$ and

$$\overline{N^\infty(T + S) \cap R^\infty(T + S)} = \overline{(N^\infty(T + S) \cap R^\infty(T + S) + F) \cap R^\infty(T + S)}.$$

Since, $R^\infty(T + S)$ is a closed subspace, then

$$(\overline{N^\infty(T + S) \cap R^\infty(T + S)} + F_1) \cap R^\infty(T + S) = \overline{N^\infty(T + S) \cap R^\infty(T + S)}.$$

Now, to prove Lemma 3.2, it is sufficient to show that, for $k \in \mathbb{N}$,

$$R^\infty(T + S) \bigcap N(T + S)^k \subset R^\infty(T).$$

We will do this by induction on k . The statement is clear for $k = 0$. Let $k \in \mathbb{N}^*$ and assume that the inclusion holds for $k - 1$. Let $x_0 \in R^\infty(T + S) \bigcap N(T + S)^k$. Then we have $x_0 \in R^\infty(T + S) \bigcap N^\infty(T + S)$. Using the hypothesis (A_5) we can find an infinite sequence x_0, x_1, \dots in $R^\infty(T + S) \bigcap N^\infty(T + S)$ such that $x_{j-1} = g_{T,S}(x_j)$. By Lemma 3.1, $(x_j)_{j \in \mathbb{N}} \in \overline{N^\infty(T)}$. Since T is essentially Kato, then $\dim(\overline{N^\infty(T)}/R^\infty(T) \bigcap \overline{N(T)}) = m$ is finite. Thus x_0, \dots, x_m are linearly dependent, i.e., there exists a non-trivial combination

$$x := \sum_{i=0}^m \alpha_i x_i \in R^\infty(T).$$

let l be such that $\alpha_l \neq 0$ and $\alpha_j = 0$ for $j = l + 1, \dots, m$. We obtain:

$$g_{T,S}^l x = \alpha_l x_0 + \sum_{j=0}^{l-1} \alpha_j g_{T,S}^l x_j \in \alpha_l x_0 + (R^\infty(T + S) \bigcap N(T + S)^{k-1}).$$

Finally, we obtain $x_0 \in R^\infty(T)$.

Q.E.D.

Proof of Theorem 3.1

By Lemma 3.1 we have $\overline{N^\infty(T + S)} \subset \overline{N^\infty(T)}$, then, the use of Lemma 3.2 leads to

$$R^\infty(T + S) \cap \overline{N^\infty(T + S)} \subset R^\infty(T)$$

and the theorem is proved.

Q.E.D.

4 Stability of the ascent of n -tuple

In the following we consider some classes of n -tuples commuting operators.

(i) For $\alpha > 0$, consider $\mathcal{G}_{n\alpha}(X) := \mathcal{K}_{n\alpha}^e(X) \cap \mathcal{C}_n(X)$.

(ii) Let $\mathcal{G}_n(X) := \bigcup_{\alpha > 0} \mathcal{G}_{n\alpha}(X)$.

(iii) For $\alpha > 0$, let $\mathcal{P}(\mathcal{K}_{n\alpha}^e(X))$ the set of all n -tuples $T = (T_1, \dots, T_n) \in \mathcal{B}_n(X)$ such that, for all $S \in \mathcal{K}_{n\alpha}^e(X)$ and mutually commuting with T , $T + S \in \mathcal{K}_{n\alpha}^e(X)$.

(iv) $\mathcal{P}(\mathcal{G}_{n\alpha}(X))$ the set of all n -tuple T such that, for all $S \in \mathcal{G}_{n\alpha}(X)$ and mutually commuting with T , $T + S \in \mathcal{G}_{n\alpha}(X)$.

(v) Let $\mathcal{P}(\mathcal{G}_n(X)) := \bigcap_{\alpha > 0} \mathcal{P}(\mathcal{G}_{n\alpha}(X))$.

Remark 4.1 (i) It is clear that, for $n = 1$, $\mathcal{G}_n(X) = \Phi_+(X)$.

(ii) According to Proposition 3.1, the set of n -tuples defined in example 2.2 is included in $\mathcal{G}_n(X)$.

(iii) Notice that for $n = 1$ and according to [7, Theorem 9, Section 21] $\mathcal{P}(\mathcal{K}_{n\alpha}^e(X))$ contains the set of all compact and quasinilpotent operators.

Proposition 4.1 (i) If $0 < \alpha \leq \beta$, then $\mathcal{P}(\mathcal{K}_{n\alpha}^e(X)) \subset \mathcal{P}(\mathcal{K}_{n\beta}^e(X))$.

(ii) If $T \in \mathcal{P}(\mathcal{K}_{n\alpha}^e(X))$, then for all $\lambda \in]0, 1]$, $\lambda T \in \mathcal{P}(\mathcal{K}_{n\alpha}^e(X))$. ◇

Proof. Follows from Proposition 2.2 (i) and (iii). Q.E.D.

Theorem 4.1 Let $T = (T_1, \dots, T_n)$ be in $\mathcal{K}_{n\alpha}^e(X)$ and let S be in $\mathcal{P}(\mathcal{K}_{n\alpha}^e(X))$ and mutually commuting with T . Then

$$R^\infty(T + S) \cap \overline{N^\infty(T + S)} \subset R^\infty(T) \cap \overline{N^\infty(T)}. \quad \diamond$$

Proof. Let $\mu \in [0, 1]$. By Proposition 4.1, $T_\mu = T + \mu S \in \mathcal{K}_n^e(X)$. According to Theorem 3.1 there exists $\varepsilon > 0$ such that, $\forall \lambda \in]\mu, \mu + \frac{\varepsilon}{\|S\|}[$, we have

$$R^\infty(T_\lambda) \cap \overline{N^\infty(T_\lambda)} \subset R^\infty(T_\mu) \cap \overline{N^\infty(T_\mu)}.$$

If we apply the above procedure with $\mu_0 = 0 < \mu_1 < \dots < \mu_p = 1$, then, for all $i \in \{0, \dots, p-1\}$, there exists $\varepsilon_i > 0$ such that $\mu_{i+1} \in]\mu_i, \mu_i + \frac{\varepsilon_i}{\|S\|}[$ and we have

$$R^\infty(T_{\mu_{i+1}}) \cap \overline{N^\infty(T_{\mu_{i+1}})} \subset R^\infty(T_{\mu_i}) \cap \overline{N^\infty(T_{\mu_i})}.$$

Thus,

$$R^\infty(T_{\mu_p}) \cap \overline{N^\infty(T_{\mu_p})} \subset R^\infty(T_{\mu_{p-1}}) \cap \overline{N^\infty(T_{\mu_{p-1}})} \subset \dots \subset R^\infty(T_{\mu_0}) \cap \overline{N^\infty(T_{\mu_0})},$$

which implies that

$$R^\infty(T + S) \cap \overline{N^\infty(T + S)} \subset R^\infty(T) \cap \overline{N^\infty(T)}. \quad \text{Q.E.D.}$$

Now, we are ready to state the main result of this paper.

Theorem 4.2 Let $T = (T_1, \dots, T_n) \in \mathcal{B}_n(X)$. Then

$$T \in \mathcal{G}_{n\alpha}(X), \ a(T) < +\infty \text{ and } S \in \mathcal{P}(\mathcal{G}_{n\alpha}(X)), \ \alpha > 0 \implies a(T + S) < +\infty. \quad \diamond$$

Proof. Applying Theorem 4.1 , we have

$$R^\infty(T + S) \cap \overline{N^\infty(T + S)} \subset R^\infty(T) \cap \overline{N^\infty(T)}.$$

According to Corollary 2.1 we infer that

$$R^\infty(T + S) \cap \overline{N^\infty(T + S)} = \{0\}.$$

Thus, by the same corollary we get $a(T + S) < +\infty$.

Q.E.D.

Corollary 4.1 *Let $T = (T_1, \dots, T_n) \in \mathcal{B}_n(X)$. Then*

$$T \in \mathcal{G}_n(X), \ a(T) < +\infty \text{ and } S \in \mathcal{P}(\mathcal{G}_n(X)) \implies a(T + S) < +\infty.$$

◇

Proof. The result follows since $\mathcal{G}_n(X) = \bigcup_{\alpha > 0} \mathcal{G}_{n\alpha}(X)$.

Q.E.D.

Remark 4.2 *Observe that for $n = 1$, we obtain the well known Theorem 1 established by V. Rakocevic in [7] indeed we have $\mathcal{G}_1(X) = \Phi_+(X)$.*

References

- [1] B. Abdelmoumen and H. Baklouti, *Fredholm Perturbations and Seminorms Related to Upper Semi-Fredholm Perturbations*, Filomat 27:6 (2013), 11471155 DOI DOI: 10.2298/FIL1306147A
- [2] B. Abdelmoumen, H. Baklouti, *Perturbation results on semi-Fredholm operators set and applications*, J. Ineq. Appl., (2009) , Article ID 284526, 13 pages doi:10.1155/2009/284526.
- [3] M. Burgos, A. Kaidi, M. Mbekhta, and M. Oudghiri, *The Descent Spectrum and Perturbations*, J. Oper. Theo. 56:2 (2006), 259-271.
- [4] R. Gelca, *Compact perturbations of Fredholm n -tuples*, Proc. Amer. Math. Soci, Volume 122, Number 1. September 1994.
- [5] M. Gonzalez, *The perturbation class problem in Fredholm theory*, Jour. Func. Anal, 200 (2003) 65-70.
- [6] T. Kato, *Perturbation theory for nullity, deficiency and other quantities of linear operators*, J. Anal. Math. 6 (1958) 261-322.
- [7] V. Müller, *Spectral theory of linear operator and spectral system in Banach algebras*, Operator theory advance and application vol. 139, (2003).
- [8] V. Rakoćević, *Semi-Fredholm operators with finite ascent or descent and perturbations*, Proc. Amer. Math. Soc. Volume 123, Number 12, December 1995.
- [9] T. T. West, *A Riesz-Schauder theorem for semi-Fredholm operators*, Proc. Roy. Irish Acad. Sect. A 87 (1987), 137-146.