

Grouped GEE Analysis for Longitudinal Data

Tsubasa Ito¹ and Shonosuke Sugasawa²

¹M&D Data Science Center, Tokyo Medical and Dental University

²Center for Spatial Information Science, The University of Tokyo

Abstract

Generalized estimating equation (GEE) is widely adopted for regression modeling for longitudinal data, taking account of potential correlations within the same subjects. Although the standard GEE assumes common regression coefficients among all the subjects, such assumption is not realistic when there are potential heterogeneity in regression coefficients among subjects. In this paper, we develop a flexible and interpretable approach, called grouped GEE analysis, to modeling longitudinal data with allowing heterogeneity in regression coefficients. The proposed method assumes that the subjects are divided into a finite number of groups and subjects within the same group share the same regression coefficient. We provide a simple algorithm for grouping subjects and estimating the regression coefficients simultaneously, and show asymptotic properties of the proposed estimator. The number of groups can be determined by the cross validation with averaging method. We demonstrate the proposed method through simulation studies and an application to a real dataset.

Key words: Estimating equation; Grouping; Information criterion; k -means algorithm; Unobserved heterogeneity

1 Introduction

Longitudinal data in which responses (repeated measurements) within the same subject are correlated is appeared in many scientific applications such as biomedical statistics and social science. For analyzing longitudinal data, it is typically difficult to correctly specify the underlying correlation structures among response variables within the same subject, and one of the standard approaches is the generalized estimating equations (GEE) developed by Liang and Zeger (1986), which uses “working” correlation structures specified by users. The advantage of the GEE approach is that the estimator is still consistent even when the working correlation is misspecified. However, the existing GEE methods assumes homogeneous regression coefficients that are common to all the subject, which could be restrictive in practical applications since there might be potential heterogeneity among subjects or clusters as confirmed in several applications (e.g. Barban and Billari, 2012; Lin and Ng, 2012; Nagin et al., 2018).

In this work, we extend the standard GEE analysis to take account of potential heterogeneity in longitudinal data. Specifically, we develop grouped GEE analysis by adopting the grouping approach that is widely adopted in literatures for panel data analysis (e.g. Bonhomme and Manresa, 2015; Zhang et al., 2019; Liu et al., 2020). We assume that subjects in longitudinal data can be classified to a finite number of groups, and subjects within the same group share the same regression coefficients, that is, the regression coefficients are homogeneous over subjects in the same groups. Since the grouping assignment of subjects are unknown, we treat it as unknown parameters and estimate them and group-wise regression coefficient simultaneously. Given the grouping parameters, the standard GEE can be performed to obtain group-wise estimators of regression coefficients. On the other hand, given the group-wise regression coefficients, we consider estimating the grouping parameters using a kind of Mahalanobis distance between response variables and predictors with taking account of potential correlations via working correlation matrix. In other words, we employ the working correlation not only in performing GEE analysis in each group but also

estimating the grouping assignment. We will show that the grouped GEE method can be easily carried out by a simple iterative algorithm similar to *k-means* algorithm that combines the existing algorithm for the standard GEE and simple optimization steps for grouping assignment. Moreover, we adopt the cross validation with averaging method proposed in Wang (2010) to carry out data-dependent selection of the number of groups.

We derive the statistical properties of the grouped GEE estimator in an asymptotic framework where both n (the number of subjects) and T (the number of repeated measurements) tend to infinity, but we here allow T to grow considerably slower than n , namely, $n/T^\nu \rightarrow 0$ for some large ν . Hence, our method can be applicable when T is much smaller n as observed in many applications using longitudinal data. As theoretical difficulties of the grouped estimation in longitudinal data analysis, the true correlations within the same subject can be considerably high, so the existing theoretical argument assuming negligibly small correlations imposed typically by mixing conditions (e.g. Bonhomme and Manresa, 2015; Zhang et al., 2019; Gu and Volgushev, 2019) for the underlying true correlations is no more applicable. To overcome the limitation of the existing theoretical argument, we consider grouping assignment using a kind of Mahalanobis distance with working correlation, and we will show that such grouping strategy leads to consistent estimation of the grouping parameters as long as the working correlation is relatively close to the true one. Therefore, even when the underlying correlations within the same subject is not weak, we can successfully estimate the grouping parameters using a reasonable working correlation matrix. Then, we will also establish consistency and asymptotic normality of the grouped GEE estimator of the regression coefficients, and also provide a consistent estimator of asymptotic variances.

In the context of longitudinal data or clustered data analysis, several methods to take account of the potential heterogeneity among subjects have been proposed. Rubin and Wu (1997), Sun et al. (2007), Ng and McLachlan (2014) and Sugawara et al. (2019) proposed a mixture modeling based on random effects, but the estimation algorithms can be computationally very intensive since the algorithms include

iteration steps that entail numerical integration. On the other hand, Rosen et al. (2000) and Tang and Qu (2016) proposed a mixture modeling based on the GEE, but the primary interest in these works is estimation of the component distributions in the mixture rather than grouping subjects. Hajjem et al. (2011) and Fokkema et al. (2018) employed regression tree techniques for grouping observations, but the tree-based methods can handle grouping based on covariate information rather than regression coefficients. Moreover, Coffey et al. (2014), Vogt and Linton (2017) and Zhu and Qu (2018) proposed grouping methods for longitudinal curves, and Tang et al. (2020) developed covariate-specific grouping methods via regularization. To the best of our knowledge, this paper is the first one to consider grouped estimation in the GEE analysis with valid theoretical argument.

This paper is organized as follows. In Section 2, we illustrate the proposed GEE analysis, and provide an iterative estimation algorithm. We also propose the averaging method for selecting the number of groups. In Section 3, we give the asymptotic properties of the grouped GEE estimator. In Section 4, we demonstrate the grouped GEE analysis through simulation studies and an application to a real longitudinal dataset. We give some discussions in Section 5. All the technical details and the proofs of the theorems are given in the Supplementary Materials. R code implementing the proposed method is available at GitHub repository (<https://github.com/sshonosuke/GGEE>).

2 Grouped GEE Analysis

2.1 Grouped models for longitudinal data

For longitudinal data, let y_{it} be the response of interest and x_{it} be a p -dimensional vector of covariate information of subject i at time t , where $i = 1, \dots, n$ and $t = 1, \dots, T_i$. For ease of notation, we set $T_i = T$ for all i , representing balanced data case, but the extension to unbalanced case is straightforward. We consider a generalized

linear model for y_{it} , given by

$$f(y_{it}|x_{it}; \beta_i, \phi) = \exp\left(\{y_{it}\theta_{it} - a(\theta_{it}) + b(y_{it})\}/\phi\right), \quad (1)$$

where $a(\cdot)$ and $b(\cdot)$ are known functions, and $\theta_{it} = u(x_{it}^t/\beta_i)$ for a known monotone function $u(\cdot)$. A commonly used link function is the canonical link function, that is, $u(x) = x$. Here β_i is the regression parameters of interest which can be heterogeneous among subjects, and ϕ is a known scale parameter that is common to all the subjects. Under the model (1), the first two moments of y_{it} are given by $m(x_{it}^t/\beta_i) = a'(\theta_{it})$ and $\sigma^2(x_{it}^t/\beta_i) = a''(\theta_{it})\phi$, respectively. For example, under binary response, it follows that $a(x) = \log\{1 + \exp(x)\}$, leading to the logistic model given by $m(x_{it}^t/\beta_i) = \{1 + \exp(-x_{it}^t/\beta_i)\}^{-1}$.

In the standard GEE analysis, the regression parameters are homogeneous, that is, $\beta_i = \beta$, but we allow potential heterogeneity among the subjects. However, the number of β_i increases as the number of subjects, so β_i cannot be estimated with reasonable accuracy as long as T is not large, which is the standard situation in longitudinal data analysis. Hence, we consider a grouped structure for the subjects, that is, the n subjects are divided into G groups and subjects within the same group share the same regression coefficients. Specifically, we introduce an unknown grouping variable $g_i \in \{1, \dots, G\}$ which determines the group that i th subject belongs to. Then, we define $\beta_i = \beta_{g_i}$ under which the unknown regression parameters are β_1, \dots, β_G . Therefore, if G is not large compared with n and T , then β_1, \dots, β_G can be accurately estimated. Moreover, due to the grouping nature, the estimation results of g_i gives grouping of subjects in terms of regression coefficients, so the estimation result is easily interpretable for users. We treat G as unknown parameter as well, but we assume that G is known for a while. The estimation will be discussed in Section 2.3.

2.2 Estimation algorithm

Define $y_i = (y_{i1}, \dots, y_{iT})^t$ as a $T \times 1$ response vector and $X_i = (x_{i1}, \dots, x_{iT})^t$ as a $T \times p$ covariate matrix. We also define $m(X_i\beta_g) = (m(x_{i1}^t/\beta_g), \dots, m(x_{iT}^t/\beta_g))^t$, $A_i(\beta_g) =$

$\text{diag}(\sigma^2(x_{i1}^t\beta_g), \dots, \sigma^2(x_{iT}^t\beta_g))$, $\Delta_i(\beta_g) = \text{diag}(u'(x_{i1}^t\beta_g), \dots, u'(x_{iT}^t\beta_g))$ and $D_i(\beta_g) = A_i(\beta_g)\Delta_i(\beta_g)X_i$. In what follows, we might abbreviate the explicit dependence on the parameters for notational simplicity when there seems no confusion. We here introduce “working” correlation matrix $R(\alpha)$ to approximate the true underlying correlation matrix of y_i , which one can be chosen freely, where it might include the nuisance unknown parameter α . Then, we define working covariance matrix $V_i(\beta)$ as $V_i(\beta) = A_i^{1/2}(\beta)\widehat{R}A_i^{1/2}(\beta)$ with $\widehat{R} = R(\widehat{\alpha})$. If \widehat{R} is consistent to the true correlation matrix R^0 , $V_i(\beta^0)$ with the true parameter β^0 is also consistent to the true covariance matrix of y_i .

Given the grouping parameter $\bar{g} = (g_1, \dots, g_n)$, we can estimate β_g by performing the standard GEE estimation (Liang and Zeger, 1986) for each group, namely, solving the following estimating equation:

$$S_g(\beta_g) \equiv \sum_{i=1}^n \mathbf{I}(g_i = g) S_i(\beta_g) = 0, \quad S_i(\beta_g) \equiv D_i^t(\beta_g) V_i^{-1}(\beta_g) \{y_i - m(X_i \beta_g)\}, \quad (2)$$

which is the GEE based on the subjects classified to the g th group. We can employ existing numerical algorithm for the standard GEE to obtain the solution of (2). On the other hand, given $\beta = (\beta_1^t, \dots, \beta_G^t)^t$, it is quite reasonable to classify the subjects to groups having the most preferable regression structures to explain the variation of y_i . Thus, we propose estimating the unknown \bar{g} based on the following minimization problem:

$$\widehat{g}_i(\beta) = \underset{g=1, \dots, G}{\text{argmin}} \{y_i - m(X_i \beta_g)\}^t \widehat{R}^{-1}(\beta, \bar{g}) \{y_i - m(X_i \beta_g)\}, \quad (3)$$

where $\widehat{R}(\beta, \bar{g}) = R(\widehat{\alpha}, \beta, \bar{g})$ is an estimator of a working correlation matrix given β and \bar{g} . The objective function in (3) can be seen as a kind of the Mahalanobis distance with taking the working correlation structure into account. Such estimation strategy for the grouping variable has not been paid attention very much, but the use of the working correlation in the grouping step is shown to be quite important to expand our theoretical argument given in Section 3. Note that the above minimization problem

can be carried out separately for each subject, thus (3) can be easily solved by simply evaluating all the values of the objective function over $g \in \{1, \dots, G\}$.

Regarding the estimation of the nuisance parameter α in the working correlation, we suggest using a moment-based method. Given β and \bar{g} , one can estimate α by solving the following minimization problem:

$$\hat{\alpha}(\beta, \bar{g}) = \underset{\alpha}{\operatorname{argmin}} \left\| R(\alpha) - \frac{1}{n} \sum_{i=1}^n A_i^{-1/2} \{y_i - m(X_i \beta_{g_i})\} \{y_i - m(X_i \beta_{g_i})\}^t A_i^{-1/2} \right\|_F, \quad (4)$$

where $\|\cdot\|_F$ is the Frobenius norm. This method can be easily extended to the heterogeneous correlation structures among different groups. Let $\alpha_1, \dots, \alpha_G$ be different correlation parameters. Then, α_g can be estimated by minimizing

$$\left\| R(\alpha_g) - n_g^{-1} \sum_{i=1}^n \mathbf{I}(g_i = g) A_i^{-1/2} \{y_i - m(X_i \beta_{g_i})\} \{y_i - m(X_i \beta_{g_i})\}^t A_i^{-1/2} \right\|_F,$$

where n_g is the number of subjects classified to the g th group.

The estimating equation (2) and two optimization problems (3) and (4) define the grouped GEE estimator of β and \bar{g} , and the estimator can be easily computed by the following iterative algorithm:

Algorithm 1 (grouped GEE estimation).

Starting from some initial values $\beta^{(0)}$, $\bar{g}^{(0)}$ and $\alpha^{(0)}$, we repeat the following procedure until algorithm converges:

- Update $\bar{g}^{(r)}$ to get $\bar{g}^{(r+1)}$ by solving (3) with $\beta = \beta^{(r)}$ and $\alpha = \alpha^{(r)}$.
- Update $\beta^{(r)}$ to get $\beta^{(r+1)}$ by solving (2) with $\bar{g} = \bar{g}^{(r+1)}$ and $\alpha = \alpha^{(r)}$.
- Update $\alpha^{(r)}$ to get $\alpha^{(r+1)}$ by solving (4) with $\beta = \beta^{(r+1)}$ and $\bar{g} = \bar{g}^{(r+1)}$.

Since there might be multiple solutions for the grouped GEE estimator, the above algorithm might be sensitive to the setting of initial values. A reasonable starting value for α would be one that induces independent correlation matrix of R , for example $\alpha = 0$ in the exchangeable working correlation. Regarding β and \bar{g} , we suggest two simple methods to determine their initial values. First method is to apply the

finite mixture models with G components of the form: $y_{it}|(z_{it} = k) \sim h_k(y_{it}; x_{it}^t \beta_k)$ and $P(z_{it} = k) = \pi_k$, for $k = 1, \dots, G$, where h_k is the distribution having mean $m(x_{it}^t \beta)$. Then, we set the initial values of β_k and g_i to the estimates of β_k and the maximizer of $\sum_{t=1}^T P_*(z_{it} = k)$ over $k \in \{1, \dots, G\}$, respectively, where $P_*(z_{it} = k)$ is the conditional probability that y_{it} belongs to the k th group. The second approach is separately fitting the regression model with mean structure $m(x_{it}^t \beta_i)$ for each subject. Based on the estimates $\hat{\beta}_i$ of β_i , we apply the k -means clustering algorithm with G clusters to the n -points $\{\hat{\beta}_1, \dots, \hat{\beta}_n\}$, and set the initial values of β_k and g_i to the center of the resulting clusters and clustering assignment, respectively. Note that the second method is only applicable when T is sufficiently larger than p to get stable estimates of β_i .

2.3 Selecting the number of groups

Since the number of groups is typically unknown in practice, we need to estimate it based on some appropriate criterion. One possible strategy is to adopt a criterion using quasi-likelihood (Wedderburn, 1974) and to use a penalty term in view of Bayesian-type information criterion in GEE analysis (e.g. Wang and Qu, 2009). However, theoretical asymptotic properties of such approaches are not necessarily clear even under the standard GEE settings, so the theoretical investigation would be more complicated under the grouping structure. Instead, we here adopt the cross validation with averaging method (CVA) proposed in Wang (2010), which is shown to have the selection consistency when the clusters are properly separated into subgroups. The same strategy is adopted in Zhang et al. (2019) in the context of quantile regression for panel data.

The CVA criterion is concerned with clustering instability under given G . For $c = 1, \dots, C$, we randomly divide n subjects into three subsets: two training datasets with sizes M and one testing set with size $n - 2M$, where the subject indices included in the three subsets are denoted by Z_1^c, Z_2^c and Z_3^c , respectively, that is, $|Z_1^c| = |Z_2^c| = M$, $|Z_3^c| = n - 2M$, $Z_h^c \cap Z_{h'}^c = \emptyset$ for $h \neq h'$ and $\cup_{h=1,2,3} Z_h^c = \{1, \dots, n\}$. We first apply the proposed grouped GEE method to the two training datasets, which gives

us the estimates of regression coefficients and working correlation matrices. Then, we can compute the optimal grouping assignment in the test data as

$$\widehat{g}_i^{(h)} = \underset{g=1, \dots, G}{\operatorname{argmin}} \{y_i - m(X_i \widehat{\beta}_g^{(h)})\}^t \widehat{R}^{(h)-1} \{y_i - m(X_i \widehat{\beta}_g^{(h)})\}, \quad i \in Z_3^c,$$

where $\widehat{\beta}_g^{(h)}$ and $\widehat{R}^{(h)}$ are estimates of regression coefficients and working correlation based on the h th training data for $h = 1, 2$. Based on the grouping assignment, grouping instability can be quantified as

$$\widehat{s}^c(G) = \sum_{i,j \in Z_3^c} \mathbf{I}\{\widehat{g}_i^{(1)} = \widehat{g}_j^{(1)}\} + \mathbf{I}\{\widehat{g}_i^{(2)} = \widehat{g}_j^{(2)}\} = 1\}.$$

By averaging the above values over $c = 1, \dots, C$, we have $\widehat{s}(G) = C^{-1} \sum_{c=1}^C \widehat{s}^c(G)$, and we select G as the minimizer of the criterion among some candidates of G . Finally, regarding the choice of M , we simply set $M = \lfloor n/3 \rfloor$, so the three subsets have almost the same numbers of subjects.

3 Asymptotic Properties

We here provide the asymptotic properties of the grouped GEE estimators, that is, the grouping parameter \bar{g} can be consistently estimated, and $\widehat{\beta}_g$ admits both consistency and asymptotic normality. Our asymptotic framework is that both n and T tends to infinity, but we allow T to grow considerably slower than n as discussed in later.

We first prepare some notations before assumptions. Let $M_g(\beta_g) = \operatorname{Cov}(S_g(\beta_g)) = \sum_{i=1}^n \mathbf{I}\{g_i = g\} D_i^t V_i^{-1} \Sigma_i(\beta_g) V_i^{-1} D_i$ and $H_g(\beta_g) = -\mathbf{E}[\partial S_g(\beta_g) / \partial \beta_g] = \sum_{i=1}^n \mathbf{I}\{g_i = g\} D_i^t V_i^{-1} D_i$. Let $\bar{R}(\beta, \bar{g}) = R(\bar{\alpha}, \beta, \bar{g})$ be a constant positive definite matrix and $\bar{\alpha}$ be a nonrandom constant to which $\widehat{\alpha}$ converges. We do not require $\bar{R}(\beta, \bar{g})$ to be the true correlation matrix R^0 . Next, we denote $\bar{V}_i(\beta_g)$ by replacing $\widehat{R}(\beta, \bar{g})$ with $\bar{R}(\beta, \bar{g})$ in $V_i(\beta_g)$. $\bar{S}_i(\beta_g)$, $\bar{S}_g(\beta_g)$, $\bar{M}_g(\beta_g)$ and $\bar{H}_g(\beta_g)$ are defined similarly. To facilitate the Taylor expansion of the estimating function of GEE, we denote the negative gradient function of $S_i(\beta_g)$ as $\mathcal{D}_i(\beta_g) = -\partial S_i(\beta_g) / \partial \beta_g^t$. $\bar{\mathcal{D}}_i(\beta_g)$ is defined as $\bar{V}_i(\beta_g)$. For $g = 1, \dots, G$, let β_g^0 be a true value of β_g and g_i^0 be a group variable

which i th cluster actually belongs to. Then, we also define the oracle score function for β_g under the true grouping assignment as $S_g^*(\beta_g) = \sum_{i=1}^n \mathbf{I}\{g_i^0 = g\} S_i(\beta_g)$. $\bar{S}_g^*(\beta_g)$, $\bar{M}_g^*(\beta_g)$ and $\bar{H}_g^*(\beta_g)$ are similarly defined. As discussed in Xie and Yang (2003), to prove the existence and weak consistency of the clustered GEE estimators, we need assumptions given later in Assumption (A2), that is, for all $g = 1, \dots, G$, $\bar{H}_g^*(\beta_g^0)$'s or $\lambda_{\min}(\bar{H}^*) \equiv \min_{1 \leq g \leq G} \lambda_{\min}(\bar{H}_g^*(\beta_g^0))$ are divergent at a rate faster than $\tau \equiv \sup_{\beta, \bar{g}} \lambda_{\max}(\{\bar{R}(\beta, \bar{g})\}^{-1} R^0)$. In order to make further assumptions, we need to introduce some notations similar to those in Xie and Yang (2003) and Wang (2011). The following notations are similar to those in Xie and Yang (2003) and are needed to provide assumptions assuring a sufficient conditions for the conditions (I*), (L*) and (CC) in Xie and Yang (2003), under which the existence, weak consistency and asymptotic normality of the GEE estimator hold:

$$\begin{aligned} \kappa &= \max_{1 \leq i \leq n, 1 \leq t \leq T} \{u'(x_{it}^t, \beta_{g_i^0}^0)\}^2, \quad \pi = \sup_{\beta, \bar{g}} \lambda_{\max}(\bar{R}^{-1}(\beta, \bar{g})) / \lambda_{\min}(\bar{R}^{-1}(\beta, \bar{g})), \\ \gamma &= \tau \max_{1 \leq i \leq n, 1 \leq t \leq T} \max_{1 \leq g \leq G} x_{it}^t \{\bar{H}_g^*(\beta_g^0)\}^{-1} x_{it}. \end{aligned}$$

We denote a local neighborhood of $\beta^0 = (\beta_1^{0t}, \dots, \beta_G^{0t})^t$ as $\mathcal{B}_{nT} = \{\beta = (\beta_1^t, \dots, \beta_G^t)^t : \max_{g=1, \dots, G} \|\{\bar{H}_g^*(\beta_g^0)\}^{1/2}(\beta_g - \beta_g^0)\| = C\tau^{1/2}\}$. Lastly, we denote $\varepsilon_{it} = A_{it}^{-1/2}(\beta_{g_i^0}^0)\{y_{it} - m(x_{it}^t, \beta_{g_i^0}^0)\}$ and $\varepsilon_i = (\varepsilon_{i1}, \dots, \varepsilon_{iT})^t$ for all $i = 1, \dots, n$ and $t = 1, \dots, T$.

There are many assumptions for the grouped GEE. Then, we here give some key assumptions and the other regularity assumptions are given in the Supplementary Materials.

Assumption

- (A1) (i) For all $g = 1, \dots, G$, $\lim_{n \rightarrow \infty} (1/n) \sum_{i=1}^n \mathbf{I}\{g_i^0 = g\} = \pi_g > 0$ and (ii) for all $g, g' = 1, \dots, G$ such that $g \neq g'$ and $c > 0$, $\min_{1 \leq g, g' \leq G} \|\beta_g^0 - \beta_{g'}^0\| > c$.
- (A2) $T^6 \tau \lambda_{\min}^{-1}(\bar{H}^*) \rightarrow 0$ and $n/T^\nu \rightarrow 0$ for some large $\nu > 0$.
- (A3) (i) $\pi^2 \kappa \gamma \rightarrow 0$ and (ii) $v \pi \kappa \gamma \rightarrow 0$ for $v = (\sqrt{nT} \wedge T \pi / \min_{1 \leq i \leq n, 1 \leq t \leq T} \{\sigma^2(x_{it}^t, \beta_{g_i^0}^0)\})$.
- (A4) For all $i = 1, \dots, n$ and $t = 1, \dots, T$, $E[\varepsilon_{it}^{2+2/\zeta}] \leq M$ for some $0 < \zeta \leq 1$.

(A5) (i) the eigenvalues of the true correlation matrix R^0 are bounded away from 0 and the eigenvalues of $\bar{R}(\beta, \bar{g})$ are bounded away from 0 uniformly for any β and \bar{g} , and (ii) for any \bar{g} , the eigenvalues of $\bar{R}^{-2}(\beta^0, \bar{g})R^0$ are bounded away from ∞ .

(A6) (i) $\sup_{\beta \in \mathcal{B}_{nT}} \max_{1 \leq k, l \leq T} | \{ R(\alpha, \beta, \bar{g}) - R(\alpha, \beta^0, \bar{g}) \}_{k,l} | = O_p(\lambda_{\min}^{-1/2}(\bar{H}^*)\tau^{1/2})$ for any α and \bar{g} , (ii) for any \bar{g} , $\sup_{\beta \in \mathcal{B}_{nT}} \max_{1 \leq k, l \leq T} | \{ \hat{R}(\beta, \bar{g}) - \bar{R}(\beta, \bar{g}) \}_{k,l} | = O_p(n^{-1/2} \sqrt{\lambda_{\min}^{-1/2}(\bar{H}^*)\tau^{1/2}})$ and $\max_{1 \leq k, l \leq T} | \{ \hat{R}(\beta^0, \bar{g}) - \bar{R}(\beta^0, \bar{g}) \}_{k,l} | = O_p(n^{-1/2})$, and (iii) for any α, β and \bar{g}_{i^*} whose only i th component differs from that of \bar{g} , $\max_{1 \leq k, l \leq T} | \{ R(\alpha, \beta, \bar{g}_{i^*}) - R(\alpha, \beta, \bar{g}) \}_{kl} | = O_p(1/n)$.

Assumption (A1) is typically imposed in the literatures on the grouping approach in panel data models (e.g. Bonhomme and Manresa, 2015), which ensures that the G subgroups are well separated so that the parameters β_g 's and \bar{g} can be identifiable. Assumption (A2) is apparently much stronger than the condition (L*) in Xie and Yang (2003), which says $\tau \lambda_{\min}^{-1}(\bar{H}^*) \rightarrow 0$. This difference comes from the fact that we estimate \bar{g} taking into account the worst case when the correlations of $\{\varepsilon_{it}\}_{t=1, \dots, T}$ do not decay at all. Then, in most cases, such a large T is not required when estimating β_s 's and \bar{g} . Assumption (A3) is the technical assumption similar to the assumptions in Lemma A.2 (ii) and A.3 (ii) of Xie and Yang (2003), which ensure the sufficient conditions for the conditions (I*) and (CC) in Xie and Yang (2003). Assumption (A4) is slightly stronger than the condition in Lemma 2 of Xie and Yang (2003) since we require the fourth moment of error terms to be finite. Assumption (A5)-(i) is the same assumption imposed well in the literatures of GEE with large cluster sizes. Assumption (A5)-(i) is a much weaker assumption than one typically adopted in existing literatures on the grouped estimation (e.g. Bonhomme and Manresa, 2015; Zhang et al., 2019; Gu and Volgushev, 2019) in which $\{\varepsilon_{it}\}_{t=1, \dots, T}$ is assumed to satisfy some strong mixing conditions with a faster-than-polynomial decay rate. Such assumptions are quite unrealistic in longitudinal data analysis, so we do not impose no restriction for the correlation strength of $\{\varepsilon_{it}\}_{t=1, \dots, T}$, which is essentially related

to the use of a kind of Mahalanobis distance for grouping assignment given in (3). On the other hand, Assumption (A5)-(ii) is slightly strict and important to ensure the classification consistency of the clustering method given in (3). This assumption requires that the eigenvalues of $\bar{R}^{-2}(\beta^0, \bar{g})R^0$ are not divergent, which is satisfied if the working correlation matrix reasonably approximate the true correlation matrix. The idea behind Assumption (A6) is similar to that of the condition (A4) in Wang (2011), that is, it is essential to approximate $S_g(\beta_g)$ by $\bar{S}_g^*(\beta_g)$ whose moments are easier to evaluate. For this, Assumption (A6)-(i) and -(ii) say that the estimated working correlation matrix can be approximated by $\bar{R}(\beta^0, \bar{g})$ in a local neighborhood of β_g^0 's and $\bar{\alpha}$. Assumption (A6)-(iii) says that each cluster is linearly additive for the estimation of the working correlation matrix. Then, this is an intuitively reasonable assumption which most of estimators of the working correlation matrix satisfies. In the Supplementary Materials, we provide the accuracy of these approximations under the unstructured working correlation matrix.

We now give our main theorems. We first establish the existence and weak consistency of the grouped GEE estimators and the classification consistency of the grouping variables.

Theorem 1. *Suppose the Assumptions (A1)-(A6) and the Assumptions in the Supplementary Materials hold. For all $g = 1, \dots, G$, $S_g(\beta_g) = 0$ has a root $\hat{\beta}_g$ such that $\hat{\beta}_g \rightarrow \beta_g^0$ in probability. Moreover, as n and T tend to infinity, it holds that $P(\max_{1 \leq i \leq n} |\hat{g}_i(\hat{\beta}) - g_i^0| > 0) = o(1) + O(nT^{-\delta})$ for all $\delta > 0$ for $\hat{g}_i(\hat{\beta})$'s are obtained by (3).*

We next establish the asymptotic normality of $\hat{\beta}_g$ for $g = 1, \dots, G$. We denote $c^* = \max_{1 \leq g \leq G} \lambda_{\max}(\bar{M}_g^*(\beta_g^0)^{-1} \bar{H}_g^*(\beta_g^0))$ and

$$\gamma^* = \max_{1 \leq g \leq G} \max_{i: g_i^0 = g} \lambda_{\max}(\bar{H}_g^*(\beta_g^0)^{-1/2} D_i^t(\beta_g^0) \bar{V}_i^{-1}(\beta_g^0) D_i(\beta_g^0) \bar{H}_g^*(\beta_g^0)^{-1/2}).$$

The following result is a direct consequence of Theorem 4 in Xie and Yang (2003) combined with Lemma S9 in the Supplementary Materials.

Theorem 2. *Suppose the Assumptions (A1)-(A6) and the Assumptions in the Supplementary Materials hold. Moreover, suppose that, for all $g = 1, \dots, G$, there exists a constant ζ such that $(c^*T)^{1+\zeta}\gamma^* \rightarrow 0$ as $n \rightarrow \infty$. Moreover, suppose the marginal distribution of each individual observation has a density of the form from (1). Then, as $n \rightarrow \infty$, we have $\overline{M}_g^*(\beta_g^0)^{-1/2}\overline{H}_g^*(\beta_g^0)(\widehat{\beta}_g - \beta_g^0) \rightarrow N(0, I_p)$ in distribution.*

From Theorem 2, the asymptotic variance of $\widehat{\beta}_g$ is given by $\overline{H}_g^*(\beta_g^0)\overline{M}_g^*(\beta_g^0)^{-1}\overline{H}_g^*(\beta_g^0)$, which involves $\overline{M}_g^*(\beta_g^0)$ depending on the unknown covariance matrix Σ_i . Following Liang and Zeger (1986), we suggest estimating $\overline{M}_g^*(\beta_g^0)$ by

$$\sum_{i=1}^n \mathbf{I}(\widehat{g}_i = g) D_i^t(\beta_g) V_i^{-1}(\beta_g) \{y_i - m(X_i \beta_g)\} \{y_i - m(X_i \beta_g)\}^t V_i^{-1}(\beta_g) D_i(\beta_g) \Big|_{\beta_g = \widehat{\beta}_g},$$

which is consistent as $n \rightarrow \infty$.

4 Numerical studies

4.1 Simulation studies

We investigate the finite sample performance of the proposed grouped GEE method based on simulation studies. We first consider estimation and classification accuracy of the grouped GEE estimator. To this end, we generated two dimensional covariate vector (x_{1it}, x_{2it}) form a two-dimensional normal distribution with mean 0, marginal variance 1 and correlation 0.4, for $i = 1, \dots, n$ and $t = 1, \dots, T$. We considered the logistic model for the marginal expectation of Y_{it} , namely, $Y_{it} \sim \text{Ber}(\pi_{it})$ and $\text{logit}(\pi_{it}) = X_{it}^t \beta_{g_i}$, where $X_{it} = (1, x_{1it}, x_{2it})^t$, $g_i \in \{1, \dots, G\}$ and $\beta_g = (\beta_{g0}, \beta_{g1}, \beta_{g2})$ is a vector of unknown regression coefficients. Here we set $G = 3$ and $\beta_1 = (0, -2, 0)$, $\beta_2 = (0, 1, 2)$ and $\beta_3 = (0, 1, -2)$. For the true grouping assignment, we set $g_i = 1$ for $i = 1, \dots, n/3$, $g_2 = 2$ for $i = n/3 + 1, \dots, 2n/3$ and $g_i = 3$ for $i = 2n/3 + 1, \dots, n$. Based on the probability π_{it} , we generated (Y_{i1}, \dots, Y_{iT}) from a correlated binary vector using R package “bindata” with two scenarios of correlation matrix, exchangeable correlation matrix with 0.5 correlation parameter, and AR(1) correlation matrix with 0.7 correlation parameter. We then applied the proposed grouped GEE method

with $G = 3$ and four options of correlation matrices, independent, exchangeable correlation, AR(1) and unstructured correlation matrices, and unknown parameters in these correlation matrices were also estimated. We evaluated the performance of the estimation of β_g by using the squared error loss defined as $\text{SEL}_g = \sum_{k=1}^2 (\hat{\beta}_{gk} - \beta_{gk})^2$, and assessed the classification accuracy via the classification error given by $\text{CE} = n^{-1} \sum_{i=1}^n \text{I}(\hat{g}_i \neq g_i)$. In Tables 1 and 2, we reported the average values of SEL and CE using 5000 Monte Carlo replications, respectively, under four combinations of (n, T) . From Table 1, we can see that the correct specification of working correlation matrices induces most efficient estimation of the regression coefficient while the use of the other working correlations that are not necessarily equal to the true correlation structures can still provide more efficient estimation than the independence working structure. We also note that the US working correlation includes both EX and AR although the number of unknown parameters is much larger than these structures. Hence, the estimation performance under the moderate sample size such as $(n, T) = (180, 10)$ is not very satisfactory, but the performance is getting improved as the sample size increases. From Table 2, it is observed that introducing working correlation structures in the classification step (3) achieves more accurate classification than the common classification strategy using the standard sum of squared residuals as adopted in existing literatures, when observations within the same subject are correlated. Moreover, the results reveals that the correct specification of the working correlation leads to the most accurate classification.

We next investigate the performance of the CVA selection strategy given in Section 2.3 by adopting the same data generating process with exchangeable correlation structure. For the simulated dataset, we selected the number of component G using the CVA criteria from the candidate $G \in \{2, 3, \dots, 7\}$, noting that the true number of component is 3. We employed four working correlations, ID, EX, AR and US, to carry out the grouped GEE analysis for each G . Based on Monte Carlo replications, we obtained selection probabilities of each G , which are reported in Table 3. It is observed that the use of independent working correlations under significant correlations within the same individual does not necessarily provide satis-

factory selection performance when the number of samples is limited. We can also see that the selection probabilities of the true number of components based on EX and US working correlations tend to be larger than those of using AR working correlation structure since the true correlation is EX. Moreover, when the sample sizes are large such as $(n, T) = (270, 20)$, the adopted CVA strategy can select the true number of component with probability almost 1, which would be compatible with the selection consistency of the strategy.

Finally, we compare the proposed grouped GEE method with the standard random effete models under situations where the subjects do not necessarily admit perfect grouping. To this end, we considered the following underlying scenarios for the subject-specific regression coefficients:

$$(S1) \quad \beta_i = (0, -2, 0)I(g_i = 1) + (0, 1, 2)I(g_i = 2) + (0, 1, -2)I(g_i = 3)$$

$$(S2) \quad \beta_i = (0, -2, 0)I(g_i = 1) + (0, 1, 2)I(g_i = 2) + (0, 1, -2)I(g_i = 3) + U([-0.5, 0.5]^3)$$

$$(S3) \quad \beta_{i0} \sim U([-0.2, 0.2]), \quad \beta_{i1} \sim U([-2, 2]), \quad \beta_{i2} \sim U([0, 2]),$$

where $U(A)$ denotes the uniform distribution on the region A , and g_i is the grouping variable defined as $g_i = 1$ for $i = 1, \dots, n/3$, $g_i = 2$ for $i = n/3 + 1, \dots, 2n/3$ and $g_i = 3$ for $i = 2n/3 + 1, \dots, n$. Hence, the scenario (S1) is the same as one used in the previous simulation study, in which the subjects can be completely clarified into three groups. On the other hand, in scenarios (S2) and (S3), the subjects do not admit complete classification since the regression coefficients are different among subjects. We also note that in scenario (S2), the subjects may admit approximate classification based on g_i , but there seems no trivial classification in scenario (S3) as the regression coefficients are completely random. We generated the binary response variable Y_{it} in the same way as the previous study with the exchangeable correlation structure with 0.5 correlation parameter. For the simulated dataset, we applied the proposed grouped GEE method with the estimated number of groups to estimate β_i by $\widehat{\beta}_{\widehat{g}_i}$. For comparison, we also applied the logistic random coefficient (RC) models by using R package “lme4” (Bates et al., 2016) to estimate β_i via random effects.

We evaluate the estimation performance via the average squared error loss given by $ASL = n^{-1} \sum_{i=1}^n \sum_{k=1}^2 (\hat{\beta}_{ik} - \beta_{ik})^2$, where $\hat{\beta}_{ik}$ is the estimator obtained from grouped GEE or RC methods. In Table 4, we report the average values of ASL based on 5000 Monte Carlo replications. In scenario (S1), since the subject-specific regression coefficients can be perfectly grouped, the proposed methods provide much better estimation accuracy than the RC model. Moreover, in scenario (S2), the subjects do not hold exact grouping structures but can be approximately grouped, and the proposed method still works better than the RC method. On the other hand, the regression coefficients are completely random in scenario (S3), so it is quite natural that the RC method performs better than the proposed method. However, it should be noted that the difference between the grouped GEE and RC methods are relatively comparable, which would indicate that the proposed grouped GEE method can reasonably approximate the subject-specific random coefficients by grouping subjects having similar regression coefficients. Finally, comparing the two working correlations, the EX correlation provides better performance than the US correlation since the EX is the true underlying correlation structure within the same subject, whereas the US correlation is quite comparable with EX.

4.2 Application to election panel study

We apply the proposed method to the 2007-2008 AP YAHOO NEWS election panel study (<http://www.knowledgenetworks.com/GANP/election2008/index.html>). The study intends to measure opinion changes starting with the primary elections through the presidential election in November 2008. The data consist of 4965 participants over 11 waves from November 2007 to December 2008, with nine waves before the election. The main goal of the study is investigating the change of participants' interest in the election over time and relevant factors associated with their interest. Since interest in election news would be a reasonable proxy variable to represent the interest in election, we chose "Question LV3" that among the survey questions, in which five categories of opinions are recorded as an ordinal response variable from 1 to 5, noting that a smaller value corresponds to a high level of interest in the election news. We

Table 1: Average values of squared error loss of the regression coefficients in three groups based on the grouped GEE method with independent (ID), exchangeable correlation (EX), first-order autoregressive (AR) and unstructured (US) working correlation matrices, averaged over 5000 Monte Carlo replications. All values are multiplied by 100.

(n, T)	Group	true correlation: EX				true correlation: AR			
		ID	EX	AR	US	ID	EX	AR	US
(180, 10)	1	23.5	10.2	17.0	13.1	12.3	10.2	9.3	10.8
	2	23.9	10.9	17.3	13.0	13.3	10.0	8.6	11.6
	3	24.4	10.8	17.4	12.8	13.8	10.4	9.0	11.9
(180, 20)	1	13.0	6.3	8.8	7.6	6.2	5.3	5.2	5.9
	2	13.1	6.3	8.6	7.2	5.5	5.6	5.2	5.9
	3	13.4	6.5	8.5	7.4	5.7	6.1	5.6	5.6
(270, 10)	1	18.9	8.1	12.5	9.4	9.1	7.3	6.8	6.8
	2	19.1	7.0	12.2	9.6	10.0	7.3	6.7	7.5
	3	19.1	7.5	12.6	8.9	9.8	7.3	6.5	7.2
(270, 20)	1	8.9	5.2	6.7	4.7	4.7	4.4	4.4	5.6
	2	10.4	6.0	6.6	4.9	5.4	5.0	5.9	4.3
	3	9.9	5.0	6.5	4.7	5.5	4.4	5.1	4.8

Table 2: Average values of classification error (%) of the grouping parameters in the grouped GEE analysis with independent (ID), exchangeable correlation (EX), first-order autoregressive (AR) and unstructured (US) working correlation matrices, averaged over 5000 Monte Carlo replications.

(n, T)	true correlation: EX				true correlation: AR			
	ID	EX	AR	US	ID	EX	AR	US
(180, 10)	9.6	4.4	6.6	5.3	6.5	4.8	4.0	4.8
(180, 20)	4.3	1.5	2.3	1.8	1.9	1.6	1.2	1.5
(270, 10)	8.5	4.3	6.0	4.9	6.1	4.6	4.0	4.4
(270, 20)	3.7	1.5	2.1	1.4	1.8	1.4	1.3	1.4

Table 3: Selection probabilities (%) of the number of groups (G) obtained from the CVA criteria in Section 2.3 with independent (ID), exchangeable (EX), first-order autoregressive (AR) and unstructured (US) working correlation matrices, based on 200 Monte Carlo replications.

(n, T)	working correlation	G					
		2	3	4	5	6	7
(180, 10)	ID	0.5	61.0	8.0	10.0	3.5	17.0
	EX	3.0	95.0	0.5	1.0	0.5	0.0
	AR	3.0	78.0	5.0	7.5	2.5	4.0
	US	0.0	89.5	2.0	2.5	1.0	5.0
(180, 20)	ID	0.0	94.0	3.0	1.0	0.5	1.5
	EX	0.0	100.0	0.0	0.0	0.0	0.0
	AR	0.0	100.0	0.0	0.0	0.0	0.0
	US	0.0	93.5	5.5	0.0	0.0	1.0
(270, 10)	ID	3.0	77.5	3.5	9.5	0.5	6.0
	EX	2.0	98.0	0.0	0.0	0.0	0.0
	AR	2.0	97.0	0.0	0.5	0.0	0.5
	US	0.5	98.5	0.5	0.0	0.5	0.0
(270, 20)	ID	0.0	96.5	1.0	1.5	0.0	1.0
	EX	0.5	99.5	0.0	0.0	0.0	0.0
	AR	0.0	100.0	0.0	0.0	0.0	0.0
	US	0.0	100.0	0.0	0.0	0.0	0.0

Table 4: Average values of average square error loss of estimators of subject-specific regression coefficients based on the grouped GEE (GGEE) with the estimated number of groups and exchangeable (EX) and unstructured (US) working correlation matrices and logistic random coefficients (RC) methods under three scenarios with $n = 180$. The number of Monte Carlo replications is 5000.

Method	(S1)		(S2)		(S3)	
	$T = 10$	$T = 20$	$T = 10$	$T = 20$	$T = 10$	$T = 20$
GGEE-EX	34.7	12.2	46.0	18.9	56.3	45.5
GGEE-US	39.0	17.7	50.7	23.7	58.2	48.4
RC	72.5	70.7	74.4	69.9	49.0	38.1

then dichotomized the response by setting values of 1 or 2 to “much interest” (1) and the other values to “little interest” (0). In this analysis, we only used subjects without no responses in the 9 waves, which results in 1169 subjects. As auxiliary information, we adopted indicator variables of gender (1:male, 0:female), indicators of black and others, respectively, time from 1 to 9, and the interaction term between the gender and time. Among the individuals, it would be reasonable to assume that different types of individuals exist, that is, some individuals are always interested in the election whereas some individuals are not or their interest changes during the term. Therefore, we applied the proposed grouped GEE approach to take account of such heterogeneity.

Let y_{it} be the binary response variable, and x_{it} be the vector of five covariates and an intercept, for $i = 1, \dots, n (= 1169)$ and $t = 1, \dots, T (= 9)$. We consider the mean structure $E[y_{it}|x_{it}] = m(x_{it}^t \beta_{g_i})$ with $m(x) = \exp(x) / \{1 + \exp(x)\}$ and $g_i \in \{1, \dots, G\}$. We first selected the number of groups, G , from candidates $\{2, 3, \dots, 10\}$ using CVA strategy under exchangeable working correlation, and $G = 8$ was selected. Then, we carried out the grouped GEE analysis with $G = 8$ using the three working correlations, exchangeable (EX), the first-order autoregressive (AR) and unstructured (US) correlations. The estimated regression coefficients and their standard errors in 8 groups based on the EX working correlation are shown in Table 5, and the results with AR and US working correlations are reported in the Supplementary Materials. From Table 5, it is observed that regression coefficients of the 8 groups are very different each other. We can also observe that the results with different working correlations does not have much impact on the estimation results, that is, the results are quite comparable among the three working correlations. In order to see the grouping performance, we divided the individuals into 8 groups according to the estimated grouping assignment based on the EX working correlation. In each group, we computed average values of y_{it} for $t = 1, \dots, T$, where the results are presented in Figure 1. From the result, we can easily understand the characteristics of 8 groups. For example, individuals in groups 5 and 2 have high and low interest in the election, respectively, in the whole period. On the other hand, individuals in group 6 and 7

are not so interested in the election at the first period, but their interest increases during the period, although the increasing speed in two groups are quite different. Therefore, we can conclude that the classical GEE analysis assuming homogeneity in the regression coefficients is not a appropriate strategy for the dataset whereas the proposed grouped GEE analysis can successfully capture the potential heterogeneity among individuals.




Figure 1: Estimated probabilities of being interested in the election changing over time in 8 groups identified by the proposed grouped GEE analysis.

5 Concluding Remarks

This paper developed a new statistical approach to analyzing longitudinal data. The proposed method, called grouped GEE analysis, carries out grouping subjects and

Table 5: Point estimates (PE) and its standard errors (SE) of the regression coefficients of gender (G), indicators of black (BL) and others (OT), time (T) and the interaction term of gender and time (T×G) in 8 groups.

		Group							
		1	2	3	4	5	6	7	8
Intercept	PE	-0.37	-0.45	-5.03	0.29	7.89	-3.89	-2.46	2.71
	SE	0.14	0.23	0.45	0.18	1.06	0.33	0.33	0.21
G	PE	2.87	-2.37	0.80	-0.52	-3.29	-0.89	0.85	-3.04
	SE	0.30	0.42	0.60	0.24	1.06	0.65	0.37	0.27
BL	PE	0.12	-2.55	17.10	-0.35	-2.69	-0.40	-1.73	-1.02
	SE	0.15	0.93	0.13	0.23	0.28	0.21	0.30	0.23
OT	PE	-0.73	-0.80	2.34	-0.65	-2.22	-14.04	5.15	-2.07
	SE	0.13	0.09	0.25	0.16	0.38	0.11	1.05	0.20
T	PE	0.27	-0.18	0.26	-0.05	-0.94	0.65	1.05	0.24
	SE	0.03	0.05	0.06	0.03	0.15	0.05	0.12	0.05
T×G	PE	-0.64	-0.01	0.73	0.25	0.95	-0.03	-0.17	-0.32
	SE	0.06	0.09	0.12	0.04	0.16	0.09	0.14	0.06

estimating the regression coefficients simultaneously to take account of potential heterogeneity. We employ working correlations both in estimation and grouping steps, and provided a simple iterative algorithm to obtain grouped GEE estimator. We also developed asymptotic properties of the proposed method. The simulation studies and an application to the election panel study suggest the usefulness of the proposed approach.

The proposed method has some useful extensions. First, we can introduce a penalty term in the grouping step as considered in Sugasawa (2020), which can make subjects have similar characteristics or covariates tend to be classified to the same group. This might make the estimation results more interpretable. Moreover, when the dimension of the regression coefficients is large, it would be better to conduct variable selection, which can be done by introducing penalty function in the estimating equation as considered in Wang et al. (2012). Finally, instead of using working correlation matrices, it would be beneficial to consider quadratic inference functions (Qu et al., 2000), and develop the grouped GEE method with theoretical justifications. We leave the detailed investigation of these issues for interesting future works.

Acknowledgement

This work is partially supported by the Japan Society for the Promotion of Science (JSPS KAKENHI) grant numbers: 18K12757 and 19K23242.

References

- Barban, N. and F. C. Billari (2012). Classifying life course trajectories: a comparison of latent class and sequence analysis. *Journal of the Royal Statistical Society: Series C* 61, 765–784.
- Bates, D., M. Machler, B. Bolker, and S. Walker (2016). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67, 1–48.
- Bonhomme, S. and E. Manresa (2015). Grouped pattern of heterogeneity in panel data. *Econometrica* 83, 1147–1184.
- Coffey, N., J. Hinde, and E. Holian (2014). Clustering longitudinal profiles using p-splines and mixed effects models applied to time-course gene expression data. *Computational Statistics & Data Analysis* 71, 14–29.
- Fokkema, M., N. Smits, A. Zeileis, T. Hothorn, and H. Kelderman (2018). Detecting treatment-subgroup interactions in clustered data with generalized linear mixed-effects model trees. *Behavior Research Methods* 50, 2016–2034.
- Gu, J. and S. Volgushev (2019). Panel data quantile regression with grouped fixed effects. *Journal of Econometrics* 213, 68–91.
- Hajjem, A., F. Bellavance, and D. Larocque (2011). Mixed effects regression trees for clustered data. *Statistics & Probability Letters* 81, 451–459.
- Liang, L. and S. L. Zeger (1986). Longitudinal data analysis using generalized linear models. *Biometrika* 73, 13–22.

- Lin, C. C. and S. Ng (2012). Estimation of panel data models with parameter heterogeneity when group membership is unknown. *Journal of Econometric Methods* 1, 42–55.
- Liu, R., Z. Shang, Y. Zhang, and Q. Zhou (2020). Identification and estimation in panel models with overspecified number of groups. *Journal of Econometrics*, to appear.
- Nagin, D. S., B. L. Jones, V. L. Passos, and R. E. Tremblay (2018). Group-based multi-trajectory modeling. *Statistical Methods in Medical Research* 27, 2015–2023.
- Ng, S. K. and G. J. McLachlan (2014). Mixture models for clustering multilevel growth trajectories. *Computational Statistics & Data Analysis* 71, 43–51.
- Qu, A., B. G. Lindsay, and B. Li (2000). Improving generalised estimating equations using quadratic inference functions. *Biometrika* 87(4), 823–836.
- Rosen, O., W. Jiang, and M. A. Tanner (2000). Mixtures of marginal models. *Biometrika* 87, 391–404.
- Rubin, D. B. and Y. Wu (1997). Modeling schizophrenic behavior using general mixture components. *Biometrics* 53, 243–261.
- Sugasawa, S. (2020). Grouped heterogeneous mixture modeling for clustered data. *Journal of the American Statistical Association*, to appear.
- Sugasawa, S., G. Kobayashi, and Y. Kawakubo (2019). Latent mixture modeling for clustered data. *Statistics and Computing* 29, 537–548.
- Sun, Z., O. Rosen, and A. R. Sampson (2007). Multivariate bernoulli mixture models with application to postmortem tissue studies in schizophrenia. *Biometrics* 63, 901–909.
- Tang, X. and A. Qu (2016). Mixture modeling for longitudinal data. *Journal of Computational and Graphical Statistics* 25, 1117–1137.

- Tang, X., F. Xue, and A. Qu (2020). Individualized multidirectional variable selection. *Journal of the American Statistical Association*, to appear.
- Vogt, M. and O. Linton (2017). Classification of non-parametric regression functions in longitudinal data models. *Journal of the Royal Statistical Society: Series B* 79, 5–27.
- Wang, J. (2010). Consistent selection of the number of clusters via crossvalidation. *Biometrika* 97, 893–904.
- Wang, L. (2011). GEE analysis of clustered binary data with diverging number of covariates. *The Annals of Statistics* 39, 389–417.
- Wang, L. and A. Qu (2009). Consistent model selection and data-driven smooth tests for longitudinal data in the estimating equations approach. *Journal of the Royal Statistical Society: Series B* 71(1), 177–190.
- Wang, L., J. Zhou, and A. Qu (2012). Penalized generalized estimating equations for highdimensional longitudinal data analysis. *Biometrics* 68, 353–360.
- Wedderburn, R. W. (1974). Quasi-likelihood functions, generalized linear models, and the Gauss-newton method. *Biometrika* 61, 439–447.
- Xie, M. and Y. Yang (2003). Asymptotics for generalized estimating equations with large cluster sizes. *The Annals of Statistics* 31(1), 310–347.
- Zhang, Y., J. Wang, and Z. Zhu (2019). Quantile-regression-based clustering for panel data. *Journal of Econometrics* 213, 54–67.
- Zhu, X. and A. Qu (2018). Cluster analysis of longitudinal profiles with subgroups. *Electronic Journal of Statistics* 12, 171–193.

Supplementary Materials for “Grouped GEE Analysis for Longitudinal Data”

Tsubasa Ito and Shonosuke Sugasawa

This Supplementary Materials gives the proofs omitted from the main text. In Section S1, we show Theorem 1, that is, the existence and the weak consistency of the grouped GEE estimator and the classification consistency of the clustering method in (3). At first, we will show that all the clusters are consistently classified to their true groups on average in Lemma S1. Lemma S2 - S8 including the reproduction of lemmas in Xie and Yang (2003) are provided to evaluate the errors from the approximation of the generalized estimating equation by the estimating equation whose moments are easier to evaluate. In Section S2, we show Theorem 2, that is, the asymptotic normality of the grouped GEE estimator. In Lemma S9, we will show the asymptotic equivalence of the proposed estimator and the infeasible estimator obtained by the estimating equation composed of the clusters belonging to their true groups. Lemma S10 and the proof of Theorem 2 are almost the same with those of Xie and Yang (2003). In Section S3, we show that the proposed estimated working correlation matrix satisfy the Assumptions (A5) (ii), (A6) in the main text and (A11) below. Finally, in Section S4, we report the additional results of the application of election panel study discussed in Section 4.2.

In addition to the Assumption (A1)-(A6), we assume the following regularity assumptions for the grouped GEE:

Assumption

(A7) For all $g = 1, \dots, G$, the unknown parameter β_g belongs to a compact subset

$\mathcal{B} \in \mathbb{R}^p$, the true parameter value β_g^0 lies in the interior of \mathcal{B} .

(A8) The covariates $\{x_{it}, i = 1, \dots, n, t = 1, \dots, T\}$ are in a compact set \mathcal{X} .

- (A9) For all $i = 1, \dots, n$ and $t = 1, \dots, T$, $a'(\theta_{it})$ is uniformly three times continuously differentiable, $a''(\theta_{it})$ is uniformly bounded away from 0, and $u(\eta_{it})$ is uniformly four times continuously differentiable and $u'(\eta_{it})$ is uniformly bounded away from 0.
- (A10) For all $i = 1, \dots, n$, there exist positive constants, b_1, b_2, b_3 and b_4 , such that for any $\beta \in \mathcal{B}$ and \bar{g} , $b_1 \leq \lambda_{\min}((nT)^{-1} \sum_{i=1}^n X_i^t X_i) \leq \lambda_{\max}((nT)^{-1} \sum_{i=1}^n X_i^t X_i) \leq b_2$ and $b_3 \leq \lambda_{\min}(T^{-1} X_i^t X_i) \leq \lambda_{\max}(T^{-1} X_i^t X_i) \leq b_4$.
- (A11) For any $\beta \in \mathcal{B}$ and all $\delta > 0$, $\max_{1 \leq k, l \leq T} |\{\widehat{R}(\beta, \bar{g}) - \widehat{R}(\beta, \bar{g}^0)\}_{kl}| = o_p(T^{-\delta})$ for $\bar{g} \in \Gamma$, where $\Gamma = \{\bar{g} = (g_1, \dots, g_n) : \sup_{\beta \in \mathcal{B}_{nT}} n^{-1} \sum_{i=1}^n \mathbf{I}\{g_i \neq g_i^0\} = o_p(T^{-\delta}) \text{ for all } \delta > 0\}$.

Xie and Yang (2003) does not restrict the parameter space of β and the set of all possible covariate variables to be compact. However, we need Assumptions (A7) and (A8) because in the proof of the consistency of our grouped GEE estimators, we need to bound $a''(\theta_{it})$ and $u'_{it}(\eta_{it})$ uniformly on the whole parameter space for all $i = 1, \dots, n$ and $t = 1, \dots, T$. Assumption (A9) requires that the marginal variance of y_{it} is uniformly larger than 0 for any $\beta \in \mathcal{B}$ and $x_{it} \in \mathcal{X}$ for all $i = 1, \dots, n$ and $t = 1, \dots, T$. The boundedness of $a^{(k)}(\theta_{it})$ and $u^{(k)}(\eta_{it})$ for β_g 's in a local neighborhood around β_g^0 is also required to ensure the asymptotic properties of GEE estimators, which is satisfied from Assumptions (A7) and (A8). Assumption (A10) is also imposed well and ensures combined with Assumptions (A1) (i) that $\overline{H}_g(\beta_g)$, $\overline{M}_g(\beta_g)$ and so on are invertible when n or T is sufficiently large. Assumption (A11) says that the estimated working correlation matrix can be approximated by $\overline{R}(\beta, \bar{g}^0)$ if groups are consistently classified to their true groups on average.

Throughout this Supplementary Materials, we use the following notations. The notation $a_{nT} \lesssim b_{nT}$ means that $a_{nT} \leq C b_{nT}$ for all n and T , for some constant C that does not depends on n and T . For a column vector a , we use a^t to denote the transpose of a and $\|a\|$ to denote the Euclidean norm of a . For a matrix A , $\{A\}_{kl}$ denotes the (k, l) -element of A , $\lambda_{\min}(A)$ ($\lambda_{\max}(A)$) denotes the smallest (largest) eigenvalue of A ,

A^t denotes the transpose of A and $\|A\|_F = \{\text{tr}(A^t A)\}^{1/2}$ is the Frobenius norm of A .

We use the notation $a \vee b = \max(a, b)$ and $a \wedge b = \min(a, b)$.

S1 Proof of Theorem 1

First of all, we need to show the next lemma.

Lemma S1. *Under the Assumptions (A1)-(A11), it holds that for all $\delta > 0$,*

$$\sup_{\beta \in \mathcal{B}_{nT}} \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{\widehat{g}_i(\beta) \neq g_i^0\} = o_p(T^{-\delta}),$$

where $\widehat{g}_i(\beta)$ is obtained by (3).

Proof. For any \bar{g} , \bar{g}_{i0} is defined to be obtained by replacing only its i th element with g_i^0 , that is $\bar{g}_{i0} = (g_1, \dots, g_{i-1}, g_i^0, g_{i+1}, \dots, g_n)$. Note that, from the definition of $\widehat{g}_i(\beta)$, we have, for all $g = 1, \dots, G$,

$$\begin{aligned} \mathbb{I}\{\widehat{g}_i(\beta) = g\} &\leq \mathbb{I}\left\{\{y_i - m(X_i \beta_g)\}^t \widehat{R}^{-1}(\beta, \bar{g}) \{y_i - m(X_i \beta_g)\}\right. \\ &\quad \left. \leq \{y_i - m(X_i \beta_{g_i^0})\}^t \widehat{R}^{-1}(\beta, \bar{g}_{i0}) \{y_i - m(X_i \beta_{g_i^0})\}\right\}. \end{aligned}$$

Then, we can write

$$\frac{1}{n} \sum_{i=1}^n \mathbb{I}\{\widehat{g}_i(\beta) \neq g_i^0\} = \sum_{g=1}^G \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{g_i^0 \neq g\} \mathbb{I}\{\widehat{g}_i(\beta) = g\} \leq \sum_{g=1}^G \frac{1}{n} \sum_{i=1}^n Z_{ig}(\beta_g),$$

where

$$\begin{aligned} Z_{ig}(\beta_g) &= \mathbb{I}\{g_i^0 \neq g\} \mathbb{I}\left\{\{y_i - m(X_i \beta_g)\}^t \widehat{R}^{-1}(\beta, \bar{g}) \{y_i - m(X_i \beta_g)\}\right. \\ &\quad \left. \leq \{y_i - m(X_i \beta_{g_i^0})\}^t \widehat{R}^{-1}(\beta, \bar{g}_{i0}) \{y_i - m(X_i \beta_{g_i^0})\}\right\}. \end{aligned}$$

Similar to the proof of Lemma B.4 in Bonhomme and Manresa (2015), we start by

bounding $Z_{ig}(\beta_g)$ on $\beta \in \mathcal{B}_{nT}$ by a quantity that does not depend on β . Denote

$$\begin{aligned} W_{ig}(\beta) = & \{y_i - m(X_i\beta_g)\}^t \widehat{R}^{-1}(\beta, \bar{g}) \{y_i - m(X_i\beta_g)\} \\ & - \{y_i - m(X_i\beta_{g_i^0})\}^t \widehat{R}^{-1}(\beta, \bar{g}_{i0}) \{y_i - m(X_i\beta_{g_i^0})\}, \end{aligned}$$

then we have

$$Z_{ig}(\beta_g) = \mathbf{I}\{g_i^0 \neq g\} \mathbf{I}\{W_{ig}(\beta) \leq 0\} \leq \mathbf{I}\{g_i^0 \neq g\} \mathbf{I}\{W_{ig}(\beta^0) \leq |W_{ig}(\beta^0) - W_{ig}(\beta)|\}.$$

We have

$$\begin{aligned} |W_{ig}(\beta^0) - W_{ig}(\beta)| & \leq \left| \{y_i - m(X_i\beta_{g_i^0}^0)\}^t \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) \{y_i - m(X_i\beta_{g_i^0}^0)\} \right. \\ & \quad \left. - \{y_i - m(X_i\beta_{g_i^0})\}^t \widehat{R}^{-1}(\beta, \bar{g}_{i0}) \{y_i - m(X_i\beta_{g_i^0})\} \right| \\ & \quad + \left| \{y_i - m(X_i\beta_g^0)\}^t \widehat{R}^{-1}(\beta^0, \bar{g}) \{y_i - m(X_i\beta_g^0)\} \right. \\ & \quad \left. - \{y_i - m(X_i\beta_g)\}^t \widehat{R}^{-1}(\beta, \bar{g}) \{y_i - m(X_i\beta_g)\} \right| \\ & \equiv K_{ig}^{(1)}(\beta) + K_{ig}^{(2)}(\beta). \end{aligned}$$

Denote $\tilde{\varepsilon}_i = \Sigma_i^{-1/2} \{y_i - m(X_i\beta_{g_i^0}^0)\}$, then we can write

$$\begin{aligned} K_{ig}^{(1)}(\beta) & \leq |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \{\widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) - \widehat{R}^{-1}(\beta, \bar{g}_{i0})\} \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\ & \quad + 2|\{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_{g_i^0})\}^t \widehat{R}^{-1}(\beta, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\ & \quad + \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_{g_i^0})\}^t \widehat{R}^{-1}(\beta, \bar{g}_{i0}) \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_{g_i^0})\} \\ & \equiv \sum_{j=1}^3 I_j. \end{aligned}$$

For I_1 , we can write

$$\begin{aligned}
& I_1 \\
&= |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) \{ \widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0}) \} \widehat{R}^{-1}(\beta, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\leq |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) \{ \widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0}) \} \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\quad + |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) \{ \widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0}) \} \{ \widehat{R}^{-1}(\beta, \bar{g}_{i0}) - \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) \} \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\equiv I_{11} + I_{12}.
\end{aligned}$$

For I_{11} , we have

$$\begin{aligned}
& I_{11} \\
&= |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \{ \widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0}) \} \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\quad + 2 |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \{ \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \} \{ \widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0}) \} \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\quad + |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \{ \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \} \{ \widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0}) \} \\
&\quad \quad \times \{ \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \} \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\equiv \sum_{j=1}^3 I_{11j}.
\end{aligned}$$

Since we have $\lambda_{\max}(\bar{R}^{-2}(\beta^0, \bar{g}_{i0}) \Sigma_i) \leq \lambda_{\max}(\bar{R}^{-2}(\beta^0, \bar{g}_{i0}) R^0) \lambda_{\max}(A_i(\beta_{g_i^0}^0))$ for all $i = 1, \dots, n$ and $\max_{1 \leq i \leq n} \max_{1 \leq t \leq T} A_{it}(\beta_g) < \infty$ for any $\beta \in \mathcal{B}$ from Assumptions (A7), (A8) and (A9), we have

$$I_{111} \lesssim \| \widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0}) \|_F \lambda_{\max}(\bar{R}^{-2}(\beta^0, \bar{g}_{i0}) R^0) \tilde{\varepsilon}_i^t \tilde{\varepsilon}_i.$$

Then, from Assumptions (A6) (i) and (A11), there is a constant C_{111} , independent of n and T such that

$$\sup_{\beta \in \mathcal{B}_{nT}} I_{111} = C_{111} C T^2 \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \right).$$

For I_{112} , we have

$$\begin{aligned}
I_{112}/2 &\leq |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \bar{R}(\beta^0, \bar{g}_{i0})\} \\
&\quad \times \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \{\widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0})\} \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\quad + |\tilde{\varepsilon}_i^t \Sigma_i^{1/2} \{\widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g}_{i0})\} \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \bar{R}(\beta^0, \bar{g}_{i0})\} \\
&\quad \times \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \{\widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0})\} \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\equiv I_{1121} + I_{1122}.
\end{aligned}$$

For I_{1121} , we have from Cauchy-Schwarz inequality

$$\begin{aligned}
I_{1121} &\lesssim \|\widehat{R}(\beta^0, \bar{g}_{i0}) - \bar{R}(\beta^0, \bar{g}_{i0})\|_F \|\widehat{R}(\beta, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g}_{i0})\|_F \\
&\quad \times \lambda_{\min}^{-1}(\bar{R}(\beta^0, \bar{g}_{i0})) \lambda_{\max}(\bar{R}^{-2}(\beta^0, \bar{g}_{i0}) R^0) \tilde{\varepsilon}_i^t \tilde{\varepsilon}_i.
\end{aligned}$$

From Assumptions (A5), (A6) (i), (ii) and (A11), there is a constant C_{1121} , independent of n and T such that

$$\sup_{\beta \in \mathcal{B}_{nT}} I_{1121} = C_{1121} C T^2 \sqrt{\frac{T^2}{n}} \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \right),$$

which is dominated by I_{111} since $T^2/n \rightarrow 0$ from Assumption (A2). Similarly, I_{1122} , I_{113} and I_{12} are dominated by I_{111} . Then, there is a constant C_1 , independent of n and T such that

$$\sup_{\beta \in \mathcal{B}_{nT}} I_1 = C_1 C T^2 \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \left(\frac{1}{T} \sum_{j=1}^T \tilde{\varepsilon}_{jt}^2 \right).$$

For I_2 , we have

$$\begin{aligned}
I_2/2 &\leq |\{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_{g_i^0})\}^t \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\quad + |\{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_{g_i^0})\}^t \{\widehat{R}^{-1}(\beta, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g}_{i0})\} \Sigma_i^{1/2} \tilde{\varepsilon}_i| \\
&\equiv I_{21} + I_{22}.
\end{aligned}$$

From Taylor expansion, for $\beta_{g_i}^*$ between $\beta_{g_i}^0$ and $\beta_{g_i}^0$, we have

$$m(X_i\beta_{g_i}^0) - m(X_i\beta_{g_i}^0) = \phi A_i(\beta_{g_i}^*)\Delta(\beta_{g_i}^*)X_i(\beta_{g_i}^0 - \beta_{g_i}^0). \quad (\text{S1})$$

Since $\max_{1 \leq i \leq n} \max_{1 \leq t \leq T} u'(x_{it}\beta_g) < \infty$ from Assumptions (A7), (A8) and (A9), we have

$$\begin{aligned} & I_{21} \\ & \leq \|\Sigma_i^{1/2}\bar{R}^{-1}(\beta^0, \bar{g}_{i0})\{m(X_i\beta_{g_i}^0) - m(X_i\beta_{g_i}^0)\}\| \cdot \|\tilde{\varepsilon}_i\| \\ & \lesssim \lambda_{\max}^{1/2}(\bar{R}^{-2}(\beta^0, \bar{g}_{i0})R^0)\{(\beta_{g_i}^0 - \beta_{g_i}^0)X_i^t\Delta(\beta_{g_i}^*)A_i^2(\beta_{g_i}^*)\Delta(\beta_{g_i}^*)X_i(\beta_{g_i}^0 - \beta_{g_i}^0)\}^{1/2}\|\tilde{\varepsilon}_i\| \\ & \lesssim \lambda_{\max}^{1/2}(\bar{R}^{-2}(\beta^0, \bar{g}_{i0})R^0)\lambda_{\max}^{1/2}(X_i^tX_i)\|\beta_{g_i}^0 - \beta_{g_i}^0\|(\tilde{\varepsilon}_i^t\tilde{\varepsilon}_i)^{1/2}. \end{aligned}$$

Then, from Assumptions (A5), (A10) there is a constant C_{21} , independent of n and T such that

$$\sup_{\beta \in \mathcal{B}_{nT}} I_{21} \leq C_{21}CT\lambda_{\min}^{-1/2}(\bar{H}^*)\tau^{1/2}\left(\frac{1}{T}\sum_{t=1}^T \tilde{\varepsilon}_{it}^2\right)^{1/2}.$$

For I_{22} , we have

$$\begin{aligned} & I_{22} \\ & \leq |\{m(X_i\beta_{g_i}^0) - m(X_i\beta_{g_i}^0)\}^t \bar{R}^{-1}(\beta^0, \bar{g}_{i0})\{\bar{R}(\beta^0, \bar{g}_{i0}) - \hat{R}(\beta, \bar{g}_{i0})\}\bar{R}^{-1}(\beta^0, \bar{g}_{i0})\Sigma_i^{1/2}\tilde{\varepsilon}_i| \\ & \quad + |\{m(X_i\beta_{g_i}^0) - m(X_i\beta_{g_i}^0)\}^t \{\hat{R}^{-1}(\beta, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g}_{i0})\} \\ & \quad \quad \times \{\bar{R}(\beta^0, \bar{g}_{i0}) - \hat{R}(\beta, \bar{g}_{i0})\}\bar{R}^{-1}(\beta^0, \bar{g}_{i0})\Sigma_i^{1/2}\tilde{\varepsilon}_i| \\ & \equiv I_{221} + I_{222}. \end{aligned}$$

As is the same with I_{21} and from Assumption (A6) (i) and (ii), there is a constant C_{221} , independent of n and T such that

$$\sup_{\beta \in \mathcal{B}_{nT}} I_{221} \leq C_{221}CT\left\{T\lambda_{\min}^{-1/2}(\bar{H}^*)\tau^{1/2} \vee \sqrt{\frac{T^2}{n}}\right\}\left(\frac{1}{T}\sum_{t=1}^T \tilde{\varepsilon}_{it}^2\right)^{1/2},$$

which dominates I_{21} . Since I_{222} is dominated by I_{221} from Assumption (A6) (i) and (ii), there is a constant C_2 , independent of n and T such that

$$\sup_{\beta \in \mathcal{B}_{nT}} I_2 \leq C_2 C T \left\{ T \lambda_{\min}^{-1/2}(\overline{H}^*) \tau^{1/2} \vee \sqrt{\frac{T^2}{n}} \right\} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \right)^{1/2}.$$

For I_3 , we have

$$\begin{aligned} I_3 &\leq \{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_{g_i^0})\}^t \overline{R}^{-1}(\beta^0, \overline{g}_{i0}) \{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_{g_i^0})\} \\ &\quad + \{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_{g_i^0})\}^t \{\widehat{R}^{-1}(\beta, \overline{g}_{i0}) - \overline{R}^{-1}(\beta^0, \overline{g}_{i0})\} \{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_{g_i^0})\} \\ &\equiv I_{31} + I_{32}. \end{aligned}$$

As is the case in I_{21} and from Assumption (A5), there is a constant C_{31} , independent of n and T such that $\sup_{\beta \in \mathcal{B}_{nT}} I_{31} \leq C_{31} C^2 T \lambda_{\min}^{-1}(\overline{H}^*) \tau$. From Assumption (A6) (i) and (ii), there is a constant C_{32} , independent of n and T such that

$$\sup_{\beta \in \mathcal{B}_{nT}} I_{32} \leq C_{32} C^2 T \left\{ T \lambda_{\min}^{-1/2}(\overline{H}^*) \tau^{1/2} \vee \sqrt{\frac{T^2}{n}} \right\} \lambda_{\min}^{-1}(\overline{H}^*) \tau,$$

which is dominated by I_{31} from Assumption (A2). then, there is a constant C_3 , independent of n and T such that $\sup_{\beta \in \mathcal{B}_{nT}} I_3 \leq C_3 C^2 T \lambda_{\min}^{-1}(\overline{H}^*) \tau$. For $K_{ig}^{(2)}(\beta)$, we can write

$$\begin{aligned} K_{ig}^{(2)}(\beta) &\leq |\{y_i - m(X_i \beta_g^0)\}^t \widehat{R}^{-1}(\beta^0, \overline{g}) \{y_i - m(X_i \beta_g^0)\}| \\ &\quad + |\{y_i - m(X_i \beta_g)\}^t \widehat{R}^{-1}(\beta, \overline{g}) \{y_i - m(X_i \beta_g)\}| \\ &\equiv I_4 + I_5. \end{aligned}$$

For I_4 , we can write

$$\begin{aligned} I_4 &\leq |\{y_i - m(X_i \beta_g^0)\}^t \overline{R}^{-1}(\beta^0, \overline{g}) \{y_i - m(X_i \beta_g^0)\}| \\ &\quad + |\{y_i - m(X_i \beta_g^0)\}^t \{\widehat{R}^{-1}(\beta^0, \overline{g}) - \overline{R}^{-1}(\beta^0, \overline{g})\} \{y_i - m(X_i \beta_g^0)\}| \\ &\equiv I_{41} + I_{42}. \end{aligned}$$

Since it is obvious that $J_{41} = O_p(T)$ and I_{42} is dominated by I_{41} from Assumption (A6) (i) and (ii), we have $I_4 = O_p(T)$. Since we can also show that $I_5 = O_p(T)$, there is a constant C_4 , independent of n and T such that $\sup_{\beta \in \mathcal{B}_{nT}} K_{ig}^{(2)}(\beta) \leq C_4 T$. Next, we will bound $W_{ig}(\beta^0, \bar{g})$ from below. It can be written as

$$\begin{aligned} W_{ig}(\beta^0, \bar{g}) &= \{y_i - m(X_i \beta_{g_i^0}^0)\}^t \{\widehat{R}^{-1}(\beta^0, \bar{g}) - \widehat{R}^{-1}(\beta^0, \bar{g}_{i0})\} \{y_i - m(X_i \beta_{g_i^0}^0)\} \\ &\quad + \{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_g^0)\}^t \widehat{R}^{-1}(\beta^0, \bar{g}) \{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_g^0)\} \\ &\quad + 2\{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_g^0)\}^t \widehat{R}^{-1}(\beta^0, \bar{g}) \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &\equiv \sum_{j=1}^3 J_j. \end{aligned}$$

For J_1 , we have

$$\begin{aligned} J_1 &= \{y_i - m(X_i \beta_{g_i^0}^0)\}^t \widehat{R}^{-1}(\beta^0, \bar{g}) \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g})\} \widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) \{y_i - m(X_i \beta_{g_i^0}^0)\} \\ &= \tilde{\varepsilon}_i^t \Sigma_i^{1/2} \bar{R}^{-1}(\beta^0, \bar{g}) \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g})\} \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &\quad + \tilde{\varepsilon}_i^t \Sigma_i^{1/2} \{\widehat{R}^{-1}(\beta^0, \bar{g}) - \bar{R}^{-1}(\beta^0, \bar{g})\} \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g})\} \bar{R}^{-1}(\beta^0, \bar{g}_{i0}) \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &\quad + \tilde{\varepsilon}_i^t \Sigma_i^{1/2} \bar{R}^{-1}(\beta^0, \bar{g}) \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g})\} \{\widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g}_{i0})\} \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &\quad + \tilde{\varepsilon}_i^t \Sigma_i^{1/2} \{\widehat{R}^{-1}(\beta^0, \bar{g}) - \bar{R}^{-1}(\beta^0, \bar{g})\} \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g})\} \\ &\quad \times \{\widehat{R}^{-1}(\beta^0, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g}_{i0})\} \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &= \sum_{j=1}^4 J_{1j}. \end{aligned}$$

For J_{11} , we have

$$\begin{aligned} J_{11} &= \tilde{\varepsilon}_i^t \Sigma_i^{1/2} \bar{R}^{-1}(\beta^0, \bar{g}) \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g})\} \bar{R}^{-1}(\beta^0, \bar{g}) \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &\quad + \tilde{\varepsilon}_i^t \Sigma_i^{1/2} \bar{R}^{-1}(\beta^0, \bar{g}) \{\widehat{R}(\beta^0, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g})\} \{\bar{R}^{-1}(\beta^0, \bar{g}_{i0}) - \bar{R}^{-1}(\beta^0, \bar{g})\} \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &\equiv J_{111} + J_{112}. \end{aligned}$$

For J_{111} , we have $J_{111} \gtrsim -\lambda_{\max}(\bar{R}^{-2}(\beta^0, \bar{g}) R^0) \|\widehat{R}(\beta^0, \bar{g}_{i0}) - \widehat{R}(\beta^0, \bar{g})\|_F \tilde{\varepsilon}_i^t \tilde{\varepsilon}_i$. From Assumptions (A6) (iii) and (A11), there is a constant C_{511} , independent of C and T , such that $J_{111} \geq -C_{511}(T/n) \tilde{\varepsilon}_i^t \tilde{\varepsilon}_i$. From Assumptions (A2), (A6) (i) and (ii), it can

be shown that J_{112} , J_{12} , J_{13} and J_{14} are dominated by J_{111} , then there is a constant C_5 , independent of C and T , such that $J_1 \geq -C_5(T^2/n)(\sum_{t=1}^T \tilde{\varepsilon}_{it}^2/T)$. For J_2 , we have

$$\begin{aligned} J_2 &= \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_g^0)\}^t \bar{R}^{-1}(\beta^0, \bar{g}) \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_g^0)\} \\ &\quad + \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_g^0)\}^t \{\widehat{R}^{-1}(\beta^0, \bar{g}) - \bar{R}^{-1}(\beta^0, \bar{g})\} \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_g^0)\} \\ &\equiv J_{21} + J_{22}. \end{aligned}$$

For J_{21} , by using (S1), we have for $\beta_{g_i^*}^*$ between $\beta_{g_i^0}^0$ and β_g^0

$$J_{21} = (\beta_{g_i^0}^0 - \beta_g^0) X_i^t \Delta(\beta_{g_i^*}^*) A_i(\beta_{g_i^*}^*) \bar{R}^{-1}(\beta^0, \bar{g}) A_i(\beta_{g_i^*}^*) \Delta(\beta_{g_i^*}^*) X_i (\beta_{g_i^0}^0 - \beta_g^0).$$

Since $\lambda_{\max}(X_i^t \bar{R}^{-1}(\beta^0, \bar{g}) X_i)$ is at least of order $O_p(T)$ from Assumption (A10). Then, from Assumption (A1) (ii) there is a constant C_6^* , independent of C and T , such that $J_{21} \geq C_6^* T$. From Assumptions (A5) and (A6) (ii), it can be shown that J_{22} is dominated by J_{21} , then there is a constant C_6 , independent of C and T , such that $J_2 \geq C_6 T$. For J_3 , we have

$$\begin{aligned} J_3 &= \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_g^0)\}^t \bar{R}^{-1}(\beta^0, \bar{g}) \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &\quad + \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_g^0)\}^t \{\widehat{R}^{-1}(\beta^0, \bar{g}) - \bar{R}^{-1}(\beta^0, \bar{g})\} \Sigma_i^{1/2} \tilde{\varepsilon}_i \\ &\equiv J_{31} + J_{32}. \end{aligned}$$

From Assumption (A6) (i) and (ii), J_{32} is dominated by J_{31} . Let $U\Lambda U^t$ be the eigendecomposition of $\bar{R}^{-1}(\beta^0, \bar{g}) \Sigma_i^{1/2}$, where $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_T)$ for $\lambda_1 \geq \dots, \lambda_T$ is a diagonal matrix formed from the eigenvalues and U is the corresponding eigenvectors of $\bar{R}^{-1}(\beta^0, \bar{g}) \Sigma_i^{1/2}$. Then there is a constant C_7 , independent of C and T , such that

$$J_3 = C_7 \{m^*(X_i\beta_{g_i^0}^0) - m^*(X_i\beta_g^0)\}^t \Lambda \tilde{\varepsilon}_i^* = C_7 \sum_{t=1}^T \lambda_t \{m^*(x_{it}^t \beta_{g_i^0}^0) - m^*(x_{it}^t \beta_g^0)\} \tilde{\varepsilon}_{it}^*,$$

for $m^*(X_i\beta_g) = U m(X_i\beta_g)$ and $\tilde{\varepsilon}_i^* = U \tilde{\varepsilon}_i$. Combined with the above results, we thus

obtain

$$\begin{aligned}
& \sup_{\beta \in \mathcal{B}_{nT}} Z_{ig}(\beta_g) \\
& \leq \mathbf{I}\{g_i^0 \neq g\} \\
& \quad \times \mathbf{I}\left\{ -C_5 \frac{T^2}{n} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \right) + C_6 T + C_7 \sum_{t=1}^T \lambda_t \{m^*(x_{it}^t, \beta_{g_i^0}^0) - m^*(x_{it}^t, \beta_g^0)\} \tilde{\varepsilon}_{it}^* \right. \\
& \quad \leq C_1 C T^2 \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \right) \\
& \quad \quad + C_2 C T \left\{ T \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee \sqrt{\frac{T^2}{n}} \right\} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \right)^{1/2} \\
& \quad \quad \left. + C_3 C^2 T \lambda_{\min}^{-1}(\bar{H}^*) \tau \right\}.
\end{aligned}$$

Since the right-hand side of the above inequality does not depend on β_g for $g = 1, \dots, G$, we can denote it as \tilde{Z}_{ig} . As a result, we have

$$\sup_{\beta \in \mathcal{B}_{nT}} \frac{1}{n} \sum_{i=1}^n \mathbf{I}\{\hat{g}_i(\beta) \neq g_i^0\} \leq \frac{1}{n} \sum_{i=1}^n \sum_{g=1}^G \tilde{Z}_{ig}.$$

Using standard probability algebra, we have for all g and M in Assumption (A4),

$$\begin{aligned}
& P(\tilde{Z}_{ig} = 1) \\
& \leq P\left(-C_5 \frac{T^2}{n} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2\right) + C_6 T + C_7 \sum_{t=1}^T \lambda_t \{m^*(x_{it}^t \beta_{g_i}^0) - m^*(x_{it}^t \beta_g^0)\} \tilde{\varepsilon}_{it}^*\right) \\
& \quad \leq C_1 C T^2 \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2\right) \\
& \quad \quad + C_2 C T \left\{T \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee \sqrt{\frac{T^2}{n}}\right\} \left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2\right)^{1/2} \\
& \quad \quad + C_3 C^2 T \lambda_{\min}^{-1}(\bar{H}^*) \tau \\
& \leq P\left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \geq \frac{n}{T^2} M\right) + P\left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \geq T^{-2} \lambda_{\min}^{1/2}(\bar{H}^*) \tau^{-1/2} M\right) \\
& \quad + P\left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \geq T^{-2} \left\{T^{-2} \lambda_{\min}(\bar{H}^*) \tau^{-1} \wedge n/T^2\right\} M\right) \\
& \quad \quad + P\left(C_7 \sum_{t=1}^T \lambda_t \{m^*(x_{it}^t \beta_{g_i}^0) - m^*(x_{it}^t \beta_g^0)\} \tilde{\varepsilon}_{it}^*\right) \\
& \quad \leq C_5 M - C_6 T + C_1 C M T + C_2 C \sqrt{M} T + C_3 C^2 T \lambda_{\min}^{-1}(\bar{H}^*) \tau.
\end{aligned}$$

From Markov's inequality, we have for any $\delta > 0$,

$$P\left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \geq \frac{n}{T^2} M\right) \leq \exp\left(-\frac{n}{T^2} M\right) E\left[\exp\left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2\right)\right] = o_p(T^{-\delta}),$$

Since the order of $T^{-1} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2$ is at most $O_p(T)$, we have $P(T^{-1} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \geq (n/T^2)M) = o_p(T^{-\delta})$ for any $\delta > 0$ from Assumption (A2). Similarly, we have

$$\begin{aligned}
& P\left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \geq T^{-2} \lambda_{\min}^{1/2}(\bar{H}^*) \tau^{-1/2} M\right) \\
& \leq \exp\left(-T^{-2} \lambda_{\min}^{1/2}(\bar{H}^*) \tau^{-1/2} M\right) E\left[\exp\left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2\right)\right] = o_p(T^{-\delta}),
\end{aligned}$$

where the second inequality follows from Assumption (A2). Similarly, we have

$$P\left(\frac{1}{T} \sum_{t=1}^T \tilde{\varepsilon}_{it}^2 \geq T^{-2} \left\{ T^{-2} \lambda_{\min}(\overline{H}^*) \tau^{-1} \wedge n/T^2 \right\} M\right) = o_p(T^{-\delta}).$$

For the last probability, we can write

$$\begin{aligned} & P\left((1/T)C_7 \sum_{t=1}^T \lambda_t \{m^*(x_{it}^t, \beta_{g_0}^0) - m^*(x_{it}^t, \beta_g^0)\} \tilde{\varepsilon}_{it}^* \right. \\ & \quad \left. \leq (1/T)C_5 M - C_6 + C_1 C M + C_2 C \sqrt{M} + C_3 C^2 \lambda_{\min}^{-1}(\overline{H}^*) \tau\right). \end{aligned}$$

For the right-hand side of the inequality in the above probability, both of the first and last terms are asymptotically dominated by other terms. Then, by taking $C < C_6/(C_1 M + C_2 \sqrt{M})$, the right-hand side of the inequality can be negative. Moreover, it is noted that $\lambda_1 = O_p(1)$. Then, the left-hand side of the inequality is a linear combination of $\tilde{\varepsilon}_{it}^*$, and its expectation is 0 and the order of its variance is at most $O(T^{-1})$. Since $\tilde{\varepsilon}_{it}^*$ for $t = 1, \dots, T$ are uncorrelated, we can use Theorem 6.2 in Rio (2000), in which the second term of the right-hand side of the equation (6.5) vanishes in this case due to the uncorrelatedness of $\tilde{\varepsilon}_{it}^*$'s. Thus, by using the consequence of Theorem 6.2 in Rio (2000) for $\lambda = T$, $r = T^{1/2}$ and $s_n^2 = T$, the probability above is bounded above by $4(1 + T^{1/2})^{-T^{1/2}} = o_p(T^{-\delta})$ for any $\delta > 0$. This ends the proof. \square

Similar to Wang (2011), in order to prove the consistency it is essential to approximate $S_g(\beta_g)$, $H_g(\beta_g)$ and so on by $\overline{S}_g^*(\beta_g)$ and $\overline{H}_g^*(\beta_g)$ whose moments are easier to evaluate. The following lemmas S2-S8 establish the accuracy of these approximations, which play important roles in deriving the asymptotic normality.

Lemma S2. *Under Assumptions (A1)-(A11), it holds that, for all $g = 1, \dots, G$ and all $\delta > 0$,*

$$\begin{aligned} & \sup_{\beta \in \mathcal{B}_{nT}, \overline{g} \in \Gamma} \|\{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \{S_g(\beta_g) - S_g^*(\beta_g)\}\| = O_p(\lambda_{\min}^{-1/2}(\overline{H}^*) n T) o_p(T^{-\delta}), \\ & \sup_{\beta \in \mathcal{B}_{nT}, \overline{g} \in \Gamma} \|\{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \{\overline{S}_g(\beta_g) - \overline{S}_g^*(\beta_g)\}\| = O_p(\lambda_{\min}^{-1/2}(\overline{H}^*) n T) o_p(T^{-\delta}). \end{aligned}$$

Proof. We will show the second part of the lemma. Form Assumption (A6) (ii), the first part of the lemma can be shown similarly by replacing $\bar{R}(\beta, \bar{g})$ and $\bar{R}(\beta, \bar{g}^0)$ with $\hat{R}(\beta, \bar{g})$ and $\hat{R}(\beta, \bar{g}^0)$ respectively. It can be written as

$$\begin{aligned}
& \bar{S}_g(\beta_g) - \bar{S}_g^*(\beta_g) \\
&= \sum_{i=1}^n \mathbb{I}\{g_i = g\} X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \bar{R}^{-1}(\beta, \bar{g}) A_i^{-1/2}(\beta_g) \{y_i - m(X_i \beta_g)\} \\
&\quad - \sum_{i=1}^n \mathbb{I}\{g_i^0 = g\} X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \bar{R}^{-1}(\beta, \bar{g}^0) A_i^{-1/2}(\beta_g) \{y_i - m(X_i \beta_g)\} \\
&= \sum_{i=1}^n \mathbb{I}\{g_i^0 = g\} X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \{\bar{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g}^0)\} A_i^{-1/2}(\beta_g) \{y_i - m(X_i \beta_g)\} \\
&\quad + \sum_{i=1}^n (\mathbb{I}\{g_i = g\} - \mathbb{I}\{g_i^0 = g\}) X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \bar{R}^{-1}(\beta, \bar{g}) A_i^{-1/2}(\beta_g) \{y_i - m(X_i \beta_g)\} \\
&\equiv I_1 + I_2.
\end{aligned}$$

For I_1 , we have

$$\begin{aligned}
I_1 &= \sum_{i: g_i^0 = g} \sum_{t_1, t_2=1}^T \{\bar{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g}^0)\}_{t_1, t_2} A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) \{y_{it_2} - m(x_{it_2}^t \beta_g)\} x_{it_1} \\
&= \sum_{t_1=1}^T \sum_{t_2=1}^T \{\bar{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g}^0)\}_{t_1, t_2} \\
&\quad \times \left[\sum_{i: g_i^0 = g} A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) \{A_{it_2}^{1/2}(\beta_g^0) \varepsilon_{it_2} + m(x_{it_2}^t \beta_g^0) - m(x_{it_2}^t \beta_g)\} x_{it_1} \right].
\end{aligned}$$

It is noted that we have

$$E \left[\left\| \sum_{i: g_i^0 = g} A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) A_{it_2}^{1/2}(\beta_g^0) \varepsilon_{it_2} x_{it_1} \right\|^2 \right] \lesssim \sum_{i: g_i^0 = g} x_{it_1}^t x_{it_1} = O(n),$$

and

$$\begin{aligned}
& \sup_{\beta \in \mathcal{B}_{nT}} \left\| \sum_{i: g_i^0 = g} A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) \{m(x_{it_2}^t \beta_g^0) - m(x_{it_2}^t \beta_g)\} x_{it_1} \right\|^2 \\
& \leq \sup_{\beta \in \mathcal{B}_{nT}} \sum_{i: g_i^0 = g} \left\| A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) \dot{m}(\{x_{it_2}^t \beta_g\}^*) x_{it_2}^t (\beta_g^0 - \beta_g) x_{it_1} \right\|^2 \\
& \lesssim \sup_{\beta \in \mathcal{B}_{nT}} \sum_{i: g_i^0 = g} (\beta_g^0 - \beta_g) x_{it_2} x_{it_2}^t (\beta_g^0 - \beta_g) x_{it_1}^t x_{it_1} \\
& = O_p(n \lambda_{\min}^{-1}(\bar{H}^*) \tau).
\end{aligned}$$

Since it can be written as

$$\begin{aligned}
& \{\widehat{R}(\beta, \bar{g}) - \widehat{R}(\beta, \bar{g}^0)\}_{t_1, t_2} \\
& = \sum_{k=1}^T \sum_{l=1}^T \{\widehat{R}(\beta, \bar{g})\}_{t_1, k} \{\widehat{R}^{-1}(\beta, \bar{g}^0) - \widehat{R}^{-1}(\beta, \bar{g})\}_{kl} \{\widehat{R}(\beta, \bar{g}^0)\}_{l, t_2},
\end{aligned}$$

we have

$$\begin{aligned}
& \max_{1 \leq k, l \leq T} \{\widehat{R}^{-1}(\beta, \bar{g}^0) - \widehat{R}^{-1}(\beta, \bar{g})\}_{kl} \\
& = \{\widehat{R}(\beta, \bar{g}) - \widehat{R}(\beta, \bar{g}^0)\}_{t_1, t_2} / \sum_{k=1}^T \sum_{l=1}^T \{\widehat{R}(\beta, \bar{g})\}_{t_1, k} \{\widehat{R}(\beta, \bar{g}^0)\}_{l, t_2},
\end{aligned}$$

which implies that $\max_{1 \leq k, l \leq T} \{\widehat{R}^{-1}(\beta, \bar{g}^0) - \widehat{R}^{-1}(\beta, \bar{g})\}_{kl} = o_p(T^{-\delta})$ for $\bar{g} \in \Gamma$ from Assumption (A12) and the fact that the denominator in the right hand side of the above equality is at most of order $O_p(1)$. Then, we have $\sup_{\beta \in \mathcal{B}_{nT}} \|I_1\| = O_p(n^{1/2} T^2) o_p(T^{-\delta})$. For I_2 , we have from the triangle inequality

$$\|I_2\|^2 \leq \sum_{i=1}^n \mathbf{I}\{g_i \neq g_i^0\} \sum_{i=1}^n \|X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \bar{R}^{-1}(\beta, \bar{g}) A_i^{-1/2}(\beta_g) \{y_i - m(X_i \beta_g)\}\|^2$$

Since we have

$$\begin{aligned}
& \|X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \bar{R}^{-1}(\beta, \bar{g}) A_i^{-1/2}(\beta_g) \{y_i - m(X_i \beta_g)\}\|^2 \\
& \lesssim \lambda_{\max}(X_i^t X_i) \|y_i - m(X_i \beta_g)\|^2 = O_p(T^2),
\end{aligned}$$

we have $\sup_{\bar{g} \in \Gamma} \|I_2\| = O_p(nT)o_p(T^{-\delta})$, which ends the proof. \square

Lemma S3. *Under Assumptions (A1)-(A11), it holds that, for all $g = 1, \dots, G$,*

$$\sup_{\beta \in \mathcal{B}_{nT}} \|\{\bar{H}_g^*(\beta_g^0)\}^{-1/2}\{S_g(\beta_g) - \bar{S}_g(\beta_g)\}\| = O_p(\lambda_{\min}^{-1/2}(\bar{H}^*)T^2).$$

Proof. From Lemma S2, it is enough to show that

$$\|\{\bar{H}_g^*(\beta_g^0)\}^{-1/2}\{S_g^*(\beta_g^0) - \bar{S}_g^*(\beta_g^0)\}\| = O_p(\lambda_{\min}^{-1/2}(\bar{H}^*)T^2).$$

The proof is almost the same with that of Lemma 3.1 in Wang (2011). Let $Q = \{q_{j_1, j_2}\}_{1 \leq j_1, j_2 \leq T}$ denote the matrix $\hat{R}^{-1}(\beta, \bar{g}^0) - \bar{R}^{-1}(\beta, \bar{g}^0)$. Then,

$$\begin{aligned} & S_g^*(\beta_g) - \bar{S}_g^*(\beta_g) \\ &= \sum_{i=1}^n \sum_{t_1=1}^T \sum_{t_2=1}^T \{\hat{R}^{-1}(\beta, \bar{g}^0) - \bar{R}^{-1}(\beta, \bar{g}^0)\}_{t_1, t_2} A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) \{y_{it_2} - m(x_{it_2}^t \beta_g)\} x_{it_1} \\ &= \sum_{t_1=1}^T \sum_{t_2=1}^T \{\hat{R}^{-1}(\beta, \bar{g}^0) - \bar{R}^{-1}(\beta, \bar{g}^0)\}_{t_1, t_2} \\ & \quad \times \left[\sum_{i=1}^n A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) \{A_{it_2}^{1/2}(\beta_g^0) \varepsilon_{it_2} + m(x_{it_2}^t \beta_g^0) - m(x_{it_2}^t \beta_g)\} x_{it_1} \right] \end{aligned}$$

Note that

$$E \left[\left\| \sum_{i=1}^n A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) A_{it_2}^{1/2}(\beta_g^0) \varepsilon_{it_2} x_{it_1} \right\|^2 \right] \lesssim \sum_{i=1}^n x_{it_1}^t x_{it_1} = O(n),$$

and

$$\begin{aligned} & \sup_{\beta \in \mathcal{B}_{nT}} \left\| \sum_{i=1}^n A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) \{m(x_{it_2}^t \beta_g^0) - m(x_{it_2}^t \beta_g)\} x_{it_1} \right\|^2 \\ &= \sup_{\beta \in \mathcal{B}_{nT}} \sum_{i=1}^n \left\| A_{it_1}^{1/2}(\beta_g) A_{it_2}^{-1/2}(\beta_g) \dot{m}(\{x_{it_2}^t \beta_g\}^*) x_{it_2}^t (\beta_g^0 - \beta_g) x_{it_1} \right\|^2 \\ &\lesssim \sup_{\beta \in \mathcal{B}_{nT}} \sum_{i=1}^n (\beta_g^0 - \beta_g)^t x_{it_2} x_{it_2}^t (\beta_g^0 - \beta_g) x_{it_1}^t x_{it_1} \\ &= C^2 \lambda_{\min}^{-1}(\bar{H}^*) \tau O_p(n). \end{aligned}$$

Similar to the proof of Lemma S2, we have $\max_{1 \leq k, l \leq T} \{\widehat{R}^{-1}(\beta, \bar{g}^0) - \bar{R}^{-1}(\beta, \bar{g}^0)\}_{kl} = O_p(n^{-1/2})$ from Assumption (A6) (ii). Then, we have $\sup_{\beta \in \mathcal{B}_{nT}} \|S_g^*(\beta_g) - \bar{S}_g^*(\beta_g)\| = O_p(T^2)$, which proves the lemma. \square

The following Lemma is from Remark 1 in Xie and Yang (2003).

Lemma S4. *It holds that, for all $i = 1, \dots, n$,*

$$\overline{\mathcal{D}}_i(\beta_g) = \overline{H}_i(\beta_g) + \overline{B}_i(\beta_g) + \overline{\mathcal{E}}_i(\beta_g),$$

for $\overline{B}_i(\beta_g) = \overline{B}_i^{[1]}(\beta_g) + \overline{B}_i^{[2]}(\beta_g)$ and $\overline{\mathcal{E}}_i(\beta_g) = \overline{\mathcal{E}}_i^{[1]}(\beta_g) + \overline{\mathcal{E}}_i^{[2]}(\beta_g)$, where

$$\begin{aligned} \overline{B}_i^{[1]}(\beta_g) &= X_i^t \text{diag}[\overline{R}^{-1} A_i^{-1/2}(\beta_g) \{m(X_i \beta_{g_0}^0) - m(X_i \beta_g)\}] G_i^{[1]}(\beta_g) X_i, \\ \overline{B}_i^{[2]}(\beta_{g_i}) &= X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \overline{R}^{-1} \text{diag}[m(X_i \beta_{g_0}^0) - m(X_i \beta_g)] G_i^{[2]}(\beta_g) X_i, \\ \overline{\mathcal{E}}_i^{[1]}(\beta_g) &= X_i^t \text{diag}[\overline{R}^{-1} A_i^{-1/2}(\beta_g) A_i^{1/2}(\beta_{g_0}^0) \varepsilon_i] G_i^{[1]}(\beta_g) X_i, \end{aligned}$$

and

$$\overline{\mathcal{E}}_i^{[2]}(\beta_{g_i}) = X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \overline{R}^{-1} \text{diag}[A_i^{1/2}(\beta_{g_0}^0) \varepsilon_i] G_i^{[2]}(\beta_g) X_i.$$

Here, $G_i^{[\ell]}(\beta_g) = \text{diag}(q_{it}^{[\ell]}(\beta_g), \dots, q_{it}^{[\ell]}(\beta_g))$, for $\ell = 1, 2$, where

$$q_{it}^{[1]}(\beta_g) = [a''(\theta_{it})]^{-1/2} m'(\eta_{it}), \quad q_{it}^{[2]}(\beta_g) = [a''(\theta_{it})]^{-1/2},$$

and

$$q_{it}^{[1]}(\beta_g) = -\frac{1}{2} \frac{a^{(3)}(\theta_{it})}{[a''(\theta_{it})]^{5/2}} \{m'(\eta_{it})\}^2 + \frac{m''(\eta_{it})}{[a''(\theta_{it})]^{1/2}}, \quad q_{it}^{[2]}(\beta_{g_i}) = -\frac{1}{2} \frac{a^{(3)}(\theta_{it})}{[a''(\theta_{it})]^{5/2}} m'(\eta_{it}).$$

Lemma S5. *Under Assumptions (A1)-(A11), it holds that, for any $\lambda \in \mathbb{R}^p$ and $g = 1, \dots, G$,*

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [\mathcal{D}_g^*(\beta_g) - \overline{\mathcal{D}}_g^*(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\overline{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n).$$

Proof. By Lemma S4, it is sufficient to prove the following three results:

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [H_g^*(\beta_g) - \bar{H}_g^*(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n),$$

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [B_g^*(\beta_g) - \bar{B}_g^*(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n),$$

and

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [\mathcal{E}_g^*(\beta_g) - \bar{\mathcal{E}}_g^*(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n).$$

We have

$$\begin{aligned} |\lambda^t [H_g^*(\beta_g) - \bar{H}_g^*(\beta_g)] \lambda| &= \left| \sum_{i: g_i^0 = g} \lambda^t X_i^t \Delta_i(\beta_g^0) A_i^{1/2}(\beta_g^0) \hat{R}^{-1}(\beta, \bar{g}) \{ \hat{R}(\beta, \bar{g}) - \bar{R}(\beta, \bar{g}) \} \right. \\ &\quad \left. \times \bar{R}^{-1}(\beta, \bar{g}) A_i^{1/2}(\beta_g^0) \Delta_i(\beta_g^0) X_i \lambda \right| \\ &\lesssim \| \hat{R}(\beta, \bar{g}) - \bar{R}(\beta, \bar{g}) \|_F \lambda_{\max} \left(\sum_{i: g_i^0 = g} X_i^t X_i \right), \end{aligned}$$

which implies that $\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [H_g^*(\beta_g) - \bar{H}_g^*(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n)$ from Assumptions (A1) (i), (A10) and (A6) (ii). Next, we will verify

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [B_g^{[1]*}(\beta_g) - \bar{B}_g^{[1]*}(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n),$$

and

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [B_g^{[2]*}(\beta_g) - \bar{B}_g^{[2]*}(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n).$$

We have from Cauchy-Schwarz inequality

$$\begin{aligned}
& |\lambda^t [B_g^{[1]*}(\beta_g) - \bar{B}_g^{[1]*}(\beta_g)] \lambda| \\
&= \left| \sum_{i: g_i^0 = g} \lambda^t X_i^t \text{diag}[\{\widehat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} A_i^{-1/2}(\beta_g)] \right. \\
&\quad \left. \times \{m(X_i \beta_g^0) - m(X_i \beta_g)\} G_i^{[1]}(\beta_g) X_i \lambda \right| \\
&= \left| \sum_{i: g_i^0 = k} \lambda^t X_i^t G_i^{[1]}(\beta_g) \text{diag}[X_i \lambda] \{\widehat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} A_i^{-1/2}(\beta_g) \right. \\
&\quad \left. \times \{m(X_i \beta_g^0) - m(X_i \beta_g)\} \right| \\
&\leq \sum_{i: g_i^0 = g} \|\text{diag}[X_i \lambda] G_i^{[1]}(\beta_g) X_i \lambda\| \\
&\quad \times \|\{\widehat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} A_i^{-1/2}(\beta_g) \{m(X_i \beta_g^0) - m(X_i \beta_g)\}\|.
\end{aligned}$$

We have

$$\lambda^t X_i^t G_i^{[1]}(\beta_g) \text{diag}^2[X_i \lambda] G_i^{[1]}(\beta_g) X_i \lambda \leq \max_{1 \leq t \leq T} |x_{it}^t \lambda|^2 \max_{1 \leq t \leq T} |q_{it}^{[1]}(\beta_g)|^2 \lambda_{\max}(X_i^t X_i),$$

and, by using (S1), we have for β_g^* between β_g^0 and β_g ,

$$\begin{aligned}
& \{m(X_i \beta_g^0) - m(X_i \beta_g)\}^t A_i^{-1/2}(\beta_g) \{\widehat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\}^2 \\
& \quad \times A_i^{-1/2}(\beta_g) \{m(X_i \beta_g^0) - m(X_i \beta_g)\} \\
&= (\beta_g^0 - \beta_g)^t X_i^t \Delta(\beta_g^*) A_i(\beta_g^*) A_i^{-1/2}(\beta_g) [\widehat{R}^{-1}(\beta, \bar{g}) \{\bar{R}^{-1}(\beta, \bar{g}) - \widehat{R}^{-1}(\beta, \bar{g})\} \\
& \quad \times \bar{R}^{-1}(\beta, \bar{g})]^2 A_i^{-1/2}(\beta_g) A_i(\beta_g^*) \Delta(\beta_g^*) X_i (\beta_g^0 - \beta_g) \\
&\lesssim \|\widehat{R}(\beta, \bar{g}) - \bar{R}(\beta, \bar{g})\|_F^2 \lambda_{\max}(X_i^t X_i) \lambda_{\min}^{-1}(\bar{H}^*) \|\{\bar{H}_g^*(\beta_g^0)\}^{1/2} (\beta_g - \beta_g^0)\|.
\end{aligned}$$

Then, from Assumptions (A13) (ii) and (A10) we have

$$\begin{aligned}
& \sup_{\beta_k \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [B_{nk}^{[1]*}(\beta_k) - \bar{B}_{nk}^{[1]*}(\beta_k)] \lambda| \\
&= n O_p(T^{1/2}) O_p(\{T \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee T n^{-1/2}\}) O_p(T^{1/2}) \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \\
&= O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n) \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2},
\end{aligned}$$

which proves $\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [B_g^{[1]*}(\beta_g) - \bar{B}_g^{[1]*}(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n)$ since $\lambda_{\min}^{-1}(\bar{H}^*) \tau \rightarrow 0$. Moreover, we have from Cauchy-Schwarz inequality

$$\begin{aligned}
& |\lambda^t [B_g^{[2]*}(\beta_g) - \bar{B}_g^{[2]*}(\beta_g)] \lambda| \\
&= \left| \sum_{i: g_i^0 = g} \lambda^t X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \{\hat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} \right. \\
&\quad \left. \times \text{diag}[m(X_i \beta_g^0) - m(X_i \beta_g)] G_i^{[2]}(\beta_g) X_i \lambda \right| \\
&= \left| \sum_{i: g_i^0 = g} \lambda^t X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \{\hat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} G_i^{[2]}(\beta_g) \right. \\
&\quad \left. \times \text{diag}[X_i \lambda] \{m(X_i \beta_g^0) - m(X_i \beta_g)\} \right| \\
&\leq \sum_{i: g_i^0 = g} \|\text{diag}[X_i \lambda] G_i^{[2]}(\beta_g) \{\hat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} A_i^{1/2}(\beta_g) \Delta_i(\beta_g) X_i \lambda\| \\
&\quad \times \|m(X_i \beta_g^0) - m(X_i \beta_g)\|.
\end{aligned}$$

We have

$$\begin{aligned}
& \lambda^t X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \{\hat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} G_i^{[2]}(\beta_g) \text{diag}^2[X_i \lambda] G_i^{[2]}(\beta_g) \\
&\quad \times \{\hat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} A_i^{1/2}(\beta_g) \Delta_i(\beta_g) X_i \lambda \\
&\lesssim \max_{1 \leq t \leq T} |x_{it}^t \lambda|^2 \max_{1 \leq t \leq T} |q_{it}^{[2]}(\beta_g)|^2 \|\hat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\|_F^2 \lambda_{\max}(X_i^t X_i),
\end{aligned}$$

and for β_g^* between β_g and β_g^0 , we have

$$\begin{aligned}
\|m(X_i \beta_g^0) - m(X_i \beta_g)\|^2 &= (\beta_g^0 - \beta_g)^t X_i^t A_i(\beta_g^*) \Delta^2(\beta_g^*) A_i(\beta_g^*) X_i (\beta_g^0 - \beta_g) \\
&\lesssim \lambda_{\max}(X_i^t X_i) \lambda_{\min}^{-1}(\bar{H}^*) \|\{\bar{H}_g^*(\beta_g^0)\}^{1/2}(\beta_g - \beta_g^0)\|.
\end{aligned}$$

Then, from Assumption (A6) (ii) and (A10) we have

$$\begin{aligned}
& \sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [B_g^{[2]*}(\beta_g) - \bar{B}_g^{[2]*}(\beta_g)] \lambda| \\
&= n O_p(\{T \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee T n^{-1/2}\}) O_p(T^{1/2}) O_p(T^{1/2}) \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \\
&= O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n) \lambda_{\max}^{-1/2}(\bar{H}^*) \tau^{1/2},
\end{aligned}$$

which proves $\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [B_g^{[2]*}(\beta_g) - \bar{B}_g^{[2]*}(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n)$ since $\lambda_{\min}^{-1}(\bar{H}^*) \tau \rightarrow 0$. Lastly, we will verify

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [\mathcal{E}_g^{[1]*}(\beta_g) - \bar{\mathcal{E}}_g^{[1]*}(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n),$$

and

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [\mathcal{E}_g^{[2]*}(\beta_g) - \bar{\mathcal{E}}_g^{[2]*}(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n).$$

We have from Cauchy-Schwarz inequality

$$\begin{aligned}
& |\lambda^t [\mathcal{E}_g^{[1]*}(\beta_g) - \bar{\mathcal{E}}_g^{[1]*}(\beta_g)] \lambda| \\
&= \left| \sum_{i: g_i^0 = g} \lambda^t X_i^t G_i^{[1]}(\beta_g) \text{diag}[X_i \lambda] \{\hat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} A_i^{-1/2}(\beta_g) A_i^{1/2}(\beta_g^0) \varepsilon_i \right| \\
&\leq \sum_{i: g_i^0 = g} \|G_i^{[1]}(\beta_g) \text{diag}[X_i \lambda] X_i \lambda\| \cdot \|\{\hat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} A_i^{-1/2}(\beta_g) A_i^{1/2}(\beta_g^0) \varepsilon_i\| \\
&\lesssim \sum_{i: g_i^0 = g} \max_{1 \leq j \leq T} \{\|x_{it}^t \lambda\|\} \lambda_{\max}^{1/2}(X_i^t X_i) \|\hat{R}(\beta, \bar{g}) - \bar{R}(\beta, \bar{g})\|_F \|\varepsilon_i\|.
\end{aligned}$$

Then, from Assumption (A6) (ii) and (A10) we have we have

$$\begin{aligned}
& \sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [\mathcal{E}_{nk}^{[1]}(\beta_k) - \bar{\mathcal{E}}_{nk}^{[1]*}(\beta_k)] \lambda| \\
&= n O_p(T^{1/2}) O_p(\{T \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee T n^{-1/2}\}) O_p(T^{1/2}) \\
&= O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n),
\end{aligned}$$

which proves $\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [\mathcal{E}_g^{[1]*}(\beta_g) - \bar{\mathcal{E}}_g^{[1]*}(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n)$. Moreover, we have from Cauchy-Schwarz inequality

$$\begin{aligned}
& |\lambda^t [\mathcal{E}_g^{[2]*}(\beta_g) - \bar{\mathcal{E}}_g^{[2]*}(\beta_g)] \lambda| \\
&= \left| \sum_{i: g_i^0 = g} \lambda^t X_i^t \Delta_i(\beta_g) A_i^{1/2}(\beta_g) \{\widehat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} \text{diag}[A_i^{1/2}(\beta_{g_i^0}^0) \varepsilon_i] G_i^{[2]}(\beta_g) X_i \lambda \right| \\
&\leq \left(\sum_{i: g_i^0 = g} \|\widehat{R}^{-1}(\beta, \bar{g}) - \bar{R}^{-1}(\beta, \bar{g})\} A_i^{1/2}(\beta_g) \Delta_i(\beta_g) X_i \lambda \|^2 \right)^{1/2} \\
&\quad \times \left(\sum_{i: g_i^0 = g} \|\text{diag}[A_i^{1/2}(\beta_{g_i^0}^0) \varepsilon_i] G_i^{[2]}(\beta_g) X_i \lambda \|^2 \right)^{1/2} \\
&\lesssim \|\widehat{R}(\beta, \bar{g}) - \bar{R}(\beta, \bar{g})\|_F \max_{1 \leq j \leq T} \{ |A_{it}^{1/2}(\beta_{g_i^0}^0) \varepsilon_{it}| \} \lambda_{\max} \left(\sum_{i: g_i^0 = g} X_i^t X_i \right).
\end{aligned}$$

Then, from Assumption (A6) (ii) and (A10) we have

$$\begin{aligned}
& \sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [\mathcal{E}_g^{[2]*}(\beta_g) - \bar{\mathcal{E}}_g^{[2]*}(\beta_g)] \lambda| \\
&= O_p(\{T \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee T n^{-1/2}\}) O_p(nT) \\
&= O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n),
\end{aligned}$$

which proves $\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t [\mathcal{E}_g^{[2]*}(\beta_g) - \bar{\mathcal{E}}_g^{[2]*}(\beta_g)] \lambda| = O_p(\{\lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} \vee n^{-1/2}\} T^2 n)$. \square

The following three lemmas are from Lemma A.1. (ii), Lemma A.2. (ii), Lemma A.3. (ii) in Xie and Yang (2003), respectively. These three lemmas are hold under the assumption (AH) in Xie and Yang (2003), which is satisfied in our problem from Assumptions (A7) and (A8).

Lemma S6. *Suppose Assumption (A3) (i), (A7) and (A8) hold. It holds that, for any $\lambda \in \mathbb{R}^p$ and $g = 1, \dots, G$,*

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t \{\bar{H}_g^*(\beta_g^0)\}^{-1/2} \bar{H}_g^*(\beta_g) \{\bar{H}_g^*(\beta_g^0)\}^{-1/2} \lambda - 1| = o_p(1).$$

Lemma S7. *Suppose Assumptions (A3) (i), (A7) and (A8) hold. It holds that, for any $\lambda \in \mathbb{R}^p$ and $g = 1, \dots, G$,*

$$\sup_{\beta \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \overline{B}_g^*(\beta_g) \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \lambda| = o_p(1).$$

Lemma S8. *Suppose Assumptions (A3) (ii), (A7) and (A8) hold. It holds that, for any $\lambda \in \mathbb{R}^p$ and $g = 1, \dots, G$,*

$$\sup_{\beta_g \in \mathcal{B}_{nT}} \sup_{\|\lambda\|=1} |\lambda^t \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \overline{\mathcal{E}}_g^*(\beta_g) \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \lambda| = o_p(1).$$

The proof is based on that of Theorem 3.6 in Wang (2011). We will verify the following condition: for any $\epsilon > 0$, there exists a constant $C > 0$ such that for all n and T sufficiently large,

$$P\left(\sup_{\beta \in \mathcal{B}_{nT}, \bar{g} \in \Gamma} (\beta_g - \beta_g^0)^t S_g(\beta_g) < 0\right) \geq 1 - \epsilon,$$

where $\mathcal{B}_{nT} = \{\beta : \max_{g=1, \dots, G} \|\{\overline{H}_g^*(\beta_g^0)\}^{1/2}(\beta_g - \beta_g^0)\| = C\tau^{1/2}\}$ and $\Gamma = \{\bar{g} = (g_1, \dots, g_n) : \sup_{\beta \in \mathcal{B}_{nT}} n^{-1} \sum_{i=1}^n \mathbf{I}\{g_i \neq g_i^0\} = o_p(T^{-\delta}) \text{ for all } \delta > 0\}$. This is a sufficient condition to ensure the existence of a sequence of roots $\hat{\beta}_g$ of the equation $S_g(\beta_g) = 0$ for $g = 1, \dots, G$ such that $\hat{\beta} \in \mathcal{B}_{nT}$ for $\bar{g} \in \Gamma$. Then, Theorem follows from Lemma S1.

From Taylor expansion, we can write

$$\begin{aligned} (\beta_g - \beta_g^0)^t S_g(\beta_g) &= (\beta_g - \beta_g^0)^t S_g(\beta_g^0) - (\beta_g - \beta_g^0)^t \sum_{i=1}^n \mathbf{I}\{g_i = g\} \mathcal{D}_i(\beta_{g_i}^*) (\beta_{g_i} - \beta_g^0) \\ &\equiv I_1 + I_2, \end{aligned}$$

where $\beta_{g_i}^*$ lies between β_{g_i} and β_g^0 for $i = 1, \dots, n$. Next, we write

$$I_1 = (\beta_g - \beta_g^0)^t \overline{S}_g^*(\beta_g^0) + (\beta_g - \beta_g^0)^t \{S_g(\beta_g^0) - \overline{S}_g^*(\beta_g^0)\} \equiv I_{11} + I_{12}.$$

For $\ell = 1, \dots, p$, denote $e_\ell \in \mathbb{R}^p$ with ℓ th element equal to 1 and the others equal to

0. Then, we have

$$\begin{aligned}
& E[\{e_\ell^t \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \overline{S}_g^*(\beta_g^0)\}^2] \\
&= e_\ell^t \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \sum_{i=1}^n \mathbb{I}\{g_i^0 = g\} X_i^t \Delta_i(\beta_g^0) A_i^{1/2}(\beta_g^0) \overline{R}^{-1}(\beta^0, \overline{g}^0) R^0 \overline{R}^{-1}(\beta^0, \overline{g}^0) \\
&\quad \times A_i^{1/2}(\beta_g^0) \Delta_i(\beta_g^0) X_i \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} e_\ell \\
&\leq \lambda_{\max}(R^0 \overline{R}^{-1}(\beta^0, \overline{g}^0)).
\end{aligned}$$

Thus, we can bound $|I_{11}|$ by

$$\sup_{\beta \in \mathcal{B}_{nT}} |I_{11}| \leq \|\{\overline{H}_g^*(\beta_g^0)\}^{1/2}(\beta_g - \beta_g^0)\| \cdot \|\{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \overline{S}_g^*(\beta_g^0)\| \leq C\tau.$$

From the Lemma S2 and S3, we have

$$\begin{aligned}
\sup_{\beta \in \mathcal{B}_{nT}} |I_{12}| &\leq \|\{\overline{H}_g^*(\beta_g^0)\}^{1/2}(\beta_g - \beta_g^0)\| \cdot \|\{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \{S_g(\beta_g^0) - \overline{S}_g^*(\beta_g^0)\}\| \\
&\leq \tau^{1/2} O_p(\lambda_{\min}^{-1/2}(\overline{H}^*) T^2).
\end{aligned}$$

Since $\tau^{-1/2} \lambda_{\min}^{-1/2}(\overline{H}^*) T^2 \rightarrow 0$ from Assumption (A2), $\sup_{\beta \in \mathcal{B}_{nT}} |I_{12}| = o_p(\tau)$. Hence, we have $\sup_{\beta \in \mathcal{B}_{nT}} |I_1| \leq C\tau$. In what follows, we will evaluate I_2 . It can be written as

$$\begin{aligned}
I_2 &= -(\beta_g - \beta_g^0)^t \sum_{i=1}^n \mathbb{I}\{g_i = g\} \overline{\mathcal{D}}_i(\beta_{g_i}^*) (\beta_{g_i} - \beta_g^0) \\
&\quad - (\beta_g - \beta_g^0)^t \sum_{i=1}^n \mathbb{I}\{g_i = g\} \{\mathcal{D}_i(\beta_{g_i}^*) - \overline{\mathcal{D}}_i(\beta_{g_i}^*)\} (\beta_{g_i} - \beta_g^0) \\
&\equiv I_{21} + I_{22}.
\end{aligned}$$

For $g_i^0 = g_i = g$, $\beta_{g_i}^*$ lies between β_g and β_g^0 , and then we write $\beta_{g_i}^* \equiv \beta_g^*$ for such i .

Hence, we can write

$$\begin{aligned}
I_{21} &= -(\beta_g - \beta_g^0)^t \overline{\mathcal{D}}_g^*(\beta_g^*)(\beta_g - \beta_g^0) \\
&\quad - (\beta_g - \beta_g^0)^t \sum_{i=1}^n (\mathbf{I}\{g_i = g\} - \mathbf{I}\{g_i^0 = g\}) \overline{\mathcal{D}}_i(\beta_{g_i}^*)(\beta_{g_i} - \beta_g^0) \\
&\equiv I_{211} + I_{212}.
\end{aligned}$$

For I_{211} , we write

$$\begin{aligned}
I_{211} &= -(\beta_g - \beta_g^0)^t \overline{H}_g^*(\beta_g^*)(\beta_g - \beta_g^0) - (\beta_g - \beta_g^0)^t \{\overline{\mathcal{D}}_g^*(\beta_g^*) - \overline{H}_g^*(\beta_g^*)\}(\beta_g - \beta_g^0) \\
&\equiv I_{2111} + I_{2112}.
\end{aligned}$$

For I_{2111} , we can write

$$\begin{aligned}
I_{2111} &= -(\beta_g - \beta_g^0)^t \overline{H}_g^*(\beta_g^0)(\beta_g - \beta_g^0) \\
&\quad - (\beta_g - \beta_g^0)^t \{\overline{H}_g^*(\beta_g^0)\}^{1/2} \left[\{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \overline{H}_g^*(\beta_g^*) \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} - I_p \right] \\
&\quad \times \{\overline{H}_g^*(\beta_g^0)\}^{1/2} (\beta_g - \beta_g^0) \\
&\equiv I_{21111} + I_{21112}.
\end{aligned}$$

For $\beta \in \mathcal{B}_{nT}$, we have $I_{21111} = -C^2\tau$. Moreover, for $g_i^0 = g_i = g$, $\beta_{g_i}^* \equiv \beta_g^*$ is contained in a local neighborhood of β_g^0 . Then, for I_{21112} , we have from Lemma S6,

$$\begin{aligned}
|I_{21112}| &\leq \sup_{\beta \in \mathcal{B}_{nT}} \max \left\{ \left| \lambda_{\min} \left(\left[\{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \overline{H}_g^*(\beta_g^*) \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} - I_p \right] \right) \right|, \right. \\
&\quad \left. \left| \lambda_{\max} \left(\left[\{\overline{H}_g^*(\beta_g^0)\}^{-1/2} \overline{H}_g^*(\beta_g^*) \{\overline{H}_g^*(\beta_g^0)\}^{-1/2} - I_p \right] \right) \right| \right\} \\
&\quad \times \|\{\overline{H}_g^*(\beta_g^0)\}^{1/2} (\beta_g - \beta_g^0)\|^2 \\
&= o(1)C^2\tau,
\end{aligned}$$

which is dominated by I_{21111} . Hence, for $\beta \in \mathcal{B}_{nT}$ we have $I_{2111} = -C^2\tau$. Next, we

verify I_{2112} . For $g_i^0 = g_i = g$, we have from Lemma S4, S7 and S8

$$\begin{aligned}
|I_{2112}| &= |(\beta_g - \beta_g^0)^t \{\bar{B}_g^*(\beta_g^*) + \bar{\mathcal{E}}_g^*(\beta_g^*)\} (\beta_g - \beta_g^0)| \\
&\leq \sup_{\beta \in \mathcal{B}_{nT}} \{ \lambda_{\max}(\{\bar{H}_g^*(\beta_g^0)\}^{-1/2} \bar{B}_g^*(\beta_g^*) \{\bar{H}_g^*(\beta_g^0)\}^{-1/2}) \\
&\quad + \lambda_{\max}(\{\bar{H}_g^*(\beta_g^0)\}^{-1/2} \bar{\mathcal{E}}_g^*(\beta_g^*) \{\bar{H}_g^*(\beta_g^0)\}^{-1/2}) \| \{\bar{H}_g^*(\beta_g^0)\}^{1/2} (\beta_g - \beta_g^0) \|^2 \\
&= o(1) C^2 \tau,
\end{aligned}$$

which is dominated by I_{2111} . Hence, for $\beta \in \mathcal{B}_{nT}$ we have $I_{211} = -C^2 \tau$. Next, we verify I_{212} .

$$\begin{aligned}
|I_{212}| &\leq \left| (\beta_g - \beta_g^0)^t \sum_{i=1}^n (\mathbb{I}\{g_i = g\} - \mathbb{I}\{g_i^0 = g\}) \bar{H}_i(\beta_{g_i}^*) (\beta_{g_i} - \beta_{g_i^0}^0) \right| \\
&\quad + \left| (\beta_g - \beta_g^0)^t \sum_{i=1}^n (\mathbb{I}\{g_i = g\} - \mathbb{I}\{g_i^0 = g\}) \bar{B}_i(\beta_{g_i}^*) (\beta_{g_i} - \beta_{g_i^0}^0) \right| \\
&\quad + \left| (\beta_g - \beta_g^0)^t \sum_{i=1}^n (\mathbb{I}\{g_i = g\} - \mathbb{I}\{g_i^0 = g\}) \bar{\mathcal{E}}_i(\beta_{g_i}^*) (\beta_{g_i} - \beta_{g_i^0}^0) \right| \\
&\equiv I_{2121} + I_{2122} + I_{2123}.
\end{aligned}$$

From Cauchy-Schwarz inequality, $\beta \in \mathcal{B}_{nT}$ we have

$$\begin{aligned}
|I_{2121}| &\lesssim \lambda_{\min}^{-1/2}(\bar{H}^*) \| \{\bar{H}_g^*(\beta_g^0)\}^{1/2} (\beta_g - \beta_g^0) \| \sum_{i=1}^n \mathbb{I}\{g_i \neq g_i^0\} n \{ \max_{1 \leq i \leq n} \sup_{\beta \in \mathcal{B}} \lambda_{\max}(\bar{H}_i(\beta_{g_i})) \} \\
&\lesssim C \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} n \left(\frac{1}{n} \sum_{i=1}^n \mathbb{I}\{g_i \neq g_i^0\} \right) n \{ \max_{1 \leq i \leq n} \sup_{\beta \in \mathcal{B}} \lambda_{\max}(\bar{H}_i(\beta_{g_i})) \}.
\end{aligned}$$

From Assupmtions (A7), (A8) and (A9), for $i = 1, \dots, n$ we have

$$\max_{\beta \in \mathcal{B}} \{ \lambda_{\max}(\bar{H}_i(\beta_{g_i})) \} \lesssim \max_{\beta \in \mathcal{B}} \max_{t=1, \dots, T} [a''(\theta_{it}(\beta_{g_i})) \{u'(x_{it}^t \beta_{g_i})\}^2] \lambda_{\max}(X_i^t X_i) = O_p(T),$$

for $\beta_{g_i}^*$ between $\beta_{g_i^0}^0$ and β_{g_i} , which implies that

$$\sup_{\beta \in \mathcal{B}_{nT}, \bar{g} \in \Gamma} |I_{2121}| = C \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} n^2 T o_p(T^{-\delta}) = o_p(\tau).$$

Similarly, Cauchy-Schwarz inequality we have

$$\begin{aligned} \sup_{\beta \in \mathcal{B}_{nT}, \bar{g} \in \Gamma} |I_{2122}| &\lesssim C \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} n \left(\frac{1}{n} \sum_{i=1}^n \mathbb{I}\{g_i \neq g_i^0\} \right) \\ &\quad \times n \left[\left\{ \max_{1 \leq i \leq n} \sup_{\beta \in \mathcal{B}} \|\bar{B}_i^{[1]}(\beta_{g_i})\|_F \right\} + \left\{ \max_{1 \leq i \leq n} \sup_{\beta \in \mathcal{B}} \|\bar{B}_i^{[2]}(\beta_{g_i})\|_F \right\} \right]. \end{aligned}$$

It is noted that we have from Cauchy-Schwarz inequality

$$\begin{aligned} &\{\bar{B}_i^{[1]}(\beta_{g_i})\}_{jk} \\ &= e_j^t X_i^t \text{diag}[\bar{R}^{-1}(\beta, \bar{g}) A_i^{-1/2}(\beta_{g_i}) \{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_{g_i})\}] G_i^{[1]}(\beta_{g_i}) X_i e_k \\ &\leq \lambda_{\max}(X_i^t X_i) \lambda_{\max}(\text{diag}[\bar{R}^{-1}(\beta, \bar{g}) A_i^{-1/2}(\beta_{g_i}) \{m(X_i \beta_{g_i^0}^0) - m(X_i \beta_{g_i})\}]) \lambda_{\max}(G_i^{[1]}(\beta_{g_i})) \\ &= \lambda_{\max}(X_i^t X_i) \max_{1 \leq k \leq T} \left\{ \sum_{t=1}^T \{\bar{R}^{-1}(\beta, \bar{g})\}_{kj} A_{it}^{-1/2}(\beta_{g_i}) \{m(x_{it}^t \beta_{g_i^0}^0) - m(x_{it}^t \beta_{g_i})\} \right\} \\ &\quad \times \lambda_{\max}(G_i^{[1]}(\beta_{g_i})) \\ &= O_p(T^2). \end{aligned}$$

Similarly $\{\bar{B}_i^{[2]}(\beta_{g_i})\}_{jk} = O_p(T^2)$, then we have

$$\sup_{\beta \in \mathcal{B}_{nT}, \bar{g} \in \Gamma} |I_{2122}| = \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} n o_p(T^{-\delta}) n T^{5/2} = o_p(\tau).$$

Similarly, we have

$$\begin{aligned} \sup_{\beta \in \mathcal{B}_{nT}} |I_{2123}| &\lesssim C \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} n \left(\frac{1}{n} \sum_{i=1}^n \mathbb{I}\{g_i \neq g_i^0\} \right) \\ &\quad \times n \left[\left\{ \max_{1 \leq i \leq n} \sup_{\beta \in \mathcal{B}} \|\bar{\mathcal{E}}_i^{[1]}(\beta_{g_i})\|_F \right\} + \left\{ \max_{1 \leq i \leq n} \sup_{\beta \in \mathcal{B}} \|\bar{\mathcal{E}}_i^{[2]}(\beta_{g_i})\|_F \right\} \right]. \end{aligned}$$

It is noted that we have

$$\begin{aligned}
& E[\|\bar{\mathcal{E}}_i^{[1]}(\beta_{g_i})\|_F^2] \\
&= \sum_{\ell=1}^T E[e_\ell^t \bar{\mathcal{E}}_i^{[1]}(\beta_{g_i})^t \bar{\mathcal{E}}_i^{[1]}(\beta_{g_i}) e_\ell] \\
&= \sum_{\ell=1}^T E \left[\varepsilon_i^t A_i^{1/2}(\beta_{g_i^0}) A_i^{-1/2}(\beta_{g_i}) \bar{R}^{-1}(\beta, \bar{g}) \text{diag}[X_i e_\ell] G_i^{[1]}(\beta_{g_i}) X_i \right. \\
&\quad \left. \times X_i^t G_i^{[1]}(\beta) \text{diag}[X_i e_\ell] \bar{R}^{-1}(\beta, \bar{g}) A_i^{-1/2}(\beta_{g_i}) A_i^{1/2}(\beta_{g_i^0}) \varepsilon_i \right] \\
&\leq \sum_{\ell=1}^T \lambda_{\max}(X_i^t X_i) \max_{1 \leq i \leq n, 1 \leq t \leq T} \max_{\beta \in \mathcal{B}} |q'_{it}^{[1]}(\beta_{g_i})| \max_{1 \leq i \leq n, 1 \leq t \leq T} |x_{it}^t e_\ell|^2 \\
&\quad \times \max_{\beta \in \mathcal{B}} \left\{ \max_{1 \leq i \leq n, 1 \leq t \leq T} A_{it}^{-1}(\beta_{g_i}) A_{it}(\beta_{g_i^0}) \right\} E[\varepsilon_i^t \varepsilon_i] \\
&= O(T^3),
\end{aligned}$$

which implies that $\|\bar{\mathcal{E}}_i^{[1]}(\beta_{g_i})\|_F = O_p(T^{3/2})$. Similarly $\|\bar{\mathcal{E}}_i^{[2]}(\beta_{g_i})\|_F = O_p(T^{3/2})$, then we have

$$\sup_{\beta \in \mathcal{B}_{nT}, \bar{g} \in \Gamma} |I_{2123}| = \lambda_{\min}^{-1/2}(\bar{H}^*) \tau^{1/2} n o_p(T^{-\delta}) n T^{3/2} = o_p(\tau).$$

Thus, I_{2121} , I_{2122} and I_{2123} are dominated by I_{211} for $\beta \in \mathcal{B}_{nT}$ and $\bar{g} \in \Gamma$. Hence $I_{21} = -C^2 \tau$ for $\beta \in \mathcal{B}_{nT}$ and $\bar{g} \in \Gamma$. Lastly, we verify I_{22} . We can write

$$\begin{aligned}
I_{22} &= -(\beta_g - \beta_g^0)^t \sum_{i=1}^n \mathbb{I}\{g_i = g\} \mathbb{I}\{g_i = g_i^0\} \{\mathcal{D}_i(\beta_{g_i}^*) - \bar{\mathcal{D}}_i(\beta_{g_i}^*)\} (\beta_{g_i} - \beta_g^0) \\
&\quad - (\beta_g - \beta_g^0)^t \sum_{i=1}^n \mathbb{I}\{g_i = g\} \mathbb{I}\{g_i \neq g_i^0\} \{\mathcal{D}_i(\beta_{g_i}^*) - \bar{\mathcal{D}}_i(\beta_{g_i}^*)\} (\beta_{g_i} - \beta_g^0) \\
&\equiv I_{221} + I_{222}.
\end{aligned}$$

For I_{221} , we can write, from Lemma S5,

$$\begin{aligned} |I_{221}| &\leq \sup_{\beta \in \mathcal{B}_{nT}} \max\{|\lambda_{\max}(D\tilde{S}_g(\beta_g) - \overline{\mathcal{D}}_g^*(\beta_g))|, |\lambda_{\min}(D\tilde{S}_g(\beta_g) - \overline{\mathcal{D}}_g^*(\beta_g))|\} \\ &\quad \times \lambda_{\min}^{-1}(\overline{H}^*) \|\{\overline{H}_g^*(\beta_g^0)\}^{1/2}(\beta_g - \beta_g^0)\|^2 \\ &= O_p(\{\lambda_{\min}^{-1/2}(\overline{H}^*)\tau^{1/2} \vee n^{-1/2}\}T^2n)C^2\lambda_{\min}^{-1}(\overline{H}_g^*)\tau. \end{aligned}$$

Since $\lambda_{\min}(\overline{H}^*)$ is at least of order larger than $O_p(nT)$, and from definition, we have $\tau = \sup_{\beta \in \mathcal{B}, \bar{g}} \lambda_{\max}(\{\overline{R}(\beta, \bar{g})\}^{-1}R^0) \leq \sup_{\beta \in \mathcal{B}, \bar{g}} \lambda_{\max}(\{\overline{R}(\beta, \bar{g})\}^{-1})\lambda_{\max}(R^0) \leq O_p(T)$ from Assumption (A5), the order of $\tau\lambda_{\min}^{-2}(\overline{H}^{orac})n^2$ is at most $O_p(T^{-1})$. Then, from Assumption (A2) we have $\sup_{\beta \in \mathcal{B}_{nT}} |I_{221}| = \tau o_p(1)$. As for I_{222} , we have

$$\begin{aligned} |I_{222}| &\leq \lambda_{\min}^{-1/2}(\overline{H}^*) \|\{\overline{H}_g^*(\beta_g^0)\}^{1/2}(\beta_g - \beta_g^0)\| \\ &\quad \times \sum_{i=1}^n \mathbf{I}\{g_i = g\} \mathbf{I}\{g_i \neq g_i^0\} \cdot \|\mathcal{D}_i(\beta_{g_i}^*) - \overline{\mathcal{D}}_i(\beta_{g_i}^*)\|_F \cdot \|\beta_{g_i} - \beta_g^0\| \\ &\leq \lambda_{\min}^{-1/2}(\overline{H}^*)\tau n \left(\sum_{i=1}^n \mathbf{I}\{g_i \neq g_i^0\} \right) \sum_{i=1}^n \|\mathcal{D}_i(\beta_{g_i}^*) - \overline{\mathcal{D}}_i(\beta_{g_i}^*)\|_F \cdot \|\beta_{g_i} - \beta_g^0\|. \end{aligned}$$

It is noted that the order of $\|\mathcal{D}_i(\beta_{g_i}) - \overline{\mathcal{D}}_i(\beta_{g_i})\|_F$ is at most $O_p(T)$. Then, from Lemma S1, $\sup_{\beta \in \mathcal{B}_{nT}, \bar{g} \in \Gamma} |I_{222}| = o_p(T^{-\delta})$, which implies that I_{22} is dominated by I_{21} . Thus, $(\beta_g - \beta_g^0)^t S_g(\beta_g)$ on $\beta \in \mathcal{B}_{nT}$ and $\bar{g} \in \Gamma$ is asymptotically dominated in probability by $I_{11} + I_{21} = C\tau - C^2\tau$, which is negative for C large enough, which proves the first part of the Theorem.

Next, we show the second part of the Theorem. We have

$$\begin{aligned} &P\left(\max_{1 \leq i \leq n} |\widehat{g}_i(\widehat{\beta}) - g_i^0| > 0\right) \\ &\leq G \max_{1 \leq g \leq G} P(\widehat{\beta}_g \notin \mathcal{B}_{nT}) + n \max_{1 \leq i \leq n} P(\widehat{\beta}_g \in \mathcal{B}_{nT}, \widehat{g}_i(\widehat{\beta}) \neq g_i^0). \end{aligned}$$

The order of the first term is $o(1)$ from the first part of the Theorem. We have

$\sup_{\beta \in \mathcal{B}_{nT}} \mathbb{I}\{\widehat{g}_i(\beta) \neq g_i^0\} \leq \sum_{g=1}^G \widetilde{Z}_{ig}$. Then,

$$\begin{aligned} \max_{1 \leq i \leq n} P\left(\widehat{\beta}_g \in \mathcal{B}_{nT}, \widehat{g}_i(\widehat{\beta}) \neq g_i^0\right) &= \max_{1 \leq i \leq n} E[\mathbb{I}\{\widehat{\beta}_g \in \mathcal{B}_{nT}\} \mathbb{I}\{\widehat{g}_i \neq g_i^0\}] \\ &\leq \max_{1 \leq i \leq n} E\left[\mathbb{I}\{\widehat{\beta}_g \in \mathcal{B}_{nT}\} \sum_{g=1}^G \widetilde{Z}_{ig}\right] \\ &\leq \max_{1 \leq i \leq n} \sum_{g=1}^G P(\widetilde{Z}_{ig} = 1) = o(T^{-\delta}), \end{aligned}$$

which proves the Theorem.

S2 Proof of Theorem 2

In order to show Theorem 2, we need to show the next lemmas.

Let $\widetilde{\beta}_g$ denote a root of $\overline{S}_g^*(\beta_g) = 0$. The next result shows that the group GEE estimator and the infeasible estimator with known population groups are asymptotically equivalent.

Lemma S9. *Suppose the Assumptions (A1)-(A11) hold. As n and T tend to infinity, we have $\widehat{\beta}_g = \widetilde{\beta}_g + o_p(1)$ for $g = 1, \dots, G$.*

Proof. We have

$$\begin{aligned} &\sup_{\beta \in \mathcal{B}_{nT}, \overline{g} \in \Gamma} \|S_g(\beta_g) - \overline{S}_g^*(\beta_g)\| \\ &\leq \sup_{\beta \in \mathcal{B}_{nT}} \|S_g(\beta_g) - \overline{S}_g(\beta_g)\| + \sup_{\beta \in \mathcal{B}_{nT}, \overline{g} \in \Gamma} \|\overline{S}_g(\beta_g) - \overline{S}_g^*(\beta_g)\|. \end{aligned}$$

Then, we have $\sup_{\beta \in \mathcal{B}_{nT}, \overline{g} \in \Gamma} \|S_g(\beta_g) - \overline{S}_g^*(\beta_g)\| = O_p(T^2)$ from Lemmas S2 and S3. Since $\widehat{\beta}_g \in \mathcal{B}_{nT}$ for $\overline{g} \in \Gamma$ from Theorem 1 and $\widetilde{\beta}_g \in \mathcal{B}_{nT}$ from Theorem 2 in Xie and Yang (2003), this implies

$$\sup_{\overline{g} \in \Gamma} |(\widehat{\beta}_g - \widetilde{\beta}_g)^t \{S_g(\widehat{\beta}_g) - \overline{S}_g^*(\widehat{\beta}_g)\}| = |(\widehat{\beta}_g - \widetilde{\beta}_g)^t \overline{S}_g^*(\widehat{\beta}_g)| = O_p(T^2).$$

From Taylor expansion, for β_g^* between $\widehat{\beta}_g$ and $\widetilde{\beta}_g$ we have

$$\begin{aligned}\overline{S}_g^*(\widehat{\beta}_g) &= \overline{S}_g^*(\widetilde{\beta}_g) - \overline{\mathcal{D}}_g^*(\beta_g^*)(\widehat{\beta}_g - \widetilde{\beta}_g) \\ &= -\overline{H}_g^*(\beta_g^*)(\widehat{\beta}_g - \widetilde{\beta}_g) - \{\overline{\mathcal{D}}_g^*(\beta_g^*) - \overline{H}_g^*(\beta_g^*)\}(\widehat{\beta}_g - \widetilde{\beta}_g).\end{aligned}$$

Then, we have, from Lemmas S6 - S8,

$$|(\widehat{\beta}_g - \widetilde{\beta}_g)^t \{S_g(\widehat{\beta}_g) - \overline{S}_g^*(\widehat{\beta}_g)\}| = (\widehat{\beta}_g - \widetilde{\beta}_g)^t \overline{H}_g^*(\beta_g^*)(\widehat{\beta}_g - \widetilde{\beta}_g) + o_p(1).$$

Hence, we have

$$\sup_{\overline{g} \in \Gamma} \inf_{\beta \in \mathcal{B}_{nT}} \lambda_{\min}(\overline{H}_g^*(\beta)) \|\widehat{\beta}_g - \widetilde{\beta}_g\|^2 \leq O_p(T^2) + o_p(1),$$

which implies $\|\widehat{\beta}_g - \widetilde{\beta}_g\| = o_p(1)$, since the order of $\lambda_{\min}(\overline{H}_g^*(\beta))$ is at least $O_p(nT)$.

The Lemma follows from Lemma S1. \square

Next lemma is almost the same with Lemma 2 in Xie and Yang (2003).

Lemma S10. *Suppose the Assumptions (A11)-(A11) hold. Moreover, suppose that, for all $g = 1, \dots, G$, there exists a constant ζ such that $(c^*T)^{1+\zeta}\gamma^* \rightarrow 0$ as $n \rightarrow \infty$. Moreover, suppose the marginal distribution of each individual observation has a density of the form from (1). Then, when $n \rightarrow \infty$, we have*

$$\{\overline{M}_g^*(\beta_g^0)\}^{-1/2} \overline{S}_g^*(\beta_g^0) \rightarrow N(0, I_p) \quad \text{in distribution.}$$

Proof. Let $\lambda^t \{\overline{M}_g^*(\beta_g^0)\}^{-1/2} \overline{S}_g^*(\beta_g^0) = \sum_{i:g_i^0=g} Z_{nTi}$, where, for any $p \times 1$ vector λ such that $\|\lambda\| = 1$, $Z_{nTi} = \lambda^t \{\overline{M}_g^*(\beta_g^0)\}^{-1/2} X_i^t \Delta_i(\beta_g^0) A_i^{1/2}(\beta_g^0) \overline{R}_i^{-1}(\beta^0, \overline{g}^0) \varepsilon_i$. To establish the asymptotic normality, it suffices to check the Lindeberg condition, that is, for any $\epsilon > 0$,

$$\sum_{i:g_i^0=g} E[Z_{nTi}^2 \mathbf{I}\{|Z_{nTi}| > \epsilon\}] \rightarrow 0,$$

which is shown in the proof of Lemma 2 in Xie and Yang (2003). \square

We will show

$$\{\overline{M}_g^*(\beta_g^0)\}^{-1/2}\overline{H}_g^*(\beta_g^0)(\tilde{\beta}_g - \beta_g^0) \rightarrow N(0, I_p) \quad \text{in distribution.}$$

The theorem follows from Lemma S9.

For $\beta_g^* \in \mathcal{B}_{nT}$ between $\tilde{\beta}_g$ and β_g^0 , we have

$$\begin{aligned} & \{\overline{H}_g^*(\beta_g^0)\}^{-1/2}\overline{S}_g^*(\beta_g^0) \\ &= -\{\overline{H}_g^*(\beta_g^0)\}^{1/2}(\tilde{\beta}_g - \beta_g^0) \\ & \quad - \left[\{\overline{H}_g^*(\beta_g^0)\}^{-1/2}\overline{\mathcal{D}}_g^*(\beta_g^*)\{\overline{H}_g^*(\beta_g^0)\}^{-1/2} - I_p \right] \{\overline{H}_g^*(\beta_g^0)\}^{1/2}(\tilde{\beta}_g - \beta_g^0). \end{aligned}$$

From Lemmas S4 and S6 - S8, the second term in the right hand side of the above equation is $o_p(1)$, which implies that $\{\overline{M}_g^*(\beta_g^0)\}^{-1/2}\overline{S}_g^*(\beta_g^0)$ and $\{\overline{M}_g^*(\beta_g^0)\}^{-1/2}\overline{H}_g^*(\beta_g^0)(\tilde{\beta}_g - \beta_g^0)$ are asymptotically identically distributed. Hence, the theorem follows from Lemma S10.

S3 Property of $\overline{R}^*(\beta, \overline{g})$

In this section, we denote the estimated unstructured working correlation matrix as $\widehat{R}^*(\beta, \overline{g}) = R^*(\widehat{\alpha}(\beta, \overline{g}), \beta, \overline{g})$ for $\widehat{\alpha}(\beta, \overline{g})$ given in (4). Then, it follows that

$$\begin{aligned} \overline{R}^*(\beta, \overline{g}) &= \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}) A_i^{1/2}(\beta_{g_i}^0) R^0 A_i^{1/2}(\beta_{g_i}^0) A_i^{-1/2}(\beta_{g_i}) \\ & \quad + \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}) \{m(X_i \beta_{g_i}^0) - m(X_i \beta_{g_i})\} \{m(X_i \beta_{g_i}^0) - m(X_i \beta_{g_i})\}^t A_i^{-1/2}(\beta_{g_i}). \end{aligned}$$

The next lemma shows that $\overline{R}^*(\beta, \overline{g})$ satisfies Assumption (A5) (ii).

Lemma S11. *Under Assumptions (A1)-(A5) (i) and (A7)-(A10), it holds that for any \overline{g} , $\lambda_{\max}(\{\overline{R}^{mo}(\beta^0, \overline{g})\}^{-2} R^0) = O_p(1)$.*

Proof. Since the eigenvalues of $\overline{R}^*(\beta^0, \overline{g})(R^0)^{-1/2}$ and $(R^0)^{-1/4}\overline{R}^*(\beta^0, \overline{g})(R^0)^{-1/4}$ are the same, we will show that $\lambda_{\min}((R^0)^{-1/4}\overline{R}^*(\beta^0, \overline{g})(R^0)^{-1/4})$ is bounded away from

zero. It can be written as

$$\begin{aligned}
& \lambda_{\min}((R^0)^{-1/4}\bar{R}^*(\beta^0, \bar{g})(R^0)^{-1/4}) \\
& \geq \lambda_{\min}\left((R^0)^{-1/4}\frac{1}{n}\sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0)A_i^{1/2}(\beta_{g_i^0}^0)R^0 A_i^{1/2}(\beta_{g_i^0}^0)A_i^{-1/2}(\beta_{g_i}^0)(R^0)^{-1/4}\right) \\
& \quad + \lambda_{\min}\left((R^0)^{-1/2}\frac{1}{n}\sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0)\{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_{g_i}^0)\}\right. \\
& \quad \quad \quad \left. \times \{m(X_i\beta_{g_i^0}^0) - m(X_i\beta_{g_i}^0)\}^t A_i^{-1/2}(\beta_{g_i}^0)(R^0)^{-1/2}\right).
\end{aligned}$$

Since the smallest eigenvalue does not diverge to infinity, it is enough to show that the first term of the right hand side of the above inequality is bounded away from zero. Then, we have

$$\begin{aligned}
& \lambda_{\min}\left((R^0)^{-1/4}\frac{1}{n}\sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0)A_i^{1/2}(\beta_{g_i^0}^0)R^0 A_i^{1/2}(\beta_{g_i^0}^0)A_i^{-1/2}(\beta_{g_i}^0)(R^0)^{-1/2}\right) \\
& \geq \frac{1}{n}\sum_{i=1}^n \lambda_{\min}\left((R^0)^{-1/4}A_i^{-1/2}(\beta_{g_i}^0)A_i^{1/2}(\beta_{g_i^0}^0)R^0 A_i^{1/2}(\beta_{g_i^0}^0)A_i^{-1/2}(\beta_{g_i}^0)(R^0)^{-1/4}\right) \\
& \geq \frac{1}{n}\sum_{i=1}^n \lambda_{\min}^2\left((R^0)^{-1/4}A_i^{-1/2}(\beta_{g_i}^0)A_i^{1/2}(\beta_{g_i^0}^0)(R^0)^{1/2}\right) \\
& \geq \frac{1}{n}\sum_{i=1}^n \min_{1 \leq t \leq T} \{A_{it}^{-1}(\beta_{g_i}^0)A_{it}(\beta_{g_i^0}^0)\} \lambda_{\min}^{1/4}(R^0) > 0,
\end{aligned}$$

where the last inequality follows from Assumption (A5) (i). \square

The next lemma shows that $\widehat{R}^*(\beta, \bar{g})$ satisfies Assumption (A6) (i).

Lemma S12. *Under Assumptions (A1)-(A5) (i) and (A7)-(A10), it holds that for any \bar{g} ,*

$$\sup_{\beta \in \mathcal{B}_{nT}} \max_{1 \leq k, l \leq T} \{\widehat{R}^*(\beta, \bar{g}) - \widehat{R}^*(\beta^0, \bar{g})\}_{kl} = O_p(\lambda_{\min}^{-1/2}(\bar{H}^*)\tau^{1/2}).$$

Proof. For any \bar{g} , we can write

$$\begin{aligned}
& \widehat{R}^*(\beta, \bar{g}) - \widehat{R}^*(\beta^0, \bar{g}) \\
&= \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}) \{y_i - m(X_i \beta_{g_i})\} \{y_i - m(X_i \beta_{g_i})\}^t A_i^{-1/2}(\beta_{g_i}) \\
&\quad - \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \{y_i - m(X_i \beta_{g_i}^0)\} \{y_i - m(X_i \beta_{g_i}^0)\}^t A_i^{-1/2}(\beta_{g_i}^0) \\
&= \frac{1}{n} \sum_{i=1}^n \{A_i^{-1/2}(\beta_{g_i}) - A_i^{-1/2}(\beta_{g_i}^0)\} \{y_i - m(X_i \beta_{g_i})\} \\
&\quad \times \{y_i - m(X_i \beta_{g_i})\}^t \{A_i^{-1/2}(\beta_{g_i}) - A_i^{-1/2}(\beta_{g_i}^0)\} \\
&\quad + \frac{1}{n} \sum_{i=1}^n \{A_i^{-1/2}(\beta_{g_i}) - A_i^{-1/2}(\beta_{g_i}^0)\} \{y_i - m(X_i \beta_{g_i})\} \{y_i - m(X_i \beta_{g_i})\}^t A_i^{-1/2}(\beta_{g_i}^0) \\
&\quad + \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \{y_i - m(X_i \beta_{g_i})\} \{y_i - m(X_i \beta_{g_i})\}^t \{A_i^{-1/2}(\beta_{g_i}) - A_i^{-1/2}(\beta_{g_i}^0)\} \\
&\quad + \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \left[\{y_i - m(X_i \beta_{g_i})\} \{y_i - m(X_i \beta_{g_i})\}^t \right. \\
&\quad \quad \left. - \{y_i - m(X_i \beta_{g_i}^0)\} \{y_i - m(X_i \beta_{g_i}^0)\}^t \right] A_i^{-1/2}(\beta_{g_i}^0) \\
&\equiv \sum_{j=1}^4 I_j.
\end{aligned}$$

From Taylor expansion, for $\beta_{g_i}^*$ between β_{g_i} and $\beta_{g_i}^0$, we have

$$\begin{aligned}
1 - A_{it}^{1/2}(\beta_{g_i}) A_{it}^{-1/2}(\beta_{g_i}^0) &= 1 - \sqrt{\frac{a''(x_{it}^t \beta_{g_i})}{a''(x_{it}^t \beta_{g_i}^0)}} \\
&= -\frac{1}{2} \{a''(x_{it}^t \beta_{g_i}^*) a''(x_{it}^t \beta_{g_i}^0)\}^{-1/2} u'(x_{it}^t \beta_{g_i}^*) x_{it}^t (\beta_{g_i} - \beta_{g_i}^0).
\end{aligned}$$

Then, the (k, l) -element of I_1 can be written as

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \{A_{ik}^{-1/2}(\beta_{g_i}) - A_{ik}^{-1/2}(\beta_{g_i}^0)\} \{A_{il}^{-1/2}(\beta_{g_i}) - A_{il}^{-1/2}(\beta_{g_i}^0)\} \\
& \quad \times \{y_{ik} - m(x_{ik}^t \beta_{g_i})\} \{y_{il} - m(x_{il}^t \beta_{g_i})\} \\
& = \frac{1}{4n} \sum_{i=1}^n \{a''(x_{ik}^t \beta_{g_i}^*) a''(x_{ik}^t \beta_{g_i}^0)\}^{-1/2} u'(x_{ik}^t \beta_{g_i}^*) x_{ik}^t (\beta_{g_i} - \beta_{g_i}^0) \\
& \quad \times \{a''(x_{il}^t \beta_{g_i}^*) a''(x_{il}^t \beta_{g_i}^0)\}^{-1/2} u'(x_{il}^t \beta_{g_i}^*) x_{il}^t (\beta_{g_i} - \beta_{g_i}^0) \\
& \quad \times A_{ik}^{-1/2}(\beta_{g_i}) \{y_{ik} - m(x_{ik}^t \beta_{g_i})\} \{y_{il} - m(x_{il}^t \beta_{g_i})\} A_{il}^{-1/2}(\beta_{g_i}) \\
& \lesssim \left(\frac{1}{n} \sum_{i=1}^n (\beta_{g_i} - \beta_{g_i}^0)^t x_{ik} x_{ik}^t (\beta_{g_i} - \beta_{g_i}^0) \{y_{ik} - m(x_{ik}^t \beta_{g_i})\}^2 \right)^{1/2} \\
& \quad \times \left(\frac{1}{n} \sum_{i=1}^n (\beta_{g_i} - \beta_{g_i}^0)^t x_{il} x_{il}^t (\beta_{g_i} - \beta_{g_i}^0) \{y_{il} - m(x_{il}^t \beta_{g_i})\}^2 \right)^{1/2} \\
& \leq \{ \max_{1 \leq t \leq T} \lambda_{\max}(x_{it} x_{it}^t) \} \frac{1}{n} \sum_{i=1}^n \|\beta_{g_i} - \beta_{g_i}^0\|^2 \{y_{ik} - m(x_{ik}^t \beta_{g_i})\}^2,
\end{aligned}$$

where the second last inequality follows from Cauchy-Schwarz inequality. Since we have for all $t = 1, \dots, T$, $\lambda_{\max}(x_{it} x_{it}^t) = O_p(1)$ and $\frac{1}{n} \sum_{i=1}^n \{y_{it} - m(x_{it}^t \beta_{g_i})\}^2 = O_p(1)$, this implies that the order of $\{I_1\}_{kl}$ is $O_p(\lambda_{\min}^{-1}(\overline{H}^*) \tau)$ for $\beta \in \mathcal{B}_{nT}$. Similarly, the order of $\{I_2\}_{kl}$ and $\{I_3\}_{kl}$ are $O_p(\lambda_{\min}^{-1/2}(\overline{H}^*) \tau^{1/2})$ for $\beta \in \mathcal{B}_{nT}$. For I_4 , we can write

$$\begin{aligned}
I_4 & = \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \{m(X_i \beta_{g_i}^0) - m(X_i \beta_{g_i})\} \{y_i - m(X_i \beta_{g_i}^0)\}^t A_i^{-1/2}(\beta_{g_i}^0) \\
& \quad + \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \{y_i - m(X_i \beta_{g_i}^0)\} \{m(X_i \beta_{g_i}^0) - m(X_i \beta_{g_i})\}^t A_i^{-1/2}(\beta_{g_i}^0) \\
& \quad + \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \{m(X_i \beta_{g_i}^0) - m(X_i \beta_{g_i})\} \{m(X_i \beta_{g_i}^0) - m(X_i \beta_{g_i})\}^t A_i^{-1/2}(\beta_{g_i}^0) \\
& \equiv \sum_{j=1}^3 I_{4j}.
\end{aligned}$$

By using (S1) for $\beta_{g_i}^*$ between β_{g_i} and $\beta_{g_i}^0$, the (k, l) -element of I_{41} can be written as

from Cauchy-Schwarz inequality,

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n A_{ik}^{-1/2}(\beta_{g_i}^0) A_{il}^{-1/2}(\beta_{g_i}^0) \{m(x_{ik}^t, \beta_{g_i}^0) - m(x_{ik}^t, \beta_{g_i})\} \{y_{il} - m(x_{il}^t, \beta_{g_i}^0)\} \\
&= \frac{1}{n} \sum_{i=1}^n A_{ik}^{-1/2}(\beta_{g_i}^0) A_{il}^{-1/2}(\beta_{g_i}^0) \phi A_{ik}(\beta_{g_i}^*) u'(x_{ik}^t, \beta_{g_i}^*) x_{ik}^t (\beta_{g_i}^0 - \beta_{g_i}) \{y_{il} - m(x_{il}^t, \beta_{g_i}^0)\} \\
&\lesssim \left(\frac{1}{n} \sum_{i=1}^n (\beta_{g_i}^0 - \beta_{g_i})^t x_{ik} x_{ik}^t (\beta_{g_i}^0 - \beta_{g_i}) \right)^{1/2} \\
&\quad \times \left(\frac{1}{n} \sum_{i=1}^n A_{ik}^{-1}(\beta_{g_i}^0) A_{il}^{-1}(\beta_{g_i}^0) A_{ik}^2(\beta_{g_i}^*) \{u'(x_{ik}^t, \beta_{g_i}^*)\}^2 \{y_{il} - m(x_{il}^t, \beta_{g_i}^0)\}^t \{y_{il} - m(x_{il}^t, \beta_{g_i}^0)\} \right)^{1/2},
\end{aligned}$$

which implies that the order of $\{I_{41}\}_{kl}$ is $O_p(\lambda_{\min}^{-1/2}(\overline{H}^*)\tau^{1/2})$ for $\beta \in \mathcal{B}_{nT}$. Similarly, the order of $\{I_{42}\}_{kl}$ and $\{I_{43}\}_{kl}$ are $O_p(\lambda_{\min}^{-1/2}(\overline{H}^*)\tau^{1/2})$ and $O_p(\lambda_{\min}^{-1}(\overline{H}^*)\tau)$, respectively for $\beta \in \mathcal{B}_{nT}$, which proves the lemma. \square

The next lemma shows that $\widehat{R}^*(\beta, \bar{g})$ satisfies Assumption (A6) (ii).

Lemma S13. *Under Assumptions (A1)-(A5) (i) and (A7)-(A10), it holds that for any \bar{g} ,*

$$\sup_{\beta \in \mathcal{B}_{nT}} \max_{1 \leq k, l \leq T} |\{\widehat{R}^*(\beta, \bar{g}) - \overline{R}^*(\beta, \bar{g})\}_{kl}| = O_p(n^{-1/2} \vee \lambda_{\min}^{-1/2}(\overline{H}^*(\beta^0))\tau^{1/2}),$$

Proof. From Lemma S12, it is enough to show that

$$\max_{1 \leq k, l \leq T} \{\widehat{R}^*(\beta^0, \bar{g}) - \overline{R}^*(\beta^0, \bar{g})\}_{kl} = O_p(n^{-1/2}).$$

We can write

$$\begin{aligned}
& \widehat{R}^*(\beta^0, \bar{g}) - \bar{R}^*(\beta^0, \bar{g}) \\
&= \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \left\{ \{y_i - m(X_i \beta_{g_i}^0)\} \{y_i - m(X_i \beta_{g_i}^0)\}^t - \Sigma_i \right\} A_i^{-1/2}(\beta_{g_i}^0) \\
&\quad + \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \{m(X_i \beta_{g_i}^0) - m(X_i \beta_{g_i}^0)\} \{y_i - m(X_i \beta_{g_i}^0)\}^t A_i^{-1/2}(\beta_{g_i}^0) \\
&\quad + \frac{1}{n} \sum_{i=1}^n A_i^{-1/2}(\beta_{g_i}^0) \{y_i - m(X_i \beta_{g_i}^0)\} \{m(X_i \beta_{g_i}^0) - m(X_i \beta_{g_i}^0)\}^t A_i^{-1/2}(\beta_{g_i}^0) \\
&= I_1 + I_2 + I_3.
\end{aligned}$$

For $\sigma_{ikl} = \{\Sigma_i\}_{kl}$, the (k, l) -element of I_1 can be written as

$$\{I_1\}_{kl} = \frac{1}{n} \sum_{i=1}^n A_{ik}^{-1/2}(\beta_{g_i}^0) A_{il}^{-1/2}(\beta_{g_i}^0) [\{y_{ik} - m(x_{ik}^t \beta_{g_i}^0)\} \{y_{il} - m(x_{il}^t \beta_{g_i}^0)\} - \sigma_{ikl}].$$

Then, it is obvious $E[\{I_1\}_{kl}] = 0$ and

$$\begin{aligned}
\text{Var}(\{I_1\}_{kl}) &= \frac{1}{n^2} \sum_{i=1}^n A_{ik}^{-1}(\beta_{g_i}^0) A_{il}^{-1}(\beta_{g_i}^0) A_{ik}(\beta_{g_i}^0) A_{il}(\beta_{g_i}^0) \text{Var}(\varepsilon_{ik} \varepsilon_{il}) \\
&\leq \frac{1}{n^2} \sum_{i=1}^n A_{ik}^{-1}(\beta_{g_i}^0) A_{il}^{-1}(\beta_{g_i}^0) A_{ik}(\beta_{g_i}^0) A_{il}(\beta_{g_i}^0) (E[\varepsilon_{ik}^4] E[\varepsilon_{il}^4])^{1/2} = O_p(1/n),
\end{aligned}$$

where the last equality follows from Assumptions (A4), (A7) and (A8). Then, this implies that the order of the (k, l) -element of I_1 is $O_p(n^{-1/2})$. Similarly, both of the (k, l) -elements of I_2 and I_3 are $O_p(n^{-1/2})$, which implies the lemma. \square

The next lemma shows that $\widehat{R}^*(\beta, \bar{g})$ satisfies Assumption (A6) (iii).

Lemma S14. *Under Assumptions (A1)-(A5) (i) and (A7)-(A10), it holds that for any $\beta \in \mathcal{B}$, \bar{g} and \bar{g}_{i^*} whose only i th component differs from that of \bar{g} ,*

$$\max_{1 \leq k, l \leq T} |\{\widehat{R}^*(\beta, \bar{g}_{i^*}) - \widehat{R}^*(\beta, \bar{g})\}_{kl}| = O_p(1/n).$$

Proof. The lemma immediately holds since we can write

$$\begin{aligned} & \{\widehat{R}^*(\beta, \bar{g}_{i*}) - \widehat{R}^*(\beta, \bar{g})\}_{kl} \\ &= \frac{1}{n} \left\{ A_{ik}^{-1/2}(\beta_{g_i^*}) A_{il}^{-1/2}(\beta_{g_i^*}) \{y_{ik} - m(x_{ik}^t \beta_{g_i^*})\} \{y_{il} - m(x_{il}^t \beta_{g_i^*})\} \right. \\ & \quad \left. - A_{ik}^{-1/2}(\beta_{g_i^*}) A_{il}^{-1/2}(\beta_{g_i^*}) \{y_{ik} - m(x_{ik}^t \beta_{g_i^*})\} \{y_{il} - m(x_{il}^t \beta_{g_i^*})\} \right\}, \end{aligned}$$

which is of order $O_p(1/n)$. \square

The next lemma shows that $\widehat{R}^*(\beta, \bar{g})$ satisfies Assumption (A11).

Lemma S15. *Under Assumptions (A1)-(A5) (i) and (A7)-(A10), it holds that for any $\beta \in \mathcal{B}$, any \bar{g} satisfying $\sup_{\beta \in \mathcal{B}_{nT}} n^{-1} \sum_{i=1}^n \mathbb{I}\{g_i \neq g_i^0\} = o_p(T^{-\delta})$ and all $\delta > 0$,*

$$\max_{1 \leq k, l \leq T} |\{\widehat{R}^*(\beta, \bar{g}) - \widehat{R}^*(\beta, \bar{g}^0)\}_{kl}| = o_p(T^{-\delta}).$$

Proof. From Cauchy-Schwarz inequality, we can write

$$\begin{aligned} & \{\widehat{R}^*(\beta, \bar{g}) - \widehat{R}^*(\beta, \bar{g}^0)\}_{kl} \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{g_i \neq g_i^0\} \left\{ A_{ik}^{-1/2}(\beta_{g_i}) A_{il}^{-1/2}(\beta_{g_i}) \{y_{ik} - m(x_{ik}^t \beta_{g_i})\} \{y_{il} - m(x_{il}^t \beta_{g_i})\} \right. \\ & \quad \left. - A_{ik}^{-1/2}(\beta_{g_i^0}) A_{il}^{-1/2}(\beta_{g_i^0}) \{y_{ik} - m(x_{ik}^t \beta_{g_i^0})\} \{y_{il} - m(x_{il}^t \beta_{g_i^0})\} \right\} \\ & \leq \left(\frac{1}{n} \sum_{i=1}^n \mathbb{I}\{g_i \neq g_i^0\} \right)^{1/2} \\ & \quad \times \left(\frac{1}{n} \sum_{i=1}^n \left\{ A_{ik}^{-1/2}(\beta_{g_i}) A_{il}^{-1/2}(\beta_{g_i}) \{y_{ik} - m(x_{ik}^t \beta_{g_i})\} \{y_{il} - m(x_{il}^t \beta_{g_i})\} \right. \right. \\ & \quad \left. \left. - A_{ik}^{-1/2}(\beta_{g_i^0}) A_{il}^{-1/2}(\beta_{g_i^0}) \{y_{ik} - m(x_{ik}^t \beta_{g_i^0})\} \{y_{il} - m(x_{il}^t \beta_{g_i^0})\} \right\}^2 \right)^{1/2}. \end{aligned}$$

Since we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \left\{ A_{ik}^{-1/2}(\beta_{g_i}) A_{il}^{-1/2}(\beta_{g_i}) \{y_{ik} - m(x_{ik}^t \beta_{g_i})\} \{y_{il} - m(x_{il}^t \beta_{g_i})\} \right. \\ & \quad \left. - A_{ik}^{-1/2}(\beta_{g_i^0}) A_{il}^{-1/2}(\beta_{g_i^0}) \{y_{ik} - m(x_{ik}^t \beta_{g_i^0})\} \{y_{il} - m(x_{il}^t \beta_{g_i^0})\} \right\}^2 = O_p(1), \end{aligned}$$

the lemma follows from Lemma S1. □

S4 Additional results of the application to election panel study

We here provide additional results of the regression coefficients obtained by the proposed grouped GEE analysis with first-order autoregressive (AR) and unstructured (UN) correlation structures with $G = 8$. The results are shown in Table S1, which shows that the estimates of regression coefficients are generally close to the ones based on the exchangeable working correlation given in Table 1 in the main text.

References

- Bonhomme, S. and Manresa, E. (2015). Grouped pattern of heterogeneity in panel data. *Econometrica* 83, 1147–1184.
- Rio, E. (2000). *Théorie asymptotique des processus aléatoires faiblement dépendants*. Berlin: Springer. 1158–1176.
- Wang, L. (2011). GEE analysis of clustered binary data with diverging number of covariates. *The Annals of Statistics* 39, 389–417.
- Xie, M. and Yang, Y. (2003). Asymptotics for generalized estimating equations with large cluster sizes. *The Annals of Statistics* 31, 310–347.

Table S1: Point estimates (PE) and its standard errors (SE) of the regression coefficients of gender (G), indicators of black (BL) and others (OT), time (T) and the interaction term of gender and time (T×G) in 8 groups under AR and US working correlations.

		Group							
AR		1	2	3	4	5	6	7	8
Intercept	PE	-0.26	-0.60	-5.67	0.08	7.79	-4.12	-2.38	2.67
	SE	0.14	0.26	0.54	0.17	1.04	0.34	0.33	0.22
G	PE	2.76	-2.21	1.43	-0.32	-3.09	-0.66	0.77	-3.00
	SE	0.29	0.44	0.67	0.23	1.04	0.66	0.36	0.28
BL	PE	0.13	-2.34	17.46	0.00	-2.58	-0.40	-1.71	-1.04
	SE	0.15	0.93	0.15	0.25	0.27	0.21	0.29	0.24
OT	PE	-0.74	-0.59	2.42	-0.64	-2.21	-14.10	4.45	-2.20
	SE	0.12	0.08	0.29	0.18	0.39	0.12	0.76	0.21
T	PE	0.24	-0.20	0.34	-0.06	-0.93	0.69	1.02	0.27
	SE	0.03	0.06	0.07	0.03	0.14	0.06	0.12	0.06
T×G	PE	-0.61	0.01	0.65	0.26	0.92	-0.07	-0.14	-0.36
	SE	0.06	0.10	0.13	0.04	0.16	0.09	0.14	0.07

		Group							
US		1	2	3	4	5	6	7	8
Intercept	PE	-0.71	-0.02	-4.52	0.20	7.65	-4.04	-2.36	2.77
	SE	0.14	0.22	0.32	0.17	1.10	0.33	0.33	0.20
G	PE	3.36	-2.77	0.55	-0.48	-3.61	-1.00	1.13	-3.07
	SE	0.31	0.41	0.48	0.20	1.10	0.75	0.39	0.27
BL	PE	0.52	-12.41	16.79	-0.54	-2.48	-0.56	-1.92	-1.31
	SE	0.18	0.10	0.11	0.26	0.28	0.27	0.29	0.20
OT	PE	-0.89	0.00	0.69	-0.18	-2.31	-13.85	4.37	-2.40
	SE	0.13	0.01	0.62	0.14	0.24	0.13	0.78	0.21
T	PE	0.33	-0.30	0.23	-0.05	-0.92	0.67	1.01	0.22
	SE	0.03	0.06	0.05	0.03	0.16	0.05	0.12	0.05
T×G	PE	-0.72	0.10	0.66	0.21	0.98	-0.02	-0.22	-0.32
	SE	0.06	0.09	0.10	0.03	0.17	0.10	0.14	0.06