

Sample complexity and effective dimension for regression on manifolds

Andrew D. McRae Justin Romberg Mark A. Davenport

School of Electrical and Computer Engineering
Georgia Institute of Technology
Atlanta, GA 30332

admcr@gatech.edu, jrom@ece.gatech.edu, mdav@gatech.edu

Abstract

We consider the theory of regression on a manifold using reproducing kernel Hilbert space methods. Manifold models arise in a wide variety of modern machine learning problems, and our goal is to help understand the effectiveness of various implicit and explicit dimensionality-reduction methods that exploit manifold structure. Our first key contribution is to establish a novel nonasymptotic version of the Weyl law from differential geometry. From this we are able to show that certain spaces of smooth functions on a manifold are effectively finite-dimensional, with a complexity that scales according to the manifold dimension rather than any ambient data dimension. Finally, we show that given (potentially noisy) function values taken uniformly at random over a manifold, a kernel regression estimator (derived from the spectral decomposition of the manifold) yields minimax-optimal error bounds that are controlled by the effective dimension.

1 Introduction

High-dimensional data is ubiquitous in modern machine learning. Examples include images (2-D and 3-D), document texts, DNA, and neural recordings. In many cases, the number of dimensions in the data is much larger than the number of actual data samples. Traditional statistical methods cannot handle such cases, so researchers have turned to a variety of explicit dimensionality-reduction techniques—which make inference more tractable—and to tools such as neural networks that often implicitly transform the data into a much lower-dimensional feature space. These techniques inherently assume that the data have an *intrinsic* dimension that is much lower than that of the data’s original representation. Our goal in this paper is to show that the difficulty of a supervised learning problem depends only on this intrinsic dimension and not on the (potentially much larger) ambient dimension. In particular, we consider the common assumption that the data lie on a low-dimensional *manifold* embedded in Euclidean space (see [1–4] for some of the many example applications).

As an illustration of the kind of results we hope to obtain, we first consider a simple example: a function on the circle S^1 (or, equivalently, a periodic function on the real line). Specifically, suppose that we want to estimate a function f^* on the circle from random samples. In general, it is intractable to estimate an arbitrary function from finitely many samples, but it becomes possible if we assume f^* is structured. For example, f^* may exhibit a degree of smoothness, which can be readily characterized via the *Fourier series* for f^* . Specifically, recall that we can write f^* as the Fourier series sum $f^*(x) = a_0 + \sum_{\ell \geq 1} (a_\ell \cos(2\pi\ell x) + b_\ell \sin(2\pi\ell x))$. One common notion of smoothness in signal processing is that f^* is *bandlimited*, meaning that this sum can be truncated at some largest frequency Ω . In this case, f^* lies in a subspace of dimension at most $p(\Omega) = 2\lfloor \Omega/2\pi \rfloor + 1$. We know (see, e.g., [5, Chapter 12] or [6]) that we can recover such a function exactly, with high probability, from $n \gtrsim p(\Omega) \log p(\Omega)$ samples placed uniformly at random. If there is measurement noise, the

squared L_2 error due to noise scales like $\frac{p(\Omega)}{n}\sigma^2$. In higher dimensions (say, on the torus T^m), an Ω -bandlimited function lies in a space of dimension $p(\Omega) = O(\Omega^m)$, and the number of random samples required scales accordingly.

Another model for smoothness is that f^* , rather than being bandlimited, has exponentially-decaying frequency components. For example, suppose the Fourier coefficients satisfy $\sum_{\ell} e^{t\ell^2} (a_{\ell}^2 + b_{\ell}^2) < \infty$ for some $t > 0$ (this is roughly equivalent to f^* being the convolution of a Gaussian function with an arbitrary function in L_2). The space of such functions is infinite-dimensional, but any function in it can be approximated as Ω -bandlimited to within an error of size $O(e^{-c\Omega^2 t})$, which should enable us to recover a close approximation to f^* from $O(p(\Omega) \log p(\Omega))$ samples.

In this paper, we provide precise analogs of these sample complexity results in the general case of a function on an arbitrary manifold \mathcal{M} with dimension m . As on the circle or torus, an L_2 function $f(x)$ on a Riemannian manifold has a *spectral decomposition* into modes $u_{\ell}(x)$ corresponding to vibrational frequencies ω_{ℓ} for all non-negative integers ℓ ; these modes are the eigenfunctions of the Laplace-Beltrami operator on \mathcal{M} . Our first key contribution (described in Theorem 2) is a nonasymptotic version of the *Weyl law* from differential geometry: this states that, for large enough Ω , the set $\mathcal{H}_{\Omega}^{\text{bl}}$ of Ω -bandlimited functions on \mathcal{M} (functions composed of modes with frequencies below Ω) has dimension $\dim(\mathcal{H}_{\Omega}^{\text{bl}}) \leq C_m \text{vol}(\mathcal{M}) \Omega^m =: p(\Omega)$. Thus the number of degrees of freedom scales according to the *manifold dimension* m rather than a larger ambient dimension.

Our second key contribution is an error bound for recovering functions on \mathcal{M} from randomly-placed samples using kernel regression. We show in Theorem 3 that if we take $n \gtrsim p(\Omega) \log p(\Omega)$ samples of f^* , we can recover any Ω -bandlimited function with error

$$\frac{\|\hat{f} - f^*\|_{L_2}^2}{\text{vol}(\mathcal{M})} \lesssim \frac{p(\Omega)}{n} \sigma^2,$$

which is precisely the error rate for parametric regression in a $D(\Omega)$ -dimensional space. Our results extend further to approximately-bandlimited functions: for example, if f^* satisfies $\sum_{\ell} a_{\ell}^2 e^{t\omega_{\ell}^2} < \infty$, where $f^* = \sum_{\ell} a_{\ell} u_{\ell}$, then, again with $n \gtrsim p(\Omega) \log p(\Omega)$ samples, we get (Theorem 4)

$$\frac{\|\hat{f} - f^*\|_{L_2}^2}{\text{vol}(\mathcal{M})} \lesssim \frac{p(\Omega)}{n} \sigma^2 + O(e^{-c\Omega^2 t}).$$

Both bounds are minimax optimal in the presence of noise.

These results follow from our Theorem 1, which is a more general result on regression in a reproducing kernel Hilbert space. Theorems 3 and 4 adapt this result to a specific choice of kernel.

The paper is organized as follows. Sections 2 and 3 describe our framework, survey the relevant literature, and compare it to our results. Section 4 contains our main theoretical results. The proofs are in the appendices in the supplementary material. The key technical results are Theorem 1, which is proved via empirical risk minimization and operator concentration inequalities, and Lemma 1 (used to prove Theorem 2), which is proved via heat kernel comparison results on manifolds of bounded curvature.

2 Framework and notation

2.1 Kernel regression and interpolation

Kernels provide a convenient and popular framework for nonparametric function estimation. They allow us to treat the evaluation of a nonlinear function as a *linear* operator on a Hilbert space, and they give us a computationally feasible way to estimate such a function (which is often in an infinite-dimensional space) from a finite set of samples. Here, we review some of the key ideas that we will need in analyzing kernel methods.

Let S be an arbitrary set, and suppose $k: S \times S \rightarrow \mathbf{R}$ is a positive definite kernel. Let \mathcal{H} be its associated reproducing kernel Hilbert space (RKHS), characterized by the identity $f(x) = \langle f, k(\cdot, x) \rangle_{\mathcal{H}}$ for all $f \in \mathcal{H}$ and $x \in S$.

Now, suppose we have $X_1, \dots, X_n \in S$, $f^* \in \mathcal{H}$ is an unknown function, and we observe $Y_i = f^*(X_i) + \xi_i$ for $i = 1, \dots, n$, where the ξ_i 's represent noise. A common estimator for f^* is the

regularized empirical risk minimizer

$$\hat{f} = \arg \min_{f \in \mathcal{H}} \frac{1}{n} \sum_{i=1}^n (Y_i - f(X_i))^2 + \alpha \|f\|_{\mathcal{H}}^2, \quad (1)$$

where $\alpha \geq 0$ is a regularization parameter. The solution to the optimization problem (1) is

$$\hat{f}(x) = \sum_{i=1}^n a_i k(x, X_i), \quad (2)$$

where $\mathbf{a} = (a_1, \dots, a_n) \in \mathbf{R}^n$ is given by

$$\mathbf{a} = (n\alpha \mathbf{I}_n + \mathbf{K})^{-1} \mathbf{Y},$$

where $\mathbf{Y} = (Y_1, \dots, Y_n) \in \mathbf{R}^n$, \mathbf{K} is the kernel matrix on X_1, \dots, X_n defined by $K_{ij} = k(X_i, X_j)$, and \mathbf{I}_n is the $n \times n$ identity matrix.

In general, \hat{f} corresponds to a *ridge regression* estimate of f^* . The limiting case $\alpha = 0$ can be recast as the problem

$$\hat{f} = \arg \min_{f \in \mathcal{H}} \|f\|_{\mathcal{H}} \text{ s.t. } Y_i = f(X_i), \quad i = 1, \dots, n.$$

In this case, if the X_i 's are distinct, then \hat{f} interpolates the measured values of f^* .

2.2 Kernel integral operator and eigenvalue decomposition

A common tool for analyzing kernel interpolation and regression, which will play a central role in our analysis in Section 4, is the eigenvalue decomposition of a kernel's associated integral operator. The integral operator \mathcal{T} is defined for functions f on S by

$$(\mathcal{T}(f))(x) = \int_S k(x, y) f(y) d\mu(y),$$

where μ is a measure on S . Under certain assumptions¹ on S , μ , and k , \mathcal{T} is a well-defined operator on $L_2(S)$, is compact and positive definite with respect to the L_2 inner product, and has eigenvalue decomposition

$$\mathcal{T}(f) = \sum_{\ell=1}^{\infty} t_{\ell} \langle f, v_{\ell} \rangle_{L_2} v_{\ell}, \quad f \in L_2(S),$$

where the eigenvalues $\{t_{\ell}\}$ are arranged in decreasing order and converge to 0, and the eigenfunctions $\{v_{\ell}\}$ are an orthonormal basis for $L_2(S)$. We also have $k(x, y) = \sum_{\ell=1}^{\infty} t_{\ell} v_{\ell}(x) v_{\ell}(y)$, where the convergence is uniform and in L_2 .

This eigendecomposition plays an important role in characterizing the RKHS \mathcal{H} associated with the kernel k . Combining this expression for k with the identity $\langle f, k(\cdot, x) \rangle_{\mathcal{H}} = f(x)$, we can derive the fact that, for all $f, g \in \mathcal{H}$,

$$\langle f, g \rangle_{\mathcal{H}} = \sum_{\ell=1}^{\infty} \frac{\langle f, v_{\ell} \rangle_{L_2} \langle g, v_{\ell} \rangle_{L_2}}{t_{\ell}}.$$

This implies that $\langle f, g \rangle_{L_2} = \langle \mathcal{T}^{1/2}(f), \mathcal{T}^{1/2}(g) \rangle_{\mathcal{H}}$ for all $f, g \in L_2(S)$. Thus $\mathcal{T}^{1/2}$ is an isometry from $L_2(S)$ to \mathcal{H} , and so for any $f \in \mathcal{H}$, we can write $f = \mathcal{T}^{1/2}(f_0)$, where $\|f_0\|_{L_2} = \|f\|_{\mathcal{H}}$. This implies that, for any $p \geq 1$, the projection of f onto $(\text{span}\{v_1, \dots, v_p\})^{\perp}$ has L_2 norm at most $\sqrt{t_{p+1}} \|f\|_{\mathcal{H}}$. Hence the decay of the eigenvalues $\{t_{\ell}\}$ of \mathcal{T} characterizes the “effective dimension” of \mathcal{H} in L_2 , which will be a fundamental building block for our analysis.

¹E.g., S is a compact metric space; μ is strictly positive, finite, and Borel; and k is continuous [7].

2.3 Spectral decomposition of a manifold and related kernels

We now turn to our specific problem of regression on a manifold, considering how an RKHS framework can help us. The book [8] is an excellent reference for the material in this section.

A smooth, compact Riemannian manifold \mathcal{M} (without boundary) can be analyzed via the *spectral decomposition* of its Laplace-Beltrami operator $\Delta_{\mathcal{M}}$ (we will often call it the Laplacian for short). This operator is defined as $\Delta_{\mathcal{M}}f := -\operatorname{div}(\nabla f)$. In \mathbf{R}^m , it is simply the operator $-\sum_{i=1}^m \frac{\partial^2}{\partial x_i^2}$. The Laplacian can be diagonalized as

$$\Delta_{\mathcal{M}}f = \sum_{\ell=0}^{\infty} \lambda_{\ell} \langle f, u_{\ell} \rangle_{L_2} u_{\ell},$$

where $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots$, the sequence $\lambda_{\ell} \rightarrow \infty$ as $\ell \rightarrow \infty$, and $\{u_{\ell}\}$ is an orthonormal basis for $L_2(\mathcal{M})$ (all integrals are with respect to the standard volume measure on \mathcal{M}).

The eigenvalues $\{\lambda_{\ell}\}$ are the squared resonant frequencies of \mathcal{M} , and the eigenfunctions $\{u_{\ell}\}$ are the vibrating modes, since solutions to the wave equation $f_{tt} + \Delta_{\mathcal{M}}f = 0$ on \mathcal{M} have the form

$$f(t, x) = \sum_{\ell=0}^{\infty} (a_{\ell} \sin \sqrt{\lambda_{\ell}} t + b_{\ell} \cos \sqrt{\lambda_{\ell}} t) u_{\ell}(x).$$

The classical Weyl law (e.g., [8, p. 9]) says that, if \mathcal{M} has dimension m , then, asymptotically,

$$|\{\ell : \lambda_{\ell} \leq \lambda\}| \sim c_m \operatorname{vol}(\mathcal{M}) \lambda^{m/2}$$

as $\lambda \rightarrow \infty$, where $c_m = (2\pi)^{-m} V_m$, with V_m denoting the volume of the unit ball in \mathbf{R}^m .

Using the spectral decomposition of the Laplacian, any number of kernels can be defined by

$$k(x, y) = \sum_{\ell=0}^{\infty} g(\lambda_{\ell}) u_{\ell}(x) u_{\ell}(y)$$

for some function g . With this construction, the integral operator of k has eigenvalue decomposition $\mathcal{T}(f) = \sum_{\ell \geq 0} g(\lambda_{\ell}) \langle f, u_{\ell} \rangle_{L_2} u_{\ell}$, hence, per Section 2.2, $\|f\|_{\mathcal{H}}^2 = \sum_{\ell \geq 0} \langle f, u_{\ell} \rangle^2 / g(\lambda_{\ell})$.

Our results could, in principle, apply to many kernels with the above form, but we will primarily consider *bandlimited kernels* and the *heat kernel*. The bandlimited kernel with bandlimit $\Omega > 0$ is

$$k_{\Omega}^{\text{bl}}(x, y) = \sum_{\lambda_{\ell} \leq \Omega^2} u_{\ell}(x) u_{\ell}(y),$$

which is the reproducing kernel of the space of bandlimited functions on \mathcal{M} :

$$\mathcal{H}_{\Omega}^{\text{bl}} = \{f \in L_2(\mathcal{M}) : f \in \operatorname{span}\{u_{\ell} : \lambda_{\ell} \leq \Omega^2\}\}$$

with $\|f\|_{\mathcal{H}_{\Omega}^{\text{bl}}} = \|f\|_{L_2}$ for $f \in \mathcal{H}_{\Omega}^{\text{bl}}$. The heat kernel is a natural counterpart to the common Gaussian radial basis function on \mathbf{R}^m . Detailed treatments can be found in [8, 9]. We will define it for $t > 0$ as

$$k_t^{\text{h}}(x, y) = \sum_{\ell=0}^{\infty} e^{-\lambda_{\ell} t / 2} u_{\ell}(x) u_{\ell}(y).$$

Its corresponding RKHS is

$$\mathcal{H}_t^{\text{h}} = \left\{ f \in L_2(\mathcal{M}) : \|f\|_{\mathcal{H}_t^{\text{h}}}^2 = \sum_{\ell=0}^{\infty} e^{\lambda_{\ell} t / 2} \langle f, u_{\ell} \rangle_{L_2}^2 < \infty \right\}.$$

The heat kernel k_t^{h} gets its name from the fact that it is the fundamental solution to the heat equation $f_t + \frac{1}{2} \Delta_{\mathcal{M}} f = 0$ on \mathcal{M} . The heat kernel on \mathbf{R}^m is $k_t^{\text{h}}(x, y) = \frac{1}{(2\pi t)^{m/2}} e^{-\|x-y\|^2 / 2t}$.

3 Related work

3.1 Dimensionality reduction and low-dimensional structure

There is an extensive literature on the use of low-dimensional manifold structure in machine learning. Perhaps most prominently, nonlinear dimensionality-reduction techniques that exploit manifold structure have been developed, such as [10–14]. More recently, there has been explicit inclusion of manifold models into neural network architectures [15–18]. However, none of this research provides nonasymptotic performance guarantees.

On the other hand, the field of high-dimensional statistics provides many theoretical guarantees for low-dimensional data models. For example, there are extensive bodies of theory for models such as sparsity [19, 20] and low-rank structure [21]. One can view low-dimensional manifold models as a more powerful generalization of such structures. One interesting work that bridges the gap between manifold models and high-dimensional statistics is [22], which is another explicit dimensionality-reduction technique. Another similar line of work is the study of algebraic variety models (e.g., [23]), which are also nonlinear and low-dimensional.

While the great success of the many implicit and explicit dimensionality-reducing methods provides empirical evidence for the possibility of exploiting manifold structure, there are still very large gaps in our theoretical understanding of when and why these methods can be effective.

3.2 Manifold regression and kernels

Regression on manifold domains has been explored in a number of previous works. The closely-related problem of density estimation is considered in [24, 25]. Particularly relevant to our paper, [24] uses the same bandlimited kernel and heat kernel that we highlight (and it analyzes the spectral decomposition of these kernels via the asymptotic Weyl law). It is primarily interested in the power of the error rate that can be obtained by assuming the function (density) of interest has a certain number of derivatives; in particular, it shows that $\|\hat{f} - f^*\|_{L_2}^2 \lesssim n^{-2s/(m+2s)}$ if f has s bounded derivatives. Both works, like ours, assume explicit knowledge of the manifold.

Perhaps more relevant to practical applications, [26] seeks to provide a manifold-agnostic algorithm via local linear approximations to the data manifold; however, it is also primarily interested in asymptotic error rates. The paper [27] examines related methods asymptotically in more detail. Another manifold-agnostic method similar in spirit to ours is that of [28], who consider kernel estimation with (Euclidean) Gaussian radial basis functions. They obtain the optimal $n^{-2s/(m+2s)}$ rate for s -smooth regression functions; however, their assumptions are quite different from ours in that their regression functions must have *smooth extensions* to (a neighborhood in) the embedding space. Similarly, [29] obtain the optimal rate for functions that are s -smooth (in the manifold calculus, similarly to our assumptions) using a neural-network-type architecture. However, they implicitly assume that the manifold is C^∞ -embedded in Euclidean space.

In [30, 31], the authors explore Gaussian process models (which are closely related to kernel methods) on a manifold.

The error rate $\|\hat{f} - f^*\|_{L_2}^2 \lesssim n^{-2s/(m+2s)}$ is standard (and minimax optimal) in nonparametric statistics. However, our function model and results are quite different in nature. The regression functions we consider are *infinitely* smooth, and we show that the estimation of these functions is much like a *finite-dimensional* regression problem; not only do we get an n^{-1} error rate (as we do when we take $s \rightarrow \infty$ above), but the constant in front of this rate and the minimum number of samples needed are proportional to the finite effective dimension.

Finally, we also note that the idea of using a kernel that can be expressed in terms of the spectral decomposition of a manifold’s Laplacian also has precedent. In addition to [24], the paper [32] suggests using such kernels for interpolation in Sobolev spaces on a manifold.

3.3 General kernel interpolation and regression

Regression is a strict superset of interpolation; interpolation typically assumes that we sample function values exactly (i.e., there is no noise), while regression allows for (and often assumes) noise.

There is a substantial literature on the use of a kernel for interpolation of functions in an RKHS (often, in this literature, referred to as the “native space” of the kernel). A fairly comprehensive survey can be found in [33]. Distinct from our work, most of this literature considers *deterministic* samples of the function of interest. Given (deterministic) sample locations $\{X_1, \dots, X_n\} \subset S$, results in this literature tend to have the form $\|\hat{f} - f^*\|_\infty \leq g(h_X) \|f^*\|_{\mathcal{H}}$, where $h_X = \max_{x \in S} \min_{i \in \{1, \dots, n\}} d(x, X_i)$, and $g(h)$ is a function that decreases to 0 as $h \rightarrow 0$ at a rate that depends on the properties of the kernel k (typically as a power or exponentially). Some recent work applying kernel interpolation theory to manifolds is [34–36].

Much of the literature on (noisy) RKHS regression primarily considers the case when the eigenvalues of the integral operator (described in Section 2.1) decay as $t_\ell \lesssim \ell^{-b}$. In [37–39], it is shown that the minimax optimal error rate is $\|f^* - \hat{f}\|_{L_2}^2 \lesssim n^{-b/(b+1)}$. Many other recent papers have explored this rate of convergence in a variety of settings [40–43]. Several of these include more general spectral regularization algorithms, suggested by [44]. Some interesting recent extensions consider a variety of algorithms that may be more practical for large-data situations. These include iterative methods [45–47] and distributed algorithms [48–50].

Another set of results (which are the most similar to ours) uses a regularized effective dimension $p_\alpha = \sum_\ell \frac{t_\ell}{\alpha + t_\ell}$, where α is the regularization parameter. This is considered in [51] and greatly refined in [52]. Variations on these results can be found in [53]. See Section 4.1 for further discussion and comparison to our results. The earlier report [54] resembles our work in its analysis of truncated operators. We note that in the case of power-law eigenvalue decay, these results (and ours) recover the $n^{-b/(b+1)}$ error rate.

It is interesting to note that the squared error rate $n^{-b/(b+1)}$ can recover the standard rate for regression of s -smooth functions on manifolds. The Sobolev space of order s is the RKHS of the kernel $\sum_\ell (1 + \lambda_\ell)^{-s} u_\ell(x) u_\ell(y)$. By the Weyl law, its eigenvalues decay according to $t_\ell \approx \ell^{-2s/m}$; plugging $2s/m$ in for b recovers the standard rate $n^{-2s/(m+2s)}$.

4 Main theoretical results

4.1 Dimensionality in RKHS regression

Here we present our main results for general regression and interpolation in an RKHS. Our results also apply to the slightly more general setting of learning in an arbitrary Hilbert space (see, e.g., [52]), but we do not explore this here. We continue to use the notation established in Sections 2.1 and 2.2, and we further assume that $\mu(S) = 1$ (since μ is finite, we can always obtain this by a rescaling). We assume that the function samples we take are uniformly distributed on S :

Assumption 1. The sample locations X_1, \dots, X_n are i.i.d. according to μ .

Since \mathcal{H} is, in general, infinite-dimensional, there is typically no hope of recovering an arbitrary $f^* \in \mathcal{H}$ to within a small error in \mathcal{H} -norm from a finite number of measurements. However, the discussion in Section 2.2 suggests a more feasible goal. Since any set of functions bounded in \mathcal{H} -norm can be approximated within an arbitrarily small L_2 error in a finite-dimensional subspace of L_2 , as long as the number of measurements is proportional to this loosely-defined “effective dimension” of \mathcal{H} , we have hope of recovering f^* accurately in an L_2 sense.

Let $p > 0$ be a fixed integer dimension. Let $G = \text{span}\{v_1, \dots, v_p\} \subset \mathcal{H} \cap L_2(S)$, and let G^\perp be its orthogonal complement in $L_2(S)$ and \mathcal{H} . We denote by \mathcal{T}_G and \mathcal{T}_{G^\perp} the restrictions of \mathcal{T} onto G and G^\perp , respectively. We make the following assumptions on the eigenvalues and eigenfunctions of \mathcal{T} :

Assumption 2. For some constants K_p and R_p , we have $\sum_{\ell=1}^p v_\ell^2(x) \leq K_p$ and $\sum_{\ell=p+1}^\infty t_\ell v_\ell^2(x) \leq R_p$ for almost every $x \in S$.

This says that the energy of the eigenfunctions of \mathcal{T} is reasonably spread out over the domain S —for the basis $\{v_1, \dots, v_p\}$, this is a type of incoherence assumption. If the eigenfunctions are well-behaved, we can expect $K_p \approx p$ and $R_p \approx \text{tr } \mathcal{T}_{G^\perp}$. This holds in our original example of the Fourier series on the circle, since the sinusoid basis functions are bounded by an absolute constant. Our “pointwise” Weyl law in Theorem 2 shows that we have similar behavior for the spectral de-

composition of a manifold. Note that K_p in Assumption 2 is identical to the quantity $K(p)$ in [6], which uses similar methods to handle a much simpler problem.

Assumption 3. For some $\gamma, \gamma' \geq 0$, we have $\frac{\text{tr } \mathcal{T}_{G^\perp}}{t_{p+1}} \leq \gamma p$ and $\frac{R_p}{t_{p+1}} \leq \gamma' K_p$.

This assumption greatly simplifies the notation of our results and is always true with an appropriate choice of γ and γ' . γ is often small when t_{p+1} is in the decaying “tail” of eigenvalues. If the eigenvalues decay like $t_\ell \approx \ell^{-b}$, we can take $\gamma \approx (b-1)^{-1}$. Note that a similar assumption appears in [55]. If $K_p \approx p$ and $R_p \approx \text{tr } \mathcal{T}_{G^\perp}$, then $\gamma \approx \gamma'$.

With these assumptions in place, we can state our main theorem for RKHS regression:

Theorem 1. Suppose Assumptions 1 to 3 hold. Let $\delta \in (0, 1)$. If

$$n \geq (7 \vee 3\gamma') K_p \log \frac{(2 \vee 4\gamma)p}{\delta},$$

then the following hold for the kernel estimate \hat{f} with regularization parameter $\alpha \geq 0$:

1. If there is no noise, that is, $Y_i = f^*(X_i)$ for each i , then, with probability at least $1 - \delta$, uniformly in f^* ,

$$\|\hat{f} - f^*\|_{L_2} \leq (\sqrt{2\alpha} + 6\sqrt{t_{p+1}}) \|f^*\|_{\mathcal{H}}.$$

2. Now suppose that $Y_i = f(X_i) + \xi_i$, where the ξ_i 's are i.i.d., zero-mean, sub-exponential random variables with variance σ^2 and are independent of the X_i 's. If we additionally have

$$\frac{n}{\log^2 n} \geq C(1 \vee \gamma') \frac{K_p \|\xi\|_{\psi_1}^2}{\sigma^2},$$

where C is a universal constant, and $\alpha \geq 54t_{p+1}$, then, with probability at least $1 - 2\delta$, uniformly in f^* ,

$$\|\hat{f} - f^*\|_{L_2} \leq (\sqrt{2\alpha} + 6\sqrt{t_{p+1}}) \|f^*\|_{\mathcal{H}} + 4 \left(1 + \frac{\sqrt{\gamma}}{8}\right) \frac{\sqrt{p} + 2\sqrt{\log 4/\delta}}{\sqrt{n}} \sigma.$$

Our results guarantee an L_2 recovery error bounded by two terms: (1) a “bias” depending on the next tail eigenvalue t_{p+1} and the regularization coefficient α , and (2) a “variance” term that behaves similarly to the error found in p -dimensional regression. When $K_p \approx p$, this result yields the $n \gtrsim p \log p$ sample complexity that we expect. If \mathcal{H} is, in fact, p -dimensional (which our framework can handle with $t_\ell = 0$ for $\ell > p$), this result recovers standard p -dimensional regression bounds such as those in [6].

We assume i.i.d. noise for simplicity, but our result could easily be extended beyond this case. Note that if the noise is Gaussian, the ratio $\|\xi\|_{\psi_1}^2/\sigma^2$ is an absolute constant.

For interpolation ($\alpha = 0$) in the noiseless case, this theorem yields $\|\hat{f} - f^*\|_{L_2} \leq 6\sqrt{t_{p+1}} \|f^*\|_{\mathcal{H}}$. In the noisy case, the lower bound on α can be relaxed to get a result with worse constants. We obtain qualitatively similar results whenever $\alpha \gtrsim t_{p+1}$. The assumptions and results of [52] (specialized to our setting) are comparable to Theorem 1 when we set $\alpha \approx t_{p+1}$. However, our results have the advantage of applying even in infinite-dimensional settings with no regularization: the regularized effective dimension $p_\alpha = \sum_\ell \frac{t_\ell}{\alpha + t_\ell}$ from their work would be infinite if $\alpha = 0$.

Although we do not explore it here, we note that one could generalize our approach to the case where the sampling measure differs from that under which the L_2 norm is calculated. We could simply bound the ratio (Radon-Nikodym derivative) between the two measures, or we could perform leverage-score sampling to mitigate the need for bounding the eigenfunctions (see, e.g., [55] for similar ideas).

In the presence of noise, Theorem 1 is minimax optimal over the set $\{f \in \mathcal{H} : \|f\|_{\mathcal{H}} \leq r\}$ for any $r > 0$ if p is chosen so that $\frac{p}{n}\sigma^2 \approx t_{p+1}r^2$. In this case,

$$\left\{f \in \text{span}\{v_1, \dots, v_p\} : \|f\|_{L_2} \lesssim \sqrt{\frac{p}{n}}\sigma\right\} \subset \{f \in \mathcal{H} : \|f\|_{\mathcal{H}} \leq r\},$$

and the minimax rate (with, say, Gaussian noise) over the left-hand set is well-known to be $\sqrt{\frac{p}{n}}\sigma$.

4.2 Manifold function estimation

We now describe how we can leverage Theorem 1 to establish sample complexity bounds for regression on a manifold. Suppose, again, that \mathcal{M} is an m -dimensional smooth, compact Riemannian manifold. To study the eigenvalues and eigenfunctions of the Laplacian $\Delta_{\mathcal{M}}$, we consider the heat kernel k_t^h . Our key tool is the following fact:

Lemma 1. *Let $\epsilon \in (0, 2/3)$. Suppose the sectional curvature of \mathcal{M} is bounded above by κ . For $t \leq \frac{6\epsilon}{(m-1)^2\kappa}$ and all $x \in \mathcal{M}$,*

$$k_t^h(x, x) \leq \frac{1 + \epsilon}{(2\pi t)^{m/2}}.$$

This is a precise quantification of the well-known asymptotic behavior of the heat kernel as $t \rightarrow 0$ (see, e.g., [8, Section VI]). It is derived in Appendix B from a novel set of more general upper and lower bounds for the heat kernel on a manifold of bounded curvature; we note that these may be of independent interest.

Our nonasymptotic Weyl law is a simple consequence of Lemma 1:

Theorem 2. *If \mathcal{M} has sectional curvature bounded above by κ , and $\epsilon \in (0, 2/3)$, then, for all $x \in \mathcal{M}$ and $\lambda \geq \frac{m(m-1)^2\kappa}{6\epsilon}$,*

$$N_x(\lambda) := \sum_{\lambda_\ell \leq \lambda} u_\ell^2(x) \leq \frac{2(1 + \epsilon)\sqrt{m}}{(2\pi)^m} V_m \lambda^{m/2}.$$

With appropriate rescaling by $\text{vol}(\mathcal{M})$, this gives us a bound on the constant K_p from Section 4.1. Since this result bounds the eigenfunctions, it is a type of “local Weyl law” (see, e.g., [56]). Integrating this result over \mathcal{M} gives a nonasymptotic version of the traditional Weyl law. Our bound is within the modest factor $2(1 + \epsilon)\sqrt{m}$ of the optimal asymptotic law. For simplicity, we will take $\epsilon = 1/2$ in what follows, but slightly better constants could be obtained with smaller ϵ .

The following result for the finite-dimensional bandlimited kernel is a straightforward consequence of Theorems 1 and 2:

Theorem 3. *Suppose the sectional curvature of \mathcal{M} is bounded above by κ . Let $\Omega^2 \geq \frac{m(m-1)^2\kappa}{3}$, and suppose $f^* \in \mathcal{H}_\Omega^{\text{bl}}$. Let \hat{f} be the kernel regression estimate with kernel k_Ω^{bl} .²*

Let $\delta \in (0, 1)$, and suppose $n \geq 7p \log \frac{2p}{\delta}$, where

$$p = p(\Omega) := \frac{3\sqrt{m} V_m}{(2\pi)^m} \text{vol}(\mathcal{M}) \Omega^m. \quad (3)$$

Under the same noise assumptions as in Theorem 1, if $\frac{n}{\log^2 n} \geq C \|\xi\|_{\psi_1}^2 / \sigma^2$, then, with probability at least $1 - 2\delta$, uniformly in f^ ,*

$$\frac{\|\hat{f} - f^*\|_{L_2}}{\sqrt{\text{vol}(\mathcal{M})}} \leq 4 \frac{\sqrt{p} + 2\sqrt{\log 4/\delta}}{\sqrt{n}} \sigma.$$

To analyze the heat kernel, which has an infinite number of nonzero eigenvalues, we need the following additional corollary of Lemma 1, which will let us bound the constant R_p from Section 4.1:

Lemma 2. *For $\epsilon \in (0, 2/3)$, $t \leq \frac{6\epsilon}{(m-1)^2\kappa}$, $\lambda \geq m/t$, and all $x \in \mathcal{M}$,*

$$\sum_{\lambda_\ell \geq \lambda} e^{-\lambda_\ell t/2} u_\ell^2(x) \leq e^{-\lambda t/2} \frac{2(1 + \epsilon)\sqrt{m}}{(2\pi)^m} V_m \lambda^{m/2}.$$

From this, we obtain the following result:

²The calculation of this estimate is somewhat different than usual, since the rank of the kernel matrix \mathbf{K} is at most the dimension of $\mathcal{H}_\Omega^{\text{bl}}$. We do not use regularization, but we use the Moore-Penrose pseudoinverse of \mathbf{K} instead of \mathbf{K}^{-1} .

Theorem 4. Suppose the sectional curvature of \mathcal{M} is bounded above by κ . Let $t \leq \frac{3}{(m-1)^2\kappa}$, and suppose $f^* \in \mathcal{H}_t^h$. Fix $\Omega^2 \geq m/t$, and let \hat{f} be the kernel regression estimate of f^* with kernel k_t^h and regularization parameter $\alpha \geq 54 \frac{e^{-\Omega^2 t/2}}{\text{vol}(\mathcal{M})}$.

Let $\delta \in (0, 1)$, and suppose $n \geq 7p \log \frac{4p}{\delta}$, with p defined as in (3).

Under the same noise assumptions as in Theorem 1, if $\frac{n}{\log^2 n} \geq C \|\xi\|_{\psi_1}^2 / \sigma^2$, then, with probability at least $1 - 2\delta$, uniformly in f^* ,

$$\frac{\|\hat{f} - f^*\|_{L_2}}{\sqrt{\text{vol}(\mathcal{M})}} \leq \left(\sqrt{2\alpha} + 6 \sqrt{\frac{e^{-\Omega^2 t/2}}{\text{vol}(\mathcal{M})}} \right) \|f^*\|_{\mathcal{H}_t^h} + \frac{9}{2} \frac{\sqrt{p} + 2\sqrt{\log 4/\delta}}{\sqrt{n}} \sigma.$$

These results illustrate how we can exploit the effective finite dimension of spaces of smooth functions on manifolds in regression. This function space dimension (and hence the sample complexity of regression) grows exponentially in the *manifold dimension*, rather than in the larger ambient data dimension, if \mathcal{M} is embedded in a higher-dimensional space. In practice, the true bandlimited or heat kernels may be difficult to compute. It is an interesting open question whether we can obtain similar results for manifold-agnostic algorithms (the work of [26], although it does not apply to our function classes, is an interesting potential starting point).

As discussed in Section 4.1, our general regression result Theorem 1 is similar to prior results [51, 52], but it has the advantage of applying even without regularization in the noiseless case. However, we note that one could obtain results in many ways comparable (minus this advantage) to Theorems 3 and 4 by plugging Theorem 2 and Lemma 2 into those previous regression results. We could not do this with classical power-law results such as [37–39], since our eigenvalue decay is *exponential* rather than power-law.

Since the (classical) Weyl law also *lower* bounds the complexity of spaces of bandlimited functions, then, as discussed in Section 4.1, Theorems 3 and 4 (for the optimally chosen value of Ω) are minimax optimal when there is noise. Furthermore, the requirement $n \gtrsim p \log p$ is necessary in general: if we consider the torus T^m , recovering arbitrary Ω -bandlimited functions requires every point on T^m to be within distance $O(1/\Omega)$ of a sample point; considering a uniform grid on T^m and a coupon collector argument makes it clear that $n \gtrsim O(\Omega^m \log \Omega^m)$ randomly sampled points are required.

As mentioned in Section 4.1 for general kernel learning, these results could be extended to consider nonuniform sampling over the manifold.

There are also some very interesting connections between kernel methods and neural networks. The recent works [57, 58] show that trained multi-layer neural networks approach, in the infinite-width limit, a kernel regression function with a “neural tangent kernel” that depends on the initialization distribution of the weights and the network architecture. This follows literature on the connections between Gaussian processes (closely related to kernel methods) and wide neural networks (see, e.g., [59, 60]). It would be very interesting to explore any potential connections between these and the kernels considered in this paper, which are derived from a manifold’s spectral decomposition.

Broader Impact

The results in this paper further illuminate the role of low-dimensional structure in machine learning algorithms. An improved theoretical understanding of the performance of these algorithms is increasingly important as tools from machine learning become ever-more-widely adopted in a range of applications with significant societal implications. Although, in general, there are well-known ethical issues that can arise from inherent biases in the way data are sampled and presented to regression and classification algorithms, we do not have reason to believe that the methods presented in this paper would either enhance or diminish these issues. Our analysis is abstract and, for better or for worse, assumes a completely neutral sampling model (uniform over a manifold).

Acknowledgments and Disclosure of Funding

This work was supported, in part, by National Science Foundation grant CCF-1350616, a gift from the Alfred P. Sloan Foundation, and the Georgia Tech ARC-TRIAD student fellowship.

References

- [1] D. Donoho and C. Grimes, “Image manifolds which are isometric to Euclidean space,” *J. Math. Imaging Vis.*, vol. 23, no. 1, pp. 5–24, 2005.
- [2] G. Peyré, “Manifold models for signals and images,” *Comput. Vis. Image Underst.*, vol. 113, pp. 249–260, 2009.
- [3] B. Zhu, J. Liu, S. Cauley, B. Rosen, and M. Rosen, “Image reconstruction by domain-transform manifold learning,” *Nature*, vol. 555, pp. 487–492, 2018.
- [4] S. Ganguli and H. Sompolinsky, “Compressed sensing, sparsity, and dimensionality in neuronal information processing and data analysis,” *Annu. Rev. Neurosci.*, vol. 35, pp. 485–508, 2012.
- [5] S. Foucart and H. Rauhut, *A Mathematical Introduction to Compressive Sensing*. New York: Birkhäuser, 2013.
- [6] A. Cohen, M. A. Davenport, and D. Leviatan, “On the stability and accuracy of least squares approximations,” *Found. Comput. Math.*, vol. 13, pp. 819–834, 2013.
- [7] I. Steinwart and C. Scovel, “Mercer’s theorem on general domains: On the interaction between measures, kernels, and RKHSs,” *Constr. Approx.*, vol. 35, pp. 363–417, 2012.
- [8] I. Chavel, *Eigenvalues in Riemannian Geometry*. Academic Press, 1984.
- [9] E. Hsu, *Stochastic Analysis on Manifolds*. Providence, RI: American Mathematical Society, 2002.
- [10] S. T. Roweis and L. K. Saul, “Nonlinear dimensionality reduction by locally linear embedding,” *Science*, vol. 290, no. 5500, pp. 2323–2326, 2000.
- [11] J. B. Tenenbaum, V. de Silva, and J. C. Langford, “A global geometric framework for nonlinear dimensionality reduction,” *Science*, vol. 290, no. 5500, pp. 2319–2323, 2000.
- [12] M. Belkin and P. Niyogi, “Laplacian eigenmaps for dimensionality reduction and data representation,” *Neural Comput.*, vol. 15, pp. 1373–1396, 2003.
- [13] R. R. Coifman and S. Lafon, “Diffusion maps,” *Appl. Comput. Harmon. Anal.*, vol. 21, pp. 5–30, 2006.
- [14] L. van der Maaten and G. Hinton, “Visualizing data using t-SNE,” *J. Mach. Learn. Res.*, vol. 9, pp. 2579–2605, 2008.
- [15] F. Monti, D. Boscaini, J. Masci, E. Rodolà, J. Svoboda, and M. Bronstein, “Geometric deep learning on graphs and manifolds using mixture model CNNs,” in *Proc. Conf. Comput. Vis. Pattern Recog. (CVPR)*, Honolulu, Hawaii, United States, Jul. 2017, pp. 5425–5434.
- [16] J. Masci, D. Boscaini, M. Bronstein, and P. Vandergheynst, “Geodesic convolutional neural networks on riemannian manifolds,” in *Proc. IEEE Int. Conf. Comput. Vis. (ICCV) Workshop*, Santiago, Chile, Dec. 2015, pp. 832–840.
- [17] D. Boscaini, J. Masci, E. Rodolà, and M. Bronstein, “Learning shape correspondence with anisotropic convolutional neural networks,” in *Proc. Conf. Neural Inf. Process. Syst. (NeurIPS)*, Barcelona, Spain, Dec. 2016.
- [18] H. Shao, A. Kumar, and P. T. Fletcher, “The Riemannian geometry of deep generative models,” in *Proc. Conf. Comput. Vis. Pattern Recog. Workshops (CVPRW)*, Salt Lake City, Utah, United States, Jun. 2018.
- [19] D. Donoho, “Compressed sensing,” *IEEE Trans. Inf. Theory*, vol. 52, no. 4, pp. 1289–1306, 2006.
- [20] E. J. Candès, J. Romberg, and T. Tao, “Robust uncertainty principles: Exact signal reconstruction from highly incomplete frequency information,” *IEEE Trans. Inf. Theory*, vol. 52, no. 2, pp. 489–509, 2006.
- [21] E. J. Candès and B. Recht, “Exact matrix completion via convex optimization,” *Found. Comput. Math.*, vol. 9, pp. 717–772, 2009.

- [22] R. Baraniuk and M. Wakin, “Random projections of smooth manifolds,” *Found. Comput. Math.*, vol. 9, pp. 51–77, 2009.
- [23] G. Ongie, R. Willett, R. Nowak, and L. Balzano, “Algebraic variety models for high-rank matrix completion,” in *Proc. Int. Conf. Mach. Learn. (ICML)*, Sydney, Australia, Aug. 2017, pp. 2691–2700.
- [24] H. Hendriks, “Nonparametric estimation of a probability density on a Riemannian manifold using Fourier expansions,” *Ann. Stat.*, vol. 18, no. 2, pp. 832–849, 1990.
- [25] B. Pelletier, “Kernel density estimation on riemannian manifolds,” *Stat. Probab. Lett.*, vol. 73, no. 3, pp. 297–304, 2005.
- [26] P. J. Bickel and B. Li, “Local polynomial regression on unknown manifolds,” in *Complex Datasets and Inverse Problems*. Beachwood, OH: Institute of Mathematical Statistics, 2007, pp. 177–186.
- [27] A. Aswani, P. Bickel, and C. Tomlin, “Regression on manifolds: Estimation of the exterior derivative,” *Ann. Stat.*, vol. 39, no. 1, pp. 48–81, 2011.
- [28] T. Hamm and I. Steinwart, “Adaptive learning rates for support vector machines working on data with low intrinsic dimension,” 2020. arXiv: [2003.06202 \[math.ST\]](#).
- [29] M. Chen, H. Jiang, W. Liao, and T. Zhao, “Nonparametric regression on low-dimensional manifolds using deep ReLU networks,” 2019. arXiv: [1908.01842 \[cs.LG\]](#).
- [30] R. Guhaniyogi and D. B. Dunson, “Compressed Gaussian process for manifold regression,” *J. Mach. Learn. Res.*, vol. 17, 2016.
- [31] R. Calandra, J. Peters, C. E. Rasmussen, and M. P. Deisenroth, “Manifold Gaussian processes for regression,” in *Proc. Joint Int. Conf. Neural Netw. (ICJNN)*, Vancouver, BC, Canada, Jul. 2016.
- [32] N. Dyn, F. J. Narcowich, and J. D. Ward, “Variational principles and Sobolev-type estimates for generalized interpolation on a riemannian manifold,” *Constr. Approx.*, vol. 15, no. 2, pp. 175–208, 1999.
- [33] H. Wendland, *Scattered Data Approximation*. Cambridge, 2005.
- [34] T. Hangelbroek, F. J. Narcowich, and J. D. Ward, “Kernel approximation on manifolds I: Bounding the Lebesgue constant,” *SIAM J. Math. Anal.*, vol. 42, no. 4, pp. 1732–1760, 2010.
- [35] T. Hangelbroek, F. J. Narcowich, X. Sun, and J. D. Ward, “Kernel approximation on manifolds II: The L_∞ norm of the L_2 projector,” *SIAM J. Math. Anal.*, vol. 43, no. 2, pp. 662–684, 2011.
- [36] T. Hangelbroek, F. J. Narcowich, and J. D. Ward, “Polyharmonic and related kernels on manifolds: Interpolation and approximation,” *Found. Comput. Math.*, vol. 12, no. 5, pp. 625–670, 2012.
- [37] A. Caponetto and E. De Vito, “Optimal rates for the regularized least-squares algorithm,” *Found. Comput. Math.*, vol. 7, no. 3, pp. 331–368, 2007.
- [38] I. Steinwart, D. Hush, and C. Scovel, “Optimal rates for regularized least squares regression,” in *Proc. Conf. Learn. Theory (COLT)*, Montreal, Canada, Jun. 2009.
- [39] S. Mendelson and J. Neeman, “Regularization in kernel learning,” *Ann. Stat.*, vol. 38, no. 1, pp. 526–565, 2010.
- [40] G. Blanchard and N. Mücke, “Optimal rates for regularization of statistical inverse learning problems,” *Found. Comput. Math.*, vol. 18, pp. 971–1013, 2018.
- [41] —, “Kernel regression, minimax rates, and effective dimensionality: Beyond the regular case,” *Anal. Appl.*, 2020, in press.
- [42] J. Lin, A. Rudi, L. Rosasco, and V. Cevher, “Optimal rates for spectral algorithms with least-squares regression over Hilbert spaces,” *Appl. Comput. Harmon. Anal.*, vol. 48, no. 3, pp. 868–890, 2020, in press.
- [43] S. Fischer and I. Steinwart, “Sobolev norm learning rates for regularized least-squares algorithms,” 2017. arXiv: [1702.07254 \[stat.ML\]](#).
- [44] F. Bauer, S. Pereverzev, and L. Rosasco, “On regularization in learning theory,” *J. Complexity*, vol. 23, pp. 52–72, 2007.
- [45] G. Blanchard and N. Krämer, “Optimal learning rates for kernel conjugate gradient regression,” in *Proc. Conf. Neural Inf. Process. Syst. (NeurIPS)*, vol. 23, Vancouver, Canada, 2010.
- [46] A. Dieuleveut, N. Flammarion, and F. Bach, “Harder, better, faster, stronger convergence rates for least-squares regression,” *J. Mach. Learn. Res.*, vol. 18, 2017.

- [47] A. Dieuleveut and F. Bach, “Nonparametric stochastic approximation with large step-sizes,” *Ann. Stat.*, vol. 44, no. 4, pp. 1363–1399, 2016.
- [48] Y. Zhang, J. Duchi, and M. Wainwright, “Divide and conquer kernel ridge regression: A distributed algorithm with minimax optimal rates,” *J. Mach. Learn. Res.*, vol. 16, pp. 3299–3340, 2015.
- [49] S.-B. Lin, X. Guo, and D.-X. Zhou, “Distributed learning with regularized least squares,” *J. Mach. Learn. Res.*, vol. 18, 2017.
- [50] Z.-C. Guo, S.-B. Lin, and D.-X. Zhou, “Learning theory of distributed spectral algorithms,” *Inverse Probl.*, vol. 33, 2017.
- [51] T. Zhang, “Learning bounds for kernel regression using effective data dimensionality,” *Neural Comput.*, vol. 17, pp. 2077–2098, 2005.
- [52] D. Hsu, S. M. Kakade, and T. Zhang, “Random design analysis of ridge regression,” *Found. Comput. Math.*, vol. 14, pp. 569–600, 2014.
- [53] L. H. Dicker, D. P. Foster, and D. Hsu, “Kernel ridge vs. principal component regression: Minimax bounds and the qualification of regularization operators,” *Electron. J. Stat.*, vol. 11, pp. 1022–1047, 2017.
- [54] —, “Kernel methods and regularization techniques for nonparametric regression: Minimax optimality and adaptation,” Technical report, Rutgers University, Tech. Rep., 2015.
- [55] F. Bach, “On the equivalence between kernel quadrature rules and random feature expansions,” *J. Mach. Learn. Res.*, vol. 18, 2017.
- [56] Y. Canzani and B. Hanin, “Scaling limit for the kernel of the spectral projector and remainder estimates in the pointwise Weyl law,” *Anal. PDE*, vol. 8, no. 7, pp. 1707–1731, 2015.
- [57] A. Jacot, F. Gabriel, and C. Hongler, “Neural tangent kernel: Convergence and generalization in neural networks,” in *Proc. Conf. Neural Inf. Process. Syst. (NeurIPS)*, Montréal, Canada, Dec. 2018.
- [58] S. Arora, S. S. Du, W. Hu, Z. Li, R. Salakhutdinov, and R. Wang, “On exact computation with an infinitely wide neural net,” in *Proc. Conf. Neural Inf. Process. Syst. (NeurIPS)*, Vancouver, Canada, Dec. 2019.
- [59] R. M. Neal, “Bayesian learning for neural networks,” Ph.D. dissertation, University of Toronto, 1995.
- [60] J. Lee, Y. Bahri, R. Novak, S. S. Schoenholz, J. Pennington, and J. Sohl-Dickstein, “Deep neural networks as Gaussian processes,” in *Proc. Int. Conf. Learn. Representations (ICLR)*, Vancouver, Canada, Apr.–May 2018.

A Proof of general RKHS results (Theorem 1)

We write \mathcal{P}_G and \mathcal{P}_{G^\perp} for the projections in L_2 and \mathcal{H} onto G and its orthogonal complement G^\perp , respectively.

For brevity, we denote by P_n the empirical measure given by the n independent samples of the variables (X, ξ) , i.e., $P_n w = \frac{1}{n} \sum_{i=1}^n w(X_i, \xi_i)$. For example, if $h: S \rightarrow \mathbf{R}$ is a function, $P_n h^2 = \frac{1}{n} \sum_{i=1}^n h^2(X_i)$, and $P_n \xi h = \frac{1}{n} \sum_{i=1}^n \xi_i h(X_i)$.

We use the following lemmas in our proof of Theorem 1:

Lemma 3. Let $\delta \in (0, 1)$. If

$$n \geq \max\{7, 3\gamma'\} K_p \log \frac{\max\{2, 4\gamma\} p}{\delta},$$

then, with probability at least $1 - \delta$,

$$P_n f^2 \geq \frac{1}{2} \|f\|_{L_2}^2 - 3\sqrt{t_{p+1}} \|f\|_{L_2} \|f\|_{\mathcal{H}}$$

for all $f \in \mathcal{H}$.

Lemma 4. There is a universal constant C such that, if

$$\frac{n}{\log^2 n} \geq C(1 \vee \gamma') \frac{K_p}{p} \frac{\|\xi\|_{\psi_1}^2}{\sigma^2},$$

then, with probability at least $1 - \delta$,

$$\begin{aligned} |P_n \xi f| &\leq \frac{3}{2} \sigma \cdot \left(\frac{\sqrt{p} + 2\sqrt{\log 4/\delta}}{\sqrt{n}} \|f\|_{L_2} + \frac{\sqrt{\text{tr } T_{G^\perp}} + 2\sqrt{t_{p+1} \log 4/\delta}}{\sqrt{n}} \|f\|_{\mathcal{H}} \right) \\ &\leq \frac{3}{2} \sigma \cdot \left(\frac{\sqrt{p} + 2\sqrt{\log 4/\delta}}{\sqrt{n}} \right) (\|f\|_{L_2} + \sqrt{\gamma t_{p+1}} \|f\|_{\mathcal{H}}). \end{aligned}$$

for all $f \in \mathcal{H}$.

With these, we prove the main result:

Proof of Theorem 1. We write our objective function as

$$F(f) = \frac{1}{n} \sum_{i=1}^n (Y_i - f(X_i))^2 + \alpha \|f\|_{\mathcal{H}}^2.$$

\hat{f} satisfies $\nabla F(\hat{f}) = 0$. Noting that

$$\frac{1}{2} \nabla F(f) = -\frac{1}{n} \sum_{i=1}^n (Y_i - f(X_i)) k(\cdot, X_i) + \alpha f,$$

we have

$$\begin{aligned} 0 &= \frac{1}{2} \langle \nabla F(\hat{f}), f^* - \hat{f} \rangle_{\mathcal{H}} \\ &= \left\langle \alpha \hat{f} - \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{f}(X_i)) k(\cdot, X_i), f^* - \hat{f} \right\rangle_{\mathcal{H}} \\ &= \alpha \langle \hat{f}, f^* - \hat{f} \rangle_{\mathcal{H}} - \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{f}(X_i)) (f^*(X_i) - \hat{f}(X_i)) \\ &= \alpha \langle \hat{f}, f^* - \hat{f} \rangle_{\mathcal{H}} + \frac{1}{n} \sum_{i=1}^n \left[(Y_i - f^*(X_i)) (\hat{f}(X_i) - f^*(X_i)) - (\hat{f}(X_i) - f^*(X_i))^2 \right] \\ &= \alpha \langle \hat{f}, f^* - \hat{f} \rangle_{\mathcal{H}} + P_n \xi (\hat{f} - f^*) - P_n (\hat{f} - f^*)^2. \end{aligned} \tag{4}$$

Let E_1 and E_2 denote the events of Lemmas 3 and 4. For part 1 of the theorem, we assume that E_1 holds, which occurs with probability at least $1 - \delta$. For part 2, we assume $E_1 \cap E_2$ holds, which occurs with probability at least $1 - 2\delta$. In what follows, we treat the two cases the same (and assume $\alpha > 0$), since we can simply take $\sigma = 0$ and the limit $\alpha \downarrow 0$ for part 1.

Let $e_2 = \|\hat{f} - f^*\|_{L_2}$ and $e_{\mathcal{H}} = \|\hat{f} - f^*\|_{\mathcal{H}}$. On $E_1 \cap E_2$, (4) implies

$$\frac{1}{2}e_2^2 \leq \sigma(ae_2 + be_{\mathcal{H}}) + ce_2e_{\mathcal{H}} + \alpha\langle \hat{f}, f^* - \hat{f} \rangle_{\mathcal{H}},$$

where $a = \frac{3}{2} \frac{\sqrt{p+2}\sqrt{\log 4/\delta}}{\sqrt{n}}$, $b = \sqrt{\gamma t_{p+1}}a$, and $c = 3\sqrt{t_{p+1}}$. First, note that

$$\langle \hat{f}, f^* - \hat{f} \rangle_{\mathcal{H}} = \langle f^*, f^* - \hat{f} \rangle_{\mathcal{H}} - e_{\mathcal{H}}^2 \leq \|f^*\|_{\mathcal{H}}e_{\mathcal{H}} - e_{\mathcal{H}}^2,$$

so

$$\sigma be_{\mathcal{H}} + \alpha\langle \hat{f}, f^* - \hat{f} \rangle_{\mathcal{H}} \leq (\sigma b + \alpha\|f^*\|_{\mathcal{H}})e_{\mathcal{H}} - \alpha e_{\mathcal{H}}^2 \leq \frac{(\sigma b + \alpha\|f^*\|_{\mathcal{H}})^2}{\alpha}.$$

To control the error term $ce_2e_{\mathcal{H}}$, we need a more explicit bound on $e_{\mathcal{H}}$. Because $P_n(\hat{f} - f^*)^2 \geq 0$, (4) gives

$$e_{\mathcal{H}}^2 \leq \|f^*\|_{\mathcal{H}}e_{\mathcal{H}} + \frac{1}{\alpha}P_n\xi(\hat{f} - f^*) \leq \|f^*\|_{\mathcal{H}}e_{\mathcal{H}} + \frac{\sigma}{\alpha}(ae_2 + be_{\mathcal{H}}).$$

Because $x^2 \leq a + bx$ implies $x \leq \sqrt{a} + b$, we then have

$$e_{\mathcal{H}} \leq \|f^*\|_{\mathcal{H}} + \frac{\sigma b}{\alpha} + \sqrt{\frac{\sigma ae_2}{\alpha}}.$$

Putting everything together, we have

$$\frac{1}{2}e_2^2 \leq \frac{(\sigma b + \alpha\|f^*\|_{\mathcal{H}})^2}{\alpha} + \sigma ae_2 + ce_2 \left(\|f^*\|_{\mathcal{H}} + \frac{\sigma b}{\alpha} + \sqrt{\frac{\sigma ae_2}{\alpha}} \right).$$

$x^2 \leq a + bx + cx^{3/2}$ implies $x \leq \sqrt{a} + b + c^2$, so

$$\begin{aligned} e_2 &\leq \sqrt{2} \frac{\sigma b}{\sqrt{\alpha}} + \sqrt{2\alpha}\|f^*\|_{\mathcal{H}} + 2\sigma a + 2c\|f^*\|_{\mathcal{H}} + 2\frac{\sigma cb}{\alpha} + 4\frac{c^2\sigma a}{\alpha} \\ &= (\sqrt{2\alpha} + 2c)\|f^*\|_{\mathcal{H}} + 2\sigma \left(a + \frac{b}{\sqrt{2\alpha}} + \frac{bc}{\alpha} + 2\frac{ac^2}{\alpha} \right). \end{aligned}$$

The result immediately follows by substituting our choices of a , b , and c and, if $\sigma \neq 0$, using the assumption that $\alpha \geq 54t_{p+1}$. \square

A.1 Proofs of key lemmas

Lemma 3 follows quickly from the following two concentration results:

Lemma 5. *If $\delta \in (0, 1)$, and $n \geq 7K_p \log \frac{p}{\delta}$, then, with probability at least $1 - \delta$, for all $f \in G$,*

$$P_n f^2 \geq \frac{1}{2} \|f\|_{L_2}^2.$$

Proof. Note that for all $f \in G$,

$$\begin{aligned} P_n f^2 &= \frac{1}{n} \sum_{i=1}^n f^2(X_i) \\ &= \frac{1}{n} \sum_{i=1}^n \langle Z(X_i), f \rangle_{L_2}^2 \\ &= \langle (P_n(Z \otimes_{L_2} Z))f, f \rangle_{L_2}, \end{aligned}$$

where we define $Z(X) = \sum_{\ell=1}^p v_\ell(X) v_\ell \in G$. The lemma will follow from a concentration result on $P_n(Z \otimes_{L_2} Z)$. Note that the operator $Z(X) \otimes_{L_2} Z(X) \succeq 0$ for all X , and, by Assumption 2, we have

$$\|Z(X) \otimes_{L_2} Z(X)\|_{L_2} = \|Z(X)\|_{L_2}^2 = \sum_{\ell=1}^p v_\ell^2(X) \leq K_p$$

almost surely. Also, $\mathbf{E} P_n(Z \otimes_{L_2} Z) = \mathbf{E} Z(X) \otimes_{L_2} Z(X) = \mathcal{I}_G$. The matrix Chernoff bound [1, Theorem 5.1.1] implies that, for all $\epsilon \in [0, 1)$,

$$\mathbf{P}(P_n(Z \otimes_{L_2} Z) \succeq (1 - \epsilon) \mathcal{I}_G) \geq 1 - p \left(\frac{e^{-\epsilon}}{(1 - \epsilon)^{1-\epsilon}} \right)^{n/K_p}.$$

Choosing $\epsilon = 1/2$ gives the result. \square

Lemma 6. *If $\delta \in (0, 1)$, and $n \geq \frac{3R_p}{t_{p+1}} \log \frac{2 \operatorname{tr} T_{G^\perp}}{t_{p+1}\delta}$, then, with probability at least $1 - \delta$, for all $f \in G^\perp$,*

$$P_n f^2 \leq 2t_{p+1} \|f\|_{\mathcal{H}}^2.$$

Proof. Similarly to the proof of Lemma 5, for all $f \in G^\perp$,

$$P_n f^2 = \langle (P_n(W \otimes_{\mathcal{H}} W))f, f \rangle_{\mathcal{H}},$$

where $W(X) = \sum_{\ell > p} t_\ell v_\ell(X) v_\ell$. Note that $\mathbf{E} W(X) \otimes_{\mathcal{H}} W(X) = T_{G^\perp}$. By Assumption 2,

$$\|W(X) \otimes_{\mathcal{H}} W(X)\|_{\mathcal{H}} = \|W(X)\|_{\mathcal{H}}^2 = \sum_{\ell > p} t_\ell v_\ell^2(X) \leq R_p$$

almost surely. By [1, Theorem 7.2.1], if $\epsilon \geq R_p/nt_{p+1}$, then

$$\mathbf{P}(\|P_n(W \otimes_{\mathcal{H}} W)\|_{\mathcal{H}} \leq (1 + \epsilon)t_{p+1}) \geq 1 - 2d_p \left(\frac{e^\epsilon}{(1 + \epsilon)^{1+\epsilon}} \right)^{nt_{p+1}/R_p},$$

where $d_p = \operatorname{tr} T_{G^\perp}/t_{p+1}$. Choosing $\epsilon = 1$ gives the result. \square

Proof of Lemma 3. Applying Lemmas 5 and 6 (with $\delta/2$ substituted for δ) and a union bound, we have, with probability at least $1 - \delta$,

$$\begin{aligned} \sqrt{P_n f^2} &\geq \sqrt{P_n(\mathcal{P}_G f)^2} - \sqrt{P_n(\mathcal{P}_{G^\perp} f)^2} \\ &\geq \frac{1}{\sqrt{2}} \|\mathcal{P}_G f\|_{L_2} - \sqrt{2t_{p+1}} \|\mathcal{P}_{G^\perp} f\|_{\mathcal{H}}, \end{aligned}$$

so

$$\begin{aligned} P_n f^2 &\geq \frac{1}{2} \|\mathcal{P}_G f\|_{L_2}^2 - 2\sqrt{t_{p+1}} \|\mathcal{P}_G f\|_{L_2} \|\mathcal{P}_{G^\perp} f\|_{\mathcal{H}} \\ &\geq \frac{1}{2} \|f\|_{L_2}^2 - \frac{1}{2} \|\mathcal{P}_{G^\perp} f\|_{L_2}^2 - 2\sqrt{t_{p+1}} \|f\|_{L_2} \|f\|_{\mathcal{H}} \\ &\geq \frac{1}{2} \|f\|_{L_2}^2 - 3\sqrt{t_{p+1}} \|f\|_{L_2} \|f\|_{\mathcal{H}}. \end{aligned}$$

\square

Proof of Lemma 4. Let B_2^G denote the L_2 -unit ball in G , and let $B_{\mathcal{H}}^{G^\perp}$ denote the \mathcal{H} -unit ball in G^\perp . Note that for all $f \in \mathcal{H}$, we have

$$f \in \|f\|_{L_2} B_2^G + \|f\|_{\mathcal{H}} B_{\mathcal{H}}^{G^\perp},$$

where the plus sign denotes Minkowski addition. Therefore, because $|P_n \xi f|$ is sublinear in f , it suffices to bound

$$Z_1 := \sup_{f \in B_2^G} |P_n \xi f|$$

and

$$Z_2 := \sup_{f \in B_{\mathcal{H}}^{G^\perp}} |P_n \xi f|.$$

We present a complete proof for the bound of Z_1 ; the proof for Z_2 is similar.

First, note that

$$\begin{aligned} Z_1 &= \sup_{f \in B_2^G} |P_n \xi f| \\ &= \sup_{\sum_{\ell=1}^p a_\ell^2 \leq 1} \left| P_n \left(\xi \sum_{\ell=1}^p a_\ell v_\ell \right) \right| \\ &= \sup_{\sum_{\ell=1}^p a_\ell^2 \leq 1} \left| \sum_{\ell=1}^p a_\ell P_n(\xi v_\ell) \right| \\ &= \left(\sum_{\ell=1}^p P_n^2(\xi v_\ell) \right)^{1/2}, \end{aligned}$$

so

$$\mathbf{E} Z_1 \leq \sqrt{\mathbf{E} Z_1^2} = \sqrt{\sum_{\ell=1}^p \mathbf{E} P_n^2(\xi v_\ell)} = \sigma \sqrt{\frac{p}{n}}.$$

We also have

$$\sup_{f \in B_2^G} \sum_{i=1}^n \mathbf{E}(\xi_i f(x_i))^2 = n\sigma^2.$$

Finally, note that

$$\sup_{f \in B_2^G} \|f\|_\infty \leq \sqrt{K_p},$$

so

$$\left\| \max_i \sup_{f \in B_2^G} |\xi_i f(x_i)| \right\|_{\psi_1} \leq \sqrt{K_p} \|\xi\|_{\psi_1} \log n.$$

Let $\eta \in (0, 1)$. [2, Theorem 4] (with, in the notation of that paper, $\delta = 1$) implies that, with probability at least $1 - \delta/2$,

$$Z_1 \leq \sigma \left((1 + \eta) \sqrt{\frac{p}{n}} + 2 \sqrt{\frac{\log 4/\delta}{n}} \right) + \frac{C'_\eta \sqrt{K_p} \|\xi\|_{\psi_1} (\log n) (\log 12/\delta)}{n}$$

for a constant C'_η that only depends on η . By a similar argument, we have, with probability at least $1 - \delta/2$,

$$Z_2 \leq \sigma \left((1 + \eta) \sqrt{\frac{\text{tr } T_{G^\perp}}{n}} + 2 \sqrt{\frac{t_{p+1} \log 4/\delta}{n}} \right) + \frac{C'_\eta \sqrt{R_p} \|\xi\|_{\psi_1} (\log n) (\log 12/\delta)}{n}.$$

Fixing $\eta \in (0, 1/2)$ and choosing a suitable constant C to ensure n is large enough completes the proof. \square

B Proof of heat kernel approximation (Lemma 1)

In this appendix, we prove upper and lower bounds on the heat kernel diagonal values. Although we only use the upper bound in our paper, we include the lower bound also as both may be of independent interest.

The concepts from differential geometry used in this section can be found in, for example, [3, 4]. The key tools we will use in our analysis of how well the heat kernel is approximated by a Gaussian RBF are the following *comparison theorems*:

Lemma 7 ([5, Theorem 4.5.1]). *If the sectional curvature of an m -dimensional manifold \mathcal{M} is bounded above by $K > 0$, then, for all $x, y \in \mathcal{M}$, $k_t^h(x, y) \leq k_t^{h,K}(d_{\mathcal{M}}(x, y))$, where $k_t^{h,K}(r)$ is the (radially symmetric) heat kernel on the m -dimensional space of constant curvature K , and, if $K > 0$, we set $k_t^{h,K}(r) = k_t^{h,K}(\pi/\sqrt{K})$ for $r \geq \pi/\sqrt{K}$.*

Lemma 8 ([5, Theorem 4.5.2]). *If the Ricci curvature of \mathcal{M} is bounded below by $(m-1)K$ for some constant K , then, for all $x, y \in \mathcal{M}$, $k_t^h(x, y) \geq k_t^{h,K}(d_{\mathcal{M}}(x, y))$, where $k_t^{h,K}(r)$ is the heat kernel on the space of constant curvature K .*

A lower bound of K on sectional curvature implies a lower bound of $(m-1)K$ on the Ricci curvature tensor (see, e.g., the formula for $\text{Ric}(v, v)$ in [4, p. 38]), so Lemma 8 also holds under the (stronger) assumption of a lower bound of K on sectional curvature.

The space of constant curvature $K > 0$ is the sphere $S_K^m = S^m/\sqrt{K}$, while the space of constant curvature $-K < 0$ is the scaled hyperbolic space $H_K^m = H^m/\sqrt{K}$. To apply Lemmas 7 and 8, we need to find bounds for the heat kernel on the sphere and on hyperbolic space.

We will use the following result:

Lemma 9 ([6, Theorem 1]). *The heat kernel in hyperbolic space H^m has the radial representation*

$$k_t^{h,H^m}(r) = e^{-\frac{(m-1)^2 t}{8}} \left(\frac{r}{\sinh r} \right)^{\frac{m-1}{2}} \frac{e^{-r^2/2t}}{(2\pi t)^{m/2}} \\ \times \mathbf{E}_r \exp \left(-\frac{(m-1)(m-3)}{8} \int_0^t \left(\frac{1}{\sinh^2 R_s} - \frac{1}{R_s^2} \right) ds \right),$$

where R_s is an m -dimensional Bessel process, and \mathbf{E}_r denotes expectation conditioned on $R_t = r$.

A nearly identical argument to that in [6] gives a corresponding result for the sphere S^m for $m \geq 2$:

Lemma 10. *For all $m \geq 2$, the heat kernel on the sphere S^m has the radial representation*

$$k_t^{h,S^m}(r) = e^{\frac{(m-1)^2 t}{8}} \left(\frac{r}{\sin r} \right)^{\frac{m-1}{2}} \frac{e^{-r^2/2t}}{(2\pi t)^{m/2}} \\ \times \mathbf{E}_r \exp \left(-\frac{(m-1)(m-3)}{8} \int_0^t \left(\frac{1}{\sin^2 R_s} - \frac{1}{R_s^2} \right) ds \right),$$

where, again, R_s is an m -dimensional Bessel process, and \mathbf{E}_r denotes expectation conditioned on $R_t = r$.

For $m \geq 3$, the exponent in the integrands in the formula of Lemma 9 (resp. Lemma 10) is always positive (resp. negative), so we have the following simple bounds on the heat kernels on the standard spaces of constant curvature:

$$k_t^{h,H^m}(r) \geq e^{-\frac{(m-1)^2 t}{8}} \left(\frac{r}{\sinh r} \right)^{\frac{m-1}{2}} \frac{e^{-r^2/2t}}{(2\pi t)^{m/2}}, \quad (5)$$

and

$$k_t^{h,S^m}(r) \leq e^{\frac{(m-1)^2 t}{8}} \left(\frac{r}{\sin r} \right)^{\frac{m-1}{2}} \frac{e^{-r^2/2t}}{(2\pi t)^{m/2}}. \quad (6)$$

It is easily verified that $p_t^{S_K^m}(r) = p_{Kt}^{S^m}(\sqrt{K}r)$, with a similar formula for scaled hyperbolic space. We can summarize this in the following result:

Lemma 11. *Suppose \mathcal{M} is an m -dimensional complete Riemannian manifold for $m \geq 3$.*

1. *Suppose \mathcal{M} has Ricci curvature bounded below by $-(m-1)K_1$. Then, for all $x, y \in \mathcal{M}$, denoting $r = d(x, y)$,*

$$k_t^h(x, y) \geq e^{-\frac{(m-1)^2}{8} K_1 t} \left(\frac{\sqrt{K_1} r}{\sinh(\sqrt{K_1} r)} \right)^{\frac{m-1}{2}} \frac{e^{-r^2/2t}}{(2\pi t)^{m/2}}.$$

2. Suppose \mathcal{M} has sectional curvature bounded above by K_2 . Then, for $r < \pi/\sqrt{K_2}$, and for all $x, y \in \mathcal{M}$ such that $d(x, y) \geq r$,

$$k_t^h(x, y) \leq e^{\frac{(m-1)^2}{8}K_2 t} \left(\frac{\sqrt{K_2}r}{\sin(\sqrt{K_2}r)} \right)^{\frac{m-1}{2}} \frac{e^{-r^2/2t}}{(2\pi t)^{m/2}}.$$

We note that, for $r = 0$ and t small, these results are comparable to the well-known asymptotic expansion for the heat kernel, which depends on the scalar curvature at x (see, e.g., [7, Section VI.4]).

Finally, we specialize to the case $r = 0$ and simplify:

Proposition 1. *Let $\epsilon \leq 2/3$.*

1. *Under the conditions of Lemma 11.1, for $t \leq \frac{8\epsilon}{(m-1)^2 K_1}$ and all $x \in \mathcal{M}$,*

$$k_t^h(x, x) \geq \frac{1 - \epsilon}{(2\pi t)^{m/2}}.$$

2. *Under the conditions of Lemma 11.2, for $t \leq \frac{6\epsilon}{(m-1)^2 K_2}$ and all $x \in \mathcal{M}$,*

$$k_t^h(x, x) \leq \frac{1 + \epsilon}{(2\pi t)^{m/2}}.$$

Proof. From Lemma 11, we have

$$e^{-\frac{(m-1)^2}{8}K_1 t} \leq (2\pi t)^{-m/2} k_t^h(x, x) \leq e^{\frac{(m-1)^2}{8}K_2 t}.$$

The result follows from noting that $e^{-s} \geq 1 - s$ for all $s \geq 0$, and $e^s \leq 1 + \frac{4}{3}s$ for $0 \leq s \leq 1/2$. \square

Lemma 1 is a case of this last result, taking $K_2 = \kappa$.

C Proof of non-asymptotic Weyl law estimates (Theorem 2 and Lemma 2)

Proof of Theorem 2. By Lemma 1, for all $\lambda \geq 0$ and $t \leq \frac{6\epsilon}{(m-1)^2 \kappa}$,

$$\begin{aligned} e^{-\lambda t/2} N_x(\lambda) &= e^{-\lambda t/2} \sum_{\lambda_\ell \leq \lambda} v_\ell^2(x) \\ &\leq \sum_{\ell=0}^{\infty} e^{-\lambda_\ell t/2} v_\ell^2(x) \\ &= k_t^h(x, x) \\ &\leq \frac{1 + \epsilon}{(2\pi t)^{m/2}}. \end{aligned}$$

Taking $t = m/\lambda$, we get

$$\begin{aligned} N_x(\lambda) &\leq \frac{(1 + \epsilon)e^{\lambda t/2}}{(2\pi t)^{m/2}} \\ &= \frac{1 + \epsilon}{(4\pi)^{m/2}} \frac{e^{m/2}}{(m/2)^{m/2}} \lambda^{m/2} \\ &\leq \frac{1 + \epsilon}{(4\pi)^{m/2}} \frac{2\sqrt{m}}{\Gamma(\frac{m}{2} + 1)} \lambda^{m/2} \\ &= \frac{2(1 + \epsilon)\sqrt{m}}{(2\pi)^m} V_m \lambda^{m/2}, \end{aligned}$$

where the second inequality uses Stirling's approximation. \square

Proof of Lemma 2. For $c \in (0, 1)$, note that

$$\begin{aligned}
\sum_{\lambda_\ell \geq \lambda} e^{-\lambda_\ell t/2} v_\ell^2(x) &\leq e^{-(1-c)\lambda t/2} \sum_{\lambda_\ell \geq \lambda} e^{-c\lambda_\ell t/2} v_\ell^2(x) \\
&\leq e^{-(1-c)\lambda t/2} \sum_{k=0}^{\infty} e^{-c\lambda_\ell t/2} v_\ell^2(x) \\
&= e^{-(1-c)\lambda t/2} p_{ct}^{\mathcal{M}}(x, x) \\
&\leq e^{-\lambda t/2} (1 + \epsilon) \frac{e^{c\lambda t/2}}{(2\pi ct)^{m/2}}.
\end{aligned}$$

Choosing $c = m/\lambda t$, the remainder of the proof is identical to that of Theorem 2. \square

D Proof of manifold regression results (Theorems 3 and 4)

Proof of Theorems 3 and 4. To apply the framework of Sections 2.1 and 4.1, which assumes the set S has measure 1, we consider the normalized volume measure $d\tilde{V} = dV / \text{vol } \mathcal{M}$. With respect to \tilde{V} , k_t^h has the eigenvalue decomposition

$$k_t^h(x, y) = \frac{1}{\text{vol } \mathcal{M}} \sum_{\ell} e^{-\lambda_\ell t/2} \tilde{u}_\ell(x) \tilde{u}_\ell(y),$$

where $\tilde{u}_\ell = \sqrt{\text{vol } \mathcal{M}} u_\ell$. A similar normalized expansion holds for k_Ω^{bl} .

Note that Theorem 2 and Lemma 2 only give us bounds on the constants K_p and R_p in Assumption 2. For k_Ω^{bl} , this holds with $K_p = p(\Omega)$ (taking $\epsilon = 1/2$ in Theorem 2) and $R_p = 0$. Assumption 3 holds trivially with $\gamma = \gamma' = 0$.

For k_t^h , we can again take $K_p = p(\Omega)$ (again taking $\epsilon = 1/2$), and we get a bound on R_p such that $\gamma = \gamma' = 1$.

Finally, for both kernels, we take into account the fact that $\|\cdot\|_{L_2(\mathcal{M}, \tilde{V})} = \|\cdot\|_{L_2(\mathcal{M}, V)} / \sqrt{\text{vol } \mathcal{M}}$. With these considerations in mind, the results follow from Theorem 1. \square

References

- [1] J. Tropp, “An introduction to matrix concentration inequalities,” *Found. Trends Mach. Learn.*, vol. 8, no. 1-2, pp. 1–230, 2015.
- [2] R. Adamczak, “A tail inequality for suprema of unbounded empirical processes with applications to Markov chains,” *Electron. J. Probab.*, vol. 13, pp. 1000–1034, 2008.
- [3] J. M. Lee, *Introduction to Riemannian Manifolds*, 2nd ed. Springer, 2018.
- [4] P. Petersen, *Riemannian Geometry*, 3rd ed. Springer, 2016.
- [5] E. Hsu, *Stochastic Analysis on Manifolds*. Providence, RI: American Mathematical Society, 2002.
- [6] X. Cheng and T.-H. Wang, “Bessel bridge representation for the heat kernel in hyperbolic space,” *Proc. Amer. Math. Soc.*, vol. 146, no. 4, pp. 1781–1792, 2018.
- [7] I. Chavel, *Eigenvalues in Riemannian Geometry*. Academic Press, 1984.