

ASYMPTOTIC REGULARITY AND ATTRACTORS FOR SLIGHTLY COMPRESSIBLE BRINKMAN-FORCHEIMER EQUATIONS

VARGA KALANTAROV^{1,2} AND SERGEY ZELIK^{3,4}

ABSTRACT. Slightly compressible Brinkman-Forchheimer equations in a bounded 3D domain with Dirichlet boundary conditions are considered. These equations model fluids motion in porous media. The dissipativity of these equations in higher order energy spaces is obtained and regularity and smoothing properties of the solutions are studied. In addition, the existence of a global and an exponential attractors for these equations in a natural phase space is verified.

CONTENTS

1. Introduction	1
2. Well-posedness, dissipativity and partial smoothing	4
3. Asymptotic regularity	9
4. Attractors	17
5. Generalizations and concluding remarks	21
Appendix A. An auxiliary linear problem	24
References	28

1. INTRODUCTION

We give a comprehensive study of slightly compressible Brinkman-Forchheimer equations in the following form:

$$(1.1) \quad \begin{cases} \partial_t u - \Delta_x u + \nabla_x p + f(u) = g, & u|_{\partial\Omega} = 0, \quad u|_{t=0} = u_0, \\ \partial_t p + \operatorname{div}(Du) = 0, & p|_{t=0} = p_0 \end{cases}$$

in a bounded domain $\Omega \subset \mathbb{R}^3$ with sufficiently smooth boundary $\partial\Omega$. Here $u = (u^1(t, x), u^2(t, x), u^3(t, x))$ and $p = p(t, x)$ are unknown velocity vector field and pressure respectively, D is a given positive self-adjoint matrix, f is a given nonlinearity and g is the external force.

Equations of the form (1.1) arise in the mathematical theory of fluids in porous media and are of a big permanent interest from both theoretical and

2010 *Mathematics Subject Classification.* 35B40, 35B45, 35K10.

Key words and phrases. Brinkman-Forchheimer equations, compressible fluid, tidal equations, dissipativity, global attractor, exponential attractor, regularity of solutions, localization.

This work is partially supported by the RSF grant 19-71-30004 as well as the EPSRC grant EP/P024920/1.

applied points of view, see [1, 3, 10, 12, 14, 18, 19, 25, 29, 32, 34, 36, 38] and references therein. The first equation of (1.1) is usually interpreted as a generalization of the Darcy law:

$$\tau \partial_t v - \beta \Delta_x v + f(v) = -D \nabla_x p,$$

where D is a normalized permeability tensor, $f(v)$ is a Forchheimer nonlinearity which typically has a form

$$(1.2) \quad f(v) = \alpha v + \beta (\mathcal{C}v \cdot v)^l v + \gamma \sqrt{(\mathcal{C}v \cdot v)} v,$$

where $\mathcal{C}$ is another positive self-adjoint matrix and α, β, γ and $l \geq \frac{1}{2}$ are some constants, see e.g., [1] and references therein, $\beta \Delta_x v$ is a Brinkman term with effective viscosity parameter $\beta > 0$, see [3] and $\tau \partial_t v$ is time relaxation term which is especially important in the case of non-monotone f or/and presence of the inertial term $(v, \nabla_x)v$ to provide the unique expression of v through $\nabla_x p$. The second equation

$$\partial_t p + \operatorname{div}(v) = 0$$

is just a standard slightly compressible approximation of the continuity equation, see e.g., [36, 22, 5] and references therein. Making the change of variables $v = Du$, we end up with a system of the form (1.1) with a slightly unusual term $\operatorname{div}(Du)$.

We also mention that the equations (1.1) in 2D case naturally arise in the dynamic theory of tides as a generalization of the classical Laplace tidal equations. In this case, $u := (u^1, u^2)$ is the horizontal transport vector (the horizontal velocity averaged over the vertical axis) and the scalar p is a vertical tidal elevation, see, e.g., [13, 15, 21, 24, 28] and references therein.

Equations (1.1) have a non-trivial structure which is interesting also from purely mathematical point of view. Indeed, in the simplest case $f = g = 0$ $D = 1$, we may introduce a new variable $\omega = \operatorname{curl} u$ and reduce the system to the following equations

$$\partial_t \omega - \Delta_x \omega = 0, \quad \partial_t^2 p - \Delta_x \partial_t p - \Delta_x p = 0,$$

so we see a combination of a heat equation with the so-called strongly damped wave equation. This system is decoupled in the case of periodic boundary conditions, but in the case of Dirichlet boundary conditions we have a non-trivial coupling already on the level of linear equations through boundary conditions. Thus, in contrast to the incompressible case, one cannot expect instantaneous smoothing property for u and p , but similarly to damped wave equation, one can expect that some components of the solution may have this property, see [17] for more details. Of course, for non-zero nonlinearity f , we also have coupling through nonlinear terms.

Another possibility is to differentiate the first equation in time and exclude the pressure using the second equation. This gives the second order in time equation:

$$\partial_t^2 u - \Delta_x \partial_t u + f'(u) \partial_t u - \nabla_x \operatorname{div}(Du) = 0$$

which is again a sort of strongly damped wave equation with the nonlinearity of Van der Pol type, see e.g., [16] for the regularity and longtime behavior of such equations in the scalar case. However, this form of equations (1.1)

is not convenient especially for the study of longtime behavior since the operator $\nabla_x \operatorname{div}(Du)$ is degenerate.

The longtime behavior of solutions to *incompressible* Brinkman-Forchheimer or Brinkman-Forchheimer-Navier-Stokes equations (often also referred as tamed Navier-Stokes equations) is studied in many papers, see [14, 18, 25, 39, 40] and references therein. However, the slightly compressible case is essentially less understood. To the best of our knowledge, similar problems have been considered only in 2D case only for slightly compressible Navier-Stokes equations, see [12] and [10] for global and exponential attractors respectively, but even in this case, Dirichlet's boundary conditions were out of consideration because of the problems with obtaining dissipative estimates for the pressure in $H^1(\Omega)$ which are caused by "bad" boundary terms in higher energy estimates.

The aim of the present paper is to verify the global well-posedness and dissipativity of the problem (1.1) in the initial phase space

$$(u_0, p_0) \in E := H_0^1(\Omega) \times \bar{L}^2(\Omega), \quad \bar{L}^2(\Omega) := \{p_0 \in L^2(\Omega), \langle p_0 \rangle = 0\},$$

where $\langle v \rangle$ is a mean value of the function $v(x)$ as well as in the higher energy space

$$E^1 = E \cap (H^2(\Omega) \times H^1(\Omega))$$

and to prove the existence of global and exponential attractors for the associated solution semigroup. Note that, similarly to [12], we are unable to verify the dissipativity in E^1 using the energy-type estimates because of the appearance of "bad" boundary integrals. We overcome this problem using the combination of partial instantaneous smoothing property and localization technique inspired by [17]. Actually, the localization technique is used here in a bit non-standard way, since it is usually applied to verify the higher regularity. In our situation, this higher regularity is more or less straightforward and the localization is used in order to get the dissipative estimate only, see Appendix A for more details.

Throughout of the paper, we assume that the external force $g \in L^2(\Omega)$ and the nonlinearity $f(u)$ has the following form

$$(1.3) \quad f(u) := \varphi(|u|^2)u,$$

where $\varphi \in C^1((0, \infty))$ and satisfies the conditions:

$$(1.4) \quad \begin{cases} 1. & K - Cz^{-1/2} \leq \varphi'(z) \leq C_1 z^{-1/2}(1 + z^{3/2}), \\ 2. & -C + \alpha z^l \leq \varphi(z) \leq C(1 + z^l), \quad z \in \mathbb{R}_+ \end{cases}$$

for some positive constants α, K, C, C_1 and the exponent $l \in (0, 2]$.

Clearly these conditions are satisfied for the typical nonlinearity (1.2) if $\mathcal{C} = 1$ (or $DCD = 1$ if we take into the account the change of variables mentioned above). The case of a general self-adjoint positive $\mathcal{C}$ is completely analogous, we only need to take $\mathcal{C}u \cdot u$ instead of $|u|^2$ in (1.3), we assume that $\mathcal{C} = 1$ only for simplicity. In contrast to this, the extra assumption that $D = 1$ somehow oversimplifies the problem since some additional energy type identities hold in this particular case, so we prefer to keep a general matrix D . We also mention that the exponent $-1/2$ is fixed in (1.4) in order to handle the term $\sqrt{\mathcal{C}u \cdot u}u$ in (1.2). Of course, if $l = \frac{1}{2}$, we need

to assume that $\gamma + \beta > 0$ in (1.2) in order to get dissipativity. Analogously, for $l > \frac{1}{2}$, we need to assume that $\beta > 0$.

The paper is organized as follows. In §1 we derive the basic dissipative estimate for problem (1.1) in the energy phase space E , verify the existence and uniqueness of solutions and prove, some instantaneous regularization for the u component of the solution. Namely, we establish that, starting from $(u(0), p(0)) \in E$, at the next time moment t , we will have $\partial_t u(t) \in L^2(\Omega)$ and $\nabla_x u(t) \in L^2(\Omega)$. This regularization allows us, similarly to the case of strongly damped wave equations, to truncate system (1.1) and reduce the analysis to simpler equations:

$$(1.5) \quad \partial_t p + \operatorname{div}(Du) = 0, \quad p|_{t=0} = p_0, \quad -\Delta_x u + \nabla_x p + f(u) = g(t),$$

where $g \in L^\infty(\mathbb{R}_+, L^2(\Omega))$ is new given external force (of course, the relation of this system to the initial equations (1.1) is given by $g(t) := g - \partial_t u(t)$).

The detailed analysis of this truncated system is presented in §3. In particular, we prove there that this system is well-posed and dissipative in higher energy space $p \in \bar{L}^2(\Omega) \cap H^1(\Omega)$ and also establish the exponential smoothing property for this system, namely, we check that the ball in the space $H^1(\Omega)$ attracts exponentially fast the trajectories $p(t)$ of (1.5) starting from bounded sets of $\bar{L}^2(\Omega)$. Returning back to the full system (1.1), we establish after that its well-posedness and dissipativity in higher energy space E^1 as well as the fact that the proper ball in E^1 is an exponentially attracting set for the solutions of (1.1) starting from E . This fact, in turn, is crucial for our study of global and exponential attractors.

Note also that the analysis presented in this section is heavily based on the study of linear problem (1.5) (which corresponds to $f = 0$) presented in Appendix A and, in particular, on the dissipativity of this linear problem in higher energy space $H^1(\Omega)$. This dissipativity is proved using the localization technique and is of independent interest.

In §4 we verify the existence of a global and exponential attractors for the solution semigroup associated with problem (1.1). These results are more or less standard corollaries of the asymptotic regularity and exponential attraction proved in §3, see [2, 4, 6, 8, 27, 35] for more details.

Finally, in §5, we also consider briefly some generalizations of the proved results, including the case of the extra convective terms in the initial Brinkman-Forchheimer equation and discuss some open problems for further research.

2. WELL-POSEDNESS, DISSIPATIVITY AND PARTIAL SMOOTHING

In this section, we verify the global well-posedness and dissipativity of slightly compressible Brinkman-Forchheimer equations:

$$(2.1) \quad \begin{cases} \partial_t u - \Delta_x u + \nabla_x p + f(u) = g, & u|_{\partial\Omega} = 0, \quad u|_{t=0} = u_0, \\ \partial_t p + \operatorname{div}(Du) = 0, & p|_{t=0} = p_0 \end{cases}$$

in the energy space $(u_0, p_0) \in E$ as well as establish some partial smoothing results for the solutions of this system which are crucial for what follows. We start with the basic a priori estimate in the phase space E .

Theorem 2.1. *Let $g \in L^2(\Omega)$, $D = D^* > 0$ and the nonlinearity f satisfy (1.3) and (1.4). Let also $(u(t), p(t))$ be a sufficiently smooth solution of (2.1). Then, the following estimate holds:*

$$(2.2) \quad \|(u, p)(t)\|_E^2 + \int_t^{t+1} (\|\nabla_x u(s)\|_{L^2}^2 + (|f(u(s)) \cdot Du(s)|, 1)) ds \leq \\ \leq Q(\|(u, p)(0)\|_E^2) e^{-\alpha t} + Q(\|g\|_{L^2}^2),$$

for some monotone function Q and positive constant α independent on u and t .

Proof. We multiply the first equation of (2.1) by Du and integrate over Ω . Then, integrating by parts and using the second equation, we arrive at

$$(2.3) \quad \frac{1}{2} \frac{d}{dt} \left(\|u\|_{L_D^2}^2 + \|p\|_{\bar{L}^2}^2 \right) + \|\nabla_x u\|_{L_D^2}^2 + (f(u), Du) = (g, Du),$$

where $\|u\|_{L_D^2}^2 := \int_{\Omega} Du(x) \cdot u(x) dx$. Here and below $\xi \cdot \eta$ stands for the standard dot product of vectors $\xi, \eta \in \mathbb{R}^3$.

This energy identity is still not enough to get the dissipative estimate since it does not contain the term $\|p\|_{L^2}^2$ without time differentiation. To get this term we use the so-called Bogovski operator:

$$(2.4) \quad \mathfrak{B} : \bar{L}^2(\Omega) \rightarrow H_0^1(\Omega), \quad \bar{L}^2(\Omega) := \{p \in L^2(\Omega), \langle p \rangle = 0\}, \quad \operatorname{div} \mathfrak{B}p = p.$$

It is well-known that such an operator exists as a linear continuous operator if Ω is smooth enough, see e.g., [33]. Multiplying the first equation of (2.1) by $\mathfrak{B}p$, integrating with respect to x and using the second equation, we get

$$(2.5) \quad \frac{d}{dt} (u, \mathfrak{B}p) + \|p\|_{\bar{L}^2}^2 = -(p, \mathfrak{B} \operatorname{div}(Du)) - \\ - (\nabla_x u, \nabla_x \mathfrak{B}p) - (f(u), \mathfrak{B}p) + (g, \mathfrak{B}p).$$

Multiplying (2.5) by a small $\varepsilon > 0$ and taking a sum with equation (2.3), after using the Hölder inequality and Sobolev embedding $H^1 \subset L^6$, we get

$$(2.6) \quad \frac{d}{dt} \left(\|u\|_{L_D^2}^2 + \|p\|_{\bar{L}^2}^2 + 2\varepsilon(u, \mathfrak{B}p) \right) + \\ + \|\nabla_x u\|_{L_D^2}^2 + \varepsilon\|p\|_{\bar{L}^2}^2 + (f(u), Du) \leq \varepsilon\|f(u)\|_{L^{6/5}}\|p\|_{\bar{L}^2} + C(\|g\|_{L^2}^2 + 1).$$

Using our assumptions (1.4) on functions f and φ , it is not difficult to verify that

$$(2.7) \quad |f(u)|^{6/5} \leq C(|f(u) \cdot Du| + 1).$$

This gives us the following differential inequality:

$$(2.8) \quad \frac{d}{dt} \mathcal{E}_{\varepsilon}(u, p) + \varepsilon \mathcal{E}_{\varepsilon}(u, p) \leq C\varepsilon^6 \mathcal{E}_{\varepsilon}^3(u, p) + C(\|g\|_{L^2}^2 + 1),$$

where

$$(2.9) \quad \mathcal{E}_{\varepsilon}(u, p) := \|u\|_{L_D^2}^2 + \|p\|_{\bar{L}^2}^2 + 2\varepsilon(u, \mathfrak{B}p).$$

Moreover, for sufficiently small $\varepsilon > 0$, we have

$$\frac{1}{2} \|(u, p)\|_E^2 \leq \mathcal{E}_{\varepsilon}(u, p) \leq \frac{3}{2} \|(u, p)\|_E^2.$$

Thus, applying the Gronwall lemma with a parameter to (2.8), see [11, 31] and also [42] for details, we end up with the desired estimate (2.2) and finishes the proof of the theorem. $\square$

At the next step we define a weak energy solution for problem (2.1).

Definition 2.2. A pair of function $(u, p) \in C_w(0, T; E)$ is a weak energy solution of problem (2.1) if, in addition,

$$(2.10) \quad u \in L^2(0, T; H_0^1(\Omega)) \cap L^{2(l+2)}(0, T; L^{2(l+2)}(\Omega))$$

and the equations (2.1) are satisfied in the sense of distributions.

Remark 2.3. From the first equation (2.1), we see that

$$(2.11) \quad \partial_t u \in L^2(0, T; H^{-1}(\Omega)) + L^{l'}(0, T; L^{l'}(\Omega)) = \\ = [L^2(0, T; H_0^1(\Omega)) \cap L^{2(l+1)}(0, T; L^{2(l+1)}(\Omega))]^*,$$

where $\frac{1}{l'} + \frac{1}{2(l+1)} = 1$. Thus, multiplication of the first equation of (2.1) by u or $\mathfrak{B}p$ is justified on the level of weak energy solutions and we have, in addition, that $u \in C(0, T; L^2(\Omega))$, see e.g., [4] for the details. The situation with p -component is even simpler since we have

$$\partial_t p \in L^2(0, T; \bar{L}^2(\Omega))$$

and multiplication on p is allowed. So, we also have that $p \in C(0, T; \bar{L}^2(\Omega))$. In particular, all manipulations done for the derivation of the key estimate (2.8) are actually justified for weak energy solutions, so all such solutions satisfy the dissipative estimate (2.2).

We now turn to the uniqueness.

Theorem 2.4. *Suppose that the assumptions of Theorem 2.1 are satisfied and let $(u_1(t), p_1(t))$ and $(u_2(t), p_2(t))$ be two weak energy solutions of problem (2.1). Then, the following estimate holds:*

$$(2.12) \quad \|(u_1(t) - u_2(t), p_1(t) - p_2(t))\|_E^2 \leq \\ \leq Ce^{Kt} \|(u_1(0) - u_2(0), p_1(0) - p_2(0))\|_E^2,$$

where the constants C and K depend only on f and D .

Proof. We first note that it suffices to verify (2.12) for $t \leq T$ for some small, but positive T . Then, to get the general estimate, it will be enough to iterate (2.1). Let $\bar{u}(t) := u_1(t) - u_2(t)$ and $\bar{p}(t) := p_1(t) - p_2(t)$. Then, these functions solve

$$(2.13) \quad \partial_t \bar{u} - \Delta_x \bar{u} + \nabla_x \bar{p} + [f(u_1) - f(u_2)] = 0, \quad \partial_t \bar{p} + \operatorname{div}(D\bar{u}) = 0.$$

Integrating the second equation, we get

$$(2.14) \quad \bar{p}(t) = \bar{p}(0) - \int_0^t \operatorname{div}(D\bar{u}(s)) ds.$$

Multiplying now the first equation by $\bar{u}(t)$ and using that, due to assumptions (1.3) and (1.4), $f'(u) \geq -L$, after the standard transformations, we

end up with

$$(2.15) \quad \frac{d}{dt} \|\bar{u}(t)\|_{L^2}^2 + \|\nabla_x \bar{u}\|_{L^2}^2 + 2 \left(\int_0^t \operatorname{div}(D\bar{u}(s)) \, ds, \operatorname{div} \bar{u}(t) \right) \leq C \|\bar{p}(0)\|_{L^2}^2 + 2L \|\bar{u}(t)\|_{L^2}^2.$$

Assuming that T is small enough, we estimate

$$(2.16) \quad \left| 2 \left(\int_0^t \operatorname{div}(D\bar{u}(s)) \, ds, \operatorname{div} \bar{u}(t) \right) \right| \leq \int_0^t \|\operatorname{div}(D\bar{u}(s))\|_{L^2}^2 \, ds + T \|\operatorname{div}(\bar{u}(t))\|_{L^2}^2 \leq C \int_0^t \|\nabla_x \bar{u}(s)\|_{L^2}^2 \, ds + \frac{1}{2} \|\nabla_x \bar{u}(t)\|_{L^2}^2$$

and, therefore,

$$(2.17) \quad \frac{d}{dt} \|\bar{u}(t)\|_{L^2}^2 + \frac{1}{2} \|\nabla_x \bar{u}\|_{L^2}^2 - C \int_0^t \|\nabla_x \bar{u}(s)\|_{L^2}^2 \, ds \leq C \|\bar{p}(0)\|_{L^2}^2 + 2L \|\bar{u}(t)\|_{L^2}^2.$$

Integrating this inequality in time, we end up with

$$(2.18) \quad \|\bar{u}(t)\|_{L^2}^2 + \int_0^t \left(\frac{1}{2} - C(t-s) \right) \|\nabla_x \bar{u}(s)\|_{L^2}^2 \, ds \leq C \|(\bar{u}(0), \bar{p}(0))\|_E^2 + 2L \int_0^t \|\bar{u}(s)\|_{L^2}^2 \, ds.$$

Fixing now T small enough that the integral in the left-hand side is positive and applying the Gronwall inequality, we finally arrive at

$$(2.19) \quad \|\bar{u}(t)\|_{L^2}^2 + \int_0^t \|\nabla_x \bar{u}(s)\|_{L^2}^2 \, ds \leq K \|(\bar{u}(0), \bar{p}(0))\|_{L^2}^2, \quad t \leq T.$$

The corresponding estimate for the p -component follows now from (2.14). Thus, the estimate (2.12) is verified and the theorem is proved. $\square$

Corollary 2.5. *Let the assumptions of Theorem 2.1 hold. Then, equations (2.1) generate a dissipative globally Lipschitz continuous semigroup $S(t)$ in the phase space E :*

$$(2.20) \quad S(t)(u_0, p_0) := (u(t), p(t)),$$

where $(u(t), p(t))$ is a unique energy solution of (2.1) with the initial data $(u_0, p_0) \in E$.

Proof. According to theorems 2.1 and 2.4, we only need to verify the existence of a weak solution. This can be done in many standard ways, one of the is to use vanishing viscosity method. Namely, we may approximate (2.1) by a family of parabolic equations:

$$\partial_t u - \Delta_x u + \nabla_x p = g, \quad \partial_t p + \operatorname{div}(Du) = \nu \Delta_x p, \quad u|_{\partial\Omega} = 0, \quad \partial_n p|_{\partial\Omega} = 0,$$

where $\nu > 0$ is a small parameter. The solution of this parabolic problem can be obtained using e.g., the Galerkin approximations and the passage to the limit as $\nu \rightarrow 0$ is also straightforward since the analogue of (2.3) gives the necessary uniform with respect to $\nu \rightarrow 0$ estimates (although they are non-dissipative, this is not important for the existence of a solution on a finite time interval). So, we omit the details here. $\square$

By the analogy with strongly damped wave equation (see [30, 17] and references therein), one may expect that (2.1) partially possesses instantaneous smoothing property. The next results shows that such a smoothing indeed holds.

Theorem 2.6. *Let the assumptions of Theorem 2.1 hold and let (u, p) be a weak energy solution of problem (2.1). Then the following partial smoothing property holds:*

$$(2.21) \quad t\|\nabla_x u(t)\|_{L^2}^2 + t^2\|\partial_t u(t)\|_{L^2}^2 + t\|\partial_t p(t)\|_{L^2}^2 + \int_0^t s^2\|\nabla_x \partial_t u(s)\|_{L^2}^2 ds \leq C(1 + \|g\|_{L^2}^2 + \|(u(0), p(0))\|_E^2),$$

where $t \in [0, 1]$ and the constant C is independent of t and u .

Proof. Let us first multiply the first equation of (2.1) by $t\partial_t u$ and integrate over Ω . Then, using the gradient structure of nonlinearity f , we arrive at

$$(2.22) \quad \frac{d}{dt} \left(\frac{t}{2} \|\nabla_x u\|_{L^2}^2 + t(F(u), 1) - t(p, \operatorname{div} u) \right) + t\|\partial_t u\|_{L^2}^2 = - (p, \operatorname{div} u) + \frac{1}{2} \|\nabla_x u\|_{L^2}^2 + (F(u), 1) + t(\operatorname{div}(Du), \operatorname{div} u) + t(g, \partial_t u).$$

where $F(u) := \frac{1}{2} \int_0^{|u|^2} \varphi(z) dz$. Integrating this identity in time and using the estimate (2.2), we arrive at the following smoothing property:

$$(2.23) \quad t\|\nabla_x u(t)\|_{L^2}^2 + t\|u(t)\|_{L^{2(l+1)}}^{2(l+1)} + t\|\partial_t p(t)\|_{L^2}^2 + \int_0^t s\|\partial_t u(s)\|_{L^2}^2 ds \leq C(\|(u(0), p(0))\|_E^2 + 1 + \|g\|_{L^2}^2),$$

where $t \in [0, 1]$ and C is independent of u , p and t .

Let us now differentiate equations (2.1) in time and denote $v := \partial_t u$ and $q = \partial_t p$. Then, we end up with the following equations

$$(2.24) \quad \partial_t v - \Delta_x v + \nabla_x q + f'(u)v = 0, \quad \partial_t q + \operatorname{div}(Dv) = 0.$$

Multiplying the first equation of (2.24) by $t^2 v$ and integrating over Ω , we get

$$(2.25) \quad \frac{1}{2} \frac{d}{dt} (t^2 \|v(t)\|_{L^2}^2) + t^2 \|\nabla_x v\|_{L^2}^2 + t^2 (f'(u)v, v) = t^2 (\partial_t p, \operatorname{div} v) + t\|\partial_t u\|_{L^2}^2.$$

Integrating this equality in time and using (2.23) together with the assumption $f'(u) \geq -L$, we get the desired smoothing property in the form

$$(2.26) \quad t^2 \|\partial_t u(t)\|_{L^2}^2 + \int_0^t s^2 \|\nabla_x \partial_t u(s)\|_{L^2}^2 ds \leq \\ \leq C(1 + \|g\|_{L^2}^2 + \|(u(0), p(0))\|_E^2),$$

where $t \in [0, 1]$ and finish the proof of the theorem. $\square$

Combining smoothing estimate (2.21) with the dissipative estimate (2.2), we get the following result.

Corollary 2.7. *Let the assumptions of Theorem 2.1 hold and let (u, p) be a weak energy solution of problem (2.1). Then, we have the following dissipative estimate for higher norms:*

$$(2.27) \quad \|\nabla_x u(t)\|_{L^2}^2 + \|\partial_t u(t)\|_{L^2}^2 + \int_t^{t+1} \|\nabla_x \partial_t u(s)\|_{L^2}^2 ds \leq \\ \leq \frac{t^2 + 1}{t^2} (Q(\|(u(0), p(0))\|_E) e^{-\alpha t} + Q(\|g\|_{L^2})),$$

where the positive constant α and monotone function Q are independent of t , u and p .

This estimate, in turn, allows us (analogously to the case of strongly damped wave equations, see [30, 17]) to reduce the study of the asymptotic smoothness for solutions to the following truncated auxiliary problem

$$(2.28) \quad \begin{cases} -\Delta_x u + \nabla_x p + f(u) = g(t), & u|_{\partial\Omega} = 0, \\ \partial_t p + \operatorname{div}(Du) = 0, & p|_{t=0} = p_0, \end{cases}$$

where the external force $g(t) = g - \partial_t u(t)$ satisfies the estimate

$$(2.29) \quad \|g\|_{L^\infty(\mathbb{R}_+, L^2(\Omega))} \leq C$$

which will be studied in the next sections. We also mention here that, in order to restore the u -component of a solution (p, u) of this problem in a unique way by the p -component, we need to assume in addition that

$$(2.30) \quad f'(u) \geq 0.$$

This assumption however, is not restrictive since, in a general case, the extra term Lu can be added to the nonlinearity and also to the external force $g(t)$ and the $L^2(\Omega)$ -norm of this term is under the control.

3. ASYMPTOTIC REGULARITY

In this section, we study the asymptotic smoothing for the truncated system (2.28) which is also of independent interest. We will mainly concentrate here on the case of critical quintic growth rate of the nonlinearity ($f(u) \sim u^5$). The subcritical case is essentially simpler since the standard linear splitting of the solution semigroup on a contracting and compact components works. In contrast to this, we need a *nonlinear* splitting in the critical case. Moreover, due to specific structure of our problem, we

need a combination of different decompositions. We start with the following splitting

$$p = q + r, \quad u = v + w,$$

where

$$(3.1) \quad \begin{cases} \partial_t q + \operatorname{div}(Dv) = 0, & q|_{t=0} = p|_{t=0}, \\ -\Delta_x v + \nabla_x q + f(v) + Lv = 0, & v|_{\partial\Omega} = 0 \end{cases}$$

and

$$(3.2) \quad \begin{cases} \partial_t r + \operatorname{div}(Dr) = 0, & r|_{t=0} = 0, \\ -\Delta_x w + \nabla_x r + [f(u) - f(v)] = Lv + g(t), & w|_{\partial\Omega} = 0. \end{cases}$$

According to the results of previous section, we may assume without loss of generality that $p(0)$ belongs to the absorbing ball in $\bar{L}^2(\Omega)$. Then, from the analogues of dissipative estimates for equation (3.1), we conclude that

$$(3.3) \quad \|p(t)\|_{L^2} + \|q(t)\|_{\bar{L}^2} + \|r(t)\|_{\bar{L}^2} + \|u(t)\|_{H^1} + \|v(t)\|_{H^1} + \|w(t)\|_{H^1} \leq R$$

for all $t \geq 0$. We start with the contracting part (q, v) .

Proposition 3.1. *Let the function f satisfy (1.4), (1.3) and (2.30), $D = D^* > 0$ and estimates (3.3) and (2.29) hold. Then, there exists $L = L(R)$ such that the solution $r(t)$ of the problem (3.2) satisfies the estimate:*

$$(3.4) \quad \|q(t)\|_{\bar{L}^2}^2 + \|v(t)\|_{H^1}^2 \leq Ce^{-\alpha t} \|p(0)\|_{\bar{L}^2}^2,$$

where positive constants C and α are independent of t , u and p .

Proof. We fix $L > 0$ in such a way that

$$f(v) \cdot Dv + Lv \cdot Dv \geq 0, \quad v \in \mathbb{R}^3$$

(it is possible to do so since $f(0) = 0$ and $f(v) \cdot Dv \geq -C$). Then, multiplying the first and second equations of (3.1) by q and Dv respectively and integrating over Ω , we end up with

$$(3.5) \quad \frac{1}{2} \frac{d}{dt} \|q\|_{\bar{L}^2}^2 + \|\nabla_x v\|_{L_D^2}^2 \leq 0.$$

Multiplying now the second equation of (3.1) by $\mathfrak{B}q$ and using the inequality

$$\|f(v)\|_{H^{-1}} \leq C(1 + \|v\|_{H^1}^4) \|v\|_{H^1} \leq C_R \|\nabla_x v\|_{L_D^2},$$

we infer that

$$\|q\|_{\bar{L}^2}^2 \leq C'_R \|\nabla_x v\|_{L_D^2}^2$$

and, therefore,

$$\frac{1}{2} \frac{d}{dt} \|q\|_{\bar{L}^2}^2 + \alpha_R \|q\|_{\bar{L}^2}^2 \leq 0,$$

for some positive α_R depending only on R . Applying the Gronwall inequality, we arrive at the desired estimate for q :

$$\|q(t)\|_{\bar{L}^2}^2 \leq e^{-\alpha_R t} \|p(0)\|_{\bar{L}^2}^2.$$

To get the desired estimate for $\|v\|_{H^1}^2$, it remains to note that multiplication of the second equation of (3.1) by Dv gives

$$\|\nabla_x v(t)\|_{L_D^2}^2 \leq C \|q(t)\|_{\bar{L}^2}^2.$$

Thus, the proposition is proved. $\square$

We now turn to the smooth part $(w(t), r(t))$ of the solution generated by the problem (3.2). At the first step, we derive *exponentially growing* estimate for this part in higher norms which will be improved later.

Proposition 3.2. *Let the assumptions of Proposition 3.1 hold and let $\delta \in (0, \frac{1}{2})$. Then, the following estimate for the solution $(w(t), r(t))$ of (3.2) is valid:*

$$(3.6) \quad \|r(t)\|_{H^\delta}^2 + \|w(t)\|_{H^{1+\delta}}^2 \leq Ce^{Kt},$$

where $K > 0$ and the constant C depends on g (through assumption (2.29)) and R , but is independent of t , p and u .

Proof. To verify this estimate we need the following standard lemma.

Lemma 3.3. *Let $a(x) \geq 0$ be a symmetric measurable matrix and the function $w \in H_0^1(\Omega) \cap L_a^2(\Omega)$ be a solution of the following problem:*

$$(3.7) \quad -\Delta_x w + a(x)w = \nabla_x r + g, \quad w|_{\partial\Omega} = 0,$$

where $L_a^2(\Omega)$ is a weighted Lebesgue space determined by the semi-norm

$$\|w\|_{L_a^2}^2 := \int_{\Omega} a(x)w(x) \cdot w(x) dx < \infty,$$

$r \in \bar{H}^\delta(\Omega) := H^\delta(\Omega) \cap \bar{L}^2(\Omega)$ for some $\delta \in (0, \frac{1}{2})$, and $g \in L^2(\Omega)$. Then, the following estimate holds:

$$(3.8) \quad \|w\|_{L^s} \leq C(\|r\|_{\bar{H}^\delta} + \|g\|_{L^2}),$$

where the constant C is independent of a , g , w and r and $s = \frac{6}{1-2\delta}$ is the Sobolev embedding exponent for $H^{1+\delta} \subset L^s$.

Proof of the lemma. Since g is more regular than $\nabla_x r$, it suffices to verify the estimate for $g = 0$ only. We give below only the formal derivation of (3.8) which can be justified by standard approximation arguments. To this end, we multiply equation (3.7) by $w|w|^n$, where the exponent n will be fixed later and integrate over Ω . This gives

$$(|\nabla_x w|^2, |w|^n) + \|\nabla_x(|w|^{\frac{n+2}{2}})\|_{L^2}^2 \leq C(|r|(|\nabla_x w||w|^{n/2}), |w|^{n/2}).$$

Using the proper Hölder inequality together with Sobolev embeddings, we get

$$(|\nabla_x w|^2, |w|^n) + \|w\|_{L^{3(n+2)}}^{n+2} \leq C\|r\|_{\bar{H}^\delta}(|\nabla_x w|^2, |w|^n)^{1/2} \|w\|_{L^{mn/2}}^{n/2},$$

where $\frac{1}{2} - \frac{\delta}{3} + \frac{1}{2} + \frac{1}{m} = 1$, i.e., $m = \frac{3}{\delta}$. Therefore, we have

$$\|w\|_{L^{3(n+2)}}^{n+2} \leq C\|r\|_{\bar{H}^\delta}^{n+2} + \frac{1}{2}\|w\|_{L^{\frac{3n}{2\delta}}}^{n+2}.$$

Fixing now n in such a way that $3(n+2) = \frac{3n}{2\delta}$, we see that $3(n+2) = s$ and the last estimate finishes the proof of the lemma. $\square$

We now return to the proof of the proposition. First, applying the lemma to the second equation of (3.2) with

$$a(x) := \int_0^1 f'(\kappa u(x) + (1-\kappa)v(x)) d\kappa \geq 0,$$

we end up with

$$(3.9) \quad \|w(t)\|_{L^s} \leq C(\|r(t)\|_{\bar{H}^\delta} + \|g(t)\|_{L^2} + L\|v(t)\|_{L^2}) \leq C(\|r(t)\|_{\bar{H}^\delta} + 1).$$

Second, using the growth restriction on f and Sobolev embedding theorems, it is not difficult to see that

$$(3.10) \quad \|f(u) - f(v)\|_{H^{-1+\delta}} \leq C(1 + \|u\|_{H^1}^4 + \|v\|_{H^1}^4)\|u - v\|_{L^s}.$$

Therefore,

$$(3.11) \quad \|f(u(t)) - f(v(t))\|_{H^{-1+\delta}} \leq C_R\|w(t)\|_{L^s} \leq C_R(\|r(t)\|_{\bar{H}^\delta} + 1).$$

Third, we multiply the second equation of (3.2) by $(-\Delta_x)^\delta w$ and integrate over Ω . This gives

$$(3.12) \quad \|w\|_{H^{1+\delta}}^2 = ((-\Delta_x)^{-1+\delta/2} \nabla_x r, (-\Delta_x)^{1+\delta/2} w) + \\ + (-\Delta_x)^{-1+\delta/2} [f(u) - f(v)], (-\Delta_x)^{1+\delta/2} w) - (Lv(t) + g(t), (-\Delta_x)^\delta w)$$

and therefore

$$(3.13) \quad \|w\|_{H^{1+\delta}} \leq C(\|f(u) - f(v)\|_{H^{-1+\delta}} + \|r\|_{\bar{H}^\delta} + 1) \leq C_R(1 + \|r\|_{\bar{H}^\delta}).$$

Finally, from the first equation of (3.2), we get

$$(3.14) \quad \|r(t)\|_{\bar{H}^\delta} \leq \int_0^t \|\operatorname{div}(Dw(\tau))\|_{H^\delta} d\tau \leq \\ \leq C \int_0^t \|w(\tau)\|_{H^{1+\delta}} d\tau \leq C_R \int_0^t (\|r(\tau)\|_{\bar{H}^\delta} + 1) d\tau$$

and the Gronwall inequality finishes the proof of the proposition. $\square$

At the next step, we split following [41] (see also [26] for some improvements) the solution $u(t)$ of (2.28) on uniformly small $(\bar{u}(t))$ and smooth $(\tilde{u}(t))$ parts.

Proposition 3.4. *Let $\beta > 0$ be arbitrary and $\delta \in (0, \frac{1}{2})$. Let also $(p(t), u(t))$ be a solution of (2.28) satisfying (3.3). Then, there exists $T = T_\delta$ such that the function $u(t)$ can be split in a sum*

$$(3.15) \quad u(t) = \bar{u}(t) + \tilde{u}(t),$$

where for every $t \geq T$

$$(3.16) \quad \|\bar{u}(t)\|_{H^1} \leq \beta, \quad \|\tilde{u}(t)\|_{H^{1+\delta}} \leq C_\beta$$

and the constant C_β depends only on β , δ and R .

Proof. This splitting is an almost immediate corollary of the proved Propositions 3.1 and 3.2. Indeed, let us fix $T = T_\beta$ from the equation

$$Ce^{-\alpha T} R^2 = \beta^2,$$

where all of the constants are the same as in Proposition 3.1. Then, for the v -component of the solution u , we will have the estimate

$$\|v(t)\|_{H^1} \leq \beta, \quad t \geq T.$$

Moreover, if we fix C_β from $Ce^{2KT} = C_\beta^2$ where the constants are the same as in (3.6), we get

$$\|w(t)\|_{H^{1+\delta}} \leq C_\beta$$

if $t \leq 2T$. Thus, functions $v(t)$ and $w(t)$ give the desired splitting of $u(t)$ for $t \in [T, 2T]$.

To construct the desired splitting for all $t \geq T$, we define functions $(q_n(t), v_n(t))$ and $(r_n(t), w_n(t))$ for all $n \in \mathbb{N}$ as solutions of (3.1) and (3.2) respectively, but starting from $t = T(n-1)$ with the initial conditions

$$q_n|_{t=T(n-1)} = 0, \quad r_n|_{t=T(n-1)} = p|_{t=T(n-1)}.$$

Then, arguing analogously, we see that $u(t) = v_n(t) + w_n(t)$ gives the required splitting on the interval $t \in [Tn, T(n+1)]$. Finally, to get the desired splitting for all $t \geq T$, we define $\bar{u}$ and $\tilde{u}(t)$ as hybrid piece-wise continuous functions:

$$\bar{u}(t) = v_n(t), \quad t \in [Tn, T(n+1)), \quad \tilde{u}(t) = w_n(t), \quad t \in [Tn, T(n+1)), \quad n \in \mathbb{Z}.$$

This finishes the proof of the proposition. $\square$

We are now ready to refine Proposition 3.2 and get the dissipative estimate for $(r(t), w(t))$.

Proposition 3.5. *Let the assumptions of Proposition 3.2 hold. Then the solution $(r(t), w(t))$ of problem (3.2) satisfies the estimate*

$$(3.17) \quad \|r(t)\|_{\bar{H}^\delta} + \|w(t)\|_{H^{1+\delta}} \leq C,$$

where the constant C depends on R , but is independent of u , p and t .

Proof. Without loss of generality, we may assume that estimates (3.16) hold for $t \geq 0$. The general case is reduced to this particular one by the proper time shift. The only difference is that we need to put non-zero initial data for $r(t)$. Since the H^δ norm of $r(t)$ on the interval $t \in [0, T]$ can be controlled by (3.6), we just need to assume that

$$(3.18) \quad r|_{t=0} = r_0, \quad \|r_0\|_{H^\delta} \leq C_\beta.$$

This also gives that

$$(3.19) \quad \|v(t)\|_{H^1} \leq \beta, \quad t \geq 0.$$

Moreover, again without loss of generality, we may assume that $f'(0) = 0$. In a general case, the term $f'(0)w(t)$ is lower order and can be treated as a part of $g(t)$.

The idea of the proof is to refine estimate (3.11) using the result of Proposition 3.4. First, we refine (3.10) using the fact that $f'(0) = 0$, namely, this assumption gives us that

$$(3.20) \quad \|f(u) - f(v)\|_{H^{-1+\delta}} \leq C(\|u\|_{H^1} + \|v\|_{H^1})(1 + \|u\|_{H^1}^3 + \|v\|_{H^1}^3)\|u - v\|_{L^s}$$

for some constant C depending only on f . Second, we write

$$f(u) - f(v) = [f(\bar{u} + \tilde{u}) - f(\bar{u})] + [f(\bar{u}) - f(v)]$$

and apply (3.20) to both terms on the right-hand side. Indeed, since $H^{1+\delta} \subset L^s$ and the function $\tilde{u}$ is bounded in $H^{1+\delta}$, we have

$$\|f(\bar{u} + \tilde{u}) - f(\bar{u})\|_{H^{-1+\delta}} \leq C(1 + \|\bar{u}\|_{H^1}^4 + \|\tilde{u}\|_{H^1}^4)\|\tilde{u}\|_{H^{1+\delta}} \leq C_1$$

for some $C_1 > 0$ which depends on β and R . Applying estimate (3.20) to the second term and using inequalities (3.16) and (3.19), we get

$$\|f(\bar{u}) - f(v)\|_{H^{-1+\delta}} \leq C\beta\|\bar{u} - v\|_{L^s}$$

and using that

$$\|\tilde{u} - v\|_{L^s} = \|\tilde{u} - w\|_{L^s} \leq \|w\|_{L^s} + C\|\tilde{u}\|_{H^{1+\delta}} \leq \|w\|_{L^s} + C$$

we get

$$(3.21) \quad \|f(u) - f(v)\|_{H^{-1+\delta}} \leq C\beta\|w\|_{L^s} + C_\beta,$$

where the constant C is independent of $\beta > 0$. Together with the result of Lemma 3.3, we finally arrive at the refined estimate

$$(3.22) \quad \|f(u(t)) - f(v(t))\|_{H^{-1+\delta}} \leq C\beta\|r(t)\|_{H^\delta} + C_\beta.$$

Crucial for us is that the constant C is independent of β , so the coefficient in front of $\|r(t)\|_{H^\delta}$ can be made arbitrary small by the choice of δ .

We are now ready to complete the proof of the proposition. To this end, we treat equation (3.2) as a linear (A.1) interpreting the term $f(u(t)) - f(v(t))$ as a part of external force $g(t)$ and use estimate (A.7) with $K_\delta = -\alpha < 0$, see Corollary A.5. This gives

$$(3.23) \quad \|r(t)\|_{\bar{H}^\delta} \leq C\|r(0)\|_{\bar{H}^\delta} e^{-\alpha t} + C_\beta + C\beta \int_0^t e^{-\alpha(t-\tau)} \|r(\tau)\|_{\bar{H}^\delta} d\tau.$$

Fixing now $\beta > 0$ in such a way that $C\beta = \frac{\alpha}{2}$ and applying the Gronwall inequality, we end up with the desired estimate

$$\|r(t)\|_{\bar{H}^\delta} \leq C\|r(0)\|_{\bar{H}^\delta} e^{-\alpha t/2} + C_1.$$

Combining this estimate with (3.13), we end up with (3.17) and finish the proof of the proposition. $\square$

We now summarize our results concerning the truncated system (2.28) under the assumptions (2.30) and (2.29) for the nonlinearity f and the external force $g(t)$. We first mention that the global well-posedness and dissipativity of this problem in the space $\bar{L}^2(\Omega)$ can be obtained exactly as in Theorems 2.1 and 2.4, so we have the estimate

$$(3.24) \quad \|p(t)\|_{\bar{L}^2}^2 + \|u(t)\|_{H^1}^2 \leq Q(\|p(0)\|_{\bar{L}^2}) e^{-\alpha t} + Q(\|g\|_{L^\infty}),$$

where positive constant α and monotone function Q are independent of p and t .

Thus, problem (2.28) can be considered independently of problem (2.1) on the whole phase space $\bar{L}^2(\Omega)$ and estimate (3.24) gives us the existence of an absorbing ball in $\bar{L}^2(\Omega)$, so the key assumptions (3.3) will be automatically satisfied if we take the initial data from this absorbing ball.

Let us denote by $\mathcal{U}(t) : \bar{L}^2(\Omega) \rightarrow \bar{L}^2(\Omega)$ the solution operator for problem (2.28):

$$(3.25) \quad \mathcal{U}(t)p(0) := p(t),$$

where $p(t)$ is a solution of (2.28). Then, taking into the account that the $u(t)$ -component of the solution can be restored in a unique way (due to Lemma 3.3) if the $p(t)$ -component is known, we can reformulate the results of Propositions 3.5 and 3.1 as follows.

Corollary 3.6. *Let the nonlinearity f satisfy (2.30), (1.4) and (1.3) and the function g satisfy (2.29). Then, for a sufficiently large R , the R -ball $\mathcal{B}_R^\delta$ of radius R in $\bar{H}^\delta(\Omega)$ is an exponentially attracting for the solution operator*

$\mathcal{U}(t)$, i.e., there exists positive constant $\alpha > 0$ and a monotone function Q such that, for every bounded set $B \subset \bar{L}^2(\Omega)$,

$$(3.26) \quad \text{dist}_{\bar{L}^2}(\mathcal{U}(t)B, \mathcal{B}_R^\delta) \leq Q(\|B\|_{\bar{L}^2})e^{-\alpha t},$$

where $\text{dist}_H(A, B)$ stands for the non-symmetric Hausdorff distance between the sets A and B in a Banach space H .

We also have the analogue of the dissipative estimate (3.24) in the space H^δ for any exponent $\delta \in [0, \frac{1}{2})$.

Corollary 3.7. *Let the assumptions of Corollary 3.6 hold and let $p(0) \in \bar{H}^\delta(\Omega)$ for some $\delta \in [0, \frac{1}{2})$. Then the following dissipative estimate holds for the solution of problem (2.28):*

$$(3.27) \quad \|p(t)\|_{\bar{H}^\delta}^2 + \|u(t)\|_{H^{1+\delta}}^2 \leq Q(\|p(0)\|_{\bar{H}^\delta})e^{-\alpha t} + Q(\|g\|_{L^\infty(\mathbb{R}_+, L^2)})$$

for some positive α and monotone function Q which are independent of p and t .

Indeed, this estimate can be proved analogously to the proof of Proposition 3.5, but even simpler since we may take $q(t) = v(t) = 0$, so we leave the details to the reader.

Thus, we have verified that the solution operator $\mathcal{U}(t)$ is well-defined and dissipative in $\bar{H}^\delta(\Omega)$ for any $0 \leq \delta < \frac{1}{2}$. It also worth to note that all of the estimates obtained so far uses only that

$$(3.28) \quad \|g\|_{L^\infty(\mathbb{R}_+, H^{-1+\delta})} \leq C, \quad \delta \in (0, \frac{1}{2}).$$

The natural next step is to extend this result to $\delta = 1$ using bootstrapping arguments. The situation here is much simpler than for the first step since the nonlinearity f is subcritical in the phase space $\bar{H}^\delta(\Omega)$, so the linear splitting may be used. Moreover, due to the embedding theorem $H^{1+\frac{1}{5}} \subset L^{10}$ and the growth restrictions on f , we have

$$(3.29) \quad \|f(u)\|_{L^2} \leq C(1 + \|u\|_{H^{1+\delta}}^5), \quad \delta \geq \frac{1}{5}$$

and, therefore, only one more step of iterations is necessary to reach $\delta = 1$. Namely, we split the solution (p, u) as follows:

$$p(t) = p_1(t) + p_2(t), \quad u(t) = u_1(t) + u_2(t),$$

where the decaying component $(p_1(t), u_1(t))$ solves

$$(3.30) \quad \partial_t p_1 + \text{div}(Du_1) = 0, \quad -\Delta_x u_1 + \nabla_x p_1 = 0, \quad p_1|_{t=0} = p|_{t=0}$$

and the smooth component $(p_2(t), u_2(t))$ is a solution of

$$(3.31) \quad \partial_t p_2 + \text{div}(Du_2) = 0, \quad -\Delta_x u_2 + \nabla_x p_2 = g(t) - f(u(t)), \quad p_2|_{t=0} = 0.$$

Then, the following proposition holds.

Proposition 3.8. *Let $\delta \in [\frac{1}{5}, \frac{1}{2})$ and let the initial data $p(0)$ belongs to the absorbing ball $\mathcal{B}_R^\delta$. Then the following estimates hold for the solutions of (3.30) and (3.31):*

$$(3.32) \quad \|p_1(t)\|_{\bar{H}^\delta} + \|u_1(t)\|_{H^{1+\delta}} \leq C\|p(0)\|_{\bar{H}^\delta}e^{-\alpha t}$$

and

$$(3.33) \quad \|p_2(t)\|_{\bar{H}^1} + \|u_2(t)\|_{H^2} \leq C\|p(0)\|_{\bar{H}^\delta} e^{-\alpha t} + C_R(1 + \|g\|_{L^\infty(L^2)}),$$

where $\alpha > 0$ and C, C_R are independent of u, p and t .

Indeed, these estimates follow immediately from estimate (A.7) with $K_\delta = -\alpha < 0$ for the linear equation, dissipative estimate (3.27) and estimate (3.29).

Analogously to Corollary 3.7, this result gives the dissipativity in the phase space $\bar{H}^1$.

Corollary 3.9. *Let the assumptions of Corollary 3.6 hold and let $p(0) \in \bar{H}^1(\Omega)$. Then the following dissipative estimate holds for the solution of problem (2.28):*

$$(3.34) \quad \|p(t)\|_{\bar{H}^1}^2 + \|u(t)\|_{H^2}^2 \leq Q(\|p(0)\|_{\bar{H}^1})e^{-\alpha t} + Q(\|g\|_{L^\infty(\mathbb{R}_+, L^2)})$$

for some positive α and monotone function Q which are independent of p and t .

Indeed, to get this estimate, it is enough to estimate the L^2 -norm of $f(u)$ using Corollary 3.7 and get the desired estimate for the H^1 -norm from the linear equation (A.1) treating $f(u(t))$ as a part of the external forces.

Analogously to Corollary 3.6 the result of Proposition 3.8 can be rewritten in the following form.

Corollary 3.10. *Let the assumptions of Corollary 3.6 hold. Then, for a sufficiently large R , the R -ball $\mathcal{B}_R^1$ of radius R in $\bar{H}^1(\Omega)$ is an exponentially attracting for the solution operator $\mathcal{U}(t)$ in $\bar{H}^\delta$, i.e., there exists positive constant $\alpha > 0$ and a monotone function Q such that, for every bounded set $B \subset \bar{H}^\delta(\Omega)$,*

$$(3.35) \quad \text{dist}_{\bar{H}^\delta}(\mathcal{U}(t)B, \mathcal{B}_R^1) \leq Q(\|B\|_{\bar{H}^\delta})e^{-\alpha t}.$$

Moreover, using the Lipschitz continuity of $\mathcal{U}(t)$ in $\bar{L}^2(\Omega)$, exponential attractions (3.26) and (3.35) together with the transitivity of exponential attraction (see [9]), we arrive at the following result.

Corollary 3.11. *Let the assumptions of Corollary 3.6 hold. Then, for a sufficiently large R , the R -ball $\mathcal{B}_R^1$ of radius R in $\bar{H}^1(\Omega)$ is an exponentially attracting for the solution operator $\mathcal{U}(t)$ in $\bar{L}^2(\Omega)$, i.e., there exists positive constant $\alpha > 0$ and a monotone function Q such that, for every bounded set $B \subset \bar{L}^2(\Omega)$,*

$$(3.36) \quad \text{dist}_{\bar{L}^2}(\mathcal{U}(t)B, \mathcal{B}_R^1) \leq Q(\|B\|_{\bar{L}^2})e^{-\alpha t}.$$

We conclude this section by translating the obtained results for the truncated system (2.28) to the initial problem (2.1). The next result can be considered as the main result of this section.

Theorem 3.12. *Let the assumptions of Theorem 2.1 hold. Then the R -ball $\mathbb{B}_R^1$ in the higher energy space*

$$E^1 := [H^2(\Omega) \cap H_0^1(\Omega)] \times \bar{H}^1(\Omega)$$

is an exponentially attracting set for the solution semigroup $S(t) : E \rightarrow E$ generated by the problem (2.1) if R is large enough, i.e., there exists $\alpha > 0$ and monotone Q such that, for every bounded set $B \subset E$,

$$(3.37) \quad \text{dist}_E(S(t)B, \mathbb{B}_R^1) \leq Q(\|B\|_E)e^{-\alpha t}.$$

Moreover, the problem (2.1) is well-posed and dissipative in the space E^1 as well, i.e., if $(u(0), p(0)) \in E^1$ then the following estimate holds:

$$(3.38) \quad \|(u(t), p(t))\|_{E^1} \leq Q(\|(u(0), p(0))\|_{E^1})e^{-\alpha t} + Q(\|g\|_{L^2})$$

for some positive α and monotone Q .

Proof. Indeed, the exponential attraction (3.37) follows immediately from Corollary 3.11 and smoothing property of Corollary 2.7.

To get the dissipative estimate (3.38), we note that if the initial data $(u(0), p(0)) \in E^1$, we have from equations (2.1) that

$$\|u(0)\|_C + \|\partial_t u(0)\|_{L^2} + \|\partial_t p(0)\|_{\bar{L}^2} \leq Q(\|(u(0), p(0))\|_{E^1}),$$

so, we need not to use multiplication by t and t^2 in the estimates given in the proof of Theorem 2.6 in order to remove the initial data and this gives us better analogue of estimate (2.27):

$$(3.39) \quad \|\nabla_x u(t)\|_{L^2} + \|\partial_t u(t)\|_{L^2} \leq Q(\|(u(0), p(0))\|_{E^1})e^{-\alpha t} + Q(\|g\|_{L^2}).$$

This, in turn, allows to use the truncated system (2.28) starting from $t = 0$. Then the desired dissipative estimate follows from the analogous estimate (3.34) for the truncated system. Thus, the theorem is proved. $\square$

4. ATTRACTORS

In this section, we use the results obtained above for constructing global and exponential attractors for problem (2.1). We start with a global attractor.

Definition 4.1. Let $S(t) : E \rightarrow E$, $t \geq 0$ be a semigroup. Then, a set $\mathcal{A} \subset E$ is a global attractor for $S(t)$ in E if

1. $\mathcal{A}$ is compact in E ;
2. $\mathcal{A}$ is strictly invariant, i.e., $S(t)\mathcal{A} = \mathcal{A}$ for all $t \geq 0$.
3. $\mathcal{A}$ is an attracting set for $S(t)$ in E . The latter means that for every bounded set B in E and every neighbourhood $\mathcal{O}(\mathcal{A})$ of the set $\mathcal{A}$ there exist $T = T(B, \mathcal{O})$ such that

$$(4.1) \quad S(t)B \subset \mathcal{A}, \quad \forall t \geq T.$$

If $S(t)$ is a solution semigroup related with an evolutionary equation, then the attractor $\mathcal{A}$ of $S(t)$ is often called an attractor of this evolutionary equation, see [2, 4, 20, 27, 35] for more details.

Theorem 4.2. Let the assumptions of Theorem 2.1 hold. Then equation (2.1) possesses an attractor $\mathcal{A}$ in E which is a bounded set of E^1 . Moreover, this attractor possesses the following description:

$$(4.2) \quad \mathcal{A} = \mathcal{K}|_{t=0},$$

where $\mathcal{K} \subset L^\infty(\mathbb{R}, E)$ is a set of all complete (=defined for all $t \in \mathbb{R}$) bounded in E solutions of equation (2.1).

Proof. According to the abstract attractor's existence theorem, see e.g., [2], we need to verify two properties:

1. The operators $S(t)$ are continuous for every fixed t as operators from E to E ;
2. The semigroup $S(t)$ possesses a compact attracting set in E .

The first property is verified in Theorem 2.4 and the second one follows from Theorem 3.12. Since the attractor is always a subset of a compact attracting set, we get the boundedness of $\mathcal{A}$ in E^1 and the representation formula (4.2) also follows from the abstract attractor's existence theorem. Thus, the theorem is proved. $\square$

We now turn to exponential attractors. These objects have been introduced in [6] in order to overcome the major drawback of the theory of global attractors, namely, the fact that the rate of attraction to a global attractor may be arbitrarily slow and that there is no way in general to control this rate of attraction in terms of physical parameters of the considered equation. This makes the global attractor sensitive to perturbations and it becomes in a sense unobservable in finite-time simulations, see [6, 7, 8, 27] for more details. We start with the formal definition.

Definition 4.3. A set $\mathcal{M} \subset E$ is an exponential attractor for the semigroup $S(t) : E \rightarrow E$, $t \geq 0$, if

1. $\mathcal{M}$ is a compact set in E ;
2. $\mathcal{M}$ is semi-invariant $S(t)\mathcal{M} \subset \mathcal{M}$ for $t \geq 0$;
3. $\mathcal{M}$ has a finite box-counting dimension in E :

$$\dim_F(\mathcal{A}, E) \leq C < \infty;$$

4. There exist positive constant α and monotone function Q such that, for every bounded set $B \subset E$, we have

$$(4.3) \quad \text{dist}_E(S(t)B, \mathcal{M}) \leq Q(\|B\|_E)e^{-\alpha t}$$

for all $t \geq 0$.

The next theorem can be considered as the main result of this section.

Theorem 4.4. *Let the assumptions of Theorem 2.1 hold. Then equation (2.1) possesses an exponential attractor $\mathcal{M}$ in E which is a bounded set in the space E^1 .*

Proof. Following the general strategy, see [7, 8, 9, 27], we first construct a discrete exponential attractor $\mathcal{M}_d \subset E^1$ for the semigroup $S_n = S_1^n$ generated by the map $S(T)$ restricted to the R -ball $\mathcal{B}_R^1$ in E^1 . Here we fix $T > 0$ in such a way that

$$S(T) : \mathbb{B}_R^1 \rightarrow \mathbb{B}_R^1.$$

It is possible to do due to estimate (3.38). If the discrete attractor $\mathcal{M}_d$ is constructed its continuous analogue $\mathcal{M} \subset E^1$ is given by the standard formula

$$(4.4) \quad \mathcal{M} := \cup_{t \in [0, T]} S(t)\mathcal{M}_d.$$

This, together with (3.38) gives us the attraction property in E for all bounded sets of E^1 . Combining this with the exponential attraction (3.37)

and transitivity of exponential attraction (see [9]), we get the desired exponential attraction of any bounded set in E . The semi-invariance follows immediately from semi-invariance of a discrete attractor and the explicit formula (4.4). The compactness and finite-dimensionality also follow from (4.4) if we know, in addition, that $(t, \xi) \rightarrow S(t)\xi$ is Lipschitz (or Hölder) continuous as a map from $[0, T] \times \mathcal{M}_d \rightarrow E$. The Lipschitz continuity with respect to the initial data is verified in Theorem 2.4 and the Lipschitz continuity in times follows from the fact that $\|\partial_t u(t)\|_{L^2}$ and $\|\partial_t p(t)\|_{\bar{L}^2}$ are uniformly bounded on $\mathbb{B}_R^1$ (due to estimate (3.39)). Thus, we only need to verify the existence of a discrete exponential attractor $\mathcal{M}_d$ on a set $\mathbb{B}_R^1$. To this end, we need the following standard result on the existence of exponential attractors, see [7, 8, 27].

Lemma 4.5. *Let E and V be two B -spaces such that V is compactly embedded in E and let $\mathbb{B} \subset E$ be a bounded set in E . Assume also that we are given a map $S : \mathbb{B} \rightarrow \mathbb{B}$ such that, for every two points $\xi_1, \xi_2 \in \mathbb{B}$, we have a splitting*

$$(4.5) \quad S(\xi_1) - S(\xi_2) = \hat{\xi} + \tilde{\xi},$$

where

$$(4.6) \quad \|\hat{\xi}\|_E \leq \kappa \|\xi_1 - \xi_2\|_E$$

for some $\kappa < \frac{1}{2}$ and

$$(4.7) \quad \|\tilde{\xi}\|_V \leq K \|\xi_1 - \xi_2\|_E,$$

where κ and K are independent of ξ_1 and ξ_2 . Then the discrete semi-group generated by iterations of the map S possesses an exponential attractor $\mathcal{M}_s \subset B$ on $B \subset E$.

To apply this lemma, we need to split the solution $(\bar{u}(t), \bar{p}(t))$ of system (2.13) for differences of two solutions of system (2.1) on a sum of contracting $(\hat{u}(t), \hat{u}(t))$ and smoothing $(\tilde{u}(t), \tilde{p}(t))$ components. The first part will solve the homogeneous linear system:

$$(4.8) \quad \begin{cases} \partial_t \hat{u} - \Delta_x \hat{u} + \nabla_x \hat{p} = 0, & \hat{u}|_{t=0} = \bar{u}|_{t=0}, \\ \partial_t \hat{p} + \operatorname{div}(D\hat{u}) = 0, & \hat{p}|_{t=0} = \bar{p}|_{t=0} \end{cases}$$

and the smoothing component is taken as a solution of

$$(4.9) \quad \partial_t \tilde{u} - \Delta_x \tilde{u} + \nabla_x \tilde{p} = -l(t)\bar{u}, \quad \partial_t \tilde{p} + \operatorname{div}(D\tilde{u}) = 0, \quad \tilde{u}|_{t=0} = \tilde{p}|_{t=0} = 0,$$

where $l(t) := \int_0^1 f'(\tau u_1 + (1 - \tau)u_2) d\tau$. We recall that, according to Theorem 2.4,

$$(4.10) \quad \|\bar{u}(t)\|_{L^2} + \|\bar{p}(t)\|_{\bar{L}^2} \leq Ce^{Kt} (\|\bar{u}(0)\|_{L^2} + \|\bar{p}(0)\|_{\bar{L}^2}).$$

Moreover, since $(u_i(0), p_i(0)) \in \mathbb{B}_R^1$, $i = 1, 2$, due to (3.39) the C -norm of $u_i(t)$ is uniformly bounded and, therefore,

$$(4.11) \quad \|l(t)\bar{u}(t)\|_{L^2} \leq Ce^{Kt} (\|\bar{u}(0)\|_{L^2} + \|\bar{p}(0)\|_{\bar{L}^2}),$$

so the term $l(t)\bar{u}(t)$ can be treated as an external force. Estimates (4.6) and (4.7) are verified in the next two lemmas.

Lemma 4.6. *Let the above assumptions hold. Then, the solution $(\hat{u}(t), \hat{p}(t))$ of problem (4.8) satisfies the estimate:*

$$(4.12) \quad \|\hat{u}(t)\|_{L^2}^2 + \|\hat{p}(t)\|_{L^2}^2 \leq Ce^{-\alpha t} (\|\bar{u}(0)\|_{L^2}^2 + \|\bar{p}(0)\|_{L^2}^2),$$

where the positive constants C and α are independent of u_i and p_i .

Proof of the lemma. Indeed, multiplying the first equation of (4.8) by $D\hat{u}$ integrating with respect to x and using the second equation, we arrive at

$$\frac{1}{2} \frac{d}{dt} (\|\hat{u}(t)\|_{L_D^2}^2 + \|\hat{p}(t)\|_{L^2}^2) + \|\nabla_x \hat{u}(t)\|_{L_D^2}^2 = 0.$$

Moreover, multiplying the first equation on $-\mathfrak{B}\hat{p}(t)$ and using again the second equation we get

$$-\frac{d}{dt} (\hat{u}(t), \mathfrak{B}\hat{p}(t)) + \|\hat{p}(t)\|_{L^2}^2 - (\hat{u}(t), \mathfrak{B} \operatorname{div}(D\hat{u}(t))) = 0.$$

Multiplying this equation by small positive ε and taking a sum with the previous equation, we finally get

$$\frac{1}{2} \frac{d}{dt} (\|\hat{u}(t)\|_{L_D^2}^2 - 2\varepsilon (\hat{u}(t), \mathfrak{B}\hat{p}(t)) + \|\hat{p}(t)\|_{L^2}^2) + \alpha \|\hat{u}(t)\|_{L_D^2}^2 + \varepsilon \|\hat{p}(t)\|_{L^2}^2 \leq 0$$

for some positive α . The Gronwall inequality applied to this relation gives the desired result if $\varepsilon > 0$ is small enough. Thus, the lemma is proved. $\square$

Lemma 4.7. *Let the above assumptions hold. Then, the solution $(\tilde{u}(t), \tilde{p}(t))$ satisfies the following estimate:*

$$(4.13) \quad \|\tilde{u}(t)\|_{H^1}^2 + \|\tilde{p}(t)\|_{\bar{H}^1}^2 \leq Ce^{Kt} (\|\bar{u}(0)\|_{L^2}^2 + \|\bar{p}(0)\|_{L^2}^2),$$

where the constants C and K depend on R , but are independent of u_i and p_i .

Proof of the lemma. Indeed, multiplying the second equation of (4.9) by $\Delta_x \tilde{u}$ and using (4.11), we get

$$\frac{1}{2} \frac{d}{dt} \|\tilde{u}(t)\|_{H^1}^2 + \|\Delta_x \tilde{u}(t)\|_{H^2}^2 \leq C \|\tilde{p}(t)\|_{H^1}^2 + Ce^{Kt} (\|\bar{u}(0)\|_{L^2}^2 + \|\bar{p}(0)\|_{L^2}^2).$$

Taking now ∇_x from the both sides of the second equation of (4.9) and multiplying it by $\nabla_x \tilde{p}(t)$, we arrive at

$$\frac{1}{2} \frac{d}{dt} \|\nabla_x \tilde{p}(t)\|_{L^2}^2 \leq \|\Delta_x \tilde{u}(t)\|_{L^2}^2 + C \|\nabla_x \tilde{p}(t)\|_{L^2}^2.$$

Taking a sum of the obtained inequalities, we finally infer that

$$(4.14) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} (\|\nabla_x \tilde{u}(t)\|_{L^2}^2 + \|\nabla_x \tilde{p}(t)\|_{L^2}^2) \\ \leq C \|\nabla_x \tilde{p}(t)\|_{L^2}^2 + Ce^{Kt} (\|\bar{u}(0)\|_{L^2}^2 + \|\bar{p}(0)\|_{L^2}^2) \end{aligned}$$

and the Gronwall inequality applied to this relation finishes the proof of the lemma. $\square$

We are now ready to complete the proof of the theorem. Indeed, estimates (4.12) and (4.13) guarantee that the assumptions of Lemma 4.5 are satisfied if we take

$$V := H_0^1(\Omega) \times \bar{H}^1(\Omega)$$

and fix T big enough that $Ce^{-\alpha T} < \frac{1}{2}$. Thus, the discrete exponential attractor $\mathcal{M}_d$ is constructed and the desired continuous exponential attractor

$\mathcal{M}$ can be constructed via (4.4) as explained above. Therefore, the theorem is proved. $\square$

5. GENERALIZATIONS AND CONCLUDING REMARKS

In this section, we briefly discuss the so-called Navier-Stokes-Brinkman-Forchheimer equation in the following form:

$$(5.1) \quad \begin{cases} \partial_t u + B(u, u) - \Delta_x u + \nabla_x p + f(u) = g, & u|_{\partial\Omega} = 0, \quad u|_{t=0} = u_0, \\ \partial_t p + \operatorname{div}(u) = 0, & p|_{t=0} = p_0, \end{cases}$$

where

$$(5.2) \quad B(u, v) = (u, \nabla_x) v + \frac{1}{2} \operatorname{div}(u) v.$$

The extra term $\frac{1}{2} \operatorname{div}(u) u$ is added to the standard Navier-Stokes inertial term in order to preserve the energy identity, see [10, 12, 36] and references therein. Indeed, in this case we have

$$(B(u, v), v) \equiv 0, \quad \forall u, v \in H_0^1(\Omega)$$

and we have the energy identity (2.3) with $D = 1$ exactly as in the case $B = 0$ considered above, namely,

$$(5.3) \quad \frac{1}{2} \frac{d}{dt} (\|u\|_{L^2}^2 + \|p\|_{L^2}^2) + \|\nabla_x u\|_{L^2}^2 + (f(u), u) = (g, u).$$

The theory of this equation is very similar to the case $B = 0$ considered above with the only difference that, in order to control the extra non-linearity B , we need to assume that $f(u)$ has a super-cubic growth rate, see [14, 18], but this assumption is already incorporated to (1.4) if $l > 1$.

We start with the analogue of dissipative estimate (2.2). The analogue of (2.3) is already obtained, so in order to get the key differential inequality (2.8), we only need to estimate the extra term

$$(5.4) \quad |\varepsilon(B(u, u), \mathfrak{B}p)| \leq C\varepsilon \|u\|_{L^3} \|\nabla_x u\|_{L^2} \|p\|_{\bar{L}^2} \leq C \|u\|_{L^3}^3 + \frac{1}{2} \|\nabla_x u\|_{L^2}^2 + C\varepsilon \|p\|_{\bar{L}^2}^6 \leq \frac{1}{2} (f(u), u) + C + \frac{1}{2} \|\nabla_x u\|_{L^2}^2 + C\varepsilon^6 \mathcal{E}_\varepsilon(u, p)^3,$$

where the constant C is independent of ε . Thus, analogously to Theorem 2.1, we have the following result.

Proposition 5.1. *Let the assumptions of Theorem 2.1 hold, $l > 1$ and let (u, p) be a weak energy solution of problem (5.1). Then, this solution satisfies the dissipative estimate (2.2).*

Let us now turn to uniqueness. This can be proved exactly as in the incompressible case (see [18]). Indeed, in comparison with Theorem 2.4, we need to estimate the extra term

$$(B(u_1, u_1) - B(u_2, u_2), u_1 - u_2) = (B(\bar{u}, u_2), \bar{u}), \quad \bar{u} = u_1 - u_2,$$

where u_1 and u_2 are two solutions of (5.1). Integrating by parts and using the Cauchy-Schwarz inequality, we get

$$|B(\bar{u}, u_2), \bar{u}| \leq \frac{1}{4} \|\nabla_x \bar{u}\|_{L^2}^2 + C \|u_2 \bar{u}\|_{L^2}^2.$$

On the other hand, using assumptions (1.4), analogously [18], we get

$$(f(u_1) - f(u_2), \bar{u}) \geq \kappa(|u_1|^{1+l} + |u_2|^{1+l}, |\bar{u}|^2) - L\|\bar{u}\|_{L^2}^2 \geq C\|u_2\bar{u}\|^2 - \tilde{L}\|\bar{u}\|_{L^2}^2$$

and, therefore,

$$(B(u_1) - B(u_2), \bar{u}) + (f(u_1) - f(u_2), \bar{u}) \geq -\tilde{L}\|\bar{u}\|_{L^2}^2.$$

Arguing further as in the proof of Theorem 2.4, we get estimate (2.12) and verify the uniqueness of the solution for problem (5.1). Thus, as in the case of $B = 0$, equation (5.1), generates a dissipative semigroup $S(t)$ in the phase space E .

We now discuss the smoothing property and start with the instantaneous smoothing (analog of Theorem 2.6).

Proposition 5.2. *Let the assumptions of Theorem 2.1 hold, $l > 1$ and let (u, p) be a weak energy solution of equations (5.1). Then the following partial smoothing property holds:*

$$(5.5) \quad t^{8/3}\|\nabla_x u(t)\|_{L^2}^2 + t^{8/3}\|\partial_t u(t)\|_{L^2}^2 + t^{8/3}\|\partial_t p(t)\|_{L^2}^2 + \\ + \int_0^t s^{8/3}\|\nabla_x \partial_t u(s)\|_{L^2}^2 ds \leq Q(\|(u(0), p(0))\|_E^2) + Q(\|g\|_{L^2}),$$

where $t \in [0, 1]$ and a function Q is independent of t and u .

Proof. Here, we have a little difference (in comparison with the proof of Theorem 2.6), namely, multiplication of the equation on $\partial_t u$ does not work since we have not enough regularity to control the term $(B(u, u), \partial_t u)$. By this reason, again similarly to the incompressible case (see [18]), we need to differentiate the first equation of (5.1) with respect to t and multiply it by $v = \partial_t u$ at the first step. The nonlinearity $B(u, v) + B(v, u)$ is controlled here by the second nonlinearity $f'(u)v$ exactly as in the proof of uniqueness, so we get the following analogue of

$$(5.6) \quad \frac{1}{2} \frac{d}{dt} (\|v(t)\|_{L^2}^2 + \|\partial_t p\|_{L^2}^2) + \|\nabla_x v\|_{L^2}^2 \leq \tilde{L}\|v\|_{L^2}^2.$$

Moreover, from the first equation of (5.1) and the dissipative estimate (2.2), we infer after the standard estimates that

$$(5.7) \quad \|v\|_{L^{6/5}(0,1;H^{-1})} \leq Q(\|(u_0, p_0)\|_E) + Q(\|g\|_{L^2}).$$

Estimate (5.7) replaces the missed control of the quantity $\int_0^t s\|v(s)\|_{L^2}^2 ds$ and allows us to get the desired smoothing property. Indeed, multiplying (5.6) by $t^{8/3}$, integrating in time and using the estimate

$$(5.8) \quad \int_0^t s^{5/3}\|v(s)\|_{L^2}^2 ds \leq \\ \leq \int_0^s (s^{4/3}\|v(s)\|_{L^2})^{1/2} (s^{4/3}\|v(s)\|_{H^1})^{3/4} \|v(s)\|_{H^{-1}}^{3/4} ds \leq \\ \leq \varepsilon \sup_{s \in [0,t]} \left\{ s^{8/3}\|v(s)\|_{L^2}^2 \right\} + \varepsilon \int_0^t s^{8/3}\|v(s)\|_{H^1}^2 ds + C_\varepsilon \|v\|_{L^{6/5}(0,1;H^{-1})}^2,$$

where $\varepsilon > 0$ can be taken arbitrarily small, we end up with the desired smoothing property for the derivatives

$$t^{4/3}\|\partial_t u(t)\|_{L^2} + t^{4/3}\|\partial_t p(t)\|_{L^2} \leq Q(\|u_0, p_0\|_E) + Q(\|g\|_{L^2})$$

for $t \in [0, 1]$ and some monotone function Q . Returning back to the first equation of (5.1), multiplying it by $u(t)$ and integrating in x , we get

$$\|\nabla_x u(t)\|_{L^2}^2 + (|f(u(t)).u(t)|, 1) \leq C(\|p(t)\|_{L^2}^2 + \|g\|_{L^2}^2 + \|\partial_t u(t)\|_{L^2}^2)$$

which together with the previous estimate and dissipative estimate (2.2) give the desired smoothing property and finishes the proof of the proposition. $\square$

As in the case $B = 0$, this instantaneous smoothing property allows us to reduce the study of the asymptotic smoothing to the truncated problem

$$(5.9) \quad \begin{cases} -\Delta_x u + \nabla_x p + B(u, u) + f(u) = g(t), & u|_{\partial\Omega} = 0 \\ \partial_t p + \operatorname{div}(Du) = 0, & p|_{t=0} = p_0, \end{cases}$$

where $g(t) := g - \partial_t u(t)$ satisfies (2.29). Moreover, using the obvious estimate

$$(5.10) \quad \|B(u(t), u(t))\|_{H^{-1/2}} \leq \|B(u(t), u(t))\|_{L^{3/2}} \leq C\|\nabla_x u(t)\|_{L^2}^2,$$

we can assume without loss of generality (due to the dissipative estimate (2.2) and smoothing property (5.5)) that the nonlinearity $B(u, u)$ is bounded in $L^\infty(\mathbb{R}_+, H^{-1/2})$. Thus, we may treat the nonlinearity $B(u, u)$ as a part of g as well. Then the new function g will satisfy (3.28) and we may treat equations (5.9) exactly as equations (2.28).

This gives us the analogues of Corollaries 3.6 and 3.7 for the truncated system (5.9). In order to make the second step of bootstrapping, we note that

$$\|B(u(t), u(t))\|_{L^2} \leq C\|u(t)\|_{H^{1+\delta}}$$

for $\delta \geq \frac{1}{4}$. Therefore, if the $H^{1+\delta}$ -regularity of $u(t)$ is verified for $\delta \geq \frac{1}{4}$, the next step of bootstrapping will give us the H^2 -regularity exactly as in the Section 3. Thus, we have proved the following analogue of Theorem 3.12.

Theorem 5.3. *Let the assumptions of Theorem 2.1 hold and $l > 1$. Then the R -ball $\mathbb{B}_R^1$ in the higher energy space E^1 is an exponentially attracting set for the solution semigroup $S(t) : E \rightarrow E$ generated by the problem (5.1) if R is large enough, i.e., there exists $\alpha > 0$ and monotone Q such that, for every bounded set $B \subset E$,*

$$(5.11) \quad \operatorname{dist}_E(S(t)B, \mathbb{B}_R^1) \leq Q(\|B\|_E)e^{-\alpha t}.$$

Moreover, the problem (5.1) is well-posed and dissipative in the space E^1 as well, i.e., if $(u(0), p(0)) \in E^1$ then the following estimate holds:

$$(5.12) \quad \|(u(t), p(t))\|_{E^1} \leq Q(\|(u(0), p(0))\|_{E^1})e^{-\alpha t} + Q(\|g\|_{L^2})$$

for some positive α and monotone Q .

Finally, we have the analogue of Theorem 4.4 on exponential attractors.

Theorem 5.4. *Let the assumptions of Theorem 2.1 hold and $l > 1$. Then equation (5.1) possesses an exponential attractor $\mathcal{M}$ in E which is a bounded set in the space E^1 .*

The proof of this result repeats word by word the proof of Theorem 4.4 (we have more than enough regularity of solutions $u_1(t)$ and $u_2(t)$ to handle the extra nonlinear term) and by this reason is omitted.

We conclude the exposition by several remarks.

Remark 5.5. We have considered equations (2.1) and (5.1) in the most complicated 3D case only. The 2D case can be treated analogously, but it is actually essentially simpler. Indeed, due to the Sobolev embedding $H^1 \subset L^q$ for all $q < \infty$, the control of the H^1 -norm of the solution u gives the control of the L^2 -norm of $f(u)$ for any growth exponent l , so the restriction $l \leq 2$ can be removed here and any polynomial nonlinearity is *subcritical* in 2D case.

Another simplification comes from the fact that in 2D case the inertial term $B(u, u)$ can be handled without the help of the nonlinearity $f(u)$, so we do not need to require the super-cubic growth rate of $f(u)$. In particular, the purely Navier-Stokes case $f(u)$ is also covered by our theory and gives some new results here as well. For instance, in comparison with [12], we get the E^1 -regularity of the attractor for the case of Dirichlet boundary conditions as well.

Remark 5.6. An interesting question is related with the supercritical case where the nonlinearity grows faster than $u|u|^4$. In the case of incompressible Brinkman-Forchheimer equations as well as in the case of strongly damped wave equations, the restriction $l \leq 2$ in (1.4) is not necessary as shown in [18, 17]. Some methods developed there can be extended to the case of equations (2.1) as well.

For, the existence of weak solutions in this case can be verified based on the energy identity (2.3), their uniqueness follows exactly as in the proof of Theorem 2.4 where only the monotonicity assumption $f'(u) \geq -L$ is actually used. Moreover, the local smoothing property and estimates for $\partial_t u$ stated in Theorem 2.6 also work for the super-critical case as well.

However, there is a problem here which prevents us to treat the supercritical case, namely, the absence of a *dissipative* estimate for the solution u in the energy norm. Indeed, the derivation of such an estimate in Theorem 2.1 is based on multiplication of the equation by $\mathfrak{B}p$, where $\mathfrak{B}$ is a Bogowski operator, but in the supercritical case we cannot do this at least in a direct way since the term $(f(u), \mathfrak{B}p)$ is out of control. We believe that this problem has a technical nature which can be overcome and are planning to return to the supercritical case somewhere else.

APPENDIX A. AN AUXILIARY LINEAR PROBLEM

In this appendix, we study the following linear problem:

$$(A.1) \quad \partial_t p + \operatorname{div}(Du) = 0, \quad -\Delta_x u + \nabla_x p = g(t), \quad p|_{t=0} = p_0, \quad u|_{\partial\Omega} = 0.$$

Note that, solving the second equation of (A.1) with respect to u , we get

$$(A.2) \quad u(t) = -(-\Delta_x)^{-1} \nabla_x p + (-\Delta_x)^{-1} g(t),$$

where the Laplacian is endowed with the homogeneous Dirichlet boundary condition. Inserting this expression to the first equation, we arrive at

$$(A.3) \quad \partial_t p + \mathfrak{A}p = \operatorname{div}(D(-\Delta_x)^{-1} g(t)),$$

where

$$(A.4) \quad \mathfrak{A}p := -\operatorname{div}(D(-\Delta_x)^{-1}\nabla_x p).$$

Thus, the key question here are the properties of the operator $\mathfrak{A}$.

Proposition A.1. *The operator $\mathfrak{A} \in \mathcal{L}(\bar{H}^\delta(\Omega), \bar{H}^\delta(\Omega))$ if $\delta > -\frac{1}{2}$. Moreover, this operator is positive definite and self-adjoint in $\bar{L}^2(\Omega)$:*

$$(A.5) \quad (\mathfrak{A}p, p) \geq \alpha \|p\|_{\bar{L}^2}^2, \quad p \in \bar{L}^2(\Omega)$$

for some $\alpha > 0$.

Proof. Indeed, the first statement is an immediate corollary of the classical elliptic regularity estimates for the Laplacian, see e.g., [37], so we only need to check the stated properties for $\delta = 0$. The fact that $\mathfrak{A}$ is self-adjoint is also straightforward, so we need to verify positiveness. Namely,

$$(A.6) \quad (\mathfrak{A}p, p) = -(\operatorname{div}(Du), p) = (Du, \operatorname{div} p) = \\ = -(Du, \Delta_x u) = (D\nabla_x u, \nabla_x u) \geq \alpha_1 \|\nabla_x u\|_{L^2}^2,$$

where $-\Delta_x u + \nabla_x p = 0$ and $\alpha_1 > 0$ is the smallest eigenvalue of the matrix D . Using, e.g., the Bogovski operator it is easy to show that $\|p\|_{\bar{L}^2}^2 \leq C\|\nabla_x u\|_{L^2}^2$ for some positive constant C . Thus, the proposition is proved. $\square$

As an immediate corollary of this proposition, we get the following result.

Corollary A.2. *Let $p_0 \in \bar{H}^\delta(\Omega)$ and $g \in L^1(0, T; H^{\delta-1}(\Omega))$, $\delta > -\frac{1}{2}$. Then, the solution $p(t)$ of equation (A.3) belongs to $\bar{H}^\delta$ for all $t \geq 0$ and the following estimate holds:*

$$(A.7) \quad \|p(t)\|_{\bar{H}^\delta} \leq C_\delta \|p(0)\|_{\bar{H}^\delta} e^{K_\delta t} + C_\delta \int_0^t e^{K_\delta(t-\tau)} \|g(\tau)\|_{H^{\delta-1}} d\tau,$$

where the constants C_δ and K_δ depend only on δ . In particular, for $\delta = 0$, the corresponding exponent $K_0 = -\alpha < 0$.

Remark A.3. The result of Corollary A.2 gives the dissipative estimate for $\delta = 0$ only. For other values of δ , the constant K_δ a priori may be positive, then the obtained estimate will be not dissipative. This is related with the fact that we do not know a priori that the spectrum of operator $\mathfrak{A}$ is the same in all Sobolev spaces $\bar{H}^\delta(\Omega)$, so if it depends on δ , then it may happen that equation (A.3) may become unstable for some values of δ . We expect that, in a fact, the spectrum of $\mathfrak{A}$ is *independent* of δ , but failed to find the proper reference. So, in order to avoid the technicalities, we restrict ourselves to the most important for our purposes case $\delta = 1$ and verify that the corresponding K_1 is also negative.

Proposition A.4. *Let $p_0 \in \bar{H}^1(\Omega)$ and $g \in L^1(0, T; L^2(\Omega))$. Then, the solution $p(t)$ of the truncated problem (A.3) satisfied the following estimate:*

$$(A.8) \quad \|p(t)\|_{\bar{H}^1} \leq C \|p(0)\|_{\bar{H}^1} e^{-\alpha t} + C \int_0^t \|g(s)\|_{L^2} ds,$$

where the positive constants C and α are independent of t and p .

Proof. In the case of *periodic* boundary conditions, the desired estimate can be obtained just by multiplying equation (A.3) by $\Delta_x p$. However, this does not work in the case of Dirichlet boundary conditions because of the presence of extra boundary integrals arising after integration by parts. So, in this case we will use the localization technique instead. Note also that we only need to verify (A.8) for $g = 0$. The general case will follow then from the Duhamel formula.

Step 1. Interior estimates. Let us fix a non-negative cut-off function $\phi(x) \in C_0^1(\mathbb{R})$ such that $\phi(x) = 0$ if x is in the $\mu/2$ -neighbourhood of the boundary $\partial\Omega$ and $\phi \equiv 1$ if $x \in \Omega$ and is outside the μ -neighbourhood of Ω . In addition, we require that

$$|\nabla_x \phi(x)| \leq C\phi(x)^{1/2}, \quad x \in \Omega.$$

It is not difficult to see that such a function exists for all $\mu > 0$ small enough.

We write equation (A.3) as a system (A.1) with $g = 0$ (in order to avoid the inverse Laplacian) and multiply the first equation by $-\text{div}(\phi \nabla_x p)$. Then, after integration by x , we get

$$\begin{aligned} \text{(A.9)} \quad & \frac{1}{2} \frac{d}{dt} (\phi, |\nabla_x p|^2) = (\text{div}(Du), \text{div}(\phi \nabla_x p)) = \\ & = \sum_{i=1}^3 (\text{div}(Du), \partial_{x_i}(\phi \partial_{x_i} p)) = - \sum_{i=1}^3 (Du, \partial_{x_i}(\phi \partial_{x_i} \nabla_x p)) - \\ & - \sum_{i=1}^3 (Du, \partial_{x_i}(\nabla_x \phi \partial_{x_i} p)) = - \sum_{i=1}^3 (\partial_{x_i}(D \partial_{x_i} u), \nabla_x p) + \\ & + (\text{div}(Du), \nabla_x \phi \cdot \nabla_x p) = - \sum_{i=1}^3 (\partial_{x_i}(D \phi \partial_{x_i} u), \Delta_x u) + \\ & + (\text{div}(Du), \nabla_x \phi \cdot \nabla_x p) = -(\phi D \Delta_x u, \Delta_x u) - (D \Delta_x u \cdot \nabla_x \phi, \text{div}(u)) + \\ & + (\text{div}(Du), \nabla_x \phi \cdot \nabla_x p) \leq -\alpha_1(\phi, |\Delta_x u|^2) + \frac{\alpha_1}{4}(\phi, |\Delta_x u|^2) + \\ & + \frac{\alpha_1}{4}(\phi, |\nabla_x p|^2) + C \|\nabla_x u\|_{L^2}^2 \leq -\frac{\alpha_1}{2}(\phi, |\nabla_x p|^2) + C \|p\|_{L^2}^2. \end{aligned}$$

Since we have already known from Corollary A.2 that

$$\text{(A.10)} \quad \|p(t)\|_{L^2} + \|\nabla_x u(t)\|_{L^2}^2 \leq C e^{-\alpha t} \|p(0)\|_{L^2},$$

then applying the Gronwall inequality to the obtained relation, we get the desired interior dissipative estimate:

$$\text{(A.11)} \quad (\phi, |\nabla_x p(t)|^2) \leq C ((\phi, |\nabla_x p(0)|^2) + \|p(0)\|_{L^2}^2) e^{-\beta t},$$

where C and β are some *positive* constants.

Step 2. Boundary estimates: tangential directions. Let us introduce in a small neighbourhood of the boundary three smooth orthonormal vector fields

$$\tau_3(x) := n = (n^1(x), n^2(x), n^3(x)), \quad \tau_1(x) := (\tau_1^1(x), \tau_1^2(x), \tau_1^3(x))$$

and $\tau_2(x) := (\tau_2^1(x), \tau_2^2(x), \tau_2^3(x))$ such that $n(x)$ coincides with the outer normal vector when $x \in \partial\Omega$ and $\tau_1(x), \tau_2(x)$ give the complement pair of tangential vectors. This triple of vector field may not exist globally near the

boundary, but only locally, so being pedantic we need to use the partition of unity near the boundary to localize them, but we ignore this standard procedure in order to avoid technicalities (this localization can be done exactly in the way how we get interior estimates). After defining the triple of vector fields near the boundary, we use the proper scalar cut-off function in order to extend these fields to the whole domain $\bar{\Omega}$.

Let us define the corresponding differentiation operators along these vector fields:

$$\partial_{\tau_i} u := \sum_{j=1}^3 \tau_i^j(x) \partial_{x_j} u$$

In contrast to the differentiation with respect to coordinate directions, these operators do not commute in general, but their commutator is a lower order operator (again first order differential operator):

$$[\partial_{\tau_i}, \partial_{\tau_j}] = \partial_{\{\tau_i, \tau_j\}},$$

where $\{\tau_i, \tau_j\}$ is a Lie bracket of vector fields τ_i and τ_j . This commutation up to lower order terms is important for our method. One more crucial fact for us is that the condition $u|_{\partial\Omega} = 0$ implies that $\partial_{\tau_i} u|_{\partial\Omega} = 0$, $i = 1, 2$ so differentiation with respect to tangential derivatives preserve the Dirichlet boundary conditions.

We are now ready to get the desired estimates for tangential derivatives. To this end, we denote $q := \partial_{\tau_i} p$ and $v = \partial_{\tau_i} u$. Then, differentiating the equations (A.1) in the direction τ_i , we arrive at

$$(A.12) \quad \partial_t q + \operatorname{div}(Dv) + M(x) \nabla_x u = 0, \quad -\Delta_x v + \nabla_x q = N(x) \nabla_x p + Ru,$$

where the matrices M and N are smooth and R is a linear second order differential operator with smooth coefficients. Multiplying the first and second equations of (A.12) by q and Dv respectively, we arrive at

$$(A.13) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|q\|_{L^2}^2 &= (Dv, \nabla_x q) - (M \nabla_x u, q) - (D \nabla_x v, \nabla_x v) - \\ &\quad - (\nabla_x q, Dv) + (N \nabla_x p, Dv) + (Ru, Dv) \leq \\ &\leq -\alpha_1 \|\nabla_x v\|_{L^2}^2 + \varepsilon \|\nabla_x v\|_{L^2}^2 + \varepsilon \|q\|_{L^2}^2 + C_\varepsilon (\|\nabla_x u\|_{L^2}^2 + \|p\|_{L^2}^2), \end{aligned}$$

where $\varepsilon > 0$ can be arbitrarily small. Moreover, multiplying the second equation by Bq after the standard estimates, we get

$$(A.14) \quad \|q\|_{L^2}^2 \leq C \|\nabla_x v\|_{L^2}^2 + C \|\nabla_x u\|_{L^2}^2 + C \|p\|_{L^2}^2.$$

Inserting this estimate in (A.13) and fixing $\varepsilon > 0$ to be small enough, we finally arrive at

$$(A.15) \quad \frac{d}{dt} \|q\|_{L^2}^2 + \bar{\alpha} \|q\|_{L^2}^2 \leq C \|\nabla_x u\|_{L^2}^2 + \|p\|_{L^2}^2$$

for some $\bar{\alpha} > 0$. Applying the Gronwall inequality to this relation and using (A.10), we have

$$(A.16) \quad \|\partial_{\tau_1} p(t)\|_{L^2}^2 + \|\partial_{\tau_2} p(t)\|_{L^2}^2 \leq C e^{-\alpha t} \|\nabla_x p(0)\|_{L^2}^2,$$

where $\alpha > 0$ and C are independent of p and t . Thus, the desired estimates for tangential derivatives are obtained.

Step 3. Boundary estimates: normal direction. We now want to estimate the normal derivative $\partial_n p$ using equations (A.1) and the already obtained estimates for the tangential derivatives. To this end, we need some preparations. Let us write the vector u in the form

$$u = u_n n + u_{\tau_1} \tau_1 + u_{\tau_2} \tau_2, \quad u_n := u \cdot n, \quad u_{\tau_i} = u \cdot \tau_i.$$

Then, multiplying the second equation of (A.1) by τ_i , $i = 1, 2$ and using the fact that the L^2 -norm of $\partial_{\tau_i} p$ as well as H^1 -norm of u are already estimated, we get

$$(A.17) \quad \|u_{\tau_1}\|_{H^2} + \|u_{\tau_2}\|_{H^2} \leq C e^{-\alpha t} \|p(0)\|_{H^1}.$$

Moreover, multiplying the second equation of (A.1) by n and using that the H^1 -norms of tangential derivatives of u are already under the control, we arrive at

$$(A.18) \quad \|\partial_n^2 u_n(t) - \partial_n p(t)\|_{L^2} \leq C e^{-\alpha t} \|p(0)\|_{H^1}.$$

We now return to the first equation of (A.1) (the equation for pressure). Taking the normal derivative from both sides of this equation and using (A.17) and the fact that the H^1 -norm of $\partial_{\tau_j} u_n$, $j = 1, 2$ are also under the control, we arrive at

$$\partial_t \partial_n p + (Dn \cdot n) \partial_n^2 u_n = h(t), \quad \|h(t)\|_{L^2} \leq C e^{-\alpha t} \|p(0)\|_{H^1}.$$

Multiplying the obtained equation by $\partial_n p$, integrating over x and using (A.18) together with positivity of the matrix D , we finally get

$$\frac{d}{dt} \|\partial_n(t)\|_{L^2}^2 + \alpha_2 \|\partial_n p(t)\|_{L^2}^2 \leq C e^{-2\alpha t} \|p(0)\|_{H^1}^2$$

and applying the Gronwall inequality, we get the desired estimate for the normal derivative:

$$\|\partial_n p(t)\|_{L^2}^2 \leq C e^{-\alpha t} \|p(0)\|_{H^1}^2$$

where $\alpha > 0$ and C are independent of t and u .

Combining together the obtained interior, tangential and normal estimates, we derive that

$$\|p(t)\|_{H^1}^2 \leq C e^{-\alpha t} \|p(0)\|_{H^1}^2$$

and finish the proof of the proposition. $\square$

Corollary A.5. *Let the assumptions of Corollary A.2 hold and let $\delta \in [0, 1]$. Then, the corresponding estimate (A.7) holds with $K_\delta \leq -\alpha < 0$.*

Indeed, we have verified this property for $\delta = 0$ and $\delta = 1$. For fractional values $0 < \delta < 1$, the result follows by the interpolation.

REFERENCES

- [1] E. Aulisa, L. Bloshanskaya, L. Hoang, and A. Ibragimov, *Analysis of Generalized Forchheimer Flows of Compressible Fluids in Porous Media*, J. of Mat. Phys., 50, 103102 (2009).
- [2] A. Babin and M. Vishik, *Attractors of Evolution Equations*, Studies in Mathematics and its Applications, 25. North-Holland Publishing Co., Amsterdam, 1992.
- [3] H. Brinkman, *A Calculation of the Viscous Force Exerted by a Flowing Fluid on a Dense Swarm of Particles*, Applied Scientific Research, vol. 1 (1949), 27–34.

- [4] V. Chepyzhov and M. Vishik, *Attractors for Equations of Mathematical Physics*, American Mathematical Society Colloquium Publications, 49. American Mathematical Society, Providence, RI, 2002.
- [5] D. Donatelli and S. Spirito, *Weak Solutions of Navier-Stokes Equations Constructed by Artificial Compressibility Method are Suitable*, Jour. Hyper. Diff. Eqns, vol. 08, no. 01 (2011), 101–113.
- [6] A. Eden, C. Foias, B. Nicolaenko, and R. Temam, *Exponential Attractors for Dissipative Evolution Equations*, Wiley/Masson, Chichester/Paris, 1994.
- [7] M. Efendiev, A. Miranville and S. Zelik, *Exponential Attractors for a Nonlinear Reaction-Diffusion System in $\mathbb{R}^3$* , C. R. Acad. Sci. Paris, vol. 330 (2000), 713–718.
- [8] M. Efendiev, A. Miranville and S. Zelik, *Exponential Attractors and Finite-Dimensional Reduction for Nonautonomous Dynamical Systems*, Proc. R. Soc. Edinburg Sect. A, vol. 135 (2005), 703–730.
- [9] P. Fabrie, C. Galushinski, A. Miranville, and S. Zelik, *Uniform Exponential Attractors for a Singular Perturbed Damped Wave Equation*, Disc. Cont. Dyn. Sys., vol. 10 (2004), 211–238.
- [10] P. Fabrie and C. Galusinski, *Exponential Attractors for the Slightly Compressible 2D-Navier-Stokes*, DCDS-A, vol. 2, no. 3 (1996), 315–348.
- [11] S. Gatti, V. Pata, and S. Zelik, *A Gronwall-Type Lemma with Parameter and Dissipative Estimates for PDEs*, Nonlinear Anal., vol. 70, no. 6, (2009), 2337–2343.
- [12] J. Ghidaglia and R. Temam, *Long Time Behavior for Partially Dissipative Equations: the Slightly Compressible 2D-Navier-Stokes Equations*, Asympt. Anal., vol. 1 (1988), 23–49.
- [13] R. G. Gordeev, *The Existence of a Periodic Solution in a Tide Dynamics Problem*, Jour. Sov. Mat., vol. 6 (1976), 1–4.
- [14] K. Hajduk and J. Robinson, *Energy Equality for the 3D Critical Convective Brinkman-Forchheimer Equations*, J. Diff. Eqns, v. 263 no. 11 (2017), 7141–7161.
- [15] V. Ipatova, *Solvability of a Tide Dynamics Model in Adjacent Seas*, Russ. J. Numer. Anal. Math. Modelling, vol. 20, no. 1 (2005), 67–79.
- [16] V. Kalantarov and S. Zelik, *A Note on a Strongly Damped Wave Equation with Fast Growing Nonlinearities*, J. Math. Phys., vol. 56, no. 1 (2015), 011501.
- [17] V. Kalantarov and S. Zelik, *Finite-Dimensional Attractors for the Quasi-Linear Strongly Damped Wave Equation*, J. Diff. Eqns., vol. 247, no. 4 (2009), 1120–1155.
- [18] V. Kalantarov and S. Zelik, *Smooth Attractors for the Brinkman-Forchheimer Equations with Fast Growing Nonlinearities*, Comm. Pure Appl. Anal., vol. 11 (2012), 2037–2054.
- [19] O. Ladyzhenskaya, *Mathematical Problems of the Dynamics of a Viscous Incompressible Fluid* (Russian), Nauka, Moscow, 1970.
- [20] O. Ladyzhenskaya, *Attractors for Semi-groups and Evolution Equations* (Lezioni Lincee), Cambridge: Cambridge University Press, 1991.
- [21] A. Likhtarnikov, *Existence and Stability of Bounded and Periodic Solutions in a Nonlinear Problem of Tidal Dynamics* (Russian), in: *The direct method in the theory of stability and its applications*, (Irkutsk, 1979), pp. 83–91, **276**, Nauka, 1981.
- [22] M. Louaked, N. Seloula, S. Sun, and S. Trabelsi, *A Pseudocompressibility Method for the Incompressible Brinkman-Forchheimer Equations*, Differential and Integral Equations, vol. 28 (2015), 361–382.
- [23] U. Manna, J. Menaldi, and S. Sritharan, *Stochastic Analysis of Tidal Dynamics Equation*, in Infinite dimensional stochastic analysis, QPPQ: Quantum Probab. White Noise Anal., vol. 22, World Sci. Publ., Hackensack, NJ, 2008, 90–113.
- [24] G. Marchuk and B. Kagan, *Dynamics of Ocean Tides*. Kluwer Academic Publ., 1989.
- [25] P. Markowich, E. Titi, and S. Trabelsi, *Continuous Data Assimilation for the Three-Dimensional Brinkman-Forchheimer-Extended Darcy Model*. Nonlinearity, vol. 29, no. 4 (2016), 1292–1328.
- [26] X. Mei, A. Savostianov, C. Sun and S. Zelik, *Infinite Energy Solutions for Weakly Damped Quintic Wave Equations in $\mathbb{R}^3$* , arXiv:2004.11864, submitted.

- [27] A. Miranville and S. Zelik, *Attractors for Dissipative Partial Differential Equations in Bounded and Unbounded Domains*, Handbook of diff. eqns: evolutionary equations. Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam IV, (2008), 103–200.
- [28] M.T. Mohan, *On the Two-Dimensional Tidal Dynamics System: Stationary Solution and Stability*, Applicable Analysis (2018), 1–32.
- [29] M. Muskat, *The Flow of Homogeneous Fluids Through Porous Media*, McGraw-Hill, New York, 1937.
- [30] V. Pata and S. Zelik, *Smooth Attractors for Strongly Damped Wave Equations*, Nonlinearity, vol. 19, no. 7 (2006), 1495–1506.
- [31] V. Pata, *Uniform Estimates of Gronwall Type*, J. Mat. Anal. Appl., vol. 373, no. 1 (2011), 264–270.
- [32] K. Rajagopal, *On a Hierarchy of Approximate Models for Flows of Incompressible Fluids through Porous Solids*, Math. Models Methods Appl. Sci., vol. 17 (2007), 215–252.
- [33] H. Sohr, *The Navier-Stokes Equations. An Elementary Functional Analytic Approach*. Birkhäuser, 2001.
- [34] B. Straughan, *Stability and Wave Motion in Porous Media*, Applied Mathematical Sciences, vol. 165, Springer, New York, 2008.
- [35] R. Temam, *Infinite Dimensional Dynamical Systems in Mechanics and Physics*, Springer Verlag, 1997.
- [36] R. Temam, *Navier-Stokes Equations and Nonlinear Functional Analysis*, vol. 66, Siam, 1995.
- [37] H. Triebel, *Interpolation Theory, Function Space, Differential Operators*, North-Holland, Amsterdam-New York, 1978.
- [38] S. Whitaker, *The Forchheimer Equation: A Theoretical Development*, Transp. Porous Media, vol. 25, (1996), 27–61.
- [39] B. Wang and S. Lin, *Existence of Global Attractors for the Three-Dimensional Brinkman-Forchheimer Equation*, Math. Meth. Appl. Sci., vol. 31 (2008), 1479–1495.
- [40] Y. You Y, C. Zhao, and S. Zhou, *The Existence of Uniform Attractors for 3D Brinkman-Forchheimer Equations*, Disc. Cont. Dyn. Syst., vol. 32 (2012), 3787–3800.
- [41] S. Zelik, *Asymptotic Regularity of Solutions of a Nonautonomous Damped Wave Equation with a Critical Growth Exponent*, Comm. Pure Appl. Anal., vol. 3, no. 4 (2004), 921–934.
- [42] S. Zelik, *Spatially Non-Decaying Solutions of 2D Navier-Stokes Equations in a Strip*, Glasgow Math Jour., vol. 49 (2007), 525–588.

¹ DEPARTMENT OF MATHEMATICS,
KOÇ UNIVERSITY, RUMELIFENERI YOLU, SARIYER, ISTANBUL, TURKEY

² DEPARTMENT OF GENERAL AND APPLIED MATHEMATICS,
AZERBAIJAN STATE OIL AND INSUSTRY UNIVERSITY, BAKU, AZERBAIJAN
E-mail address: vkalantarov@ku.edu.tr

³ DEPARTMENT OF MATHEMATICS,
UNIVERSITY OF SURREY, GU27XH, GUILDFORD, UK

⁴ SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU
730000, P.R. CHINA
E-mail address: s.zelik@surrey.ac.uk