

Estimates on Learning Rates for Multi-Penalty Distribution Regression

Zhan Yu*

Department of Mathematics, Hong Kong Baptist University
224 Waterloo Road, Kowloon Tong, Hong Kong
Email: zhanyu@hkbu.edu.hk; mathyuzhan@gmail.com

Daniel W. C. Ho

Department of Mathematics, City University of Hong Kong
83 Tat Chee Avenue, Kowloon Tong, Hong Kong
Email: madaniel@cityu.edu.hk

June 2020 (first version)

Abstract

This paper is concerned with functional learning by utilizing two-stage sampled distribution regression. We study a multi-penalty regularization algorithm for distribution regression in the framework of learning theory. The algorithm aims at regressing to real-valued outputs from probability measures. The theoretical analysis of distribution regression is far from maturity and quite challenging since only second-stage samples are observable in practical settings. In our algorithm, to transform information of distribution samples, we embed the distributions to a reproducing kernel Hilbert space $\mathcal{H}_K$ associated with Mercer kernel K via mean embedding technique. One of the primary contributions of this work is the introduction of a novel multi-penalty regularization algorithm, which is able to capture more potential features of distribution regression. Optimal learning rates of the algorithm are obtained under mild conditions. The work also derives learning rates for distribution regression in the hard learning scenario $f_\rho \notin \mathcal{H}_K$, which has not been explored in the existing literature. Moreover, we propose a new distribution-regression-based distributed learning algorithm to face large-scale data or information challenges arising from distribution data. The optimal learning rates are derived for the distributed learning algorithm. By providing new algorithms and showing their learning rates, the work improves the existing literature in various aspects.

Keywords: learning theory, distribution regression, distributed learning, integral operator, multi-penalty regularization, learning rate

1 Introduction

In the era of big data, functional data or matrix-valued data have become increasingly prevalent in practical applications within machine learning and statistics. Instead of scalar data settings,

*Corresponding author

these types of data present more challenges and difficulties in handling intricate information. As a result, there is a need for improved regression methods to effectively address these issues. The development of a suitable regression scheme for these types of problems has become a highly desirable goal. Recently, a regression method called distribution regression has emerged as an effective tool to handle data defined on some appropriate Banach spaces ([6], [16], [20], [21]). Specifically, the input data are (probability) distributions on a compact metric space $\tilde{X}$. The proposed method contains two sampling stages to learn the regressor from the distributions to the real-valued outputs. The first-stage sample is made up of distributions, and the second-stage sample is drawn according to these distributions. For the first stage, we define the data set as $\tilde{D} = \{(x_i, y_i)\}_{i=1}^{|D|} \subset X \times Y$, in which $|D|$ is the cardinality of $\tilde{D}$, and each pair (x_i, y_i) is i.i.d. sampled from a meta distribution. X is the input space of probability measures on $\tilde{X}$, and $Y = \mathbb{R}$ is the output space equipped with the standard Euclidean metric. For the second stage, the elements in the sample set $\hat{D} = \{(\{x_{i,s}\}_{s=1}^{d_i}, y_i)\}_{i=1}^{|D|}$ are obtained from the distributions $\{x_i\}_{i=1}^{|D|}$ accordingly, where $x_{i,j} \in \tilde{X}$. Many important machine learning and statistical operations have been found to be directly related to the distribution regression scheme mentioned above, for example, the multi-instance learning and point estimation problems without analytical solution.

In this paper, we consider a mean-embedding-based ridge regression method for distribution regression. Let $H = H(\tilde{K})$ be a reproducing kernel Hilbert space (RKHS) with $\tilde{K} : \tilde{X} \times \tilde{X} \rightarrow \mathbb{R}$ as the reproducing kernel. Let $(\tilde{X}, \mathcal{F})$ be a measurable space with $\mathcal{F}$ being a Borel σ -algebra on $\tilde{X}$. Denote the set of Borel probability measures on $(\tilde{X}, \mathcal{F})$ by $\mathcal{M}_1(\mathcal{F})$. Then the *mean embedding* of a distribution $x \in \mathcal{M}_1(\mathcal{F})$ to an element μ_x of RKHS H is given by

$$\mu_x = \int_{\tilde{X}} \tilde{K}(\cdot, \xi) dx(\xi).$$

In practical applications, a class of kernels of great interest are the characteristic kernels (e.g. [6], [21]), of which the mean embedding maps are injective. Some well-known universal kernels are shown to be characteristic such as Gaussian RBF kernel, exponential kernel and binomial kernel. Denote the set of the mean embeddings by $X_\mu = \{\mu_x : x \in \mathcal{M}_1(\mathcal{F})\} \subseteq H$ and the mean embeddings of $\tilde{D}$ to X_μ by $D = \{(\mu_{x_i}, y_i)\}_{i=1}^{|D|}$. Let ρ be the μ -induced probability measure on the product space $Z = X_\mu \times Y$. The regression function with respect to the measure ρ is defined by

$$f_\rho(\mu_x) = \int_Y y d\rho(y|\mu_x), \quad \mu_x \in X_\mu, \quad (1.1)$$

in which $\rho(\cdot|\mu_x)$ is the conditional probability measure of ρ induced at $\mu_x \in X_\mu$. f_ρ is just the minimizer of the least square problem

$$\mathcal{E}(f) = \int_Z (f(\mu_x) - y)^2 d\rho.$$

Generally, the measure ρ is unknown. In the scenario of distribution regression, the distributions $\{x_i\}_{i=1}^{|D|}$ are still unknown, we are only able to access their information by the random sample

$$\hat{D} = \{(\{x_{i,s}\}_{s=1}^{d_i}, y_i)\}_{i=1}^{|D|}$$

with size $d_i \in \mathbb{N}$, $i = 1, 2, \dots, |D|$ respectively. In the context of distribution regression, in a reproducing kernel Hilbert space $(\mathcal{H}_K, \|\cdot\|_K)$ associated with a Mercer kernel $K : X_\mu \times X_\mu \rightarrow \mathbb{R}$, the conventional regularization approach follows the functional optimization scheme

$$f_{\hat{D}, \lambda} = \arg \min_{f \in \mathcal{H}_K} \left\{ \frac{1}{|D|} \sum_{i=1}^{|D|} (f(\mu_{\hat{x}_i}) - y_i)^2 + \lambda \|f\|_K^2 \right\}, \quad (1.2)$$

in which $\hat{x}_i = \frac{1}{d_i} \sum_{s=1}^{d_i} \delta_{x_{i,s}}$ serves as the empirical distribution determined by the observable quantity $\hat{D} = \{(\{x_{i,s}\}_{s=1}^{d_i}, y_i)\}_{i=1}^{|D|}$, $\mu_{\hat{x}_i} = \frac{1}{d_i} \sum_{s=1}^{d_i} \tilde{K}(\cdot, x_{i,s})$ is the mean embedding of $\hat{x}_i$, and $\lambda > 0$ is the regularization parameter. (1.2) is essentially a Tikhonov regularized scheme in RKHS. It is an extension of the one-stage kernel ridge regression scheme.

In recent years, there has been extensive research focused on the setting of one-stage-sampling regularized least squares algorithm ([4], [5], [7], [8], [12], [13], [17], [22], [23], [24]). In the one-stage setting, learning rates of regularized least squares schemes have been thoroughly investigated using kernel method in learning theory (for kernel methods, refer to e.g., [4], [8], [22]). However, there is still a need for further exploration of regularized least squares algorithms with two sampling stages. The existing theoretical analysis on learning rates for distribution regression is only provided in [6] and [21]. These works demonstrate that the optimal minimax learning rates for Algorithm (1.2) can be obtained through an integral operator approach under certain mild conditions. It can be observed that, the above studies on distribution regression are still confined to a single-penalty category, where only one regularization term appears in Algorithm (1.2) with a corresponding single parameter λ . On the other hand, the multi-penalty regularization, as a stable and robust method in one stage-sampling kernel ridge regression setting (e.g. [1], [2], [7], [9], [11], [14], [17], [25]), has been demonstrated to possess many advantages over single-penalty regularization. In contrast to single penalty regularization, it can incorporate any prior information into additional penalties. It thus can simultaneously include various features in regularized solutions, such as boundedness, monotonicity, and smoothness. With these merits, the multi-penalty regularization scheme has been widely used in a variety of inspiring applications like image reconstruction, earth gravity potential reconstruction, option pricing models, data detection, and sparsity analysis. In the context of learning theory, the learning rates of multi-penalty Tikhonov regularization have been primarily studied in [1] and [7]. However, the exploration of the multi-penalty regularization scheme in the literature on distribution regression remains incomplete and requires further investigation. Inspired by the aforementioned advantages of the multi-penalty regularization scheme, we aim to explore its performance in a distribution regression setting to capture more potential features of distribution regression.

In this paper, we investigate a more general framework in two-stages distribution regression by considering the novel multi-penalty regularization scheme:

$$f_{\hat{D}, \lambda_1, \lambda_2} = \arg \min_{f \in \mathcal{H}_K} \left\{ \frac{1}{|D|} \sum_{i=1}^{|D|} (f(\mu_{\hat{x}_i}) - y_i)^2 + \lambda_1 \|f\|_K^2 + \lambda_2 \|V_D f\|_K^2 \right\}. \quad (1.3)$$

In (1.3), $V_D : \mathcal{H}_K \rightarrow \mathcal{H}_K$ is a bounded linear operator which may depend on the data set $\tilde{D}$ and its related first-stage mean embedding set D . The goal of distribution regression in this paper is to learn the regression function by utilizing Algorithm (1.3) based on the given training samples $\hat{D} = \{(\{x_{i,j}\}_{j=1}^{d_i}, y_i)\}_{i=1}^{|D|}$ with $x_{i,1}, x_{i,2}, \dots, x_{i,d_i} \sim x_i$ (i.i.d.). This paper aims to investigate the learning rates of multi-penalty distribution regression (1.3). By deriving optimal learning rates for the proposed multi-penalty distribution regression scheme via an integral operator approach, we improve the results in the literature. On the other hand, in the real world, with the development of data mining, large-scale data are always collected in various application domains, including financial engineering, medicine, business analysis, personal social network, sensor network, and monitoring. In these applications, sensitive data, such as personal data, are always trained in machine learning for different requirements. Hence, it is essential to protect data privacy. In recent years, distributed learning has been shown to be a powerful strategy for tackling privacy-preserving problems (e.g. [5], [7], [8], [12], [13], [18]). On the other hand, the unprecedentedly large data size and complexity of distribution samples would always raise the difficulties of computing in the distribution regression approach. Large-scale data would add unpredictable storage burdens and

memory capacity for a single machine. Meanwhile, it would take a huge amount of time for a single machine to process scientific computing on distribution regression. Motivated by these challenges, and building upon the proposed multi-penalty distribution regression algorithm, we introduce a novel distributed learning method to address these difficulties in the realm of distribution regression. The distributed learning method with multi-penalty distribution regression in this paper is based on a divide-and-conquer approach. Specifically, for the given data set $\tilde{D}$, our distributed learning algorithm starts with partitioning the mean-embedding data set D (related to $\tilde{D}$) and its associated $\hat{D}$ into m disjoint sub-datasets $\{D_j\}_{j=1}^m$ and $\{\hat{D}_j\}_{j=1}^m$ with corresponding disjoint union

$$D = \bigcup_{i=1}^m D_j, \quad \hat{D} = \bigcup_{i=1}^m \hat{D}_j.$$

Subsequently, we assign the corresponding local mean-embedding data set D_j and its associated second-stage local sample set $\hat{D}_j$ to one machine (processor) to produce a local estimator $f_{\hat{D}_j, \lambda_1, \lambda_2}$ in RKHS by the multi-penalty distribution regression scheme (1.3). Once these local estimators are communicated to a central processor, the global estimator $\overline{f_{\hat{D}, \lambda_1, \lambda_2}}$ is synthesized by the central processor through the following averaging process

$$\overline{f_{\hat{D}, \lambda_1, \lambda_2}} = \sum_{j=1}^m \frac{|D_j|}{|D|} f_{\hat{D}_j, \lambda_1, \lambda_2} \quad (1.4)$$

using the local estimators $\{f_{\hat{D}_j, \lambda_1, \lambda_2}\}_{j=1}^m$. For Algorithm (1.4), the learning theory analysis for $\overline{f_{\hat{D}, \lambda_1, \lambda_2}}$ is conducted through an integral operator approach.

We summarize some main contributions of the work. In this work, a novel multi-penalty regularization scheme is proposed for distribution regression. By incorporating additional penalties, Algorithm (1.3) becomes more flexible to capture features of regressor when regressing from distribution samples. By integral operator approach, we study the learning rates of Algorithm (1.3) under different restrictive conditions on regression function f_ρ . Optimal learning rates are derived when $f_\rho \in \mathcal{H}_K$. In kernel-based learning theory, when the target function does not belong to the underlying kernel space, the corresponding problem is often referred to as a hard learning problem ([15]). In this paper, we also provide rigorous analysis and derive learning rates for the hard learning case $f_\rho \notin \mathcal{H}_K$. To the best of our knowledge, this paper is the first work to study distribution regression in a hard learning scenario. The learning rates in this paper improve the existing achievable results. Finally, based on Algorithm (1.3), a novel multi-penalty distribution regression-based distributed learning algorithm (1.4) is proposed. Optimal rates are achieved for this distributed learning algorithm. We will compare the main results in this work with those in the literature in Section 3.

2 Notations, assumptions and main results

We assume that, throughout the paper, there exists a constant $M > 0$ such that $|y| \leq M$ almost surely. $\tilde{K}$ and K are bounded Mercer kernel (symmetric, continuous, positive semidefinite) with bounds $B_{\tilde{K}}$ and κ :

$$B_{\tilde{K}} = \sup_{v \in \tilde{X}} \tilde{K}(v, v) < \infty, \quad \kappa = \sup_{\mu_u \in X_\mu} \sqrt{K(\mu_u, \mu_u)} < \infty. \quad (2.1)$$

Suppose that $\alpha \in (0, 1]$ and $L > 0$. Denote $\mathcal{L}(Y, \mathcal{H}_K)$ as the Banach space of the bounded linear operators from space Y to $\mathcal{H}_K$. Denote $K_{\mu_x} = K(\mu_x, \cdot)$, $\mu_x \in X_\mu$. We treat K_{μ_x} as an element of

$\mathcal{L}(Y, \mathcal{H}_K)$ by defining the linear mapping

$$K_{\mu_x}(y) = yK_{\mu_x}, \quad y \in Y.$$

We assume that the mapping $K_{(\cdot)} : X_\mu \rightarrow \mathcal{L}(Y, \mathcal{H}_K)$ is (α, L) -Hölder continuous in the following sense

$$\|K_{\mu_x} - K_{\mu_y}\|_{\mathcal{L}(Y, \mathcal{H}_K)} \leq L \|\mu_x - \mu_y\|_H^\alpha, \quad \forall (\mu_x, \mu_y) \in X_\mu \times X_\mu. \quad (2.2)$$

Due to the fact in [21], the set of mean embeddings X_μ is a separable compact set of the continuous function space defined on $\tilde{X}$. Denote ρ_{X_μ} as the marginal distribution of ρ on X_μ . Let $L_{\rho_{X_\mu}}^2$ be the Hilbert space of square-integrable functions defined on X_μ . For $f \in L_{\rho_{X_\mu}}^2$, denote the $L_{\rho_{X_\mu}}^2$ -norm of f by

$$\|f\|_\rho = \|f\|_{L_{\rho_{X_\mu}}^2} = \langle f, f \rangle_{\rho_{X_\mu}}^{1/2} = \left(\int_{X_\mu} |f(\mu_x)|^2 d\rho_{X_\mu}(\mu_x) \right)^{1/2}.$$

Define the integral operator L_K on $L_{\rho_{X_\mu}}^2$ associated with the Mercer kernel $K : X_\mu \times X_\mu \rightarrow \mathbb{R}$ as

$$L_K(f) = \int_{X_\mu} K_{\mu_x} f(\mu_x) d\rho_{X_\mu}, \quad f \in L_{\rho_{X_\mu}}^2. \quad (2.3)$$

Because the set of mean embeddings X_μ is compact and K is a Mercer kernel, we know L_K is a positive compact operator on $L_{\rho_{X_\mu}}^2$. Then for any $r > 0$, its r -th power L_K^r is well-defined according to spectral theorem in functional calculus. Throughout the paper, we assume the following *regularity condition* for the regression function f_ρ :

$$f_\rho = L_K^r(g_\rho) \text{ for some } g_\rho \in L_{\rho_{X_\mu}}^2, \quad r > 0. \quad (2.4)$$

The above assumption means that the regression function lies in the range of operator L_K^r , the special case $r = 1/2$ corresponds to $f_\rho \in \mathcal{H}_K$. We use the *effective dimension* $\mathcal{N}(\lambda_1)$ to measure the capacity of $\mathcal{H}_K$ with respect to the measure ρ_{X_μ} , which is defined to be the trace of the operator $(\lambda_1 I + L_K)^{-1} L_K$, that is

$$\mathcal{N}(\lambda_1) = \text{Tr}((\lambda_1 I + L_K)^{-1} L_K), \quad \lambda_1 > 0. \quad (2.5)$$

For the effective dimension $\mathcal{N}(\lambda_1)$, we need the following capacity assumption, which focuses on the β -rate of $\mathcal{N}(\lambda_1)$: there exists a constant $\mathcal{C}_0$ such that for any $\lambda_1 > 0$,

$$\mathcal{N}(\lambda_1) \leq \mathcal{C}_0 \lambda_1^{-\beta}, \quad \text{for some } 0 < \beta \leq 1. \quad (2.6)$$

In this paper, we assume that the sample set $D = \{(\mu_{x_i}, y_i)\}_{i=1}^{|D|}$ is drawn independently according to the probability measure ρ . Sample $\{x_{i,s}\}_{s=1}^{d_i}$ is drawn independently according to probability distribution x_i for $i = 1, 2, \dots, |D|$. For the operator V_D , throughout the paper, we assume that there is a constant $c_V > 0$ independent of the data set such that

$$\|V_D^T V_D\| \leq c_V \quad \text{a.s.}, \quad (2.7)$$

in which V_D^T denotes the adjoint operator of V_D and $\|\cdot\|$ is the operator norm. Since $\|V_D\|^2 = \|V_D^T V_D\|$, therefore an equivalent assumption

$$\|V_D\| \leq \sqrt{c_V}, \quad \text{a.s.},$$

can also be used in this paper. In the following, we denote the quantity $\mathcal{B}_{|D|,\lambda_1}$ and $\mathcal{B}'_{|D|,\lambda_1}$ as

$$\mathcal{B}_{|D|,\lambda_1} = \frac{2\kappa}{\sqrt{|D|}} \left(\frac{\kappa}{\sqrt{|D|\lambda_1}} + \sqrt{\mathcal{N}(\lambda_1)} \right), \quad (2.8)$$

$$\mathcal{B}'_{|D|,\lambda_1} = \frac{1}{|D|\sqrt{\lambda_1}} + \frac{\sqrt{\mathcal{N}(\lambda_1)}}{\sqrt{|D|}}. \quad (2.9)$$

It can be observed that $\mathcal{B}_{|D|,\lambda_1}$ and $\mathcal{B}'_{|D|,\lambda_1}$ only differ in κ -scaling sense. $\mathcal{B}_{|D|,\lambda_1}$ and $\mathcal{B}'_{|D|,\lambda_1}$ will be used in error upper bound representation and operator estimates in subsequent analysis.

Consider the estimator $f_{\widehat{D},\lambda_1,\lambda_2}$ defined in Algorithm (1.3). We present our main results on error estimate between $f_{\widehat{D},\lambda_1,\lambda_2}$ and f_ρ as detailed in the following theorems. These results offer estimates on the expected difference between $f_{\widehat{D},\lambda_1,\lambda_2}$ and f_ρ in $L^2_{\rho_{X_\mu}}$ -norm with the expectation taken for both D and $\widehat{D}$. The first theorem presents a general error estimate without imposing decaying restrictions on the effective dimension $\mathcal{N}(\lambda_1)$.

Theorem 1. *Assume that the regularity condition (2.4) holds with $1/2 \leq r \leq 1$, $|y| \leq M$ holds almost surely, the operator boundedness condition (2.7) holds and the mapping $K_{(\cdot)} : X_\mu \rightarrow \mathcal{L}(Y, \mathcal{H}_K)$ is (α, L) -Hölder continuous with $\alpha \in (0, 1]$ and $L > 0$. If the regularization parameters $\lambda_1, \lambda_2 \in (0, 1)$ satisfy the relation $2c_V \lambda_2 = \lambda_1^{2r}$ and $\widetilde{d}$ satisfies $\frac{1}{d^{\frac{\alpha}{2}}} = \frac{1}{|D|} \sum_{i=1}^{|D|} \frac{1}{d_i^{\alpha/2}}$, then we have*

$$\begin{aligned} & E \left[\|f_{\widehat{D},\lambda_1,\lambda_2} - f_\rho\|_\rho \right] \\ & \leq 2(2\sqrt{2}(2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_K^{\frac{\alpha}{2}}}{\lambda_1 \widetilde{d}^{\frac{\alpha}{2}}} + 2) \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \frac{2^{\frac{\alpha}{2}} B_K^{\frac{\alpha}{2}}}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \left[(2 + \sqrt{\pi})^{\frac{1}{2}} LM(2\Gamma(3) + \log^2 2) \right. \\ & \quad + 2(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \left(2\sqrt{6}(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \frac{M}{\kappa} \frac{\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right) + \sqrt{3}(\lambda_1 + \lambda_2 c_V) \times \right. \\ & \quad \left. \left. \lambda_1^{r-\frac{3}{2}} 2^{r-\frac{1}{2}} \|g_\rho\|_\rho (2\Gamma(4r-1) + \log^{4r-2} 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2r-1} + \sqrt{3} \kappa^{r-\frac{1}{2}} \|g_\rho\|_\rho \right) \right] \\ & \quad + 4(2\Gamma(3) + \log^2 2)^{\frac{1}{2}} (2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \frac{M}{\kappa} \mathcal{B}_{|D|,\lambda_1} \\ & \quad + (2\Gamma(2r+1) + \log^{2r} 2) 2^r (\lambda_1^r + \lambda_1^{r-1} \lambda_2 c_V) \|g_\rho\|_\rho \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2r}. \end{aligned}$$

The next result establishes the explicit learning rates of multi-penalty distribution regression after we further assume the decaying capacity condition on effective dimension $\mathcal{N}(\lambda_1) \leq \mathcal{C}_0 \lambda_1^{-\beta}$, $\forall \lambda_1 > 0$, for some $\beta \in (0, 1]$. The following result on minimax optimal learning rates for multi-penalty distribution regression algorithm (1.3) holds.

Theorem 2. *Assume that the regularity condition (2.4) holds with $1/2 \leq r \leq 1$, $|y| \leq M$ holds almost surely, the capacity condition (2.6) holds with index $\beta \in (0, 1]$, the operator boundedness condition (2.7) holds and the mapping $K_{(\cdot)} : X_\mu \rightarrow \mathcal{L}(Y, \mathcal{H}_K)$ is (α, L) -Hölder continuous with $\alpha \in (0, 1]$ and $L > 0$. If we choose the regularization parameters $\lambda_1 = |D|^{-\frac{1}{2r+\beta}}$, $\lambda_2 = \frac{1}{2c_V} |D|^{-\frac{2r}{2r+\beta}}$ and second-stage sample size $d_1 = d_2 = \dots = d_{|D|} = |D|^{\frac{1+2r}{\alpha(2r+\beta)}}$, then we have*

$$E \left[\|f_{\widehat{D},\lambda_1,\lambda_2} - f_\rho\|_\rho \right] = \mathcal{O}(|D|^{-\frac{r}{2r+\beta}}). \quad (2.10)$$

The above results handle standard setting $r \in [1/2, 1]$, which corresponds to $f_\rho \in \mathcal{H}_K$. The following two results are concerned with error bounds and learning rates in the hard learning scenario $f_\rho \notin \mathcal{H}_K$. The investigation of this scenario does not appear in the literature on distribution regression. The following error bound holds without capacity assumption on effective dimension $\mathcal{N}(\lambda_1)$.

Theorem 3. *Assume that the regularity condition (2.4) holds with $0 < r < 1/2$, $|y| \leq M$ holds almost surely, the operator boundedness condition (2.7) holds and the mapping $K_{(\cdot)} : X_\mu \rightarrow \mathcal{L}(Y, \mathcal{H}_K)$ is (α, L) -Hölder continuous with $\alpha \in (0, 1]$ and $L > 0$. If the regularization parameters $\lambda_1, \lambda_2 \in (0, 1)$ satisfy the relation $2c_V \lambda_2 = \lambda_1$ and $\tilde{d}$ satisfies $\frac{1}{\tilde{d}^{\frac{\alpha}{2}}} = \frac{1}{|D|} \sum_{i=1}^{|D|} \frac{1}{d_i^{\alpha/2}}$, then we have*

$$\begin{aligned}
& E \left[\|f_{\widehat{D}, \lambda_1, \lambda_2} - f_\rho\|_\rho \right] \\
& \leq 2 \left(2\sqrt{2}(2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\lambda_1 \tilde{d}^{\frac{\alpha}{2}}} + 2 \right) (2 + \sqrt{\pi})^{\frac{1}{2}} L M \frac{2^{\frac{\alpha}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} (2\Gamma(3) + \log^2 2) \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \\
& + \left(2\sqrt{2}(2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\lambda_1 \tilde{d}^{\frac{\alpha}{2}}} + 2 \right) \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 (2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \kappa L (2 + \sqrt{\pi})^{\frac{1}{2}} 2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}} \times \\
& \left(2\sqrt{3}(2\Gamma(9) + \log^8 2)^{\frac{1}{4}} (2\Gamma(5) + \log^4 2)^{\frac{1}{4}} \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \left[2M(\kappa + 1) \frac{1}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} \frac{1}{\sqrt{\lambda_1}} \mathcal{B}'_{|D|, \lambda_1} \right. \right. \\
& \left. \left. + 2(\kappa^2 + \kappa) \|g_\rho\|_\rho \frac{\lambda_1^{r-1}}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} \mathcal{B}'_{|D|, \lambda_1} \right] + 2\sqrt{3} \|g_\rho\|_\rho \frac{\lambda_1^{r-\frac{1}{2}}}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} \right) \\
& + 4(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} (2\Gamma(3) + \log^2 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \left[M(\kappa + 1) \mathcal{B}'_{|D|, \lambda_1} + \|g_\rho\|_\rho (\kappa^2 + \kappa) \lambda_1^{r-\frac{1}{2}} \mathcal{B}'_{|D|, \lambda_1} \right] \\
& + 2\|g_\rho\|_\rho c_V \lambda_2 \lambda_1^{r-1} + \lambda_1^r \|g_\rho\|_\rho. \tag{2.11}
\end{aligned}$$

When the capacity condition for $\mathcal{N}(\lambda_1)$ holds, for $r \in (0, 1/2)$, by assigning new $\lambda_1, \lambda_2, d_1, d_2, \dots, d_{|D|}$, we derive following learning rates for the case $f_\rho \notin \mathcal{H}_K$.

Theorem 4. *Assume that the regularity condition (2.4) holds with $0 < r < 1/2$ and $|y| \leq M$ almost surely. Assume the capacity condition (2.6) holds with index $\beta \in (0, 1]$, the operator boundedness condition (2.7) holds and the mapping $K_{(\cdot)} : X_\mu \rightarrow \mathcal{L}(Y, \mathcal{H}_K)$ is (α, L) -Hölder continuous with $\alpha \in (0, 1]$ and $L > 0$. If we choose the regularization parameters $\lambda_1 = |D|^{-\frac{1}{1+\beta}}, \lambda_2 = \frac{1}{2c_V} |D|^{-\frac{1}{1+\beta}}$ and choose $d_1 = d_2 = \dots = d_{|D|} = |D|^{\frac{2}{\alpha(1+\beta)}}$, then we have*

$$E \left[\|f_{\widehat{D}, \lambda_1, \lambda_2} - f_\rho\|_\rho \right] = \mathcal{O}(|D|^{-\frac{r}{1+\beta}}). \tag{2.12}$$

For the multi-penalty distribution regression-based distributed learning algorithm (1.4), under a mild restriction on the machine number m , we derive the following result on optimal learning rate of the estimator $\overline{f_{\widehat{D}, \lambda_1, \lambda_2}}$ generated from our new distributed learning scheme of distribution regression in (1.4).

Theorem 5. *Assume that the regularity condition (2.4) holds with $1/2 \leq r \leq 1$, $|y| \leq M$ almost surely, the capacity condition (2.6) holds and the operator boundedness condition (2.7) holds with a constant $c_V > 0$ independent of data sets. The mapping $K_{(\cdot)} : X_\mu \rightarrow \mathcal{L}(Y, \mathcal{H}_K)$ is assumed to be (α, L) -Hölder continuous with $\alpha \in (0, 1]$ and $L > 0$. If the sample size of the local machines*

satisfies $|D_j| = |D|/m$ for $j = 1, 2, \dots, m$, the penalties satisfy $\lambda_1 = |D|^{-\frac{1}{2r+\beta}}$, $\lambda_2 = \frac{1}{2c_V}|D|^{-\frac{2r}{2r+\beta}}$, the second-stage sample sizes are taken as $d = |D|^{\frac{1+2r}{\alpha(2r+\beta)}}$ and total number m of local machines satisfies

$$m \leq |D|^{\frac{2r-1}{2r+\beta}},$$

then we have

$$E\left[\|\overline{f_{\widehat{D}, \lambda_1, \lambda_2}} - f_\rho\|_\rho\right] = \mathcal{O}(|D|^{-\frac{r}{2r+\beta}}). \quad (2.13)$$

The theoretical result in Theorem 5 indicates that our new proposed distributed estimator $\overline{f_{\widehat{D}, \lambda_1, \lambda_2}}$ can achieve optimal learning rates of $\mathcal{O}(|D|^{-\frac{r}{2r+\beta}})$. Hence our theoretical result provides a satisfactory guarantee for the nice learning ability of $\overline{f_{\widehat{D}, \lambda_1, \lambda_2}}$. Moreover, Theorem 5 provides the possibility that, when handling massive distribution samples in practical computation, we can divide the large-scale distribution data set D into several subsets to reduce the huge computational burden. Hence these facts indicate the great potential of our distributed estimator $\overline{f_{\widehat{D}, \lambda_1, \lambda_2}}$ in practical computation field on handling massive distribution data in the big data era.

3 Related work and discussions

We make some comparisons between this work and those in the existing literature. In the past decade, studies on classical regularized least squares regression are on the way toward maturity. Various approaches are utilized to analyze the minimax learning rates of regularized least squares regression. Recently, studies have just turned to the learning rates of distribution regression. Theoretical investigation on distribution regression is very limited. To the best of our knowledge, the only existing works on learning theory analysis of learning rates of distribution regression scheme (1.2) are contained in [6] and [21]. In Theorem 1 and Theorem 2, we show that an optimal learning rate of

$$E\left[\|f_{\widehat{D}, \lambda_1, \lambda_2} - f_\rho\|_\rho\right] = \mathcal{O}(|D|^{-\frac{r}{2r+\beta}})$$

for multi-penalty distribution regression is obtained for $\frac{1}{2} \leq r \leq 1$. The results improve the work of [21] and [6] to a more general setting. Specifically, the optimal rates cover the case of [21] with $r = 1/2$, in contrast to the suboptimal rate of $E[\|f_{\widehat{D}, \lambda} - f_\rho\|^2] = \mathcal{O}(|D|^{-\frac{2}{5}})$. Also, the rates coincide with the optimal rate in [6], and this work improves the result in [6] to a more general setting by considering the additional penalty with parameter λ_2 and a bounded operator V_D in its multi-penalty regularization framework. It can also be observed that the regression scheme of [6] can be treated as a degenerate case of the scheme (1.3) with $\lambda_2 = 0$. As we do not impose coercive conditions on operator V_D except for its boundedness, Algorithm (1.3) presents potential flexibility in handling distribution regression. In contrast to previous works in the literature of distribution regression, due to the introduction of an additional new penalty induced by the operator V_D , difficulties arise when we need to handle both first-stage operator estimates and second-stage estimates since the operator V_D has already participated in the operator analysis in each part where the operator decomposition process needs to be carried out. To face the analysis challenges arising from the additional penalty induced by the operator V_D , we have rigorously introduced the quantities $\mathcal{A}_{D, \lambda_1, \lambda_2, V_D}$ and $\Omega_{D, \lambda_1, \lambda_2, V_D}$ below in order to address the influence of V_D and perform operator decomposition successfully in the current setting.

On the other hand, all existing results in distribution regression literature consider the case when the regression function $f_\rho \in \mathcal{H}_K$. In kernel-based learning theory, when the target function does not live in the underlying kernel space, the corresponding problem is often referred to as a hard learning problem (see, e.g. [15]) which is very hot in recent studies in the literature. In

kernel ridge regression, the hard learning problems have been intensively studied very recently (e.g. [10], [15], [18], [19]). One of the reasons for investigating hard learning scenario is that, the assumption that f_ρ lies precisely in $\mathcal{H}_K$ is quite restrictive in many practical applications. However, for f_ρ satisfying the regularity condition $r \in (0, 1/2)$, namely $f_\rho \notin \mathcal{H}_K$, which belongs to the hard learning scenario, the learning theory has not been established for distribution regression and the convergence analysis on min-max learning rates has not been carried out. Hence it is interesting to ask whether the multi-penalty distribution regression estimator $\hat{f}_{\widehat{D}, \lambda_1, \lambda_2}$ still maintains a nice learning performance in the hard learning setting. In Theorem 3 and Theorem 4 of this work, we have answered the question by carrying out learning rate analysis and improved the analyzable regularity range from $[1/2, 1]$ for the standard setting $f_\rho \in \mathcal{H}_K$ ([6], [21]) to $r \in (0, 1/2)$ for hard learning setting $f_\rho \notin \mathcal{H}_K$. The theoretical results of Theorem 3 and Theorem 4 fill the gap of the study of hard learning problem in distribution regression.

Let us describe some core differences in theoretical approaches between the current work and previous work [6]. As a starting point of convergence analysis, the previous work [6], in fact, employed an error decomposition in terms of

$$f_{\widehat{D}, \lambda} - f_\rho = (f_{\widehat{D}, \lambda} - f_{D, \lambda}) + (f_{D, \lambda} - f_\rho)$$

in which $f_{D, \lambda}$ is the first-stage counterpart of $f_{\widehat{D}, \lambda}$. For the hard learning scenario $f_\rho \notin \mathcal{H}_K$, one of the difficulties is that such a type of error decomposition approach fails in the tough settings when f_ρ does not lie in the RKHS $\mathcal{H}_K$ since we can not directly handle $f_{D, \lambda} - f_\rho$ via operator representation and the corresponding underlying bounds for estimating $f_{\widehat{D}, \lambda} - f_{D, \lambda}$ also lose. In this work, to handle such a tough setting, we in fact, essentially carry out a further new two-stage error decomposition

$$(f_{\widehat{D}, \lambda_1, \lambda_2} - f_{D, \lambda_1, \lambda_2}) + (f_{D, \lambda_1, \lambda_2} - f_{\lambda_1}) + (f_{\lambda_1} - f_\rho)$$

for the estimator $\hat{f}_{\widehat{D}, \lambda_1, \lambda_2}$ which is generated from our more general multi-penalty regularized regression scheme (1.3), one of the technical novelties of this type two-stage error decomposition is to use the crucial data-free minimizer f_{λ_1} (associated with the first regularization parameter λ_1) defined by $f_{\lambda_1} = \arg \min_{f \in \mathcal{H}_K} \{\|f - f_\rho\|_{L^2_{\rho_{X_\mu}}}^2 + \lambda_1 \|f\|_K^2\}$ as a key bridge to realize core estimates in both the first-stage representation and the second-stage representation (also see discussions in Remark 1 and Remark 2). Fortunately, we also observe that the introduction of f_{λ_1} also performs very well when handling some main terms hidden in the estimates of the second-stage

$$f_{\widehat{D}, \lambda_1, \lambda_2} - f_{D, \lambda_1, \lambda_2}$$

(for example, see Proposition 7 and the proof of Theorem 4) where $f_{D, \lambda_1, \lambda_2}$ is defined in (4.3). These nice performances just right result in a satisfactory mini-max rate of $\mathcal{O}(|D|^{-\frac{r}{1+r}})$ ($r \in (0, 1/2)$) for $f_{\widehat{D}, \lambda_1, \lambda_2} - f_{D, \lambda_1, \lambda_2}$ after using our selection rule for regularization parameters λ_1, λ_2 (see analysis in the proof of Theorem 4). The detailed analysis has been carried out in Section 4.2 (Also see Remark 1 and Remark 2). In addition, due to the multi-penalty nature of our algorithm, we need further take the influence of the operator V_D that induces the additional penalty into consideration in each operator decomposition process. In contrast to previous works, the participation of V_D raises the difficulties of the operator decomposition processes and makes the operator norm estimates in both the first and second stage more intricate. To overcome the difficulties arising from the operator V_D , as mentioned above, we have rigorously introduced the new quantities $\mathcal{A}_{D, \lambda_1, \lambda_2, V_D}$ and $\Omega_{D, \lambda_1, \lambda_2, V_D}$ and successfully realize the operator decomposition process in our setting.

Another contribution of this paper is that, based on a divide-and-conquer approach, we present a novel two-stage multi-penalty distribution-regression-based distributed learning algorithm. It

can be witnessed that, among the existing works in the literature on distribution regression, there is no powerful method for handling large-scale or massive distribution data problems. The theoretical result is also lacking and not established. The existing models of distribution regression mainly focus on the single-machine model. When the data scale of the distribution samples is extremely large, their drawback becomes obvious since the single machine can not perform efficiently when handling massive distribution data. However, an appropriate distributed learning scheme for two-stage distribution regression has not been proposed. One of the difficulties in deriving results of learning rates is that a well-defined two-stage distributed distribution regression estimator is still lacking. Another challenge is that a suitable two-stage error decomposition for the estimator has not yet been established either before. In learning theory, establishing an appropriate error decomposition related to generalization error is often a crucial step to obtaining the desired learning rates of the estimator. The current work fills the gap by proposing the distributed learning estimator $\overline{f_{\widehat{D}, \lambda_1, \lambda_2}}$ (defined in (1.4)) for handling massive distribution data and providing a novel two-stage decomposition for

$$\overline{f_{\widehat{D}, \lambda_1, \lambda_2}} - f_{\widehat{D}, \lambda_1, \lambda_2}$$

via operator representation (See Proposition 8). The optimal rates are obtained for this method in Theorem 5. The mini-max rates coincide with the rates we establish in non-distributed settings. Therefore the new regression method presents the advantages of reducing the computational burden on computing time, storage bottleneck, and memory requirements over a single machine when processing the distribution regression algorithm. Distributed learning algorithms have been extensively studied recently (e.g., [5, 8, 12, 13]). From these existing methods, it can be observed that the distributed learning method of handling distribution or functional data is still unexplored. By presenting the new distributed learning algorithm with multi-penalty distribution regression and proving its optimal rates, this paper provides an effective distributed learning method for handling distribution data. Based on the above discussions and the satisfactory theoretical result in Theorem 5, the new proposed multi-penalty distribution-regression-based algorithm possesses great potential for handling large-scale distribution data in modern computational science and statistical science.

4 Analysis and estimates on learning rates

In this section, we provide the analysis and proofs for the main results of Theorems 1, 2, 3, 4. In order to estimate the expected learning rates of the algorithm, the subsequent estimates for our multi-penalty distribution regression estimator $f_{\widehat{D}, \lambda_1, \lambda_2}$ are based on the following basic two-stage error decomposition

$$\|f_{\widehat{D}, \lambda_1, \lambda_2} - f_\rho\|_\rho \leq \|f_{\widehat{D}, \lambda_1, \lambda_2} - f_{D, \lambda_1, \lambda_2} + f_{D, \lambda_1, \lambda_2} - f_\rho\|_\rho \leq \|f_{\widehat{D}, \lambda_1, \lambda_2} - f_{D, \lambda_1, \lambda_2}\|_\rho + \|f_{D, \lambda_1, \lambda_2} - f_\rho\|_\rho, \quad (4.1)$$

and a further two-stage error decomposition

$$\|f_{\widehat{D}, \lambda_1, \lambda_2} - f_\rho\|_\rho \leq \|f_{\widehat{D}, \lambda_1, \lambda_2} - f_{D, \lambda_1, \lambda_2}\|_\rho + \|f_{D, \lambda_1, \lambda_2} - f_{\lambda_1}\|_\rho + \|f_{\lambda_1} - f_\rho\|_\rho \quad (4.2)$$

where

$$f_{\lambda_1} = \arg \min_{f \in \mathcal{H}_K} \left\{ \|f - f_\rho\|_{L_{\rho, X_\mu}^2}^2 + \lambda_1 \|f\|_K^2 \right\}.$$

In the above decomposition, $f_{D, \lambda_1, \lambda_2}$ is the first-stage counterpart of $f_{\widehat{D}, \lambda_1, \lambda_2}$ which is the minimizer of the following classical multi-penalty regression scheme associated with the first-stage sample

$$D = \{(\mu_{x_i}, y_i)\}_{i=1}^{|D|},$$

$$f_{D, \lambda_1, \lambda_2} = \arg \min_{f \in \mathcal{H}_K} \left\{ \frac{1}{|D|} \sum_{i=1}^{|D|} (f(\mu_{x_i}) - y_i)^2 + \lambda_1 \|f\|_K^2 + \lambda_2 \|V_D f\|_K^2 \right\}. \quad (4.3)$$

This quantity serves as an important bridge in subsequent analysis.

To make a preliminary preparation, we define the sampling operator $S_D : \mathcal{H}_K \rightarrow \mathbb{R}^{|D|}$ associated with the first-stage sample as

$$S_D f = (f(\mu_{x_1}), f(\mu_{x_2}), \dots, f(\mu_{x_{|D|}}))^T, \quad f \in \mathcal{H}_K.$$

The adjoint operator $S_D^T : \mathbb{R}^{|D|} \rightarrow \mathcal{H}_K$ is given by

$$S_D^T \mathbf{c}_D = \frac{1}{|D|} \sum_{i=1}^{|D|} c_i K_{\mu_{x_i}}, \quad \mathbf{c}_D = (c_1, c_2, \dots, c_{|D|})^T \in \mathbb{R}^{|D|}.$$

Then we can define the first-stage empirical operator $L_{K,D}$ of L_K as

$$L_{K,D}(f) = S_D^T S_D(f) = \frac{1}{|D|} \sum_{i=1}^{|D|} f(\mu_{x_i}) K_{\mu_{x_i}} = \frac{1}{|D|} \sum_{i=1}^{|D|} \langle K_{\mu_{x_i}}, f \rangle_K K_{\mu_{x_i}}, \quad f \in \mathcal{H}_K.$$

We also define the sampling operator $\hat{S}_D : \mathcal{H}_K \rightarrow \mathbb{R}^{|D|}$ associated with the second-stage sample as

$$\hat{S}_D f = (f(\mu_{\hat{x}_1}), f(\mu_{\hat{x}_2}), \dots, f(\mu_{\hat{x}_{|D|}}))^T, \quad f \in \mathcal{H}_K.$$

Its adjoint operator $\hat{S}_D^T : \mathbb{R}^{|D|} \rightarrow \mathcal{H}_K$ is given by

$$\hat{S}_D^T \mathbf{c}_D = \frac{1}{|D|} \sum_{i=1}^{|D|} c_i K_{\mu_{\hat{x}_i}}, \quad \mathbf{c}_D = (c_1, c_2, \dots, c_{|D|})^T \in \mathbb{R}^{|D|}.$$

Then the empirical operator of $L_{K,D}$ can be defined accordingly by using the second-stage sample $\hat{D}$ as follows

$$L_{K,\hat{D}}(f) = \hat{S}_D^T \hat{S}_D(f) = \frac{1}{|D|} \sum_{i=1}^{|D|} f(\mu_{\hat{x}_i}) K_{\mu_{\hat{x}_i}} = \frac{1}{|D|} \sum_{i=1}^{|D|} \langle K_{\mu_{\hat{x}_i}}, f \rangle_K K_{\mu_{\hat{x}_i}}, \quad f \in \mathcal{H}_K. \quad (4.4)$$

With these notations, after taking Fréchet derivative on the right hand sides of Algorithm (1.3) and its first-stage associate (4.3), it is easy to see the following representation holds,

$$f_{\hat{D}, \lambda_1, \lambda_2} = (L_{K,\hat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \hat{S}_D^T y_D,$$

$$f_{D, \lambda_1, \lambda_2} = (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} S_D^T y_D,$$

with $y_D = (y_1, y_2, \dots, y_{|D|})^T \in \mathbb{R}^{|D|}$. In the following, we use $E_{\mathbf{z}^{|D|}}[\cdot]$ to denote the expectation w.r.t. $\mathbf{z}^{|D|} = \{z_i = (\mu_{x_i}, y_i)\}_{i=1}^{|D|}$. Use $E_{\mathbf{x}^{\mathbf{d}, |D|} | \mathbf{z}^{|D|}}$ to denote the conditional expectation w.r.t. sample $\{x_{i,s}\}_{s=1}^{d_i} \}_{i=1}^{|D|}$ conditioned on $\{z_1, z_2, \dots, z_{|D|}\}$. Namely

$$E_{\mathbf{z}^{|D|}}[\cdot] = E_{\{(\mu_{x_i}, y_i)\}_{i=1}^{|D|}}[\cdot], \quad E_{\mathbf{x}^{\mathbf{d}, |D|} | \mathbf{z}^{|D|}}[\cdot] = E_{\{x_{i,s}\}_{s=1}^{d_i} \}_{i=1}^{|D|} | \{z_i\}_{i=1}^{|D|}}[\cdot].$$

Following (13) in [20] and Theorem 15 in [3], we know that for any $i = 1, 2, \dots, |D|$,

$$\|\mu_{\hat{x}_i} - \mu_{x_i}\|_H \leq \frac{(1 + \sqrt{\delta})\sqrt{2B_{\tilde{K}}}}{\sqrt{d_i}}$$

with probability $1 - e^{-\delta}$. Set $\xi = \|\mu_{\hat{x}_i} - \mu_{x_i}\|_H^{2\alpha}$ with $\alpha \in (0, 1]$. The boundedness condition on kernel $\tilde{K}$ implies $\xi \in [0, 2^{2\alpha} B_{\tilde{K}}^\alpha]$ almost surely. Then it follows that $E[\xi] = \int_0^\infty \text{Prob}_{\{x_{i,s}\}_{s=1}^{d_i}|x_i}(\xi > t)dt = \int_0^{2^{2\alpha} B_{\tilde{K}}^\alpha} \text{Prob}_{\{x_{i,s}\}_{s=1}^{d_i}|x_i}(\|\mu_{\hat{x}_i} - \mu_{x_i}\|_H > t^{\frac{1}{2\alpha}})dt$. Then a variable change of $t = (2B_{\tilde{K}}/d_i)(1 + \sqrt{\delta})^{2\alpha}$ implies that

$$E_{\{x_{i,s}\}_{s=1}^{d_i}|x_i}[\|\mu_{\hat{x}_i} - \mu_{x_i}\|_H^{2\alpha}] \leq (2 + \sqrt{\pi}) \frac{2^\alpha B_{\tilde{K}}^\alpha}{d_i^\alpha}.$$

Then we know that

$$\begin{aligned} & \left\{ E_{\mathbf{x}^d, |D|, |\mathbf{z}|^{|D|}} \left[\|\hat{S}_D^T y_D - S_D y_D\|_K^2 \right] \right\}^{\frac{1}{2}} = \left\{ E_{\mathbf{x}^d, |D|, |\mathbf{z}|^{|D|}} \left[\left\| \frac{1}{|D|} \sum_{i=1}^{|D|} y_i (K_{\mu_{\hat{x}_i}} - K_{\mu_{x_i}}) \right\|_K^2 \right] \right\}^{\frac{1}{2}} \\ & \leq \left\{ E_{\mathbf{x}^d, |D|, |\mathbf{z}|^{|D|}} \left[\frac{1}{|D|} \sum_{i=1}^{|D|} |y_i| \|K_{\mu_{\hat{x}_i}} - K_{\mu_{x_i}}\|_{\mathcal{L}(Y, \mathcal{H}_K)}^2 \right] \right\}^{\frac{1}{2}} \leq M \left\{ E_{\mathbf{x}^d, |D|, |\mathbf{z}|^{|D|}} \left[\frac{1}{|D|} \sum_{i=1}^{|D|} L^2 \|\mu_{\hat{x}_i} - \mu_{x_i}\|_H^{2\alpha} \right] \right\}^{\frac{1}{2}} \\ & \leq ML \frac{1}{|D|} \sum_{i=1}^{|D|} \left\{ E_{\mathbf{x}^d, |D|, |\mathbf{z}|^{|D|}} \left[\|\mu_{\hat{x}_i} - \mu_{x_i}\|_H^{2\alpha} \right] \right\}^{\frac{1}{2}} \leq (2 + \sqrt{\pi})^{\frac{1}{2}} ML \frac{1}{|D|} \sum_{i=1}^{|D|} \frac{2^{\frac{\alpha}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{d_i^{\frac{\alpha}{2}}}, \end{aligned} \quad (4.5)$$

in which the first inequality follows from the convex inequality of norm $\|\cdot\|_K$ and the fact that $\|yK_{\mu_x}\|_K \leq \|y\|_Y \|K_{\mu_x}\|_{\mathcal{L}(Y, \mathcal{H}_K)}$ for any $y \in Y$, $\mu_x \in X_\mu$. Specifically, $\|\cdot\|_Y = |\cdot|$ when $Y = \mathbb{R}$ in our setting. The second inequality follows from the Hölder assumption of mapping $K_{(\cdot)}$. Recalling the structure of $L_{K, \hat{D}}$ defined in (4.4) and boundedness condition of kernel K in (2.1), using Hölder condition again to $K_{(\cdot)}$, we know

$$\left\{ E_{\mathbf{x}^d, |D|, |\mathbf{z}|^{|D|}} \left[\|L_{K, \hat{D}} - L_{K, D}\|^2 \right] \right\}^{\frac{1}{2}} \leq \kappa L (2 + \sqrt{\pi})^{\frac{1}{2}} \frac{1}{|D|} \sum_{i=1}^{|D|} \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{d_i^{\frac{\alpha}{2}}}. \quad (4.6)$$

In subsequent analysis, we need the following lemma ([7]) of one-stage learning theory to handle the terms involving the information of additional penalty.

Lemma 1. *If the regularization parameters $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2c_V \lambda_2 = \lambda_1^{\max\{2r, 1\}}$ for $r \in (0, 1]$, then the following norm bound holds almost surely:*

$$\left\| (\lambda_1 I + L_K)(\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{-1} \right\| \leq 2. \quad (4.7)$$

In following proofs, we denote

$$\mathcal{A}_{D, \lambda_1, \lambda_2, V_D} = \|(\lambda_1 I + L_K + \lambda_2 V_D^T V_D)(\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1}\|.$$

4.1 Analysis of the error bounds and rates when $r \in [1/2, 1]$

Now we come to the proof of Theorem 1 and Theorem 2, we begin with the following error estimate for $\|f_{D, \lambda_1, \lambda_2} - f_\rho\|_\rho$.

Proposition 1. Assume that $|y| \leq M$ and (2.7) hold almost surely. Let the regularity condition (2.4) hold with some index r satisfying $1/2 \leq r \leq 1$. If $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2c_V \lambda_2 = \lambda_1^{2r}$. Then there holds almost surely

$$\begin{aligned} \|f_{D, \lambda_1, \lambda_2} - f_\rho\|_\rho &\leq 2\mathcal{A}_{D, \lambda_1, \lambda_2, V_D} \|(\lambda_1 I + L_K)^{-\frac{1}{2}} (S_D^T y_D - L_{K, D} f_\rho)\|_K \\ &\quad + (\lambda_1^r + \lambda_1^{r-1} \lambda_2 c_V) 2^r \mathcal{A}_{D, \lambda_1, \lambda_2, V_D}^r \|g_\rho\|_\rho. \end{aligned}$$

Proof. Using the fact that for any $h \in L_{\rho_{X_\mu}}^2$, $\|h\|_\rho = \|L_K^{1/2} h\|_K$, we split $\|f_{D, \lambda_1, \lambda_2} - f_\rho\|_\rho$ into three parts as

$$\begin{aligned} &\|f_{D, \lambda_1, \lambda_2} - f_\rho\|_\rho \\ &= \left\| L_K^{1/2} \left\{ (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1} S_D^T y_D - (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1} (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D) f_\rho \right\} \right\|_K \\ &= \left\| L_K^{1/2} \left\{ (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1} (S_D^T y_D - L_{K, D} f_\rho - \lambda_1 f_\rho - \lambda_2 V_D^T V_D f_\rho) \right\} \right\|_K \\ &\leq \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3, \end{aligned}$$

where

$$\begin{aligned} \mathcal{T}_1 &= \|L_K^{1/2} (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1} (S_D^T y_D - L_{K, D} f_\rho)\|_K, \\ \mathcal{T}_2 &= \lambda_1 \|L_K^{1/2} (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1} L_K^r g_\rho\|_K, \\ \mathcal{T}_3 &= \|L_K^{1/2} (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1} \lambda_2 V_D^T V_D f_\rho\|_K. \end{aligned}$$

We estimate $\mathcal{T}_1$, $\mathcal{T}_2$, and $\mathcal{T}_3$ respectively, as follows, we have almost surely,

$$\begin{aligned} \mathcal{T}_1 &= \|L_K^{1/2} (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1} (S_D^T y_D - L_{K, D} f_\rho)\|_K \\ &\leq \|L_K^{\frac{1}{2}} (L_K + \lambda_1 I)^{-\frac{1}{2}}\| \| (L_K + \lambda_1 I)^{\frac{1}{2}} (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \| \| (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \\ &\quad (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \| \| (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \| \\ &\quad \| (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (\lambda_1 I + L_K)^{\frac{1}{2}} \| \| (\lambda_1 I + L_K)^{-\frac{1}{2}} (S_D^T y_D - L_{K, D} f_\rho) \|_K \\ &\leq 2\mathcal{A}_{D, \lambda_1, \lambda_2, V_D} \|(\lambda_1 I + L_K)^{-\frac{1}{2}} (S_D^T y_D - L_{K, D} f_\rho)\|_K, \end{aligned}$$

in which we have used $\|L_K^{1/2} (L_K + \lambda_1 I)^{-1/2}\| \leq 1$, Lemma 1, and the fact that, for any positive self-adjoint operators T_1 and T_2 on a Hilbert space and $s \in [0, 1]$, $\|T_1^s T_2^s\| \leq \|T_1 T_2\|^s$. For $\mathcal{T}_2$, we have

$$\begin{aligned} \mathcal{T}_2 &= \lambda_1 \|L_K^{1/2} (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-1} L_K^r g_\rho\|_K \\ &\leq \lambda_1 \|L_K^{\frac{1}{2}} (L_K + \lambda_1 I)^{-\frac{1}{2}}\| \| (L_K + \lambda_1 I)^{\frac{1}{2}} (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \| \| (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \\ &\quad (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \| \| (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{r-\frac{1}{2}} \| \\ &\quad \| (\lambda_1 I + L_{K, D} + \lambda_2 V_D^T V_D)^{-r+\frac{1}{2}} (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{r-\frac{1}{2}} \| \\ &\quad \| (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{-r+\frac{1}{2}} (\lambda_1 I + L_K)^{r-\frac{1}{2}} \| \\ &\quad \| (\lambda_1 I + L_K)^{-r+\frac{1}{2}} L_K^{r-\frac{1}{2}} \| \| L_K^{1/2} g_\rho \|_K \\ &\leq \lambda_1^r \mathcal{A}_{D, \lambda_1, \lambda_2, V_D}^r \cdot 2^r \|g_\rho\|_\rho, \end{aligned}$$

where we have used the fact that

$$\begin{aligned} & \|(\lambda_1 I + L_{K,D} + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (\lambda_1 I + L_{K,D} + \lambda_2 V_D^T V_D)^{r-\frac{1}{2}}\| \\ &= \|(\lambda_1 I + L_{K,D} + \lambda_2 V_D^T V_D)^{r-1}\| \leq (\lambda_1 + \lambda_D^2 + \lambda_2 \lambda_{V_D}^2)^{r-1} \leq \lambda_1^{r-1} \end{aligned}$$

in which

$$\lambda_D = \inf_{f \in \mathcal{H}_K} \frac{\|S_D f\|_{l^2}}{\|f\|_K}, \quad \lambda_{V_D} = \inf_{f \in \mathcal{H}_K} \frac{\|V_D f\|_K}{\|f\|_K}.$$

This fact can be treated as a direct result of Proposition 3.1 in [1]. Note that V_D is a bounded operator on $\mathcal{H}_K$, then following a similar procedure with the estimate on $\mathcal{T}_2$, we obtain

$$\mathcal{T}_3 \leq \lambda_1^{r-1} \lambda_2 c_V 2^r \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^r \|g_\rho\|_\rho.$$

Combining the three estimates, the desired bound follows. ■

The following lemma is used to prove the main results. It is based on techniques of Neumann expansion [7] and the second-order decomposition [12] for invertible operators on Banach space, namely, for any invertible operators A and B ,

$$A^{-1} - B^{-1} = B^{-1}(B - A)A^{-1}(B - A)B^{-1} + B^{-1}(B - A)B^{-1} \quad (4.8)$$

$$= B^{-1}(B - A)B^{-1}(B - A)A^{-1} + B^{-1}(B - A)B^{-1}. \quad (4.9)$$

Lemma 2. *Let the sample set D be drawn independently according to the probability measure ρ . If $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2c_V \lambda_2 = \lambda_1^{\max\{2r, 1\}}$ for $r \in (0, 1]$. Then we have,*

$$E_{\mathbf{z}^{[D]}} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^s \right] \leq (2\Gamma(2s+1) + \log^{2s} 2) \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2s}, \quad s \geq 0; \quad (4.10)$$

$$E_{\mathbf{z}^{[D]}} \left[\left\| (\lambda_1 I + L_K)^{-\frac{1}{2}} (S_D^T y_D - L_{K,D} f_\rho) \right\|_K^s \right] \leq (2\Gamma(s+1) + \log^s 2) \left(\frac{2M}{\kappa} \mathcal{B}_{|D|,\lambda_1} \right)^s, \quad s \geq 1. \quad (4.11)$$

Proof. From Proposition 4.3 in [7], we know, for any $\delta \in (0, 1)$, there holds

$$\text{Prob} \left\{ \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \leq \left(\frac{2\mathcal{B}_{|D|,\lambda_1} \log \frac{2}{\delta}}{\sqrt{\lambda_1}} + 1 \right)^2 \right\} \geq 1 - \delta. \quad (4.12)$$

Then it is easy to see, for any $\delta \in (0, 2)$,

$$\text{Prob} \left\{ \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \leq \left(\frac{2\mathcal{B}_{|D|,\lambda_1} \log \frac{4}{\delta}}{\sqrt{\lambda_1}} + 1 \right)^2 \right\} \geq 1 - \frac{\delta}{2}. \quad (4.13)$$

Denote the random variable $\xi = \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^s$, perform the variable substitution $t = \left(\frac{2\mathcal{B}_{|D|,\lambda_1} \log \frac{2}{\delta}}{\sqrt{\lambda_1}} + 1 \right)^{2s} \log^{2s} \frac{4}{\delta}$ with $s \geq 0$, we know, for $t > \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2s} \log^{2s} 2$,

$$\text{Prob} \left(\xi > t \right) = \text{Prob} \left(\xi^{\frac{1}{s}} > t^{\frac{1}{s}} \right) \leq \frac{\delta}{2} = 2 \exp \left\{ - \frac{t^{\frac{1}{2s}}}{\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1} \right\}. \quad (4.14)$$

Then by using the formula $E_{\mathbf{z}^{[D]}}[\xi] = \int_0^\infty \text{Prob}(\xi > t) dt$, it follows that

$$E_{\mathbf{z}^{[D]}}[\xi] \leq \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2s} \log^{2s} 2 + \int_{\left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2s} \log^{2s} 2}^\infty 2 \exp \left\{ - \frac{t^{\frac{1}{2s}}}{\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1} \right\} dt$$

$$\begin{aligned}
&\leq \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2s} \log^{2s} 2 + 4s \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2s} \int_{\log 2}^{\infty} e^{-x} x^{2s-1} dx \\
&\leq (2\Gamma(2s+1) + \log^{2s} 2) \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2s}.
\end{aligned}$$

For the second inequality, note that [4] implies that, for any $0 < \delta < 1$, with probability of at least $1 - \delta$,

$$\|(\lambda_1 I + L_K)^{-\frac{1}{2}}(S_D^T y_D - L_{K,D} f_\rho)\|_K \leq \frac{2M}{\kappa} \mathcal{B}_{|D|,\lambda_1} \log \frac{2}{\delta}. \quad (4.15)$$

Then the same procedure of proving the first inequality implies the result. ■

Proposition 2. Assume that $|y| \leq M$ and (2.7) hold almost surely. Assume the regularity condition (2.4) holds with some index $1/2 \leq r \leq 1$. If $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2c_V \lambda_2 = \lambda_1^{2r}$. Then we have,

$$\begin{aligned}
E_{\mathbf{z}|D|} \left[\|f_{D,\lambda_1,\lambda_2} - f_\rho\|_\rho \right] &\leq 4(2\Gamma(3) + \log^2 2)^{\frac{1}{2}} (2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \frac{M}{\kappa} \mathcal{B}_{|D|,\lambda_1} \\
&\quad + (2\Gamma(2r+1) + \log^{2r} 2) 2^r (\lambda_1^r + \lambda_1^{r-1} \lambda_2 c_V) \|g_\rho\|_\rho \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2r}. \quad (4.16)
\end{aligned}$$

Proof. Starting from Proposition 1, taking expectations on both sides and using Schwarz inequality, we have

$$\begin{aligned}
E_{\mathbf{z}|D|} \left[\|f_{D,\lambda_1,\lambda_2} - f_\rho\|_\rho \right] &\leq 2E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \|(\lambda_1 I + L_K)^{-\frac{1}{2}}(S_D^T y_D - L_{K,D} f_\rho)\|_K \right] \\
&\quad + (\lambda_1^r + \lambda_1^{r-1} \lambda_2 c_V) 2^r \|g_\rho\|_\rho E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^r \right]. \\
&\leq 2 \left\{ E_{\mathbf{z}|D|} \left[\|(\lambda_1 I + L_K)^{-\frac{1}{2}}(S_D^T y_D - L_{K,D} f_\rho)\|_K^2 \right] \right\}^{1/2} \left\{ E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^2 \right] \right\}^{1/2} \\
&\quad + (\lambda_1^r + \lambda_1^{r-1} \lambda_2 c_V) 2^r \|g_\rho\|_\rho E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^r \right]. \\
&\leq 2(2\Gamma(3) + \log^2 2)^{1/2} \frac{2M}{\kappa} \mathcal{B}_{|D|,\lambda_1} (2\Gamma(5) + \log^4 2)^{1/2} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \\
&\quad + (\lambda_1^r + \lambda_1^{r-1} \lambda_2 c_V) 2^r \|g_\rho\|_\rho (2\Gamma(2r+1) + \log^{2r} 2) \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2r},
\end{aligned}$$

in which the last inequality follows from (4.11) with index $s = 2$ and (4.11) with index $s = r$ in Lemma 2. ■

Denote

$$\Omega_{D,\lambda_1,\lambda_2,V_D} = \|L_K^{1/2}(L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}\|. \quad (4.17)$$

Now we start to bound $\|f_{\widehat{D},\lambda_1,\lambda_2} - f_{D,\lambda_1,\lambda_2}\|_\rho$ part in error decomposition of $\|f_{\widehat{D},\lambda_1,\lambda_2} - f_\rho\|_\rho$.

Proposition 3. There holds almost surely

$$\|f_{\widehat{D},\lambda_1,\lambda_2} - f_{D,\lambda_1,\lambda_2}\|_\rho \leq \Omega_{D,\lambda_1,\lambda_2,V_D} \left(\|\hat{S}_D^T y_D - S_D^T y_D\|_K + \|L_{K,D} - L_{K,\widehat{D}}\| \|f_{D,\lambda_1,\lambda_2}\|_K \right). \quad (4.18)$$

Proof. Recalling the representations of $f_{\widehat{D},\lambda_1,\lambda_2}$ in (1.3) and $f_{D,\lambda_1,\lambda_2}$ in (4.3), we have

$$\begin{aligned}
& f_{\widehat{D},\lambda_1,\lambda_2} - f_{D,\lambda_1,\lambda_2} \\
&= (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \hat{S}_D^T y_D - (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} S_D^T y_D \\
&= (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (\hat{S}_D^T y_D - S_D^T y_D) \\
&\quad + \left[(L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} - (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right] S_D^T y_D \\
&= (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (\hat{S}_D^T y_D - S_D^T y_D) \\
&\quad + (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_{K,D} - L_{K,\widehat{D}}) f_{D,\lambda_1,\lambda_2},
\end{aligned}$$

in which we have used the fact that, for any invertible operator A and B , $A^{-1} - B^{-1} = A^{-1}(B - A)B^{-1}$, with $A = L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D$ and $B = L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D$. Take $L_{\rho_{X_\mu}}^2$ -norm on both sides of the above equality and use the triangle formula, it follows that

$$\begin{aligned}
\|f_{\widehat{D},\lambda_1,\lambda_2} - f_{D,\lambda_1,\lambda_2}\|_\rho &\leq \|L_K^{1/2} (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (\hat{S}_D^T y_D - S_D^T y_D)\|_K \\
&\quad + \|L_K^{1/2} (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_{K,D} - L_{K,\widehat{D}}) f_{D,\lambda_1,\lambda_2}\|_K \\
&\leq \|L_K^{1/2} (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}\| \|\hat{S}_D^T y_D - S_D^T y_D\|_K \\
&\quad + \|L_K^{1/2} (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}\| \|L_{K,D} - L_{K,\widehat{D}}\| \|f_{D,\lambda_1,\lambda_2}\|_K \\
&= \Omega_{D,\lambda_1,\lambda_2,V_D} \left(\|\hat{S}_D^T y_D - S_D^T y_D\|_K + \|L_{K,D} - L_{K,\widehat{D}}\| \|f_{D,\lambda_1,\lambda_2}\|_K \right). \blacksquare
\end{aligned}$$

The following result provides an estimate for $\Omega_{D,\lambda_1,\lambda_2,V_D}$.

Lemma 3. Let D be a sample drawn independently according to measure ρ , and $\{x_{i,s}\}_{s=1}^{d_i}$ be a sample independently drawn according to distribution x_i , $i = 1, 2, \dots, |D|$. Let the regularity condition (2.4) hold with some index $r \in (0, 1]$ and the mapping $K_{(\cdot)} : X_\mu \rightarrow \mathcal{L}(Y, \mathcal{H}_K)$ be (α, L) -Hölder continuous with $\alpha \in (0, 1]$ and $L > 0$. If $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2c_V \lambda_2 = \lambda_1^{\max\{2r, 1\}}$, then we have

$$\begin{aligned}
\left\{ E_{\mathbf{x}^d, |D|} [\Omega_{D,\lambda_1,\lambda_2,V_D}^2] \right\}^{\frac{1}{2}} &\leq \left(\frac{2\sqrt{2}}{\lambda_1^{3/2}} (2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{1}{|D|} \sum_{i=1}^{|D|} \frac{2^{\frac{\alpha+2}{2}} B_{\frac{2}{K}}^{\frac{\alpha}{2}}}{d_i^{\frac{\alpha}{2}}} \right) \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \\
&\quad + \frac{2}{\lambda_1^{\frac{1}{2}}} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{1/2}.
\end{aligned}$$

Proof. Divide $\Omega_{D,\lambda_1,\lambda_2,V_D}$ into two parts and use triangle inequality of operator norm as follows,

$$\begin{aligned}
\Omega_{D,\lambda_1,\lambda_2,V_D} &\leq \left\| L_K^{1/2} \left\{ (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} - (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right\} \right\| \\
&\quad + \left\| L_K^{1/2} (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right\|.
\end{aligned}$$

The first term is estimated as follows

$$\left\| L_K^{1/2} \left\{ (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} - (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right\} \right\|$$

$$\begin{aligned}
&= \left\| L_K^{1/2} (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_{K,D} - L_{K,\widehat{D}}) (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right\| \\
&\leq \left\| L_K^{1/2} (L_K + \lambda_1 I)^{-\frac{1}{2}} \right\| \left\| (\lambda_1 I + L_K)^{\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \left\| (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \right. \\
&\quad \left. (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \left\| (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \right\| \\
&\quad \left\| (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_K + \lambda_1 I)^{\frac{1}{2}} \right\| \left\| (L_K + \lambda_1 I)^{-\frac{1}{2}} \right\| \left\| L_{K,D} - L_{K,\widehat{D}} \right\| \\
&\quad \left\| (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right\| \\
&\leq \frac{2}{\lambda_1^{3/2}} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \left\| L_{K,D} - L_{K,\widehat{D}} \right\|,
\end{aligned}$$

where we have used

$$\begin{aligned}
&\left\| L_K^{1/2} (L_K + \lambda_1 I)^{-1/2} \right\| \leq 1, \\
&\left\| (L_K + \lambda_1 I)^{-1/2} \right\| \leq \frac{1}{\sqrt{\lambda_1}}, \\
&\left\| (L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right\| \leq \frac{1}{\lambda_1},
\end{aligned}$$

and the fact that for two self-adjoint operators T_1, T_2 , there holds $\|T_1 T_2\| = \|T_2 T_1\|$. The second term is estimated as follows,

$$\begin{aligned}
&\left\| L_K^{1/2} (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right\| \\
&\leq \left\| L_K^{1/2} (\lambda_1 I + L_K)^{-\frac{1}{2}} \right\| \left\| (\lambda_1 I + L_K)^{\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \left\| (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \right. \\
&\quad \left. (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \left\| (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \\
&\leq \frac{\sqrt{2}}{\sqrt{\lambda_1}} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{1/2},
\end{aligned}$$

in which we have used the fact

$$\left\| (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \leq \frac{1}{\sqrt{\lambda_1}}$$

which follows from the analysis above. Then we know from the above analysis that

$$\Omega_{D,\lambda_1,\lambda_2,V_D} \leq \frac{2}{\lambda_1^{3/2}} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \left\| L_{K,D} - L_{K,\widehat{D}} \right\| + \frac{\sqrt{2}}{\sqrt{\lambda_1}} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{1/2}.$$

This directly implies

$$\Omega_{D,\lambda_1,\lambda_2,V_D}^2 \leq \frac{8}{\lambda_1^3} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^2 \left\| L_{K,D} - L_{K,\widehat{D}} \right\|^2 + \frac{4}{\lambda_1} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}.$$

Taking expectations on both sides and using (4.6), we obtain that

$$\begin{aligned}
&\left\{ E_{\mathbf{x}^d, |D|} \left[\Omega_{D,\lambda_1,\lambda_2,V_D}^2 \right] \right\} \leq \frac{8}{\lambda_1^3} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^2 E_{\mathbf{x}^d, |D|} \left[\left\| L_{K,D} - L_{K,\widehat{D}} \right\|^2 \right] + \frac{4}{\lambda_1} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \\
&\leq \left(\frac{8}{\lambda_1^3} \kappa^2 (2 + \sqrt{\pi}) L^2 \frac{1}{|D|^2} \sum_{i=1}^{|D|} \frac{2^{\alpha+2} B_{\widetilde{K}}^\alpha}{d_i^\alpha} \right) \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^2
\end{aligned}$$

$$+\frac{4}{\lambda_1}\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}.$$

Finally, we have

$$\left\{E_{\mathbf{x}^{\mathbf{d}},|\mathbf{D}|,|\mathbf{z}|,|\mathbf{D}|}\left[\Omega_{D,\lambda_1,\lambda_2,V_D}^2\right]\right\}^{\frac{1}{2}}\leq\left(\frac{2\sqrt{2}}{\lambda_1^{3/2}}(2+\sqrt{\pi})^{\frac{1}{2}}\kappa L\frac{1}{|\mathbf{D}|}\sum_{i=1}^{|\mathbf{D}|}\frac{2^{\frac{\alpha+2}{2}}B_{\tilde{K}}^{\frac{\alpha}{2}}}{d_i^{\frac{\alpha}{2}}}\right)\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}+\frac{2}{\lambda_1^{\frac{1}{2}}}\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{1/2}.\blacksquare$$

Before coming to the proof of Theorem 1, the following expected bound for $\|f_{D,\lambda_1,\lambda_2}\|_K$ in (4.18) is needed.

Proposition 4. *Suppose $|y| \leq M$ almost surely. Let the regularity condition (2.4) hold with some index $1/2 \leq r \leq 1$. If $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2c_V\lambda_2 = \lambda_1^{2r}$, then we have*

$$\begin{aligned} \left\{E_{\mathbf{z}|\mathbf{D}}\left[\|f_{D,\lambda_1,\lambda_2}\|_K^2\right]\right\}^{1/2} &\leq 2\sqrt{6}(2\Gamma(5) + \log^4 2)^{1/2}\frac{M}{\kappa}\frac{\mathcal{B}_{|\mathbf{D}|,\lambda_1}}{\sqrt{\lambda_1}}\left(\frac{2\mathcal{B}_{|\mathbf{D}|,\lambda_1}}{\sqrt{\lambda_1}} + 1\right) \\ &\quad + \sqrt{3}(\lambda_1 + \lambda_2 c_V)\lambda_1^{r-\frac{3}{2}}2^{r-\frac{1}{2}}\|g_\rho\|_\rho(2\Gamma(4r-1) + \log^{4r-2} 2)^{1/2}\left(\frac{2\mathcal{B}_{|\mathbf{D}|,\lambda_1}}{\sqrt{\lambda_1}} + 1\right)^{2r-1} + \sqrt{3}\kappa^{r-\frac{1}{2}}\|g_\rho\|_\rho \end{aligned}$$

where $\mathcal{B}_{|\mathbf{D}|,\lambda_1}$ is defined as in (2.8).

Proof. Dividing $f_{D,\lambda_1,\lambda_2}$ into two parts and using the triangle formula for RKHS norm, we have

$$\begin{aligned} \|f_{D,\lambda_1,\lambda_2}\|_K &\leq \|f_{D,\lambda_1,\lambda_2} - f_\rho\|_K + \|f_\rho\|_K \\ &= \|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}(S_D^T y_D - L_{K,D} f_\rho - \lambda_1 f_\rho - \lambda_2 V_D^T V_D f_\rho)\|_K + \|f_\rho\|_K \\ &\leq \|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}(S_D^T y_D - L_{K,D} f_\rho)\|_K \\ &\quad + (\lambda_1 + \lambda_2 c_V)\|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}L_K^r g_\rho\|_K + \|f_\rho\|_K. \end{aligned}$$

The first term is estimated as follows,

$$\begin{aligned} &\|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}(S_D^T y_D - L_{K,D} f_\rho)\|_K \\ &= \|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}}\|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}}(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{\frac{1}{2}}\| \\ &\quad \|(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}}(L_K + \lambda_1 I)^{\frac{1}{2}}\|(L_K + \lambda_1 I)^{-\frac{1}{2}}(S_D^T y_D - L_{K,D} f_\rho)\|_K \\ &\leq \frac{\sqrt{2}}{\sqrt{\lambda_1}}\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{1/2}\|(L_K + \lambda_1 I)^{-\frac{1}{2}}(S_D^T y_D - L_{K,D} f_\rho)\|_K. \end{aligned}$$

Recall that $1/2 \leq r \leq 1$, then the second term is estimated as follows,

$$\begin{aligned} &(\lambda_1 + \lambda_2 c_V)\|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}L_K^r g_\rho\|_K \\ &\leq (\lambda_1 + \lambda_2 c_V)\|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{r-\frac{3}{2}}\|(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-r+\frac{1}{2}}(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{r-\frac{1}{2}}\| \\ &\quad \|(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-r+\frac{1}{2}}(L_K + \lambda_1 I)^{r-\frac{1}{2}}\|(L_K + \lambda_1 I)^{-r+\frac{1}{2}}L_K^{r-\frac{1}{2}}\|L_K^{1/2}g_\rho\|_K \\ &\leq (\lambda_1 + \lambda_2 c_V)\lambda_1^{r-\frac{3}{2}}2^{r-\frac{1}{2}}\|g_\rho\|_\rho\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{r-\frac{1}{2}}. \end{aligned}$$

Hence

$$\|f_{D,\lambda_1,\lambda_2}\|_K \leq \frac{\sqrt{2}}{\sqrt{\lambda_1}}\|(L_K + \lambda_1 I)^{-\frac{1}{2}}(S_D^T y_D - L_{K,D} f_\rho)\|_K\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{1/2}$$

$$+(\lambda_1 + \lambda_2 c_V) \lambda_1^{r-\frac{3}{2}} 2^{r-\frac{1}{2}} \|g_\rho\|_\rho \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{r-\frac{1}{2}} + \kappa^{r-\frac{1}{2}} \|g_\rho\|_\rho.$$

After taking expectations, Jensen's inequality implies

$$\begin{aligned} E_{\mathbf{z}|D|} \left[\|f_{D,\lambda_1,\lambda_2}\|_K^2 \right] &\leq \frac{6}{\lambda_1} E_{\mathbf{z}|D|} \left[\|(L_K + \lambda_1 I)^{-\frac{1}{2}} (S_D^T y_D - L_{K,D} f_\rho)\|_K^2 \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \right] \\ &\quad + 3(\lambda_1 + \lambda_2 c_V)^2 \lambda_1^{2r-3} 2^{2r-1} \|g_\rho\|_\rho^2 E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{2r-1} \right] + 3\kappa^{2r-1} \|g_\rho\|_\rho^2 \\ &\leq \frac{6}{\lambda_1} \left\{ E_{\mathbf{z}|D|} \left[\|(L_K + \lambda_1 I)^{-\frac{1}{2}} (S_D^T y_D - L_{K,D} f_\rho)\|_K^4 \right] \right\}^{\frac{1}{2}} \left\{ E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^2 \right] \right\}^{\frac{1}{2}} \\ &\quad + 3(\lambda_1 + \lambda_2 c_V)^2 \lambda_1^{2r-3} 2^{2r-1} \|g_\rho\|_\rho^2 E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{2r-1} \right] + 3\kappa^{2r-1} \|g_\rho\|_\rho^2 \\ &\leq 24(2\Gamma(5) + \log^4 2) \frac{M^2}{\kappa^2} \frac{\mathcal{B}_{|D|,\lambda_1}^2}{\lambda_1} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \\ &\quad + 3(\lambda_1 + \lambda_2 c_V)^2 \lambda_1^{2r-3} 2^{2r-1} \|g_\rho\|_\rho^2 (2\Gamma(4r-1) + \log^{4r-2} 2) \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{4r-2} + 3\kappa^{2r-1} \|g_\rho\|_\rho^2, \end{aligned}$$

in which the second inequality follows from the Schwarz inequality, and the third inequality follows from Lemma 2 with $s = 2$ and $2r - 1$ in (4.11) and $s = 4$ in (4.11). The desired result is obtained after noting the fact $\sqrt{a_1 + a_2 + a_3} \leq \sqrt{a_1} + \sqrt{a_2} + \sqrt{a_3}$ for $a_1, a_2, a_3 \geq 0$. ■

Proof of Theorem 1. Return to Proposition 3, after taking expectations on both sides, and using Schwarz inequality, inequalities (4.5), (4.6) and Lemma 3, we have

$$\begin{aligned} &E \left[\Omega_{D,\lambda_1,\lambda_2,V_D} \|\hat{S}_D^T y_D - S_D^T y_D\| \right] \\ &\leq E_{\mathbf{z}|D|} \left[\left\{ E_{\mathbf{x}^{\mathbf{d}},|D||\mathbf{z}|D|} [\Omega_{D,\lambda_1,\lambda_2,V_D}^2] \right\}^{\frac{1}{2}} \left\{ E_{\mathbf{x}^{\mathbf{d}},|D||\mathbf{z}|D|} [\|\hat{S}_D^T y_D - S_D^T y_D\|^2] \right\}^{\frac{1}{2}} \right] \\ &\leq \left(\frac{2\sqrt{2}}{\lambda_1^{3/2}} (2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} + \frac{2}{\lambda_1^{\frac{1}{2}}} \right) (2 + \sqrt{\pi})^{\frac{1}{2}} L M \frac{2^{\frac{\alpha}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} (2\Gamma(3) + \log^2 2) \left[\left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \right. \\ &\quad \left. + (2\Gamma(2) + \log 2) \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right) \right] \\ &\leq 2 \left(\frac{2\sqrt{2}}{\lambda_1^{3/2}} (2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} + \frac{2}{\lambda_1^{\frac{1}{2}}} \right) (2 + \sqrt{\pi})^{\frac{1}{2}} L M \frac{2^{\frac{\alpha}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} (2\Gamma(3) + \log^2 2) \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \end{aligned} \quad (41)$$

and

$$\begin{aligned} &E \left[\Omega_{D,\lambda_1,\lambda_2,V_D} \|L_{K,D} - L_{K,\hat{D}}\| \|f_{D,\lambda_1,\lambda_2}\|_K \right] \\ &\leq E_{\mathbf{z}|D|} \left[E_{\mathbf{x}^{\mathbf{d}},|D||\mathbf{z}|D|} [\Omega_{D,\lambda_1,\lambda_2,V_D} \|L_{K,D} - L_{K,\hat{D}}\|] \cdot \|f_{D,\lambda_1,\lambda_2}\|_K \right] \\ &\leq E_{\mathbf{z}|D|} \left[\left\{ E_{\mathbf{x}^{\mathbf{d}},|D||\mathbf{z}|D|} [\Omega_{D,\lambda_1,\lambda_2,V_D}^2] \right\}^{\frac{1}{2}} \left\{ E_{\mathbf{x}^{\mathbf{d}},|D||\mathbf{z}|D|} [\|L_{K,D} - L_{K,\hat{D}}\|^2] \right\}^{\frac{1}{2}} \|f_{D,\lambda_1,\lambda_2}\|_K \right] \\ &\leq \left(\frac{2\sqrt{2}}{\lambda_1^{3/2}} (2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} + \frac{2}{\lambda_1^{\frac{1}{2}}} \right) \kappa L (2 + \sqrt{\pi})^{\frac{1}{2}} \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} E_{\mathbf{z}|D|} \left[\|f_{D,\lambda_1,\lambda_2}\|_K (\mathcal{A}_{D,\lambda_1,\lambda_2,V_D} + \mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^{1/2}) \right] \end{aligned}$$

$$\begin{aligned}
&\leq \left(\frac{2\sqrt{2}}{\lambda_1^{3/2}} (2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} + \frac{2}{\lambda_1^{\frac{1}{2}}} \right) \kappa L (2 + \sqrt{\pi})^{\frac{1}{2}} \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} \left\{ E_{\mathbf{z}|D|} \left[\|f_{D,\lambda_1,\lambda_2}\|_K^2 \right] \right\}^{\frac{1}{2}} \times \\
&\quad \left(\left\{ E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^2 \right] \right\}^{\frac{1}{2}} + \left\{ E_{\mathbf{z}|D|} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \right] \right\}^{\frac{1}{2}} \right) \\
&\leq 2 \left(\frac{2\sqrt{2}}{\lambda_1^{3/2}} (2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} + \frac{2}{\lambda_1^{\frac{1}{2}}} \right) \kappa L (2 + \sqrt{\pi})^{\frac{1}{2}} \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\tilde{d}^{\frac{\alpha}{2}}} \left[2\sqrt{6}(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \frac{M}{\kappa} \times \right. \\
&\quad \left. \frac{\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right) + \sqrt{3}(\lambda_1 + \lambda_2 c_V) \lambda_1^{r-\frac{3}{2}} 2^{r-\frac{1}{2}} \|g_\rho\|_\rho (2\Gamma(4r-1) + \log^{4r-2} 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2r-1} \right. \\
&\quad \left. + \sqrt{3} \kappa^{r-\frac{1}{2}} \|g_\rho\|_\rho \right] (2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2. \tag{4.20}
\end{aligned}$$

Combining (4.19) and (4.20) with Proposition 3 and taking expectations on both sides of (4.18), we have

$$\begin{aligned}
E \left[\|f_{\hat{D},\lambda_1,\lambda_2} - f_{D,\lambda_1,\lambda_2}\|_\rho \right] &\leq 2 \left(2\sqrt{2}(2 + \sqrt{2})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\lambda_1 \tilde{d}^{\frac{\alpha}{2}}} + 2 \right) \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \frac{2^{\frac{\alpha}{2}} B_{\tilde{K}}^{\frac{\alpha}{2}}}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} \times \\
&\quad \left[(2 + \sqrt{\pi})^{\frac{1}{2}} LM(2\Gamma(3) + \log^2 2) + 2(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \left(2\sqrt{6}(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \frac{M}{\kappa} \frac{\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right) \right. \right. \\
&\quad \left. \left. + \sqrt{3}(\lambda_1 + \lambda_2 c_V) \lambda_1^{r-\frac{3}{2}} 2^{r-\frac{1}{2}} \|g_\rho\|_\rho (2\Gamma(4r-1) + \log^{4r-2} 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^{2r-1} + \sqrt{3} \kappa^{r-\frac{1}{2}} \|g_\rho\|_\rho \right) \right] \tag{4.21}
\end{aligned}$$

Also, we have already obtained (4.16), noting that $\|f_{\hat{D},\lambda_1,\lambda_2} - f_\rho\|_\rho \leq \|f_{\hat{D},\lambda_1,\lambda_2} - f_{D,\lambda_1,\lambda_2}\|_\rho + \|f_{D,\lambda_1,\lambda_2} - f_\rho\|_\rho$, we complete the proof. ■

Proof of Theorem 2. Considering the selection of

$$\lambda_1 = |D|^{-\frac{1}{2r+\beta}}, \lambda_2 = \frac{1}{2c_V} |D|^{-\frac{2r}{2r+\beta}}, d_1 = d_2 = \dots = d_{|D|} = |D|^{\frac{1+2r}{\alpha(2r+\beta)}},$$

and the capacity condition $\mathcal{N}(\lambda_1) \leq \mathcal{C}_0 \lambda_1^{-\beta}$, $\beta \in (0, 1]$, we can obtain the following bounds hold,

$$\frac{1}{|D|} \sum_{i=1}^{|D|} \frac{1}{\lambda_1^{\frac{1}{2}} d_i^{\frac{\alpha}{2}}} = |D|^{-\frac{\alpha}{2} \frac{1+2r}{\alpha(2r+\beta)}} |D|^{\frac{1}{2(2r+\beta)}} = |D|^{-\frac{r}{2r+\beta}}, \tag{4.22}$$

$$\frac{1}{|D|} \sum_{i=1}^{|D|} \frac{1}{\lambda_1 d_i^{\frac{\alpha}{2}}} = |D|^{\frac{1}{2r+\beta}} |D|^{-\frac{1+2r}{2(2r+\beta)}} = |D|^{\frac{2-1-2r}{2(2r+\beta)}} = |D|^{\frac{1-2r}{2(2r+\beta)}} \leq 1 \quad (r \in [1/2, 1]), \tag{4.23}$$

$$\begin{aligned}
\mathcal{B}_{|D|,\lambda_1} &= \frac{2\kappa}{\sqrt{|D|}} \left(\frac{\kappa}{\sqrt{|D|\lambda_1}} + \sqrt{\mathcal{N}(\lambda_1)} \right) \leq 2\kappa(\kappa + \sqrt{\mathcal{C}_0}) |D|^{-\frac{r}{2r+\beta}}, \\
\frac{\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} &\leq 2\kappa(\kappa + \sqrt{\mathcal{C}_0}), \\
\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 &\leq 4\kappa(\kappa + \sqrt{\mathcal{C}_0}) + 1,
\end{aligned}$$

$$(\lambda_1 + \lambda_2 c_V) \lambda_1^{r-\frac{3}{2}} \leq (\lambda_1 + \frac{\lambda_1^{2r}}{2}) \lambda_1^{r-\frac{3}{2}} \leq (\frac{3}{2}) \lambda_1^{r-\frac{1}{2}} \leq \frac{3}{2},$$

$$\lambda_1^r + \lambda_1^{r-1} \lambda_2 c_V = \lambda_1^r + \frac{1}{2} \lambda_1^{3r-1} \leq \frac{3}{2} \lambda_1^r = \frac{3}{2} |D|^{-\frac{r}{2r+\beta}}.$$

Substitute these bounds into Theorem 1, we arrive at the following bound

$$\begin{aligned} & E \left[\left\| f_{\widehat{D}, \lambda_1, \lambda_2} - f_\rho \right\|_\rho \right] \\ & \leq \left(2\sqrt{2}(2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L 2^{\frac{\alpha+2}{2}} B_{\widetilde{K}}^{\frac{\alpha}{2}} + 2 \right) (2 + \sqrt{\pi})^{\frac{1}{2}} L M 2^{\frac{\alpha}{2}} B_{\widetilde{K}}^{\frac{\alpha}{2}} (2\Gamma(3) + \log^2 2) \times \left(4\kappa(\kappa + \sqrt{\mathcal{C}_0}) + 1 \right)^2 |D|^{-\frac{r}{2r+\beta}} \\ & + \left(2\sqrt{2}(2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L 2^{\frac{\alpha+2}{2}} B_{\widetilde{K}}^{\frac{\alpha}{2}} + 2 \right) (2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L 2^{\frac{\alpha+2}{2}} B_{\widetilde{K}}^{\frac{\alpha}{2}} \left[2\sqrt{6}(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} M(2\kappa(\kappa + \sqrt{\mathcal{C}_0})) \times \right. \\ & \left. \left(4\kappa(\kappa + \sqrt{\mathcal{C}_0}) + 1 \right) + \frac{3\sqrt{3}}{2} 2^{r-\frac{1}{2}} \|g_\rho\|_\rho (2\Gamma(4r-1) + \log^{4r-1} 2)^{\frac{1}{2}} \left(4\kappa(\kappa + \sqrt{\mathcal{C}_0}) + 1 \right)^{2r-1} + \sqrt{3} \kappa^{r-\frac{1}{2}} \|g_\rho\|_\rho \right] \times \\ & (2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \left(4\kappa(\kappa + \sqrt{\mathcal{C}_0}) + 1 \right)^2 |D|^{-\frac{r}{2r+\beta}} \\ & + 4(2\Gamma(3) + \log^2 2)^{\frac{1}{2}} (2\Gamma(5) + \log^4 2) \left(4\kappa(\kappa + \sqrt{\mathcal{C}_0}) + 1 \right)^2 \frac{M}{\kappa} 2\kappa(\kappa + \sqrt{\mathcal{C}_0}) |D|^{-\frac{r}{2r+\beta}} \\ & + (2\Gamma(2r+1) + \log^{2r} 2) 2^r \|g_\rho\|_\rho \left(4\kappa(\kappa + \sqrt{\mathcal{C}_0}) + 1 \right)^{2r} \frac{3}{2} |D|^{-\frac{r}{2r+\beta}}, \end{aligned}$$

which directly implies the desired learning rate and completes the proof. ■

4.2 Analysis of the error bounds and rates when $r \in (0, 1/2)$

Now we turn to prove the result in the nonstandard setting $r \in (0, 1/2)$ that corresponds to $f_\rho \notin \mathcal{H}_K$. In this section, we need the following norms for subsequent estimates

$$\Pi_{D, \lambda_1} = \|(\lambda_1 I + L_K)^{-\frac{1}{2}} (S_D^T y_D - L_K f_\rho)\|_K, \quad (4.24)$$

$$\Xi_{D, \lambda_1} = \|(\lambda_1 I + L_K)^{-\frac{1}{2}} (L_K - L_{K,D})\|. \quad (4.25)$$

For D i.i.d. drawn from ρ , and $|y| \leq M$ almost surely and for any $0 < \delta < 1$, [4] and [12] imply that, with probability at least $1 - \delta$, there holds,

$$\begin{aligned} \Pi_{D, \lambda_1} & \leq 2M(\kappa + 1) \mathcal{B}'_{|D|, \lambda_1} \log \frac{2}{\delta}, \\ \Xi_{D, \lambda_1} & \leq 2(\kappa^2 + \kappa) \mathcal{B}'_{|D|, \lambda_1} \log \frac{2}{\delta}. \end{aligned}$$

Then, same procedures with the proof of Lemma 2 imply the following results.

Lemma 4. *Let the sample set D be drawn independently according to probability measure ρ . If $|y| \leq M$ almost surely, then we have,*

$$\begin{aligned} E_{\mathbf{z}|D} \left[\Pi_{D, \lambda_1}^s \right] & \leq (2\Gamma(s+1) + \log^s 2) \left(2M(\kappa + 1) \mathcal{B}'_{|D|, \lambda_1} \right)^s, s \geq 1; \\ E_{\mathbf{z}|D} \left[\Xi_{D, \lambda_1}^s \right] & \leq (2\Gamma(s+1) + \log^s 2) \left(2(\kappa^2 + \kappa) \mathcal{B}'_{|D|, \lambda_1} \right)^s, s \geq 1, \end{aligned}$$

in which $\mathcal{B}'_{|D|, \lambda_1}$ is defined as in (2.9).

Now we prepare to prove the main results when f_ρ does not lie in $\mathcal{H}_K$. We use the following function

$$f_{\lambda_1} = \arg \min_{f \in \mathcal{H}_K} \{ \|f - f_\rho\|_{L_{\rho X_\mu}}^2 + \lambda_1 \|f\|_K^2 \} \quad (4.26)$$

which lies in $\mathcal{H}_K$ as a bridge to perform further operator analysis. f_{λ_1} in fact has the operator representation $f_{\lambda_1} = (L_K + \lambda_1 I)^{-1} L_K f_\rho$. We derive the following estimate.

Proposition 5. *Assume that $|y| \leq M$ and (2.7) hold almost surely. Suppose the parameters $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2c_V \lambda_2 = \lambda_1$. Then there holds almost surely*

$$\max \left\{ \|f_{D, \lambda_1, \lambda_2} - f_{\lambda_1}\|_\rho, \sqrt{\lambda_1} \|f_{D, \lambda_1, \lambda_2} - f_{\lambda_1}\|_K \right\} \leq 2\mathcal{A}_{D, \lambda_1, \lambda_2, V_D} \Pi_{D, \lambda_1} + 4\Xi_{D, \lambda_1} \mathcal{A}_{D, \lambda_1, \lambda_2, V_D} \|f_{\lambda_1}\|_K + \frac{2\lambda_2 c_V}{\sqrt{\lambda_1}} \|f_{\lambda_1}\|_K.$$

Proof. Start from the following decomposition

$$\begin{aligned} f_{D, \lambda_1, \lambda_2} - f_{\lambda_1} &= (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} S_D^T y_D - (L_K + \lambda_1 I)^{-1} L_K f_\rho \\ &= (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (S_D^T y_D - L_K f_\rho) \\ &\quad + \left[(L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} - (L_K + \lambda_1 I)^{-1} \right] L_K f_\rho \\ &= (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (S_D^T y_D - L_K f_\rho) \\ &\quad + \left[(L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} - (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \right] L_K f_\rho \\ &\quad + \left[(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} - (L_K + \lambda_1 I)^{-1} \right] L_K f_\rho. \end{aligned}$$

Then use the fact $\|h\|_\rho = \|L_K^{1/2} h\|_K$ for any $h \in L_{\rho X_\mu}^2$, we have

$$\begin{aligned} &\max \left\{ \|f_{D, \lambda_1, \lambda_2} - f_{\lambda_1}\|_\rho, \sqrt{\lambda_1} \|f_{D, \lambda_1, \lambda_2} - f_{\lambda_1}\|_K \right\} \\ &\leq \left\| (L_K + \lambda_1 I)^{\frac{1}{2}} (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (S_D^T y_D - L_K f_\rho) \right\|_K \\ &\quad + \left\| (L_K + \lambda_1 I)^{\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_K - L_{K, D}) (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} L_K f_\rho \right\|_K \\ &\quad + \left\| (L_K + \lambda_1 I)^{\frac{1}{2}} (L_K + \lambda_1 I)^{-1} \lambda_2 V_D^T V_D (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} L_K f_\rho \right\|_K \\ &:= \tilde{\mathcal{T}}_1 + \tilde{\mathcal{T}}_2 + \tilde{\mathcal{T}}_3. \end{aligned}$$

We estimate $\tilde{\mathcal{T}}_1, \tilde{\mathcal{T}}_2, \tilde{\mathcal{T}}_3$ as follows. Decompose the operators in $\tilde{\mathcal{T}}_1$, we have

$$\begin{aligned} \tilde{\mathcal{T}}_1 &= \left\| (L_K + \lambda_1 I)^{\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \right. \\ &\quad \left. (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \right. \\ &\quad \left. (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_K + \lambda_1 I)^{\frac{1}{2}} (L_K + \lambda_1 I)^{-\frac{1}{2}} (S_D^T y_D - L_K f_\rho) \right\|_K \\ &\leq \left\| (L_K + \lambda_1 I)^{\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \left\| (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \right. \\ &\quad \left. (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \left\| (L_{K, D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{\frac{1}{2}} \right\| \\ &\quad \left\| (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_K + \lambda_1 I)^{\frac{1}{2}} \right\| \left\| (L_K + \lambda_1 I)^{-\frac{1}{2}} (S_D^T y_D - L_K f_\rho) \right\|_K \\ &\leq 2^{\frac{1}{2}} \mathcal{A}_{D, \lambda_1, \lambda_2, V_D} \cdot 2^{\frac{1}{2}} \Pi_{D, \lambda_1} = 2\mathcal{A}_{D, \lambda_1, \lambda_2, V_D} \Pi_{D, \lambda_1}. \end{aligned}$$

$\tilde{\mathcal{T}}_2$ is estimated as follows,

$$\begin{aligned}\tilde{\mathcal{T}}_2 &\leq \left\| (L_K + \lambda_1 I)^{\frac{1}{2}} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} \right\| \left\| (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-\frac{1}{2}} (L_K + \lambda_1 I)^{\frac{1}{2}} \right\| \\ &\quad \left\| (L_K + \lambda_1 I)^{-\frac{1}{2}} (L_K - L_{K,D}) \right\| \left\| (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D) \right\| \\ &\quad \left\| (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_K + \lambda_1 I) \right\| \left\| (L_K + \lambda_1 I)^{-1} L_K f_\rho \right\|_K \\ &\leq 2^{\frac{1}{2}} \cdot 2^{\frac{1}{2}} \cdot 2\Xi_{D,\lambda_1} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \|f_{\lambda_1}\|_K = 4\Xi_{D,\lambda_1} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \|f_{\lambda_1}\|_K.\end{aligned}$$

$\tilde{\mathcal{T}}_3$ is estimated as follows,

$$\begin{aligned}\tilde{\mathcal{T}}_3 &\leq \left\| (L_K + \lambda_1 I)^{\frac{1}{2}} (L_K + \lambda_1 I)^{-\frac{1}{2}} (L_K + \lambda_1 I)^{-\frac{1}{2}} \lambda_2 V_D^T V_D \right. \\ &\quad \left. (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_K + \lambda_1 I) (L_K + \lambda_1 I)^{-1} L_K f_\rho \right\|_K \\ &\leq \left\| (L_K + \lambda_1 I)^{-\frac{1}{2}} \right\| \lambda_2 c_V \left\| (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_K + \lambda_1 I) \right\| \left\| (L_K + \lambda_1 I)^{-1} L_K f_\rho \right\|_K \\ &\leq \lambda_2 c_V \frac{1}{\sqrt{\lambda_1}} 2 \|f_{\lambda_1}\|_K.\end{aligned}$$

The desired result is obtained after combining the above three estimates for $\tilde{\mathcal{T}}_1, \tilde{\mathcal{T}}_2, \tilde{\mathcal{T}}_3$. ■

Remark 1. Under the regularity condition $f_\rho = L_K^r(g_\rho)$, $r \in (0, 1/2)$ for some $g_\rho \in L_{\rho_{X_\mu}}^2$, the main difference with case $r \in [1/2, 1]$ is that the regression function f_ρ no longer lies in $\mathcal{H}_K$ any more. Proposition 5 is derived to overcome the difficulties. It plays an important role in the following proofs. In fact, when $r \in (0, 1/2)$, the previous estimates in Proposition 1 for $\|f_{D,\lambda_1,\lambda_2} - f_\rho\|_\rho$ will fail for further deriving learning rates of $\|f_{\hat{D},\lambda_1,\lambda_2} - f_\rho\|_\rho$. The main gap and difficulty arise in previous second-term $\mathcal{T}_2$ and third-term $\mathcal{T}_3$ estimates. When $r \in (0, 1/2)$, if we continue handling $\|f_{D,\lambda_1,\lambda_2} - f_\rho\|_\rho$ and decomposing term $\mathcal{T}_2$ as that in the proof of Proposition 1, it can be observed that the operator norms

$$\begin{aligned}&\left\| (\lambda_1 I + L_{K,D} + \lambda_2 V_D^T V_D)^{-r+\frac{1}{2}} (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{r-\frac{1}{2}} \right\|, \\ &\left\| (\lambda_1 I + L_K + \lambda_2 V_D^T V_D)^{-r+\frac{1}{2}} (\lambda_1 I + L_K)^{r-\frac{1}{2}} \right\|,\end{aligned}$$

and

$$\left\| (\lambda_1 I + L_K)^{-r+\frac{1}{2}} L_K^{r-\frac{1}{2}} \right\|$$

do not possess effective upper bounds for further estimates. Hence, they can not be used to derive learning rates of $\|f_{\hat{D},\lambda_1,\lambda_2} - f_\rho\|_\rho$ in the hard learning scenario where $r \in (0, 1/2)$. The previous work in [6] also suffered from similar difficulty, and hence only learning rates related to standard regularity index $r \in [1/2, 1]$ can be derived. Till now, we have described one of the main reasons why previous analysis fails in the hard learning scenario. To overcome the difficulties, in Proposition 5, we introduce data-free representation f_{λ_1} and make estimates of the RKHS norm and $L_{\rho_{X_\mu}}^2$ -norm of $f_{D,\lambda_1,\lambda_2} - f_{\lambda_1}$. Due to the nice representation of f_{λ_1} , the decomposition for $\tilde{\mathcal{T}}_1, \tilde{\mathcal{T}}_2$ and $\tilde{\mathcal{T}}_3$ successfully excludes the interference of the regularity index of r in the operator decomposition process. After using the operator decomposition for $\tilde{\mathcal{T}}_1, \tilde{\mathcal{T}}_2$ and $\tilde{\mathcal{T}}_3$, the corresponding upper bounds have essentially changed. We finally derive a mild bound in Proposition 5, which can be used to estimate learning rates further. The upper bound in Proposition 5 also influences the error bounds for the second-stage estimate of $f_{\hat{D},\lambda_1,\lambda_2} - f_{D,\lambda_1,\lambda_2}$, this fact can be witnessed throughout the rest of this section (see Proposition 7 and the proof of Theorem 3).

Furthermore, in contrast to previous work, to capture more potential features of regularized distribution regression scheme, we have proposed a novel multi-penalty distribution regression scheme and considered an additional penalty induced by the operator V_D that previous works on distribution regression have not explored yet. The participation of the operator V_D also increases the difficulty of the operator analysis process in the estimates mentioned above compared with previous works [6], [21]. Compared with previous analyses in works [6], [21], we have already introduced two new crucial norms $\mathcal{A}_{D,\lambda_1,\lambda_2,V_D} = \|(\lambda_1 I + L_K + \lambda_2 V_D^T V_D)(\lambda_1 I + L_{K,D} + \lambda_2 V_D^T V_D)^{-1}\|$ and $\Omega_{D,\lambda_1,\lambda_2,V_D} = \|L_K^{1/2}(L_{K,\widehat{D}} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}\|$ to overcome the difficulties arising from the operator V_D . From the analysis above, the two quantities successfully realize effective bounds for operator decompositions in our new multi-penalty setting. Hence they overcame the current complicated multi-penalty analysis environment and finally produced satisfactory theoretical results on learning rates.

Based on the estimate in Proposition 5, the following expected error bound is obtained.

Proposition 6. Assume that $|y| \leq M$ and (2.7) hold almost surely. Let the regularity condition (2.4) hold with some index $0 < r < 1/2$. If $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2c_V \lambda_2 = \lambda_1$. Then we have,

$$E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2} - f_\rho\|_\rho \right] \leq 4(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} (2\Gamma(3) + \log^2 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \left[M(\kappa + 1)\mathcal{B}'_{|D|,\lambda_1} \right. \\ \left. + \|g_\rho\|_\rho (\kappa^2 + \kappa) \lambda_1^{r-\frac{1}{2}} \mathcal{B}'_{|D|,\lambda_1} \right] + 2\|g_\rho\|_\rho c_V \lambda_2 \lambda_1^{r-1} + \lambda_1^r \|g_\rho\|_\rho. \quad (4.27)$$

Proof. Note that the following decomposition holds,

$$E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2} - f_\rho\|_\rho \right] \leq E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2} - f_{\lambda_1}\|_\rho \right] + E_{\mathbf{z}^{[D]}} \left[\|f_{\lambda_1} - f_\rho\|_\rho \right]. \quad (4.28)$$

The one stage estimate in [22] shows that $\|f_{\lambda_1} - f_\rho\|_\rho \leq \lambda_1^r \|g_\rho\|_\rho$. Hence, the second term is bounded as follows,

$$E_{\mathbf{z}^{[D]}} \left[\|f_{\lambda_1} - f_\rho\|_\rho \right] \leq \lambda_1^r \|g_\rho\|_\rho. \quad (4.29)$$

We estimate the first term by taking expectations on both sides of Proposition 5. Since

$$\|f_{\lambda_1}\|_K = \|(\lambda_1 I + L_K)^{-1} L_K f_\rho\|_K \leq \|(\lambda_1 I + L_K)^{-1} L_K^{r+\frac{1}{2}}\| \|L_K^{1/2} g_\rho\|_K \leq \lambda_1^{r-\frac{1}{2}} \|g_\rho\|_\rho,$$

by using Schwarz inequality and using Lemma 4 with $s = 2$, we have

$$E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2} - f_{\lambda_1}\|_\rho \right] \leq 2E_{\mathbf{z}^{[D]}} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \Pi_{D,\lambda_1} \right] + 4E_{\mathbf{z}^{[D]}} \left[\Xi_{D,\lambda_1} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \right] \|f_{\lambda_1}\|_K + \frac{2\lambda_2 c_V}{\sqrt{\lambda_1}} \|f_{\lambda_1}\|_K \\ \leq 2 \left\{ E_{\mathbf{z}^{[D]}} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^2 \right] \right\}^{\frac{1}{2}} \left\{ E_{\mathbf{z}^{[D]}} \left[\Pi_{D,\lambda_1}^2 \right] \right\}^{\frac{1}{2}} \\ + 4\lambda_1^{r-\frac{1}{2}} \|g_\rho\|_\rho \left\{ E_{\mathbf{z}^{[D]}} \left[\Xi_{D,\lambda_1}^2 \right] \right\}^{\frac{1}{2}} \left\{ E_{\mathbf{z}^{[D]}} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^2 \right] \right\}^{\frac{1}{2}} + \frac{2\lambda_2 c_V}{\sqrt{\lambda_1}} \lambda_1^{r-\frac{1}{2}} \|g_\rho\|_\rho$$

which can be further bounded by

$$2(2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 (2\Gamma(3) + \log^2 2)^{\frac{1}{2}} 2M(\kappa + 1)\mathcal{B}'_{|D|,\lambda_1} \\ + 4\lambda_1^{r-\frac{1}{2}} \|g_\rho\|_\rho (2\Gamma(3) + \log^2 2)^{\frac{1}{2}} 2(\kappa^2 + \kappa)\mathcal{B}'_{|D|,\lambda_1} (2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \left(\frac{2\mathcal{B}_{|D|,\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 + 2c_V \lambda_2 \lambda_1^{r-1} \|g_\rho\|_\rho.$$

The desired result is obtained by combining these estimates. ■

For the case $f_\rho \notin \mathcal{H}_K$, we need the following new estimate for $\left\{E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2}\|_K^2 \right] \right\}^{\frac{1}{2}}$. The approach on deriving an upper bound of $\left\{E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2}\|_K^2 \right] \right\}^{\frac{1}{2}}$ relies on our new estimate in Proposition 5.

Proposition 7. *Suppose $|y| \leq M$ almost surely. Let the regularity condition (2.4) hold with some index $r \in (0, 1/2)$. If $\lambda_1, \lambda_2 \in (0, 1)$ satisfy $2\lambda_2 c_V = \lambda_1$, then we have*

$$\begin{aligned} \left\{E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2}\|_K^2 \right] \right\}^{\frac{1}{2}} &\leq 2\sqrt{3}(2\Gamma(9) + \log^8 2)^{\frac{1}{4}}(2\Gamma(5) + \log^4 2)^{\frac{1}{4}} \left(\frac{2\mathcal{B}_{[D],\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \times \\ &\times \left[2M(\kappa + 1) \frac{1}{\sqrt{\lambda_1}} \mathcal{B}'_{[D],\lambda_1} + 2(\kappa^2 + \kappa) \|g_\rho\|_\rho \lambda_1^{r-1} \mathcal{B}'_{[D],\lambda_1} \right] + 2\sqrt{3} \|g_\rho\|_\rho \lambda_1^{r-\frac{1}{2}}. \end{aligned}$$

Proof. We start with the decomposition $\|f_{D,\lambda_1,\lambda_2}\|_K \leq \|f_{D,\lambda_1,\lambda_2} - f_{\lambda_1}\|_K + \|f_{\lambda_1}\|_K$. Proposition 5 implies

$$\sqrt{\lambda_1} \|f_{D,\lambda_1,\lambda_2} - f_{\lambda_1}\|_K \leq 2\mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \Pi_{D,\lambda_1} + 4\Xi_{D,\lambda_1} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \|f_{\lambda_1}\|_K + \frac{2\lambda_2 c_V}{\sqrt{\lambda_1}} \|f_{\lambda_1}\|_K.$$

The above inequality together with the fact $\|f_{\lambda_1}\|_K \leq \lambda_1^{r-\frac{1}{2}} \|g_\rho\|_\rho$ imply that

$$\|f_{D,\lambda_1,\lambda_2} - f_{\lambda_1}\|_K \leq 2 \frac{1}{\sqrt{\lambda_1}} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \Pi_{D,\lambda_1} + 4\lambda_1^{-\frac{1}{2}} \lambda_1^{r-\frac{1}{2}} \|g_\rho\|_\rho \Xi_{D,\lambda_1} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} + 2 \|g_\rho\|_\rho c_V \lambda_2 \lambda_1^{r-1-\frac{1}{2}}.$$

Noting the condition $2\lambda_2 c_V = \lambda_1$, we obtain

$$\|f_{D,\lambda_1,\lambda_2}\|_K \leq \frac{2}{\sqrt{\lambda_1}} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} \Pi_{D,\lambda_1} + 4\lambda_1^{r-1} \|g_\rho\|_\rho \Xi_{D,\lambda_1} \mathcal{A}_{D,\lambda_1,\lambda_2,V_D} + 2\lambda_1^{r-\frac{1}{2}} \|g_\rho\|_\rho.$$

The basic Jensen's inequality and Schwarz inequality implies that

$$\begin{aligned} E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2}\|_K^2 \right] &\leq \frac{12}{\lambda_1} \left\{ E_{\mathbf{z}^{[D]}} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^4 \right] \right\}^{\frac{1}{2}} \left\{ E_{\mathbf{z}^{[D]}} \left[\Pi_{D,\lambda_1}^4 \right] \right\}^{\frac{1}{2}} \\ &+ 48\lambda_1^{2r-2} \|g_\rho\|_\rho^2 \left\{ E_{\mathbf{z}^{[D]}} \left[\Xi_{D,\lambda_1}^4 \right] \right\}^{\frac{1}{2}} \left\{ E_{\mathbf{z}^{[D]}} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^4 \right] \right\}^{\frac{1}{2}} + 12\lambda_1^{2r-1} \|g_\rho\|_\rho^2. \end{aligned}$$

Using Jensen's inequality and Lemma 4 with $s = 4$, we have

$$\begin{aligned} \left\{ E_{\mathbf{z}^{[D]}} \left[\|f_{D,\lambda_1,\lambda_2}\|_K^2 \right] \right\}^{\frac{1}{2}} &\leq \frac{2\sqrt{3}}{\sqrt{\lambda_1}} \left\{ E_{\mathbf{z}^{[D]}} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^4 \right] \right\}^{\frac{1}{4}} \left\{ E_{\mathbf{z}^{[D]}} \left[\Pi_{D,\lambda_1}^4 \right] \right\}^{\frac{1}{4}} \\ &+ 2\sqrt{3} \lambda_1^{r-1} \|g_\rho\|_\rho \left\{ E_{\mathbf{z}^{[D]}} \left[\Xi_{D,\lambda_1}^4 \right] \right\}^{\frac{1}{4}} \left\{ E_{\mathbf{z}^{[D]}} \left[\mathcal{A}_{D,\lambda_1,\lambda_2,V_D}^4 \right] \right\}^{\frac{1}{4}} + 2\sqrt{3} \lambda_1^{r-\frac{1}{2}} \|g_\rho\|_\rho \\ &\leq 2\sqrt{3}(2\Gamma(9) + \log^8 2)^{\frac{1}{4}} \left(\frac{2\mathcal{B}_{[D],\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 (2\Gamma(5) + \log^4 2)^{\frac{1}{4}} 2M(\kappa + 1) \mathcal{B}'_{[D],\lambda_1} \frac{1}{\sqrt{\lambda_1}} \\ &4\sqrt{3} \|g_\rho\|_\rho (2\Gamma(9) + \log^8 2)^{\frac{1}{4}} \left(\frac{2\mathcal{B}_{[D],\lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 (2\Gamma(5) + \log^4 2)^{\frac{1}{4}} 2(\kappa^2 + \kappa) \mathcal{B}'_{[D],\lambda_1} \lambda_1^{r-1} + 2\sqrt{3} \|g_\rho\|_\rho \lambda_1^{r-\frac{1}{2}}. \end{aligned}$$

After combining the first two terms, we finish the proof. $\blacksquare$

Now we are ready to prove Theorem 3.

Proof of Theorem 3. Under above preparations, we need to re-estimate the following expected norm,

$$\begin{aligned} & E \left[\Omega_{D, \lambda_1, \lambda_2, V_D} \|L_{K, D} - L_{K, \widehat{D}}\| \|f_{D, \lambda_1, \lambda_2}\|_K \right] \\ & \leq \left(2\sqrt{2}(2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_K^{\frac{\alpha}{2}}}{\lambda_1 \widetilde{d}^{\frac{\alpha}{2}}} + 2 \right) \kappa L (2 + \sqrt{\pi})^{\frac{1}{2}} \frac{2^{\frac{\alpha+2}{2}} B_K^{\frac{\alpha}{2}}}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \left\{ E_{\mathbf{z}|D} \left[\|f_{D, \lambda_1, \lambda_2}\|_K^2 \right] \right\}^{\frac{1}{2}} \left\{ E_{\mathbf{z}|D} \left[\mathcal{A}_{D, \lambda_1, \lambda_2, V_D}^2 \right] \right\}^{\frac{1}{2}}. \end{aligned}$$

Substitute Proposition 7 into above inequality, we know that the above terms can be further bounded by

$$\begin{aligned} & \left(2\sqrt{2}(2 + \sqrt{\pi})^{\frac{1}{2}} \kappa L \frac{2^{\frac{\alpha+2}{2}} B_K^{\frac{\alpha}{2}}}{\lambda_1 \widetilde{d}^{\frac{\alpha}{2}}} + 2 \right) \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 (2\Gamma(5) + \log^4 2)^{\frac{1}{2}} \kappa L (2 + \sqrt{\pi})^{\frac{1}{2}} 2^{\frac{\alpha+2}{2}} B_K^{\frac{\alpha}{2}} \times \\ & \left(4\sqrt{3}(2\Gamma(9) + \log^8 2)^{\frac{1}{4}} (2\Gamma(5) + \log^4 2)^{\frac{1}{4}} \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1 \right)^2 \left[2M(\kappa + 1) \frac{1}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \frac{1}{\sqrt{\lambda_1}} \mathcal{B}'_{|D|, \lambda_1} \right. \right. \\ & \left. \left. + 2(\kappa^2 + \kappa) \|g_\rho\|_\rho \frac{\lambda_1^{r-1}}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \mathcal{B}'_{|D|, \lambda_1} \right] + 2\sqrt{3} \|g_\rho\|_\rho \frac{\lambda_1^{r-\frac{1}{2}}}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \right). \end{aligned} \quad (4.30)$$

On the other hand, note that the estimate of (4.19) on $E \left[\Omega_{D, \lambda_1, \lambda_2, V_D} \|\hat{S}_D^T y_D - S_D^T y_D\| \right]$ in case $r \in (0, 1/2)$ does not change. Now combine (4.19), (4.30) and (4.27) in Proposition 6, the desired bound follows. ■

Remark 2. In contrast to previous works, for handling the tough case $r \in (0, 1/2)$, since we have employed the different decomposition approach described above, the core terms of the second-stage estimates have already changed to some new terms

$$\frac{1}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \frac{1}{\sqrt{\lambda_1}} \mathcal{B}'_{|D|, \lambda_1}, \frac{\lambda_1^{r-1}}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \mathcal{B}'_{|D|, \lambda_1}, \frac{\lambda_1^{r-\frac{1}{2}}}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}}.$$

In fact, when $r \in (0, 1/2)$, in the above estimate of $\Omega_{D, \lambda_1, \lambda_2, V_D} \|L_{K, D} - L_{K, \widehat{D}}\| \|f_{D, \lambda_1, \lambda_2}\|_K$, the derivation of the upper bound of $\|f_{D, \lambda_1, \lambda_2}\|_K$ deeply relies on our new estimates in Proposition 5 and Proposition 7.

Proof of Theorem 4. We substitute the main parameters

$$\lambda_1 = |D|^{-\frac{1}{1+\beta}}, \quad \lambda_2 = \frac{1}{2c_V} |D|^{-\frac{1}{1+\beta}}, \quad \widetilde{d} = d_1 = d_2 = \dots = d_{|D|} = |D|^{\frac{2}{\alpha(1+\beta)}}$$

into the general error bound estimate in Theorem 3. We obtain

$$\mathcal{B}_{|D|, \lambda_1} = \frac{2\kappa}{\sqrt{|D|}} \left(\frac{\kappa}{\sqrt{|D|} \lambda_1} + \sqrt{\mathcal{N}(\lambda_1)} \right) \leq 2\kappa(\kappa + \sqrt{\mathcal{C}_0}) |D|^{-\frac{1}{2(1+\beta)}}$$

and

$$\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} \leq 4\kappa(\kappa + \sqrt{\mathcal{C}_0}).$$

Now we estimate the key terms

$$\frac{1}{\lambda_1 \widetilde{d}^{\frac{\alpha}{2}}}, \frac{1}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \frac{1}{\sqrt{\lambda_1}} \mathcal{B}'_{|D|, \lambda_1}, \frac{\lambda_1^{r-1}}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}} \mathcal{B}'_{|D|, \lambda_1}, \frac{\lambda_1^{r-\frac{1}{2}}}{\lambda_1^{\frac{1}{2}} \widetilde{d}^{\frac{\alpha}{2}}}.$$

We start with the following three estimates,

$$\begin{aligned}\frac{1}{\lambda_1 \tilde{d}^{\frac{\alpha}{2}}} &= |D|^{\frac{1}{1+\beta}} \cdot |D|^{-\frac{\alpha}{2} \frac{2}{\alpha(1+\beta)}} = |D|^{\frac{1}{1+\beta} - \frac{1}{1+\beta}} = 1. \\ \frac{1}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} &= |D|^{\frac{1}{2(1+\beta)}} \cdot |D|^{-\frac{\alpha}{2} \frac{2}{\alpha(1+\beta)}} = |D|^{-\frac{1}{2(1+\beta)}}, \\ \mathcal{B}'_{|D|, \lambda_1} &= \frac{1}{|D| \sqrt{\lambda_1}} + \frac{\sqrt{\mathcal{N}(\lambda_1)}}{\sqrt{|D|}} \leq (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{1}{2(1+\beta)}}.\end{aligned}$$

The above estimate for $\mathcal{B}'_{|D|, \lambda_1}$ and the fact $r \in (0, 1/2)$ in fact also imply

$$\mathcal{B}'_{|D|, \lambda_1} \leq (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{1}{2(1+\beta)}} < (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{r}{1+\beta}}.$$

Therefore, we obtain

$$\begin{aligned}\frac{1}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} \frac{1}{\sqrt{\lambda_1}} \mathcal{B}'_{|D|, \lambda_1} &\leq |D|^{-\frac{1}{2(1+\beta)}} |D|^{\frac{1}{2(1+\beta)}} (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{1}{2(1+\beta)}} \\ &= (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{1}{2(1+\beta)}} < (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{r}{1+\beta}},\end{aligned}$$

$$\frac{\lambda_1^{r-1}}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} \mathcal{B}'_{|D|, \lambda_1} \leq |D|^{-\frac{r-1}{1+\beta}} |D|^{-\frac{1}{2(1+\beta)}} (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{1}{2(1+\beta)}} = (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{r}{1+\beta}},$$

and

$$\frac{\lambda_1^{r-\frac{1}{2}}}{\lambda_1^{\frac{1}{2}} \tilde{d}^{\frac{\alpha}{2}}} = |D|^{-\frac{r-\frac{1}{2}}{1+\beta}} |D|^{-\frac{1}{2(1+\beta)}} = |D|^{-\frac{r}{1+\beta}}.$$

We turn to estimate $\lambda_1^{r-\frac{1}{2}} \mathcal{B}'_{|D|, \lambda_1}$, $c_V \lambda_2 \lambda_1^{r-1}$ and λ_1^r . Above basic estimate for $\mathcal{B}'_{|D|, \lambda_1}$ implies

$$\lambda_1^{r-\frac{1}{2}} \mathcal{B}'_{|D|, \lambda_1} \leq |D|^{-\frac{r-\frac{1}{2}}{1+\beta}} \cdot (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{1}{2(1+\beta)}} \leq (1 + \sqrt{\mathcal{C}_0}) |D|^{-\frac{r}{1+\beta}}.$$

Also, note that the above parameters design for λ_1 and λ_2 imply

$$c_V \lambda_2 \lambda_1^{r-1} = \frac{1}{2} |D|^{-\frac{r}{1+\beta}}, \quad \lambda_1^r = |D|^{-\frac{r}{1+\beta}}.$$

After combining all the above estimates with Theorem 3, the desired rates are obtained. $\blacksquare$

5 Distributed learning with multi-penalty distribution regression and its learning rates

This section is devoted to proving the result on distributed learning with multi-penalty distribution regression. The proof is based on an observation of the relation between our former estimates in this paper and the one-stage distributed learning theory. Firstly, we make some preparations. Recall the definition of f_{λ_1} , we have

$$\lambda_1 f_{\lambda_1} = L_K(f_\rho - f_{\lambda_1}).$$

Then for any mean embedding set D associated with the given data set $\tilde{D}$, we have

$$\begin{aligned}
& f_{D, \lambda_1, \lambda_2} - f_{\lambda_1} \\
&= (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} S_D^T y_D - f_{\lambda_1} \\
&= (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} [S_D^T y_D - L_{K,D} f_{\lambda_1} - \lambda_1 f_{\lambda_1} - \lambda_2 V_D^T V_D f_{\lambda_1}] \\
&= (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \left\{ \frac{1}{|D|} \sum_{z \in D} (y - f_{\lambda_1}(\mu_x)) K_{\mu_x} - L_K(f_\rho - f_{\lambda_1}) - \lambda_2 V_D^T V_D f_{\lambda_1} \right\}
\end{aligned}$$

in which $z = (\mu_x, y) \in X_\mu \times Y$. Denote

$$\Delta_D = \frac{1}{|D|} \sum_{z \in D} (y - f_{\lambda_1}(\mu_x)) K_{\mu_x} - L_K(f_\rho - f_{\lambda_1})$$

and

$$\mathcal{Q}_D = (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} - (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}.$$

Then we can decompose $f_{\tilde{D}, \lambda_1, \lambda_2} - f_{\lambda_1}$ as

$$\begin{aligned}
f_{\tilde{D}, \lambda_1, \lambda_2} - f_{\lambda_1} &= \left[f_{\tilde{D}, \lambda_1, \lambda_2} - f_{D, \lambda_1, \lambda_2} \right] + \mathcal{Q}_D \Delta_D + (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \Delta_D \\
&\quad - (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \lambda_2 V_D^T V_D f_{\lambda_1}.
\end{aligned} \tag{5.1}$$

In the following, we will use the corresponding notations $\mathcal{Q}_{D_j}$, L_{K,D_j} , Δ_{D_j} , V_{D_j} involving the sample subset $\tilde{D}_j$ and its associated mean embedding set D_j . Then the corresponding representations for $f_{\tilde{D}_j, \lambda_1, \lambda_2} - f_{\lambda_1}$ are well-defined for local data sets. Applying second-order decomposition (4.9) to $\mathcal{Q}_D$ with

$$\begin{aligned}
A &= L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D, \\
B &= L_K + \lambda_1 I + \lambda_2 V_D^T V_D,
\end{aligned}$$

we have

$$\begin{aligned}
L_K^{1/2} \mathcal{Q}_D \Delta_D &= L_K^{1/2} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_K - L_{K,D}) (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \Delta_D \\
&\quad + L_K^{1/2} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} (L_K - L_{K,D}) (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \\
&\quad (L_K - L_{K,D}) (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \Delta_D
\end{aligned}$$

which can be further decomposed as

$$\begin{aligned}
& \{L_K^{1/2} (L_K + \lambda_1 I)^{-1/2}\} \{(L_K + \lambda_1 I)^{1/2} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1/2}\} \\
& \{(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1/2} (L_K + \lambda_1 I)^{1/2}\} \{(L_K + \lambda_1 I)^{-1/2} (L_K - L_{K,D})\} \\
& \{(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1/2}\} \{(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1/2} (L_K + \lambda_1 I)^{1/2}\} \\
& \{(L_K + \lambda_1 I)^{-1/2} \Delta_D\} \\
& + \{L_K^{1/2} (L_K + \lambda_1 I)^{-1/2}\} \{(L_K + \lambda_1 I)^{1/2} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1/2}\} \\
& \{(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1/2} (L_K + \lambda_1 I)^{1/2}\} \{(L_K + \lambda_1 I)^{-1/2} (L_K - L_{K,D})\} \\
& \{(L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1}\} \{(L_K - L_{K,D}) (L_K + \lambda_1 I)^{-1/2}\} \\
& \{(L_K + \lambda_1 I)^{1/2} (L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1/2}\} \{(L_K + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1/2} (L_K + \lambda_1 I)^{1/2}\} \\
& \{(L_K + \lambda_1 I)^{-1/2} \Delta_D\}.
\end{aligned}$$

After taking RKHS-norms on both sides, we obtain

$$\begin{aligned}\left\|L_K^{1/2}\mathcal{Q}_D\Delta_D\right\|_K &\leq 2^{\frac{3}{2}}\frac{\Xi_{D,\lambda_1}}{\sqrt{\lambda_1}}\left\|(L_K+\lambda_1I)^{-1/2}\Delta_D\right\|_K + 4\frac{\Xi_{D,\lambda_1}^2}{\lambda_1}\left\|(L_K+\lambda_1I)^{-1/2}\Delta_D\right\|_K \\ &= \left(2^{\frac{3}{2}}\frac{\Xi_{D,\lambda_1}}{\sqrt{\lambda_1}} + 4\frac{\Xi_{D,\lambda_1}^2}{\lambda_1}\right)\left\|(L_K+\lambda_1I)^{-1/2}\Delta_D\right\|_K\end{aligned}\quad (5.2)$$

where Ξ_{D,λ_1} is defined in (4.25). Considering the above representation (5.1) on mean embedding set D_j , we know that

$$\begin{aligned}f_{\widehat{D}_j,\lambda_1,\lambda_2} - f_{\lambda_1} &= \left[f_{\widehat{D}_j,\lambda_1,\lambda_2} - f_{D_j,\lambda_1,\lambda_2}\right] + \mathcal{Q}_{D_j}\Delta_{D_j} + (L_K + \lambda_1I + \lambda_2V_{D_j}^TV_{D_j})^{-1}\Delta_{D_j} \\ &\quad - (L_{K,D_j} + \lambda_1I + \lambda_2V_{D_j}^TV_{D_j})^{-1}\lambda_2V_{D_j}^TV_{D_j}f_{\lambda_1}.\end{aligned}$$

Then we have

$$\begin{aligned}\overline{f_{\widehat{D},\lambda_1,\lambda_2}} - f_{\lambda_1} &= \sum_{j=1}^m \frac{|D_j|}{|D|} \left[f_{\widehat{D}_j,\lambda_1,\lambda_2} - f_{\lambda_1}\right] \\ &= \sum_{j=1}^m \frac{|D_j|}{|D|} \left[f_{\widehat{D}_j,\lambda_1,\lambda_2} - f_{D_j,\lambda_1,\lambda_2}\right] + \sum_{j=1}^m \frac{|D_j|}{|D|} (L_{K,D_j} + \lambda_1I + \lambda_2V_{D_j}^TV_{D_j})^{-1}\Delta_{D_j} \\ &\quad - \sum_{j=1}^m \frac{|D_j|}{|D|} (L_{K,D_j} + \lambda_1I + \lambda_2V_{D_j}^TV_{D_j})^{-1}\lambda_2V_{D_j}^TV_{D_j}f_{\lambda_1}.\end{aligned}\quad (5.3)$$

Since $\Delta_D = \sum_{j=1}^m \frac{|D_j|}{|D|} \Delta_{D_j}$, subtraction between (5.3) and (5.1) yields the crucial error decomposition for two-stage multi-penalty distribution regression distributed learning scheme (1.4) in the following proposition.

Proposition 8. *For the distributed estimator $\overline{f_{\widehat{D},\lambda_1,\lambda_2}}$ and the estimator $f_{\widehat{D},\lambda_1,\lambda_2}$ associated with a single data set D , there holds the following two-stage error decomposition:*

$$\begin{aligned}\overline{f_{\widehat{D},\lambda_1,\lambda_2}} - f_{\widehat{D},\lambda_1,\lambda_2} &= \sum_{j=1}^m \frac{|D_j|}{|D|} \left[[f_{\widehat{D}_j,\lambda_1,\lambda_2} - f_{D_j,\lambda_1,\lambda_2}] - [f_{\widehat{D},\lambda_1,\lambda_2} - f_{D,\lambda_1,\lambda_2}] \right] \\ &\quad + \sum_{j=1}^m \frac{|D_j|}{|D|} \left[(L_{K,D_j} + \lambda_1I + \lambda_2V_{D_j}^TV_{D_j})^{-1} - (L_{K,D} + \lambda_1I + \lambda_2V_D^TV_D)^{-1} \right] \Delta_{D_j} \\ &\quad + \sum_{j=1}^m \frac{|D_j|}{|D|} \left[(L_{K,D} + \lambda_1I + \lambda_2V_D^TV_D)^{-1}\lambda_2V_D^TV_Df_{\lambda_1} - (L_{K,D_j} + \lambda_1I + \lambda_2V_{D_j}^TV_{D_j})^{-1}\lambda_2V_{D_j}^TV_{D_j}f_{\lambda_1} \right].\end{aligned}$$

In the following, we always denote the three terms on the right-hand side of the above error decomposition by $\mathcal{S}_1$, $\mathcal{S}_2$, $\mathcal{S}_3$.

Remark 3. *It can be observed that $\mathcal{S}_1$ presents the second-stage sampling effect to the multi-penalty distribution regression-based distributed learning algorithm. $\mathcal{S}_2$ is mainly related to one-stage distributed decomposition. $\mathcal{S}_3$ shows some additional multi-penalty influence on the proposed distributed learning algorithm. In contrast to former works, the decomposition is able to capture a more general setting when the data are functional or distribution data.*

For later use, we split $\Delta_{D_j} = \Delta'_{D_j} + \Delta''_{D_j}$ in which

$$\Delta'_{D_j} = S_{D_j}^Ty_{D_j} - L_{K,D_j}f_{\rho}, \quad \Delta''_{D_j} = L_{K,D_j}(f_{\rho} - f_{\lambda_1}) - L_K(f_{\rho} - f_{\lambda_1}). \quad (5.4)$$

We are ready to prove Theorem 5 on the optimal learning rates for distributed learning with multi-penalty distribution regression. In subsequent proof, we use the notation $f_1 \lesssim f_2$ to denote that there is an absolute constant C independent of $|D|$ and m such that $f_1 \leq C f_2$.

Proof of Theorem 5. We aim to bound $E[\|\mathcal{S}_1 + \mathcal{S}_2 + \mathcal{S}_3\|_\rho]$, in which $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$ are defined as in Proposition 8. The inequality in (4.21) implies that,

$$E\left[\|f_{\widehat{D}, \lambda_1, \lambda_2} - f_{D, \lambda_1, \lambda_2}\|_\rho\right] \lesssim \left(\frac{1}{\lambda_1 d^{\frac{\alpha}{2}}} + 1\right) \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right)^2 \frac{1}{\lambda_1^{\frac{1}{2}} d^{\frac{\alpha}{2}}} \left[2 + \frac{\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right) + (\lambda_1 + \lambda_2 c_V) \lambda_1^{r-\frac{3}{2}} \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right)^{2r-1}\right],$$

and same procedure implies that the above inequality continues to hold when D is replaced by D_j . Substituting $\lambda_1 = |D|^{-\frac{1}{2r+\beta}}$ into (2.6) yields $\mathcal{N}(\lambda_1) \leq C_0 |D|^{\frac{\beta}{2r+\beta}}$. Using the condition $m \leq |D|^{\frac{2r-1}{2r+\beta}}$ and $|D_j| = |D|/m$, $j = 1, 2, \dots, m$, we have

$$\frac{\mathcal{N}(\lambda_1)}{\lambda_1 |D_j|} \leq C_0 m |D|^{\frac{1-2r}{2r+\beta}} \leq C_0. \quad (5.5)$$

Then it follows that, for $j = 1, 2, \dots, m$,

$$\frac{\mathcal{B}_{|D_j|, \lambda_1}}{\sqrt{\lambda_1}} = \frac{2\kappa}{\sqrt{\lambda_1 |D_j|}} \left(\frac{\kappa}{\sqrt{|D_j| \lambda_1}} + \sqrt{\mathcal{N}(\lambda_1)}\right) \leq 2\kappa(\kappa + \sqrt{C_0}).$$

Note that by taking $\tilde{d} = d$, the former estimates (4.22) and (4.23) have already implied

$$\frac{1}{\lambda_1 d^{\frac{\alpha}{2}}} \leq 1, \quad \frac{1}{\lambda_1^{\frac{1}{2}} d^{\frac{\alpha}{2}}} \leq |D|^{-\frac{r}{2r+\beta}}, \quad (\lambda_1 + \lambda_2 c_V) \lambda_1^{r-\frac{3}{2}} \leq \frac{3}{2}.$$

Now we can estimate $\mathcal{S}_1$ in expectation form as

$$\begin{aligned} E\left[\|\mathcal{S}_1\|_\rho\right] &\lesssim \sum_{j=1}^m \frac{|D_j|}{|D|} \left(\frac{1}{\lambda_1 d^{\frac{\alpha}{2}}} + 1\right) \left(\frac{2\mathcal{B}_{|D_j|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right)^2 \frac{1}{\lambda_1^{\frac{1}{2}} d^{\frac{\alpha}{2}}} \left[2 + \frac{\mathcal{B}_{|D_j|, \lambda_1}}{\sqrt{\lambda_1}} \left(\frac{2\mathcal{B}_{|D_j|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right) + (\lambda_1 + \lambda_2 c_V) \lambda_1^{r-\frac{3}{2}} \left(\frac{2\mathcal{B}_{|D_j|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right)^{2r-1}\right] \\ &\quad + \left(\frac{1}{\lambda_1 d^{\frac{\alpha}{2}}} + 1\right) \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right)^2 \frac{1}{\lambda_1^{\frac{1}{2}} d^{\frac{\alpha}{2}}} \left[2 + \frac{\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right) + (\lambda_1 + \lambda_2 c_V) \lambda_1^{r-\frac{3}{2}} \left(\frac{2\mathcal{B}_{|D|, \lambda_1}}{\sqrt{\lambda_1}} + 1\right)^{2r-1}\right] \lesssim |D|^{-\frac{r}{2r+\beta}}. \end{aligned}$$

We turn to estimate $\mathcal{S}_2$. By subtracting and adding the operator $(L_K + \lambda_1 I + \lambda_2 V_{D_j}^T V_{D_j})^{-1}$, noting that $\Delta_{D_j} = \Delta'_{D_j} + \Delta''_{D_j}$, where $\Delta'_{D_j}, \Delta''_{D_j}$ are given as in (5.4), we have the decomposition

$$\mathcal{S}_2 = \sum_{j=1}^m \frac{|D_j|}{|D|} \mathcal{Q}_{D_j} \Delta'_{D_j} + \sum_{j=1}^m \frac{|D_j|}{|D|} \mathcal{Q}_{D_j} \Delta''_{D_j} - \mathcal{Q}_D \Delta_D. \quad (5.6)$$

Using the procedure of deriving (5.2) with Δ_D replaced by Δ'_{D_j} and Δ''_{D_j} , and taking RKHS-norms on both sides, we have

$$\left\|L_K^{1/2} \mathcal{Q}_{D_j} \Delta'_{D_j}\right\|_K^2 \leq \left(2^{\frac{3}{2}} \frac{\Xi_{D_j, \lambda_1}}{\sqrt{\lambda_1}} + 4 \frac{\Xi_{D_j, \lambda_1}^2}{\lambda_1}\right)^2 \left\|(L_K + \lambda_1 I)^{-1/2} \Delta'_{D_j}\right\|_K^2$$

$$\lesssim \left(\frac{\Xi_{D_j, \lambda_1}}{\sqrt{\lambda_1}} + \frac{\Xi_{D_j, \lambda_1}^2}{\lambda_1} \right)^2 \left\| (L_K + \lambda_1 I)^{-1/2} \Delta'_{D_j} \right\|_K^2$$

and

$$\left\| L_K^{1/2} \mathcal{Q}_{D_j} \Delta''_{D_j} \right\|_K \lesssim \left(\frac{\Xi_{D_j, \lambda_1}}{\sqrt{\lambda_1}} + \frac{\Xi_{D_j, \lambda_1}^2}{\lambda_1} \right) \left\| (L_K + \lambda_1 I)^{-1/2} \Delta''_{D_j} \right\|_K.$$

Now we have obtained the error bounds that we need via the decomposition for $\mathcal{S}_2$ in (5.6). We observe that the bounds $\left(\frac{\Xi_{D_j, \lambda_1}}{\sqrt{\lambda_1}} + \frac{\Xi_{D_j, \lambda_1}^2}{\lambda_1} \right) \left\| (L_K + \lambda_1 I)^{-1/2} \Delta'_{D_j} \right\|_K$ and $\left(\frac{\Xi_{D_j, \lambda_1}}{\sqrt{\lambda_1}} + \frac{\Xi_{D_j, \lambda_1}^2}{\lambda_1} \right) \left\| (L_K + \lambda_1 I)^{-1/2} \Delta''_{D_j} \right\|_K$ have already been handled in [12] for one-stage distributed learning theory. Therefore, we obtain that when $|D_j| = |D|/m$, $\mathcal{S}_2$, in fact, shares the same expected $L_{\rho_{X_\mu}}^2$ -norm bounds with the bound of Corollary 3 in [12]. Namely,

$$E_{\mathbf{z}^{(D)}} \left[\left\| \mathcal{S}_2 \right\|_\rho \right] \lesssim \sqrt{\frac{\mathcal{N}(\lambda_1)}{\lambda_1 |D|}} \left(\sqrt{\lambda_1} + \frac{m \|f_\rho - f_{\lambda_1}\|_\rho}{\sqrt{|D| \lambda_1}} \right). \quad (5.7)$$

Applying (5.5) with $|D_j|$ replaced by $|D|/m$, we know that $\frac{\mathcal{N}(\lambda_1)}{\lambda_1 |D|} \lesssim |D|^{-\frac{1-2r}{2r+\beta}}$, which shows

$$\sqrt{\frac{\mathcal{N}(\lambda_1)}{\lambda_1 |D|}} \lesssim |D|^{-\frac{\frac{1}{2}-r}{2r+\beta}}.$$

Recall the fact that $\|f_{\lambda_1} - f_\rho\|_\rho \leq \lambda_1^r \|g_\rho\|_\rho$, then we have

$$\begin{aligned} E_{\mathbf{z}^{(D)}} \left[\left\| \mathcal{S}_2 \right\|_\rho \right] &\lesssim \sqrt{\frac{\mathcal{N}(\lambda_1)}{\lambda_1 |D|}} \lambda_1^r \left(\lambda_1^{\frac{1}{2}-r} + \frac{m}{\sqrt{|D| \lambda_1}} \right) \\ &\lesssim |D|^{-\frac{r}{2r+\beta}} \left(|D|^{-\frac{\frac{1}{2}-r}{2r+\beta}} |D|^{\frac{\frac{1}{2}-r}{2r+\beta}} + |D|^{\frac{2r-1}{2r+\beta}} |D|^{-\frac{1}{2}} |D|^{\frac{\frac{1}{2}}{2r+\beta}} |D|^{\frac{\frac{1}{2}-r}{2r+\beta}} \right) \\ &\lesssim |D|^{-\frac{r}{2r+\beta}}. \end{aligned}$$

We finally estimate $\mathcal{S}_3$. Note that for any $j = 1, 2, \dots, m$,

$$\begin{aligned} &\left\| L_K^{1/2} (L_{K, D_j} + \lambda_1 I + \lambda_2 V_{D_j}^T V_{D_j})^{-1} \lambda_2 V_{D_j}^T V_{D_j} f_{\lambda_1} \right\|_K \\ &\leq \lambda_2 \left\| V_{D_j}^T V_{D_j} \right\| \left\| L_K^{1/2} (\lambda_1 I + L_K)^{-1/2} \right\| \left\| (\lambda_1 I + L_K)^{1/2} (L_K + \lambda_1 I + \lambda_2 V_{D_j}^T V_{D_j})^{-1/2} \right\| \\ &\quad \left\| (L_K + \lambda_1 I + \lambda_2 V_{D_j}^T V_{D_j})^{1/2} (L_{K, D_j} + \lambda_1 I + \lambda_2 V_{D_j}^T V_{D_j})^{-1/2} \right\| \\ &\quad \left\| (L_{K, D_j} + \lambda_1 I + \lambda_2 V_{D_j}^T V_{D_j})^{-1/2} \right\| \left\| f_{\lambda_1} \right\|_K \\ &\leq \lambda_2 c_V \sqrt{2} \mathcal{A}_{D_j, \lambda_1, \lambda_2, V_{D_j}}^{1/2} \frac{1}{\sqrt{\lambda_1}} \|f_{\lambda_1}\|_K. \end{aligned}$$

Taking expectations on both sides and noting from [22] the fact that $\|f_{\lambda_1}\|_K \leq \kappa^{2r-1} \|g_\rho\|_\rho$ when $r \in [1/2, 1]$, we have

$$\begin{aligned} &E_{\mathbf{z}^{(D)}} \left[\left\| L_K^{1/2} (L_{K, D_j} + \lambda_1 I + \lambda_2 V_{D_j}^T V_{D_j})^{-1} \lambda_2 V_{D_j}^T V_{D_j} f_{\lambda_1} \right\|_K \right] \\ &\lesssim E_{\mathbf{z}^{(D)}} \left[\mathcal{A}_{D_j, \lambda_1, \lambda_2, V_{D_j}}^{1/2} \right] \lambda_1^{2r-\frac{1}{2}} \lesssim \left(\frac{2\mathcal{B}_{|D_j|, \lambda_1}}{\sqrt{\lambda_1}} + 1 \right) |D|^{-\frac{2r-\frac{1}{2}}{2r+\beta}} \lesssim |D|^{-\frac{r}{2r+\beta}}, \end{aligned}$$

in which the last inequality follows from $\frac{1}{2} \leq r \leq 1$. Similar procedure implies

$$E_{\mathbf{z}|D} \left[\left\| L_K^{1/2} (L_{K,D} + \lambda_1 I + \lambda_2 V_D^T V_D)^{-1} \lambda_2 V_D^T V_D f_{\lambda_1} \right\|_K \right] \lesssim |D|^{-\frac{r}{2r+\beta}}.$$

Finally, we obtain that

$$E_{\mathbf{z}|D} \left[\left\| \mathcal{S}_3 \right\|_\rho \right] \lesssim |D|^{-\frac{r}{2r+\beta}}.$$

Now combine above three estimates for $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$, we arrive at

$$E \left[\left\| \overline{f_{\widehat{D}, \lambda_1, \lambda_2}} - f_{\widehat{D}, \lambda_1, \lambda_2} \right\|_\rho \right] = \mathcal{O}(|D|^{-\frac{r}{2r+\beta}}).$$

On the other hand, Theorem 1 has already shown

$$E \left[\left\| f_{\widehat{D}, \lambda_1, \lambda_2} - f_\rho \right\|_\rho \right] = \mathcal{O}(|D|^{-\frac{r}{2r+\beta}}).$$

Minkowski inequality finally implies

$$E \left[\left\| \overline{f_{\widehat{D}, \lambda_1, \lambda_2}} - f_\rho \right\|_\rho \right] = \mathcal{O}(|D|^{-\frac{r}{2r+\beta}}).$$

The proof is complete. ■

6 Conclusion

In this paper, we have proposed and studied the multi-penalty distribution regression algorithm. The learning rates are systematically studied via integral operator theory. The optimal rates are shown to be achievable under appropriate conditions. Meanwhile, the learning rates in the hard learning scenario of $f_\rho = L_K^r(g_\rho)$, $r \in (0, 1/2)$ for some $g_\rho \in L_{\rho_{X_\mu}}^2$ are first studied in the literature on two-stage distribution regression. The results improve the existing achievable rates in the literature. Moreover, based on the multi-penalty distribution regression scheme we provided, we propose a new distributed learning algorithm and derive optimal learning rates for it. It would be interesting to extend our methods in this work to other settings in learning theory, such as multi-kernel learning and deep neural network. The potential application domains are expected to be explored.

Acknowledgments

The authors would like to thank Professor Ding-Xuan Zhou for his valuable suggestions on this work. This work was partially supported by the Research Grants Council of the Hong Kong Special Administrative Region, China (CityU 11202819, 11203521).

References

- [1] Rastogi Abhishake, Sampath Sivananthan (2016). Multi-penalty regularization in learning theory. *Journal of Complexity*, 36, 141-165.
- [2] Mikhail Belkin, Partha Niyogi, and Vikas Sindhwani.(2006) Manifold regularization: A geometric framework for learning from labeled and unlabeled examples." *Journal of Machine Learning Research*. Nov (2006): 2399-2434.

- [3] Yasemin Altun, Smola Alex (2006). Unifying divergence minimization and statistical inference via convex duality. In International Conference on Computational Learning Theory (pp. 139-153). Springer, Berlin, Heidelberg.
- [4] Andrea Caponnetto, Ernesto De Vito (2007). Optimal rates for the regularized least-squares algorithm. *Foundations of Computational Mathematics*: 331-368.
- [5] Xiangyu Chang, Shao-Bo Lin, Ding-Xuan Zhou (2017). "Distributed semi-supervised learning with kernel ridge regression." *The Journal of Machine Learning Research* 18.1 (2017): 1493-1514.
- [6] Zhiying Fang, Zheng-Chu Guo, Ding-Xuan Zhou (2020). Optimal learning rates for distribution regression. *Journal of Complexity*, 56, 101426.
- [7] Zheng-Chu Guo, Shao-Bo Lin, Lei Shi (2019). Distributed learning with multi-penalty regularization. *Applied and Computational Harmonic Analysis*, 46(3), 478-499.
- [8] Zheng-Chu Guo, Shao-Bo Lin, Ding-Xuan Zhou (2017). Learning theory of distributed spectral algorithms. *Inverse Problems*, 33(7), 074009.
- [9] Kateřina Hlaváčková-Schindler, Valeriya Naumova, Sergiy Pereverzyev. (2017) Multi-penalty regularization for detecting relevant variables." *Recent Applications of Harmonic Analysis to Function Spaces, Differential Equations, and Data Science*. Birkhäuser, Cham, 2017. 889-916.
- [10] Junhong Lin, Alessandro Rudi, Lorenzo Rosasco, Volkan Cevher. Optimal rates for spectral algorithms with least-squares regression over Hilbert spaces. *Applied and Computational Harmonic Analysis* 48.3 (2020): 868-890.
- [11] Vladimir Koltchinskii, Ming Yuan (2010). Sparsity in multiple kernel learning." *The Annals of Statistics* 38.6 : 3660-3695.
- [12] Shao-Bo Lin, Xin Guo, Ding-Xuan Zhou (2017). Distributed learning with regularized least squares. *The Journal of Machine Learning Research*, 18(1), 3202-3232.
- [13] Shao-Bo Lin, Di Wang, Ding-Xuan Zhou (2020). Distributed Kernel Ridge Regression with Communications. *The Journal of Machine Learning Research*, in press.
- [14] Valeriya Naumova, Sergei V. Pereverzyev (2013). Multi-penalty regularization with a component-wise penalization *Inverse Problems* 29.7 : 075002.
- [15] Loucas Pillaud-Vivien, Rudi Alessandro, Bach Francis. Statistical optimality of stochastic gradient descent on hard learning problems through multiple passes. *Advances in Neural Information Processing Systems* 31 (2018).
- [16] Barnabás Póczos, Alessamdro Rinaldo, Aarti Singh, Larry Wasserman (2013). Distribution-free distribution regression.
- [17] Abhishake Rastogi (2017). Manifold learning via multi-penalty regularization. *International Journal of Artificial Intelligence and Applications (IJAIA)*, Vol.8, No.5, 2017.
- [18] Lei Shi. Distributed learning with indefinite kernels. *Analysis and Applications* 17.06 (2019): 947-975.
- [19] Fischer Simon, Ingo Steinwart. Sobolev norm learning rates for regularized least-squares algorithms. *The Journal of Machine Learning Research* 21.1 (2020): 8464-8501.

- [20] Zoltán Szabó, Arthur Gretton, Barnabás Póczos, Bharath K. Sriperumbudur, (2015). Two-stage sampled learning theory on distributions. In *Artificial Intelligence and Statistics* (pp. 948-957).
- [21] Zoltán Szabó, Bharath K. Sriperumbudur, Barnabás Póczos, Arthur Gretton (2016). Learning theory for distribution regression. *The Journal of Machine Learning Research*, 17(1), 5272-5311.
- [22] Steve Smale, Ding-Xuan Zhou (2007). Learning theory estimates via integral operators and their approximations. *Constructive approximation*, 26(2), 153-172.
- [23] Steve Smale, Ding-Xuan Zhou (2005). Shannon sampling II: Connections to learning theory. *Applied and Computational Harmonic Analysis* 19.3 : 285-302.
- [24] Steve Smale, Ding-Xuan Zhou (2004). Shannon sampling and function reconstruction from point values. *Bulletin of the American Mathematical Society* 41(3): 279-305.
- [25] Wei Wang, Shuai Lu, Heng Mao, Jin Cheng (2013). Multi-parameter Tikhonov regularization with the l^0 sparsity constraint. *Inverse Problems*, 29(6), 065018.