

Family of mean-mixtures of multivariate normal distributions: properties, inference and assessment of multivariate skewness

Me'raj Abdi^a, Mohsen Madadi^{a,*}, Narayanaswamy Balakrishnan^b, Ahad Jamalizadeh^a

^aDepartment of Statistics, Faculty of Mathematics and Computer, Shahid Bahonar University of Kerman, Kerman, Iran

^bDepartment of Mathematics and Statistics, McMaster University, Hamilton, Ontario, Canada

Abstract

In this paper, a new mixture family of multivariate normal distributions, formed by mixing multivariate normal distribution and a skewed distribution, is constructed. Some properties of this family, such as characteristic function, moment generating function, and the first four moments are derived. The distributions of affine transformations and canonical forms of the model are also derived. An EM-type algorithm is developed for the maximum likelihood estimation of model parameters. Some special cases of the family, using standard gamma and standard exponential mixture distributions, denoted by $MMNG$ and $MMNE$, respectively, are considered. For the proposed family of distributions, different multivariate measures of skewness are computed. In order to examine the performance of the developed estimation method, some simulation studies are carried out to show that the maximum likelihood estimates do provide a good performance. For different choices of parameters of $MMNE$ distribution, several multivariate measures of skewness are computed and compared. Because some measures of skewness are scalar and some are vectors, in order to evaluate them properly, a simulation study is carried out to determine the power of tests, based on sample versions of skewness measures as test statistics for testing the fit of the $MMNE$ distribution. Finally, two real data sets are used to illustrate the usefulness of the proposed model and the associated inferential methods.

Keywords: Canonical form, EM algorithm, Mean mixtures of normal distribution, Moments, Multivariate measures of skewness.

2010 MSC: 60E07, 60E10, 62H05, 62H10 and 62H12.

1. Introduction

The multivariate normal distribution plays a fundamental role in statistical analyses and applications. One of the most basic properties of the normal distribution is the symmetry of its density function. However, in practice, data sets do not follow the normal distribution or even possess symmetry, and for this reason, researchers search for new distributions to fit data with different features allowing flexibility in skewness, kurtosis, tails and multimodality; see for example, [18, 21]. Several new families of distributions have been introduced for modeling skewed data, including normal distribution as a special case. One such prominent distribution in the univariate case is the skew normal (SN) distribution due to [6, 7]. The multivariate version of the SN distribution has been introduced in [10]. This distribution has found diverse applications such as portfolio optimization concepts and risk measurement indices in financial markets; see [15]. A complete set of extensions of multivariate SN distributions can be found in [8, 9]. [12] calculated and compared several different measures of skewness for the multivariate SN distribution. [13] proposed a test to assess if a sample comes from a multivariate SN distribution. Here, we use $\phi_p(\cdot; \boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $\Phi_p(\cdot; \boldsymbol{\mu}, \boldsymbol{\Sigma})$ to denote the probability density function (PDF) and the cumulative distribution function (CDF) of the p -variate normal distribution, with mean $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$, respectively, and also $\phi(\cdot)$ and $\Phi(\cdot)$ to denote the PDF and CDF of the univariate standard normal distribution, respectively.

*Corresponding author. Email address: madadi@uk.ac.ir

Email addresses: me.abdi.z@gmail.com (M. Abdi), bala@mcmaster.ca (N. Balakrishnan) and a.jamalizadeh@uk.ac.ir (A. Jamalizadeh).

From [9] and [10], a p -dimensional random vector \mathbf{Y} follows a multivariate \mathcal{SN} distribution if it has the PDF

$$f(\mathbf{y}) = 2\phi_p(\mathbf{y}; \boldsymbol{\xi}, \boldsymbol{\Omega}) \Phi \left(\frac{\boldsymbol{\delta}^\top \bar{\boldsymbol{\Omega}}^{-1} \boldsymbol{\omega}^{-1} (\mathbf{y} - \boldsymbol{\xi})}{\sqrt{1 - \boldsymbol{\delta}^\top \bar{\boldsymbol{\Omega}}^{-1} \boldsymbol{\delta}}} \right),$$

with stochastic representation

$$\mathbf{Y} \stackrel{d}{=} \boldsymbol{\xi} + \boldsymbol{\omega} (\delta U + \mathbf{Z}), \quad (1)$$

where $\stackrel{d}{=}$ stands for equality in distribution, $\boldsymbol{\xi} \in \mathbb{R}^p$, $\mathbf{Z} \sim \mathcal{N}_p(\mathbf{0}, \bar{\boldsymbol{\Omega}} - \delta\delta^\top)$ and univariate random variable U has a standard normal distribution within the truncated interval $(0, \infty)$, independently of \mathbf{Z} . Truncated normal distribution in the interval $(0, \infty)$ with parameters (a, b) is denoted by $\mathcal{TN}(a, b, (0, \infty))$. The vector $\boldsymbol{\delta} = (\delta_1, \dots, \delta_p)^\top$ is the skewness parameter vector, such that $-1 < \delta_i < 1$, for $i \in \{1, \dots, p\}$. The matrix $\boldsymbol{\omega} = \text{diag}(\omega_1, \dots, \omega_p) = (\boldsymbol{\Omega} \odot \mathbf{I}_p)^{1/2} > 0$ is a diagonal matrix formed by the standard deviations of $\boldsymbol{\Omega}$ and $\bar{\boldsymbol{\Omega}} = \boldsymbol{\omega} \bar{\boldsymbol{\Omega}} \boldsymbol{\omega}$. Here, \mathbf{I}_p is identity matrix of size p . The Hadamard product of matrices $\mathbf{A} = (a_{ij}) : m \times n$ and $\mathbf{B} = (b_{ij}) : m \times n$ is given by the $m \times n$ matrix $\mathbf{A} \odot \mathbf{B} = (a_{ij} b_{ij})$. In the stochastic representation in (1), positive definite matrices $\boldsymbol{\Omega}$ and $\bar{\boldsymbol{\Omega}}$ are covariance and correlation matrices, respectively. The parameters $\boldsymbol{\xi}$, $\boldsymbol{\omega}$ and $\boldsymbol{\delta}$ are the location, scale and skewness parameters, respectively.

Upon using the stochastic representation in (1), a general new family of mixture distributions of multivariate normal distribution can be introduced based on arbitrary random variable U . A p -dimensional random vector \mathbf{Y} follows a multivariate mean mixture of normal (\mathcal{MMN}) distribution if, in (1), U is an arbitrary random variable with CDF $H(\cdot; \boldsymbol{\nu})$, independently of \mathbf{Z} , indexed by the parameter $\boldsymbol{\nu} = (\nu_1, \dots, \nu_p)^\top$. Then, we say that \mathbf{Y} has a mean mixture of multivariate normal (\mathcal{MMN}) distribution, and denote it by $\mathbf{Y} \sim \mathcal{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$.

[37] presented a new family of distributions as a mixture of normal distribution and studied its properties in the univariate and multivariate cases. These authors defined a p -dimensional random vector \mathbf{Y} to have a multivariate mean mixture of normal distribution if it has the stochastic representation $\mathbf{Y} \stackrel{d}{=} \boldsymbol{\xi} + \delta U + \mathbf{Z}$, where $\mathbf{Z} \sim \mathcal{N}_p(\mathbf{0}, \boldsymbol{\Omega})$ and U is an arbitrary positive random variable with CDF $H(\cdot; \boldsymbol{\nu})$ independently of \mathbf{Z} , indexed by the parameter vectors $\boldsymbol{\nu} = (\nu_1, \dots, \nu_p)^\top$ and $\boldsymbol{\delta} = (\delta_1, \dots, \delta_p)^\top \in \mathbb{R}^p$. The stochastic representation used by [37] is along the lines of the stochastic representation of the restricted multivariate \mathcal{SN} distribution (see [8]), but in this work, we use a different stochastic representation in (1). [37] examined some properties of this family in the univariate case for general U , and also two specific cases of the family. In the present work, we consider the multivariate form of this family and study its properties.

In (1), if the random variable U is a skewed random variable, then the p -dimensional vector \mathbf{Y} will also be skewed. In the \mathcal{MMN} family, skewness can be regulated through the parameter $\boldsymbol{\delta}$. If in (1) $\boldsymbol{\delta} = \mathbf{0}$, the \mathcal{MMN} family is reduced to the multivariate normal distribution. The extended form of the \mathcal{SN} distribution is obtained from (1) when U is distributed as $\mathcal{N}(0, 1)$ variable truncated below $-\tau$ instead of 0, for some constant τ . The representation in (1) means that the \mathcal{MMN} distribution is a “mean mixture” of the multivariate normal distribution when the mixing random variable is U . Specifically, we have the following hierarchical representation for the \mathcal{MMN} distribution:

$$\mathbf{Y} | (U = u) \sim \mathcal{N}_p(\boldsymbol{\xi} + \boldsymbol{\omega} \delta u, \boldsymbol{\Omega} - \boldsymbol{\omega} \delta \delta^\top \boldsymbol{\omega}), \quad U \sim H(\cdot; \boldsymbol{\nu}). \quad (2)$$

According to (2), in the \mathcal{MMN} model, just the mean parameter is mixed with arbitrary random variable U , and so this class can not be obtained from the Normal Mean-Variance Mixture (\mathcal{NMVM}) family. The family of multivariate \mathcal{NMVM} distributions, originated by [14], is another extension of multivariate normal distribution, with a skewness parameter $\boldsymbol{\delta} \in \mathbb{R}^p$. A p -dimensional random vector \mathbf{Y} is said to have a multivariate \mathcal{NMVM} distribution if it has the representation

$$\mathbf{Y} = \boldsymbol{\xi} + \delta U + \sqrt{U} \mathbf{Z}, \quad (3)$$

where $\mathbf{Z} \sim \mathcal{N}_p(\mathbf{0}, \boldsymbol{\Omega})$ and U is a positive random variable and the CDF of U , $H(\cdot; \boldsymbol{\nu})$, is the mean-variance mixing distribution. Both families of distributions in (1) and (3) include the multivariate normal distribution as a special case

and can be used for modeling data possessing skewness. In (3), both mean and variance are mixed with the same positive random variable U , while in (1) just the mean parameter is mixed with U , however, the class in (1) cannot be obtained from the class in (3).

Besides, skewness is a feature commonly found in the returns of some financial assets. For more information on applications of skewed distributions in finance theory, one may refer to [2]. In the presence of skewness in asset returns, the SN and skew- t (ST) distributions have been found to be useful models in both theoretical and empirical work. Their parametrization is parsimonious, and they are mathematically tractable, and in financial applications, the distributions are interpretable in terms of the efficient market hypothesis. Furthermore, they lead to theoretical results that are useful for portfolio selection and asset pricing. In actuarial science, the presence of skewness in insurance claims data is the primary motivation for using SN distribution and its extensions. In this regard, the MMN family that is developed here will also prove useful in finance, insurance science, and other applied fields.

[40] proposed that the n -dimensional vector of returns on financial assets should be represented as $\mathbf{X} = \mathbf{U} + \lambda V$. The n -dimensional vector \mathbf{U} is assumed to have a multivariate elliptically symmetric distribution, independently of the non-negative univariate random variable V , having an unspecified skewed distribution. The vector λ , whose elements may take any real values, induces skewness in the return of individual assets. [3] have described multivariate versions of the normal-exponential and normal-gamma distributions. Both of them are specific cases of the model proposed in [40]. [1] and [3] used the representation in [40], with specific choices of \mathbf{U} and V , and introduced a number of distributions such as SN , extended SN , ST , normal-exponential, and normal-gamma, and investigated the corresponding distributions and their applications in capital pricing, return on financial assets and portfolio selection.

In this paper, with an arbitrary random variable U for the MMN family with stochastic representation in (1), basic distributional properties of the class such as the characteristic function (CF), the moment generating function (MGF), the first four moments of the model, distributions of linear and affine transformations, the canonical form of the family and the mode of the model are derived in general. Also, the maximum likelihood estimation of the parameters by using an EM-type algorithm is discussed, and then different measures of multivariate skewness are obtained. The special cases when U has standard gamma and standard exponential distributions, with the corresponding distributions denoted by $MMNG$ and $MMNE$ distributions, respectively, are studied in detail. For the $MMNG$ distribution, in addition to all the above basic properties of the distribution the infinitely divisibility of the model is also discussed. For the $MMNE$ distribution, the basic properties of the distribution as well as log-concavity of the model are discussed. The maximum likelihood estimates of the parameters of the $MMNE$ distribution are evaluated using the bias and the mean square error by means of a simulation study. Moreover, various multivariate measures of skewness are computed and compared. Finally, for two real data sets, the $MMNE$ distribution is fitted and compared with the SN and ST distributions in terms of log-likelihood value as well as AIC and BIC criteria.

2. Model and Properties

In this section, some basic properties of the model are studied. From (1), if U has a PDF $h(\cdot; \nu)$, an integral form of the PDF of $\mathbf{Y} \sim MMN_p(\xi, \Omega, \delta; H)$ can be obtained as

$$f_{MMN_p}(\mathbf{y}; \xi, \Omega, \delta, \nu) = \int_{-\infty}^{+\infty} \phi_p(\mathbf{y}; \xi + \omega \delta u, \Omega - \omega \delta \delta^\top \omega) dH(u; \nu) = \int_{-\infty}^{+\infty} \phi_p(\mathbf{y}; \xi + \omega \delta u, \Omega - \omega \delta \delta^\top \omega) h(u; \nu) du. \quad (4)$$

We now present some theorems and lemmas with regard to different properties of these distributions, proofs of which are presented in Appendix A.

Remark 1. We can introduce the normalized MMN distribution through the transformation $\mathbf{X} = \omega^{-1}(\mathbf{Y} - \xi)$. It is immediate that the stochastic representation of $\mathbf{X} = \delta U + \mathbf{Z}$ has the hierarchical representation $\mathbf{X}|(U = u) \sim \mathcal{N}_p(\delta u, \Omega - \delta \delta^\top)$ and $U \sim H(\cdot; \nu)$. Then, we say that \mathbf{X} has a normalized mean mixture of multivariate normal distributions, and denote it by $\mathbf{X} \sim MMN_p(\mathbf{0}, \Omega, \delta; H)$.

Lemma 1. If $\mathbf{Y} \sim MMN_p(\xi, \Omega, \delta; H)$, the CF and MGF of \mathbf{Y} are as follows:

$$C_{\mathbf{Y}}(\mathbf{t}) = e^{it^\top \xi + \frac{1}{2} \mathbf{t}^\top \Sigma_{\mathbf{Y}} \mathbf{t}} C_U(it^\top \omega \delta; \nu), \quad M_{\mathbf{Y}}(\mathbf{t}) = e^{t^\top \xi + \frac{1}{2} \mathbf{t}^\top \Sigma_{\mathbf{Y}} \mathbf{t}} M_U(t^\top \omega \delta; \nu), \quad (5)$$

respectively, where $i = \sqrt{-1}$, $\Sigma_Y = \Omega - \omega\delta\delta^\top\omega$, and $C_U(\cdot; \nu) = C_U(\cdot)$ and $M_U(\cdot; \nu) = M_U(\cdot)$ are the CF and MGF of U , respectively.

Moreover, if $\mathbf{X} \sim \mathcal{MMN}_p(\mathbf{0}, \overline{\Omega}, \delta; H)$, the CF and MGF of \mathbf{X} are

$$C_{\mathbf{X}}(\mathbf{t}) = e^{\frac{1}{2}\mathbf{t}^\top \Sigma_{\mathbf{X}} \mathbf{t}} C_U(i\mathbf{t}^\top \delta; \nu), \quad M_{\mathbf{X}}(\mathbf{t}) = e^{\frac{1}{2}\mathbf{t}^\top \Sigma_{\mathbf{X}} \mathbf{t}} M_U(\mathbf{t}^\top \delta; \nu), \quad (6)$$

respectively, where $\Sigma_{\mathbf{X}} = \overline{\Omega} - \delta\delta^\top$. The first four moments of \mathbf{X} , presented in the following lemma, are derived by using the partial derivatives of MGF of normalized MMN distribution, and these, in turn, can be used to obtain the first four moments of \mathbf{Y} .

Lemma 2. Suppose $\mathbf{X} \sim \mathcal{MMN}_p(\mathbf{0}, \overline{\Omega}, \delta; H)$. Then, the first four moments of \mathbf{X} are as follows:

$$M_1(\mathbf{X}) = M_1^{\mathbf{X}} = E[U]\delta, \quad (7)$$

$$M_2(\mathbf{X}) = M_2^{\mathbf{X}} = \Sigma_{\mathbf{X}} + E[U^2](\delta \otimes \delta^\top), \quad (8)$$

$$M_3(\mathbf{X}) = M_3^{\mathbf{X}} = E[U] \left\{ \delta \otimes \Sigma_{\mathbf{X}} + \text{vec}(\Sigma_{\mathbf{X}}) \delta^\top + (\mathbf{I}_p \otimes \delta) \Sigma_{\mathbf{X}} \right\} + E[U^3](\mathbf{I}_p \otimes \delta)(\delta \otimes \delta^\top), \quad (9)$$

$$M_4(\mathbf{X}) = M_4^{\mathbf{X}} = (\mathbf{I}_{p^2} + \mathbf{U}_{p,p})(\Sigma_{\mathbf{X}} \otimes \Sigma_{\mathbf{X}}) + \text{vec}(\Sigma_{\mathbf{X}})(\text{vec}(\Sigma_{\mathbf{X}}))^\top + E[U^2] \left[\delta \otimes \delta^\top \otimes \Sigma_{\mathbf{X}} + \delta \otimes \Sigma_{\mathbf{X}} \otimes \delta^\top + \Sigma_{\mathbf{X}} \otimes \delta \otimes \delta^\top + \delta^\top \otimes \Sigma_{\mathbf{X}} \otimes \delta + \delta^\top \otimes \text{vec}(\Sigma_{\mathbf{X}}) \otimes \delta^\top + (\delta \otimes \delta)(\text{vec}(\Sigma_{\mathbf{X}}))^\top \right] + E[U^4] \delta \delta^\top \otimes \delta \delta^\top, \quad (10)$$

where $E(U^k) = M_U^{(k)}(0)$, with $M_U^{(k)}(\cdot)$ being the k -th derivative of $M_U(t)$ with respect to t .

The Kronecker product of matrices $\mathbf{A} = (a_{ij}) : m \times n$ and $\mathbf{B} = (b_{ij}) : p \times q$ is a $mp \times nq$ matrix $\mathbf{A} \otimes \mathbf{B} = (a_{ij}\mathbf{B})$. A matrix $\mathbf{A} = (\mathbf{a}_1, \dots, \mathbf{a}_n) : m \times n$ with columns $\mathbf{a}_1, \dots, \mathbf{a}_n$ is sometimes written as a vector and called $\text{vec}(\mathbf{A})$, defined by $\text{vec}(\mathbf{A}) = (\mathbf{a}_1^\top, \dots, \mathbf{a}_n^\top)^\top$. The matrix $\mathbf{U}_{p,p}$ is the permutation matrix (commutation matrix) associated with a $p \times p$ matrix (its size is $p^2 \times p^2$). For details about Kronecker product, permutation matrix and its properties, see [23] and [39]. We extend the results of Lemma 2, using the stochastic representation in (1), to incorporate location and scale parameters, ξ and ω , through the transformation $\mathbf{Y} = \xi + \omega\mathbf{X}$.

Theorem 1. If $\mathbf{Y} \sim \mathcal{MMN}_p(\xi, \Omega, \delta; H)$, then its first four moments are as follows:

$$M_1(\mathbf{Y}) = \xi + \omega M_1^{\mathbf{X}}, \quad (11)$$

$$M_2(\mathbf{Y}) = \xi \otimes \xi^\top + \xi \otimes (\omega M_1^{\mathbf{X}})^\top + \omega M_1^{\mathbf{X}} \otimes \xi^\top + \omega M_2^{\mathbf{X}} \omega, \quad (12)$$

$$M_3(\mathbf{Y}) = \xi \xi^\top \otimes \xi + \xi \xi^\top \otimes (\omega M_1^{\mathbf{X}}) + \xi (\omega M_1^{\mathbf{X}})^\top \otimes \xi + (\omega M_1^{\mathbf{X}}) \otimes \xi \xi^\top + (\omega M_2^{\mathbf{X}} \omega) \otimes \xi + \xi \otimes (\omega M_2^{\mathbf{X}} \omega) + (\omega \otimes \omega) \text{vec}(M_2^{\mathbf{X}}) \otimes \xi^\top + (\omega \otimes \omega) M_3^{\mathbf{X}} \omega, \quad (13)$$

$$M_4(\mathbf{Y}) = \xi \xi^\top \otimes \xi \xi^\top + \xi \xi^\top \otimes \xi (\omega M_1^{\mathbf{X}})^\top + \xi \xi^\top \otimes (\omega M_1^{\mathbf{X}}) \xi^\top + \xi \xi^\top \otimes (\omega M_2^{\mathbf{X}} \omega) + \xi (\omega M_1^{\mathbf{X}})^\top \otimes \xi \xi^\top + (\xi \otimes \xi) (\text{vec}(M_2^{\mathbf{X}}))^\top (\omega \otimes \omega) + \xi \otimes (\omega M_2^{\mathbf{X}} \omega) \otimes \xi^\top + \xi \otimes \omega (M_3^{\mathbf{X}})^\top (\omega \otimes \omega) + (\omega M_1^{\mathbf{X}}) \xi^\top \otimes \xi \xi^\top + \xi^\top \otimes (\omega M_2^{\mathbf{X}} \omega) \otimes \xi + \xi^\top \otimes (\omega \otimes \omega) \text{vec}(M_2^{\mathbf{X}}) \otimes \xi^\top + \xi^\top \otimes (\omega \otimes \omega) M_3^{\mathbf{X}} \omega + (\omega M_2^{\mathbf{X}} \omega) \otimes \xi \xi^\top + \omega (M_3^{\mathbf{X}})^\top (\omega \otimes \omega) \otimes \xi + (\omega \otimes \omega) M_3^{\mathbf{X}} \omega \otimes \xi^\top + (\omega \otimes \omega) M_4^{\mathbf{X}} (\omega \otimes \omega). \quad (14)$$

From the above expressions, we can obtain the mean vector and covariance matrix of $\mathcal{MMN}_p(\xi, \Omega, \delta; H)$ family as $E(\mathbf{Y}) = \xi + E(U)\omega\delta$ and $\text{var}(\mathbf{Y}) = \Omega + (\text{var}(U) - 1)\omega\delta\delta^\top\omega$. Multiplication of $M_{\mathbf{X}}(\mathbf{t})$ by the MGF of the $\mathcal{N}_p(\mu, \Sigma)$ distribution, $\exp(\mathbf{t}^\top \mu + \mathbf{t}^\top \Sigma \mathbf{t}/2)$, is still a function of type $M_{\mathbf{X}}(\mathbf{t})$, and we thus obtain the following result.

Theorem 2. If $\mathbf{Y}_1 \sim \mathcal{MMN}_p(\xi, \Omega, \delta; H)$ and $\mathbf{Y}_2 \sim \mathcal{N}_p(\mu, \Sigma)$ are independent variables, then $\mathbf{Y} = \mathbf{Y}_1 + \mathbf{Y}_2 \sim \mathcal{MMN}_p(\xi_Y, \Omega_Y, \delta_Y; H)$, where $\xi_Y = \xi + \mu$, $\Omega_Y = \Omega + \Sigma$, and $\delta_Y = \omega_Y^{-1}\omega\delta$, with $\omega_Y = (\Omega_Y \odot \mathbf{I}_p)^{1/2}$.

From the MGF's $M_{\mathbf{X}}(\mathbf{t})$ and $M_{\mathbf{Y}}(\mathbf{t})$, it is clear that the family of MMN distributions is closed under affine transformations, as given in the following results.

Theorem 3. If $\mathbf{X} \sim \text{MMN}_p(\mathbf{0}, \overline{\boldsymbol{\Omega}}, \boldsymbol{\delta}; H)$ and \mathbf{A} is a non-singular $p \times p$ matrix such that $\text{diag}(\mathbf{A}^\top \overline{\boldsymbol{\Omega}} \mathbf{A}) = \mathbf{I}_p$, that is, $\mathbf{A}^\top \overline{\boldsymbol{\Omega}} \mathbf{A}$ is a correlation matrix, then $\mathbf{A}^\top \mathbf{X} \sim \text{MMN}_p(\mathbf{0}, \mathbf{A}^\top \overline{\boldsymbol{\Omega}} \mathbf{A}, \mathbf{A}^\top \boldsymbol{\delta}; H)$.

Theorem 4. If $\mathbf{Y} \sim \text{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$, \mathbf{A} is a full-rank $p \times h$ matrix, with $h \leq p$, and $\mathbf{c} \in \mathbb{R}^h$, then $\mathbf{T} = \mathbf{c} + \mathbf{A}^\top \mathbf{Y} \sim \text{MMN}_h(\boldsymbol{\xi}_\mathbf{T}, \boldsymbol{\Omega}_\mathbf{T}, \boldsymbol{\delta}_\mathbf{T}; H)$, where $\boldsymbol{\xi}_\mathbf{T} = \mathbf{c} + \mathbf{A}^\top \boldsymbol{\xi}$, $\boldsymbol{\Omega}_\mathbf{T} = \mathbf{A}^\top \boldsymbol{\Omega} \mathbf{A}$, and $\boldsymbol{\delta}_\mathbf{T} = \boldsymbol{\omega}_\mathbf{T}^{-1} \mathbf{A}^\top \boldsymbol{\omega} \boldsymbol{\delta}$, with $\boldsymbol{\omega}_\mathbf{T} = (\boldsymbol{\Omega}_\mathbf{T} \odot \mathbf{I}_h)^{1/2}$.

As in the case of multivariate \mathcal{SN} distribution, (see [9]), it can be shown that, if the random vector \mathbf{Y} is partitioned into a number of random vectors, the independence occurs between its components when at least one component follows the MMN distribution and the others have normal distribution, that is, the independence between components occurs when only one component of the skewness parameter $\boldsymbol{\delta}$ is non-zero and all others are zero. Without loss of generality, from here on, it is assumed that the first element of $\boldsymbol{\delta}$ is non-zero. We now focus on a specific type of linear transformation of the MMN variable, having special relevance for theoretical developments but also to some extent for practical reasons.

Theorem 5. For a given variable $\mathbf{Y} \sim \text{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$, there exists a linear transformation $\mathbf{Z}^* = \mathbf{A}_*(\mathbf{Y} - \boldsymbol{\xi})$ such that $\mathbf{Z}^* \sim \text{MMN}_p(\mathbf{0}, \mathbf{I}_p, \boldsymbol{\delta}_{\mathbf{Z}^*}; H)$, where at most one component of $\boldsymbol{\delta}_{\mathbf{Z}^*}$ is not zero, and $\boldsymbol{\delta}_{\mathbf{Z}^*} = (\delta_*, 0, \dots, 0)^\top$ with $\delta_* = \left(\boldsymbol{\delta}^\top \overline{\boldsymbol{\Omega}}^{-1} \boldsymbol{\delta}\right)^{1/2}$.

The variable \mathbf{Z}^* , which we shall sometimes refer to as a canonical variate, consists of p independent components. The joint density is given by the product of $p - 1$ standard normal densities and at most one non-Gaussian component $\text{MMN}_1(0, 1, \delta_*; H)$; that is, the density of \mathbf{Z}^* is

$$f_{\mathbf{Z}^*}(\mathbf{z}) = f_{Z_1^*}(z_1) \prod_{i=2}^p \phi(z_i), \quad (15)$$

where $Z_1^* \sim \text{MMN}_1(0, 1, \delta_*; H)$ (for univariate MMN distribution, see [37]). Although Theorem 5 ensures that it is possible to obtain a canonical form, note that in general there are many possible ways to do so, but it is not obvious how to achieve the canonical form in practice. To find the appropriate \mathbf{A}_* in the linear transformation $\mathbf{Z}^* = \mathbf{A}_*(\mathbf{Y} - \boldsymbol{\xi})$, it is sufficient to find a \mathbf{A}_* satisfying the following two conditions: $\mathbf{A}_*^\top \boldsymbol{\Omega} \mathbf{A}_* = \mathbf{I}_p$ and $\mathbf{A}_*^\top \boldsymbol{\omega} \boldsymbol{\delta} = \boldsymbol{\delta}_{\mathbf{Z}^*} = (\delta_*, 0, \dots, 0)^\top$. The canonical form facilitates the computation of the mode of the distribution and the multivariate coefficients of skewness.

Theorem 6. If $\mathbf{Y} \sim \text{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$, the mode of \mathbf{Y} is $\mathbf{M}_0 = \boldsymbol{\xi} + \frac{m_0^*}{\delta_*} \boldsymbol{\omega} \boldsymbol{\delta}$, where $\delta_* = \left(\boldsymbol{\delta}^\top \overline{\boldsymbol{\Omega}}^{-1} \boldsymbol{\delta}\right)^{1/2}$ and m_0^* is the mode of the univariate $\text{MMN}_1(0, 1, \delta_*; H)$ distribution.

3. Likelihood Estimation through EM Algorithm

For obtaining the maximum likelihood estimates of all the parameters of $\text{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$, we propose an EM-type algorithm as in [35]. Let $\mathbf{Y} = (\mathbf{Y}_1, \dots, \mathbf{Y}_n)^\top$ be a random sample of size n from a $\text{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$ distribution. Consider the stochastic representation in (1) for $\mathbf{Y}_i, i \in \{1, \dots, n\}$. Following the EM algorithm, let $(\mathbf{Y}_i, U_i), i \in \{1, \dots, n\}$, be the complete data, where \mathbf{Y}_i is the observed data and U_i is considered as missing data. Let $\boldsymbol{\theta} = (\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}, \boldsymbol{\nu})$. Using (2), the distribution of \mathbf{Y}_i , for $i \in \{1, \dots, n\}$, can be written hierarchically as

$$\mathbf{Y}_i | (U_i = u_i) \sim \mathcal{N}_p(\boldsymbol{\xi} + \boldsymbol{\omega} \boldsymbol{\delta} u_i, \boldsymbol{\Sigma}_\mathbf{Y}), \quad U_i \stackrel{iid}{\sim} H(\cdot; \boldsymbol{\nu}),$$

where $\stackrel{iid}{\sim}$ denotes independence of random variables and $\boldsymbol{\Sigma}_\mathbf{Y} = \boldsymbol{\Omega} - \boldsymbol{\omega} \boldsymbol{\delta} \boldsymbol{\delta}^\top \boldsymbol{\omega}$. Let $\mathbf{y} = (\mathbf{y}_1^\top, \dots, \mathbf{y}_n^\top)^\top$, where \mathbf{y}_i is a realization of $\text{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$. Because

$$f(\mathbf{y}_i, u_i) = f(\mathbf{y}_i | u_i) h(u_i; \boldsymbol{\nu}), \quad (16)$$

the complete data log-likelihood function, ignoring additive constants, is obtained from (16) as

$$\ell_c(\boldsymbol{\theta}) = -\frac{n}{2} \ln |\boldsymbol{\Sigma}_\mathbf{Y}| - \frac{1}{2} \sum_{i=1}^n (\mathbf{y}_i - \boldsymbol{\xi})^\top \boldsymbol{\Sigma}_\mathbf{Y}^{-1} (\mathbf{y}_i - \boldsymbol{\xi}) + \boldsymbol{\alpha}^\top \boldsymbol{\Sigma}_\mathbf{Y}^{-1} \sum_{i=1}^n u_i (\mathbf{y}_i - \boldsymbol{\xi}) - \frac{1}{2} \boldsymbol{\alpha}^\top \boldsymbol{\Sigma}_\mathbf{Y}^{-1} \boldsymbol{\alpha} \sum_{i=1}^n u_i^2 + \sum_{i=1}^n \ln h(u_i; \boldsymbol{\nu}),$$

where $\alpha = \omega\delta$. Let us set

$$\widehat{E}_{i1}^{(k)} = E\left[U_i | \mathbf{Y}_i = \mathbf{y}_i, \widehat{\boldsymbol{\theta}}^{(k)}\right], \quad \widehat{E}_{i2}^{(k)} = E\left[U_i^2 | \mathbf{Y}_i = \mathbf{y}_i, \widehat{\boldsymbol{\theta}}^{(k)}\right], \quad (17)$$

where $\widehat{\boldsymbol{\theta}}^{(k)} = (\widehat{\boldsymbol{\xi}}^{(k)}, \widehat{\boldsymbol{\Omega}}^{(k)}, \widehat{\boldsymbol{\delta}}^{(k)}, \widehat{\boldsymbol{\nu}}^{(k)})$. After some simple algebra and using (17), the expectation with respect to U conditional on \mathbf{Y} , of the complete log-likelihood function, has the form

$$\begin{aligned} Q(\boldsymbol{\theta} | \widehat{\boldsymbol{\theta}}^{(k)}) &= \frac{n}{2} \ln |\boldsymbol{\Sigma}_{\mathbf{Y}}^{-1}| - \frac{1}{2} \sum_{i=1}^n (\mathbf{y}_i - \boldsymbol{\xi})^\top \boldsymbol{\Sigma}_{\mathbf{Y}}^{-1} (\mathbf{y}_i - \boldsymbol{\xi}) + \sum_{i=1}^n \text{tr} \left[\boldsymbol{\Sigma}_{\mathbf{Y}}^{-1} (\mathbf{y}_i - \boldsymbol{\xi}) \alpha^\top \right] \widehat{E}_{i1}^{(k)} \\ &\quad - \frac{1}{2} \text{tr} \left[\boldsymbol{\Sigma}_{\mathbf{Y}}^{-1} \alpha \alpha^\top \right] \sum_{i=1}^n \widehat{E}_{i2}^{(k)} + \sum_{i=1}^n E \left[\ln h(u_i; \nu) | \mathbf{Y}_i = \mathbf{y}_i, \widehat{\boldsymbol{\theta}}^{(k)} \right], \end{aligned} \quad (18)$$

where $\bar{\mathbf{y}} = \frac{1}{n} \sum_{i=1}^n \mathbf{y}_i$ is the sample mean vector. The EM-type algorithm for the ML estimation of $\boldsymbol{\theta} = (\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}, \nu)$ then proceeds as follows:

Algorithm 1. Based on the initial value of $\boldsymbol{\theta}^{(0)} = (\boldsymbol{\xi}^{(0)}, \boldsymbol{\Omega}^{(0)}, \boldsymbol{\delta}^{(0)}, \nu^{(0)})$, the EM-type algorithm iterates between the following E-step and M-step:

E-step: Given the estimates of model parameters at the k -th iteration, say $\boldsymbol{\theta} = \widehat{\boldsymbol{\theta}}^{(k)}$, compute $\widehat{E}_{i1}^{(k)}$ and $\widehat{E}_{i2}^{(k)}$, for $i \in \{1, 2, \dots, n\}$;

M-step 1: Maximization of (18) over parameters $\boldsymbol{\xi}$, α and $\boldsymbol{\Sigma}_{\mathbf{Y}}$ leads to the following closed-form expressions:

$$\begin{aligned} \widehat{\alpha}^{(k+1)} &= \frac{\sum_{i=1}^n \mathbf{y}_i \widehat{E}_{i1}^{(k)} - \bar{\mathbf{y}} \sum_{i=1}^n \widehat{E}_{i1}^{(k)}}{\sum_{i=1}^n \widehat{E}_{i2}^{(k)} - \frac{1}{n} \left(\sum_{i=1}^n \widehat{E}_{i1}^{(k)} \right)^2}, \quad \widehat{\boldsymbol{\xi}}^{(k+1)} = \bar{\mathbf{y}} - \frac{\widehat{\alpha}^{(k+1)}}{n} \sum_{i=1}^n \widehat{E}_{i1}^{(k)}, \\ \widehat{\boldsymbol{\Sigma}}_{\mathbf{Y}}^{(k+1)} &= \frac{1}{n} \sum_{i=1}^n \left(\mathbf{y}_i - \widehat{\boldsymbol{\xi}}^{(k+1)} \right) \left(\mathbf{y}_i - \widehat{\boldsymbol{\xi}}^{(k+1)} \right)^\top - \frac{2}{n} \sum_{i=1}^n \widehat{E}_{i1}^{(k)} \left(\mathbf{y}_i - \widehat{\boldsymbol{\xi}}^{(k+1)} \right) \widehat{\alpha}^{(k+1)\top} + \frac{1}{n} \widehat{\alpha}^{(k+1)} \widehat{\alpha}^{(k+1)\top} \sum_{i=1}^n \widehat{E}_{i2}^{(k)}. \end{aligned}$$

Therefore, we can compute $\widehat{\boldsymbol{\Omega}}^{(k+1)} = \widehat{\boldsymbol{\Sigma}}_{\mathbf{Y}}^{(k+1)} + \widehat{\alpha}^{(k+1)} \widehat{\alpha}^{(k+1)\top}$ and $\widehat{\boldsymbol{\delta}}^{(k+1)} = \widehat{\omega}^{(k+1)-1} \widehat{\alpha}^{(k+1)}$, where $\widehat{\omega} = (\widehat{\boldsymbol{\Omega}} \odot \mathbf{I}_p)^{1/2}$.

M-step 2: The update of $\widehat{\boldsymbol{\nu}}^{(k)}$ depends on the chosen distribution for U , and is obtained as

$$\widehat{\boldsymbol{\nu}}^{(k+1)} = \arg \max_{\boldsymbol{\nu}} \sum_{i=1}^n E \left[\ln h(u_i; \nu) | \mathbf{Y}_i = \mathbf{y}_i, \widehat{\boldsymbol{\theta}}^{(k)} \right].$$

Updating of $\widehat{\boldsymbol{\nu}}^{(k)}$ is strongly related to the form of $h(u_i; \nu)$. If the conditional expectation $E \left[\ln h(u_i; \nu) | \mathbf{Y}_i = \mathbf{y}_i, \widehat{\boldsymbol{\theta}}^{(k)} \right]$ is difficult to evaluate, one may resort to maximizing the restricted actual log-likelihood function, as follows:

Modified M-step 2: (Liu and Rubin [30]) Update $\widehat{\boldsymbol{\nu}}^{(k)}$ by $\widehat{\boldsymbol{\nu}}^{(k+1)} = \arg \max_{\boldsymbol{\nu}} \sum_{i=1}^n \ln f_{\text{MMN}_p}(\mathbf{y}_i; \widehat{\boldsymbol{\xi}}^{(k+1)}, \widehat{\boldsymbol{\Omega}}^{(k+1)}, \widehat{\boldsymbol{\delta}}^{(k+1)}, \nu)$. The above algorithm iterates between the E-step and M-step until a suitable convergence criterion is satisfied. We adopt the distance involving two successive evaluations of the log-likelihood function, i.e., $\left| \ell(\boldsymbol{\theta}^{(k+1)} | \mathbf{y}) / \ell(\boldsymbol{\theta}^{(k)} | \mathbf{y}) - 1 \right|$, as a convergence criterion, where $\ell(\boldsymbol{\theta} | \mathbf{y}) = \sum_{i=1}^n \ln f_{\text{MMN}_p}(\mathbf{y}_i; \boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}, \nu)$.

4. Special Case of MMN Distribution

In this section, we study in detail a special case of the MMN family. In the stochastic representation in (1), if the random variable U follows the standard gamma distribution with corresponding PDF $h(u; \nu) = u^{\nu-1} e^{-u} / \Gamma(\nu)$, $u > 0$, we denote it by $\mathbf{Y} \sim \text{MMN}\mathcal{G}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}, \nu)$. Then, the PDF of \mathbf{Y} can be obtained from (4) as follows:

$$f_{\text{MMN}\mathcal{G}_p}(\mathbf{y}) = \frac{\sqrt{2\pi}}{\eta^\nu \Gamma(\nu)} \exp\left(-\frac{A^2}{2}\right) \phi_p(\mathbf{y}; \boldsymbol{\xi}, \boldsymbol{\Sigma}_{\mathbf{Y}}) \int_{-A}^{+\infty} (z + A)^{\nu-1} \phi(z) dz, \quad \mathbf{y} \in \mathbb{R}^p,$$

where $\eta = \sqrt{\delta^\top \omega \Sigma_Y^{-1} \omega \delta}$, $A = \eta^{-1} [\delta^\top \omega \Sigma_Y^{-1} (y - \xi) - 1]$ and $\Sigma_Y = \Omega - \omega \delta \delta^\top \omega$. By using the MGF in (5), for $\mathbf{Y} \sim \text{MMNG}_p(\xi, \Omega, \delta, \nu)$, we obtain

$$M_Y(\mathbf{t}) = e^{\mathbf{t}^\top \xi + \frac{1}{2} \mathbf{t}^\top \Sigma_Y \mathbf{t}} (1 - \mathbf{t}^\top \omega \delta)^{-\nu}, \quad \mathbf{t}^\top \omega \delta \neq 1, \quad \forall \mathbf{t}.$$

From the expressions in (7)-(14), and the fact that $E(U^r) = \Gamma(\nu + r)/\Gamma(\nu)$, for positive constant r , we can compute the first four moments of \mathbf{Y} by substituting $E(U) = \nu$, $E(U^2) = \nu(\nu + 1)$, $E(U^3) = \nu(\nu + 1)(\nu + 2)$, and $E(U^4) = \nu(\nu + 1)(\nu + 2)(\nu + 3)$. Specifically, we find that $E(\mathbf{Y}) = \xi + \nu \omega \delta$ and $\text{var}(\mathbf{Y}) = \Omega + (\nu - 1) \omega \delta \delta^\top \omega$.




Fig. 1: Contour plots of MMNE_2 distribution for different choices of δ . For the first two rows, the scale matrix is $\Omega = (1, 0; 0, 1)$, while for the third row, it is $\Omega = (1, 1; 1, 1.5)$.

Definition 1. (Bose et al. [16] ; Steutel and Van Harn [42]) A random vector \mathbf{Y} (or its distribution) is said to be infinitely divisible if, for each $n \geq 1$, there exist independent and identically distributed (iid) random vectors $\mathbf{Y}_1, \dots, \mathbf{Y}_n$ such that $\mathbf{Y} \stackrel{d}{=} \mathbf{Y}_1 + \dots + \mathbf{Y}_n$.

Theorem 7. The MMNG distribution, in the multivariate case, is infinitely divisible.

Proof. Without loss of generality, let $\mathbf{X} \sim \text{MMNG}_p(\mathbf{0}, \overline{\Omega}, \delta, \nu)$ and $\mathbf{X}_i \stackrel{d}{=} \delta U_i + \mathbf{Z}_i$, where $U_i \sim \text{Gamma}(\alpha = \frac{\nu}{n}, \beta = 1)$ and $\mathbf{Z}_i \sim \mathcal{N}_p(0, \frac{1}{n}(\overline{\Omega} - \delta \delta^\top))$ be independent random variables. It is easy to show that $\sum_{i=1}^n U_i \sim \text{Gamma}(\nu, 1)$ and $\sum_{i=1}^n \mathbf{Z}_i \sim \mathcal{N}_p(0, \overline{\Omega} - \delta \delta^\top)$, and so we can write $\mathbf{X} \stackrel{d}{=} \mathbf{X}_1 + \dots + \mathbf{X}_n$. Hence, the required result. \square

In the following, a particular case of the MMNG distribution with $\nu = 1$ is considered. Upon substituting $\nu = 1$, the mixing distribution of U follows the standard exponential distribution and the distribution of \mathbf{Y} in this case is

denoted by $\mathcal{MMNE}_p(\xi, \Omega, \delta)$. Then, the PDF of \mathbf{Y} can be obtained as

$$f_{\mathcal{MMNE}_p}(\mathbf{y}) = \frac{\sqrt{2\pi}}{\eta} \exp\left(-\frac{A^2}{2}\right) \phi_p(\mathbf{y}; \xi, \Sigma_Y) \Phi(A), \quad \mathbf{y} \in \mathbb{R}^p.$$

Fig. 1 presents the PDFs of the bivariate \mathcal{MMNE} distribution for $\Omega = (1, 0; 0, 1)$ and $\Omega = (1, 1; 1, 1.5)$, and different choices of δ for $\xi = (0, 0)^\top$. Fig. 1 shows that the \mathcal{MMNE} distribution exhibits a wide variety of density shapes, in terms of skewness. The PDF of the \mathcal{MMNE} distribution clearly depends on Ω and δ . The following theorem is useful in the implementation of the EM algorithm for the ML estimation of the parameters of the \mathcal{MMNE} distribution.

Theorem 8. *If $\mathbf{Y} \sim \mathcal{MMNE}_p(\xi, \Omega, \delta)$ and the random variable U follows the standard exponential distribution, then $U|\mathbf{Y} = \mathbf{y}) \sim \mathcal{TN}(\eta^{-1}A, \eta^{-2}, (0, \infty))$. Furthermore, for $k \in \{2, 3, \dots\}$,*

$$\mathbb{E}[U|\mathbf{Y} = \mathbf{y}] = \eta^{-1} \left(A + \frac{\phi(A)}{\Phi(A)} \right), \quad \mathbb{E}[U^k|\mathbf{Y} = \mathbf{y}] = A\eta^{-1}\mathbb{E}[U^{k-1}|\mathbf{Y} = \mathbf{y}] + (k-1)\eta^{-2}\mathbb{E}[U^{k-2}|\mathbf{Y} = \mathbf{y}].$$

Proof. The proof of the conditional distribution is completed easily by the use of Bayes rule. \square

Now, we can obtain the ML estimates of the parameters of \mathcal{MMNE} distribution. By using Theorem 8 and letting

$$\widehat{E}_{i1}^{(k)} = \mathbb{E}\left[U_i|\mathbf{Y}_i = \mathbf{y}_i, \widehat{\boldsymbol{\theta}}^{(k)}\right] = \frac{1}{\widehat{\eta}^{(k)}} \left(\widehat{A}_i^{(k)} + \frac{\phi(\widehat{A}_i^{(k)})}{\Phi(\widehat{A}_i^{(k)})} \right) \text{ and } \widehat{E}_{i2}^{(k)} = \mathbb{E}\left[U_i^2|\mathbf{Y}_i = \mathbf{y}_i, \widehat{\boldsymbol{\theta}}^{(k)}\right] = \frac{1}{\widehat{\eta}^{(k)^2}} \left[\widehat{A}_i^{(k)^2} + \widehat{A}_i^{(k)} \frac{\phi(\widehat{A}_i^{(k)})}{\Phi(\widehat{A}_i^{(k)})} + 1 \right],$$

in expression (17), the EM algorithm for the \mathcal{MMNE} distribution can be performed, where $\widehat{\eta}^{(k)} = \sqrt{\widehat{\boldsymbol{\alpha}}^{(k)\top} \widehat{\Sigma}_Y^{(k)-1} \widehat{\boldsymbol{\alpha}}^{(k)}}$ and $\widehat{A}_i^{(k)} = \widehat{\eta}^{(k)-1} \left[\widehat{\boldsymbol{\alpha}}^{(k)\top} \widehat{\Sigma}_Y^{(k)-1} (\mathbf{y}_i - \widehat{\xi}^{(k)}) - 1 \right]$. Note that, in the case of \mathcal{MMNE} distribution, the distribution of U does not have any parameter, and so there is no need to estimate ν in the EM algorithm and so M-step 2 must be skipped.

By using the fact that $E(U^m) = m!$, for $m \in \{1, 2, \dots\}$, and by using expressions in (11)-(14), for random vector $\mathbf{Y} \sim \mathcal{MMNE}_p(\xi, \Omega, \delta)$, we have

$$M_1(\mathbf{Y}) = \xi + \omega\delta, \tag{19}$$

$$M_2(\mathbf{Y}) = \xi \otimes \xi^\top + \xi \otimes \delta^\top \omega + \omega\delta \otimes \xi^\top + (\Sigma_Y + 2\omega\delta\delta^\top \omega), \tag{20}$$

$$M_3(\mathbf{Y}) = \xi\xi^\top \otimes \xi + \xi\xi^\top \otimes \omega\delta + \xi\delta^\top \omega \otimes \xi + \omega\delta \otimes \xi\xi^\top + (\Sigma_Y + 2\omega\delta\delta^\top \omega) \otimes \xi + \xi \otimes (\Sigma_Y + 2\omega\delta\delta^\top \omega) \\ + \text{vec}(\Sigma_Y + 2\omega\delta\delta^\top \omega) \otimes \xi^\top + \omega\delta \otimes \Sigma_Y + \text{vec}(\Sigma_Y) \delta^\top \omega + (\mathbf{I}_p \otimes \omega\delta) [\Sigma_Y + 6\omega\delta\delta^\top \omega], \tag{21}$$

$$M_4(\mathbf{Y}) = \xi\xi^\top \otimes \xi\xi^\top + \xi\xi^\top \otimes \xi(\omega\delta)^\top + \xi\xi^\top \otimes (\omega\delta)\xi^\top + \xi(\omega\delta)^\top \otimes \xi\xi^\top + \xi\xi^\top \otimes (\Sigma_Y + 2\omega\delta\delta^\top \omega) \\ + (\xi \otimes \xi) (\text{vec}(\Sigma_Y + 2\omega\delta\delta^\top \omega))^\top + \xi \otimes (\Sigma_Y + 2\omega\delta\delta^\top \omega) \otimes \xi^\top + \xi \otimes \omega (M_3^X)^\top (\omega \otimes \omega) + (\omega\delta)\xi^\top \otimes \xi\xi^\top \\ + \xi^\top \otimes (\Sigma_Y + 2\omega\delta\delta^\top \omega) \otimes \xi + \xi^\top \otimes \text{vec}(\Sigma_Y + 2\omega\delta\delta^\top \omega) \otimes \xi^\top + \xi^\top \otimes (\omega \otimes \omega) M_3^X \omega \\ + (\Sigma_Y + 2\omega\delta\delta^\top \omega) \otimes \xi\xi^\top + \omega (M_3^X)^\top (\omega \otimes \omega) \otimes \xi + (\omega \otimes \omega) M_3^X \omega \otimes \xi^\top + (\omega \otimes \omega) M_4^X (\omega \otimes \omega), \tag{22}$$

where

$$M_3^X = \delta \otimes \Sigma_X + \text{vec}(\Sigma_X) \delta^\top + (\mathbf{I}_p \otimes \delta) \Sigma_X + 6(\mathbf{I}_p \otimes \delta)(\delta \otimes \delta^\top), \\ M_4^X = (\mathbf{I}_{p^2} + \mathbf{U}_{p,p})(\Sigma_X \otimes \Sigma_X) + \text{vec}(\Sigma_X)(\text{vec}(\Sigma_X))^\top + 2[\delta \otimes \delta^\top \otimes \Sigma_X + \delta \otimes \Sigma_X \otimes \delta^\top + \Sigma_X \otimes \delta \otimes \delta^\top \\ + \delta^\top \otimes \Sigma_X \otimes \delta + \delta^\top \otimes \text{vec}(\Sigma_X) \otimes \delta^\top + (\delta \otimes \delta)(\text{vec}(\Sigma_X))^\top] + 24\delta\delta^\top \otimes \delta\delta^\top,$$

and $\Sigma_X = \overline{\Omega} - \delta\delta^\top$. In particular, the mean vector and covariance matrix are $\mathbb{E}(\mathbf{Y}) = \xi + \omega\delta$ and $\text{var}(\mathbf{Y}) = \Omega$.

Theorem 9. *The \mathcal{MMNE} distribution, in the multivariate case, is log-concave.*

Proof. Because log-concavity is preserved by affine transformations, it is sufficient to prove this property for the canonical form $\mathbf{Z}^* \sim \mathcal{MMNE}_p(\mathbf{0}, \mathbf{I}_p, \delta_{\mathbf{Z}^*})$. From [4] and [38], if the elements of a random vector are independent, and each has a log-concave density function, then their joint density is log-concave. We know that in the canonical form with PDF in (15), the random variables Z_1, \dots, Z_p are independent of each other. Log-concavity of \mathcal{MMNE} distribution in the univariate case has been established in Proposition 3.1 of [37], and the PDF of the univariate normal distribution is also known to be log-concave. Hence, the result. \square

As shown in Section 2, to compute the mode of the \mathcal{MMNE} distribution, it is sufficient to obtain the mode of the distribution in its canonical form, and then compute the mode of the distribution using Theorem 6. To compute the mode of the distribution in its canonical form, we must calculate the value of the mode in the univariate case. Existence and uniqueness of the mode (log-concavity) of the \mathcal{MMNE} distribution in the univariate case has been discussed in Proposition 3.1 of [37]. For this purpose, we recall the density function of the univariate \mathcal{MMNE} distribution (given in [37]) as

$$f_{Z_1}(z; \xi, \omega^2, \lambda) = \frac{\sqrt{1 + \lambda^2}}{\omega|\lambda|} e^{-\frac{\sqrt{1 + \lambda^2}}{\lambda} z + \frac{1}{2\lambda^2}} \Phi\left(\frac{\lambda \sqrt{1 + \lambda^2} z - 1}{|\lambda|}\right),$$

where $z = (y - \xi)/\omega$, $\lambda = \delta / \sqrt{1 - \delta^2} \neq 0$, $y \in \mathbb{R}$, $\xi \in \mathbb{R}$ is a location parameter and $\omega > 0$ is a scale parameter. It is denoted by $\mathcal{MMNE}_1(\xi, \omega^2, \lambda)$. For obtaining the mode of \mathcal{MMNE}_1 , based on Theorem 6, we need to solve the equation $\partial f_{Z_1}(z; 0, 1, \lambda_*) / \partial z = 0$, where $\lambda_* = \delta_* / \sqrt{1 - \delta_*^2}$. The solution need to be obtained by using numerical methods.

5. Multivariate Measures of Skewness

The skewed shape of the distribution is usually captured by multivariate skewness measures. The skewness is a measure of the asymmetry of a distribution about its mean and its value far from zero indicates stronger asymmetry of the underlying distribution than that with close to zero skewness value. The best-known scalar function of the

Table 1: Multivariate measures of skewness for the \mathcal{MMN} family.

Mardia [33] & Malkovich and Afifi [32]	$\beta_{1,p} = (\gamma_1^*)^2$
Srivastava [41]	$\beta_{1,p}^2 = \frac{1}{p} \sum_{i=1}^p \left\{ \frac{E[\gamma_i^T (\mathbf{Y} - \mu)]^3}{\lambda_i^{3/2}} \right\}^2$
Móri-Rohatgi-Székely [36]	$\mathbf{s} = \sum_{i=1}^p E(\mathbf{Z}_i^2 \mathbf{Z}) = \left(\sum_{i=1}^p E(\mathbf{Z}_i^2 Z_1), \dots, \sum_{i=1}^p E(\mathbf{Z}_i^2 Z_p) \right)^T$
Kollo [27]	$\mathbf{b} = E\left(\sum_{i,j} (Z_i Z_j) \mathbf{Z}\right) = \left(\sum_{i,j} E[(Z_i Z_j) Z_1], \dots, \sum_{i,j} E[(Z_i Z_j) Z_p] \right)^T$
Balakrishnan-Brito-Quiroz [11]	$\mathbf{T} = \int_{\mathbb{R}^p} \mathbf{u} c_1(\mathbf{u}) d\lambda(\mathbf{u})$, $\mathcal{Q}^* = \mathbf{T}^T \Sigma_{\mathbf{Z}}^{-1} \mathbf{T}$ The elements of \mathbf{T} are $T_r = \frac{3}{p(p+2)} E(Z_r^3) + 3 \sum_{i \neq r} \frac{1}{p(p+2)} E(Z_i^2 Z_r)$
Isogai [25]	$s_I = \frac{[\delta_* E(U) - m_0^*]^2}{1 + \delta_*^2 (\text{var}(U) - 1)}$, $s_C = \left(E(U) - \frac{m_0^*}{\delta_*} \right) \delta$

vectorial measure of skewness proposed by [36] is its squared norm. Its sampling properties has been thoroughly discussed by [24] and have been implemented in the R package `MultiSkew`. [20] discusses the usage of `MultiSkew` and briefly review the literature related to the same squared norm. Also, the skewness measure in [32] has become an useful tool in projection pursuit ([31]).

In this work, multivariate measures of skewness by Mardia [33], Malkovich and Afifi [32], Srivastava [41], Móri et al. [36], Kollo [27], Balakrishnan et al. [11] and Isogai [25] are studied for the \mathcal{MMN} family. Table 1 presents these measures for the \mathcal{MMN} family of distributions. The relevant derivations are given in Appendix B. In Table 1, γ_1^* is the skewness of $Z_1^* \sim \mathcal{MMN}_1(0, 1, \delta_*; H)$ of the canonical form, respectively. Srivastava measure uses principal components $\mathbf{F} = \mathbf{\Gamma} \mathbf{Y}$, where $\mathbf{\Gamma} = (\gamma_1, \dots, \gamma_p)$ is the matrix of eigenvectors of the covariance matrix $\mathbf{\Lambda}$, that is, an orthogonal matrix such that $\mathbf{\Gamma}^T \mathbf{\Lambda} \mathbf{\Gamma} = \mathbf{\Lambda}$, and $\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_p)$ is diagonal matrix of corresponding eigenvalues. Here, $\mathbf{Z} = \mathbf{\Lambda}^{-1/2} (\mathbf{Y} - \mu) = (Z_1, \dots, Z_p)^T$ has the distribution $\mathcal{MMN}_p(\xi_{\mathbf{Z}}, \mathbf{\Omega}_{\mathbf{Z}}, \delta_{\mathbf{Z}}; H)$, with its parameters as $\xi_{\mathbf{Z}} = \mathbf{\Lambda}^{-1/2} (\xi - \mu)$, $\mathbf{\Omega}_{\mathbf{Z}} = \mathbf{\Lambda}^{-1/2} \mathbf{\Omega} \mathbf{\Lambda}^{-1/2}$, $\delta_{\mathbf{Z}} = \omega_{\mathbf{Z}}^{-1} \mathbf{\Lambda}^{-1/2} \omega \delta$, and $\omega_{\mathbf{Z}} = (\mathbf{\Omega}_{\mathbf{Z}} \odot \mathbf{I}_p)^{1/2}$. Also, m_0^* is the mode of the

scalar MMN distribution in the canonical form. From Table 1, and using the moments in (19)-(22), we can obtain different measures of skewness for the $MMNE_p(\xi, \Omega, \delta)$ distribution as follows:

- **Mardia and Malkovich-Afifi indices:** $\beta_{1,p} = \beta_1^* = 4\delta_*^6$;
- **Srivastava index:** $\beta_{1p}^2 = \frac{1}{p} \sum_{i=1}^p \left\{ \frac{E[\gamma_i^\top (\mathbf{Y} - \mu)]^3}{\lambda_i^{3/2}} \right\}^2$, where γ_i and λ_i are eigenvectors and corresponding eigenvalues for covariance matrix $\text{var}(\mathbf{Y}) = \Omega$, when $\mathbf{Y} \sim MMNE_p(\xi, \Omega, \delta)$;
- **Móri-Rohatgi-Székelly index:** If $\mathbf{Y} \sim MMNE_p(\xi, \Omega, \delta)$, then for the standardized variable $\mathbf{Z} = \Omega^{-1/2}(\mathbf{Y} - \mu)$, and with $\mathbf{A} = \Omega^{-1/2}$, we have (see Appendix A)

$$\begin{aligned} M_3(\mathbf{Z}) &= E[\mathbf{A}^\top (\mathbf{Y} - \mu)]^3 = (\mathbf{A}^\top \otimes \mathbf{A}^\top) M_3(\mathbf{Y}) \mathbf{A} - [\mathbf{A}^\top M_2(\mathbf{Y}) \mathbf{A}] \otimes [\mathbf{A}^\top E(\mathbf{Y})] - \mathbf{A}^\top E(\mathbf{Y}) \otimes [\mathbf{A}^\top M_2(\mathbf{Y}) \mathbf{A}] \\ &\quad - \text{vec}(\mathbf{A}^\top M_2(\mathbf{Y}) \mathbf{A}) E(\mathbf{Y})^\top \mathbf{A} + 2[\mathbf{A}^\top E(\mathbf{Y}) E(\mathbf{Y})^\top \mathbf{A}] \otimes [\mathbf{A}^\top E(\mathbf{Y})]. \end{aligned} \quad (23)$$

All the quantities in the Móri-Rohatgi-Székelly measure of skewness are specific non-central moments of third order of \mathbf{Z} , where $\mathbf{Z} = \Omega^{-1/2}(\mathbf{Y} - \mu) \sim MMNE_p(\xi_Z, \Omega_Z, \delta_Z)$, such that $\xi_Z = -\Omega^{-1/2}\omega\delta$, $\Omega_Z = \mathbf{I}_p$ and $\delta_Z = \Omega^{-1/2}\omega\delta$;

- **Kollo index:** To obtain Kollo's measure, we use the elements of non-central moments of third order of \mathbf{Z} ;
- **Balakrishnan-Brito-Quiroz index:** Upon substituting $E(U^m) = m!$ for $m = 1, 2, \dots$, the elements of \mathbf{T} in Table 1, for $r = 1, 2, \dots, p$, are $\mathbf{T}_r = \frac{3}{p(p+2)} (\mathbf{M}_3^Z[(r-1)p + r, r] + \sum_{i \neq r} \mathbf{M}_3^Z[(i-1)p + i, r])$, where $\mathbf{M}_3^Z[., .]$ denotes the elements of matrix \mathbf{M}_3^Z , third moments of $MMNE_p(\xi_Z, \Omega_Z, \delta_Z)$ distribution, and we can then compute $\mathbf{Q} = \mathbf{T}^\top \Sigma_T^{-1} \mathbf{T}$ and $\mathbf{Q}^* = \mathbf{T}^\top \mathbf{T}$;
- **Isogai index:** By substituting $E(U) = \text{var}(U) = 1$ in Isogai measure of skewness, we have $S_I = (\delta_* - m_0^*)^2$, where m_0^* is the mode of the $MMNE_1$ distribution in the canonical form. This index is location and scale invariant. The vectorial measure, given by [12], is $S_C = (1 - \frac{m_0^*}{\delta_*})\delta$. Therefore, the direction of δ can be regarded as a measure of vectorial skewness for the $MMNE$ distribution.

Table 2: The Average computational time spent (Atime), the average values (Mean), the corresponding standard deviations (Std.), Bias and MSE of the EM estimates over 1000 samples from the $MMNE$ model in Subsection 6.1.

n	Atime		ξ_1	ξ_2	ξ_3	δ_1	δ_2	δ_3	σ_{11}	σ_{22}	σ_{33}
50	0.3265	Mean	5.0804	10.1518	15.1735	0.1663	0.4940	0.2259	0.3947	0.5931	0.9800
		Std.	0.2625	0.3163	0.4202	0.3870	0.3912	0.4041	0.0846	0.1512	0.2003
		Bias	0.0804	0.1518	0.1735	-0.1337	-0.2060	-0.1741	-0.0053	-0.0069	-0.0200
		MSE	0.0753	0.1230	0.2065	0.1675	0.1953	0.1934	0.0072	0.0229	0.0405
100	0.4824	Mean	5.0409	10.0554	15.0637	0.2360	0.6221	0.3346	0.3979	0.5973	0.9871
		Std.	0.1755	0.2028	0.2814	0.2581	0.2453	0.2651	0.0584	0.1110	0.1492
		Bias	0.0409	0.0554	0.0637	-0.0640	-0.0779	-0.0654	-0.0021	-0.0027	-0.0129
		MSE	0.0324	0.0441	0.0832	0.0706	0.0662	0.0745	0.0034	0.0123	0.0224
500	1.8063	Mean	5.0006	10.0036	15.0024	0.3001	0.6973	0.3964	0.3993	0.5995	1.0004
		Std.	0.0508	0.0534	0.0733	0.0686	0.0492	0.0587	0.0268	0.0483	0.0644
		Bias	0.0006	0.0036	0.0025	0.0006	-0.0027	-0.0036	0.0007	0.0005	0.0004
		MSE	0.0026	0.0029	0.0054	0.0047	0.0024	0.0035	0.0007	0.0023	0.0041
1000	4.0388	Mean	5.0004	9.9996	15.0025	0.3006	0.6995	0.3971	0.3999	0.5997	0.9981
		Std.	0.0345	0.0355	0.0529	0.0435	0.0328	0.0424	0.0180	0.0336	0.0447
		Bias	0.0004	-0.0004	0.0024	0.0001	-0.0005	-0.0029	-0.0001	-0.0003	-0.0002
		MSE	0.0012	0.0013	0.0028	0.0019	0.0011	0.0018	0.0003	0.0011	0.0020

6. Simulation Study

6.1. Model Fitting

This subsection presents the results of a Monte Carlo simulation study carried out to examine the performance of the proposed estimation method for the $MMNE$ distribution in the trivariate case. We evaluate the estimates in

terms of Bias and MSE (mean squared error). The results are based on 1000 simulated samples from the \mathcal{MMNE} distribution with parameters $\xi = (5, 10, 15)^\top$, $\Omega = \text{diag}(0.4, 0.6, 1.0)$, $\delta = (0.3, 0.7, 0.4)^\top$ for different sample sizes $n \in \{50, 100, 500, 1000\}$. We computed the Bias and the MSE as $\text{Bias} = \frac{1}{1000} \sum_{j=1}^{1000} (\hat{\theta}_j - \theta)$ and $\text{MSE} = \frac{1}{1000} \sum_{j=1}^{1000} (\hat{\theta}_j - \theta)^2$, where θ is the true parameter (each of $\xi = (\xi_1, \xi_2, \xi_3)^\top$, $\delta = (\delta_1, \delta_2, \delta_3)^\top$ and $\Omega = \text{diag}(\sigma_{11}, \sigma_{22}, \sigma_{33})^\top$) and $\hat{\theta}_j$ is the estimate from the j -th simulated sample. Table 2 presents the average computational time spent (Atime) (in seconds) (computational time for the convergence of the EM algorithm), average values (Mean), the corresponding standard deviations (Std.), Bias and MSE of the EM estimates of all the parameters of the \mathcal{MMNE} model in 1000 simulated samples for each sample size. It can be observed from Table 2 that the Bias and MSE decrease as n increases, revealing the asymptotic unbiasedness and consistency of the ML estimates obtained through the EM algorithm. Note that the EM algorithm presented in this work, for the \mathcal{MMNE} distribution, leads to closed-form expressions, and so the computational time required for the convergence of the EM estimates of the parameters is quite short.

Table 3: Skewness measures in Section 5, for some bivariate \mathcal{MMNE} distributions with presumed parameter $\xi = \mathbf{0}$ and different choices of parameters Ω and δ .

#	Parameters	$\beta_{1,p}$	$\beta_{1,p}^2$	s	b	Q^*	T	s_I	s_C
1	$\Omega = \begin{bmatrix} 1 & 1 \\ 1 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.750 \\ 0.985 \end{bmatrix}$	3.966	1.975	$\begin{bmatrix} 0.825 \\ 1.812 \end{bmatrix}$	$\begin{bmatrix} 1.448 \\ 3.179 \end{bmatrix}$	0.558	$\begin{bmatrix} 0.310 \\ 0.680 \end{bmatrix}$	0.788	$\begin{bmatrix} 0.667 \\ 0.876 \end{bmatrix}$
2	$\Omega = \begin{bmatrix} 1 & 0 \\ 0 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.200 \\ 0.975 \end{bmatrix}$	3.889	1.718	$\begin{bmatrix} 0.396 \\ 1.932 \end{bmatrix}$	$\begin{bmatrix} 0.552 \\ 2.692 \end{bmatrix}$	0.547	$\begin{bmatrix} 0.149 \\ 0.724 \end{bmatrix}$	0.673	$\begin{bmatrix} 0.165 \\ 0.804 \end{bmatrix}$
3	$\Omega = \begin{bmatrix} 1 & 0 \\ 0 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.000 \\ 0.995 \end{bmatrix}$	3.881	1.941	$\begin{bmatrix} 0.000 \\ 1.970 \end{bmatrix}$	$\begin{bmatrix} 0.000 \\ 1.970 \end{bmatrix}$	0.546	$\begin{bmatrix} 0.000 \\ 0.739 \end{bmatrix}$	0.666	$\begin{bmatrix} 0.000 \\ 0.816 \end{bmatrix}$
4	$\Omega = \begin{bmatrix} 1 & 1 \\ 1 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.650 \\ 0.995 \end{bmatrix}$	3.890	1.774	$\begin{bmatrix} 0.562 \\ 1.890 \end{bmatrix}$	$\begin{bmatrix} 0.870 \\ 2.924 \end{bmatrix}$	0.547	$\begin{bmatrix} 0.211 \\ 0.709 \end{bmatrix}$	0.675	$\begin{bmatrix} 0.536 \\ 0.821 \end{bmatrix}$
5	$\Omega = \begin{bmatrix} 1 & 1 \\ 1 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.850 \\ 0.900 \end{bmatrix}$	3.337	1.511	$\begin{bmatrix} 1.099 \\ 1.460 \end{bmatrix}$	$\begin{bmatrix} 2.154 \\ 2.862 \end{bmatrix}$	0.469	$\begin{bmatrix} 0.412 \\ 0.547 \end{bmatrix}$	0.412	$\begin{bmatrix} 0.562 \\ 0.595 \end{bmatrix}$
6	$\Omega = \begin{bmatrix} 1 & 0 \\ 0 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.550 \\ -0.800 \end{bmatrix}$	3.349	0.580	$\begin{bmatrix} 1.037 \\ -1.508 \end{bmatrix}$	$\begin{bmatrix} 0.069 \\ -0.100 \end{bmatrix}$	0.471	$\begin{bmatrix} 0.389 \\ -0.566 \end{bmatrix}$	0.415	$\begin{bmatrix} 0.365 \\ -0.531 \end{bmatrix}$
7	$\Omega = \begin{bmatrix} 1 & 1 \\ 1 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.900 \\ 0.775 \end{bmatrix}$	2.731	0.843	$\begin{bmatrix} 1.254 \\ 1.077 \end{bmatrix}$	$\begin{bmatrix} 2.493 \\ 2.141 \end{bmatrix}$	0.384	$\begin{bmatrix} 0.470 \\ 0.404 \end{bmatrix}$	0.286	$\begin{bmatrix} 0.513 \\ 0.442 \end{bmatrix}$
8	$\Omega = \begin{bmatrix} 1 & 0 \\ 0 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.800 \\ 0.400 \end{bmatrix}$	2.048	0.532	$\begin{bmatrix} 1.280 \\ 0.640 \end{bmatrix}$	$\begin{bmatrix} 2.304 \\ 1.152 \end{bmatrix}$	0.288	$\begin{bmatrix} 0.480 \\ 0.240 \end{bmatrix}$	0.193	$\begin{bmatrix} 0.393 \\ 0.196 \end{bmatrix}$
9	$\Omega = \begin{bmatrix} 1 & 1 \\ 1 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.750 \\ 0.150 \end{bmatrix}$	1.607	0.521	$\begin{bmatrix} 1.263 \\ -0.110 \end{bmatrix}$	$\begin{bmatrix} 1.045 \\ -0.091 \end{bmatrix}$	0.226	$\begin{bmatrix} 0.473 \\ -0.041 \end{bmatrix}$	0.1451	$\begin{bmatrix} 0.333 \\ 0.066 \end{bmatrix}$
10	$\Omega = \begin{bmatrix} 1 & 1 \\ 1 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} -0.750 \\ -0.150 \end{bmatrix}$	1.607	0.521	$\begin{bmatrix} -1.263 \\ 0.110 \end{bmatrix}$	$\begin{bmatrix} -1.045 \\ 0.091 \end{bmatrix}$	0.226	$\begin{bmatrix} -0.474 \\ 0.041 \end{bmatrix}$	0.145	$\begin{bmatrix} -0.333 \\ -0.067 \end{bmatrix}$
11	$\Omega = \begin{bmatrix} 1 & 0 \\ 0 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.700 \\ 0.000 \end{bmatrix}$	0.471	0.235	$\begin{bmatrix} 0.686 \\ 0.000 \end{bmatrix}$	$\begin{bmatrix} 0.686 \\ 0.000 \end{bmatrix}$	0.066	$\begin{bmatrix} 0.257 \\ 0.000 \end{bmatrix}$	0.043	$\begin{bmatrix} 0.208 \\ 0.000 \end{bmatrix}$
12	$\Omega = \begin{bmatrix} 1 & -1 \\ -1 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.000 \\ 0.000 \end{bmatrix}$	0.000	0.000	$\begin{bmatrix} 0.000 \\ 0.000 \end{bmatrix}$	$\begin{bmatrix} 0.000 \\ 0.000 \end{bmatrix}$	0.000	$\begin{bmatrix} 0.000 \\ 0.000 \end{bmatrix}$	0.000	$\begin{bmatrix} 0.000 \\ 0.000 \end{bmatrix}$

6.2. Assessment of Skewness

To study and compare different multivariate measures of skewness for the \mathcal{MMN} distributions, we consider the \mathcal{MMNE} distribution. We compute the values of all the skewness measures for different choices of the parameters of the bivariate and trivariate \mathcal{MMNE} distributions. Tables 3 and 4 present the values of all the skewness measures. It should be noted that all the measures are location and scale invariant, a desirable property indeed for any measure of skewness. For similar work on skewness comparisons for \mathcal{SN} distribution, one may refer to [12], and also to [26] for a similar work on scale mixtures of \mathcal{SN} distributions. From Table 3, we find that in all cases with scalar measures of skewness, Mardia's measure have the highest value and Srivastava's measure is the next largest. Just as in the case of \mathcal{SN} distribution, for the bivariate \mathcal{MMNE} distribution, the vectorial measures yield very similar results in terms of skewness directions, especially when the distribution is highly asymmetric [12]. It is important to note that Cases 9 and 10 deal with reflected distributions; and in these cases, all the measures are the same and the vectorial ones are reflected as well. Table 4 presents the values of all the measures for the trivariate \mathcal{MMNE} distribution. In this case,

differences among the measures become much more pronounced. From Table 4, we find that in all cases, among the vectorial measures of skewness, Mardia's measure has the highest value. Of course, the magnitude of the measures alone does not say much; one has to know how significant the values are!

Table 4: Skewness measures in Section 5, for some trivariate $MMNE$ distributions with presumed parameter $\xi = \mathbf{0}$ and different choices of parameters Ω and δ .

#	Parameters	$\beta_{1,p}$	β_{1p}^2	s	b	Q^*	T	s_I	s_C
1	$\Omega = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2.5 & 0 \\ 0 & 0 & 2.5 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.10 \\ 0.70 \\ 0.70 \end{bmatrix}$	3.881	0.314	$\begin{bmatrix} 0.198 \\ 1.386 \\ 1.386 \end{bmatrix}$	$\begin{bmatrix} 0.450 \\ 3.150 \\ 3.150 \end{bmatrix}$	0.155	$\begin{bmatrix} 0.040 \\ 0.277 \\ 0.277 \end{bmatrix}$	$\begin{bmatrix} 0.082 \\ 0.574 \\ 0.574 \end{bmatrix}$
2	$\Omega = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2.5 & 1 \\ 1 & 1 & 10 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.75 \\ 0.75 \\ 0.65 \end{bmatrix}$	2.712	0.235	$\begin{bmatrix} 0.752 \\ 1.044 \\ 1.028 \end{bmatrix}$	$\begin{bmatrix} 2.211 \\ 3.070 \\ 3.023 \end{bmatrix}$	0.108	$\begin{bmatrix} 0.150 \\ 0.209 \\ 0.206 \end{bmatrix}$	$\begin{bmatrix} 0.426 \\ 0.426 \\ 0.369 \end{bmatrix}$
3	$\Omega = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2.5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.995 \\ 0.00 \\ 0.00 \end{bmatrix}$	3.881	1.294	$\begin{bmatrix} 1.970 \\ 0.000 \\ 0.000 \end{bmatrix}$	$\begin{bmatrix} 1.970 \\ 0.000 \\ 0.000 \end{bmatrix}$	0.155	$\begin{bmatrix} 0.394 \\ 0.000 \\ 0.000 \end{bmatrix}$	$\begin{bmatrix} 0.816 \\ 0.000 \\ 0.000 \end{bmatrix}$
4	$\Omega = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2.5 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.40 \\ -0.60 \\ -0.60 \end{bmatrix}$	2.726	0.130	$\begin{bmatrix} 0.704 \\ -1.056 \\ -1.056 \end{bmatrix}$	$\begin{bmatrix} 0.512 \\ -0.768 \\ -0.768 \end{bmatrix}$	0.109	$\begin{bmatrix} 0.141 \\ -0.211 \\ -0.211 \end{bmatrix}$	$\begin{bmatrix} 0.228 \\ -0.342 \\ -0.342 \end{bmatrix}$
5	$\Omega = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2.5 & 1 \\ 1 & 1 & 10 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.55 \\ 0.05 \\ -0.30 \end{bmatrix}$	1.372	0.223	$\begin{bmatrix} 1.055 \\ -0.140 \\ -0.490 \end{bmatrix}$	$\begin{bmatrix} 0.139 \\ -0.018 \\ -0.064 \end{bmatrix}$	0.055	$\begin{bmatrix} 0.211 \\ -0.028 \\ -0.098 \end{bmatrix}$	$\begin{bmatrix} 0.230 \\ 0.021 \\ -0.125 \end{bmatrix}$
6	$\Omega = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.75 \\ 0.35 \\ 0.35 \end{bmatrix}$	2.106	0.242	$\begin{bmatrix} 1.211 \\ 0.565 \\ 0.565 \end{bmatrix}$	$\begin{bmatrix} 3.154 \\ 1.472 \\ 1.472 \end{bmatrix}$	0.084	$\begin{bmatrix} 0.242 \\ 0.113 \\ 0.113 \end{bmatrix}$	$\begin{bmatrix} 0.373 \\ 0.174 \\ 0.174 \end{bmatrix}$

Table 5: Upper and lower 2.5% critical values based on the test statistics $\beta_{1,p}$, β_{1p}^2 , s_{sum} , s_{max} , b_{sum} , b_{max} , Q^* , T_{sum} , T_{max} , s_I , $s_{C\text{sum}}$, and $s_{C\text{max}}$, in Subsection 6.3, obtained from 10000 simulated samples of standard multivariate normal distribution, for sample size $n = 100$ and dimensions $p \in \{2, \dots, 8\}$.

Test Statistics	Percentile	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$
$\beta_{1,p}$	0.025	0.0000	0.0000	0.0685	0.1232	0.2304	0.3170	0.4388
	0.975	1.1816	1.4552	2.1717	2.8814	3.5455	3.8266	3.9085
β_{1p}^2	0.025	0.0000	0.0000	0.0025	0.0038	0.0041	0.0035	0.0042
	0.975	0.4474	0.2918	0.2827	0.2906	0.3167	0.2311	0.2654
s_{max}	0.025	-0.3562	-0.1626	-0.0975	-0.0244	0.0160	0.1061	0.1452
	0.975	0.8894	0.9881	1.1032	1.2195	1.2737	1.3150	1.4936
s_{sum}	0.025	-0.9945	-1.3716	-1.6814	-1.9739	-2.2822	-2.5103	-2.7708
	0.975	1.0052	1.3275	1.4690	1.9442	2.1993	2.5009	3.1478
b_{max}	0.025	-0.6804	-0.3989	-0.2855	-0.0707	0.0000	0.0002	0.0018
	0.975	1.0957	1.6014	1.7865	2.4971	2.6128	3.1300	4.0002
b_{sum}	0.025	-1.7299	-2.8449	-3.9202	-5.5741	-6.4435	-7.3387	-8.0038
	0.975	1.7621	2.8721	3.3774	5.6679	6.3422	7.8015	10.2697
Q^*	0.025	0.0000	0.0000	0.0011	0.0009	0.0009	0.0007	0.0006
	0.975	0.1662	0.0582	0.0339	0.0212	0.0138	0.0087	0.0055
T_{max}	0.025	-0.1336	-0.0325	-0.0122	-0.0021	0.0010	0.0051	0.0054
	0.975	0.3335	0.1976	0.1379	0.1045	0.0796	0.0626	0.0560
T_{sum}	0.025	-0.3729	-0.2743	-0.2102	-0.1692	-0.1426	-0.1195	-0.1039
	0.975	0.3769	0.2655	0.1836	0.1666	0.1375	0.1191	0.1180
s_I	0.025	0.0000	0.0000	0.0080	0.0133	0.0229	0.0304	0.0407
	0.975	0.1045	0.1301	0.2077	0.3122	0.4770	0.6192	0.6956
$s_{C\text{max}}$	0.025	-0.1130	-0.0536	-0.0347	-0.0108	0.0109	0.0311	0.0482
	0.975	0.2710	0.2955	0.3410	0.4033	0.4830	0.5097	0.6410
$s_{C\text{sum}}$	0.025	-0.2920	-0.4116	-0.5417	-0.6276	-0.7303	-0.8984	-1.0657
	0.975	0.3036	0.3932	0.4644	0.6154	0.7311	0.8275	1.1957

6.3. Comparison and Performance of Different Skewness Measures

The measures studied in Section 5 and in the preceding subsection are not directly comparable with each other. So, for comparing them, we should have measures obtained on the same scale. To get such a set of comparable indices, we study the sample version for each of the skewness measures considered as test statistics for the hypothesis of normal distribution against $MMNE$ distribution, and then use the power of test based on different test statistics. Let Y_1, Y_2, \dots, Y_n denote a sample of $p \times 1$ observations from any p -dimensional distribution. Then, a sample version of

Table 6: Upper 5% critical values based on the test statistics $\beta_{1,p}$, β_{1p}^2 , s_{sum} , s_{max} , b_{sum} , b_{max} , Q^* , T_{sum} , T_{max} , s_I , $s_{C\text{sum}}$, and $s_{C\text{max}}$, in Subsection 6.3, obtained from 10000 simulated samples of standard multivariate normal distribution, for sample size $n = 100$ and dimensions $p \in \{2, \dots, 8\}$.

Test Statistics	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$
$\beta_{1,p}$	0.9325	1.2317	1.7024	2.4021	3.0324	3.4899	3.8939
β_{1p}^2	0.3111	0.2135	0.1997	0.1812	0.2044	0.1612	0.2039
s_{max}	0.7679	0.8177	0.9712	1.0931	1.2011	1.1774	1.3446
s_{sum}	0.8546	1.0509	1.2424	1.6682	1.8934	1.9996	2.4041
b_{max}	0.9619	1.2206	1.3468	1.9457	2.0467	2.2711	2.8953
b_{sum}	1.3001	2.1093	2.5046	3.9367	4.4979	5.1949	7.1686
Q^*	0.1311	0.0493	0.0266	0.0176	0.0118	0.0079	0.0055
T_{max}	0.2880	0.1635	0.1214	0.0937	0.0751	0.0561	0.0504
T_{sum}	0.3205	0.2102	0.1553	0.1430	0.1183	0.0952	0.0902
s_I	0.0824	0.1091	0.1549	0.2375	0.3409	0.4576	0.6792
$s_{C\text{max}}$	0.2324	0.2468	0.2959	0.3382	0.4069	0.4282	0.5509
$s_{C\text{sum}}$	0.2619	0.3224	0.3757	0.5072	0.6228	0.6621	0.8116

Table 7: Simulated values of power for all tests based on the test statistics $\beta_{1,p}$, β_{1p}^2 , s_{sum} , s_{max} , b_{sum} , b_{max} , Q^* , T_{sum} , T_{max} , s_I , $s_{C\text{sum}}$, and $s_{C\text{max}}$, in Subsection 6.3, for bivariate normal distribution against $MMNE$ distribution.

#	Parameters	$\beta_{1,p}$	β_{1p}^2	s_{max}	s_{sum}	b_{max}	b_{sum}	Q^*	T_{max}	T_{sum}	s_I	$s_{C\text{max}}$	$s_{C\text{sum}}$
1	$\Omega = \begin{bmatrix} 1 & 0 \\ 0 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.1 \\ 0.1 \end{bmatrix}$	0.030	0.041	0.034	0.040	0.038	0.035	0.030	0.034	0.040	0.030	0.030	0.044
2	$\delta = \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix}$	0.283	0.121	0.172	0.467	0.492	0.497	0.283	0.172	0.467	0.283	0.167	0.459
3	$\delta = \begin{bmatrix} 0.1 \\ 0.8 \end{bmatrix}$	0.659	0.748	0.711	0.668	0.631	0.349	0.659	0.711	0.668	0.659	0.690	0.634
4	$\delta = \begin{bmatrix} 0.8 \\ 0.1 \end{bmatrix}$	0.682	0.724	0.729	0.700	0.661	0.383	0.682	0.729	0.700	0.682	0.711	0.682
5	$\delta = \begin{bmatrix} 0.8 \\ 0.8 \end{bmatrix}$	0.994	0.994	0.994	0.994	0.994	0.994	0.994	0.994	0.994	0.994	0.994	0.994
6	$\delta = \begin{bmatrix} -0.7 \\ -0.7 \end{bmatrix}$	0.982	0.982	0.982	0.981	0.981	0.981	0.982	0.982	0.981	0.982	0.982	0.981
7	$\Omega = \begin{bmatrix} 1 & 1 \\ 1 & 2.5 \end{bmatrix}$, $\delta = \begin{bmatrix} 0.1 \\ 0.1 \end{bmatrix}$	0.032	0.035	0.043	0.057	0.060	0.059	0.032	0.043	0.057	0.032	0.087	0.140
8	$\delta = \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix}$	0.074	0.079	0.062	0.159	0.189	0.186	0.074	0.062	0.159	0.074	0.088	0.315
9	$\delta = \begin{bmatrix} 0.1 \\ 0.8 \end{bmatrix}$	0.979	0.929	0.980	0.668	0.105	0.004	0.979	0.980	0.668	0.979	0.974	0.956
10	$\delta = \begin{bmatrix} 0.8 \\ 0.1 \end{bmatrix}$	0.982	0.978	0.984	0.941	0.639	0.084	0.982	0.984	0.941	0.982	0.977	0.963
11	$\delta = \begin{bmatrix} 0.8 \\ -0.1 \end{bmatrix}$	0.956	0.958	0.959	0.816	0.076	0.002	0.956	0.959	0.816	0.956	0.960	0.938
12	$\delta = \begin{bmatrix} -0.8 \\ 0.1 \end{bmatrix}$	0.953	0.955	0.023	0.832	0.003	0.005	0.953	0.023	0.832	0.953	0.024	0.936
13	$\delta = \begin{bmatrix} -0.8 \\ -0.1 \end{bmatrix}$	0.968	0.963	0.004	0.921	0.004	0.088	0.968	0.004	0.921	0.968	0.248	0.944
14	$\delta = \begin{bmatrix} 0.8 \\ 0.8 \end{bmatrix}$	0.929	0.929	0.815	0.977	0.979	0.980	0.929	0.815	0.977	0.929	0.918	0.987

all the skewness measures described can be obtained by replacing ξ , Ω , and δ with the maximum likelihood estimates of these quantities [12]. As seen in the previous sections, the Mardia and Malkovich-Afifi measure $\beta_{1,p}$, Srivastava measure β_{1p}^2 , Isogai measure s_I , Balakrishnan-Brito-Quiroz measure Q^* are scalar indices and the Móri-Rohatgi-Székely measure s , Kollo measure b , Balakrishnan-Brito-Quiroz measure T , and Isogai measure s_C are vectorial indices. Here, we study different statistics for testing the null hypothesis and powers for each of these tests to quantify the capacity of each skewness measure to identify the specific asymmetry present in the $MMNE$ distribution. The power of the test is a probability, and its use enables us to compare different statistics, no matter what the original scales of them were. To obtain a single test statistic for the vectorial measures, we propose two different metrics, namely, the sum and the maximum (see [12], pages 82-83). For the Móri-Rohatgi-Székely measure, we compute

Table 8: Simulated values of power for all tests based on the test statistics $\beta_{1,p}$, $\beta_{1,p}^2$, s_{sum} , s_{max} , b_{sum} , b_{max} , Q^* , T_{sum} , T_{max} , s_I , $s_{C\text{sum}}$, and $s_{C\text{max}}$, in Subsection 6.3, for trivariate normal distribution against $MMNE$ distribution.

#	Parameters		$\beta_{1,p}$	$\beta_{1,p}^2$	s_{max}	s_{sum}	b_{max}	b_{sum}	Q^*	T_{max}	T_{sum}	s_I	$s_{C\text{max}}$	$s_{C\text{sum}}$
1	$\Omega = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2.5 & 0 \\ 0 & 0 & 2.5 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.1 \\ 0.1 \\ 0.1 \end{bmatrix}$	0.056	0.058	0.047	0.036	0.042	0.035	0.056	0.047	0.036	0.056	0.053	0.038
2		$\delta = \begin{bmatrix} -0.1 \\ -0.1 \\ -0.1 \end{bmatrix}$	0.057	0.061	0.057	0.035	0.048	0.052	0.057	0.057	0.035	0.057	0.056	0.039
3	$\Omega = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2.5 & 1 \\ 1 & 1 & 10 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.1 \\ 0.7 \\ 0.1 \end{bmatrix}$	0.601	0.192	0.637	0.166	0.082	0.031	0.601	0.637	0.166	0.601	0.557	0.513
4	$\Omega = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2.5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.7 \\ 0.1 \\ 0.1 \end{bmatrix}$	0.272	0.324	0.309	0.277	0.262	0.168	0.272	0.309	0.277	0.272	0.308	0.287
5	$\Omega = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2.5 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.4 \\ 0.6 \\ 0.6 \end{bmatrix}$	0.899	0.513	0.816	0.902	0.901	0.902	0.899	0.816	0.902	0.899	0.815	0.901
6	$\Omega = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2.5 & 1 \\ 1 & 1 & 10 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.7 \\ 0.05 \\ 0.3 \end{bmatrix}$	0.745	0.674	0.750	0.501	0.277	0.114	0.745	0.750	0.501	0.745	0.638	0.709
7	$\Omega = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\delta = \begin{bmatrix} 0.7 \\ 0.3 \\ 0.3 \end{bmatrix}$	0.572	0.349	0.461	0.726	0.729	0.703	0.572	0.461	0.726	0.572	0.476	0.711
8		$\delta = \begin{bmatrix} -0.7 \\ 0.3 \\ 0.1 \end{bmatrix}$	0.381	0.228	0.025	0.025	0.015	0.015	0.380	0.025	0.025	0.380	0.029	0.029
9		$\delta = \begin{bmatrix} 0.7 \\ -0.3 \\ 0.1 \end{bmatrix}$	0.415	0.248	0.400	0.061	0.034	0.013	0.415	0.401	0.061	0.415	0.414	0.081
10		$\delta = \begin{bmatrix} 0.7 \\ 0.3 \\ -0.1 \end{bmatrix}$	0.394	0.240	0.378	0.344	0.259	0.152	0.392	0.378	0.344	0.394	0.385	0.362
11		$\delta = \begin{bmatrix} -0.7 \\ -0.3 \\ 0.1 \end{bmatrix}$	0.404	0.206	0.073	0.304	0.059	0.146	0.404	0.073	0.304	0.404	0.072	0.317
12		$\delta = \begin{bmatrix} -0.7 \\ 0.3 \\ -0.1 \end{bmatrix}$	0.415	0.222	0.034	0.050	0.015	0.022	0.415	0.034	0.050	0.415	0.044	0.069
13		$\delta = \begin{bmatrix} 0.7 \\ -0.3 \\ -0.1 \end{bmatrix}$	0.393	0.257	0.356	0.031	0.020	0.015	0.393	0.356	0.031	0.393	0.370	0.037
14		$\delta = \begin{bmatrix} -0.8 \\ -0.8 \\ -0.8 \end{bmatrix}$	0.996	0.965	0.997	0.993	0.993	0.993	0.996	0.997	0.993	0.996	0.997	0.993

$s_{\text{sum}} = \sum_{r=1}^p s_r$ and $s_{\text{max}} = \max_{r \in \{1, \dots, p\}} s_r$, for the Kollo measure $b_{\text{sum}} = \sum_{r=1}^p b_r$ and $b_{\text{max}} = \max_{r \in \{1, \dots, p\}} b_r$, for the Balakrishnan-Brito-Quiroz measure $T_{\text{sum}} = \sum_{r=1}^p T_r$ and $T_{\text{max}} = \max_{r \in \{1, \dots, p\}} T_r$, for Isogai's measure $s_{C\text{sum}} = \sum_{r=1}^p s_{C_r}$ and $s_{C\text{max}} = \max_{r \in \{1, \dots, p\}} s_{C_r}$. The distributions of sample versions of measures, $\beta_{1,p}$, $\beta_{1,p}^2$, s_{sum} , s_{max} , b_{sum} , b_{max} , Q^* , T_{sum} , T_{max} , s_I , $s_{C\text{sum}}$, and $s_{C\text{max}}$ are not analytically computable easily, and so we may determine the critical values of these tests through Monte Carlo simulations. Two sets of critical values obtained by Monte Carlo simulation, based on 10000 samples from the standard multivariate normal distribution, are presented in Tables 5 and 6, for dimensions $p \in \{2, \dots, 8\}$. To get the values of critical values, we first simulated 10000 samples of size $n = 100$ from the standard multivariate normal distribution with dimensions $p \in \{2, \dots, 8\}$. We estimated the parameters and then found the values of test statistics. Then, we arranged the obtained values in increasing order and then selected the 2.5 and 5 lower and upper percentage points as critical values.

For computing the powers of the different tests, based on the above test statistics, we simulated 1000 samples from $MMNE$ distribution of size $n = 100$ for different choices of the parameters of the $MMNE$ distribution and estimated the test statistics by using the ML estimates of parameters evaluated by EM algorithm. Then, we computed the proportion of samples falling in the same rejection region. For test statistics $\beta_{1,p}$, $\beta_{1,p}^2$, Q^* and s_I , we considered the sample versions exceeding the critical values as critical regions, in the form $CR = \{Q_0 > q_\alpha\}$, and for all other test statistics, the rejection regions were the two-sided areas of the form $CR = \{Q_0 < q_{1-\alpha/2} \text{ or } Q_0 > q_{\alpha/2}\}$, where Q_0 is

test statistic under null hypothesis and q_α is upper α percentile of distribution of test statistic.

Table 9: Simulated values of power for all tests based on the test statistics $\beta_{1,p}$, $\beta_{1,p}^2$, s_{sum} , s_{max} , b_{sum} , b_{max} , Q^* , T_{sum} , T_{max} , s_I , $s_{C\text{sum}}$, and $s_{C\text{max}}$, in Subsection 6.3, for seven dimensional normal distribution against $MMNE$ distribution when $\Omega = \mathbf{I}_7$.

#	Parameter	$\beta_{1,p}$	$\beta_{1,p}^2$	s_{max}	s_{sum}	b_{max}	b_{sum}	Q^*	T_{max}	T_{sum}	s_I	$s_{C\text{max}}$	$s_{C\text{sum}}$
1	$\delta = (0.1, 0.1, 0.1, 0.1, 0.1, 0.1, 0.1)^\top$	0.061	0.060	0.061	0.038	0.042	0.046	0.062	0.061	0.044	0.061	0.059	0.045
2	$\delta = (0.7, 0.7, 0.7, 0.7, 0.7, 0.7, 0.7)^\top$	0.978	0.985	0.000	0.985	0.985	0.985	0.978	0.000	0.985	0.978	0.704	0.985
3	$\delta = (0.1, 0.7, 0.1, 0.7, 0.1, 0.7, 0.1)^\top$	0.952	0.543	0.035	0.952	0.953	0.951	0.952	0.035	0.952	0.952	0.580	0.954
4	$\delta = (0.4, 0.2, 0.5, 0.1, 0.7, 0.6, 0.3)^\top$	0.918	0.357	0.033	0.923	0.927	0.923	0.919	0.034	0.923	0.918	0.326	0.925
5	$\delta = -(0.1, 0.1, 0.1, 0.1, 0.1, 0.1, 0.1)^\top$	0.052	0.055	0.093	0.058	0.047	0.070	0.052	0.095	0.060	0.052	0.075	0.054
6	$\delta = -(0.7, 0.7, 0.7, 0.7, 0.7, 0.7, 0.7)^\top$	0.980	0.982	0.986	0.982	0.983	0.982	0.980	0.986	0.982	0.980	0.983	0.982
7	$\delta = (0.1, -0.7, 0.1, -0.7, 0.1, -0.7, 0.1)^\top$	0.959	0.516	0.010	0.684	0.000	0.080	0.959	0.010	0.771	0.959	0.007	0.841
8	$\delta = (-0.4, 0.2, -0.5, 0.1, -0.7, 0.6, -0.3)^\top$	0.889	0.413	0.015	0.009	0.006	0.005	0.891	0.015	0.014	0.889	0.102	0.156

In the simulation study, we took $\xi = \mathbf{0}$, and the parameters Ω and δ as given in Tables 7-9. Tables 7-9 present the power of the proposed tests for bivariate, trivariate and seven dimensional normal distribution against $MMNE$ distribution, respectively. The comparison of different measures may be done directly from the results in Tables 7-9. These results show clearly which are the poorer indices of skewness among those considered. Based on our empirical study, by considering different cases of the $MMNE$ distributions in two, three and seven dimensions, we make the following points: for all cases with small skewness, as expected, the power of the tests are lower for distributions more similar to the normal, and test statistics $\beta_{1,p}$, $\beta_{1,p}^2$, Q^* and s_I have better performance. From Tables 7-9, as expected, for increasing values of the elements of the skewness parameters, the power values of all tests increase. For large elements close to 1 or -1, for skewness parameters, the power of the tests are higher and have almost the same values for different test statistics.

The behaviour of test statistics $\beta_{1,p}$, Q^* and s_I , are very close to each other and have the same power. For small values of the skewness parameter, these test statistics have poorer performance. The power of the tests for s_{max} , and T_{max} statistics are the same, and the test statistics s_{sum} and T_{sum} often have similar behaviour. For large and moderate values of skewness parameters, b_{sum} and b_{max} statistics have the lowest test power and have the worst performance compared with other test statistics. For the bivariate case in Table 7, when one element of the skewness parameter is large and one is small, the statistics $\beta_{1,p}^2$, T_{max} and s_{max} perform well, but b_{sum} and b_{max} statistics have the lowest test power.

For the trivariate case in Table 8, when one element of the skewness parameter is large and two elements and small, the statistics $\beta_{1,p}^2$, T_{max} and s_{max} have better performance.

From Table 9, for case 3, the statistic $s_{C\text{sum}}$ has the best performance and T_{max} and s_{max} have lower power close to 0.05. From Table 9, for case 4, the statistic b_{max} has the best performance, but T_{max} and s_{max} have lower power close to 0.05. A result that we find from Tables 7-9 is that the test statistic $s_{C\text{max}}$ performs better than others in many cases.

7. Illustrative Examples

In this section, we fit the $MMNE$ model for two real data sets to illustrate the flexibility of the model. It is also compared with SN and ST distributions in terms of some measures of fit.

7.1. AIS data

The first example considers the Australian Institute of Sport (AIS) data [17], containing 11 biomedical measurements on 202 Australian athletes (100 female and 102 male). Here, we focus solely on the first 100, and the trivariate case corresponding to BMI, SSF and Bfat variables, where the three acronyms denote Body Mass Index, Sum of Skin Folds, and Body Fat percentage, respectively. These data are available in the R software, `sn` package. Fig. 2 presents the histograms for the three variables. Upon using the EM algorithm, we obtained the maximum likelihood estimates of parameters of the model. Table 10 presents the estimates of parameters (ξ, Ω, δ) . Table 11 presents values of all skewness measures by using the estimates of parameters, presented in Table 10. The relative difference in the fit of a number of candidate models can be compared by using the maximized log-likelihood values $\ell(\hat{\theta}|\mathbf{y})$, the Akaike information criterion (AIC) and the Bayesian information criterion (BIC). The AIC and BIC indices are defined as




Fig. 2: The histograms for the three selected variables BMI, SSF and Bfat of the AIS data set in Subsection 7.1.

$AIC = 2k - \ell(\hat{\theta}|\mathbf{y})$ and $BIC = k \ln n - 2\ell(\hat{\theta}|\mathbf{y})$, where k is the number of model parameters and $\ell(\hat{\theta}|\mathbf{y})$ is the maximized log-likelihood value of a fitted model. The larger value of $\ell(\hat{\theta}|\mathbf{y})$ and the smaller value of AIC or BIC indicates a better fit of the model to the data.

Table 12 summarizes the fitting performance of $MMNE$ model, as compared to the SN and ST distributions. From Table 12, it is seen that the $MMNE$ model provides the best fit overall as it provides the largest $\ell(\hat{\theta}|\mathbf{y})$ value and the lowest AIC and BIC values. Fig. 3 shows the scatter plots of pairs of the three variables BMI, SSF, Bfat, along with the contour plots for the fitted $MMNE$, SN and ST distributions.

Table 10: Parameter estimates of the $MMNE$ distribution by using EM Algorithm presented in Section 3, based on the three selected variables (BMI, SSF, Bfat) of the AIS data set in Subsection 7.1.

$\hat{\xi}$	$\hat{\Omega}$			$\hat{\delta}$
20.1099	7.2870	66.3650	10.2745	0.6963
56.1969	66.3650	1238.1858	191.2748	0.8747
13.6666	10.2745	191.2748	31.3535	0.7471

Table 11: Values of skewness measures in Section 5, based on the three selected variables (BMI, SSF, Bfat) of the AIS data set in Subsection 7.1.

$\beta_{1,p}$	$\beta_{1,p}^2$	s	b	Q^*	T	s_I	s_C
3.5539	0.5973	$\begin{bmatrix} 0.3182 \\ 1.7703 \\ -0.5644 \end{bmatrix}$	$\begin{bmatrix} 0.2080 \\ 1.1571 \\ -0.3689 \end{bmatrix}$	0.1422	$\begin{bmatrix} 0.0636 \\ 0.3541 \\ -0.1129 \end{bmatrix}$	0.4800	$\begin{bmatrix} 0.4920 \\ 0.6181 \\ 0.5279 \end{bmatrix}$

7.2. Italian olive oil data

As a second example, we consider the well-known data on the percentage composition of eight fatty acids found by lipid fraction of 572 Italian olive oils. These data come from three areas; within each area, there are a number of constituent regions, 9 in total. The data set includes a data frame with 572 observations and 10 columns. The




Fig. 3: Scatter plots of pairs of the three selected variables for the AIS data set, along with the contour plots for the fitted $MMNE$, skew-normal (SN) and skew-t (ST) distributions presented in Section 1 and 2.

Table 12: Comparison of fitting measures, maximized log-likelihood value $\ell(\hat{\theta}|\mathbf{y})$, Akaike information criterion AIC and Bayesian information criterion BIC, for skew-normal (SN), skew-t (ST) and $MMNE$ distributions for the three selected variables (BMI, SSF, Bfat) of the AIS data set in Subsection 7.1.

Distribution	$\ell(\hat{\theta} \mathbf{y})$	AIC	BIC
SN	-866.2725	1756.545	1787.807
ST	-852.1354	1730.271	1764.138
$MMNE$	-850.7388	1725.478	1756.740

first column gives the area (one of Southern Italy, Sardinia, and Northern Italy), the second gives the region, and the remaining 8 columns give the variables. Southern Italy consists of North Apulia, Calabria, South Apulia, and Sicily regions, Sardinia is divided into Inland Sardinia, and Coastal Sardinia, and Northern Italy consists of Umbria, East Liguria, and West Liguria regions. These data are available in the R software, pgmm package.




Fig. 4: The histograms of the two selected variables Linolenic and Arachidic fatty acids of olive oil data set in Subsection 7.2.

Table 13: Parameter estimates of the $MMNE$ distribution by using EM Algorithm presented in Section 3, based on the two selected variables (Linolenic, Arachidic) of the olive oil data set in Subsection 7.2.

ξ	Ω		δ
36.8344	63.3623	40.9481	0.1546
55.3462	40.9481	124.0575	0.6977

Table 14: Values of skewness measures in Section 5, based on the two selected variables (Linolenic, Arachidic) of the olive oil data set in Subsection 7.2.

$\beta_{1,p}$	β_{1p}^2	s	b	Q^*	T	s_I	s_C
0.5707	0.1218	$\begin{bmatrix} -0.0492 \\ 0.7538 \end{bmatrix}$	$\begin{bmatrix} -0.0428 \\ 0.6557 \end{bmatrix}$	0.0802	$\begin{bmatrix} -0.0185 \\ 0.2827 \end{bmatrix}$	0.0517	$\begin{bmatrix} 0.0486 \\ 0.2195 \end{bmatrix}$

For the purpose of illustration, we consider 323 cases from Southern Italy, and columns (8, 9), Linolenic and Arachidic fatty acids, respectively, so as to consider the bivariate case. Fig. 4 shows the histograms of the two selected variables, while Table 13 presents the estimates of parameters and Table 14 presents the values of skewness measures. Table 15 provides the fit of $MMNE$ model, as compared to those of SN and ST distributions, for the considered data. From Table 15, it is clear that the $MMNE$ model provides the best overall fit as it possesses the largest $\ell(\hat{\theta}|\mathbf{y})$ value and the lowest AIC and BIC values. Fig. 5 shows the scatter plot of the data and the contour plots of the fitted $MMNE$, SN and ST distributions.

Table 15: Comparison of fitting measures, maximized log-likelihood value $\ell(\hat{\theta}|\mathbf{y})$, Akaike information criterion AIC and Bayesian information criterion BIC, for skew-normal (SN), skew-t (ST) and $MMNE$ distributions for the two selected variables (Linolenic, Arachidic) of the olive oil data set in Subsection 7.2.

Distribution	$\ell(\hat{\theta} \mathbf{y})$	AIC	BIC
SN	-2320.039	4654.079	4680.522
ST	-2316.320	4648.640	4678.861
$MMNE$	-2314.604	4643.207	4669.651




Fig. 5: Scatter plots of the olive oil data, and the contour plots of the fitted $MMNE$, skew-normal (SN) and skew-t (ST) distributions presented in Section 1 and 2.

8. Concluding Remarks

In this paper, we have discussed the mean mixture of multivariate normal distribution (MMN), which includes the normal, SN , and extended SN distributions as particular cases. We have studied several features of this family of distributions, including the first four moments, the distributions of affine transformations and canonical forms, estimation of parameters by using an EM-type algorithm with closed-form expressions, and different measures of multivariate skewness. Two special cases of the MMN family, with standard gamma and standard exponential distributions as mixing distributions, denoted by $MMNG$ and $MMNE$ distributions, have been studied in detail. A simulation study has been performed to evaluate the performance of the MLEs of parameters of the $MMNE$ distribution. From the results in Section 7, for the AIS and olive oil data sets, the $MMNE$ distribution is shown to provide a better fit than the SN and ST distributions. Different multivariate measures of skewness have been derived for the $MMNE$ distribution, and the evaluation of tests based on these measures is carried out in terms of powers of tests.

There are several possible directions for future research. For example, the study of finite mixtures and scale mixtures of MMN family will be of great interest. In the stochastic representation in (1), if the skewness parameter is a matrix, with representation $\mathbf{Y} \stackrel{d}{=} \boldsymbol{\xi} + \boldsymbol{\omega}(\Delta\mathbf{U} + \mathbf{Z})$, then \mathbf{Y} has the unified skew normal (SUN) distribution (see [5]), wherein elements of \mathbf{U} have the standard half-normal distribution. In this connection, consideration of a general distribution for \mathbf{U} would be of interest. All the computations presented in this paper were performed by using the statistical software R, version 4.0.0. The computer program for the implementation of the proposed EM-type algorithm and comparison of the skewness measures are available as supplementary material associated with this article.

CRedit authorship contribution statement

Me'raj Abdi: Conceptualization, Methodology, Software, Writing - original draft, Writing review & editing. Investigation, Validation. **Mohsen Madadi:** Methodology, Supervision, Investigation. **Narayanaswamy Balakrishnan:** Supervision, Writing - review & editing, Methodology. **Ahad Jamalizadeh:** Conceptualization, Methodology, Supervision, Visualization.

Acknowledgments

The authors are grateful to the Editors and two anonymous reviewers who provided very helpful feedback, comments, and suggestions, based on which the paper has improved significantly.

Appendix A. Proofs

Proof of Lemma 2. By using (6), we can calculate the partial derivatives of $M_{\mathbf{X}}(\mathbf{t})$, the MGF of normalized MMN distribution, that are directly related to the moments of the MMN random vector. Suppose $\mathbf{X} \sim MMN_p(\mathbf{0}, \bar{\mathbf{\Omega}}, \delta; H)$. Then, some derivatives of $M_{\mathbf{X}}(\mathbf{t})$ in (6) are as follows:

$$\begin{aligned} \frac{\partial M_{\mathbf{X}}(\mathbf{t})}{\partial \mathbf{t}} &= e^{\frac{1}{2}\mathbf{t}^\top \Sigma_{\mathbf{X}} \mathbf{t}} \left[\Sigma_{\mathbf{X}} \mathbf{t} M_U(\mathbf{t}^\top \delta) + \delta M_U^{(1)}(\mathbf{t}^\top \delta) \right], \\ \frac{\partial^2 M_{\mathbf{X}}(\mathbf{t})}{\partial \mathbf{t} \partial \mathbf{t}^\top} &= e^{\frac{1}{2}\mathbf{t}^\top \Sigma_{\mathbf{X}} \mathbf{t}} \left\{ M_U(\mathbf{t}^\top \delta) \left[\Sigma_{\mathbf{X}} + (\Sigma_{\mathbf{X}} \mathbf{t}) \otimes (\Sigma_{\mathbf{X}} \mathbf{t})^\top \right] + M_U^{(1)}(\mathbf{t}^\top \delta) \left[(\Sigma_{\mathbf{X}} \mathbf{t}) \otimes \delta^\top + \delta \otimes (\Sigma_{\mathbf{X}} \mathbf{t})^\top \right] + M_U^{(2)}(\mathbf{t}^\top \delta) \delta \otimes \delta^\top \right\}, \\ \frac{\partial^3 M_{\mathbf{X}}(\mathbf{t})}{\partial \mathbf{t} \partial \mathbf{t}^\top \partial \mathbf{t}} &= e^{\frac{1}{2}\mathbf{t}^\top \Sigma_{\mathbf{X}} \mathbf{t}} \left\{ M_U(\mathbf{t}^\top \delta) \left[(\Sigma_{\mathbf{X}} \mathbf{t}) \otimes \Sigma_{\mathbf{X}} + \text{vec}(\Sigma_{\mathbf{X}})(\Sigma_{\mathbf{X}} \mathbf{t})^\top + (\mathbf{I}_p \otimes (\Sigma_{\mathbf{X}} \mathbf{t}))(\Sigma_{\mathbf{X}} + (\Sigma_{\mathbf{X}} \mathbf{t}) \otimes (\Sigma_{\mathbf{X}} \mathbf{t})^\top) \right] \right. \\ &\quad + M_U^{(1)}(\mathbf{t}^\top \delta) \left[\delta \otimes \Sigma_{\mathbf{X}} + \text{vec}(\Sigma_{\mathbf{X}}) \delta^\top + (\mathbf{I}_p \otimes (\Sigma_{\mathbf{X}} \mathbf{t}))[\delta \otimes (\Sigma_{\mathbf{X}} \mathbf{t})^\top + (\Sigma_{\mathbf{X}} \mathbf{t}) \otimes \delta^\top] \right. \\ &\quad + \left. (\mathbf{I}_p \otimes \delta) \left(\Sigma_{\mathbf{X}} + (\Sigma_{\mathbf{X}} \mathbf{t}) \otimes (\Sigma_{\mathbf{X}} \mathbf{t})^\top \right) \right] \\ &\quad \left. + M_U^{(2)}(\mathbf{t}^\top \delta) \left[(\mathbf{I}_p \otimes (\Sigma_{\mathbf{X}} \mathbf{t}))(\delta \otimes \delta^\top) + (\mathbf{I}_p \otimes \delta) \left(\delta \otimes (\Sigma_{\mathbf{X}} \mathbf{t})^\top + (\Sigma_{\mathbf{X}} \mathbf{t}) \otimes \delta^\top \right) \right] + M_U^{(3)}(\mathbf{t}^\top \delta)(\mathbf{I}_p \otimes \delta)(\delta \otimes \delta^\top) \right\}, \end{aligned}$$

where $M_U^{(1)}(\mathbf{t}^\top \delta) = \frac{\partial M_U(\mathbf{t}^\top \delta)}{\partial \mathbf{t}}$, $M_U^{(2)}(\mathbf{t}^\top \delta) = \frac{\partial^2 M_U(\mathbf{t}^\top \delta)}{\partial \mathbf{t} \partial \mathbf{t}^\top}$ and $M_U^{(3)}(\mathbf{t}^\top \delta) = \frac{\partial^3 M_U(\mathbf{t}^\top \delta)}{\partial \mathbf{t} \partial \mathbf{t}^\top \partial \mathbf{t}}$. Setting $\mathbf{t} = \mathbf{0}$, as in [22], we obtain the first three moments of the MMN family. To find the fourth moment, since we only need the value of fourth partial derivative of $M_{\mathbf{X}}(\mathbf{t})$ at $\mathbf{t} = \mathbf{0}$, say $M_4(\mathbf{X}) = \frac{\partial^4 M_{\mathbf{X}}(\mathbf{t})}{\partial \mathbf{t} \partial \mathbf{t}^\top \partial \mathbf{t} \partial \mathbf{t}^\top} |_{\mathbf{t}=\mathbf{0}}$, we do not need to compute the whole expression. Instead, we can simply single out all the terms in $\frac{\partial^4 M_{\mathbf{X}}(\mathbf{t})}{\partial \mathbf{t} \partial \mathbf{t}^\top \partial \mathbf{t} \partial \mathbf{t}^\top}$ that do not contain the factor \mathbf{t} or \mathbf{t}^\top . \square

Note 1: The stochastic representation $\mathbf{Y} \stackrel{d}{=} \xi + \omega(\delta U + \mathbf{Z})$ can be used directly as a way to obtain the first four moments of \mathbf{Y} in the following formulas:

$$\begin{aligned} M_1(\mathbf{Y}) &= E(\mathbf{Y}), \quad M_2(\mathbf{Y}) = E(\mathbf{Y} \otimes \mathbf{Y}^\top) = E(\mathbf{Y} \mathbf{Y}^\top), \quad M_3(\mathbf{Y}) = E(\mathbf{Y} \otimes \mathbf{Y}^\top \otimes \mathbf{Y}) = E[(\mathbf{Y} \otimes \mathbf{Y}) \mathbf{Y}^\top], \\ M_4(\mathbf{Y}) &= E(\mathbf{Y} \otimes \mathbf{Y}^\top \otimes \mathbf{Y} \otimes \mathbf{Y}^\top) = E[(\mathbf{Y} \mathbf{Y}^\top) \otimes (\mathbf{Y} \mathbf{Y}^\top)]. \end{aligned}$$

The corresponding central moments of \mathbf{Y} are then

$$\begin{aligned} \bar{M}_1(\mathbf{Y}) &= \mathbf{0}, \quad \bar{M}_2(\mathbf{Y}) = E\{[\mathbf{Y} - E(\mathbf{Y})] \otimes [\mathbf{Y} - E(\mathbf{Y})]^\top\} = \text{var}(\mathbf{Y}), \\ \bar{M}_3(\mathbf{Y}) &= E\{[\mathbf{Y} - E(\mathbf{Y})] \otimes [\mathbf{Y} - E(\mathbf{Y})]^\top \otimes [\mathbf{Y} - E(\mathbf{Y})]\}, \\ \bar{M}_4(\mathbf{Y}) &= E\{([\mathbf{Y} - E(\mathbf{Y})][\mathbf{Y} - E(\mathbf{Y})]^\top) \otimes ([\mathbf{Y} - E(\mathbf{Y})][\mathbf{Y} - E(\mathbf{Y})]^\top)\}. \end{aligned}$$

Note 2: We know that for any multivariate random vector \mathbf{Y} , the central moments of third and fourth orders are related to the non-central moments by the following relationships (see, for example, [28] and [29]):

$$\overline{M}_3(\mathbf{Y}) = M_3(\mathbf{Y}) - M_2(\mathbf{Y}) \otimes \mathbf{E}(\mathbf{Y}) - \mathbf{E}(\mathbf{Y}) \otimes M_2(\mathbf{Y}) - \text{vec}(M_2(\mathbf{Y}))\mathbf{E}(\mathbf{Y})^\top + 2\mathbf{E}(\mathbf{Y})\mathbf{E}(\mathbf{Y})^\top \otimes \mathbf{E}(\mathbf{Y}), \quad (24)$$

$$\begin{aligned} \overline{M}_4(\mathbf{Y}) = & M_4(\mathbf{Y}) - (M_3(\mathbf{Y}))^\top \otimes \mathbf{E}(\mathbf{Y}) - M_3(\mathbf{Y}) \otimes \mathbf{E}(\mathbf{Y})^\top - \mathbf{E}(\mathbf{Y}) \otimes (M_3(\mathbf{Y}))^\top - \mathbf{E}(\mathbf{Y})^\top \otimes M_3(\mathbf{Y}) \\ & + M_2(\mathbf{Y}) \otimes \mathbf{E}(\mathbf{Y})\mathbf{E}(\mathbf{Y})^\top + (\mathbf{E}(\mathbf{Y}) \otimes \mathbf{E}(\mathbf{Y}))(\text{vec}(M_2(\mathbf{Y})))^\top + \mathbf{E}(\mathbf{Y}) \otimes M_2(\mathbf{Y}) \otimes \mathbf{E}(\mathbf{Y})^\top \\ & + \mathbf{E}(\mathbf{Y})^\top \otimes M_2(\mathbf{Y}) \otimes \mathbf{E}(\mathbf{Y}) + \mathbf{E}(\mathbf{Y})^\top \otimes \text{vec}(M_2(\mathbf{Y})) \otimes \mathbf{E}(\mathbf{Y})^\top + \mathbf{E}(\mathbf{Y})\mathbf{E}(\mathbf{Y})^\top \otimes M_2(\mathbf{Y}) \\ & - 3\mathbf{E}(\mathbf{Y})\mathbf{E}(\mathbf{Y})^\top \otimes \mathbf{E}(\mathbf{Y})\mathbf{E}(\mathbf{Y})^\top. \end{aligned} \quad (25)$$

Upon using the relations for affine transformations of moments, we then obtain

$$M_1(\mathbf{A}\mathbf{Y}) = \mathbf{E}(\mathbf{A}\mathbf{Y}) = \mathbf{A}\mathbf{E}(\mathbf{Y}), \quad M_2(\mathbf{A}\mathbf{Y}) = \mathbf{E}(\mathbf{A}\mathbf{Y} \otimes (\mathbf{A}\mathbf{Y})^\top) = \mathbf{A}\mathbf{E}(\mathbf{Y} \otimes \mathbf{Y}^\top)\mathbf{A}^\top, \quad (26)$$

$$M_3(\mathbf{A}\mathbf{Y}) = \mathbf{E}[(\mathbf{A}\mathbf{Y} \otimes \mathbf{A}\mathbf{Y})(\mathbf{A}\mathbf{Y})^\top] = \mathbf{E}\{\text{vec}(\mathbf{A}\mathbf{Y}(\mathbf{A}\mathbf{Y})^\top)(\mathbf{A}\mathbf{Y})^\top\} = (\mathbf{A} \otimes \mathbf{A})M_3(\mathbf{Y})\mathbf{A}^\top, \quad (27)$$

$$M_4(\mathbf{A}\mathbf{Y}) = \mathbf{E}(\mathbf{A}\mathbf{Y}(\mathbf{A}\mathbf{Y})^\top \otimes \mathbf{A}\mathbf{Y}(\mathbf{A}\mathbf{Y})^\top) = (\mathbf{A} \otimes \mathbf{A})M_4(\mathbf{Y})(\mathbf{A} \otimes \mathbf{A})^\top. \quad (28)$$

Proof of Theorem 3. The moment generating function of $\mathbf{A}^\top \mathbf{X}$ can be written as

$$M_{\mathbf{A}^\top \mathbf{X}}(\mathbf{t}) = M_{\mathbf{X}}(\mathbf{A}\mathbf{t}) = e^{\frac{1}{2}\mathbf{t}^\top (\mathbf{A}^\top \overline{\boldsymbol{\Omega}} \mathbf{A} - \mathbf{A}^\top \boldsymbol{\delta} \boldsymbol{\delta}^\top \mathbf{A}) \mathbf{t}} M_U(\mathbf{t}^\top \mathbf{A}^\top \boldsymbol{\delta}; \nu).$$

Upon using the uniqueness property of the moment generating function, the required result is obtained. \square

Proof of Theorem 4. The moment generating function of $\mathbf{X} = \mathbf{c} + \mathbf{A}^\top \mathbf{Y}$ can be written as

$$M_{\mathbf{X}}(\mathbf{t}) = e^{\mathbf{t}^\top \mathbf{c}} M_{\mathbf{Y}}(\mathbf{A}\mathbf{t}) = e^{\mathbf{t}^\top \boldsymbol{\xi} + \frac{1}{2}\mathbf{t}^\top (\boldsymbol{\Omega}_X - \boldsymbol{\omega}_X \boldsymbol{\delta}_X \boldsymbol{\delta}_X^\top \boldsymbol{\omega}_X) \mathbf{t}} M_U(\mathbf{t}^\top \boldsymbol{\omega}_X \boldsymbol{\delta}_X; \nu),$$

which completes the proof. \square

Proof of Theorem 5. We have introduced the \mathcal{MMN} distribution by assuming $\boldsymbol{\Omega} > 0$ through the factorization $\boldsymbol{\Omega} = \boldsymbol{\omega} \overline{\boldsymbol{\Omega}} \boldsymbol{\omega}$. The matrix $\overline{\boldsymbol{\Omega}}$ is a positive definite non-singular matrix if and only if there exists some invertible (non-singular) matrix \mathbf{C} such that $\overline{\boldsymbol{\Omega}} = \mathbf{C}^\top \mathbf{C}$. If $\boldsymbol{\delta} \neq \mathbf{0}$, there exists an orthogonal matrix \mathbf{P} with the first column being proportional to $\mathbf{C} \overline{\boldsymbol{\Omega}}^{-1} \boldsymbol{\delta}$, while for $\boldsymbol{\delta} = \mathbf{0}$ we set $\mathbf{P} = \mathbf{I}_p$. Finally, define $\mathbf{A}_* = (\mathbf{C}^{-1} \mathbf{P})^\top \boldsymbol{\omega}^{-1}$. By using Theorem 4, we see that $\mathbf{Z}^* = \mathbf{A}_*(\mathbf{Y} - \boldsymbol{\xi})$ has the stated distribution with $\boldsymbol{\delta}_{\mathbf{Z}^*} = (\boldsymbol{\delta}_*, 0, \dots, 0)^\top$. \square

Proof of Theorem 6. First, consider the mode of the corresponding canonical variable $\mathbf{Z}^* \sim \mathcal{MMN}_p(\mathbf{0}, \mathbf{I}_p, \boldsymbol{\delta}_{\mathbf{Z}^*}; H)$. We find this mode by solving the following equations with respect to z_1, z_2, \dots, z_p :

$$\frac{\partial f_{\mathbf{Z}^*}(z_1)}{\partial z_1} = 0, \quad z_i f_{\mathbf{Z}^*}(z_1) = 0, \quad i \in \{2, 3, \dots, p\}.$$

The last $p - 1$ equations are fulfilled when $z_i = 0$, while the root of the first equation corresponds to the mode, m_0^* say, of the $\mathcal{MMN}_1(0, 1, \boldsymbol{\delta}_*; H)$ distribution. Therefore, the mode of \mathbf{Z}^* is $\mathbf{M}_0^* = (m_0^*, 0, \dots, 0)^\top = \frac{m_0^*}{\boldsymbol{\delta}_*} \boldsymbol{\delta}_{\mathbf{Z}^*}$. From Theorem 5, we can write $\mathbf{Y} = \boldsymbol{\xi} + \boldsymbol{\omega} \mathbf{C}^\top \mathbf{P} \mathbf{Z}^*$ and $\boldsymbol{\delta}_{\mathbf{Z}^*} = \mathbf{P}^\top \mathbf{C} \overline{\boldsymbol{\Omega}}^{-1} \boldsymbol{\delta}$. As the mode is equivariant with respect to affine transformations, the mode of \mathbf{Y} is

$$\mathbf{M}_0 = \boldsymbol{\xi} + \frac{m_0^*}{\boldsymbol{\delta}_*} \boldsymbol{\omega} \mathbf{C}^\top \mathbf{P} \boldsymbol{\delta}_{\mathbf{Z}^*} = \boldsymbol{\xi} + \frac{m_0^*}{\boldsymbol{\delta}_*} \boldsymbol{\omega} \mathbf{C}^\top \mathbf{P} \mathbf{P}^\top \mathbf{C} \overline{\boldsymbol{\Omega}}^{-1} \boldsymbol{\delta} = \boldsymbol{\xi} + \frac{m_0^*}{\boldsymbol{\delta}_*} \boldsymbol{\omega} \boldsymbol{\delta}.$$

Hence, the result. \square

Appendix B. Computation of Different Measures of Skewness

B1. Mardia Measure of Skewness

Mardia [33, 34] presented a multivariate measure of skewness of an arbitrary p -dimensional distribution F with mean vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Delta}$. Let \mathbf{X} and \mathbf{Y} be two independent and identically distributed random vectors from distribution F . Then, the measure of skewness is

$$\beta_{1,p} = E \left[\left\{ (\mathbf{X} - \boldsymbol{\mu})^\top \boldsymbol{\Delta}^{-1} (\mathbf{Y} - \boldsymbol{\mu}) \right\}^3 \right], \quad (29)$$

where $\boldsymbol{\mu} = E(\mathbf{X})$ and $\boldsymbol{\Delta} = \text{var}(\mathbf{X})$. Mardia measure of skewness is location and scale invariant (see [33]). From Theorems 4 and 5, the \mathcal{MMN} family is closed under affine transformations and have a canonical form. If $\mathbf{X} \sim \mathcal{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$, there exists a linear transformation $\mathbf{Z}^* = \mathbf{A}_*(\mathbf{Y} - \boldsymbol{\xi})$ such that $\mathbf{Z}^* \sim \mathcal{MMN}_p(\mathbf{0}, \mathbf{I}_p, \boldsymbol{\delta}_{\mathbf{Z}^*}; H)$, where at most one component of $\boldsymbol{\delta}_{\mathbf{Z}^*}$ is not zero. Without loss of any generality, we take the first component of \mathbf{Z}^* to be skewed and denote it by Z_1^* , and so for computing the measure in (29), we can use the canonical form of the \mathcal{MMN} family. Let \mathbf{X}^* and \mathbf{Y}^* be two independent and identically distributed random vectors from $\mathcal{MMN}_p(\mathbf{0}, \mathbf{I}_p, \boldsymbol{\delta}_{\mathbf{Z}^*}; H)$. Now, by using $\boldsymbol{\mu}^* = E(\mathbf{X}^*) = E(\mathbf{Y}^*) = E(U)\boldsymbol{\delta}_{\mathbf{Z}^*}$ and $\boldsymbol{\Delta}^* = \text{var}(\mathbf{X}^*) = \text{var}(\mathbf{Y}^*) = \mathbf{I}_p + (\text{var}(U) - 1)\boldsymbol{\delta}_{\mathbf{Z}^*}\boldsymbol{\delta}_{\mathbf{Z}^*}^\top$ in (29), the Mardia measure of skewness can be expressed as

$$\beta_{1,p} = E \left[\left\{ (\mathbf{X}^* - \boldsymbol{\mu}^*)^\top [\boldsymbol{\Delta}^*]^{-1} (\mathbf{Y}^* - \boldsymbol{\mu}^*) \right\}^3 \right] = \left(E \left[\frac{Z_1^* - \delta_1^*}{\sqrt{\text{var}(Z_1^*)}} \right]^3 \right)^2 = (\gamma_1^*)^2, \quad (30)$$

where γ_1^* is the univariate skewness of $Z_1^* \sim \mathcal{MMN}_1(0, 1, \delta_1^*; H)$ of the canonical form (see Theorem 5). An explicit formula of γ_1^* can be found in [37] for the univariate case.

B2. Malkovich-Afifi Measure of Skewness

Malkovich and Afifi [32] proposed a measure of multivariate skewness as a different type of generalization of the univariate measure. By denoting the unit p -dimensional sphere by $\phi_p = \{\mathbf{u} \in \mathbb{R}^p; \|\mathbf{u}\| = 1\}$, for $\mathbf{u} \in \phi_p$, the usual univariate measure of skewness in the \mathbf{u} -direction is

$$\beta_1(\mathbf{u}) = \frac{[E\{\mathbf{u}^\top (\mathbf{Y} - E(\mathbf{Y}))\}]^3]^2}{[\text{var}(\mathbf{u}^\top \mathbf{Y})]^3}, \quad (31)$$

and so the Malkovich-Afifi multivariate extension of it is defined as

$$\beta_1^* = \sup_{\mathbf{u} \in \phi_p} \beta_1(\mathbf{u}). \quad (32)$$

Malkovich-Afifi measure of multivariate skewness is also location and scale invariant. [32] then defined the measures in (31) and (32) and showed that if \mathbf{Z} is the standardized variable $\mathbf{Z} = \boldsymbol{\Delta}^{-1/2}(\mathbf{Y} - \boldsymbol{\mu})$, an equivalent version is $\beta_1^* = \sup_{\mathbf{u} \in \phi_p} (E[\mathbf{u}^\top \mathbf{Z}]^3)^2$. For obtaining β_1^* for the \mathcal{MMN} family, it is convenient to use the canonical form. If $\mathbf{Y} \sim \mathcal{MMN}_p(\boldsymbol{\xi}, \boldsymbol{\Omega}, \boldsymbol{\delta}; H)$, there exists a linear transformation $\mathbf{Z}^* = \mathbf{A}_*(\mathbf{Y} - \boldsymbol{\xi})$ such that $\mathbf{Z}^* \sim \mathcal{MMN}_p(\mathbf{0}, \mathbf{I}_p, \boldsymbol{\delta}_{\mathbf{Z}^*}; H)$, where at most one component of $\boldsymbol{\delta}_{\mathbf{Z}^*}$ is not zero. This means that the Malkovich-Afifi index, which is the maximum of the univariate skewness measures among all the directions of the unit sphere, will be, for \mathbf{Z}^* , the index of asymmetry in the only (if there is) skew direction (without loss of any generality, we take the first component of \mathbf{Z}^* to be skewed and denote it by Z_1^*):

$$\beta_1^* = \beta_1^*(\mathbf{u}) = \sup_{\mathbf{u} \in \phi_p} \frac{[E\{\mathbf{u}^\top (\mathbf{Y} - E(\mathbf{Y}))\}]^3]^2}{[\text{var}(\mathbf{u}^\top \mathbf{Y})]^3} = \frac{[E\{Z_1^* - E(Z_1^*)\}]^3]^2}{[\text{var}(Z_1^*)]^3} = (\gamma_1^*)^2. \quad (33)$$

As in the case of Mardia index, we have used γ_1^* to denote the univariate skewness measure of the unique (if any) skewed component of the canonical form \mathbf{Z}^* . As this measure is location and scale invariant, it is invariant for linear transforms and consequently (33) is also the Malkovich-Afifi measure for \mathbf{Y} , and thus it is the same as the Mardia index in (30).

B3. Srivastava Measure of Skewness

Using principal components $\mathbf{F} = \mathbf{\Gamma}\mathbf{Y}$, Srivastava [41] developed a measure of skewness for the multivariate vector \mathbf{Y} , where $\mathbf{\Gamma} = (\boldsymbol{\gamma}_1, \dots, \boldsymbol{\gamma}_p)$ is the matrix of eigenvectors of the covariance matrix $\mathbf{\Lambda}$, that is, an orthogonal matrix such that $\mathbf{\Gamma}^\top \mathbf{\Lambda} \mathbf{\Gamma} = \mathbf{\Lambda}$, and $\lambda_1, \dots, \lambda_p$ are the corresponding eigenvalues. Srivastava's measure of skewness for \mathbf{Y} may then be presented as

$$\beta_{1p}^2 = \frac{1}{p} \sum_{i=1}^p \left\{ \frac{E(F_i - \theta_i)^3}{\lambda_i^{3/2}} \right\}^2 = \frac{1}{p} \sum_{i=1}^p \left\{ \frac{E[\boldsymbol{\gamma}_i^\top (\mathbf{Y} - \boldsymbol{\mu})]^3}{\lambda_i^{3/2}} \right\}^2, \quad (34)$$

where $F_i = \boldsymbol{\gamma}_i^\top \mathbf{Y}$ and $\theta_i = \boldsymbol{\gamma}_i^\top \boldsymbol{\mu}$. The measure in (34) is based on central moments of third order $E[\boldsymbol{\gamma}_i^\top (\mathbf{Y} - \boldsymbol{\mu})]^3$. For obtaining this measure for the \mathcal{MMN} distribution, we only need to obtain the non-central moments up to third order. Upon using the relations in (24)-(27), we can obtain the third central moment by replacing \mathbf{A} by $\boldsymbol{\gamma}_i$ in (23).

B4. Móri-Rohatgi-Székelly Measure of Skewness

Móri et al. [36] suggested a vectorial measure of skewness as a p -dimensional vector. If $\mathbf{Z} = \mathbf{\Lambda}^{-1/2}(\mathbf{Y} - \boldsymbol{\mu}) = (Z_1, \dots, Z_p)^\top$ is the standardized vector, this measure can be written in terms of coordinates of \mathbf{Z} as

$$s(\mathbf{Y}) = E(\|\mathbf{Z}\|^2 \mathbf{Z}) = E(\mathbf{Z}^\top \mathbf{Z} \mathbf{Z}) = \sum_{i=1}^p E(Z_i^2 \mathbf{Z}) = \left(\sum_{i=1}^p E(Z_i^2 Z_1), \dots, \sum_{i=1}^p E(Z_i^2 Z_p) \right)^\top. \quad (35)$$

All the quantities involved in (35) are specific non-central moments of third order of \mathbf{Z} . When \mathbf{Y} has a multivariate \mathcal{MMN} distribution, \mathbf{Z} is still \mathcal{MMN} distribution, and so we can use once again the expressions in Theorem 1. Now, $\mathbf{Z} = \mathbf{\Lambda}^{-1/2}(\mathbf{Y} - \boldsymbol{\mu}) = (Z_1, \dots, Z_p)^\top$ has the distribution $\mathcal{MMN}_p(\boldsymbol{\xi}_Z, \mathbf{\Omega}_Z, \boldsymbol{\delta}_Z; H)$.

Furthermore, upon replacing \mathbf{A} by $\mathbf{\Lambda}^{-1/2}$ in the third central moment in (23), using (24)-(27), and the moments in Theorem 1, we can compute $s(\mathbf{Y})$ in (35).

B5. Kollo Measure of Skewness

Kollo [27] noticed that Móri-Rohatgi-Székelly skewness measure $s(\mathbf{Y})$ does not include all third-order mixed moments. To include all mixed moments of the third order, he defined a skewness vector of \mathbf{Y} as

$$b(\mathbf{Y}) = E\left(\sum_{i,j} (Z_i Z_j) \mathbf{Z}\right) = \left(\sum_{i,j} E[(Z_i Z_j) Z_1], \dots, \sum_{i,j} E[(Z_i Z_j) Z_p] \right)^\top. \quad (36)$$

The required moments can be obtained from Theorem 1 and the corresponding measure in (36) can then be computed.

B6. Balakrishnan-Brito-Quiroz Measure of Skewness

When reporting the skewness of a univariate distribution, it is customary to indicate skewness direction by referring to skewness 'to the left' (negative) or 'to the right' (positive). It seems natural that, in the multivariate setting, one would also like to indicate a direction for the skewness of a distribution.

Both Mardia and Malkovich-Afifi measures give an overall view of skewness measures without any specific reference to the direction of skewness. For this reason, [11] modified the Malkovich-Afifi measure to produce an overall vectorial measure of skewness as

$$\mathbf{T} = \int_{\phi_p} \mathbf{u} c_1(\mathbf{u}) d\lambda(\mathbf{u}), \quad (37)$$

where $c_1(\mathbf{u}) = E[(\mathbf{u}^\top \mathbf{Z})^3]$ is a signed measure of skewness of the standardized variable $\mathbf{Z} = \mathbf{\Lambda}^{-1/2}(\mathbf{Y} - \boldsymbol{\mu})$ in the direction of \mathbf{u} , and λ denotes the rotationally invariant probability measure on the unit p -dimensional sphere $\phi_p = \{\mathbf{u} \in \mathbb{R}^p; \|\mathbf{u}\| = 1\}$.

From [11] and [12], it turns out that the computation of \mathbf{T} is straightforward and, when the distribution of \mathbf{Y} is absolutely continuous with respect to Lebesgue measure and symmetric (in the broad sense specified below), it has,

under some moment assumptions, a Gaussian asymptotic distribution with a limiting covariance matrix, Σ_T , that can be consistently estimated from the Z_i sample.

If $c_1(\mathbf{u})$ is negative, it indicates skewness in the direction of $-\mathbf{u}$, while $\mathbf{u}c_1(\mathbf{u})$ provides a vectorial index of skewness in the \mathbf{u} (or $-\mathbf{u}$) direction. Summation of these vectors over \mathbf{u} (in the form of an integral) will then yield an overall vectorial measure of skewness presented earlier in (37).

For obtaining a single measure, [11] proposed the quantity $\mathbf{Q} = \mathbf{T}^\top \Sigma_{\mathbf{T}}^{-1} \mathbf{T}$, where \mathbf{T} is as in (37) and $\Sigma_{\mathbf{T}}$ is the covariance matrix of \mathbf{T} . However, the covariance matrix $\Sigma_{\mathbf{T}}$ depends on the moments of sixth order. Sixth order moments in this family are not in explicit form, and so as done in [12], by replacing $\Sigma_{\mathbf{T}}$ by $\Sigma_{\mathbf{Z}}$, we obtain $\mathbf{Q}^* = \mathbf{T}^\top \Sigma_{\mathbf{Z}}^{-1} \mathbf{T}$, to provide a reasonable measure of overall skewness.

In the following, evaluation of \mathbf{T} using the integrals of some monomials over the unit sphere ϕ_p are required. From [11], let u_j be the j -th coordinate of a point $\mathbf{u} \in \phi_p$. Then, the values of the integrals

$$J_4 = \int_{\phi_p} u_j^4 d\lambda(\mathbf{u}) = \frac{3}{p(p+2)}, \quad J_{2,2} = \int_{\phi_p} u_j^2 u_i^2 d\lambda(\mathbf{u}) = \frac{1}{p(p+2)},$$

for $j \neq i, 1 \leq j, i \leq p$, are obtained using Theorem 3.3 of [19]. We see that the above integrals do not depend on the particular choices of j and i . Therefore, the r -th coordinate of \mathbf{T} is simply $\mathbf{T}_r = J_4 E(Z_r^3) + 3 \sum_{i \neq r} J_{2,2} E(Z_i^2 Z_r)$. So, we must obtain the moments $E(Z_r^3)$ and $E(Z_i^2 Z_r)$. The required moments can be obtained as $E(Z_i^3) = \mathbf{M}_3^Z[(i-1)p + i, i]$ and $E(Z_i^2 Z_j) = \mathbf{M}_3(\mathbf{Z})[(i-1)p + i, j]$, where $\mathbf{M}_3(\mathbf{Z})[.,.]$ denotes the elements of $\mathbf{M}_3(\mathbf{Z})$, third moment of the $\mathcal{MMN}_p(\xi_{\mathbf{Z}}, \Omega_{\mathbf{Z}}, \delta_{\mathbf{Z}}; H)$ distribution. Upon using the above moments, we can obtain the elements of \mathbf{T} as follows:

$$\mathbf{T}_r = \frac{3}{p(p+2)} E(Z_r^3) + 3 \sum_{i \neq r} \frac{1}{p(p+2)} E(Z_i^2 Z_r). \quad (38)$$

B7. Isogai Measure of Skewness

Isogai [25] considered an overall extension of Pearson measure of skewness to a multivariate case in the form

$$S_I = (\boldsymbol{\mu} - \mathbf{M}_0)^\top g^{-1}(\Delta) (\boldsymbol{\mu} - \mathbf{M}_0),$$

where $\boldsymbol{\mu}$, \mathbf{M}_0 and Δ are the mean, mode and the covariance matrix of \mathbf{Y} , respectively. The function $g(\Delta)$ is an “appropriate” function of the covariance matrix. To derive this measure of skewness, we need to obtain the mode of the \mathcal{MMN} distribution, but the uniqueness of the mode for the family of mean mixture of normal distributions is an open problem. For obtaining this measure, we choose $g(.)$ to be the identity function. This measure is location and scale invariant, and so by using the canonical form of the \mathcal{MMN} distribution, we get

$$S_I = \frac{[\delta_* E(U) - m_0^*]^2}{1 + \delta_*^2 [\text{var}(U) - 1]},$$

where $\delta_* = (\delta^\top \bar{\Omega}^{-1} \delta)^{1/2}$ and m_0^* is the mode of the single scalar \mathcal{MMN} distribution in the canonical form. This index is essentially the Mahalanobis distance between the null vector and the vector $E(\mathbf{Y}) - \mathbf{M}_0$, and it is indeed location and scale invariant. Another vectorial measure has been given by [12] as $S_C = \omega^{-1} (\boldsymbol{\mu} - \mathbf{M}_0)$, which is a natural choice to characterize the direction of the asymmetry of the multivariate \mathcal{SN} distribution. Using the same reasoning for the \mathcal{MMN} distribution, we can consider $S_C = (E(U) - \frac{m_0^*}{\delta_*}) \delta$, and so, the direction of δ can be regarded as a measure of vectorial skewness for the \mathcal{MMN} distribution.

References

References

- [1] C.J. Adcock, Mean-variance-skewness efficient surfaces, Stein’s lemma and the multivariate extended skew-Student distribution, European Journal of Operational Research, 234 (2014) 392-401.

- [2] C.J. Adcock, M. Eling, N. Loperfido, Skewed distributions in finance and actuarial science: A review, *The European Journal of Finance*, 21 (2015) 1253-1281.
- [3] C.J. Adcock, K. Shutes, On the multivariate extended skew-normal, normal-exponential and normal-gamma distributions, *Journal of Statistical Theory and Practice*, 6 (2012) 636-664.
- [4] M.Y. An, Log-concave probability distributions: theory and statistical testing, Working Paper 96-01, CLS, Science Park Aarhus, Gustav Wieds Vej 10C, 8000 Aarhus C, Denmark, 1996.
- [5] R.B. Arellano-Valle, A. Azzalini, On the unification of families of skew-normal distributions, *Scandinavian Journal of Statistics*, 33 (2006) 561-574.
- [6] A. Azzalini, A class of distributions which includes the normal ones, *Scandinavian Journal of Statistics*, 2 (1985) 171-178.
- [7] A. Azzalini, Further results on a class of distributions which includes the normal ones. *Statistica*, 46 (1986) 199-208.
- [8] A. Azzalini, The skew-normal distribution and related multivariate families. *Scandinavian Journal of Statistics*, 32 (2005) 159-188.
- [9] A. Azzalini, A. Capitanio, *The Skew-Normal and Related Families*, Cambridge University Press, Cambridge, England, 2014.
- [10] A. Azzalini, A. Dalla Valle, The multivariate skew-normal distribution, *Biometrika*, 83 (1996) 715-726.
- [11] N. Balakrishnan, M.R. Brito, A.J. Quiroz, A vectorial notion of skewness and its use in testing for multivariate symmetry, *Communications in Statistics-Theory and Methods*, 36 (2007) 1757-1767.
- [12] N. Balakrishnan, B. Scarpa, Multivariate measures of skewness for the skew-normal distribution, *Journal of Multivariate Analysis*, 104 (2012) 73-87.
- [13] N. Balakrishnan, A. Capitanio, B. Scarpa, A test for multivariate skew-normality based on its canonical form, *Journal of Multivariate Analysis*, 128 (2014) 19-32.
- [14] O. Barndorff-Nielsen, J. Kent, M. Sørensen, Normal variance-mean mixtures and z distributions, *International Statistical Review*, 50 (1982) 145-159.
- [15] M. Bernardi, R. Cerqueti, A. Palestini, The skew normal multivariate risk measurement framework, *Computational Management Science*, 17 (2020) 105-119.
- [16] A. Bose, A. Dasgupta, H. Rubin, A contemporary review and bibliography of infinitely divisible distributions and processes, *Sankhyā, Series A*, 64 (2002) 763-819.
- [17] R.D. Cook, S. Weisberg, *An Introduction to Regression Graphics*, Wiley, New York, 1994.
- [18] M. Eling, Performance measurement in the investment industry, *Financial Analysts Journal*, 64 (2008) 54-66.
- [19] K.T. Fang, S. Kotz, K.W. Ng, *Symmetric Multivariate and Related Distributions*, Chapman and Hall, London, England, 1990.
- [20] C. Franceschini and N. Loperfido, *MaxSkew* and *MultiSkew*, Two R Packages for Detecting, Measuring and Removing Multivariate Skewness, *Symmetry* 11 (8) 970 (2019).
- [21] W. Fung, D.A. Hsieh, Performance characteristics of hedge funds and commodity funds: Natural vs. spurious biases, *Journal of Financial and Quantitative Analysis*, 35 (2000) 291-307.
- [22] M.G. Genton, L. He, X. Liu, Moments of skew-normal random vectors and their quadratic forms, *Statistics & Probability Letters*, 51 (2001) 319-325.
- [23] A. Graham, *Kronecker Products and Matrix Calculus: With Applications*, Halsted Press, New York, 1981.
- [24] N. Henze, Limit laws for multivariate skewness in the sense of Móri, Rohatgi, Székely, *Statistics & Probability Letters*, 33 (1997) 299-307.
- [25] T. Isogai, On a measure of multivariate skewness and a test for multivariate normality, *Annals of the Institute of Statistical Mathematics*, 34 (1982) 531-541.
- [26] H. M. Kim, J. Zhao, Multivariate measures of skewness for the scale mixtures of skew-normal distributions, *Communications for Statistical Applications and Methods*, 25 (2018) 109-130.
- [27] T. Kollo, Multivariate skewness and kurtosis measures with an application in ICA, *Journal of Multivariate Analysis*, 99 (2008) 2328-2338.
- [28] T. Kollo, M.S. Srivastava, Estimation and testing of parameters in multivariate Laplace distribution, *Communications in Statistics-Theory and Methods*, 33 (2004) 2363-2387.
- [29] T. Kollo, D. von Rosen, *Advanced Multivariate Statistics with Matrices*. Springer, Dordrecht, The Netherlands, 2005.
- [30] C. Liu, D.B. Rubin, The ECME algorithm: A simple extension of EM and ECM with faster monotone convergence, *Biometrika*, 81 (1994) 633-648.
- [31] N. Loperfido, Skewness-based projection pursuit: a computational approach, *Computational Statistics and Data Analysis*, 120 (2018) 42-57.
- [32] J.F. Malkovich, A.A. Afifi, On tests for multivariate normality, *Journal of the American Statistical Association*, 68 (1973) 176-179.
- [33] K.V. Mardia, Measures of multivariate skewness and kurtosis with applications, *Biometrika*, 36 (1970) 519-530.
- [34] K.V. Mardia, Applications of some measures of multivariate skewness and kurtosis in testing normality and robustness studies, *Sankhyā, Series B*, 36 (1974) 115-128.
- [35] X. L. Meng, D. B. Rubin, Maximum likelihood estimation via the ECM algorithm: A general framework, *Biometrika*, 80 (1993) 267-278.
- [36] T.F. Móri, V.K. Rohatgi, G.J. Székely, On multivariate skewness and kurtosis, *Theory of Probability and its Applications*, 38 (1993) 547-551.
- [37] H. Negarestani, A. Jamalizadeh, S. Shafiei, N. Balakrishnan, Mean mixtures of normal distributions: Properties, inference and application, *Metrika*, 82 (2019) 501-528.
- [38] A. Prékopa, On logarithmic concave measures and functions, *Acta Scientiarum Mathematicarum*, 33 (1973) 335-343.
- [39] J.R. Schott, *Matrix Analysis for Statistics*, Third edition, John Wiley & Sons, Hoboken, New Jersey, 2016.
- [40] Y. Simaan, Portfolio selection and asset pricing-three parameter framework, *Management Science*, 39 (1993) 568-587.
- [41] M.S. Srivastava, A measure of skewness and kurtosis and a graphical method for assessing multivariate normality, *Statistics & Probability Letters*, 2 (1984) 263-267.
- [42] F. W. Steutel, K. Van Harn, *Infinite Divisibility of Probability Distributions on the Real Line*, Marcel Dekker, New York, 2004.