

LIFESPAN ESTIMATES FOR 2-DIMENSIONAL SEMILINEAR WAVE EQUATIONS IN ASYMPTOTICALLY EUCLIDEAN EXTERIOR DOMAINS

NING-AN LAI, MENG-YUN LIU*, KYOUHEI WAKASA, AND CHENGBO WANG

ABSTRACT. In this paper we study the initial boundary value problem for two-dimensional semilinear wave equations with small data, in asymptotically Euclidean exterior domains. We prove that if $1 < p \leq p_c(2)$, the problem admits almost the same upper bound of the lifespan as that of the corresponding Cauchy problem, only with a small loss for $1 < p \leq 2$. It is interesting to see that the logarithmic increase of the harmonic function in 2-D has no influence to the estimate of the upper bound of the lifespan for $2 < p \leq p_c(2)$. One of the novelties is that we can deal with the problem with flat metric and general obstacles (bounded and simply connected), and it will be reduced to the corresponding problem with compact perturbation of the flat metric outside a ball.

CONTENTS

1. Introduction	2
2. Local well-posedness	7
3. Test function for the Dirichlet problem	8
3.1. Kelvin transformation	8
3.2. Test function for g_1	8
3.3. Proof of Lemma 1.3	9
3.4. An integral estimate	10
4. Proof of Theorem 1.1	10
4.1. Subcritical case	10
4.2. Critical case	12
5. Test function $\Delta_{g_1+g_3}\Phi_\lambda = \lambda^2\Phi_\lambda$	16
5.1. Spherically symmetric case	16
5.2. Derivative estimates	18
5.3. Test function $\Delta_{g_1+g_3}\Phi_\lambda = \lambda^2\Phi_\lambda$	19
5.4. Proof of Lemma 5.3	19
6. General obstacles	20
6.1. Star-shaped obstacles	20
6.2. General obstacles	21
References	22

Date: February 14, 2022.

2010 Mathematics Subject Classification. 35L05, 35L71, 35B30, 35B33, 35B40, 35B44.

Key words and phrases. exterior domain, Strauss conjecture, asymptotically Euclidean, Riemann mapping theorem, Kelvin transform.

* Corresponding author.

1. INTRODUCTION

In this paper, we are interested in the investigation of the blow-up part of the analogs of the Strauss conjecture in two dimensional asymptotically Euclidean exterior domain $(\Omega, \mathbf{g})$. We assume $\partial\Omega$ is a smooth Jordan curve. By asymptotically Euclidean exterior domain, we mean that it is a submanifold of the asymptotically Euclidean space $(\mathbb{R}^2, \mathbf{g})$. For simplicity, we assume $\Omega = \mathbb{R}^2 \setminus \overline{B_R}$ for some $R > 0$, see however, Lemma 1.6.

Recall that, for $(\mathbb{R}^2, \mathbf{g})$, the metric $\mathbf{g}$ is assumed to be of the form

$$(1.1) \quad \mathbf{g} = \mathbf{g}_1 + \mathbf{g}_2 ,$$

where $\mathbf{g}_1$ is a spherically symmetric, long range perturbation of the flat metric $\mathbf{g}_0$, and $\mathbf{g}_2$ is a short range perturbation. With possibly changing the choice of R , we could write $\mathbf{g}_1$, in terms of the polar coordinates $x = r(\cos \theta, \sin \theta) \in \Omega$, as follows

$$(1.2) \quad \mathbf{g}_1 = K^2(r)dr^2 + r^2d\theta^2 ,$$

where $d\theta^2$ is the standard metric on the unit circle $\mathbb{S}^1$, and

$$(1.3) \quad |\partial_r^m(K-1)| \lesssim \langle r \rangle^{-m-\rho_1}, m = 0, 1, 2,$$

for some given constant $\rho_1 \in (0, 1]$. Here and in what follows, $\langle x \rangle = \sqrt{1 + |x|^2}$, and we use $A \lesssim B$ ($A \gtrsim B$) to stand for $A \leq CB$ ($A \geq CB$) where the constant C may change from line to line. Concerning $\mathbf{g}_2$, we have

$$\mathbf{g} = g_{jk}(x)dx^jdx^k \equiv \sum_{j,k=1}^2 g_{jk}(x)dx^jdx^k , \quad \mathbf{g}_2 = g_{2,jk}(x)dx^jdx^k ,$$

where we have used the convention that Latin indices j, k range from 1 to 2 and the Einstein summation convention for repeated upper and lower indices. Furthermore, we assume $\mathbf{g}_2$ satisfies

$$(1.4) \quad \nabla^\beta g_{2,jk} = \mathcal{O}(\langle r \rangle^{-\rho_2-|\beta|}), |\beta| \leq 2 ,$$

for some $\rho_2 > 1$. By these assumptions, it is clear that there exists a constant $\delta_0 \in (0, 1)$ such that

$$(1.5) \quad \delta_0|\xi|^2 \leq g^{jk}(x)\xi_j\xi_k \leq \delta_0^{-1}|\xi|^2, \forall x, \xi \in \mathbb{R}^2, \quad K(r) \in (\delta_0, 1/\delta_0) ,$$

where $(g^{jk}(x))$ denotes the inverse of $(g_{jk}(x))$.

With these preparations in hand, we may write out our problem explicitly, that is, initial boundary value problem of semilinear wave equations with small initial data posed on asymptotically Euclidean manifolds $(\Omega, \mathbf{g})$ with $\mathbf{g}$ satisfies (1.1)-(1.4)

$$(1.6) \quad \begin{cases} u_{tt} - \Delta_{\mathbf{g}}u = |u|^p, & t > 0, \quad x \in \Omega, \\ u(0, x) = \varepsilon u_0(x), \quad u_t(0, x) = \varepsilon u_1(x), & x \in \Omega, \\ u(t, x) = 0, & t > 0, \quad x \in \partial\Omega, \end{cases}$$

where, $\Delta_{\mathbf{g}} = \nabla^j \partial_j$ is the standard Laplace-Beltrami operator, $\varepsilon > 0$ is a small parameter. Concerning the initial data, we assume

$$(1.7) \quad (u_0, u_1) \in H_0^1(\Omega) \times L^2(\Omega), \quad \text{supp}(u_0, u_1) \subset B_{R_0}, \quad u_0, u_1 \geq 0, u_0 \not\equiv 0, u_1 \not\equiv 0,$$

for some $R_0 > R$.

Such kind of problem is the generalization of the Strauss conjecture (see [34]): the following Cauchy problem of semilinear wave equation with small initial data with sufficient regularity and sufficient decay at infinity

$$(1.8) \quad \begin{cases} u_{tt} - \Delta u = |u|^p, & (t, x) \in (0, T) \times \mathbb{R}^n, \\ u(0, x) = \varepsilon u_0(x), \quad u_t(0, x) = \varepsilon u_1(x), & x \in \mathbb{R}^n, \end{cases}$$

admits a critical exponent $p_c(n)$ ($n \geq 2$), which means that if $1 < p \leq p_c(n)$ then problem (1.8) has no global solution in general, whereas the solution exists globally in time if $p > p_c(n)$ and $0 < \varepsilon \ll 1$. Here $p_c(n)$ is the positive root of the quadratic equation

$$(1.9) \quad (n-1)p^2 - (n+1)p - 2 = 0.$$

This conjecture has been essentially verified, we list all of the corresponding results in the following table (one can also find it in [35]):

n	$1 < p < p_c(n)$	$p = p_c(n)$	$p_c(n) < p \leq 1 + 4/(n-1)$
2	Glassey [7]	Schaeffer [28]	Glassey [8]
3	John [12]	Schaeffer [28]	John [12]
≥ 4	Sideris [29]	Yordanov-Zhang [42], Zhou [45], indep.	Georgiev-Lindblad-Sogge [5]

If there is no global solution, then it is interesting to estimate the time when the solution blows up, i.e., the lifespan. The results have been established for two cases: (I) subcritical power ($1 < p < p_c(n)$); (II) critical power ($p = p_c(n)$). For the former case, we now know that there exist two positive constants c and C such that the lifespan satisfies for $n \geq 2$ and $\max(1, 2/(n-1)) < p < p_c(n)$

$$(1.10) \quad c\varepsilon^{-\frac{2p(p-1)}{\gamma(n,p)}} \leq T(\varepsilon) \leq C\varepsilon^{-\frac{2p(p-1)}{\gamma(n,p)}},$$

where $\gamma(n, p) = 2 + (n+1)p - (n-1)p^2 > 0$. We may read the facts from the following table:

n	Lower bound	Upper bound
2	Zhou [46]	Zhou [46]
3	Lindblad [21]	Lindblad [21]
≥ 4	Lai-Zhou [14]	Takamura [35]

For relatively small power in dimension 2, we have richer results. For $(n, p) = (2, 2)$, Lindblad [21] obtained the following result

$$(1.11) \quad \begin{cases} \exists \lim_{\varepsilon \rightarrow 0^+} a(\varepsilon)^{-1} T(\varepsilon) > 0, & \text{if } \int_{\mathbb{R}^2} u_1(x) dx \neq 0, \\ \exists \lim_{\varepsilon \rightarrow 0^+} \varepsilon T(\varepsilon) > 0, & \text{if } \int_{\mathbb{R}^2} u_1(x) dx = 0, \end{cases}$$

where $a(\varepsilon)$ denotes a number satisfying

$$a^2 \varepsilon^2 \log(1+a) = 1.$$

For $1 < p < 2$ and $n = 2$, due to the results in Takamura [35] and Imai et al. [11], we have

$$(1.12) \quad \begin{cases} c\varepsilon^{-\frac{p-1}{3-p}} \leq T(\varepsilon) \leq C\varepsilon^{-\frac{p-1}{3-p}}, & \text{if } \int_{\mathbb{R}^2} u_1(x) dx \neq 0, \\ c\varepsilon^{-\frac{2p(p-1)}{\gamma(2,p)}} \leq T(\varepsilon) \leq C\varepsilon^{-\frac{2p(p-1)}{\gamma(2,p)}}, & \text{if } \int_{\mathbb{R}^2} u_1(x) dx = 0. \end{cases}$$

For the critical case ($p = p_c(n)$), the lifespan is much longer and has the form:

$$(1.13) \quad \exp(c\varepsilon^{-p(p-1)}) \leq T(\varepsilon) \leq \exp(C\varepsilon^{-p(p-1)}).$$

We have the following table:

n	Lower bound	Upper bound
2	Zhou [46]	Zhou [46]
3	Zhou [47]	Zhou [47]
≥ 4	Lindblad-Sogge [23] for $n \leq 8$ or radial solution	Takamura-Wakasa [36]

As waves propagate to infinity of the space, hence besides the Cauchy problem in the whole space, it is also interesting to consider the corresponding obstacle problem, i.e., the initial boundary value problem in exterior domain. Due to the difficulty caused by the boundary, such kind of problem has not been well understood, particularly for the global existence in high dimensions ($n \geq 5$). Anyway, we have the following results of global existence vs blow-up:

n	$1 < p < p_c(n)$	$p = p_c(n)$	$p > p_c(n)$
2	Li-Wang [19]	Lai-Zhou [17]	Smith-Sogge-Wang [31]
3	Zhou-Han [48]	Lai-Zhou [15]	Hidano et al [9]
4	Zhou-Han [48]	Sobajima-Wakasa [32]	Du et al [3], reproved by Hidano et al [9]
≥ 5	Zhou-Han [48]	Lai-Zhou [16], reproved by Sobajima-Wakasa [32]	$p = 2$, Metcalfe-Sogge [26], reproved by Wang [41]

Just like the Cauchy problem, it is meaningful to study the lifespan for the blow-up exponent. We expect the same estimate as that of the Cauchy problem, regardless of the boundary obstacle, at least when the obstacle is nontrapping. Denoting the expected sharp lower bound and upper bound by “L” and “U” respectively, we have the following known results:

n	$1 < p < p_c(n)$	$p = p_c(n)$
2	L : ? U : This work	L : ? U : This work
3	L : Du-Zhou [4] ($p = 2$), Yu [43] ($2 < p < 1 + \sqrt{2}$) Wang [41] ($2 \leq p < 1 + \sqrt{2}$) U : Zhou-Han [48]	L : Yu [43] ($T(\varepsilon) \geq \exp(c\varepsilon^{-\sqrt{2}})$), Improved by Wang [41] ($T(\varepsilon) \geq \exp(c\varepsilon^{-2\sqrt{2}})$) U : Lai-Zhou [15], reproved by Sobajima-Wakasa [32]
4	L : ? U : Zhou-Han [48]	L : Zha-Zhou [44] reproved by Wang [41] U : Sobajima-Wakasa [32]
≥ 5	L : ? U : Zhou-Han [48]	L : ? U : Lai-Zhou [16], reproved by Sobajima-Wakasa [32]

We also have to mention the generalization of problem (1.8) from Euclidean space to other manifold, such as asymptotically Euclidean manifolds (see [27], [33], [39] and references therein), and black hole spacetime (see [1], [20], [22], [25] and references therein). One can find a detailed description of such kind of generalization in a

recent survey paper [40]. Another direction is to consider the initial boundary value problem in exterior domain with big initial data (see [18], [30] and references therein).

Before stating our results, we shall make a hypothesis.

Hypothesis: There exists $\lambda_0 \in (0, 1/(2R))$, such that for any $\lambda \in (0, \lambda_0)$, we could find a solution ϕ_λ solving

$$(1.14) \quad \Delta_{\mathbf{g}} \phi_\lambda = \lambda^2 \phi_\lambda, x \in \Omega, \phi_\lambda|_{\partial\Omega} = 0,$$

which enjoys the following uniform estimates (independent of λ)

$$(H) \quad \phi_\lambda \sim \begin{cases} \frac{\ln r/R}{\ln(R\lambda)^{-1}} & r \leq \lambda^{-1}, \\ \langle r\lambda \rangle^{-1/2} e^{\lambda \int_R^r K(\tau) d\tau} & r \geq \lambda^{-1}, \end{cases}$$

for any $\lambda \in (0, \lambda_0)$.

Theorem 1.1. *Let $p \in (1, p_c(2)]$ and assume (H). Consider the problem (1.6) with initial data (1.7), posed on asymptotically Euclidean exterior domain $(\Omega, \mathbf{g})$ satisfying (1.1)-(1.4). Then we have the following*

- (1) *There exists a unique weak solution $u(t, x) \in C_t H_0^1 \cap C_t^1 L^2$ to the initial boundary value problem (1.6) on $[0, T(\varepsilon)) \times \Omega$, where $T(\varepsilon)$ denotes the lifespan, i.e., the maximal time of existence.*
- (2) *There exist constants $C, \varepsilon_0 > 0$ independent of ε such that for any $\varepsilon \in (0, \varepsilon_0)$, we have*

$$(1.15) \quad T(\varepsilon) \leq \begin{cases} \exp(C\varepsilon^{-p(p-1)}), & p = p_c(2), \\ C\varepsilon^{-\frac{2p(p-1)}{2+3p-p^2}}, & 2 \leq p < p_c(2), \\ C(\varepsilon^{-1} \ln(\varepsilon^{-1}))^{(p-1)/(3-p)}, & 1 < p < 2. \end{cases}$$

Remark 1.2. *Comparing with the upper bounds with that of the corresponding Cauchy problem, we see that we have the same upper bounds of the lifespan, when $2 < p \leq p_c(2)$ (with log loss for the $1 < p \leq 2$). It will be interesting to determine the sharp estimate of the lifespan. We expect that our upper bound is sharp, at least when the obstacle is nontrapping and $2 < p \leq p_c(2)$.*

The local well-posed result follows from a standard energy argument. For the upper bound of lifespan estimates, it relies on the so-called test function method. For the subcritical case, we have to show the proper asymptotic behaviors of two test functions, that is, (1.17) for ϕ_0 and (H) for ϕ_{λ_1} with some fixed $\lambda_1 \in (0, \lambda_0)$. However, for the critical case, we need to use a family of test function ϕ_λ with λ varying in $(0, \lambda_0)$, to construct another test function b_q with more subtle asymptotic behavior as stated in (4.8) and (4.9) below. Concerning ϕ_0 , we have

Lemma 1.3. *Let $(\Omega, \mathbf{g})$ be an asymptotically Euclidean exterior domain satisfying (1.1)-(1.4). Then there exists a solution ϕ_0 to*

$$(1.16) \quad \Delta_{\mathbf{g}} \phi_0 = 0, x \in \Omega, \phi_0|_{\partial\Omega} = 0$$

satisfying

$$(1.17) \quad \phi_0(x) \simeq \ln r/R.$$

Remark 1.4. *It will be clear from the proof that the first two upper bounds in (1.15) hold for non-negative data with either u_0 or u_1 nontrivial. While, for the last upper bound, we need only to assume non-negative data with $u_1 \neq 0$.*

Noting that Theorem 1.1 holds under the hypothesis (H). Let us review some cases where the assumption (H) is valid. It is well known for the Euclidean space $(\mathbb{R}^2, \mathfrak{g}_0)$, where ϕ_λ could be given by the spherical average of $e^{\lambda x \cdot \omega}$,

$$\phi_\lambda(x) = \int_{S^1} e^{\lambda x \cdot \omega} d\omega \sim \langle r\lambda \rangle^{-1/2} e^{\lambda r},$$

see Yordanov-Zhang [42]. When $\mathfrak{g}_3$ is an exponential perturbation, that is, there exists $\alpha > 0$ so that

$$(1.18) \quad |\nabla g_{3,jk}| + |g_{3,jk}| \lesssim e^{-\alpha \int_R^r K(\tau) d\tau},$$

the corresponding estimate for $(\mathbb{R}^2, \mathfrak{g}_1 + \mathfrak{g}_3)$ is recently obtained for $\mathfrak{g} = \mathfrak{g}_1 + \mathfrak{g}_3$ by Liu-Wang [24], while the case $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_3$ was obtained by Wakasa-Yordanov in [38]. Based on these results, we could verify the hypothesis (H), in the case of $\mathfrak{g} = \mathfrak{g}_1 + \mathfrak{g}_3$, and thus obtain the following

Theorem 1.5. *Let $\mathfrak{g} = \mathfrak{g}_1 + \mathfrak{g}_3$. Then the hypothesis (H) holds and we have the same results as that in Theorem 1.1.*

In the above, we have considered exclusively on the asymptotically Euclidean exterior domains to disk. It is natural to ask what happens for the general asymptotically Euclidean exterior domains $(D, \mathfrak{g})$. It turns out that, when ∂D is a smooth Jordan curve, the general problem could be reduced to the case of disk exterior. Actually, with the help of the Riemann mapping theorem and a cut-off argument, we could prove the following

Lemma 1.6. *Let $D \subset \mathbb{R}^2$ be exterior domain to a smooth Jordan curve ∂D . Then there exists a diffeomorphism mapping preserving the boundary*

$$(1.19) \quad \mathcal{A} : D \rightarrow \mathbb{R}^2 \setminus \overline{B_R},$$

for some $R > 0$, which is an identity map for $r \gg 1$.

With the help of this lemma, we could translate the problem (1.6) in the domain $(D, \mathfrak{g})$, into the corresponding problem for $(\Omega, \tilde{\mathfrak{g}})$, by change of variables with the new unknown function $w(x) = u(\mathcal{A}^{-1}(x)) : \Omega \rightarrow \mathbb{R}$. Thus, whenever we have (H) for $(\Omega, \tilde{\mathfrak{g}})$, the lifespan estimates (1.15) apply for $(D, \mathfrak{g})$.

In particular, the result applies for any exterior domains to smooth Jordan curves.

More precisely, let $\mathcal{K} \subset \mathbb{R}^2$ be a domain interior to a smooth Jordan curve, (that is, $\mathcal{K}$ is a nonempty, open, bounded, smooth, simply connected domain), we consider the following nonlinear wave equations with obstacle $\mathcal{K}$

$$(1.20) \quad \begin{cases} u_{tt} - \Delta u = |u|^p, & t > 0, \ x \in D = \mathbb{R}^2 \setminus \bar{\mathcal{K}}, \\ u(0, x) = \varepsilon u_0(x), \ u_t(0, x) = \varepsilon u_1(x), & x \in D, \\ u|_{\partial \mathcal{K}} = 0. \end{cases}$$

With the help of the transform $w(x) = u(\mathcal{A}^{-1}(x)) : \Omega \rightarrow \mathbb{R}$, we need only to consider the corresponding problem for $(\Omega, \tilde{\mathfrak{g}}_0)$. Thanks to Lemma 1.6, $\tilde{\mathfrak{g}}_0$ is a compact perturbation of $\mathfrak{g}_0$, which allows us to apply Theorem 1.5 to obtain the following result.

Theorem 1.7. *Let $\mathcal{K} \subset \mathbb{R}^2$ be nonempty bounded smooth simply connected domain. Consider (1.20) with initial data (1.7). Then we have the same lifespan estimates (1.15) for energy solutions.*

Remark 1.8. *When the spatial dimension is not greater than 4, all of the previous blow-up results and lifespan estimates for exterior problem with critical power heavily rely on the assumption that the obstacle is a ball (see [15], [17], [32]), under which they can construct some special test functions explicitly. In contrast, our results hold for very general obstacle in 2-D.*

Outline. Our paper is organized as follows. In the next section, we sketch the proof of local well-posedness for the energy solutions, by a standard energy argument. In particular, it shows the finite speed of propagation (2.5), for the solution. In Section 3, we prove the existence of test function ϕ_0 , Lemma 1.3, for the Dirichlet problem of the Laplace equation on $(\Omega, \mathbf{g})$. With the help of ϕ_0 , as well as the hypothesis (H), in Section 4, we present the proof of Theorem 1.1. In Section 5, we show that the hypothesis (H) holds, at least when the metric $\mathbf{g}$ is exponential perturbation of a spherically symmetric, long range asymptotically Euclidean metric $\mathbf{g} = \mathbf{g}_1 + \mathbf{g}_3$. At last, in Section 6, with the help of the Riemann mapping theorem, we prove Lemma 1.6, which enables us to reduce the problems with general obstacles to the problem exterior to a disk (keeping the metric near the spatial infinity).

2. LOCAL WELL-POSEDNESS

Before going to the proof of blow-up results, we first show the local well-posedness for problem (1.6), based on energy estimates. By multiplying the equation in (1.6) with $\partial_t u$, we get the energy estimate

$$(2.1) \quad \frac{1}{2} \frac{d}{dt} \int_{\Omega} (u_t^2 + g^{ij}(x) u_{x_i} u_{x_j}) dV_{\mathbf{g}} = \int_{\Omega} |u|^p u_t dV_{\mathbf{g}},$$

where $dV_{\mathbf{g}} = \sqrt{\det(\mathbf{g}_{jk}(x))} dx$ is the volume form with respect to the metric. Due to the assumption of $g^{ij}(x)$ in (1.5), this implies

$$(2.2) \quad \|\partial_t u\|_{L^\infty L^2([0,T] \times \Omega)} \lesssim \varepsilon + T \| |u|^p \|_{L^\infty L^2([0,T] \times \Omega)}.$$

On the other hand, it is easy to get

$$(2.3) \quad \begin{aligned} \|u\|_{L^\infty L^2([0,T] \times \Omega)} &\lesssim \|u_0\|_{L^2(\Omega)} + T \|u_t\|_{L^\infty L^2([0,T] \times \Omega)} \\ &\lesssim \varepsilon(1+T) + T^2 \| |u|^p \|_{L^\infty L^2([0,T] \times \Omega)}. \end{aligned}$$

It then follows by combining (2.2) and (2.3) that

$$(2.4) \quad \|u\|_{L^\infty H^1([0,T] \times \Omega)} \lesssim \varepsilon(1+T) + (T+T^2) \| |u|^p \|_{L^\infty L^2([0,T] \times \Omega)}.$$

Recalling that

$$H^1(\Omega) \hookrightarrow L^{2p}(\Omega), \quad 1 < p < \infty,$$

with the help of all these estimates, a standard contraction mapping argument yields the desired local well-posed result.

In particular, as the solution is obtained through iteration, we see that the solution enjoys the finite speed of propagation:

$$(2.5) \quad \text{supp } u(t) \subset \left\{ x \in \Omega : \int_R^t K(\tau) d\tau \leq t + R_1 \right\} := D_1(t) \subset \left\{ x : r \leq \frac{t + R_2}{\delta_0} \right\},$$

for any data satisfying (1.7), where $R_2 > R_1 = \int_R^{R_0} K(\tau) d\tau > 0$.

3. TEST FUNCTION FOR THE DIRICHLET PROBLEM

In this section, we consider the Laplace equation posed on the asymptotically Euclidean exterior domain $(\Omega, \mathfrak{g})$, that is, Lemma 1.3.

We observe that, for the purpose of the lemma, we need only to prove the following weaker estimate

$$\phi_0(x) \simeq \ln r, \quad r \gg 1.$$

Actually, by the strong maximum principle, we know that $\phi_0(x)$ is positive everywhere in Ω . Moreover, by Hopf's lemma, we have

$$\partial_r \phi_0(r, \theta)|_{r \rightarrow R_+} > 0,$$

which gives us that

$$\phi_0(r, \theta) \sim r - R \sim \ln r/R,$$

for any $R \leq r \lesssim 1$.

3.1. Kelvin transformation. At first, inspired by the classical Kelvin transform, we use the spatial inversion to introduce the conformal compactization. More specifically, we introduce new coordinate in the $\bar{B}_{1/R} \setminus \{0\}$ for Ω

$$(3.1) \quad \Phi(x) = \phi_0(|x|^{-2}x) = \phi_0(x^*), \quad x \in \bar{B}_{1/R} \setminus \{0\}.$$

Let $ds^2 = \mathfrak{g}_{jk}(x^*)dx^{*j}dx^{*k} = \tilde{h}_{jk}(x)dx^jdx^k$, where $x = r\omega$, $\mathfrak{g}_{jk}(x^*) = \delta_{jk} + f(x^*)\omega_j\omega_k + \mathfrak{g}_{2,jk}(x^*)$ and $f(x^*) = K^2(x^*) - 1$. By (1.3)-(1.4), we have

$$(3.2) \quad f(x^*) = \mathcal{O}(|x|^{\rho_1}), \mathfrak{g}_{2,jk}(x^*) = \mathcal{O}(|x|^{\rho_2}), \quad \rho_1 \in (0, 1], \rho_2 > 1,$$

as $x \rightarrow 0$. As

$$\frac{\partial x^{*j}}{\partial x^k} = \frac{\delta^{jk} - 2\omega^j\omega^k}{r^2},$$

we get

$$\tilde{h}_{jk}(x) = \frac{1}{r^4}(\delta^{ji} - 2\omega^j\omega^i)\mathfrak{g}_{il}(x^*)(\delta^{lk} - 2\omega^l\omega^k) := \frac{h_{jk}}{r^4},$$

where

$$(3.3) \quad \begin{aligned} h_{jk}(x) &= (\delta^{ji} - 2\omega^j\omega^i)\mathfrak{g}_{il}(x^*)(\delta^{lk} - 2\omega^l\omega^k) \\ &= \delta_{jk} + f(x^*)\omega_j\omega_k + \mathfrak{g}_{2,il}(x^*)(\delta^{ji} - 2\omega^j\omega^i)(\delta^{lk} - 2\omega^l\omega^k). \end{aligned}$$

Then we obtain

$$(3.4) \quad \Delta_{\mathfrak{g}}\phi_0(x^*) = \Delta_{\tilde{h}}\Phi(x) = r^4\Delta_h\Phi(x).$$

In particular, if we have $\mathfrak{g} = \mathfrak{g}_1$, there is h_0 such that

$$(3.5) \quad \Delta_{\mathfrak{g}_1}\phi(x^*) = r^4\Delta_{h_0}\Phi_0(x), \quad h_{0,jk}(x) = \delta_{jk} + f(x^*)\omega_j\omega_k.$$

3.2. Test function for $\mathfrak{g}_1$. Considering

$$\Delta_{\mathfrak{g}_1}\phi = 0, \quad x \in \Omega; \quad \phi = 0, \quad x \in \partial\Omega,$$

it is easy to find a spherically symmetric solution

$$(3.6) \quad \phi(r) = \int_R^r \frac{K(s)}{s} ds \simeq \ln \frac{r}{R},$$

as

$$(3.7) \quad \Delta_{\mathfrak{g}_1} = K^{-1}r^{-1}\partial_r K^{-1}r\partial_r + r^{-2}\partial_\theta^2.$$

By (3.5), we have

$$(3.8) \quad \Phi_0(x) = \phi\left(\frac{x}{|x|^2}\right) = \phi\left(\frac{1}{r}\right) = \int_R^{\frac{1}{r}} \frac{K(s)}{s} ds$$

satisfying

$$(3.9) \quad \begin{cases} \Delta_{h_0} \Phi_0 = 0, & x \in B_{1/R} \setminus \{0\}, \\ \Phi_0 = 0, & x \in \partial B_{1/R}. \end{cases}$$

3.3. Proof of Lemma 1.3. By (3.1)-(3.4), we are reduced to finding a solution in the region $B_{1/R} \setminus \{0\}$

$$(3.10) \quad \begin{cases} \Delta_h \Phi = 0, & x \in B_{1/R} \setminus \{0\}, \\ \Phi = 0, & x \in \partial B_{1/R} \end{cases}$$

satisfying $\Phi(x) \sim \ln \frac{1}{r}$ near $r = 0$. Let $u(x) = \Phi(x) - \Phi_0(x)$, it remains to construct a solution $u \in L^\infty(B_{1/R} \setminus \{0\})$, due to the fact that $\Phi_0 \sim \ln(1/r)$ near $r = 0$. Concerning u , it satisfies

$$\begin{cases} \Delta_h u = (\Delta_{h_0} - \Delta_h) \Phi_0, & x \in B_{1/R} \setminus \{0\}, \\ u = 0, & x \in \partial B_{1/R}, \end{cases}$$

and we would like to view it as the Dirichlet problem in the ball $B_{1/R}$.

For that purpose, we set $h_{jk}(0) = \delta_{jk}$ so that it is continuous in $B_{1/R}$, in view of (3.2)-(3.3). Moreover, we claim that

$$(3.11) \quad (\Delta_h - \Delta_{h_0}) \Phi_0 = \mathcal{O}(r^{\rho_2-2}),$$

which could be written in the form $\partial_1 F$ for some $F = \mathcal{O}(r^{\rho_2-1}) \in L^\infty(B_{1/R})$. With the help of the claim, by standard elliptic existence theorems, there is a unique solution $u \in H_0^1(B_{1/R})$. In addition, as the equation is of divergence form,

$$\partial_j (|h|^{1/2} h^{jk} \partial_k u) = |h|^{1/2} (\Delta_h - \Delta_{h_0}) \Phi_0 = \partial_1 F,$$

with $F \in L^\infty(B_{1/R})$, an application of Meyer's theorem (see, e.g., Taylor [37, Chapter 14, Proposition 12.2]) gives us $u \in W^{1,q}(B_{1/R})$ for some $q > 2$, which in turn gives us the desired result $u \in L^\infty(B_{1/R})$.

We are left to give the proof of the claim (3.11). By calculation, we have

$$(\Delta_h - \Delta_{h_0}) \Phi_0 = (h^{jk} - h_0^{jk}) \partial_j \partial_k \Phi_0 + [|h|^{-1/2} \partial_j (|h|^{1/2} h^{jk}) - |h_0|^{-1/2} \partial_j (|h_0|^{1/2} h_0^{jk})] \partial_k \Phi_0.$$

By (3.8) and (1.3), we get

$$\partial_j \Phi_0 = \mathcal{O}(r^{-1}), \partial_j \partial_k \Phi_0 = \mathcal{O}(r^{-2}).$$

Similarly, by (1.3)-(1.4), (3.3) and (3.5), we have

$$|\nabla^\beta (h_{jk} - h_{0,jk})| + |\nabla^\beta (h^{jk} - h_0^{jk})| + |\nabla^\beta (|h| - |h_0|)| = \mathcal{O}(r^{\rho_2-|\beta|}), |\beta| = 0, 1.$$

In summary, it is easy to see that

$$|(\Delta_h - \Delta_{h_0}) \Phi_0| \lesssim \sum_{j,k} \sum_{1 \leq |\beta| \leq 2, |\alpha|+|\beta|=2} |\nabla^\alpha \mathfrak{g}_2^{jk}(x^*)| |\nabla^\beta \Phi_0| = \mathcal{O}(r^{\rho_2-2}),$$

which completes the proof.

3.4. An integral estimate. With ϕ_0 and its asymptotic behavior, together with (H) for ϕ_λ , we will need the following

Lemma 3.1. *Let $\lambda_1 \in (0, \lambda_0)$, then we have*

$$(3.12) \quad \int_{\Omega \cap (\int_R^r K(s) ds \leq t+R_1)} \phi_0^{-\frac{1}{p-1}} e^{-p'\lambda_1 t} \phi_{\lambda_1}^{p'} dV_{\mathbf{g}} \lesssim (\ln(t+1))^{-\frac{1}{p-1}} (t+1)^{1-\frac{p'}{2}},$$

where $p' = \frac{p}{p-1}$ and the implicit constant may depend on λ_1 .

Proof. Let us begin with the region with $r \leq \lambda_1^{-1}$, in that case, by (1.17) and the first estimate in (H), we have $\phi_0^{-\frac{1}{p-1}} \phi_{\lambda_1}^{p'} \lesssim \phi_0(r) (\phi_0(\lambda_1^{-1}))^{-\frac{p}{p-1}} \lesssim \ln(r/R)$ and the integral is controlled by

$$\int_{R \leq r \leq \lambda_1^{-1}} \phi_0(r) e^{-p'\lambda_1 t} r dr \lesssim e^{-p'\lambda_1 t}.$$

For the remained case $r \geq \lambda_1^{-1}$, we could use (1.17) and the second estimate in (H). Thus we have

$$\int_{\{r \geq \lambda_1^{-1}\} \cap \{\int_R^r K(s) ds \leq \frac{t+R_1}{2}\}} \phi_0^{-\frac{1}{p-1}} e^{-p'\lambda_1 t} \phi_{\lambda_1}^{p'} dV_{\mathbf{g}} \lesssim \int_{r \lesssim t+R_2} e^{-\frac{p'\lambda_1 t}{2}} r dr \lesssim e^{-\frac{p'\lambda_1 t}{2}} (t+1)^2,$$

and

$$(3.13) \quad \begin{aligned} & \int_{\frac{t+R_1}{2} \leq \int_R^r K(s) ds \leq t+R_1} \phi_0^{-\frac{1}{p-1}} e^{-p'\lambda_1 t} \phi_{\lambda_1}^{p'} dV_{\mathbf{g}} \\ & \lesssim (\ln(t+1))^{-\frac{1}{p-1}} \int_{\frac{t+R_1}{2} \leq \int_R^r K(s) ds \leq t+R_1} e^{-p'\lambda_1 t + p'\lambda_1 \int_R^r K(s) ds} \langle \lambda_1 r \rangle^{1-\frac{p'}{2}} dr \\ & \lesssim (\ln(t+1))^{-\frac{1}{p-1}} (t+1)^{1-\frac{p'}{2}}. \end{aligned}$$

In summary, we obtain (3.12). $\square$

4. PROOF OF THEOREM 1.1

In this section, we are devoted to the proof for Theorem 1.1, under the assumption (H).

4.1. Subcritical case. Let $\eta(t) \in C^\infty([0, \infty))$ be a decreasing function satisfying

$$\eta(t) = \begin{cases} 1 & t \leq \frac{1}{2}, \\ 0 & t \geq 1, \end{cases} \quad \eta^*(t) = \eta(t) \chi_{[1/2, 1]}(t)$$

and

$$\eta_T(t) = \eta(t/T), \eta_T^*(t) = \eta^*(t/T), \quad T \in (1, T(\varepsilon)).$$

As $u \in C_t H_0^1 \cap C_t^1 L^2$ is the energy solution to (1.6), we have finite speed of propagation (2.5) and $u \in C_t^2 H^{-1}$ based on the equation. Using $\eta_T^{2p'}(t) \phi_0(x)$ as a test function, where $\phi_0(x)$ is the test function in Lemma 1.3, we obtain

$$\begin{aligned} & \langle |u|^p, \eta_T^{2p'}(t) \phi_0(x) \rangle_{\mathbf{g}} \\ &= \langle u_{tt} - \Delta_{\mathbf{g}} u, \eta_T^{2p'}(t) \phi_0(x) \rangle_{\mathbf{g}} \\ &= \frac{d}{dt} (\langle u_t, \eta_T^{2p'} \phi_0(x) \rangle_{\mathbf{g}} - \langle u, \partial_t \eta_T^{2p'} \phi_0(x) \rangle_{\mathbf{g}}) + \langle u, \square_{\mathbf{g}} (\eta_T^{2p'}(t) \phi_0(x)) \rangle_{\mathbf{g}} \\ &= \frac{d}{dt} (\langle u_t, \eta_T^{2p'} \phi_0(x) \rangle_{\mathbf{g}} - \langle u, \partial_t \eta_T^{2p'} \phi_0(x) \rangle_{\mathbf{g}}) + \langle u, \partial_t^2 \eta_T^{2p'}(t) \phi_0(x) \rangle_{\mathbf{g}} \end{aligned}$$

where $\square_{\mathfrak{g}} = \partial_t^2 - \Delta_{\mathfrak{g}}$, $\langle u, v \rangle_{\mathfrak{g}}$ denotes the dual relation between distribution u and test function v , which agrees with $\int_{\Omega} u(x)v(x)\sqrt{\det(\mathfrak{g}_{jk}(x))}dx = \int_{\Omega} u(x)v(x)dV_{\mathfrak{g}}$ for usual functions u, v . Integrating over $[0, T]$, and observing that $\eta_T(T) = \partial_t \eta_T(0) = \partial_t \eta_T(T) = 0$, we get

$$(4.1) \quad \int_0^T \int_{\Omega} |u|^p \eta_T^{2p'} \phi_0 dV_{\mathfrak{g}} dt + \varepsilon \int_{\Omega} u_1 \phi_0 dV_{\mathfrak{g}} = \int_0^T \int_{\Omega} u \partial_t^2 \eta_T^{2p'} \phi_0 dV_{\mathfrak{g}} dt .$$

Noting that

$$\partial_t^2 \eta_T^{2p'} = 2p'(2p' - 1) \eta_T^{2p'-2} (\partial_t \eta_T)^2 + 2p' \eta_T^{2p'-1} (\partial_t^2 \eta_T),$$

since $(\partial_t \eta_T)^2, |\partial_t^2 \eta_T| \leq CT^{-2}$ for some $C > 0$, and $\eta_T \leq 1$, then we have

$$\partial_t^2 \eta_T^{2p'} = \mathcal{O}(T^{-2} \eta_T^{2p'-2}).$$

Thus we obtain

$$\begin{aligned} & \int_0^T \int_{\Omega} u \partial_t^2 \eta_T^{2p'} \phi_0 dV_{\mathfrak{g}} dt \\ & \leq C \left(\int_0^T \int_{\Omega} |u|^p \eta_T^{2p'} \phi_0 dV_{\mathfrak{g}} dt \right)^{\frac{1}{p}} \left(\int_{\frac{T}{2}}^T \int_{D_1(t)} T^{-2p'} \phi_0 dV_{\mathfrak{g}} dt \right)^{\frac{1}{p'}} \\ & \leq CT^{3-2p'} \ln T + \frac{1}{3} \int_0^T \int_{\Omega} |u|^p \eta_T^{2p'} \phi_0 dV_{\mathfrak{g}} dt, \end{aligned}$$

where we have used Hölder's inequality, (1.17) and Young's inequality. Plugging it to (4.1), we know that

$$(4.2) \quad \varepsilon \int_{\Omega} u_1(x) \phi_0 dV_{\mathfrak{g}} + \int_0^T \int_{\Omega} |u|^p \eta_T^{2p'} \phi_0 dV_{\mathfrak{g}} dt \leq CT^{3-2p'} \ln T.$$

This inequality implies the lifespan estimate for $1 < p < 2$ in (1.15). Actually, from the last inequality we conclude that $T^{3-2p'} \ln T \gtrsim \varepsilon$ for any $T \in (1, T(\varepsilon))$, which gives us, for $1 < p < 3$,

$$(4.3) \quad T(\varepsilon) \lesssim (\varepsilon^{-1} \ln(\varepsilon^{-1}))^{(p-1)/(3-p)} .$$

But, comparing to the one

$$(4.4) \quad T(\varepsilon) \lesssim \varepsilon^{-\frac{2p(p-1)}{2+3p-p^2}},$$

which we will get for $1 < p < p_c(2)$, the lifespan estimate (4.3) is better than (4.4) only for $1 < p < 2$.

To proceed, we introduce another test function

$$\eta_T^{2p'} \psi(t, x) = \eta_T^{2p'} e^{-\lambda_1 t} \phi_{\lambda_1}(x)$$

to get

$$\begin{aligned}
& \int_0^T \int_{\Omega} |u|^p \eta_T^{2p'} \psi(t, x) dV_{\mathfrak{g}} dt \\
&= \int_0^T \langle u_{tt} - \Delta_{\mathfrak{g}} u, \eta_T^{2p'} \psi(t, x) \rangle_{\mathfrak{g}} dt, \\
&= \left(\langle u_t, \eta_T^{2p'} \psi(t, x) \rangle_{\mathfrak{g}} - \langle u, \partial_t (\eta_T^{2p'} \psi(t, x)) \rangle_{\mathfrak{g}} \right) \Big|_{t=0}^T + \int_0^T \langle u, \square_{\mathfrak{g}} (\eta_T^{2p'} \psi(t, x)) \rangle_{\mathfrak{g}} dt \\
&= -\varepsilon \int_{\Omega} (\lambda_1 u_0(x) + u_1(x)) \phi_{\lambda_1}(x) dV_{\mathfrak{g}} \\
&\quad + \int_0^T \int_{\Omega} u \psi \partial_t^2 \eta_T^{2p'} dV_{\mathfrak{g}} dt + 2 \int_0^T \int_{\Omega} u (\partial_t \eta_T^{2p'}) (\partial_t \psi) dV_{\mathfrak{g}} dt,
\end{aligned}$$

which yields

$$\begin{aligned}
(4.5) \quad & \varepsilon \int_{\Omega} (\lambda_1 u_0(x) + u_1(x)) \phi_{\lambda_1}(x) dV_{\mathfrak{g}} + \int_0^T \int_{\Omega} |u|^p \eta_T^{2p'} e^{-\lambda_1 t} \phi_{\lambda_1}(x) dV_{\mathfrak{g}} dt \\
&= \int_0^T \int_{\Omega} u e^{-\lambda_1 t} \phi_{\lambda_1}(x) (\partial_t^2 \eta_T^{2p'} - 2\lambda_1 \partial_t \eta_T^{2p'}) dV_{\mathfrak{g}} dt.
\end{aligned}$$

Noticing that

$$\partial_t^2 \eta_T^{2p'} - 2\lambda_1 \partial_t \eta_T^{2p'} = \mathcal{O}(T^{-1}(\eta_T^*)^{2p'-2}),$$

As above, by combining Hölder's inequality and (3.12), the right hand side of (4.5) is controlled by

$$\begin{aligned}
(4.6) \quad & CT^{-1} \int_0^T \int_{\Omega} |u| (\eta_T^*)^{2p'-2} \phi_0^{1/p} \phi_0^{-1/p} e^{-\lambda_1 t} \phi_{\lambda_1}(x) dV_{\mathfrak{g}} dt \\
&\leq CT^{-1} \|u (\eta_T^*)^{2p'-2} \phi_0^{1/p}\|_{L^p([0, T] \times \Omega)} \|\phi_0^{-1/p} e^{-\lambda_1 t} \phi_{\lambda_1}\|_{L^{p'}(\cup_{t \in [T/2, T]} D_1(t))} \\
&\leq CT^{-1+(2-\frac{p'}{2})\frac{1}{p'}} (\ln T)^{-\frac{1}{p}} \left(\int_0^T \int_{\Omega} |u|^p (\eta_T^*)^{2p'} \phi_0 dV_{\mathfrak{g}} dt \right)^{\frac{1}{p}}.
\end{aligned}$$

We then conclude from (4.5), (4.6) and (4.2) that

$$(4.7) \quad \varepsilon^p T^{2-\frac{p}{2}} \ln T \lesssim \int_0^T \int_{\Omega} |u|^p (\eta_T^*)^{2p'} \phi_0 dV_{\mathfrak{g}} dt \lesssim T^{3-2p'} \ln T,$$

which gives us the desired lifespan estimate (4.4) for $1 < p < p_c(2)$.

4.2. Critical case. For the lifespan estimate of the critical power, we need one more lemma, which is similar to Lemma 4.2 in [13]. For completeness, we give a proof.

Lemma 4.1. *Let $q > 0$, $\lambda_1 \in (0, \lambda_0)$ and*

$$b_q(t, x) = \int_0^{\lambda_1} e^{-\lambda t} \phi_{\lambda}(x) \lambda^{q-1} d\lambda,$$

then we have

(1) $b_q(t, x)$ *satisfies following identities*

$$\partial_t b_q(t, x) = -b_{q+1}(t, x), \quad \partial_t^2 b_q(t, x) = b_{q+2}(t, x) = \Delta_{\mathfrak{g}} b_q(t, x).$$

(2) For $x \in D_1(t)$, we have

$$(4.8) \quad b_q(t, x) \lesssim \begin{cases} (t + R_1)^{-q} & \text{if } 0 < q < \frac{1}{2}, \\ (t + R_1)^{-\frac{1}{2}} (t + R_1 + 1 - \int_R^r K(s) ds)^{\frac{1}{2}-q} & \text{if } q > \frac{1}{2}. \end{cases}$$

and

$$(4.9) \quad b_q(t, x) \gtrsim \frac{\phi_0}{\ln(t + R_1)} (t + R_1)^{-q}, \quad \forall q > 0.$$

Proof. The first part is trivial. Concerning (4.9), let $\delta_1 < \min(\delta_0 R_1 / R_2, \lambda_1 R_1)$, such that $\frac{\delta_1}{t + R_1} \leq \frac{\delta_1}{R_1} \leq \lambda_1$, then for any $\lambda \leq \frac{\delta_1}{t + R_1}$, we have $r \leq (t + R_2) / \delta_0 < (t + R_1) / \delta_1 \leq \lambda^{-1}$ for $x \in D_1(t)$, so that we could apply (H) and (1.17) to obtain

$$\begin{aligned} b_q(t, x) &\gtrsim \int_{\frac{\delta_1}{2(t+R_1)}}^{\frac{\delta_1}{t+R_1}} e^{-\lambda t} \frac{\phi_0(x)}{\ln(\lambda R)^{-1}} \lambda^{q-1} d\lambda \\ &\gtrsim \frac{\phi_0(x)}{\ln(t + R_1)} \int_{\frac{\delta_1}{2(t+R_1)}}^{\frac{\delta_1}{t+R_1}} e^{-\lambda(t+R_1)} \lambda^{q-1} d\lambda \\ &\gtrsim \frac{\phi_0(x)}{\ln(t + R_1)} (t + R_1)^{-q} \int_{\frac{\delta_1}{2}}^{\delta_1} e^{-\theta} \theta^{q-1} d\theta \gtrsim \frac{\phi_0(x)}{\ln(t + R_1)} (t + R_1)^{-q}. \end{aligned}$$

Turning to the upper bound, (4.8), we know from (H) that

$$(4.10) \quad b_q(t, x) \lesssim \int_0^{\lambda_1} e^{-\lambda(t+R_1)} e^{\lambda \int_R^r K(s) ds} \langle \lambda r \rangle^{-\frac{1}{2}} \lambda^{q-1} d\lambda.$$

If $\int_R^r K(s) \leq \frac{t+R_1}{2}$, we get

$$(4.11) \quad b_q(t, x) \lesssim \int_0^{\lambda_1} e^{-\frac{\lambda(t+R_1)}{2}} \lambda^{q-1} d\lambda \lesssim (t + R_1)^{-q} \int_0^\infty e^{-\lambda} \lambda^{q-1} d\lambda \lesssim (t + R_1)^{-q}.$$

For the case $\frac{t+R_1}{2} \leq \int_R^r K(s) \leq t + R_1$, we have $r \sim t + R_1$. If $q < 1/2$, the estimate for (4.10) becomes

$$b_q(t, x) \lesssim \int_0^{\lambda_1} \langle \lambda(t + R_1) \rangle^{-\frac{1}{2}} \lambda^{q-1} d\lambda \lesssim (t + R_1)^{-q} \int_0^\infty \langle s \rangle^{-\frac{1}{2}} s^{q-1} ds \lesssim (t + R_1)^{-q}.$$

It remains to consider the case $q > \frac{1}{2}$ and $r \sim t + R_1$, for which we have

$$\begin{aligned} b_q(t, x) &\lesssim (t + R_1)^{-\frac{1}{2}} \int_0^{\lambda_1} e^{-\lambda[t+R_1+1-\int_R^r K(s) ds]} \lambda^{q-3/2} d\lambda \\ &\lesssim (t + R_1)^{-\frac{1}{2}} \left(t + R_1 + 1 - \int_R^r K(s) ds \right)^{\frac{1}{2}-q} \int_0^\infty e^{-\lambda} \lambda^{q-\frac{3}{2}} d\lambda \\ &\lesssim (t + R_1)^{-\frac{1}{2}} \left(t + R_1 + 1 - \int_R^r K(s) ds \right)^{\frac{1}{2}-q}. \end{aligned}$$

This completes the proof of (4.8). $\square$

For $M \in [2, T] \subset [2, T(\varepsilon))$, we set

$$Y[|u|^p b_q](M) = \int_1^M \left(\int_0^T \int_\Omega |u|^p(t, x) b_q(t, x) (\eta_\sigma^*(t))^{2p'} dV_{\mathfrak{g}} dt \right) \sigma^{-1} d\sigma,$$

with $q = \frac{1}{2} - \frac{1}{p}$. Then from (4.9) and (4.7), we know that

$$\begin{aligned}
 (4.12) \quad M \frac{dY}{dM} &= \int_0^T \int_{\Omega} |u|^p b_q (\eta_M^*)^{2p'} dV_{\mathfrak{g}} dt \\
 &\geq \int_{M/2}^M \int_{\Omega} |u|^p \frac{\phi_0}{\ln M} M^{-q} (\eta_M^*)^{2p'} dV_{\mathfrak{g}} dt \\
 &\gtrsim \varepsilon^p M^{2-\frac{p}{2}} \ln M \frac{M^{\frac{1}{p}-\frac{1}{2}}}{\ln M} = \varepsilon^p M^{\frac{3}{2}-\frac{p}{2}+\frac{1}{p}} = \varepsilon^p,
 \end{aligned}$$

where we used the fact that $p = p_c(2)$. On the other hand, as in (4.10) in [13], we have

$$\begin{aligned}
 Y(M) &= \int_1^M \left(\int_0^T \int_{\Omega} b_q |u|^p (\eta_{\sigma}^*(t))^{2p'} dV_{\mathfrak{g}} dt \right) \sigma^{-1} d\sigma \\
 &= \int_0^T \int_{\Omega} b_q |u|^p \int_1^M (\eta_{\sigma}^*(t))^{2p'} \sigma^{-1} d\sigma dV_{\mathfrak{g}} dt \\
 &= \int_0^T \int_{\Omega} b_q |u|^p \int_{\frac{t}{M}}^t (\eta^*(s))^{2p'} s^{-1} ds dV_{\mathfrak{g}} dt \\
 &= \int_0^T \int_{\Omega} b_q |u|^p \int_{\max(\frac{t}{M}, \frac{1}{2})}^1 (\eta(s))^{2p'} s^{-1} ds dV_{\mathfrak{g}} dt \\
 &\leq \int_0^T \int_{\Omega} b_q |u|^p (\eta(t/M))^{2p'} \int_{\frac{1}{2}}^1 s^{-1} ds dV_{\mathfrak{g}} dt \leq 2 \int_0^T \int_{\Omega} \eta_M^{2p'} b_q |u|^p dV_{\mathfrak{g}} dt.
 \end{aligned}$$

As $(\partial_t^2 - \Delta_{\mathfrak{g}})b_q = 0$, we get

$$\begin{aligned}
 (4.13) \quad Y(M) &\lesssim \int_0^T \langle |u|^p, \eta_M^{2p'} b_q \rangle_{\mathfrak{g}} dt = \int_0^T \langle \partial_t^2 u - \Delta_{\mathfrak{g}} u, b_q \eta_M^{2p'} \rangle_{\mathfrak{g}} dt \\
 &= - \left(\varepsilon \int_{\Omega} u_1(x) b_q(0, x) dV_{\mathfrak{g}} + \varepsilon \int_{\Omega} u_0(x) b_{q+1}(0, x) dV_{\mathfrak{g}} \right) \\
 &\quad + \int_0^T \int_{\Omega} u (\partial_t^2 - \Delta_{\mathfrak{g}}) (b_q \eta_M^{2p'}) dV_{\mathfrak{g}} dt \\
 &\leq \int_0^T \int_{\Omega} u \left(2\partial_t b_q \partial_t \eta_M^{2p'} + b_q \partial_t^2 \eta_M^{2p'} \right) dV_{\mathfrak{g}} dt := I_1 + I_2.
 \end{aligned}$$

By (4.8) and (4.9), we know that

$$\begin{aligned}
& \int_{\frac{M}{2}}^M \int_{D_1(t)} b_q^{-\frac{1}{p-1}} b_{q+1}^{\frac{p}{p-1}} dV_{\mathfrak{g}} dt \\
& \lesssim \int_{\frac{M}{2}}^M \int_{D_1(t)} b_q^{-\frac{1}{p-1}} b_{q+1}^{\frac{p}{p-1}} r dr dt \\
& \lesssim \int_{\frac{M}{2}}^M \int_{D_1(t)} \left(\frac{\ln(t+R_1)}{\ln r/R} \right)^{\frac{1}{p-1}} \frac{(t+R_1)^{\frac{1}{2}-\frac{1}{p(p-1)}}}{t+R_1+1-\int_R^r K(s)ds} dr dt \\
& \lesssim \int_{\frac{M}{2}}^M (\ln(t+R_1))^{\frac{1}{p-1}} (t+R_1)^{\frac{1}{2}-\frac{1}{p(p-1)}} \int_{\int_R^r K(s)ds \leq \frac{t+R_1}{2}} r^{-1} (\ln(r/R))^{-\frac{1}{p-1}} dr dt \\
& \quad + \int_{\frac{M}{2}}^M (t+R_1)^{\frac{1}{2}-\frac{1}{p(p-1)}} \int_{\frac{t+R_1}{2}}^{t+R_1} (t+R_1+1-s)^{-1} ds dt \\
& \lesssim M^{\frac{3p(p-1)-2}{2p(p-1)}} \ln M = M^{\frac{p}{p-1}} \ln M,
\end{aligned}$$

then I_1, I_2 can be estimated as

$$\begin{aligned}
I_1 & \lesssim M^{-1} \left(\int_0^T \int_{\Omega} |u|^p b_q (\eta_M^*)^{2p'} dV_{\mathfrak{g}} dt \right)^{\frac{1}{p}} \left(\int_{\frac{M}{2}}^M \int_{D_1(t)} b_q^{-\frac{1}{p-1}} b_{q+1}^{\frac{p}{p-1}} dV_{\mathfrak{g}} dt \right)^{\frac{p-1}{p}} \\
& \lesssim (\ln M)^{\frac{p-1}{p}} \left(\int_0^T \int_{\Omega} |u|^p b_q (\eta_M^*)^{2p'} dV_{\mathfrak{g}} dt \right)^{\frac{1}{p}}
\end{aligned}$$

and

$$\begin{aligned}
I_2 & \lesssim M^{-2} \left(\int_0^T \int_{\Omega} |u|^p b_q \eta_M^{*2p'} dV_{\mathfrak{g}} dt \right)^{\frac{1}{p}} \left(\int_{\frac{M}{2}}^M \int_{D_1(t)} b_q dV_{\mathfrak{g}} dt \right)^{\frac{p-1}{p}} \\
& \lesssim M^{-2} \left(\int_0^T \int_{\Omega} |u|^p b_q \eta_M^{*2p'} dV_{\mathfrak{g}} dt \right)^{\frac{1}{p}} \left(\int_{\frac{M}{2}}^M \int_{\int_R^r K(s)ds \leq t+R_1} (t+R_1)^{-q} r dr dt \right)^{\frac{p-1}{p}} \\
& \lesssim \left(\int_0^T \int_{\Omega} |u|^p b_q \eta_M^{*2p'} dV_{\mathfrak{g}} dt \right)^{\frac{1}{p}}.
\end{aligned}$$

In summary, recalling (4.12), we obtain

$$MY'(M) = \int_0^T \int_{\Omega} |u|^p b_q (\eta_M^*)^{2p'} dV_{\mathfrak{g}} dt \gtrsim (\log M)^{1-p} Y^p(M), \quad MY'(M) \gtrsim \varepsilon^p,$$

for any $M \in [2, T] \subset [2, T(\varepsilon)]$. Let $M = T$, we see that

$$(4.14) \quad Y^p(T) \leq CT (\log T)^{p-1} Y'(T), \quad \varepsilon^p \lesssim TY'(T),$$

for any $T \in [2, T(\varepsilon)]$. With the help of (4.14), the desired lifespan estimate

$$T(\varepsilon) \leq \exp(c\varepsilon^{-p(p-1)})$$

for $p = p_c(2)$ follows directly from the following lemma with $p_1 = p_2 = p_c(2)$ and $\delta = \varepsilon^p$.

Lemma 4.2. (Lemma 3.10 in [10]) *Let $2 < t_0 < T$, $0 \leq \phi \in C^1([t_0, T])$. Assume that*

$$(4.15) \quad \begin{cases} \delta \leq K_1 t \phi'(t), & t \in (t_0, T), \\ \phi(t)^{p_1} \leq K_2 t (\log t)^{p_2-1} \phi'(t), & t \in (t_0, T) \end{cases}$$

with $\delta, K_1, K_2 > 0$ and $p_1, p_2 > 1$. If $p_2 < p_1 + 1$, then there exist positive constants δ_0 and K_3 (independent of δ) such that

$$(4.16) \quad T \leq \exp \left(K_3 \delta^{-\frac{p_1-1}{p_1-p_2+1}} \right)$$

when $0 < \delta < \delta_0$.

5. TEST FUNCTION $\Delta_{\mathbf{g}_1+\mathbf{g}_3} \Phi_\lambda = \lambda^2 \Phi_\lambda$

In this section, we consider the generalized eigenvalue problem of the Laplacian operator on $(\Omega, \mathbf{g})$, Theorem 1.5, when the metric $\mathbf{g}$ is an exponential perturbation of the spherically symmetric, long range asymptotically Euclidean metric $\mathbf{g}_1$.

5.1. Spherically symmetric case. Let us begin with the spherically symmetric case, considering (1.14) with $\Omega = \overline{B_R}^c$ and $\mathbf{g} = \mathbf{g}_1$:

$$\Delta_{\mathbf{g}_1} \phi_\lambda = \lambda^2 \phi_\lambda, x \in \Omega; \quad \phi_\lambda = 0, x \in \partial\Omega.$$

If we look at the corresponding problem on $\mathbb{R}^2$

$$\Delta_{\mathbf{g}_1} h_\lambda = \lambda^2 h_\lambda, x \in \mathbb{R}^2; \quad h_\lambda = 1, r = \lambda^{-1},$$

it is known from Liu-Wang [24] that there exists a small positive constant λ_2 such that

$$h_\lambda(r) \simeq \langle r\lambda \rangle^{-1/2} e^{\lambda \int_R^r K(s) ds}, \quad \forall \lambda \in (0, \lambda_2).$$

By comparing the desired solution with h_λ , we shall prove that there exists ϕ_λ such that

$$(5.1) \quad \phi_\lambda \leq h_\lambda, \phi_\lambda \geq h_\lambda - h_\lambda(R),$$

$$(5.2) \quad \phi_\lambda \sim \begin{cases} \frac{\phi_0(r)}{\phi_0(\lambda^{-1})} \simeq \frac{\ln r/R}{\ln(R\lambda)^{-1}} & r \leq \lambda^{-1}, \\ h_\lambda \simeq \langle r\lambda \rangle^{-1/2} e^{\lambda \int_R^r K(s) ds} & r \geq \lambda^{-1}, \end{cases}$$

for $\lambda \in (0, \min(1/(2R), \lambda_2))$.

Lemma 5.1. *Let $n = 2$. Then for any $\lambda \in (0, \lambda_2)$, the following boundary value problem*

$$(5.3) \quad \begin{cases} \Delta_{\mathbf{g}_1} \phi_\lambda(x) = \lambda^2 \phi_\lambda(x), & x \in \overline{B_R}^c, \\ \phi_\lambda(x) \Big|_{\partial B_R} = 0, \\ \phi_\lambda(x) - h_\lambda(x) \rightarrow 0, & |x| \rightarrow \infty \end{cases}$$

admits a spherically symmetric solution satisfying

$$(5.4) \quad 0 \leq h_\lambda(r) - h_\lambda(R) \leq \phi_\lambda(x) \leq h_\lambda(r), \quad x \in B_R^c.$$

Proof. Firstly we consider the following boundary value problem

$$(5.5) \quad \begin{cases} \Delta_{\mathbf{g}_1} \tilde{\phi}_\lambda(x) = \lambda^2 \tilde{\phi}_\lambda(x), & x \in \overline{B_R^c}, \\ \tilde{\phi}_\lambda(x) \Big|_{\partial B_R} = -h_\lambda(x) \Big|_{\partial B_R}, \\ \tilde{\phi}_\lambda(x) \rightarrow 0, & |x| \rightarrow \infty. \end{cases}$$

By standard variational argument with functional $I[u] = \int_{\overline{B_R^c}} (\lambda^2 |u|^2 + g_1^{jk} \nabla_j u \nabla_k u) dV_{\mathbf{g}_1}$, we see that there admits a unique solution $\tilde{\phi}_\lambda(x) \in H^1(\overline{B_R^c}) \cap C^\infty(\overline{B_R^c})$. Let

$$\phi_\lambda(x) = \tilde{\phi}_\lambda(x) + h_\lambda(x),$$

then $\phi_\lambda(x)$ satisfies the boundary value problem (5.3).

It remains to prove

$$\tilde{\phi}_\lambda(x) \in [-h_\lambda(R), 0], \quad x \in \overline{B_R^c}.$$

For any $C_0 > 0$, by (5.5), we know that

$$\begin{cases} -\Delta_{\mathbf{g}_1} (\tilde{\phi}_\lambda(x) - C_0) + \lambda^2 (\tilde{\phi}_\lambda(x) - C_0) = -\lambda^2 C_0 < 0, & x \in \overline{B_R^c}, \\ (\tilde{\phi}_\lambda(x) - C_0) \Big|_{\partial B_R} = (-h_\lambda(x) - C_0) \Big|_{\partial B_R} < 0, \\ \tilde{\phi}_\lambda(x) - C_0 \rightarrow -C_0 < 0, & |x| \rightarrow \infty. \end{cases}$$

By maximum principle, we have

$$\tilde{\phi}_\lambda(x) - C_0 \leq 0, \quad x \in \overline{B_R^c},$$

for any $C_0 > 0$, which means

$$\tilde{\phi}_\lambda(x) \leq 0, \quad x \in \overline{B_R^c}.$$

Similarly, it is clear that

$$\begin{cases} -\Delta_{\mathbf{g}_1} (\tilde{\phi}_\lambda(x) + h_\lambda(R)) + \lambda^2 (\tilde{\phi}_\lambda(x) + h_\lambda(R)) = \lambda^2 h_\lambda(R) > 0, & x \in \overline{B_R^c}, \\ (\tilde{\phi}_\lambda(x) + h_\lambda(R)) \Big|_{\partial B_R} = (-h_\lambda(x) + h_\lambda(R)) \Big|_{\partial B_R} = 0, \\ \tilde{\phi}_\lambda(x) + h_\lambda(R) \rightarrow h_\lambda(R) > 0, & |x| \rightarrow \infty. \end{cases}$$

By maximum principle, we have

$$\tilde{\phi}_\lambda(x) + h_\lambda(R) \geq 0, \quad x \in \overline{B_R^c},$$

and this completes the proof of Lemma 5.1. $\square$

With the help of Lemma 5.1, we get

$$(5.6) \quad \phi_\lambda(x) \simeq h_\lambda(r),$$

for any $r \geq \lambda^{-1}$ with $\lambda < \min(\lambda_2, (2R)^{-1})$.

Lemma 5.2. *Let $\phi_\lambda(x)$ be the function in Lemma 5.1 and $0 < \lambda < \min(\lambda_2, (2R)^{-1})$. Then we have*

$$(5.7) \quad \phi_\lambda(x) \simeq \frac{\ln r/R}{\ln 1/(\lambda R)}, \quad \forall R \leq r \leq \lambda^{-1}.$$

Proof. Let $\delta > 0$ to be determined and

$$F_\lambda(x) = \delta \frac{\phi_0(x)}{\phi_0(\lambda^{-1})} h_\lambda(x) - \phi_\lambda(x),$$

then, by (3.7),

$$\begin{cases} -\Delta_{\mathfrak{g}_1} F_\lambda + \lambda^2 F_\lambda(x) = -2 \frac{\delta}{K^2(r) \ln(\lambda^{-1})} \partial_r h_\lambda(x) \partial_r \phi_0 < 0, & x \in \overline{B_R^c}, \\ F_\lambda(x) = 0, & |x| = R, \\ F_\lambda(x) = \delta - \phi_\lambda(x) \leq 0, & |x| = \lambda^{-1}, \end{cases}$$

when $\delta > 0$ is sufficiently small such that $\phi_\lambda(x) \geq \delta = \delta h_\lambda(r)$ for $r = \lambda^{-1}$, in view of (5.6). By the maximum principle, one obtain

$$F_\lambda(x) \leq 0, \phi_\lambda(x) \geq \delta \frac{\phi_0(x)}{\phi_0(\lambda^{-1})} h_\lambda(x) \geq \delta h_\lambda(0) \frac{\phi_0(x)}{\phi_0(\lambda^{-1})} \geq \delta' \frac{\ln r/R}{\ln(\lambda R)^{-1}}, \quad \forall R \leq r \leq \lambda^{-1},$$

for some $\delta' > 0$, which yields the desired lower bound.

On the other hand, let

$$G_\lambda(x) = \phi_\lambda(x) - \frac{\phi_0}{\phi_0(\lambda^{-1})},$$

then

$$\begin{cases} -\Delta_{\mathfrak{g}_1} G_\lambda + \lambda^2 G_\lambda(x) = -\lambda^2 \frac{\phi_0}{\phi_0(\lambda^{-1})} < 0, & x \in \overline{B_R^c}, \\ G_\lambda(x) = 0, & |x| = R, \\ G_\lambda(x) = \phi_\lambda(x) - 1 \leq 0, & |x| = \lambda^{-1}, \end{cases}$$

which gives us $G_\lambda(x) \leq 0$. □

5.2. Derivative estimates. For future reference, we need to obtain more information concerning the behavior of ϕ_λ . At first, we claim that we have the following derivative estimates

$$(5.8) \quad \partial_r \phi_\lambda(r) \leq D_0 \lambda^2 (r - R) \phi_\lambda(r), |\partial_r^2 \phi_\lambda(r)| \leq D_0 \lambda^2 \phi_\lambda(r), \quad \forall r \geq R$$

for some constant D_0 independent of $\lambda \in (0, 1/(2R)]$.

As ϕ_λ is radial, by (1.2) and (3.7), we have

$$\Delta_{\mathfrak{g}_1} \phi_\lambda = K^{-1} r^{-1} \partial_r (K^{-1} r \partial_r \phi_\lambda),$$

and so

$$(5.9) \quad \partial_r (K^{-1} r \partial_r \phi_\lambda) = \lambda^2 K r \phi_\lambda.$$

As ϕ_λ is increasing, we get

$$K^{-1} r \partial_r \phi_\lambda \leq \int_R^r \lambda^2 \tau K \phi_\lambda(\tau) d\tau \leq \frac{1}{2} \lambda^2 (r^2 - R^2) \|K\|_{L^\infty} \phi_\lambda(r),$$

that is, $\partial_r \phi_\lambda \leq \|K\|_{L^\infty}^2 \lambda^2 (r - R) \phi_\lambda(r)$.

For the second order derivative of ϕ_λ , by (5.9), we have

$$|\partial_r^2 \phi_\lambda| = |\lambda^2 K^2 \phi_\lambda - \left(\frac{1}{r} - \frac{K'}{K}\right) \partial_r \phi_\lambda| \leq \lambda^2 K^2 \phi_\lambda + \frac{C}{r} \partial_r \phi_\lambda \lesssim \lambda^2 \phi_\lambda$$

for some $C > 0$ due to (1.3).

5.3. Test function $\Delta_{\mathbf{g}_1+\mathbf{g}_3}\Phi_\lambda = \lambda^2\Phi_\lambda$. Turning to the exponential perturbation of the spherically symmetric asymptotically Euclidean metric, we consider the following problem

$$\Delta_{\mathbf{g}_1+\mathbf{g}_3}\Phi_\lambda = \lambda^2\Phi_\lambda, x \in \Omega; \Phi_\lambda = 0, x \in \partial\Omega.$$

We would like to obtain similar bounds as that for $\mathbf{g}_1$.

Lemma 5.3. *There exists $\lambda_3 > 0$ such that for any $\lambda \in (0, \lambda_3)$, we can construct a solution Φ_λ such that*

$$(5.10) \quad \Phi_\lambda \sim \phi_\lambda \sim \begin{cases} \frac{\phi_0(r)}{\phi_0(\lambda^{-1})} & r \leq \lambda^{-1}, \\ h_\lambda \simeq \langle r\lambda \rangle^{-1/2} e^{\lambda \int_R^r K(s)ds} & r \geq \lambda^{-1}. \end{cases}$$

5.4. Proof of Lemma 5.3. As in [24, Lemma 3.1], see also Wakasa-Yordanov [38, Lemma 2.2], we introduce $\psi = \phi_\lambda - \Phi_\lambda$ where $\Delta_{\mathbf{g}_1}\phi_\lambda = \lambda^2\phi_\lambda$. Then we are reduced to prove existence of ψ and show smallness of $\|\psi\|_{L^\infty}$.

In fact, ψ satisfies

$$(5.11) \quad \Delta_{\mathbf{g}}\psi = \Delta_{\mathbf{g}}(\phi_\lambda - \Phi_\lambda) = \lambda^2\phi_\lambda - \lambda^2\Phi_\lambda - (\Delta_{\mathbf{g}_1} - \Delta_{\mathbf{g}})\phi_\lambda = \lambda^2\psi - W\phi_\lambda,$$

and $\psi|_{\partial B_R} = 0$. We claim that, for any $\lambda \in (0, \min(\alpha/2, 1/(2R), \lambda_2))$,

$$(5.12) \quad \|W\phi_\lambda\|_{L^q} \lesssim \lambda^2, \forall q \in [1, \infty].$$

Actually, we have

$$W\phi_\lambda = (\mathbf{g}_1^{jk} - \mathbf{g}^{jk})\partial_j\partial_k\phi_\lambda + [g_1^{-1/2}\partial_j(g_1^{1/2}\mathbf{g}_1^{jk}) - g^{-1/2}\partial_j(g^{1/2}\mathbf{g}^{jk})]\partial_k\phi_\lambda.$$

By (5.8) and (1.18) with $\lambda < \min(\alpha/2, 1/(2R), \lambda_2)$, it is easy to see that

$$|W\phi_\lambda| \lesssim \sum_{j,k} \sum_{1 \leq |\beta| \leq 2} |\nabla^{\leq 1} g_3^{jk}| |\nabla^\beta \phi_\lambda| \lesssim \lambda^2 \langle r \rangle e^{(\lambda-\alpha) \int_R^r K(\tau)d\tau} \lesssim \lambda^2 \langle r \rangle e^{-\frac{\alpha}{2}\delta_0 r},$$

which gives us (5.12).

Standard elliptic theory ensures that there exists a unique weak H_0^1 solution ψ to (5.11), which satisfies $\psi \in H^3 \cap C^\infty$. To show the smallness of $\|\psi\|_{L^\infty}$, we first take the (natural) inner product of (5.11) with ψ to get

$$-\langle \Delta_{\mathbf{g}}\psi, \psi \rangle_{\mathbf{g}} + \lambda^2 \langle \psi, \psi \rangle_{\mathbf{g}} = -\langle W\phi_\lambda, \psi \rangle_{\mathbf{g}}.$$

Thus by the Cauchy-Schwarz inequality and uniform elliptic condition (1.5) we have

$$\delta_0 \|\nabla \psi\|_{L_{\mathbf{g}}^2}^2 + \lambda^2 \|\psi\|_{L_{\mathbf{g}}^2}^2 \lesssim \lambda^2 \|\psi\|_{L_{\mathbf{g}}^2},$$

which yields

$$\|\psi\|_{L_x^2} \lesssim 1, \|\nabla \psi\|_{L_x^2} \lesssim \lambda.$$

In view of the equation, we obtain

$$\|\Delta_{\mathbf{g}}\psi\|_{L_x^2} \lesssim \lambda^2.$$

As $\psi \in H^2 \cap H_0^1$, we get from [6, Theorem 9.11-13] that

$$\|\nabla^2 \psi\|_{L_x^2} \lesssim \|\psi\|_{H^1} + \|\Delta_{\mathbf{g}}\psi\|_{L_x^2} \lesssim 1.$$

Applying the Gagliardo-Nirenberg inequality (for the exterior domain, see, e.g., [2]), we get

$$(5.13) \quad \|\psi\|_{L_x^4} \lesssim \|\psi\|_{L_x^2}^{1/2} \|\nabla \psi\|_{L_x^2}^{1/2} \lesssim \lambda^{1/2}, \|\nabla \psi\|_{L_x^4} \lesssim \|\nabla \psi\|_{L_x^2}^{1/2} \|\nabla^2 \psi\|_{L_x^2}^{1/2} \lesssim \lambda^{1/2}.$$

By Sobolev embedding, we obtain

$$(5.14) \quad \|\psi\|_{L_x^\infty} \lesssim \|\psi\|_{L_x^4} + \|\nabla \psi\|_{L_x^4} \lesssim \lambda^{1/2}.$$

Notice that ϕ_λ tends to zero as $r \rightarrow R$, the estimate (5.14) is not good enough to be controlled by ϕ_λ near the boundary. For that purpose, we observe that, in view of the equations, (5.12) and (5.13), we have

$$\|\Delta_{\mathfrak{g}} \psi\|_{L_x^4} \lesssim \lambda^2.$$

As $\psi \in W^{2,4} \cap W_0^{1,4}$, we get from [6, Theorem 9.11-13] that

$$\|\nabla^2 \psi\|_{L_x^4} \lesssim \|\psi\|_{W^{1,4}} + \|\Delta_{\mathfrak{g}} \psi\|_{L_x^4} \lesssim \lambda^{1/2}.$$

Another application of the Sobolev embedding gives us

$$(5.15) \quad \|\nabla \psi\|_{L_x^\infty} \lesssim \|\nabla \psi\|_{L_x^4} + \|\nabla \nabla \psi\|_{L_x^4} \lesssim \lambda^{1/2},$$

which, together with the boundary condition $\psi = 0$ for $r = R$, yields

$$(5.16) \quad \|\psi(r, \theta)\|_{L_\theta^\infty} \lesssim \lambda^{1/2}(r - R).$$

In summary, by (5.14), (5.16) and (5.2) we have obtained

$$(5.17) \quad |\psi(r, \theta)| \lesssim \lambda^{1/2} \min\{|x| - R, R\} \lesssim \lambda^{1/2} \phi_0(r) \ll \phi_\lambda,$$

which gives us the desired estimate (5.10) for $r \geq R$ and $\lambda < \lambda_3 \ll \min(1, \lambda_2, \alpha/2, 1/(2R))$.

6. GENERAL OBSTACLES

In this section, we present the proof of Lemma 1.6, which enables us to reduce the problems with general obstacles to the problem exterior to a disk.

6.1. Star-shaped obstacles. It turns out that we could construct an explicit diffeomorphism, when the obstacle $\mathcal{K}$ (the interior of the Jordan curve ∂D) is star-shaped. As $\mathcal{K}$ is star-shaped, there exists a smooth $R : \mathbb{S}^1 \rightarrow \mathbb{R}_+$:

$$\mathcal{K} = \{(r \cos \theta, r \sin \theta), r < R(\theta) \in (0, \infty)\}.$$

At first, we set $\delta_2 > 0$ such that

$$R(\theta) \in [2\delta_2, \delta_2^{-1}/2],$$

$R_3 = \delta_2$ and $R_4 = \delta_2^{-1}$. Let μ be a decreasing cut-off function such that $\mu = 0$ for $r > 2$ and $\mu = 1$ for $r < 1$. We set

$$(6.1) \quad f(r, \theta) = \mu\left(\frac{r}{R_4}\right) \frac{R_3}{R(\theta)} r + (1 - \mu\left(\frac{r}{R_4}\right)) r$$

and introduce

$$(6.2) \quad \mathcal{A} : (r, \theta) \rightarrow (f(r, \theta), \theta).$$

Notice that $f(R(\theta), \theta) = R_3$, $f(r, \theta) = r$ for any $r > 2R_4$, and

$$\partial_r f(r, \theta) = \mu\left(\frac{r}{R_4}\right) \frac{R_3}{R(\theta)} + (1 - \mu\left(\frac{r}{R_4}\right)) - r \mu'\left(\frac{r}{R_4}\right) \frac{R(\theta) - R_3}{R_4 R(\theta)} \in \left[\frac{R_3}{R(\theta)}, 1 + \frac{(2R_4)C}{R_4}\right]$$

and so

$$(6.3) \quad \partial_r f(r, \theta) \in [2\delta_2^2, 1 + 2C], \quad \forall r, \theta,$$

where we have denoted $C = \|\mu'\|_{L^\infty}$ and used the fact that $\mu' = 0$ unless $r \in [R_4, 2R_4]$.

It follows then that the map

$$\mathcal{A} : D \rightarrow \Omega := \mathbb{R}^2 \setminus \overline{B_{R_3}}$$

is a smooth diffeomorphism mapping preserving the boundary, which is an identity map for $r > 2R_4$.

6.2. General obstacles. Let γ be a Jordan curve in $\mathbb{R}^2$ and D be the unbounded component of the two components divided by γ . Without loss of generality, we assume $0 \in \mathbb{R}^2 \setminus \bar{D}$. Let

$$\hat{D} = \{z; z = x + iy, (x, y) \in D\} \subset \mathbb{C}, \check{D} = \{z; \frac{1}{z} \in \hat{D}\}.$$

Then by the Riemann mapping theorem (see, e.g., [37, Chapter 5, Theorem 4.1]), there exist $R > 0$ and (holomorphic) diffeomorphism $\mathcal{B}$ preserving the boundary such that

$$\mathcal{B} : \overline{\check{D} \cup \{0\}} \rightarrow \overline{B_{1/R}}, \mathcal{B}(0) = 0, \mathcal{B}'(0) = 1,$$

which gives us a diffeomorphism mapping

$$\mathcal{B} : \check{D} \rightarrow B_{1/R} \setminus \{0\}.$$

Let $\delta \in (0, 1)$ be a small parameter and $\mu(z)$ be a smooth cut-off function such that $\mu = 0$ when $|z| \leq 2$ and $\mu = 1$ when $|z| \geq 3$. Then we define the map

$$(6.4) \quad \mathcal{H}(z) = \mu_\delta(z)\mathcal{B}(z) + (1 - \mu_\delta(z))z, \quad \mu_\delta(z) = \mu\left(\frac{z}{\delta}\right).$$

We claim that there exists a small $\delta_1 > 0$ such that

$$(6.5) \quad \mathcal{H}(z) : \overline{\check{D}} \rightarrow \overline{B_{1/R} \setminus \{0\}}$$

is a diffeomorphism mapping when $\delta < \delta_1$. Thus it is easy to see that

$$\mathcal{A}(x, y) = \frac{1}{\mathcal{H}(\frac{1}{x+iy})} : D \rightarrow \Omega$$

is a desired diffeomorphism mapping. Hence we are reduced to showing the claim (6.5).

By definition, we have $\mathcal{H} = \mathcal{B}$ when $|z| \geq 3\delta$ and $\mathcal{H}(z) = z$ when $|z| \leq 2\delta$. So we need only to consider the case $2\delta < |z| < 3\delta$. Let $\mathcal{B} : (x, y) \rightarrow (\tilde{F}, \tilde{G})$, then we have

$$(6.6) \quad \mathcal{J}(\tilde{F}, \tilde{G})(0) = \begin{pmatrix} \tilde{F}_x(0) & \tilde{F}_y(0) \\ \tilde{G}_x(0) & \tilde{G}_y(0) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Let $\mathcal{H} : (x, y) \rightarrow (F, G)$, by definition (6.4) we have

$$F = x + \mu_\delta(\tilde{F} - x), \quad G = y + \mu_\delta(\tilde{G} - y)$$

and by (6.6), for $r = \sqrt{x^2 + y^2} < 4\delta$ small enough, we get that

$$F_x = 1 + \mathcal{O}(\delta), F_y = \mathcal{O}(\delta), G_x = \mathcal{O}(\delta), G_y = 1 + \mathcal{O}(\delta).$$

Then there exists $\delta_3 > 0$, such that for any $\delta \leq \delta_3$, we have $\mathcal{H}(B_{3\delta}) \subset B_{1/R}$ and

$$\mathcal{J}(F, G)(x, y) \in \left(\frac{1}{2}, \frac{3}{2}\right), \quad \forall r < 4\delta_3.$$

By inverse function theorem and compactness of $\bar{B}_{3\delta_3}$, there exists a uniform $\varepsilon_1 > 0$ such that $\mathcal{H}$ is a local diffeomorphism on $B_{\varepsilon_1}(r_0, \theta_0)$ for any $r_0 \in [0, 3\delta_3]$ and $\theta_0 \in [0, 2\pi]$. Thus we are reduced to proving bijection of $\mathcal{H}$.

We firstly prove the injection. For that purpose, we replace (x, y) by polar coordinates (r, θ) and (F, G) by (R, φ) . Recall that $x\partial_y - y\partial_x = \partial_\theta$ and $x\partial_x + y\partial_y = r\partial_r$, we get

$$(6.7) \quad \varphi_\theta = 1 + \mathcal{O}(\delta), R_\theta = \mathcal{O}(\delta^2), \varphi_r = \mathcal{O}(1), R_r = 1 + \mathcal{O}(\delta), r < 4\delta.$$

Let $\Gamma_r = \mathcal{H}(\partial B_r)$ ($r > 0$) be closed curves. With possibly shrinking $\delta_3 > 0$, we could assume, for some $C > 0$,

$$(6.8) \quad |\mathcal{O}(t)| \leq Ct, C\delta \leq \frac{1}{4}.$$

Then by (6.7) we have $\varphi_\theta \in [3/4, 5/4]$ and thus the winding number of Γ_r about the origin 0 must be 1, which shows that Γ_r is diffeomorphic to the unit circle $\mathbb{S}^1$. To complete the proof of injection, we need only to show $\{\Gamma_r\}_{0 < r \leq 3\delta}$ are disjoint, for which it suffices to prove

$$\Gamma_{r_1} \cap \Gamma_{r_2} = \emptyset, \quad 0 < |r_1 - r_2| < \frac{\delta\varepsilon}{2} < \frac{\varepsilon}{2}.$$

In fact, if $\Gamma_{r_1} \cap \Gamma_{r_2} \neq \emptyset$ for some $r_1 \neq r_2$ then $\varphi(r_1, \theta_1) = \varphi(r_2, \theta_2)$ for some θ_1, θ_2 . Without loss of generality, we assume $\theta_2 > \theta_1$. With $\theta_t = (1-t)\theta_0 + t\theta_1$, $r_t = (1-t)r_0 + tr_1$, by (6.7) and (6.8), we have

$$0 = \frac{d}{dt} \int_1^2 \varphi(r_t, \theta_t) dt = \int_1^2 \varphi_r \frac{d}{dt} r_t + \varphi_\theta \frac{d}{dt} \theta_t dt \geq (1 - C\delta)(\theta_2 - \theta_1) - C(r_2 - r_1),$$

which yields $\theta_2 - \theta_1 \leq \frac{\varepsilon\delta}{2} \frac{C}{1-C\delta} < \frac{\varepsilon}{2}$, that is

$$(r_2, \theta_2) \in B_\varepsilon(r_1, \theta_1).$$

This is a contradiction to the diffeomorphism of $\mathcal{H}$ on $B_\varepsilon(r_1, \theta_1)$.

Let Γ_r° denote the inner points of Γ_r , as $\Gamma_0 = \{0\}$ and $\Gamma_{3\delta} = \mathcal{B}(\partial B_{3\delta})$, we see that Γ_r is increasing about r ($\Gamma_{r_1}^\circ \subset \Gamma_{r_2}^\circ$, if $r_1 < r_2$). As Γ_r depends continuously on r , then it is clear that we have $\mathcal{H}(B_{3\delta}) = \Gamma_{3\delta}^\circ = \mathcal{B}(B_{3\delta})$. This completes the proof of (6.5).

ACKNOWLEDGMENTS

The first author is supported by NSF of Zhejiang Province(LY18A010008) and NSFC 11771194. The second and fourth author were supported in part by NSFC 11971428. The third author has been partially supported by Grant-in-Aid for Scientific Research (B) (No.18H01132) and Grant-in-Aid for Scientific Research (B) (No. 19H01795) and Young Scientists Research (No. 20K14351), Japan Society for the Promotion of Science.

REFERENCES

- [1] D. Catania, V. Georgiev, Blow-up for the semilinear wave equation in the Schwarzschild metric, *Differential Integral Equations*, 19 (2006) 799-830.
- [2] F. Crispo, P. Maremonti, An Interpolation Inequality in Exterior Domains, *REND. SEM. MAT. UNIV. PADOVA*, 112 (2004).
- [3] Y. Du, J. Metcalfe, C. D. Sogge, Y. Zhou, Concerning the Strauss conjecture and almost global existence for nonlinear Dirichlet-wave equations in 4-dimensions, *Comm. Partial Differential Equations* 33(7-9)(2008) 1487-1506.
- [4] Y. Du, Y. Zhou, The lifespan for nonlinear wave equation outside of star-shaped obstacle in three space dimensions, *Comm. Partial Differential Equations* 33 (7-9)(2008) 1455-1486.
- [5] V. Georgiev, H. Lindblad, C. D. Sogge, Weighted Strichartz estimates and global existence for semilinear wave equations, *Amer. J. Math.* 119(6) (1997) 1291-1319.

- [6] D. Gilbarg, N. S. Trudinger, Elliptic partial differential equations of second order, Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [7] R.T. Glassey, Finite-time blow-up for solutions of nonlinear wave equations, *Math. Z.* 177 (1981) 323-340.
- [8] R.T. Glassey, Existence in the large for $\square u = F(u)$ in two space dimensions, *Math. Z.* 178 (1981) 233-261.
- [9] K. Hidano, J. Metcalfe, H. F. Smith, C. D. Sogge, Y. Zhou, On abstract Strichartz estimates and the Strauss conjecture for nontrapping obstacles, *Trans. Amer. Math. Soc.* 362(5) (2010) 2789-2809.
- [10] M. Ikeda, M. Sobajima, K. Wakasa, Blow-up phenomena of semilinear wave equations and their weakly coupled systems, *J. Differential Equations*, 267(9) (2019) 5165-5201.
- [11] T. Inai, M. Kato, H. Takamura, K. Wakasa, The sharp lower bound of the lifespan of solutions to semilinear wave equations with low powers in two space dimensions. In *Asymptotic analysis for nonlinear dispersive and wave equations. Proceedings of the international conference on asymptotic analysis for nonlinear dispersive and wave equations, Osaka University, Osaka, Japan, September 6-9, 2014*, pages 31-53. Tokyo: Mathematical Society of Japan, 2019.
- [12] F. John, Blow-up of solutions of nonlinear wave equations in three space dimensions, *Manuscripta Math.* 28 (1-3)(1979) 235-268.
- [13] N. A. Lai, Z. H. Tu, Strauss exponent for semilinear wave equations with scattering space dependent damping, *J. Math. Anal. Appl.* 489 (2020) 124189.
- [14] N. A. Lai, Y. Zhou, An elementary proof of Strauss conjecture, *J. Funct. Anal.* 267 (5)(2014) 1364-1381.
- [15] N. A. Lai, Y. Zhou, Finite time blow up to critical semilinear wave equation outside the ball in 3-D, *Nonlinear Anal.* 125 (2015) 550-560.
- [16] N. A. Lai, Y. Zhou, Nonexistence of global solutions to critical semilinear wave equations in exterior domain in high dimensions, *Nonlinear Anal.* 143 (2016), 89-104.
- [17] N. A. Lai, Y. Zhou, Blow up for initial boundary value problem of critical semilinear wave equation in 2-D, *Commun. Pure Appl. Anal.* 17 (4)(2018) 1499-1510.
- [18] N. A. Lai, Y. Zhou, Global existence of solutions of the critical semilinear wave equations with variable coefficients outside obstacles, *Sci. China Math.* 54(2) (2011) 205-220.
- [19] X. Li, G. Wang, Blow up of solutions to nonlinear wave equations in 2D exterior domains, *Arch. Math.* 98 (2012) 265-275.
- [20] Y. H. Lin, N. A. Lai, S. Ming, Lifespan estimate for semilinear wave equation in Schwarzschild spacetime, *Appl. Math. Lett.* 99 (2020) 105997, 4 pp.
- [21] H. Lindblad, Blow-up for solutions of $\square u = |u|^p$ with small initial data, *Comm. Partial Differential Equations* 15 (6) (1990) 757-821.
- [22] H. Lindblad, J. Metcalfe, C. D. Sogge, M. Tohaneanu, C. B. Wang, The Strauss conjecture on Kerr black hole backgrounds, *Math. Ann.*, 359(3-4) (2014) 637-661.
- [23] H. Lindblad, C. D. Sogge, Long-time existence for small amplitude semilinear wave equations, *Amer. J. Math.* 118 (5) (1996) 1047-1135.
- [24] M. Y. Liu, C. B. Wang, The blow up of solutions to semilinear wave equations on asymptotically Euclidean manifolds, Preprint. [arXiv:1912.02540](https://arxiv.org/abs/1912.02540), 2019.
- [25] J. Marzuola, J. Metcalfe, D. Tataru, M. Tohaneanu, Strichartz estimates on Schwarzschild black hole backgrounds, *Comm. Math. Phys.* (293) 2010 37-83.
- [26] J. Metcalfe, C. D. Sogge, Global existence for high dimensional quasilinear wave equations exterior to star-shaped obstacles, *Discrete Contin. Dyn. Syst.* 28(4) (2010) 1589-1601.
- [27] J. Metcalfe, C. B. Wang, The Strauss conjecture on asymptotically flat space-times, *SIAM J Math. Anal.* 49 (2017) 4579-4594.
- [28] J. Schaeffer, The equation $\square u = |u|^p$ for the critical value of p , *Proc. Roy. Soc. Edinburgh Sect. A* 101 (1-2)(1985) 31-44.
- [29] T.C. Sideris, Nonexistence of global solutions to semilinear wave equations in high dimensions, *J. Differential Equations* 52 (1984) 378-406.
- [30] H. F. Smith, C. D. Sogge, On the critical semilinear wave equation outside convex obstacles, *J. Amer. Math. Soc.* 8 (4)(1995) 879-916.
- [31] H. F. Smith, C. D. Sogge, C. B. Wang, Strichartz estimates for Dirichlet-wave equations in two dimensions with applications, *Trans. Amer. Math. Soc.* 364 (2012) 3329-3347.
- [32] M. Sobajima, K. Wakasa, Finite time blowup of solutions to semilinear wave equation in an exterior domain, *J. Math. Anal. Appl.* 484 (2020), 123667.

- [33] C. D. Sogge, C. B. Wang, Concerning the wave equation on asymptotically Euclidean manifolds, *J Anal Math* 112 (2010) 1-32.
- [34] W. A. Strauss, Nonlinear scattering theory at low energy, *J. Funct. Anal.* 41 (1981) 110-133.
- [35] H. Takamura, Improved Kato's lemma on ordinary differential inequality and its application to semilinear wave equations, *Nonlinear Analysis* 125 (2015) 227-240.
- [36] H. Takamura, K. Wakasa, The sharp upper bound of the lifespan of solutions to critical semilinear wave equations in high dimensions, *J. Differential Equations* 251 (2011) 1157-1171.
- [37] M. E. Taylor, *Partial differential equations I-III. Second edition. Applied Mathematical Sciences*, 115-117. Springer, New York, 2011.
- [38] K. Wakasa, B. Yordanov, Blow-up of solutions to critical semilinear wave equations with variable coefficients, *J. Differential Equations* 266(9) (2019) 5360-5376.
- [39] C. B. Wang, X. Yu, Concerning the Strauss conjecture on asymptotically Euclidean manifolds, *J. Math. Anal. Appl.* 379 (2011) 549-566.
- [40] C. B. Wang, Recent progress on the Strauss conjecture and related problems (in Chinese), *Sci. Sin. Math.* 48 (2018) 111-130.
- [41] C. B. Wang, Long time existence for semilinear wave equations on asymptotically flat space-times, *Comm. Partial Differential Equations* 42 (7) (2017) 1150-1174.
- [42] B. Yordanov, Q. S. Zhang, Finite time blow up for critical wave equations in high dimensions, *J. Funct. Anal.* 231(2) (2006) 361-374.
- [43] X. Yu, Generalized Strichartz estimates on perturbed wave equation and applications on Strauss conjecture, *Differential and Integral Equations* 24(5-6) (2011) 443-468.
- [44] D. B. Zha, Y. Zhou, Lifespan of classical solutions to quasilinear wave equations outside of a star-shaped obstacle in four space dimensions, *J. Math. Pures Appl.* 103 (2015) 788-808.
- [45] Y. Zhou, Blow up of solutions to semilinear wave equations with critical exponent in high dimensions, *Chinese Ann. Math. Ser. B* 28(2)(2007) 205-212.
- [46] Y. Zhou, Lifespan of classical solutions to $\square u = |u|^p$ in two space dimensions, *Chin. Ann. Math. Ser. B* 14 (1993) 225-236.
- [47] Y. Zhou, Blow up of classical solutions to $\square u = |u|^{1+\alpha}$ in three space dimensions, *J. Partial Differential Equations* 5 (1992) 21-32.
- [48] Y. Zhou, W. Han, Blow-up of solutions to semilinear wave equations with variable coefficients and boundary, *J. Math. Anal. Appl.* 374 (2011) 585-601.

INSTITUTE OF NONLINEAR ANALYSIS AND DEPARTMENT OF MATHEMATICS, LISHUI UNIVERSITY,
LISHUI 323000, P. R. CHINA

Email address: ninganlai@lsu.edu.cn

DEPARTMENT OF MATHEMATICS, ZHEJIANG SCI-TECH UNIVERSITY, HANGZHOU 310018, P. R. CHINA

Email address: mengyunliu@zstu.edu.cn

DEPARTMENT OF CREATIVE ENGINEERING, NATIONAL INSTITUTE OF TECHNOLOGY, KUSHIRO COLLEGE, 2-32-1 OTANOSHIKE-NISHI, KUSHIRO-SHI, HOKKAIDO 084-0916, JAPAN

Email address: wakasa@kushiro-ct.ac.jp

SCHOOL OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU 310027, P. R. CHINA

Email address: wangcbo@zju.edu.cn