

Relationship of Two Discrete Dynamical Models: One-dimensional Cellular Automata and Integral Value Transformations

Sreeya Ghosh^{a,b}, Sudhakar Sahoo^c, Sk. Sarif Hassan^{d,*}, Jayanta Kumar Das^{e,b}, Pabitra Pal Choudhury^a

^a*Department of Applied Mathematics, University of Calcutta, Kolkata-700009, India*

^b*Applied Statistics Unit, Indian Statistical Institute, Kolkata-108, India*

^c*Institute of Mathematics and Applications, Bhubaneswar-751029, India*

^d*Dept. of Mathematics, Pingla Thana Mahavidyalaya, Paschim Medinipur-721140, India*

^e*School of Medicine, Johns Hopkins University, MD-21210, USA*

Abstract

Cellular Automaton(CA) and an Integral Value Transformation(IVT) are two well established mathematical models which evolve in discrete time steps. Theoretically, studies on CA suggest that CA is capable of producing a great variety of evolution patterns. However computation of non-linear CA or higher dimensional CA maybe complex, whereas IVTs can be manipulated easily.

The main purpose of this paper is to study the link between a transition function of a one-dimensional CA and IVTs. Mathematically, we have also established the algebraic structures of a set of transition functions of a one-dimensional CA as well as that of a set of IVTs using binary operations. Also DNA sequence evolution has been modeled using IVTs.

Keywords: Cellular Automaton, Integral Value Transformations

1. Introduction

Cellular Automaton(pl. cellular automata, abbrev. CA) is a discrete model which has applications in computer science, mathematics, physics, complexity science, theoretical biology and microstructure modeling. This model was introduced by J.von Neumann and S.Ulam in 1940 for designing self replicating systems [1, 2, 3].

A CA consists of a finite/countably infinite number of finite-state semi-automata [4, 5] known as ‘cells’ arranged in an ordered n -dimensional grid. Each cell receives input from the neighbouring cells and changes according to the transition function. The transitions at each of the cells together induces a change of the grid pattern. The simplest CA is a CA where the grid is a one-dimensional line. Stephen Wolfram’s work in the 1980s contributed to a systematic study of one-dimensional CA, providing the first qualitative classification of their behaviour [6]. CA has been studied for solving many interesting problems on utilizing mathematical bases such as polynomial, matrix algebra, Boolean derivative etc.[7, 8]. Further, dynamic behaviour of CA can also be studied using various mathematical tools [9, 10, 11, 12], that help to understand the crucial properties and modeling of various classes of discrete dynamical system [12].

An Integral Value Transformation (abbrev. IVT), a class of discrete dynamical system, were first introduced during 2009-10 ([13]). A IVT of k -dimension is a function defined using p -adic numbers over \mathbb{N}_0^k where $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $p \in \mathbb{N}$.

*Corresponding author

Email addresses: sreeya135@gmail.com (Sreeya Ghosh), sudhakar.sahoo@gmail.com (Sudhakar Sahoo), sarimif@gmail.com (Sk. Sarif Hassan), dasjayantakumar89@gmail.com (Jayanta Kumar Das), pabitra@isical.ac.in (Pabitra Pal Choudhury)

The IVTs have been studied in different viewpoints including the understanding of the evolution of integer sequences and behavioral patterns of integers in discrete time points [14, 15].

In this paper, Wolfram code of an elementary CA and global transition function of a one-dimensional CA has been represented through IVTs. The algebraic structure of a set of transition functions of a one-dimensional CA as well as the algebraic structure of a set of IVTs under some binary operations have been studied. In the last section, some specific applications have been discussed in which we have seen certain novelty lies in IVTs over one-dimensional CA.

2. Mathematical preliminaries

Definition 1. Let Q be a finite set of memory elements also called the **state set**.

A **global configuration** is a mapping from the group of integers \mathbb{Z} to the set Q given by $C : \mathbb{Z} \rightarrow Q$. The set $Q^{\mathbb{Z}}$ is the set of all global configurations where $Q^{\mathbb{Z}} = \{C | C : \mathbb{Z} \rightarrow Q\}$.

Definition 2. A mapping $\tau : Q^{\mathbb{Z}} \rightarrow Q^{\mathbb{Z}}$ is called a **global transition function**.

A **CA**(denoted by \mathcal{C}_{τ}^Q)(reported in [16, 17]) is a triplet $(Q, Q^{\mathbb{Z}}, \tau)$, where,

- Q is the finite state set
- $Q^{\mathbb{Z}}$ is the set of all configurations
- τ is the global transition function

Definition 3. The set $\mathcal{Q}^{\mathbb{Z}} = \{\tau | \tau : Q^{\mathbb{Z}} \rightarrow Q^{\mathbb{Z}}\}$ is the set of all possible global transition functions of CA having state set Q .

A mapping τ is invertible if $\forall C_i, C_j \in Q^{\mathbb{Z}}, \exists \tau^{-1}$ such that

$$\tau(C_i) = C_j \Leftrightarrow \tau^{-1}(C_j) = C_i$$

Definition 4. For $i \in \mathbb{Z}, r \in \mathbb{N}$, let $S_i = \{i - r, \dots, i - 1, i, i + 1, \dots, i + r\} \subseteq \mathbb{Z}$. S_i is the neighbourhood of the i^{th} cell, r is the radius of the neighbourhood of a cell. It follows that $\mathbb{Z} = \bigcup_i S_i$.

A restriction from \mathbb{Z} to S_i induces a restriction of C to $\overline{c_i}$ given by $\overline{c_i} : S_i \rightarrow Q$; where $\overline{c_i}$ may be called **local configuration** of the i^{th} cell.

The mapping $\mu_i : Q^{S_i} \rightarrow Q$ is known as a **local transition function** for the i^{th} cell having radius r . Thus $\forall i \in \mathbb{Z}, \mu_i(\overline{c_i}) \in Q$ and it follows that,

$$\tau(C) = \tau(\dots, c_{i-1}, c_i, c_{i+1}, \dots) = \dots \mu_{i-1}(\overline{c_{i-1}}) \cdot \mu_i(\overline{c_i}) \cdot \mu_{i+1}(\overline{c_{i+1}}) \dots$$

Definition 5. The set $M = \{\mu_i | \mu_i : Q^{S_i} \rightarrow Q, i \in \mathbb{Z}\}$ is the set of all possible local transition functions of CA having state set Q .

Definition 6. If for a particular CA, $|Q| = 2$ so that we can write $Q = \{0, 1\}$, then the CA is said to be a **binary CA** or a **Boolean CA**. A Boolean CA having radius 1 is known as an **Elementary CA(ECA)**.

Definition 7. **Wolfram code** is a naming system often used for a one-dimensional CA introduced by Stephen Wolfram (see[6]).

For a one-dimensional CA with Q states, the local rule μ_i for some i^{th} cell, $i \in \mathbb{Z}$ of radius r (neighbourhood $2r + 1$) can be specified by an Q^{2r+1} -bit sequence. The decimal equivalent form of this sequence is known as the Wolfram code.

Thus, the Wolfram code for a particular rule is a number in the range from 0 to $Q^{Q^{2r+1}} - 1$, converted from Q -ary to decimal notation.

Example 2.1. Let the local rule of an ECA for some i^{th} cell, $i \in \mathbb{Z}$ be,

$$\mu_i(\overline{c_i}) = \mu_i(c_{i-1}, c_i, c_{i+1}) = (c_{i-1} \vee c_{i+1}) \wedge c_i$$

where ' \vee ' stands for OR operation, ' \wedge ' stands for AND operation, c_j is the j^{th} cell configuration for $j = i - 1, i, i + 1$. Then we get the 2^3 -bit sequence as,

$$\mu(111)\mu(110)\mu(101)\mu(100)\mu(011)\mu(010)\mu(001)\mu(000) = 1\ 1\ 0\ 0\ 1\ 0\ 0\ 0$$

The decimal equivalent number for 11001000 is 200 and so the Wolfram code is **RULE 200**.

Definition 8. A p -adic k -dimensional **Integral Value Transformation (IVT)** denoted by $IVT_j^{p,k}$ for $p \in \mathbb{N}$, $k \in \mathbb{N}$ is a function of p -base numbers from \mathbb{N}_0^k to \mathbb{N}_0 defined (in [18]) as

$$IVT_j^{p,k}(n_1, n_2, \dots, n_k) = (f_j(a_0^{n_1}, \dots, a_0^{n_k})f_j(a_1^{n_1}, \dots, a_1^{n_k}) \dots f_j(a_{l-1}^{n_1}, \dots, a_{l-1}^{n_k}))_p = m$$

where $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $p \in \mathbb{N}$, $n_s = (a_0^{n_s} a_1^{n_s} \dots a_{l-1}^{n_s})_p$ for $s = 1, 2, \dots, k$.

f_j is a function from $\{0, 1, \dots, p-1\}^k$ to $\{0, 1, \dots, p-1\}$ for $j = 0, 1, \dots, p^k - 1$,

m is the decimal conversion of the p -base number.

Remark 2.1. In particular (discussed in [19, 20, 21, 22]), if $\forall i = 0, 1, \dots, l-1$, the function f_j be defined as

1. $f_j(a_i^{n_1}, \dots, a_i^{n_k}) = \lfloor (a_i^{n_1} + \dots + a_i^{n_k})/p \rfloor$, then $IVT_j^{p,k}$ is known as a **Modified Carry Value Transformation (MCVT)**.

Again, $MCVT$ with a 0 padding at the right end is known as a **Carry Value Transformation (CVT)**.

2. $f_j(a_i^{n_1}, \dots, a_i^{n_k}) = (a_i^{n_1} + \dots + a_i^{n_k}) \bmod p$, then $IVT_j^{p,k}$ is known as an **Exclusive OR Transformation (XORT)**.
3. $f_j(a_i^{n_1}, \dots, a_i^{n_k}) = \max(a_i^{n_1}, \dots, a_i^{n_k})$, then $IVT_j^{p,k}$ is known as an **Extreme Value Transformation (EVT)**.

3. Wolfram code of an ECA and IVT

For an ECA, Wolfram code for a local rule μ is the decimal number $j \in \{0, 1, 2, \dots, 255\}$ obtained from the 8-bit sequence

$$\mu(111)\mu(110)\mu(101)\mu(100)\mu(011)\mu(010)\mu(001)\mu(000) = j$$

Therefore, Wolfram code for each local transition function of a 3-neighbourhood Boolean CA can be represented by a 2 base 3-dimensional IVT.

The function μ in the above 8-bit sequence can be represented by a function $f_j : \{0, 1\}^3 \rightarrow \{0, 1\}$ which gives

$$(f_j(111)f_j(110)f_j(101)f_j(100)f_j(011)f_j(010)f_j(001)f_j(000))_2$$

$$= IVT_j^{2,3}(11110000_2, 11001100_2, 10101010_2) = IVT_j^{2,3}(240_{10}, 204_{10}, 170_{10})$$

Hence it follows that for a 3-neighbourhood Boolean CA, any Wolfram code $j \in \{0, 1, 2, \dots, 255\}$ can be equivalently represented by $IVT_j^{2,3}(240_{10}, 204_{10}, 170_{10})$.

Example 3.1. Wolfram code 200 can be equivalently represented as

$$200_{10} = (11001000)_2 = IVT_{200}^{2,3}(240_{10}, 204_{10}, 170_{10})$$

However the following example shows that for $j \in \{0, 1, 2, \dots, 255\}$, any $IVT_j^{2,3}$ may not correspond to a Wolfram code.

Example 3.2.

$$\begin{aligned} IVT_j^{2,3}(240_{10}, 204_{10}, 171_{10}) &= IVT_j^{2,3}(11110000_2, 11001100_2, 10101011_2) \\ &= (f_j(111)f_j(110)f_j(101)f_j(100)f_j(011)f_j(010)f_j(001)f_j(000))_2 \end{aligned}$$

In this 8-bit sequence, $f_j(000)$ is missing and so this cannot correspond to any Wolfram code.

3.1. Some Particular Wolfram codes and Particular IVTs

We know that transformations such as MCVT, XOR, EVT are particular cases of IVTs. Again, any Wolfram code can be represented by an IVT. Therefore some particular Wolfram codes which can be represented by an MCVT, XOR or EVT are as follows.

1. Let the local rule of an ECA for some i^{th} cell, $i \in \mathbb{Z}$ be,

$$\mu_i(\overline{c_i}) = \mu_i(c_{i-1}, c_i, c_{i+1}) = (c_{i-1} \wedge c_i) \vee ((c_{i-1} \vee c_i) \wedge c_{i+1})$$

Then we get the 2³-bit sequence as 11101000 and Wolfram code 232.

Here, μ_i can be represented by function $f_j : \{0, 1\}^3 \rightarrow \{0, 1\}$ given by

$$f_j(c_{i-1}, c_i, c_{i+1}) = (c_{i-1} \wedge c_i) \vee ((c_{i-1} \vee c_i) \wedge c_{i+1}) = \lfloor (c_{i-1} + c_i + c_{i+1})/2 \rfloor$$

Thus, Wolfram code 232 can be represented as

$$\begin{aligned} &(f_{232}(111)f_{232}(110)f_{232}(101)f_{232}(100)f_{232}(011)f_{232}(010)f_{232}(001)f_{232}(000))_2 \\ &= (\lfloor 3/2 \rfloor \lfloor 2/2 \rfloor \lfloor 2/2 \rfloor \lfloor 1/2 \rfloor \lfloor 2/2 \rfloor \lfloor 1/2 \rfloor \lfloor 1/2 \rfloor \lfloor 0/2 \rfloor)_2 \\ &= MCVT_{232}^{2,3}(11110000_2, 11001100_2, 10101010_2) = MCVT_{232}^{2,3}(240_{10}, 204_{10}, 170_{10}) \end{aligned}$$

2. Let the local rule of an ECA for some i^{th} cell, $i \in \mathbb{Z}$ be,

$$\mu_i(\overline{c_i}) = \mu_i(c_{i-1}, c_i, c_{i+1}) = (c_{i-1} \vee c_i \vee c_{i+1})$$

Then we get the 2³-bit sequence as 10010110 and Wolfram code 150.

Here, μ_i can be represented by function $f_j : \{0, 1\}^3 \rightarrow \{0, 1\}$ given by

$$f_j(c_{i-1}, c_i, c_{i+1}) = (c_{i-1} \vee c_i \vee c_{i+1}) = (c_{i-1} + c_i + c_{i+1}) \bmod 2$$

Thus, Wolfram code 150 can be represented as

$$\begin{aligned}
& (f_{150}(111)f_{150}(110)f_{150}(101)f_{150}(100)f_{150}(011)f_{150}(010)f_{150}(001)f_{150}(000))_2 \\
& = ((3 \bmod 2)(2 \bmod 2)(2 \bmod 2)(1 \bmod 2)(2 \bmod 2)(1 \bmod 2)(1 \bmod 2)(0 \bmod 2))_2 \\
& = XOR_{150}^{2,3}(11110000_2, 11001100_2, 10101010_2) = XOR_{150}^{2,3}(240_{10}, 204_{10}, 170_{10})
\end{aligned}$$

3. Let the local rule of an ECA for some i^{th} cell, $i \in \mathbb{Z}$ be,

$$\mu_i(\overline{c_i}) = \mu_i(c_{i-1}, c_i, c_{i+1}) = (c_{i-1} \vee c_i \vee c_{i+1})$$

Then we get the 2^3 -bit sequence as 11111110 and Wolfram code 254.

Here, μ_i can be represented by function $f_j : \{0, 1\}^3 \rightarrow \{0, 1\}$ given by

$$f_j(c_{i-1}, c_i, c_{i+1}) = (c_{i-1} \vee c_i \vee c_{i+1}) = \max\{c_{i-1}, c_i, c_{i+1}\}$$

Thus, Wolfram code 254 can be represented as

$$\begin{aligned}
& (f_{254}(111)f_{254}(110)f_{254}(101)f_{254}(100)f_{254}(011)f_{254}(010)f_{254}(001)f_{254}(000))_2 \\
& = (11111110)_2 \\
& = EVT_{254}^{2,3}(11110000_2, 11001100_2, 10101010_2) = EVT_{254}^{2,3}(240_{10}, 204_{10}, 170_{10})
\end{aligned}$$

4. Transition Function of a CA and IVT

Definition 9. The set $T^{p,k} = \{IVT_j^{p,k} | IVT_j^{p,k} : \mathbb{N}_0^k \rightarrow \mathbb{N}_0\}$ is the set of all p -base k -dimensional IVTs for $p \in \mathbb{N}, k \in \mathbb{N}$.

Definition 10. The v -fold cartesian product $\underbrace{T^{p,k} \times \dots \times T^{p,k}}_{v \text{ times}}$ for $v \in \mathbb{N}$, denoted by \mathcal{T}^v is given by

$$\mathcal{T}^v = \{(IVT_{j_1}^{p,k}, IVT_{j_2}^{p,k}, \dots, IVT_{j_v}^{p,k})\}$$

where $IVT_{j_s}^{p,k} : \mathbb{N}_0^k \rightarrow \mathbb{N}_0 \in T^{p,k}$, for $s = 1, 2, \dots, v(\in \mathbb{N})$

Definition 11. Let a restriction on $IVT_j^{p,k}$ from \mathbb{N}_0^k to $Q^k = \{0, 1, \dots, p-1\}^k$ for $p \in \mathbb{N}$ be denoted by $\underline{IVT}_j^{p,k}$. The set $T_{|Q}^{p,k} = \{\underline{IVT}_j^{p,k} | \underline{IVT}_j^{p,k} : Q^k \rightarrow Q\}$ is the set of all p -base k -dimensional IVTs when \mathbb{N}_0^k is restricted to the subset Q^k . Therefore for $(\eta_1, \eta_2, \dots, \eta_k) \in Q^k$ we get,

$$\underline{IVT}_j^{p,k}(\eta_1, \eta_2, \dots, \eta_k) = f_j(\eta_1, \eta_2, \dots, \eta_k)_p = m \in Q$$

Definition 12. The set $\mathcal{T}_{|Q}^v = \{(\underline{IVT}_{j_1}^{p,k}, \underline{IVT}_{j_2}^{p,k}, \dots, \underline{IVT}_{j_v}^{p,k})\}$ is the v -fold cartesian product $\underbrace{T_{|Q}^{p,k} \times \dots \times T_{|Q}^{p,k}}_{v \text{ times}}$ when \mathbb{N}_0^k is restricted to subset Q^k where $\underline{IVT}_{j_s}^{p,k} : Q^k \rightarrow Q \in T_{|Q}^{p,k}$ for $s = 1, 2, \dots, v(\in \mathbb{N})$

Now, a local transition function of any i^{th} cell of a one-dimensional CA having $p(\in \mathbb{N})$ states and radius $r(\in \mathbb{N})$ will be of the form

$$\mu_i(\overline{c_i}) = \mu_i(c_{i-r}, \dots, c_i, \dots, c_{i+r}) = c_i^*$$

where $c_{i-r}, \dots, c_i, \dots, c_{i+r}, c_i^* \in \{0, 1, \dots, p-1\}$.

This can be represented by some $\underline{IVT}_{j_i}^{p,k}$, where $p \in \mathbb{N}, k = 2r + 1$ and $j_i \in \{0, 1, 2, \dots, (p^{p^k} - 1)\}$ is the underlying Wolfram code (which is in turn equal to $IVT_{j_i}^{p,k}(240_{10}, 204_{10}, 170_{10})$).

Thus $\forall i \in \mathbb{Z}$,

$$\mu_i(\overline{c_i}) \equiv \underline{IVT}_{j_i}^{p,k}(\overline{c_i})$$

Now if τ be the global transition function of any $v(\in \mathbb{N} \geq k)$ -celled CA, then for a global configuration $C = (c_1, c_2, \dots, c_v)$, we get

$$\tau(C) = \tau(c_1, c_2, \dots, c_v) = \mu_1(\overline{c_1}) \cdot \mu_2(\overline{c_2}) \dots \mu_v(\overline{c_v})$$

Therefore it follows that,

$$\tau(c_1, c_2, \dots, c_v) \equiv (\underline{IVT}_{j_1}^{p,k}(\overline{c_1}), \underline{IVT}_{j_2}^{p,k}(\overline{c_2}), \dots, \underline{IVT}_{j_v}^{p,k}(\overline{c_v}))$$

Hence for any v -celled ECA it follows that,

$$\tau(C) \equiv (\underline{IVT}_{j_1}^{2,3}(\overline{c_1}), \underline{IVT}_{j_2}^{2,3}(\overline{c_2}), \dots, \underline{IVT}_{j_v}^{2,3}(\overline{c_v}))$$

where $j_1, j_2, \dots, j_v \in \{0, 1, \dots, 255\}$.

Now if τ be the global transition function of a countably infinite celled CA with cells having k neighbourhood, then for any global configuration

$C = (\dots c_{i-1}, c_i, c_{i+1}, \dots)$, we get

$$\tau(C) = \dots \mu_{i-1}(\overline{c_{i-1}}) \cdot \mu_i(\overline{c_i}) \cdot \mu_{i+1}(\overline{c_{i+1}}) \dots$$

Therefore it follows that,

$$\tau(C) \equiv (\dots, \underline{IVT}_{j_{i-1}}^{p,k}(\overline{c_{i-1}}), \underline{IVT}_{j_i}^{p,k}(\overline{c_i}), \underline{IVT}_{j_{i+1}}^{p,k}(\overline{c_{i+1}}), \dots)$$

where $\dots j_{i-1} j_i, j_{i+1}, \dots \in \{0, 1, 2, \dots, (p^{p^k} - 1)\}$.

Example 4.1. Let the initial configuration of a CA having state set $\{0, 1, 2\}$ be $C_0 = (01201)_3$ and the transition function be given by

$$\tau(01201)_3 = \mu_1(101)\mu_2(012)\mu_3(120)\mu_4(201)\mu_5(010) = (02001)_3$$

where μ_1, μ_3, μ_5 follow Wolfram code 377 and μ_2, μ_4 follow Wolfram code 588 for 3-state CA.

A local transition function can be equivalently represented by some $\underline{IVT}_j^{3,3}$ and thus it follows that $\tau(01201)_3$ is equivalent to

$$\underline{IVT}_{377}^{3,3}(1, 0, 1) \underline{IVT}_{588}^{3,3}(0, 1, 2) \underline{IVT}_{377}^{3,3}(1, 2, 0) \underline{IVT}_{588}^{3,3}(2, 0, 1) \underline{IVT}_{377}^{3,3}(0, 1, 0)$$

Conversely, for any $\underline{IVT}_j^{p,k}$ if k be odd, i.e. if $k = 2r + 1$ for some $r < k(\in \mathbb{N}_0)$, then $\underline{IVT}_j^{p,k}(\eta_1, \eta_2, \dots, \eta_{2r+1})$ will be equivalent to a local transition function of the $(r + 1)^{th}$ cell in a CA with $p(\in \mathbb{N})$ states and $2r + 1$ neighbourhood whose underlying Wolfram code is $j \in \{0, 1, \dots, p^{p^k} - 1\}$, given by

$$\underline{IVT}_j^{q,l}(\eta_1, \eta_2, \dots, \eta_{2r+1}) = \underline{IVT}_j^{p,k}(\overline{\eta}_{r+1}) \equiv \mu_{r+1}(\overline{\eta}_{r+1})$$

Remark 4.1. For a $p(\in \mathbb{N})$ -state $k(= 2r + 1 \in \mathbb{N})$ -neighbourhood CA, clearly $\forall \overline{c_i} \in Q^{S_i} \subseteq Q^{\mathbb{Z}}$ $\mu_j(\overline{c_i}) \equiv \underline{IVT}_j^{p,k}(\overline{c_i})$ where $j \in \{0, 1, \dots, p^{p^k} - 1\}$ is the underlying Wolfram Code. It follows that the set of local transition functions M is equivalent to the set $T_{|Q}^{p,k}$ and vice-versa.

Hence a function $\phi : M \rightarrow T_{|Q}^{p,k}$ defined by $\phi(\mu_j)(\overline{c_i}) = \underline{IVT}_{j_i}^{p,k}(\overline{c_i})$ is an isomorphism.

Moreover, for any $v(\in \mathbb{N} \geq k)$ -celled CA, having a global configuration $C = (c_1, c_2, \dots, c_v)$, if $\mathcal{Q}^v = \{\tau | \tau : Q^v \rightarrow Q^v\}$, then $\forall, \overline{c_i} \in Q^k \subseteq Q^v, i = 1, 2, \dots, v$, it follows that a function $\phi : \mathcal{Q}^v \rightarrow \mathcal{T}_{|Q}^v$ defined by

$$\phi(\tau)(c_1, c_2, \dots, c_v) \equiv (\underline{IVT}_{j_1}^{p,k}(\overline{c_1}), \underline{IVT}_{j_2}^{p,k}(\overline{c_2}), \dots, \underline{IVT}_{j_v}^{p,k}(\overline{c_v}))$$

is an isomorphism.

5. Some Algebraic Results on CA and IVT

Theorem 5.1. $(\mathcal{Q}^{\mathbb{Z}}, \circ)$ forms a monoid w.r.t. composition of global transition functions.

Proof. Clearly $\mathcal{Q}^{\mathbb{Z}}$ is closed and associative under composition of global transition functions.

The transition function μ_e such that $\forall c_i \in Q, \mu_e(\overline{c_i}) = c_i$ is the local identity and it follows that $\tau_e \in \mathcal{Q}^{\mathbb{Z}}$ is the global identity such that $\forall C \in Q^{\mathbb{Z}}, i \in \mathbb{Z}$,

$$\tau_e(C) = \dots \mu_e(\overline{c_{i-1}}) \mu_e(\overline{c_i}) \mu_e(\overline{c_{i+1}}) \dots = C$$

Hence the theorem. □

Corollary 5.1. $(\overline{\mathcal{Q}^{\mathbb{Z}}}, \circ)$ forms a group w.r.t. composition of global transition functions w $\overline{\mathcal{Q}^{\mathbb{Z}}} \subseteq \mathcal{Q}^{\mathbb{Z}}$ is the set of all invertible global transition functions of CA having state set Q .

Proof. Since any $\tau \in \overline{\mathcal{Q}^{\mathbb{Z}}} \subseteq \mathcal{Q}^{\mathbb{Z}}$ is invertible, the corollary holds true. □

Theorem 5.2. (\mathcal{T}^v, \circ) forms a monoid w.r.t. composition of IVTs.

Proof. The set \mathcal{T}^v , for $v \in \mathbb{N}$, is closed under composition ' \circ ' since for any

$(IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}), (IVT_{j_1}^{p,k}, IVT_{j_2}^{p,k}, \dots, IVT_{j_v}^{p,k}) \in \mathcal{T}^v$ and for any

$(n_1, n_2, \dots, n_v) \in \mathbb{N}_0^v \exists (m_1, m_2, \dots, m_v) \in \mathbb{N}_0^v$ such that,

$$\begin{aligned} & \left((IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}) \circ (IVT_{j_1}^{p,k}, IVT_{j_2}^{p,k}, \dots, IVT_{j_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \\ &= \left((IVT_{i_1}^{p,k} \circ IVT_{j_1}^{p,k}), (IVT_{i_2}^{p,k} \circ IVT_{j_2}^{p,k}), \dots, (IVT_{i_v}^{p,k} \circ IVT_{j_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \\ &= \left((IVT_{i_1}^{p,k} \circ IVT_{j_1}^{p,k})(\overline{n_1}), (IVT_{i_2}^{p,k} \circ IVT_{j_2}^{p,k})(\overline{n_2}), \dots, (IVT_{i_v}^{p,k} \circ IVT_{j_v}^{p,k})(\overline{n_v}) \right) \\ &= \left(IVT_{i_1}^{p,k}(IVT_{j_1}^{p,k}(\overline{n_1})), IVT_{i_2}^{p,k}(IVT_{j_2}^{p,k}(\overline{n_2})), \dots, IVT_{i_v}^{p,k}(IVT_{j_v}^{p,k}(\overline{n_v})) \right) \\ &= IVT_{i_1}^{p,k}(\overline{n_1}^*) IVT_{i_2}^{p,k}(\overline{n_2}^*) \dots IVT_{i_v}^{p,k}(\overline{n_v}^*) = (m_1, m_2, \dots, m_v)_p \end{aligned}$$

where $n_s^* = IVT_{j_s}^{p,k}(\overline{n_s}) = IVT_{j_s}^{p,k}(\eta_1^s, \eta_2^s, \dots, \eta_k^s)$ for $s = 1, 2, \dots, v$.

\mathcal{T}^v is associative under ' \circ ' since composition of functions are associative.

Again, for $(n_1, n_2, \dots, n_v) \in \mathbb{N}_0^v \exists (IVT_{e_1}^{p,k}, IVT_{e_2}^{p,k}, \dots, IVT_{e_v}^{p,k}) \in \mathcal{T}^v$ such that

$$\begin{aligned} & (IVT_{e_1}^{p,k}, IVT_{e_2}^{p,k}, \dots, IVT_{e_v}^{p,k})(n_1, n_2, \dots, n_v) \\ &= IVT_{e_1}^{p,k}(\overline{n_1}) IVT_{e_2}^{p,k}(\overline{n_2}) \dots IVT_{e_v}^{p,k}(\overline{n_v}) = (n_1, n_2, \dots, n_v) \end{aligned}$$

Since $IVT_{e_s}^{p,k}(\overline{n_s}) = n_s \forall s = 1, \dots, v$, it follows that $e_1 = e_2 = \dots = e_v = e(\text{say})$

Thus $(IVT_e^{p,k}, IVT_e^{p,k}, \dots, IVT_e^{p,k})$ is the identity element of \mathcal{T}^v

Hence the theorem. □

Definition 13. Modular addition and multiplication of two local transition functions for a $p(\in \mathbb{N})$ -state CA are defined $\forall i \in \mathbb{Z}$ as

$$\begin{aligned}(\mu_i^1 \oplus_p \mu_i^2)(\overline{c_i}) &= \mu_i^1(\overline{c_i}) \oplus_p \mu_i^2(\overline{c_i}) \quad \text{and} \\ (\mu_i^1 \otimes_p \mu_i^2)(\overline{c_i}) &= \mu_i^1(\overline{c_i}) \otimes_p \mu_i^2(\overline{c_i})\end{aligned}$$

Definition 14. Modular addition and multiplication of two $p(\in \mathbb{N})$ -base $k(\in \mathbb{N})$ -dimensional IVTs(see [18]) are defined $\forall (n_1, n_2, \dots, n_k) \in \mathbb{N}_0^k$ and $j_1, j_2 \in \{0, 1, \dots, p^{p^k} - 1\}$, as

$$\begin{aligned}(IVT_{j_1}^{p,k} \oplus_p IVT_{j_2}^{p,k})(n_1, n_2, \dots, n_k) &= IVT_{j_1}^{p,k}(n_1, n_2, \dots, n_k) \oplus_p IVT_{j_2}^{p,k}(n_1, n_2, \dots, n_k) \\ &= (f_{j_1}(a_0^{n_1}, \dots, a_0^{n_k}) \oplus_p f_{j_2}(a_0^{n_1}, \dots, a_0^{n_k}) \dots f_{j_1}(a_{l-1}^{n_1}, \dots, a_{l-1}^{n_k}) \oplus_p f_{j_2}(a_{l-1}^{n_1}, \dots, a_{l-1}^{n_k}))_p \quad \text{and}, \\ (IVT_{j_1}^{p,k} \otimes_p IVT_{j_2}^{p,k})(n_1, n_2, \dots, n_k) &= IVT_{j_1}^{p,k}(n_1, n_2, \dots, n_k) \otimes_p IVT_{j_2}^{p,k}(n_1, n_2, \dots, n_k) \\ &= (f_{j_1}(a_0^{n_1}, \dots, a_0^{n_k}) \otimes_p f_{j_2}(a_0^{n_1}, \dots, a_0^{n_k}) \dots f_{j_1}(a_{l-1}^{n_1}, \dots, a_{l-1}^{n_k}) \otimes_p f_{j_2}(a_{l-1}^{n_1}, \dots, a_{l-1}^{n_k}))_p\end{aligned}$$

where, $n_s = (a_0^{n_s}, a_1^{n_s}, \dots, a_{l-1}^{n_s})_p$ for $s = 1, 2, \dots, k$.

Theorem 5.3. $(\mathcal{Q}^{\mathbb{Z}}, \oplus_p, \otimes_p)$ forms a commutative ring with identity under the operations \oplus_p and \otimes_p defined as

$$\begin{aligned}(\tau_1 \oplus_p \tau_2)(C) &= \dots(\mu_{i-1}^1 \oplus_p \mu_{i-1}^2)(\overline{c_{i-1}}) \cdot (\mu_i^1 \oplus_p \mu_i^2)(\overline{c_i}) \cdot (\mu_{i+1}^1 \oplus_p \mu_{i+1}^2)(\overline{c_{i+1}}) \dots \quad \text{and} \\ (\tau_1 \otimes_p \tau_2)(C) &= \dots(\mu_{i-1}^1 \otimes_p \mu_{i-1}^2)(\overline{c_{i-1}}) \cdot (\mu_i^1 \otimes_p \mu_i^2)(\overline{c_i}) \cdot (\mu_{i+1}^1 \otimes_p \mu_{i+1}^2)(\overline{c_{i+1}}) \dots\end{aligned}$$

where $\tau_1, \tau_2 \in \mathcal{Q}^{\mathbb{Z}}$, $C \in Q^{\mathbb{Z}}$, \oplus_p denotes addition modulo p and \otimes_p denotes multiplication modulo p for $|Q| = p(\in \mathbb{N})$.

Proof. In a $p(\in \mathbb{N})$ -state CA, for any $\tau_1, \tau_2 \in \mathcal{Q}^{\mathbb{Z}}$,

$$(\tau_1 \oplus_p \tau_2)(C) \in Q^{\mathbb{Z}} \quad \text{and} \quad (\tau_1 \otimes_p \tau_2)(C) \in Q^{\mathbb{Z}}$$

since $\forall i \in \mathbb{Z}$,

$$(\mu_i^1 \oplus_p \mu_i^2)(\overline{c_i}) \in Q \quad \text{and} \quad (\mu_i^1 \otimes_p \mu_i^2)(\overline{c_i}) \in Q$$

Therefore $\mathcal{Q}^{\mathbb{Z}}$ is closed w.r.t. \oplus_p and \otimes_p .

Associativity follows from associativity of \oplus_p and \otimes_p .

The local transition function μ_0 such that $\forall c_i \in Q$, $\mu_0(\overline{c_i}) = 0$ is the additive identity since $\forall i \in \mathbb{Z}$

$$(\mu_0 \oplus_p \mu_i)(\overline{c_i}) = (\mu_i \oplus_p \mu_0)(\overline{c_i}) = \mu_i(\overline{c_i})$$

It follows that $\tau_0 \in \mathcal{Q}^{\mathbb{Z}}$ is the additive identity such that $\forall C \in Q^{\mathbb{Z}}$,

$$\tau_0(C) = \dots \mu_0(\overline{c_{i-1}}) \mu_0(\overline{c_i}) \mu_0(\overline{c_{i+1}}) \dots$$

The local transition function μ_{id} such that $\forall c_i \in Q$, $\mu_{id}(\overline{c_i}) = 1$ is the multiplicative identity since $\forall i \in \mathbb{Z}$

$$(\mu_{id} \otimes_p \mu_i)(\overline{c_i}) = (\mu_i \otimes_p \mu_{id})(\overline{c_i}) = \mu_i(\overline{c_i})$$

It follows that $\tau_{id} \in \mathcal{Q}^{\mathbb{Z}}$ is the multiplicative identity such that $\forall C \in Q^{\mathbb{Z}}$,

$$\tau_{id}(C) = \dots \mu_{id}(\overline{c_{i-1}}) \mu_{id}(\overline{c_i}) \mu_{id}(\overline{c_{i+1}}) \dots$$

Now any local transition function μ_i , is a p^k -bit sequence $b_1 b_2 \dots b_{p^k}$ (say), for a $p(\in \mathbb{N})$ -state $k(\in \mathbb{N})$ -neighbourhood CA where $b_1, \dots, b_{p^k} \in Q = \{0, 1, \dots, p\}$. Clearly for any $b_s \in Q$, $\exists b_s^{-1} \in Q$ such that $\forall s = 1, 2, \dots, p^k$, $b_s \oplus_p b_s^{-1} = 0$. Thus for any $\mu_i (\equiv b_1 \dots b_{p^k})$, $\exists \mu_i^{-1} (\equiv b_1^{-1} \dots b_{p^k}^{-1})$ such that $\forall c_i \in Q$,

$$(\mu_i^{-1} \oplus_p \mu_i)(\overline{c_i}) = (\mu_i \oplus_p \mu_i^{-1})(\overline{c_i}) = \mu_0(\overline{c_i}) = 0$$

It follows that for any $\tau \in \mathcal{Q}^{\mathbb{Z}}$, $\exists \tau^{-1} \in \mathcal{Q}^{\mathbb{Z}}$ such that $\forall C \in Q$,

$$(\tau \oplus_p \tau^{-1})(C) = \dots (\mu_{i-1} \oplus_p \mu_{i-1}^{-1})(\overline{c_{i-1}}) (\mu_i \oplus_p \mu_i^{-1})(\overline{c_i}) (\mu_{i+1} \oplus_p \mu_{i+1}^{-1})(\overline{c_{i+1}}) \dots = \tau_0(C),$$

$$(\tau^{-1} \oplus_p \tau)(C) = \dots (\mu_{i-1}^{-1} \oplus_p \mu_{i-1})(\overline{c_{i-1}}) (\mu_i^{-1} \oplus_p \mu_i)(\overline{c_i}) (\mu_{i+1}^{-1} \oplus_p \mu_{i+1})(\overline{c_{i+1}}) \dots = \tau_0(C),$$

Thus for any global transition function τ , its additive inverse exists in $\mathcal{Q}^{\mathbb{Z}}$.

Commutativity follows from commutativity of \oplus_p and \otimes_p .

Now for any local transitions $\mu_i^1, \mu_i^2, \mu_i^3, \forall c_i \in Q, i \in \mathbb{Z}$ we get,

$$(\mu_i^1 \otimes_p (\mu_i^2 \oplus_p \mu_i^3))(\overline{c_i}) = (\mu_i^1 \otimes_p \mu_i^2)(\overline{c_i}) \oplus_p (\mu_i^1 \otimes_p \mu_i^3)(\overline{c_i}) \text{ and,}$$

$$((\mu_i^2 \oplus_p \mu_i^3) \otimes_p \mu_i^1)(\overline{c_i}) = (\mu_i^2 \otimes_p \mu_i^1)(\overline{c_i}) \oplus_p (\mu_i^3 \otimes_p \mu_i^1)(\overline{c_i})$$

Therefore for $\tau_1, \tau_2, \tau_3 \in (Q)^{\mathbb{Z}}$, $\forall C \in Q^{\mathbb{Z}}$ we get,

$$(\tau_1 \otimes_p (\tau_2 \oplus_p \tau_3))(C) = (\tau_1 \otimes_p \tau_2)(C) \oplus_p (\tau_1 \otimes_p \tau_3)(C) \text{ and,}$$

$$((\tau_2 \oplus_p \tau_3) \otimes_p \tau_1)(C) = (\tau_2 \otimes_p \tau_1)(C) \oplus_p (\tau_3 \otimes_p \tau_1)(C)$$

Hence the theorem. \square

Remark 5.1. The following example shows that $(\mathcal{Q}^{\mathbb{Z}}, \oplus_p, \otimes_p)$ will not form a field under the operations \oplus_p and \otimes_p even if $|Q| = p$ is prime.

Example 5.1. For initial configuration $C = (10001)_2$, let

$$\tau(10001)_2 = (\mu(110)\mu(100)\mu(000)\mu(001)\mu(011))_2$$

Let μ follow Wolfram code $3(\neg(c_{i-1} \vee c_i))$.

Then $\tau(10001)_2 = (\mu_3(110)\mu_3(100)\mu_3(000)\mu_3(001)\mu_3(011))_2 = (00110)_2$.

Now, $\tau_{id}(10001)_2 = (\mu_{id}(110)\mu_{id}(100)\mu_{id}(000)\mu_{id}(001)\mu_{id}(011))_2 = (11111)_2$.

But $\nexists \tau^{-1}$ such that $(\tau \otimes_2 \tau^{-1})(10001)_2 = (11111)_2$ since $\nexists \mu^{-1}$ such that for all local configurations $\mu \otimes_2 \mu^{-1} = 1$.

Theorem 5.4. $(\mathcal{T}^v, \oplus_p, \otimes_p)$ forms a commutative ring under the operations \oplus_p and \otimes_p defined as

$$\begin{aligned} & \left((IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}) \oplus_p (IVT_{j_1}^{p,k}, IVT_{j_2}^{p,k}, \dots, IVT_{j_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \\ &= \left((IVT_{i_1}^{p,k} \oplus_p IVT_{j_1}^{p,k}), (IVT_{i_2}^{p,k} \oplus_p IVT_{j_2}^{p,k}), \dots, (IVT_{i_v}^{p,k} \oplus_p IVT_{j_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \\ &= \left((IVT_{i_1}^{p,k} \oplus_p IVT_{j_1}^{p,k})(\overline{n_1}), (IVT_{i_2}^{p,k} \oplus_p IVT_{j_2}^{p,k})(\overline{n_2}), \dots, (IVT_{i_v}^{p,k} \oplus_p IVT_{j_v}^{p,k})(\overline{n_v}) \right) \\ & \text{and, } \left((IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}) \otimes_p (IVT_{j_1}^{p,k}, IVT_{j_2}^{p,k}, \dots, IVT_{j_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \\ &= \left((IVT_{i_1}^{p,k} \otimes_p IVT_{j_1}^{p,k}), (IVT_{i_2}^{p,k} \otimes_p IVT_{j_2}^{p,k}), \dots, (IVT_{i_v}^{p,k} \otimes_p IVT_{j_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \\ &= \left((IVT_{i_1}^{p,k} \otimes_p IVT_{j_1}^{p,k})(\overline{n_1}), (IVT_{i_2}^{p,k} \otimes_p IVT_{j_2}^{p,k})(\overline{n_2}), \dots, (IVT_{i_v}^{p,k} \otimes_p IVT_{j_v}^{p,k})(\overline{n_v}) \right) \end{aligned}$$

where $i_s, j_s \in \{0, 1, \dots, p^{p^k} - 1\}$, $p, k \in \mathbb{N}$, $\overline{n_s} = (\eta_1^s, \eta_2^s, \dots, \eta_k^s)$ for $s = 1, 2, \dots, v$

Proof. For any $(IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}), (IVT_{j_1}^{p,k}, IVT_{j_2}^{p,k}, \dots, IVT_{j_v}^{p,k}) \in \mathcal{T}^v$, and $(n_1, n_2, \dots, n_v) \in \mathbb{N}_0^v$,

$$\begin{aligned} & \left((IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}) \oplus_p (IVT_{j_1}^{p,k}, IVT_{j_2}^{p,k}, \dots, IVT_{j_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \in \mathbb{N}^v \\ & \text{and, } \left((IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}) \otimes_p (IVT_{j_1}^{p,k}, IVT_{j_2}^{p,k}, \dots, IVT_{j_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \in \mathbb{N}^v \end{aligned}$$

since for every $s = 1, 2, \dots, v$,

$$(IVT_{i_s}^{p,k} \oplus_p IVT_{j_s}^{p,k})(\overline{n_s}) \in \mathbb{N}_0 \quad \text{and} \quad (IVT_{i_s}^{p,k} \otimes_p IVT_{j_s}^{p,k})(\overline{n_s}) \in \mathbb{N}_0$$

where $\overline{n_s} = (\eta_1^s, \eta_2^s, \dots, \eta_k^s)$

Therefore \mathcal{T}^v is closed w.r.t. \oplus_p and \otimes_p for $p \in \mathbb{N}$.

Associativity follows from associativity of \oplus_p and \otimes_p .

$IVT_0^{p,k} \in T^{p,k}$ is such that $\forall \overline{n_s} \in \mathbb{N}_0^k$, $IVT_0^{p,k}(\overline{n_s}) = 0$ is the additive identity of $T^{p,k}$ since $\forall s = 1, 2, \dots, v$

$$(IVT_0^{p,k} \oplus_p IVT_s^{p,k})(\overline{n_s}) = (IVT_s^{p,k} \oplus_p IVT_0^{p,k})(\overline{n_s}) = IVT_s^{p,k}(\overline{n_s})$$

where, for each $(\overline{n_s}) = (\eta_1, \eta_2, \dots, \eta_k) \in \mathbb{N}_0^k$,

$$IVT_0^{p,k}(\eta_1, \eta_2, \dots, \eta_k) = (f_0(a_0^{\eta_1}, \dots, a_0^{\eta_k}), \dots, f_0(a_{l-1}^{\eta_1}, \dots, a_{l-1}^{\eta_k})) = 0$$

where, $\eta_u = (a_0^{\eta_u} a_1^{\eta_u} \dots a_{l-1}^{\eta_u})_p$ for $u = 1, 2, \dots, k$.

It follows that $(IVT_0^{p,k}, IVT_0^{p,k}, \dots, IVT_0^{p,k})$ is the additive identity of \mathcal{T}^v such that for any $(n_1, n_2, \dots, n_v) \in \mathbb{N}_0^v$

$$(IVT_0^{p,k}, IVT_0^{p,k}, \dots, IVT_0^{p,k})(n_1, n_2, \dots, n_v) = (IVT_0^{p,k}(\overline{n_1}), IVT_0^{p,k}(\overline{n_2}), \dots, IVT_0^{p,k}(\overline{n_v}))$$

For any $IVT_s^{p,k} \in T^{p,k} \exists IVT_{-s}^{p,k} \in T^{p,k}$ such that $\forall \overline{n_s} \in \mathbb{N}_0^k$,

$$(IVT_s^{p,k} \oplus_p IVT_{-s}^{p,k})(\overline{n_s}) = (IVT_{-s}^{p,k} \oplus_p IVT_s^{p,k})(\overline{n_s}) = IVT_0^{p,k}(\overline{n_s}) = 0$$

It follows that for any $(IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}) \in \mathcal{T}^v$, $\exists (IVT_{-i_1}^{p,k}, IVT_{-i_2}^{p,k}, \dots, IVT_{-i_v}^{p,k}) \in \mathcal{T}^v$ such that $\forall (n_1, n_2, \dots, n_v) \in \mathbb{N}_0^v$,

$$\begin{aligned} & \left((IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}) \oplus_p (IVT_{-i_1}^{p,k}, IVT_{-i_2}^{p,k}, \dots, IVT_{-i_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \\ & = (IVT_0^{p,k}, IVT_0^{p,k}, \dots, IVT_0^{p,k})(n_1, n_2, \dots, n_v) \\ & = \left((IVT_{-i_1}^{p,k}, IVT_{-i_2}^{p,k}, \dots, IVT_{-i_v}^{p,k}) \oplus_p (IVT_{i_1}^{p,k}, IVT_{i_2}^{p,k}, \dots, IVT_{i_v}^{p,k}) \right) (n_1, n_2, \dots, n_v) \end{aligned}$$

Thus for any element of \mathcal{T}^v its additive inverse exists in \mathcal{T}^v .

Commutativity follows from commutativity of \oplus_p and \otimes_p .

Now for any $IVT_h^{p,k}, IVT_i^{p,k}, IVT_j^{p,k} \in T^{p,k}$, $\forall \overline{n_s} \in \mathbb{N}_0^k$, we get,

$$(IVT_h^{p,k} \otimes_p (IVT_i^{p,k} \oplus_p IVT_j^{p,k}))(\overline{n_s}) = (IVT_h^{p,k} \otimes_p IVT_i^{p,k})(\overline{n_s}) \oplus_p (IVT_h^{p,k} \otimes_p IVT_j^{p,k})(\overline{n_s}),$$

$$((IVT_i^{p,k} \oplus_p IVT_j^{p,k}) \otimes_p IVT_h^{p,k})(\overline{n_s}) = (IVT_i^{p,k} \otimes_p IVT_h^{p,k})(\overline{n_s}) \oplus_p (IVT_j^{p,k} \otimes_p IVT_h^{p,k})(\overline{n_s})$$

Thus for any $(IVT_{h_1}^{p,k}, \dots, IVT_{h_v}^{p,k}), (IVT_{i_1}^{p,k}, \dots, IVT_{i_v}^{p,k}), (IVT_{j_1}^{p,k}, \dots, IVT_{j_v}^{p,k}) \in \mathcal{T}^v$, $\forall (n_1, \dots, n_v) \in \mathbb{N}_0^v$,

$$\begin{aligned}
& \left(IVT_{h_1}^{p,k}, \dots, IVT_{h_v}^{p,k} \right) \otimes_p \left(\left(IVT_{i_1}^{p,k}, \dots, IVT_{i_v}^{p,k} \right) \oplus_p \left(IVT_{j_1}^{p,k}, \dots, IVT_{j_v}^{p,k} \right) \right) (n_1, \dots, n_v) \\
&= \left(\left(IVT_{h_1}^{p,k}, \dots, IVT_{h_v}^{p,k} \right) \otimes_p \left(IVT_{i_1}^{p,k}, \dots, IVT_{i_v}^{p,k} \right) \right) (n_1, \dots, n_v) \\
&\oplus_p \left(\left(IVT_{h_1}^{p,k}, \dots, IVT_{h_v}^{p,k} \right) \otimes_p \left(IVT_{j_1}^{p,k}, \dots, IVT_{j_v}^{p,k} \right) \right) (n_1, \dots, n_v), \text{ and} \\
&\left(\left(\left(IVT_{i_1}^{p,k}, \dots, IVT_{i_v}^{p,k} \right) \oplus_p \left(IVT_{j_1}^{p,k}, \dots, IVT_{j_v}^{p,k} \right) \right) \otimes_p \left(IVT_{h_1}^{p,k}, \dots, IVT_{h_v}^{p,k} \right) \right) (n_1, \dots, n_v) \\
&= \left(\left(IVT_{i_1}^{p,k}, \dots, IVT_{i_v}^{p,k} \right) \otimes_p \left(IVT_{h_1}^{p,k}, \dots, IVT_{h_v}^{p,k} \right) \right) (n_1, \dots, n_v) \\
&\oplus_p \left(\left(IVT_{j_1}^{p,k}, \dots, IVT_{j_v}^{p,k} \right) \otimes_p \left(IVT_{h_1}^{p,k}, \dots, IVT_{h_v}^{p,k} \right) \right) (n_1, \dots, n_v)
\end{aligned}$$

Hence the theorem. \square

Theorem 5.5. $(\mathcal{T}_{|Q}^v, \oplus_p, \otimes_p)$ forms a commutative ring with identity under the operations \oplus_p and \otimes_p .

Proof. Clearly $(\mathcal{T}_{|Q}^v, \oplus_p, \otimes_p)$ for $p, k, v \in \mathbb{N}$ is a commutative ring since $\mathcal{T}_{|Q}^v \subseteq \mathcal{T}^v$ when \mathbb{N}_0^v is restricted to Q^v .

Now, for any $(\eta_1, \eta_2, \dots, \eta_k) \in Q^k$, \exists multiplicative identity $\underline{IVT}_{id}^{p,k} \in T_{|Q}^{p,k}$ such that

$$\underline{IVT}_{id}^{p,k}(\eta_1, \eta_2, \dots, \eta_k) = (f_{id}(\eta_1, \eta_2, \dots, \eta_k))_p = 1$$

$$\text{since } (\underline{IVT}_j^{p,k} \otimes_p \underline{IVT}_{id}^{p,k})(\eta_1, \eta_2, \dots, \eta_k) = \underline{IVT}_j^{p,k}(\eta_1, \eta_2, \dots, \eta_k) = (f_j(\eta_1, \eta_2, \dots, \eta_k))_p,$$

$$\text{and } (\underline{IVT}_{id}^{p,k} \otimes_p \underline{IVT}_j^{p,k})(\eta_1, \eta_2, \dots, \eta_k) = \underline{IVT}_j^{p,k}(\eta_1, \eta_2, \dots, \eta_k) = (f_j(\eta_1, \eta_2, \dots, \eta_k))_p$$

It follows that $(\underline{IVT}_{id}^{p,k}, \underline{IVT}_{id}^{p,k}, \dots, \underline{IVT}_{id}^{p,k}) \in \mathcal{T}_{|Q}^v$ is the multiplicative identity of $\mathcal{T}_{|Q}^v$ such that $\forall (n_1, n_2, \dots, n_v) \in Q^v$,

$$(\underline{IVT}_{id}^{p,k}, \underline{IVT}_{id}^{p,k}, \dots, \underline{IVT}_{id}^{p,k})(n_1, n_2, \dots, n_v) = \underline{IVT}_{id}^{p,k}(\overline{n_1}) \underline{IVT}_{id}^{p,k}(\overline{n_2}) \dots \underline{IVT}_{id}^{p,k}(\overline{n_v})$$

where $\underline{IVT}_{id}^{p,k}(\overline{n_s}) = \underline{IVT}_{id}^{p,k}(\eta_1^s, \eta_2^s, \dots, \eta_k^s) = 1$ for each $s = 1, 2, \dots, v$.

Hence the theorem. \square

6. Modelling of DNA Sequence Evolution

DNA can be modelled as a one-dimensional CA using rule matrix multiplication for linear CA (reported in [23]). The DNA sequence corresponds to the CA lattice and the deoxyribose sugars to the CA cells. The 4 sugar bases A, C, T and G correspond to 4 possible states of the CA cell.

Now, 1-neighbourhood CA transition rule is practically absurd since the transition of cell states of a CA is dependent on neighbouring cells. Consequently, computation using 4-state, $k \in \mathbb{N} \geq 2$ -neighbourhood CA maybe complex. However, in the domain of IVTs, computations using one-dimensional IVTs hold good. Moreover, for non-linear transition rules of a CA, the corresponding rule matrix maynot be obtained in general.

Hence, modelling DNA sequence evolution in terms of one-dimensional IVTs of 4-base is a suitable alternative (see [24]).

For any one-variable system having base 4, there can be $4^{4^1} = 256$ different functions of which $4^1 = 4$ are linear functions and the others non-linear. The following example depicts DNA sequence evolution in terms of IVTs where the sugar bases have been represented with numbers as follows:

$$A \rightarrow 0, C \rightarrow 1, T \rightarrow 2, G \rightarrow 3$$

Example 6.1. Let CTCTAGAGGGAA be a particular DNA strand of length 12 at some time. This strand corresponds to the number sequence 121203033300.

Evolution of this strand at the next time step can be represented by any $(IVT_{j_1}^{4,1}, \dots, IVT_{j_{12}}^{4,1}) \in \tau^{12}$ where $j_1, \dots, j_{12} \in \{0, 1, \dots, 4^4 - 1\}$.

Let the DNA strand be broken into 2 blocks of 6 sugar bases each.

If the first block evolves according to non-linear function $f_{78}(0 \rightarrow 2, 1 \rightarrow 3, 2 \rightarrow 0, 3 \rightarrow 1)$ and the second block evolves according to linear function $f_{108}(0 \rightarrow 0, 1 \rightarrow 3, 2 \rightarrow 2, 3 \rightarrow 1)$ then,

$$\begin{aligned} & (IVT_{j_1}^{4,1}, \dots, IVT_{j_{12}}^{4,1}) (\underbrace{121203}_{\text{block1}} \underbrace{033300}_{\text{block2}})_4 \\ &= (f_{78}(1)f_{78}(2)f_{78}(1)f_{78}(2)f_{78}(0)f_{78}(3)f_{108}(0)f_{108}(3)f_{108}(3)f_{108}(3)f_{108}(0)f_{108}(0))_4 \\ &= (303021011100)_4 \end{aligned}$$

Thus the DNA strand evolved at the next step would be GAGATCACCCAA. Hence the evolution pattern at any later time step can be obtained recursively.

7. Conclusion

In this paper Wolfram code and transition function of a one-dimensional CA has been represented in terms of an IVT. Moreover, if the domain of IVTs is restricted from positive integers to $p(\in \mathbb{N})$ -base numbers, then an odd dimensional IVT is equivalent to a local transition function of a CA. It could be proved that the set of all transition functions of a $p(\in \mathbb{N})$ -state CA, forms a commutative ring with identity, whereas the set of p -base IVTs forms only a commutative ring under modular addition and multiplication. However, for the set of p -base IVTs multiplicative identity under modular addition and multiplication exists, when the domain of IVTs is restricted from positive integers to $p(\in \mathbb{N})$ -base numbers. The novelty of using IVTs instead of one-dimensional CA, for modelling DNA sequence evolution has been discussed and depicted in the last section.

References

- [1] J. von Neumann, A. W. Burks, et al., Theory of self-reproducing automata, IEEE Transactions on Neural Networks 5 (1) (1966) 3–14.
- [2] J. L. Schif, Cellular automata: a discrete view of the world, Vol. 45, John Wiley & Sons, 2011.
- [3] S. Ulam, On some mathematical problems connected with patterns of growth of figures, in: Proceedings of Symposia in Applied Mathematics, Vol. 14, Am. Math. Soc. Vol. 14, Providence, 1962, pp. 215–224.
- [4] S. Ghosh, S. Basu, Evolution patterns of some boolean cellular automata having atmost one active cell to model simple dynamical systems, Bull. Cal. Math Soc. 108 (6) (2016) 449–464.
- [5] S. Ghosh, S. Basu, Evolution patterns of finite celled synchronous cellular automata having atmost one active cell, Proceedings of The 10th International Conference MSAST 2016 (2016) 154–164.
- [6] S. Wolfram, A new kind of science, Wolfram Media, Inc., Champaign, IL, USA, 2002.

- [7] P. P. Choudhury, S. Sahoo, M. Chakraborty, S. K. Bhandari, A. Pal, Investigation of the global dynamics of cellular automata using boolean derivatives, *Computers & Mathematics with Applications* 57 (8) (2009) 1337–1351.
- [8] A. K. Das, A. Sanyal, P. Palchaudhuri, On characterization of cellular automata with matrix algebra, *Information sciences* 61 (3) (1992) 251–277.
- [9] A. Wuensche, Attractor basins of discrete networks, *Cognitive Science Research Paper*.
- [10] A. Wuensche, Discrete dynamics lab: Tools for investigating cellular automata and discrete dynamical networks, *Kybernetes*.
- [11] X. Xu, Y. Song, S. P. Banks, On the dynamical behavior of cellular automata, *International Journal of Bifurcation and Chaos* 19 (04) (2009) 1147–1156.
- [12] R. Edwards, A. Maignan, A class of discrete dynamical systems with properties of both cellular automata and l-systems, *Natural Computing* (2019) 1–33.
- [13] S. S. Hassan, P. P. Choudhury, S. Sahoo, R. B. Singh, B. K. Nayak, Collatz function like integral value transformations, *Alexandria journal of Mathematics* 1 (2) (2012) 30–35.
- [14] S. S. Hassan, P. P. Choudhury, B. K. Nayak, A. Ghosh, J. Banerjee, Integral value transformations: a class of affine discrete dynamical systems and an application, *arXiv preprint arXiv:1110.0724*.
- [15] J. K. Das, S. Sahoo, S. Hassan, P. P. Choudhury, Two dimensional discrete dynamics of integral value transformations, *arXiv preprint arXiv:1709.05205*.
- [16] S. Ghosh, S. Basu, Some algebraic properties of linear synchronous cellular automata, *arXiv preprint arXiv:1708.09751*.
- [17] J. Kari, Theory of cellular automata: A survey, *Theoretical computer science* 334 (1-3) (2005) 3–33.
- [18] S. S. Hassan, B. K. Nayak, P. P. Choudhury, One-dimensional p-adic integral value transformation, *Journal of Orissa Mathematical Society* 31 (2) (2012) 116–124.
- [19] P. P. Choudhury, S. Sahoo, B. K. Nayak, S. Hassan, Theory of carry value transformation (cvt) and its application in fractal formation, *arXiv preprint arXiv:0904.0534*.
- [20] P. P. Choudhury, S. S. Hassan, S. Sahoo, B. K. Nayak, Act of cvt and evt in the formation of number theoretic fractals, *International Journal of Computational Cognition* 9 (1) (2011) 1–8.
- [21] J. K. Das, P. P. Choudhury, S. Sahoo, Multi-number cvt-xor arithmetic operations in any base system and its significant properties, in: *2016 IEEE 6th International Conference on Advanced Computing (IACC)*, IEEE, 2016, pp. 769–773.
- [22] S. Pal, S. Sahoo, B. K. Nayak, Properties of carry value transformation, *International Journal of Mathematics and Mathematical Sciences* 2012.

- [23] C. Mizas, G. C. Sirakoulis, V. Mardiris, I. Karafyllidis, N. Glykos, R. Sandaltzopoulos, Reconstruction of dna sequences using genetic algorithms and cellular automata: Towards mutation prediction?, *Biosystems* 92 (1) (2008) 61–68.
- [24] S. S. Hassan, P. P. Choudhury, R. Guha, S. Chakraborty, A. Goswami, Dna sequence evolution through integral value transformations, *Interdisciplinary Sciences: Computational Life Sciences* 4 (2) (2012) 128–132.