# Comparison of correlation-based measures of concordance in terms of asymptotic variance

Takaaki Koike<sup>a,\*</sup>, Marius Hofert<sup>b</sup>

<sup>a</sup>Graduate School of Economics, Hitotsubashi University, Naka, Kunitachi, Tokyo, Japan. <sup>b</sup>Department of Statistics and Actuarial Science, Faculty of Science, The University of Hong Kong, Pokfulam, Hong Kong.

# Abstract

We compare measures of concordance that arise as Pearson's linear correlation coefficient between two random variables transformed so that they follow the so-called concordance-inducing distributions. The class of such transformed rank correlations includes Spearman's rho, Blomqvist's beta and van der Waerden's coefficient. When only the standard axioms of measures of concordance are required, it is not always clear which transformed rank correlation is most suitable to use. To address this question, we compare measures of concordance in terms of their best and worst asymptotic variances of some canonical estimators over a certain set of dependence structures. A simple criterion derived from this approach is that concordance-inducing distributions with smaller fourth moment are more preferable. In particular, we show that Blomqvist's beta is the optimal transformed rank correlation in this sense, and Spearman's rho outperforms van der Waerden's coefficient. Moreover, we find that Kendall's tau, although it is not a transformed rank correlation of that nature, shares a certain optimal structure with Blomqvist's beta.

#### Keywords:

Blomqvist's beta, Copula, Correlation coefficient, Kendall's tau, Measure of concordance, Spearman's rho. 2020 MSC: 62H20, 62H12

# 1. Introduction


A pair of random variables is said to be more concordant (discordant) if large values of one random variable are more likely to correspond to large (small) values of the other random variable. When the random variables are continuous, concordance and discordance are the properties of their (bivariate) *copula C*, which is a bivariate distribution function with standard uniform univariate marginal distributions. A measure of concordance quantifies concordance or discordance of *C* by a single number  $\kappa$  in [-1, 1]; see Definition 1 below.

In this paper we consider *G*-transformed rank correlations, a subclass of measures of concordance which can be represented as Pearson's linear correlation coefficient  $\rho$  between two random variables transformed so that they follow the so-called concordance-inducing distribution *G*. For a distribution function *G* on  $\mathbb{R}$ , the *G*-transformed rank correlation of a copula *C* is given by

$$\kappa_G(U, V) = \rho(G^{-1}(U), G^{-1}(V)), \quad (U, V) \sim C,$$

where  $G^{-1}$  is the quantile function of *G*. This measure  $\kappa_G$  satisfies the axioms of a measure of concordance [14] by taking *G* to be concordance-inducing; see Definition 2. This class contains popular measures of concordance, such as Spearman's rho  $\rho_S$ , Blomqvist's beta  $\beta$  and van der Waerden's coefficient  $\zeta$ ; see Example 1 below. Such a correlation representation of a measure of concordance  $\kappa$  is of great benefit to intuitively understand and explain the construction and ideas behind  $\kappa$  (see Fig. 1), to construct its estimators and investigate their asymptotic properties, and to analyze robustness [12] and matrix compatibility [7] of  $\kappa$ . Moreover, [9] showed that the class of transformed rank correlations exhausts all measures of concordance of the form  $\kappa(U, V) = \rho(g_1(U), g_2(V))$  for two possibly discontinuous functions  $g_1, g_2 : (0, 1) \rightarrow \mathbb{R}$ .

<sup>\*</sup>Corresponding author. Email address: takaaki.koike@r.hit-u.ac.jp



**Fig. 1:** Illustration of the process of calculating the *G*-transformed rank correlation  $\kappa_G$  where *G* is the standard normal distribution. The original data in (a) is first transformed so that their margins follow the standard uniform distribution. The transformed data in (b) is then converted so that their margins have the distribution *G*. Finally, the *G*-transformed rank correlation  $\kappa_G$  is calculated as a Pearson's linear correlation coefficient of the transformed data in (c).

When only the axioms of a measure of concordance are required, it is not always clear which transformed rank correlation is most suitable to use. For the related literature, [3] compares Pearson's linear correlation coefficient and Spearman's rho by numerical experiments in terms of bias, variance and robustness to outliers. Rank correlations are also extensively compared in [18]. Various measures of concordance are compared in terms of their power in tests of independence; see, for example, [6], [10] and [13].

To address this natural question of how to choose the concordance-inducing distribution, the aim of the present paper is to compare *G*-transformed rank correlations in terms of their stability concerning statistical estimation. Estimation of  $\kappa_G$  is often inevitable since an explicit form of  $\kappa_G(C)$  is not always available. For the purpose of comparison, we consider a simplified setting where independent and identically distributed (i.i.d.) samples from the underlying copula *C* are available, and the *G*-transformed rank correlation  $\kappa_G$  is estimated by the so-called canonical estimator  $\hat{\kappa}_G$ (Definition 3). Although this simplified setting may be rarely the case, the asymptotic variance  $\sigma_G^2(C)$  of the canonical estimator  $\hat{\kappa}_G$  is obtained in a tractable form. For an underlying copula *C*, a concordance-inducing distribution *G* can be more preferable to another one *G'* in terms of the stability of statistical estimation if  $\sigma_G^2(C) \leq \sigma_{G'}^2(C)$ . Since this comparison is valid only for a specific copula *C*, we consider a set of copulas  $\mathcal{D}$  and compare concordance-inducing distributions by the largest and smallest values of  $\sigma_G^2(C)$  over  $\mathcal{D}$ . Namely, a concordance-inducing distribution *G* is more preferable to another one *G'* in  $\mathcal{D}$  if the worst and best asymptotic variances of a canonical estimator  $\hat{\kappa}_G$  of  $\kappa_G$  on  $\mathcal{D}$  are smaller than those of *G'*. For such *G* and *G'*, if  $\mathcal{D}$  represents possible dependence structures which the analyst is interested in quantifying and comparing, she may be more willing to use  $\kappa_G$  instead if  $\kappa'_G$  since the former is expected to be estimated more accurately than the latter.

To illustrate the motivation of this work, let us consider the problem of quantifying the dependence among two exchange rates JPY/USD and CAD/USD from 2000 to 2015. Following [1] and [5], we model the daily log increments of each exchange rate in each year by a GARCH(1,1) model with skew-*t* innovation distribution. We then capture the dependence among the two time series by the copula of the bivariate standardized residuals. Based on the filtered i.i.d. samples, a transformed rank correlation  $\kappa_G$  is estimated by the canonical estimator  $\hat{\kappa}_G$  for different choices of *G*. Other than  $\beta$ ,  $\rho_S$  and  $\zeta$ , a standardized Student *t* distribution with  $\nu = 10$  degrees of freedom and a beta distribution with parameter vector (0.5, 0.5) are also considered as concordance-inducing distribution *G*. The corresponding asymptotic variances  $\sigma_G^2$  are estimated from samples as in Section 6. The results are plotted in Fig. 2, which also shows that the yearly trend of dependence is captured similarly by all the chosen measures of concordance, although some deviations are observed for  $\beta$ . Therefore, one may be willing to choose measures that are more stable to estimate. From the right plot in Fig. 2,  $\zeta$  and *t*-based rank correlations tend to have larger asymptotic variances than others, and  $\beta$  often attains relatively small  $\sigma_G^2$ . From these observations,  $\beta$ ,  $\rho_S$  and the beta distribution-based rank correlations may be preferable over  $\zeta$  and the *t* distribution-based rank correlation, at least in terms of the stability of their statistical estimation.

From this approach for comparing transformed rank correlations, we derive a criterion under some reasonable assumptions on  $\mathcal{D}$  that a concordance-inducing distribution *G* with smaller variance  $\operatorname{Var}_G(X^2)$ , where  $X \sim G$ , is more




Fig. 2: Estimates of *G*-transformed rank correlations (left) and their asymptotic variances (right) of dependence among the daily log increments of exchange rates JPY/USD and CAD/USD filtered by skew *t* GARCH(1,1) model each year in the period of 2000–2015.

preferable. Therefore, heavy-tailed concordance-inducing distributions, such as the Student *t* distribution, are not preferable (at least in terms of asymptotic variance) as a choice of *G* in comparison to a normal distribution, which leads to van der Waerden's coefficient  $\zeta$ . It is also deduced that  $\rho_S$  outperforms  $\zeta$ , and the beta distribution-based transformed rank correlations are more preferable to  $\rho_S$  in this sense. In particular, we prove that Blomqvist's beta is the unique optimal transformed rank correlation attaining the optimal best and worst asymptotic variances under certain conditions on  $\mathcal{D}$ . As stated by [15], one of the advantages of  $\beta$  over other measures of concordance is that it admits an explicit form if the copula can be written explicitly. The optimality of  $\beta$  provides its additional advantage in terms of accuracy of its estimation.

Despite the advantages of  $\beta$ , its major drawback is that it depends only on the local value C(1/2, 1/2) of the underlying copula *C*. Considering this drawback, we also compare *G*-transformed rank correlations with Kendall's tau  $\tau$ . Based on the representation of  $\tau$  in terms of Pearson's linear correlation coefficient, we find that  $\tau$  also attains the optimal best and worst asymptotic variances that  $\beta$  does. Therefore,  $\tau$  can be a good alternative to  $\beta$  in terms of the accuracy of its estimation even though the optimality of  $\tau$  is partly violated if the sample size required to construct its estimator is taken into account (see Remark 4 below). Finally, in a simulation study, we find that the choice of concordance-inducing distribution *G* and the strength of dependence of the underlying copula *C* affect the asymptotic variance of  $\hat{\kappa}_G$  more than the model of dependence.

This paper is organized as follows. In Section 2, we review measures of concordance, the class of *G*-transformed rank correlations  $\kappa_G$  and their basic properties. In Section 3 we introduce the approach for comparing *G*-transformed rank correlations in terms of their asymptotic variances. A canonical estimator of  $\kappa_G$  is presented in Section 3.1, and the properties of its asymptotic variance  $\sigma_G^2(C)$  are studied in Section 3.2. Section 3.3 addresses effects of location-scale transforms of *G* on  $\sigma_G^2(C)$ . In Section 4, we study optimal best and worst asymptotic variances and their attaining concordance-inducing distributions on  $\mathcal{D}$ . The case when  $\mathcal{D}$  is a set of fundamental or Fréchet copulas is analyzed in Section 4.1. The optimality of Blomqvist's beta is proved in Section 4.2, and its uniqueness is discussed in Section 4.3. Kendall's tau and *G*-transformed rank correlations are compared in Section 5. In Section 6, a simulation study is conducted to compare the asymptotic variances for various parametric copulas and concordance-inducing distributions. Section 7 concludes this work with discussions about directions for future research. Proofs of the statements are given in Section 8.

# 2. Correlation-based measures of concordance

Let  $C_2$  be the set of all bivariate copulas, that is, all bivariate distribution functions with standard uniform marginal distributions. We call  $C' \in C_2$  more concordant than  $C \in C_2$ , denoted by  $C \leq C'$ , if  $C(u, v) \leq C'(u, v)$  for all

 $(u, v) \in [0, 1]^2$ . The survival function of *C* is denoted by  $\overline{C}(u, v) = \Pr(U > u, V > v)$ ,  $(u, v) \in [0, 1]^2$ , where  $(U, V) \sim C$ . The comonotonicity, counter-monotonicity and independence copulas are denoted by  $M(u, v) = \min(u, v)$ ,  $W(u, v) = \max(u + v - 1, 0)$  and  $\Pi(u, v) = uv$ ,  $(u, v) \in [0, 1]^2$ , respectively. By the Fréchet–Hoeffding inequalities, it holds that  $W \leq C \leq M$  for all  $C \in C_2$ .

Consider a group of transforms on  $C_2$ :

$$\mathcal{T} = \{\iota, v_1, v_2, v_1 \circ v_2, \pi, \pi \circ v_1, \pi \circ v_2, \pi \circ v_1 \circ v_2\},\$$

where  $\iota : C_2 \to C_2$  is the identity  $\iota(C) = C$ ;  $\nu_1 : C_2 \to C_2$  and  $\nu_2 : C_2 \to C_2$  are the partial reflections defined, respectively, by

$$v_1(C)(u, v) = v - C(1 - u, v), \quad v_2(C)(u, v) = u - C(u, 1 - v), \quad C \in C_2;$$

their composition is given by  $v_1 \circ v_2(C)(u, v) = u + v - 1 + C(1 - u, 1 - v)$ ; and  $\pi : C_2 \to C_2$  is the permutation  $\pi(C)(u, v) = C(v, u)$ , for  $(u, v) \in [0.1]^2$ . Let  $C_{\varphi} = \varphi(C)$  for  $\varphi \in \mathcal{T}$ , and denote by

$$(U_{\nu_1}, V_{\nu_1}) \stackrel{\text{a.s.}}{=} (1 - U, V) \sim C_{\nu_1}, \qquad (U_{\nu_2}, V_{\nu_2}) \stackrel{\text{a.s.}}{=} (U, 1 - V) \sim C_{\nu_2}, (U_{\pi}, V_{\pi}) \stackrel{\text{a.s.}}{=} (V, U) \sim C_{\pi}, \qquad (U_{\nu_1 \circ \nu_2}, V_{\nu_1 \circ \nu_2}) \stackrel{\text{a.s.}}{=} (1 - U, 1 - V) \sim C_{\nu_1 \circ \nu_2},$$

for  $(U, V) \sim C$ , where "a.s." stands for almost surely.

For any map  $\kappa : C_2 \to \mathbb{R}$ , we identify  $\kappa(C)$  with  $\kappa(U, V)$  for a random vector  $(U, V) \sim C$  defined on a fixed atomless probability space  $(\Omega, \mathcal{A}, Pr)$ . A map  $\kappa$  on  $C_2$  is called a measure of concordance if it satisfies the following seven axioms [14].

**Definition 1** (Axioms for measures of concordance). A map  $\kappa : C_2 \to \mathbb{R}$  is called a measure of concordance if it satisfies the following seven axioms:

- 1. *Domain*:  $\kappa(C)$  is defined for any  $C \in C_2$ ;
- 2. *Symmetry*:  $\kappa(C_{\pi}) = \kappa(C)$  for any  $C \in C_2$ ;
- 3. *Monotonicity*: If  $C \leq C'$  for  $C, C' \in C_2$ , then  $\kappa(C) \leq \kappa(C')$ ;
- 4. *Range*:  $-1 \le \kappa(C) \le 1$  for any  $C \in C_2$ ,  $\kappa(M) = 1$  and  $\kappa(W) = -1$ ;
- 5. *Independence*:  $\kappa(\Pi) = 0$ ;
- 6. *Change of sign*:  $\kappa(C_{\nu_1}) = \kappa(C_{\nu_2}) = -\kappa(C)$  for any  $C \in C_2$ ;
- 7. *Continuity*: Let  $C_n \in C_2$ ,  $n \in \mathbb{N}$ , and  $C \in C_2$  with  $C_n$  converging pointwise to C as  $n \to \infty$ . Then  $\lim_{n\to\infty} \kappa(C_n) = \kappa(C)$ .

Consider a class of maps on  $C_2$  written as  $\kappa_{g_1,g_2}(U,V) = \rho(g_1(U),g_2(V))$ ,  $(U,V) \sim C$ , for two left-continuous functions  $g_1, g_2 : (0,1) \rightarrow \mathbb{R}$ . [9] showed that for  $\kappa_{g_1,g_2}$  to be a measure of concordance, it must be the so-called *G*-transformed rank correlation defined as follows. For a univariate distribution function  $G : \mathbb{R} \rightarrow [0,1]$ , the quantile function of *G* is defined by

$$G^{-1}(p) = \inf\{x \in \mathbb{R} : G(x) \ge p\}, p \in (0, 1).$$

A multivariate distribution *H* on  $\mathbb{R}^d$  with finite first moment is called radially symmetric if  $X - \mu \stackrel{d}{=} \mu - X$  for some  $\mu \in \mathbb{R}^d$ , where  $X \sim H$  and  $\stackrel{d}{=}$  stands for equality in distribution.

**Definition 2** (*G*-transformed rank correlation). For a univariate distribution function  $G : \mathbb{R} \to [0, 1]$ , the *G*-transformed rank correlation of  $C \in C_2$  is defined by

$$\kappa_G(C) = \rho(G^{-1}(U), G^{-1}(V)), \quad (U, V) \sim C.$$

We call G concordance-inducing if it is nondegenerate, radially symmetric with finite second moment. The set of all concordance-inducing distributions is denoted by G.

The following proposition summarizes basic properties of  $\kappa_G$ ; see [7].

**Proposition 1** (Basic properties of  $\kappa_G$ ). For any  $G \in G$ , the *G*-transformed rank correlation  $\kappa_G$  satisfies the following properties:

- 1.  $\kappa_G$  is a measure of concordance;
- 2.  $\kappa_G$  is invariant under location-scale transforms of G, that is,  $\kappa_{G_{\mu,\sigma}}(C) = \kappa_G(C)$  for all  $C \in C_2$ , where  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  and  $G_{\mu,\sigma}(x) = G((x \mu)/\sigma)$ ,  $x \in \mathbb{R}$ ;
- 3. For  $n \in \mathbb{N}$ , let  $C_1, \ldots, C_n \in C_2$  and  $\alpha_1, \ldots, \alpha_n$  be non-negative numbers such that  $\alpha_1 + \cdots + \alpha_n = 1$ . Then

$$\kappa_G\left(\sum_{i=1}^n \alpha_i C_i\right) = \sum_{i=1}^n \alpha_i \kappa_G(C_i).$$

The class of G-transformed rank correlations includes popular measures of concordance as special cases.

Example 1 (Examples of G-transformed rank correlations).

- 1) Spearman's rho:  $\kappa_G$  reduces to Spearman's rho  $\rho_S(U, V) = 12\mathbb{E}[UV] 3$  [17] if G is the standard uniform distribution Unif(0, 1).
- 2) Blomqvist's beta:  $\beta(C) = 4C(1/2, 1/2) 1$  is called Blomqvist's beta [2] (also known as median correlation), which is a *G*-transformed rank correlation with *G* being a symmetric Bernoulli distribution Bern(1/2) on {0, 1}.
- 3) van der Waerden's coefficient: When *G* is the standard normal distribution N(0, 1), then  $\kappa_G$  is known as van der Waerden's coefficient [16] (also known as normal score correlation and Gaussian rank correlation)  $\zeta(U, V) = \rho(\Phi^{-1}(U), \Phi^{-1}(V))$  where  $\Phi$  is the distribution function of N(0, 1).

## 3. Estimation of $\kappa_G$ and their comparison

In this section, we propose a novel approach for comparing *G*-transformed rank correlations to address the question which concordance-inducing distribution is most preferable to use. In the proposed approach, transformed rank correlations are compared in terms of the asymptotic variances of their canonical estimators, and one concordanceinducing distribution  $G \in \mathcal{G}$  is considered more preferable to another  $G' \in \mathcal{G}$  if the worst and best asymptotic variances of an estimator  $\hat{\kappa}_G$  of  $\kappa_G$  among a set of copulas  $\mathcal{D} \subseteq C_2$  are smaller than those of  $\kappa_{G'}$ .

#### 3.1. Canonical estimator of $\kappa_G$

Based on Proposition 1 Part 2, we first consider standardized concordance-inducing distributions *G* such that  $\mathbb{E}_G[X] = 0$  and  $\operatorname{Var}_G(X) = 1$  where  $X \sim G$ . Suppose that a data-generating i.i.d. process  $(U_1, V_1), (U_2, V_2), \ldots \stackrel{\text{iid}}{\sim} C$  on the probability space  $(\Omega, \mathcal{A}, \operatorname{Pr})$  is available to estimate a *G*-transformed rank correlation  $\kappa_G$ . This situation corresponds to the case when marginal distributions of the i.i.d. data are known. Although it may be unrealistic, this assumption is imposed throughout the paper to simplify the analysis. We then consider the following natural estimator of  $\kappa_G$ .

**Definition 3** (Canonical estimator of  $\kappa_G$ ). For  $G \in \mathcal{G}$ , the canonical estimator of  $\kappa_G$  is given by

$$\hat{\kappa}_G = \hat{\kappa}_G^{[n]}(C) = \frac{1}{n} \sum_{i=1}^n G^{-1}(U_i) G^{-1}(V_i), \quad (U_1, V_1), \dots, (U_n, V_n) \stackrel{\text{iid}}{\sim} C, \quad n \in \mathbb{N}$$

Although this simplified setting may be rarely the case, an asymptotic variance of this canonical estimator is obtained in a tractable form. To this end, let

$$\mathcal{G}_4 = \{ G \in \mathcal{G} : \mathbb{E}_G[X] = 0, \quad \operatorname{Var}_G(X) = 1, \quad \mathbb{E}_G[X^4] < \infty \text{ for } X \sim G \}.$$

If the fourth moment exists, the canonical estimator  $\hat{\kappa}_G$  satisfies the asymptotic normality by the classical central limit theorem: as  $n \to \infty$ ,

$$\sqrt{n} \{ \hat{\kappa}_G - \kappa_G(C) \} \stackrel{d}{\longrightarrow} \mathrm{N}(0, \sigma_G^2(C)),$$

where the asymptotic variance of  $\hat{\kappa}_G$  is given by

$$\sigma_G^2(C) = \operatorname{Var}(G^{-1}(U)G^{-1}(V)), \quad C \in \mathcal{C}_2$$

**Example 2** (Discrete concordance-inducing distributions). For  $m \in \mathbb{N}$ ,  $z = (z_1, \ldots, z_m) \in \mathbb{R}^m$  and  $p = (p_0, p_1, \ldots, p_m) \in \mathbb{R}^{m+1}$  such that  $0 = z_0 < z_1 < \cdots < z_m$ ,  $p_0 + 2\sum_{i=1}^m p_i = 1$  and  $\sum_{i=1}^m p_i z_i^2 = 1/2$ , consider a discrete distribution  $G_{m,z,p}$  supported on  $-z_m, \ldots, -z_1, z_0, z_1, \ldots, z_m$  with corresponding probabilities  $p_m, \ldots, p_1, p_0, p_1, \ldots, p_m \ge 0$ . Then  $G_{m,z,p}$  is a concordance-inducing distribution with mean zero and variance one. As a special case, Blomqvist's beta arises when m = 1,  $z_1 = 1$  and  $(p_0, p_1) = (0, 1/2)$ . Let  $p_+ = p_1 + \cdots + p_m$ ,  $I_{-i} = [p_+ - \sum_{j=1}^i p_j, p_+ - \sum_{j=1}^{i-1} p_j]$ ,  $I_0 = [p_+, p_+ + p_0]$  and  $I_i = [p_+ + p_0 + \sum_{j=1}^{i-1} p_j, p_+ + p_0 + \sum_{j=1}^i p_j]$  for  $i \in \{1, \ldots, m\}$ . Then

$$\kappa_{G_{m,z,p}}(C) = \mathbb{E}[G_{m,z,p}^{-1}(U)G_{m,z,p}^{-1}(V)] = \sum_{(i,j)\in\{-m,\dots,m\}} z_i z_j V_C(I_i \times I_j),$$

and

$$\sigma_{G_{m,z,p}}^{2}(C) = \operatorname{Var}(XY) = \mathbb{E}[(XY)^{2}] - (\mathbb{E}[XY])^{2}$$
$$= \sum_{(i,j)\in\{-m,\dots,m\}} z_{i}^{2} z_{j}^{2} V_{C}(I_{i} \times I_{j}) - \left(\sum_{(i,j)\in\{-m,\dots,m\}} z_{i} z_{j} V_{C}(I_{i} \times I_{j})\right)^{2},$$

where  $z_{-i} = -z_i$  for  $i \in \{1, ..., m\}$  and  $V_C(A), A \subseteq [0, 1]^2$ , is a volume of A measured by C.

For an underlying copula *C*, a concordance-inducing distribution *G* can be more preferable to another one *G'* in terms of the stability of statistical estimation if  $\sigma_G^2(C) \leq \sigma_{G'}^2(C)$ . Since this comparison is valid only for a specific copula *C*, we introduce a set of copulas  $\mathcal{D}$  as possible dependence structures which the analyst is interested in quantifying and comparing. Concordance-inducing distributions are then compared by the largest and smallest values of  $\sigma_G^2(C)$  over  $\mathcal{D}$ .

**Definition 4** (Best and worst asymptotic variances for  $\kappa_G$ ). For  $G \in \mathcal{G}_4$  and  $\mathcal{D} \subseteq C_2$ , the best and worst asymptotic variances are given by

$$\underline{\sigma}_{G}^{2}(\mathcal{D}) = \inf_{C \in \mathcal{D}} \sigma_{G}^{2}(C), \quad \overline{\sigma}_{G}^{2}(\mathcal{D}) = \sup_{C \in \mathcal{D}} \sigma_{G}^{2}(C), \tag{1}$$

respectively. If the infimum and supremum in (1) are attainable, the sets of their attaining copulas on D are denoted, respectively, by

$$\underline{C}_{G}(\mathcal{D}) = \operatorname*{argmin}_{C \in \mathcal{D}} \sigma_{G}^{2}(C), \quad \overline{C}_{G}(\mathcal{D}) = \operatorname*{argmax}_{C \in \mathcal{D}} \sigma_{G}^{2}(C).$$

Suppose that  $\mathcal{H} \subseteq \mathcal{G}_4$  represents the set of candidates among which the analyst chooses a concordance-inducing distribution. In terms of the stability of statistical estimation, we are interested in concordance-inducing distributions that minimize  $G \mapsto \underline{\sigma}_G^2(\mathcal{D})$  and/or  $G \mapsto \overline{\sigma}_G^2(\mathcal{D})$ .

**Definition 5** (Optimal best and worst asymptotic variances). For  $\mathcal{H} \subseteq \mathcal{G}_4$  and  $\mathcal{D} \subseteq \mathcal{C}_2$ , the optimal best and worst asymptotic variances on  $(\mathcal{H}, \mathcal{D})$  are defined, respectively, by

$$\underline{\sigma}^2_*(\mathcal{H}, \mathcal{D}) = \inf_{G \in \mathcal{H}} \underline{\sigma}^2_G(\mathcal{D}), \quad \overline{\sigma}^2_*(\mathcal{H}, \mathcal{D}) = \inf_{G \in \mathcal{H}} \overline{\sigma}^2_G(\mathcal{D}).$$
(2)

If the infima in (2) are attainable, the sets of their attaining concordance-inducing distributions are denoted, respectively, by

$$\underline{G}_*(\mathcal{H}, \mathcal{D}) = \operatorname*{argmin}_{G \in \mathcal{H}} \underline{\sigma}_G^2(\mathcal{D}), \quad \overline{G}_*(\mathcal{H}, \mathcal{D}) = \operatorname*{argmin}_{G \in \mathcal{H}} \overline{\sigma}_G^2(\mathcal{D}).$$

Finally, the set of optimal concordance-inducing distributions on  $(\mathcal{H}, \mathcal{D})$  is given by

$$G_*(\mathcal{H}, \mathcal{D}) = G_*(\mathcal{H}, \mathcal{D}) \cap \overline{G}_*(\mathcal{H}, \mathcal{D}).$$

Regarding the attainability in (1) and (2), we will see in Section 4 that the attaining elements in  $\mathcal{H}$  and  $\mathcal{D}$  can be described explicitly under certain assumptions on  $\mathcal{H}$  and  $\mathcal{D}$ .

The comparison of optimal best and worst asymptotic variances leads to the preference order among concordanceinducing distributions as follows.

**Definition 6** (Preference of concordance-inducing distributions). We say that  $G \in \mathcal{G}_4$  (or  $\kappa_G$ ) is more preferable to  $G' \in \mathcal{G}_4$  (or  $\kappa_{G'}$ ) on  $\mathcal{D} \subseteq \mathcal{C}_2$ , denoted by  $G' \leq_{\mathcal{D}} G$  (or  $\kappa_{G'} \leq_{\mathcal{D}} \kappa_G$ ), if

$$\underline{\sigma}_{G}^{2}(\mathcal{D}) \leq \underline{\sigma}_{G'}^{2}(\mathcal{D}), \quad \overline{\sigma}_{G}^{2}(\mathcal{D}) \leq \overline{\sigma}_{G'}^{2}(\mathcal{D}).$$

By definition, the preference order  $G' \leq_{\mathcal{D}} G$  is a partial order except some restricted cases of  $\mathcal{D}$  as seen in Corollary 3 below. If  $G \in G_*(\mathcal{H}, \mathcal{D})$ , then G may be considered as the most preferable choice among  $\mathcal{H} \subseteq \mathcal{G}_4$  to accurately estimate  $\kappa_G$  if the analyst believes that  $\mathcal{D}$  is the set of underlying copulas on which she wants to quantify and compare dependence.

Other than  $\mathcal{H} = \mathcal{G}_4$ , one may be interested, for example, in  $\mathcal{H} = \mathcal{G}_4^c$  where  $\mathcal{G}_4^c$  is the set of continuous concordanceinducing distributions in  $\mathcal{G}_4$ , and in  $\mathcal{H} = \mathcal{G}_4^b$  where  $\mathcal{G}_4^b$  is the set of concordance-inducing distributions in  $\mathcal{G}_4$  with bounded supports. Note that one-sided distributions  $X \sim G$  such that  $essup(X) = \infty$  and  $essinf(X) < \infty$ , or  $esssup(X) < \infty$  and  $essinf(X) = -\infty$ , cannot be concordance-inducing since they cannot be radially symmetric. Therefore,  $\mathcal{G}_4 \setminus \mathcal{G}_4^b$  is a set of concordance-inducing distributions supported on  $\mathbb{R}$ .

## 3.2. Properties of the asymptotic variance

We study properties of the asymptotic variance and its optimal best and worst counterparts.

**Proposition 2** (Basic properties of  $\sigma_G^2(C)$ ). Let  $G \in \mathcal{G}_4$ . Then the map  $C \mapsto \sigma_G(C)$  satisfies the following properties:

1. Covariance formula: For  $(U, V) \sim C \in C_2$  and  $(X, Y) = (G^{-1}(U), G^{-1}(V))$ , we have that

$$\sigma_G^2(C) = \text{Cov}(X^2, Y^2) + 1 - \text{Cov}(X, Y)^2;$$
(3)

- 2. Values at fundamental copulas:  $\sigma_G^2(\Pi) = 1$  and  $\sigma_G^2(M) = \sigma_G^2(W) = \operatorname{Var}_G(X^2)$ ; 3. Bounds:  $0 \le \sigma_G^2(C) \le 1 + \operatorname{Var}_G(X^2)$  for  $X \sim G$ . In particular,  $\sigma_G^2((M+W)/2) = 1 + \operatorname{Var}_G(X^2)$ ;
- 4. Boundedness:  $\sigma_G^2(C) < \infty$  for all  $C \in C_2$ ; 5. Reflection invariance:  $\sigma_G^2(C) = \sigma_G^2(C_{\nu_1}) = \sigma_G^2(C_{\nu_2}) = \sigma_G^2(C_{\nu_1 \circ \nu_2})$ .

**Proposition 3** (Concavity of  $\sigma_G^2(C)$ ). For  $G \in \mathcal{G}_4$ , the map  $C \mapsto \sigma_G^2(C)$  is concave with respect to convex combinations of copulas. In particular, for  $k \in [-1, 1]$ , the map  $C \mapsto \sigma_G^2(C)$  is linear on  $C_G(k) = \{C \in C_2 : \kappa_G(C) = k\}$ .

Concavity of  $C \mapsto \sigma_G^2(C)$  can be helpful to simplify the problem of deriving  $\underline{\sigma}_G^2(\mathcal{D})$  in some cases.

**Corollary 1** ( $\underline{\sigma}_G^2(C)$ ) over convex combinations of copulas). For  $L \in \mathbb{N}$  and  $C_1, \ldots, C_L \in C_2$ , let

$$\mathcal{D} = \operatorname{conv}(C_1, \ldots, C_L) \subseteq C_2$$

be the set of convex combinations of  $C_1, \ldots, C_L$ . Then

$$\underline{\sigma}_{G}^{2}(\mathcal{D}) = \min_{l=1,\dots,L} \sigma_{G}^{2}(C_{l}), \quad \underline{C}_{G}(\mathcal{D}) \supseteq \operatorname*{argmin}_{l=1,\dots,L} \sigma_{G}^{2}(C_{l}).$$

#### 3.3. Optimal location shift of G

Although  $\kappa_G$  is invariant under location-scale transforms of G by Proposition 1 Part 2, the asymptotic variance  $\sigma_G^2(C)$  of its canonical estimator  $\hat{k}_G$  may not be the case. To see this, let  $G_0 \in \mathcal{G}_4$  be a standardized concordanceinducing distribution with mean zero and variance one, and let  $G_{\mu,\sigma}(x) = G_0((x-\mu)/\sigma)$  be the corresponding concordance-inducing distribution of the same type as  $G_0$  but with mean  $\mu \in \mathbb{R}$  and variance  $\sigma^2 > 0$ . Since

$$\kappa_{G_{\mu,\sigma}}(C) = \rho(G_{\mu,\sigma}^{-1}(U), G_{\mu,\sigma}^{-1}(V)) = \frac{\mathbb{E}\left[G_{\mu,\sigma}^{-1}(U)G_{\mu,\sigma}^{-1}(V)\right] - \mu^2}{\sigma^2}, \quad (U,V) \sim C,$$

for known  $\mu$  and  $\sigma$ , a canonical estimator of  $\kappa_{G_{\mu,\sigma}}$  can be given by

$$\hat{\kappa}_{G_{\mu,\sigma}} = \hat{\kappa}_{G_{\mu,\sigma}}^{[n]}(C) = \frac{1}{n} \sum_{i=1}^{n} \frac{G_{\mu,\sigma}^{-1}(U_i)G_{\mu,\sigma}^{-1}(V_i)}{\sigma^2} - \left(\frac{\mu}{\sigma}\right)^2,$$

for  $(U_1, V_1), (U_2, V_2), \dots \stackrel{\text{iid}}{\sim} C$ , which reduces to (3) when  $\mu = 0$  and  $\sigma^2 = 1$ . By the CLT, asymptotic normality follows for  $\hat{\kappa}_{G_{\mu\sigma}}$  with the asymptotic variance given by

$$\sigma_{G_{\mu,\sigma}}^2(C) = \operatorname{Var}\left(\frac{G_{\mu,\sigma}^{-1}(U)G_{\mu,\sigma}^{-1}(V)}{\sigma^2}\right)$$

Since  $G_{\mu,\sigma}^{-1}(U)/\sigma = G_{\mu/\sigma,1}^{-1}(U)$  and  $G_{\mu,\sigma}^{-1}(V)/\sigma = G_{\mu/\sigma,1}^{-1}(V)$ , one can assume that  $\sigma = 1$  without changing the asymptotic variance  $\sigma_{G_{\mu,\sigma}}^2(C)$ , that is,  $\sigma_G^2(C)$  is invariant under scale transforms of *G*. On the other hand,  $\sigma_G^2(C)$  changes under location transforms of *G* since shifting  $G^{-1}$  by  $\mu \in \mathbb{R}$  leads to the asymptotic variance  $\operatorname{Var}((X + \mu)(Y + \mu)) = \operatorname{Var}(XY + \mu(X + Y))$  for  $X = G^{-1}(U)$  and  $Y = G^{-1}(V)$ , which is in general not equal to  $\operatorname{Var}(XY)$ .

Since the canonical estimator  $\hat{\kappa}_{G_{\mu,\sigma}}$  estimates the same quantity  $\kappa_{G_0}$  regardless of the mean  $\mu$  and variance  $\sigma^2$  of G, a natural choice of  $(\mu, \sigma)$  is such that  $\sigma = 1$  and  $\mu$  minimizes the asymptotic variance  $\sigma^2_{G_{\mu,1}}(C)$ . For a fixed concordance-inducing distribution  $G_0 \in \mathcal{G}_4$  with mean zero and variance one, denote by  $G_{\mu}(x) = G_0(x-\mu)$  the concordance-inducing distribution of the same type as  $G_0$  but with mean  $\mu \in \mathbb{R}$ . For  $X = X_0 + \mu \sim G_{\mu}$  and  $Y = Y_0 + \mu \sim G_{\mu}$  with  $X_0 = G_0^{-1}(U)$  and  $Y_0 = G_0^{-1}(V)$ , the asymptotic variance is given by

$$\sigma_{G_{\mu}}^{2}(C) = \operatorname{Var}(XY) = \operatorname{Var}((X_{0} + \mu)(Y_{0} + \mu)) = \operatorname{Var}(X_{0}Y_{0} + \mu(X_{0} + Y_{0}))$$
$$= \operatorname{Var}(X_{0}Y_{0}) + 2\mu \operatorname{Cov}(X_{0}Y_{0}, X_{0} + Y_{0}) + \mu^{2} \operatorname{Var}(X_{0} + Y_{0}).$$

Therefore the desired  $\mu$  is given as follows.

**Definition 7** (Optimal shift of  $G_0$ ). For  $G_0 \in \mathcal{G}_4$  and  $C \in C_2$ , the minimizer of  $\mu \mapsto \sigma^2_{G_{\mu}}(C)$  is called the optimal (location) shift of  $G_0$  under C, and is given by

$$\mu_* = \mu_*(G_0, C) = \begin{cases} -\frac{\operatorname{Cov}(X_0Y_0, X_0 + Y_0)}{\operatorname{Var}(X_0 + Y_0)}, & \text{if } \operatorname{Var}(X_0 + Y_0) > 0, \\ 0, & \text{if } \operatorname{Var}(X_0 + Y_0) = 0, \end{cases}$$

where  $(U, V) \sim C$  and  $(X_0, Y_0) = (G_0^{-1}(U), G_0^{-1}(V))$ . The optimal asymptotic variance is then given by

$$\sigma_{G_{\mu_*}}^2(C) = \begin{cases} \operatorname{Var}(X_0 Y_0) - \frac{\operatorname{Cov}(X_0 Y_0, X_0 + Y_0)^2}{\operatorname{Var}(X_0 + Y_0)}, & \text{if } \operatorname{Var}(X_0 + Y_0) > 0, \\ \operatorname{Var}(X_0 Y_0), & \text{if } \operatorname{Var}(X_0 + Y_0) = 0. \end{cases}$$

For  $G_0 \in \mathcal{G}_4$ , the degenerate case  $Var(X_0 + Y_0) = 0$  occurs if and only if  $X_0 + Y_0 \stackrel{a.s.}{=} 0$ . In this case, we have that

$$\sigma_{G_{\mu}}^{2}(C) = \operatorname{Var}(X_{0}Y_{0}) + 2\mu \operatorname{Cov}(X_{0}Y_{0}, 0) + \mu^{2} \operatorname{Var}(0) = \operatorname{Var}(X_{0}Y_{0}) = \sigma_{G_{0}}^{2}(C)$$

for every  $\mu \in \mathbb{R}$ .

The following proposition states that  $\mu_* = 0$  for a certain class of copulas.

**Proposition 4** (Sufficient condition for  $\mu_* = 0$ ). Let  $C \in C_2$  be a copula and  $G_0 \in \mathcal{G}_4$  be a concordance-inducing distribution with mean zero and variance one. Then  $\mu_*(G_0, C) = 0$  holds if C is radially symmetric  $C = C_{\nu_1 \circ \nu_2}$ , that is,  $(U, V) \stackrel{d}{=} (1 - U, 1 - V)$  for  $(U, V) \sim C$ .

By Proposition 4, a location shift of  $G_0$  does not change the asymptotic variance  $\sigma_{G_0}^2(C)$  when *C* is, for example, *M*, *W*,  $\Pi$ , a Gaussian copula, *t* copula or their mixtures. On the other hand, shifting  $G_0$  may improve  $\sigma_{G_0}^2(C)$  if *C* is, for example, a Clayton or Gumbel copula since they are not radially symmetric in general. The next proposition

states that the optimal asymptotic variance can be obtained analytically when  $(X, Y) = (G^{-1}(U), G^{-1}(V))$  is a normal variance mixture, that is,

$$(X,Y) = (\mu_1,\mu_2) + \sqrt{W}(Z_1,Z_2), \quad (\mu_1,\mu_2) \in \mathbb{R}^2, \quad (Z_1,Z_2) \sim \mathcal{N}_2(\mathbf{0}_2,\Sigma), \tag{4}$$

where  $\Sigma$  is a 2-dimensional square positive definite matrix and  $W \ge 0$  is a non-negative random variable independent of  $(Z_1, Z_2)$  and such that  $\Pr(W = 0) < 1$ .

**Proposition 5** ( $\sigma_{G_{\mu_*}}^2(C)$ ) for normal variance mixture). For  $G \in \mathcal{G}_4$  and  $(U, V) \sim C \in C_2$ , suppose that  $(X, Y) = (G^{-1}(U), G^{-1}(V))$  is a normal variance mixture specified by (4). Then the optimal asymptotic variance is given by

$$\sigma_{G_{\mu^*}}^2(C) = \left(2\frac{\mathbb{E}[W^2]}{\mathbb{E}[W]^2} - 1\right)\rho^2(X, Y) + \frac{\mathbb{E}[W^2]}{\mathbb{E}[W]^2}.$$
(5)

By (5), the function  $\rho \mapsto \sigma_{G_{\mu_*}}^2(C)$  is convex since  $\mathbb{E}[W^2] \ge \mathbb{E}[W]^2$ . For normal distributions, W = 1 and thus  $\sigma_{G_{\mu_*}}^2(C) = \rho^2(X, Y) + 1$ . Finally, as we will see in Section 6, the variance  $\operatorname{Var}(X^2)$ ,  $X \sim G$ , is observed to affect convexity or concavity of  $\rho \mapsto \sigma_{G_{\mu_*}}^2(C)$ . For the case of normal variance mixtures, we have that

$$\sigma_{G_{\mu^*}}^2(C) = \left(\operatorname{Var}(X^2) - \frac{\mathbb{E}[W^2]}{\mathbb{E}[W]^2}\right) \rho^2(X, Y) + \frac{\mathbb{E}[W^2]}{\mathbb{E}[W]^2},\tag{6}$$

since

$$\operatorname{Var}(X^{2}) = \mathbb{E}[W^{2}Z_{1}^{4}] - (\mathbb{E}[WZ_{1}^{2}])^{2} = \mathbb{E}[W^{2}] \left(\frac{3}{\mathbb{E}[W]^{2}}\right) - \mathbb{E}[W]^{2} \left(\frac{1}{\mathbb{E}[W]^{2}}\right) = 3\frac{\mathbb{E}[W^{2}]}{\mathbb{E}[W]^{2}} - 1$$

# 4. Optimal concordance-inducing distributions

In this section we investigate optimal best and worst asymptotic variances and their attaining concordanceinducing distributions for certain choices of  $\mathcal{D} \subseteq C_2$ .

#### 4.1. Asymptotic variance for fundamental and Fréchet copulas

We first consider the case when  $\mathcal{D} \subset C_2$  is a set of fundamental copulas M,  $\Pi$  and W, or their mixtures since these copulas play important roles in the discussion of the best and worst asymptotic variances. By radial symmetry of these copulas, the optimal shift  $\mu_*$  is zero and thus it suffices to consider standardized concordance-inducing distributions in  $\mathcal{G}_4$ .

Definition 8 (Fréchet copula). A bivariate Fréchet copula is defined by

$$C_{\boldsymbol{p}}^{\mathrm{F}} = p_M M + p_\Pi \Pi + p_W W, \quad \boldsymbol{p} = (p_M, p_\Pi, p_W) \in \Delta_3,$$

where  $\Delta_3 = \{(p_1, p_2, p_3) \in \mathbb{R}^3 : p_1, p_2, p_3 \ge 0, p_1 + p_2 + p_3 = 1\}$  is the standard unit simplex on  $\mathbb{R}^3$ . The set of all Fréchet copulas is denoted by  $C^F = \{C_p^F : p \in \Delta_3\}$ .

In addition to their financial applications, Fréchet copulas can be used to approximate bivariate copulas; see [19]. Moreover, for any  $G \in \mathcal{G}_4$ , the transformed rank correlation  $\kappa_G$  can take any value in [-1, 1] since, by Proposition 1 Part 3, it holds that

$$\kappa_G(C_p^F) = p_M \kappa_G(M) + p_\Pi \kappa_G(\Pi) + p_W \kappa_G(W) = p_M - p_W \in [-1, 1].$$
(7)

The following proposition is an immediate consequence from Proposition 2 Part 2.

**Proposition 6** (Optimal asymptotic variances for fundamental copulas). Let  $\mathcal{H} \subseteq \mathcal{G}_4$ .

1.  $\underline{\sigma}^2_*(\mathcal{H}, \{\Pi\}) = \overline{\sigma}^2_*(\mathcal{H}, \{\Pi\}) = 1 \text{ and } \underline{G}_*(\mathcal{H}, \{\Pi\}) = \overline{G}_*(\mathcal{H}, \{\Pi\}) = \mathcal{H}.$ 

2. Suppose  $\mathcal{D} = \{M\}, \{W\} \text{ or } \{M, W\}$ . Then

$$\underline{\sigma}^2_*(\mathcal{H},\mathcal{D}) = \overline{\sigma}^2_*(\mathcal{H},\mathcal{D}) = \inf_{G \in \mathcal{H}} \operatorname{Var}_G(X^2).$$

If this infimum is attainable, then

$$\underline{G}_*(\mathcal{H}, \mathcal{D}) = \overline{G}_*(\mathcal{H}, \mathcal{D}) = \operatorname*{argmin}_{G \in \mathcal{H}} \operatorname{Var}_G(X^2).$$

3. Suppose that  $\mathcal{D} = \{\Pi, M, W\}$ . Then

$$\underline{\sigma}^2_*(\mathcal{H}, \{M, \Pi, W\}) = 1 \land \inf_{G \in \mathcal{H}} \operatorname{Var}_G(X^2), \quad \overline{\sigma}^2_*(\mathcal{H}, \{M, \Pi, W\}) = 1 \lor \inf_{G \in \mathcal{H}} \operatorname{Var}_G(X^2).$$

If the infima above are attainable, then

$$\underline{G}_{*}(\mathcal{H}, \{M, \Pi, W\}) = \begin{cases} \operatorname{argmin} \operatorname{Var}_{G}(X^{2}), & \text{if } \min_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}) < 1, \\ \mathcal{H}, & \text{if } \min_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}) \geq 1, \end{cases}$$
$$\overline{G}_{*}(\mathcal{H}, \{M, \Pi, W\}) = \begin{cases} \mathcal{H}, & \text{if } \min_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}) < 1, \\ \operatorname{argmin}_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}), & \text{if } \min_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}) \geq 1. \end{cases}$$

The next proposition provides the best and worst asymptotic variances and their attainers when  $\mathcal{D} = C^{F}$ .

**Proposition 7** (Best and worst asymptotic variances for Fréchet copulas). For a concordance-inducing distribution  $G \in \mathcal{G}_4$ , the best and worst asymptotic variances on  $C^F$  are given by

$$\underline{\sigma}_{G}^{2}(C^{\mathrm{F}}) = 1 \wedge \operatorname{Var}_{G}(X^{2}), \quad \overline{\sigma}_{G}^{2}(C^{\mathrm{F}}) = 1 + \operatorname{Var}_{G}(X^{2})$$

with the sets of attaining copulas given by

$$\begin{split} \underline{C}_{G}(C^{\mathrm{F}}) &= \begin{cases} \{M, W\}, & \text{if } 0 \leq \mathrm{Var}_{G}(X^{2}) < 1, \\ \{M, \Pi, W\}, & \text{if } \mathrm{Var}_{G}(X^{2}) = 1, \\ \{\Pi\}, & \text{if } \mathrm{Var}_{G}(X^{2}) > 1, \end{cases} \\ \overline{C}_{G}(C^{\mathrm{F}}) &= \begin{cases} \left\{\frac{M+W}{2}\right\}, & \text{if } \mathrm{Var}_{G}(X^{2}) > 0, \\ \left\{p\frac{M+W}{2} + (1-p)\Pi : p \in [0,1]\right\}, & \text{if } \mathrm{Var}_{G}(X^{2}) = 0. \end{cases} \end{split}$$

Note that the result in Proposition 7 is consistent with Corollary 1. In the proof of Proposition 7, although  $(p_M, p_W) = (1/2, 1/2)$  is the unique point attaining the maximum v + 1 of f when v > 0, f takes the value v at the points  $(p_M, p_W) = (1, 0)$  and (0, 1), and is greater than v on  $\{(p_M, p_W) \in [0, 1]^2 : p_M + p_W = 1\}$ . Therefore, if  $v = \operatorname{Var}_G(X^2)$  is sufficiently large, the asymptotic variance  $\sigma_G^2(C)$  takes large values in  $[\operatorname{Var}_G(X^2), \operatorname{Var}_G(X^2) + 1]$  if C = pM + (1 - p)W for  $p \in [0, 1]$ .

**Remark 1** (Restrictions of  $C^{F}$ ). For a concordance-inducing distribution  $G \in \mathcal{G}_{4}$ , consider the set of Fréchet copulas such that its transformed rank correlation  $\kappa_{G}$  takes values in  $[\underline{k}, \overline{k}]$  for  $-1 \leq \underline{k} \leq \overline{k} \leq 1$ , that is,

$$C^{\mathrm{F}}_{k,\overline{k}}(G) = \{ C \in C^{\mathrm{F}} : \underline{k} \le \kappa_G(C) \le \overline{k} \}.$$

By (7), the restriction  $\underline{k} \leq \kappa_G(C) \leq \overline{k}$  reduces to  $\underline{k} \leq p_M - p_W \leq \overline{k}$  and thus  $C_{\underline{k},\overline{k}}^F(G)$  does not depend on the choice of *G*. Consequently, the maximum and minimum of the asymptotic variance  $\sigma_G^2(C)$  on  $C_{\underline{k},\overline{k}}^F(G)$  can be found by calculating max  $f(p_M, p_W)$  and min  $f(p_M, p_W)$  subject to the constraints

$$\{(p_M, p_W) \in \mathbb{R}^2 : 0 \le p_M, p_W, p_M + p_W \le 1 \text{ and } \underline{k} \le p_M - p_W \le k\}$$

This maximum and minimum always exist since  $(p_M, p_W) \mapsto f(p_M, p_W)$  is bounded, concave and the feasible set is compact in  $\mathbb{R}^2$ .

Proposition 7 immediately leads to the optimal best and worst asymptotic variances on  $\mathcal{D} = C^{F}$  as stated in the following corollary.

**Corollary 2** (Optimal best and worst asymptotic variances for Fréchet copulas). For  $\mathcal{H} \subseteq \mathcal{G}_4$ , the optimal best and worst asymptotic variances are given by

$$\underline{\sigma}^2_*(\mathcal{H}, C^{\mathsf{F}}) = 1 \land \inf_{G \in \mathcal{H}} \operatorname{Var}_G(X^2), \quad \overline{\sigma}^2_*(\mathcal{H}, C^{\mathsf{F}}) = 1 + \inf_{G \in \mathcal{H}} \operatorname{Var}_G(X^2).$$

If the infima above are attainable, then the sets of attaining concordance-inducing distributions are given, respectively, by

$$\underline{G}_{*}(\mathcal{H}, C^{\mathrm{F}}) = \begin{cases} \operatorname{argmin}_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}), & \text{if } \min_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}) < 1, \\ \mathcal{H}, & \text{if } \min_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}) \geq 1, \end{cases}$$
$$\overline{G}_{*}(\mathcal{H}, C^{\mathrm{F}}) = \operatorname{argmin}_{G \in \mathcal{H}} \operatorname{Var}_{G}(X^{2}).$$

Compared with the optimal best and worst asymptotic variances from Proposition 6 Part 3, the lower bound  $\sigma_*^2(\mathcal{H}, \mathcal{D})$  obtained in Proposition 2 remains unchanged whereas the upper bound  $\overline{\sigma}_*^2(\mathcal{H}, \mathcal{D})$  increases since the attaining copulas  $p(M + W)/2 + (1 - p)\Pi$ ,  $p \in [0, 1]$ , are not included in the set  $\mathcal{D}$  in Proposition 6. Nevertheless, the best and worst asymptotic variances are the functions of  $\operatorname{Var}_G(X^2)$  when  $\mathcal{D}$  is a set of fundamental or Fréchet copulas, and thus we have the following result.

**Corollary 3** (Optimal concordance-inducing distributions on fundamental or Fréchet copulas). Suppose that  $\mathcal{D}$  is a set of fundamental or Fréchet copulas, that is,  $\mathcal{D} = \{\Pi\}, \{M\}, \{W\}, \{M, W\}, \{M, \Pi, W\}$  or  $C^F$ . Then

- 1.  $\leq_{\mathcal{D}}$  is a total order;
- 2.  $G' \leq_{\mathcal{D}} G$  if  $\operatorname{Var}_G(X^2) \leq \operatorname{Var}_{G'}(X^2)$ ;
- 3.  $G_*(\mathcal{H}, \mathcal{D}) = \operatorname{argmin} \operatorname{Var}_G(X^2)$  provided that  $\inf_{G \in \mathcal{H}} \operatorname{Var}_G(X^2)$  is attainable.

Corollary 3 states that concordance-inducing distributions having a smaller variance of  $X^2$  for  $X \sim G$  are more preferable in terms of best and worst asymptotic variances when  $\mathcal{D}$  is a set of fundamental or Fréchet copulas. As a consequence, heavy-tailed concordance-inducing distributions, such as a Student *t* distribution with degrees of freedom  $4 < v < \infty$ , are not recommendable choices at least in terms of accuracy of statistical estimation. In particular, popular measures of concordance introduced in Example 1 are ordered as follows.

**Corollary 4** (Preference orders for  $\rho_S$ ,  $\beta$  and  $\zeta$ ). Suppose that  $\mathcal{D}$  is a set of fundamental or Fréchet copulas. Then  $\zeta \leq_{\mathcal{D}} \rho_S \leq_{\mathcal{D}} \beta$ .

# 4.2. Optimality of Blomqvist's beta

In this section, we show that Blomqvist's beta is an optimal *G*-transformed rank correlation under some conditions on  $\mathcal{D} \in C_2$ . For  $G \in \mathcal{G}_4$ , we identify  $G \in \mathcal{G}_4$  with  $\kappa_G$ , and thus we allow  $\sigma_G^2$  to be written as  $\sigma_\beta^2$  for the standardized symmetric Bernoulli distribution  $G \in \mathcal{G}_4$ .

Definition 9 (Balancedness of copulas). Let

$$p(C) = C(1/2, 1/2) + \overline{C}(1/2, 1/2), \quad C \in C_2.$$

A copula  $C \in C_2$  is called

- (i) balanced if p(C) = 1/2,
- (ii) imbalanced if  $p(C) \neq 1/2$ ,
- (iii) totally positively imbalanced (TPI) if p(C) = 1,
- (iv) totally negatively imbalanced (TNI) if p(C) = 0.

It is straightforward to check that  $\Pi$  and (M + W)/2 are balanced, M is TPI and W is TNI.

**Proposition 8** (Asymptotic variance of Blomqvist's beta). Let  $C \in C_2$ . Then the following properties hold for  $\beta$ .

- 1.  $\mu_*(\beta, C) = 0;$
- 2.  $\sigma_{\beta}^{2}(C) = 4p(C)(1 p(C)) = 1 \beta^{2}(C);$
- 3.  $0 \le \sigma_{\beta}^2(C) \le 1;$
- 4.  $\sigma_{\beta}^{2}(C) = 0$  if and only if C is a TPI or TNI copula;
- 5.  $\sigma_{\beta}^{2}(C) = 1$  if and only if C is balanced.

**Remark 2** (Asymptotic variance of  $\beta$  for elliptical copulas). Blomqvist's beta admits an explicit form  $\beta(C) = \frac{2}{\pi} \arcsin(\rho)$  when *C* is an elliptical copula with correlation parameter  $\rho \in [-1, 1]$ . Therefore, by Proposition 8 Part 2, we have that  $\sigma_{\beta}^2(C) = 1 - (\frac{2}{\pi} \arcsin(\rho))^2$ , which coincides with the result derived in [15, Proposition 9].

Next we prove the optimality of Blomqvist's beta under certain conditions on  $\mathcal{D}$ .

**Proposition 9** (Optimality of Blomqvist's beta). Let  $\mathcal{D} \subseteq C_2$  and  $\beta \in \mathcal{H} \subseteq \mathcal{G}_4$ .

- 1. If  $C^* \in \mathcal{D}$  for some TPI or TNI copula  $C_*$ , then  $\beta \in \underline{G}_*(\mathcal{H}, \mathcal{D})$ .
- 2. If  $\Pi \in \mathcal{D}$ , then  $\beta \in \overline{G}_*(\mathcal{H}, \mathcal{D})$ .
- 3. If  $\Pi$ ,  $C_* \in \mathcal{D}$  for some TPI or TNI copula  $C_*$ , then  $\beta \in G_*(\mathcal{H}, \mathcal{D})$ .

Statements 1, 2 and 3 remain valid if  $\sigma_G^2(C)$ ,  $G \in \mathcal{G}_4$ , in Definitions 4, 5 and 6 is replaced by the optimally shifted asymptotic variance  $\sigma_{\mathcal{G}_{uc}}^2(C)$ .

Proposition 9 states that Blomqvist's beta is an optimal choice of *G*-transformed rank correlation for possibly typical choices of  $\mathcal{D}$ , such as  $C_2, C_2^{\geq} = \{C \in C_2 : C \geq \Pi\}$  or  $C_2^{\leq} = \{C \in C_2 : C \leq \Pi\}$ .

# 4.3. Uniqueness of the optimality of $\beta$

In this section we investigate whether Blomqvist's beta is the unique optimal *G*-transformed rank correlation, that is, whether  $G_*(\mathcal{H}, \mathcal{D}) = \{\beta\}$ . The next proposition states that this uniqueness holds under some condition on  $\mathcal{D}$ .

**Proposition 10** (Uniqueness of  $\beta$  for  $\overline{G}_*(\mathcal{H}, \mathcal{D})$  and  $G_*(\mathcal{H}, \mathcal{D})$ ). Let  $\mathcal{H} \subseteq \mathcal{G}_4$  and  $\mathcal{D} \subseteq \mathcal{C}_2$  be such that

- 1.  $\beta \in \mathcal{H}$ ;
- 2.  $\Pi$ ,  $(M + W)/2 \in \mathcal{D}$  and  $\mathcal{D}$  contains a TPI or TNI copula.

Then  $\overline{G}_*(\mathcal{H}, \mathcal{D}) = \{\beta\}$  and thus  $G_*(\mathcal{H}, \mathcal{D}) = \{\beta\}$ . The statement remains valid if  $\sigma_G^2(C), G \in \mathcal{G}_4$ , in Definitions 4, 5 and 6 is replaced by the optimally shifted asymptotic variance  $\sigma_{G_4}^2(C)$ .

Proposition 10 does not address whether  $\underline{G}_*(\mathcal{H}, \mathcal{D}) = \{\beta\}$ . This, however, is rarely the case as we will see in what follows.

For given  $\mathcal{H} \subseteq \mathcal{G}_4$  and  $\mathcal{D} \subseteq \mathcal{C}_2$ , assume that  $\beta \in \mathcal{H}$  and that  $\mathcal{D}$  contains at least one TPI or TNI copula. Then the following equivalence relations hold by Proposition 8 Part 4:

$$G \in \underline{G}_{*}(\mathcal{H}, \mathcal{D}) \quad \Leftrightarrow \quad \text{there exists } C \in \mathcal{D} \text{ such that } \sigma_{G}^{2}(C) = 0$$
$$\Leftrightarrow \quad G^{-1}(U)G^{-1}(V) \stackrel{\text{a.s.}}{=} a \text{ for some } a \in \mathbb{R} \text{ and } (U, V) \sim C \in \mathcal{D}. \tag{8}$$

The next proposition provides necessary conditions on  $a \in \mathbb{R}$ ,  $G \in \mathcal{H}$  and  $C \in \mathcal{D}$  in (8).

**Proposition 11** (Necessary conditions on  $G \in \underline{G}_*(\mathcal{H}, \mathcal{D})$ ). For  $\mathcal{H} \subseteq \mathcal{G}_4$  and  $\mathcal{D} \subseteq C_2$ , suppose that  $\beta \in \mathcal{H}$  and that  $\mathcal{D}$  contains at least one TPI or TNI copula. If  $G \in \underline{G}_*(\mathcal{H}, \mathcal{D})$ , then  $C \in \mathcal{D}$  and  $a \in \mathbb{R}$  in (8) satisfy the following conditions:

1. If Pr(X = 0) > 0 for  $X \sim G$ , then a = 0 and  $Pr(X = 0) \ge 1/2$ ;

2. If Pr(X = 0) = 0, then  $a \neq 0$  and the copula *C* is either TPI or TNI with  $0 < a \le 1$  if *C* is TPI and  $-1 \le a < 0$  if *C* is TNI. Moreover, the conditional distribution function

$$G_+(x) = \Pr(X \le x \mid X \ge 0) = 2G(x) - 1, \quad x > 0$$

satisfies

$$\mathbb{E}_{G_{+}}[Z] \ge |a|^{1/2}, \quad Z \sim G_{+} \quad and \quad G_{+}(x) = 1 - G_{+}\left(\frac{|a|}{x}\right), \quad x > 0.$$
(9)

In particular, it holds that  $Pr(Z > |a|^{1/2}) = Pr(Z < |a|^{1/2})$  for  $Z \sim G_+$ .

By Proposition 11, not all  $G \in \mathcal{G}_4$  and  $C \in C_2$  can attain the optimal best asymptotic variance  $\sigma_G^2(C) = 0$ . The following examples show non-Bernoulli concordance-inducing distributions attaining this lower bound. Let  $M(n, \{J_i\}, \pi, w)$  denote a shuffle-of-M with n being the number of connected components in its support,  $\{J_i\} = \{J_1, \ldots, J_n\}$  being a finite partition of [0, 1] into n closed subintervals,  $\pi$  being a permutation of  $\{1, \ldots, n\}$  and  $w : \{1, \ldots, n\} \rightarrow \{-1, 1\}^n$  being a function indicating whether the strip  $J_i \times J_{\pi(i)}$  is flipped (w(i) = 1) or not (w(i) = -1); see [11, Section 3.2.3].

**Example 3** (Non-Bernoulli concordance-inducing distributions in  $G_{*}(\mathcal{G}_{4}, \mathcal{D})$ ).

- 1. The case when  $\Pr(X = 0) > 0$ : Let  $X \sim G$  be an equally weighted mixture of 0 and  $\operatorname{Unif}(-\sqrt{6}, \sqrt{6})$ . Then  $\mathbb{E}[X] = 0$ ,  $\operatorname{Var}(X) = 1$  and  $\mathbb{E}[X^4] < \infty$ , and thus  $G \in \mathcal{G}_4$ . This is Case 1 of Proposition 11 since  $\Pr(X = 0) = 1/2$ . Consider  $C_1 = M(4, \bigcup_{i=1}^4 [(i-1)/4, i/4], \{2, 1, 4, 3\}, \mathbf{1}_4)$ ,  $C_2 = M(4, \bigcup_{i=1}^4 [(i-1)/4, i/4], \{3, 4, 1, 2\}, \mathbf{1}_4)$  and  $C_3 = M(4, \bigcup_{i=1}^4 [(i-1)/4, i/4], \{2, 4, 1, 3\}, \mathbf{1}_4)$ , where  $\mathbf{1}_d = (1, \ldots, 1) \in \mathbb{R}^d$ . Then  $C_1$  is TPI,  $C_2$  is TNI and  $C_3$  is neither TPI nor TNI. Moreover, they satisfy  $\sigma_G^2(C_k) = 0$ , k = 1, 2, 3, since  $G^{-1}(U)G^{-1}(V) \stackrel{\text{a.s.}}{=} 0$  with  $(U, V) \sim C_k$  for k = 1, 2, 3.
- 2. The case when  $\Pr(X = 0) = 0$ : Let  $X \sim G$  be a discrete uniform distribution on the four points  $\{-a/b, -b, b, a/b\}$  where  $a = 1/2^{1/2}$  and  $b = (1-2^{-1/2})^{1/2}$  with  $b \approx 0.541$  and  $a/b \approx 1.307$ . Then it is straightforward to check that  $G \in \mathcal{G}_4$ . Define  $(X, Y) = (G^{-1}(U), G^{-1}(V))$  with  $(U, V) \sim C_4 = M(4, \bigcup_{i=1}^4 [(i-1)/4, i/4], \{2, 1, 4, 3\}, -\mathbf{1}_4)$ . Then (X, Y) = (-a/b, -b), (-b, -a/b), (b, a/b) and (a/b, b) are equiprobable, and thus  $\sigma_G^2(C_4) = 0$  since  $XY \stackrel{\text{s}}{=} a$ . This case belongs to Case 2 of Proposition 11 since  $C_4$  is TPI,  $0 < a \le 1$  and  $\mathbb{E}_{G_4}[Z] \approx 0.924 > 0.841 \approx \sqrt{a}$ .

# 5. Comparison of $\kappa_G$ and Kendall's tau

In Section 4.2, we showed that Blomqvist's beta is an optimal *G*-transformed rank correlation under some conditions on  $\mathcal{D} \subseteq C_2$ . In this section, we show that Kendall's tau, although it is not a transformed rank correlation, shares certain optimal structure of Blomqvist's beta.

Kendall's tau  $\tau : C_2 \to \mathbb{R}$  is defined by

$$\tau(C) = 4 \int_{[0,1]^2} C(u,v) \, \mathrm{d}C(u,v) - 1, \tag{10}$$

and is a measure of concordance; see [14]. Moreover, it is not a *G*-transformed rank correlation since  $\tau$  is not linear with respect to a mixture of copulas. Since  $\tau(C) = \rho(\mathbf{1}_{\{U > \tilde{U}\}}, \mathbf{1}_{\{V > \tilde{V}\}})$  where  $(U, V) \sim C$  and  $(\tilde{U}, \tilde{V}) \sim C$  is an independent copy of (U, V),  $\tau$  can also be written as

$$\tau(C) = \rho(g(U, \tilde{U}), g(V, \tilde{V})), \quad g(\ell, m) = \begin{cases} 1 & \text{if } \ell > m, \\ -1 & \text{if } \ell \le m, \end{cases}$$
(11)

by invariance of  $\rho$  under location-scale transforms. Assuming that the data-generating i.i.d. process from *C* is available, we consider the following estimator of  $\tau(C)$  to estimate a *G*-transformed rank correlation  $\kappa_G$  based on the representation (11).

**Definition 10** (Canonical estimator of  $\tau$ ). For  $C \in C_2$ , the canonical estimator of  $\tau$  is given by

$$\hat{\tau} = \hat{\tau}(C; n) = \frac{1}{n} \sum_{i=1}^{n} g(U_i, \tilde{U}_i) g(V_i, \tilde{V}_i), \quad (U_1, V_1), \dots, (U_n, V_n), (\tilde{U}_1, \tilde{V}_1), \dots, (\tilde{U}_n, \tilde{V}_n) \stackrel{\text{id}}{\sim} C, \quad n \in \mathbb{N}.$$

By the classical central limit theorem, the following asymptotic normality follows

$$\sqrt{n}\left\{\hat{\tau}-\tau(C)\right\}\overset{d}{\longrightarrow} \mathrm{N}(0,\sigma_{\tau}^{2}(C)), \quad n\to\infty,$$

where the asymptotic variance of  $\hat{\tau}$  is given by

$$\sigma_{\tau}^2(C) = \operatorname{Var}(g(U, \tilde{U})g(V, \tilde{V})), \quad C \in C_2.$$

For (U, V),  $(\tilde{U}, \tilde{V}) \stackrel{\text{id}}{\sim} C$ , write  $X = g(U, \tilde{U})$  and  $Y = g(V, \tilde{V})$ . As discussed in Section 3.3, location shift of X and Y does not change  $\tau(C) = \rho(X, Y)$  but may affect  $\sigma_{\tau}^2(C)$ . Thus we define the optimal location shift as follows.

**Definition 11** (Optimal shift of *g*). For  $C \in C_2$ , the optimal (location) shift of *g* under *C* is given by

$$\mu_* = \mu_*(\tau, C) = \begin{cases} -\frac{\text{Cov}(XY, X+Y)}{\text{Var}(X+Y)}, & \text{if } \text{Var}(X+Y) > 0, \\ 0, & \text{if } \text{Var}(X+Y) = 0. \end{cases}$$

where  $X = g(U, \tilde{U})$  and  $Y = g(V, \tilde{V})$  for (U, V),  $(\tilde{U}, \tilde{V}) \stackrel{\text{iid}}{\sim} C$ .

Basic properties of  $\mu_*$  and  $\sigma_{\tau}^2$  are collected in the next proposition.

**Proposition 12** (Basic properties of  $\mu_* \sigma_{\tau}^2(C)$ ). Let  $C \in C_2$ . Then the following properties hold for  $\tau$ :

- 1.  $\mu_*(\tau, C) = 0;$
- 2.  $\sigma_{\tau}^2(C) = 4p_{\tau}(C)(1 p_{\tau}(C)) = 1 \tau^2(C)$  where  $p_{\tau}(C) = 2 \int_{[0,1]^2} C(u, v) dC(u, v);$
- 3.  $0 \le \sigma_{\tau}^2(C) \le 1;$
- 4.  $\sigma_{\tau}^2(C) = 0$  if and only if  $\tau(C) = 1$  or -1, that is, C = M or W, respectively;
- 5.  $\sigma_{\tau}^2(C) = 1$  if and only if  $\tau(C) = 0$ . In particular,  $\tau(C) = 0$  when  $C = C_{\nu_1}$  or  $C = C_{\nu_2}$ .

**Remark 3** (Asymptotic variance of  $\tau$  for elliptical copulas). When *C* is an elliptical copula with correlation parameter  $\rho \in [-1, 1]$ , we have that  $\tau(C) = 2 \arcsin(\rho)/\pi$ ; see [8]. Therefore, by Proposition 13 Part 2, we have that  $\sigma_{\tau}^2(C) = 1 - (2 \arcsin(\rho)/\pi)^2$ , which also equals  $\sigma_{\beta}^2(C)$  as derived in Remark 2.

Similar to the case of G-transformed rank correlations, we consider the following best and worst asymptotic variances.

**Definition 12** (Best and worst asymptotic variances for  $\tau$ ). For  $\mathcal{D} \subseteq C_2$ , the best and worst asymptotic variances are defined by

$$\underline{\sigma}_{\tau}^{2}(\mathcal{D}) = \inf_{C \in \mathcal{D}} \sigma_{\tau}^{2}(C), \quad \overline{\sigma}_{\tau}^{2}(\mathcal{D}) = \sup_{C \in \mathcal{D}} \sigma_{\tau}^{2}(C),$$

respectively. If the infimum and supremum above are attainable, then their attaining copulas on  $\mathcal{D}$  are defined, respectively, by

$$\underline{C}_{\tau}(\mathcal{D}) = \operatorname*{argmin}_{C \in \mathcal{D}} \sigma^2_{\tau}(C) \quad \overline{C}_{\tau}(\mathcal{D}) = \operatorname*{argmax}_{C \in \mathcal{D}} \sigma^2_{\tau}(C)$$

Properties of the best and worst asymptotic variances of  $\tau$  are summarized as follows.

**Proposition 13** (Best and worst asymptotic variances of Kendall's tau). Let  $\mathcal{D} \subseteq C_2$  and  $\beta \in \mathcal{H} \subseteq \mathcal{G}_4$ .

1. If  $M \in \mathcal{D}$  or  $W \in \mathcal{D}$ , then  $\underline{\sigma}^2_{\tau}(\mathcal{D}) = \underline{\sigma}^2_*(\mathcal{H}, \mathcal{D}) = 0$  and  $\underline{C}_{\tau}(\mathcal{D}) = \{M, W\} \cap \mathcal{D}$ .

2. If  $\Pi \in \mathcal{D}$ , then  $\overline{\sigma}_{\tau}^{2}(\mathcal{D}) = \overline{\sigma}_{*}^{2}(\mathcal{H}, \mathcal{D}) = 1$  and  $\Pi \in \overline{C}_{\tau}(\mathcal{D})$ . 3. If  $\mathcal{D} = C^{F}$ , then  $\overline{\sigma}_{\tau}^{2}(\mathcal{D}) = 1$  and  $\overline{C}_{\tau}(\mathcal{D}) = \{p(M + W)/2 + (1 - p)\Pi : p \in [0, 1]\}$ .

Proposition 13 states that Kendall's tau attains the optimal best and worst asymptotic variances of transformed rank correlations, which are also attained by Blomqvist's beta. Taking into account the drawback of Blomqvist's beta that it depends only on the local value C(1/2, 1/2) of a copula C, Kendall's tau can be a good alternative of Blomqvist's beta in terms of best and worst asymptotic variances.

**Remark 4** (Comparability of  $\kappa_G$  and  $\tau$ ). Since Representation (11) of Kendall's tau in terms of Pearson's correlation coefficient depends on two independent copies  $(U, V) \sim C$  and  $(\tilde{U}, \tilde{V}) \sim C$ , the canonical estimator  $\hat{\tau}$  requires twice more samples from *C* than  $\hat{\kappa}$  does. Therefore, if the estimators  $\hat{\tau}$  and  $\hat{\kappa}_G$  are compared based on their actual variances (instead of their asymptotic variances), then  $\operatorname{Var}(\hat{\tau}) = \sigma_{\tau}^2(C)/n$  should be multiplied by 2 to be compared with  $\operatorname{Var}(\hat{\kappa}_G) = \sigma_G^2(C)/n$ . With this modification, Kendall's tau still attains the optimal best asymptotic variance  $\sigma_{\pi}^2(\mathcal{G}_4, \mathcal{D}) = 0$  since  $2\sigma_{\tau}^2(\mathcal{D}) = 0$ . On the other hand, Kendall's tau fails to attain the optimal worst asymptotic variance  $\overline{\sigma}_{\pi}^2(\mathcal{G}_4, \mathcal{D}) = 1$  since  $2\overline{\sigma}_{\tau}^2(\mathcal{D}) = 2 > 1$ .

# 6. Simulation study

In this section, we conduct a simulation study to compare the asymptotic variance  $\sigma_G^2(C)$  for various copulas  $C \in C_2$  and concordance-inducing distributions  $G \in \mathcal{G}_4$ . Not only Spearman's rho  $\rho_S$ , Blomqvist's beta  $\beta$  and van der Waerden's coefficient  $\zeta$ , we also consider *G*-transformed rank correlations with *G* given by a Student *t* distribution t(v) with v = 10 degrees of freedom and a beta distribution with shape parameters (0.5, 0.5). Note that both are radially symmetric and have finite fourth moments, and thus belong to  $\mathcal{G}_4$  after standardization (mean zero and variance one). The Beta(0.5, 0.5) concordance-inducing distribution has a different shape from the others since it puts an increasing probability mass as locations farther away from the center 1/2. Kendall's tau is also considered for comparison. Besides standardized concordance-inducing distributions, we also consider optimally shifted ones as introduced in Section 3.3. As underlying copulas, we consider Gaussian  $C_{\rho}^{Ga}$ , Student  $t C_{\rho,\nu}^t$  and Clayton copulas  $C_{\theta}^{Cl}$  where  $\rho \in [-1, 1]$  is a correlation parameter,  $\nu > 0$  is a degree of freedom and  $\theta \ge -1$  is a shape parameter. The experiment consists of the following three steps.

- 1. Set  $\rho = -0.99 + 1.98k/49$  for  $k \in \{0, 1, ..., 49\}$ ,  $\nu = 5$  and  $\theta = 2\rho/(1-\rho)$  (which yields  $\tau(C_{\theta}^{\text{Cl}}) = \rho$ ) in  $C = C_{\rho}^{\text{Ga}}$ ,  $C_{\rho,\nu}^{t}$  and  $C_{\theta}^{\text{Cl}}$ .
- 2. For each copula C in Step 1, simulate  $(U_1, V_1), \ldots, (U_n, V_n) \stackrel{\text{iid}}{\sim} C$  with  $n = 10^5$ .
- 3. Based on the samples generated in Step 2, estimate  $\sigma_G^2(C)$  and  $\sigma_\tau^2(C)$  by the sample variances of  $G^{-1}(U_i)G^{-1}(V_i)$ , i = 1, ..., n, and of  $g(U_i, U_{i+n/2})g(V_i, V_{i+n/2})$ , i = 1, ..., n/2, where *G* is a standardized, and optimally shifted uniform, Beta(0.5, 0.5), normal, t(10) and symmetric Bernoulli distribution function, and *g* is as defined in (11).

The estimates of  $\sigma_G^2(C)$  and  $\sigma_\tau^2(C)$  computed in Step 3 are plotted in Fig. 3. Observations from these plots are summarized as follows. For  $G \in \mathcal{G}$ , we denote by V(G) (or  $V(\kappa_G)$ ) the variance of  $X^2$  where X follows the standarzied distribution of G so that  $\mathbb{E}[X] = 0$  and  $\operatorname{Var}(X) = 1$ . In addition, we write  $V(\tau) = \operatorname{Var}(g(U, \tilde{U}))$  for  $U, \tilde{U} \stackrel{\text{id}}{\sim} \operatorname{Unif}(0, 1)$ .

Symmetry: For all copulas *C*, the curves of  $\sigma_G^2(C)$  and  $\sigma_\tau^2(C)$  against the correlation parameter  $\rho$  were almost symmetric around  $\rho = 0$ . For  $C = C_{\rho}^{\text{Ga}}$  and  $C_{\rho,\nu}^t$ , the symmetry is a consequence from Proposition 2 Part 5 since  $C_{-\rho}^{\text{Ga}} = v_1(C_{\rho}^{\text{Ga}})$  and  $C_{-\rho,\nu}^t = v_1(C_{\rho,\nu}^t)$ . This argument does not apply to Clayton copulas, and thus the curves  $\rho \mapsto \sigma_G^2(C_{2\rho/(1-\rho)}^{\text{Cl}})$  and  $\rho \mapsto \sigma_\tau^2(C_{2\rho/(1-\rho)}^{\text{Cl}})$  are not precisely symmetric.

*Convexity and concavity*: At least in this experiment, the curves of  $\sigma_G^2(C)$  and  $\sigma_\tau^2(C)$  are convex when V(G) > 1 (which holds if *G* is normal or t(10)), and concave when V(G) < 1 (which holds if *G* is Bernoulli, uniform and Beta(0.5, 0.5), and if Kendall's tau is considered). This observation is consistent with (6), and with the asymptotic variances obtained in Remarks 2 and 3.

Best and worst asymptotic variances: For all cases of  $C = C_{\rho}^{\text{Ga}}$ ,  $C_{\rho,\nu}^{t}$  and  $C_{\theta}^{\text{Cl}}$ , the best and worst asymptotic variances were approximately  $1 \wedge V(G)$  and  $1 \vee V(G)$ , respectively. When V(G) > 1 (normal or t(10)), the lower bound  $\sigma_{G}^{2}(C) = 1$  was attained at  $\rho = 0$  and the upper bound  $\sigma_{G}^{2}(C) = V(G)$  was attained at  $\rho = \pm 1$ . When V(G) < 1 (Bernoulli, uniform, Beta(0.5, 0.5) and Kendall), the lower bound  $\sigma_{G}^{2}(C) = V(G)$  was attained at  $\rho = \pm 1$  and the upper

bound  $\sigma_G^2(C) = 1$  was attained at  $\rho = 0$ . Note that, for  $C = C_{\rho,\nu}^t$ , the curve was slightly above these lower and upper bounds at  $\rho = 0$  since  $C_{0,\nu}^t \neq \Pi$ . Since we only consider specific classes of copulas, the global upper bound 1 + V(G)as presented in Proposition 2 Part 3 was not attained except in the cases of Blomqvist's beta and Kendall's tau where  $V(\beta) = V(\tau) = 0$ .

*Choice of G; normal or Student, and uniform or Beta*: As seen for the best and worst asymptotic variances, the variance V(G) may play important roles in determining the best and worst asymptotic variances. As theoretically indicated, concordance-inducing distributions with smaller V(G) are more preferable at least in terms of asymptotic variance. Therefore, the normal concordance-inducing distribution is more preferable to t(10) since  $V(\zeta) = 2 < 3 = V(t(10))$ . In fact, for all copulas considered,  $\zeta$  had a smaller asymptotic variance than  $\kappa_{t(10)}$ . Interestingly,  $\kappa_{\text{Beta}(0.5,0.5)}$  typically had smaller asymptotic variance than  $\rho_{\text{S}}$  as expected from  $V(\text{Beta}) = 0.5 < 0.8 = V(\rho_{\text{S}})$ .

Blomqvist's beta and Kendall's tau: As indicated in Remark 3, the curves of  $\sigma_{\beta}^2(C)$  and  $\sigma_{\tau}^2(C)$  seem to coincide when  $C = C_{\rho}^{\text{Ga}}$  and  $C_{\rho,\nu}^t$ . On the other hand,  $\sigma_{\beta}^2(C_{\theta}^{\text{Cl}})$  and  $\sigma_{\tau}^2(C_{\theta}^{\text{Cl}})$  are in general different since  $\sigma_{\beta}^2(C_{\theta}^{\text{Cl}}) = 1 - \beta^2(C_{\theta}^{\text{Cl}})$  and  $\sigma_{\tau}^2(C_{\theta}^{\text{Cl}}) = 1 - \tau^2(C_{\theta}^{\text{Cl}}) = 1 - \tau^2(C_{\theta}^{\text{Cl}})$  by Proposition 8 Part 2 and Proposition 12 Part 2, but  $\beta(C_{\theta}^{\text{Cl}}) = 4(2^{\theta+1} - 1)^{-1/\theta} - 1$  and  $\tau(C_{\theta}^{\text{Cl}}) = \theta/(\theta + 2)$ .

Strength and the model of dependence: For any concordance-inducing distribution, the difference of  $\sigma_G^2(C)$  among different copulas  $C = C_{\rho}^{\text{Ga}}$ ,  $C_{\rho,\nu}^t$  and  $C_{\theta}^{\text{Cl}}$  was typically smaller than the difference of  $\sigma_G^2(C)$  among different levels of dependence, which is controlled by  $\rho$  in this experiment. Therefore, one might say that the choice of C is less influential on the asymptotic variance  $\sigma_G^2(C)$  compared with the choice of concordance-inducing distribution and the strength of dependence.

*Effect of optimal shifts*: As theoretically indicated in Proposition 4, Proposition 8 Part 1 and Proposition 12 Part 1, the solid and dotted curves of asymptotic variances overlapped when  $C = C_{\rho}^{\text{Ga}}$  or  $C_{\rho,\nu}^{t}$ , and when  $\beta$  and  $\tau$  are considered. For other cases, the optimal shift reduced the asymptotic variance. However, even when the copula is  $C_{\theta}^{\text{Cl}}$ , only a small reduction by the optimal shift was observed in this experiment.

# 7. Concluding remarks and discussion

To answer the question which measure of concordance is best to use, we proposed a comparison of G-transformed rank correlations  $\kappa_G$  in terms of their best and worst asymptotic variances  $\sigma_G^2(C)$  on a set of copulas  $\mathcal{D}$ . When  $\mathcal{D}$  is a set of fundamental copulas or Fréchet copulas, we showed that the best and worst asymptotic variances are increasing functions of  $\operatorname{Var}_G(X^2)$ ,  $X \sim G$ , which leads to the comparison criterion that concordance-inducing distributions G having smaller  $\operatorname{Var}_G(X^2)$  are more preferable. Since  $\operatorname{Var}_G(X^2)$  attains its minimum  $\operatorname{Var}_G(X^2) = 0$  if and only if G is a standardized symmetric Bernoulli distribution, we proved that Blomqvist's beta  $\beta$  uniquely attains the optimal best and worst asymptotic variances among all transformed rank correlations under certain conditions on  $\mathcal{D}$ . Considering the drawback of  $\beta$  that it depends only on the local value C(1/2, 1/2) of a copula C, we also compared G-transformed rank correlations with Kendall's tau  $\tau$ . Based on the representation of  $\tau$  in terms of Pearson's linear correlation coefficient, we found that  $\tau$  also attains the optimal best and worst asymptotic variances that  $\beta$  does, although  $\tau$  is not a transformed rank correlation. Since the estimator of  $\tau$  requires twice more samples than that of  $\kappa_G$  does, some optimality of  $\tau$  is violated if this required sample size is taken into account. In a simulation study, we investigated the curve of the asymptotic variance of  $\kappa_G$  and  $\tau$  against the strength of dependence of the underlying copula. We observed that the curve of  $\sigma_G^2(C)$  was typically symmetric and convex or concave with the best and worst asymptotic variances give by  $1 \wedge \operatorname{Var}_G(X^2)$  and  $1 \vee \operatorname{Var}_G(X^2)$  depending on  $\operatorname{Var}_G(X^2) > 1$  or  $\operatorname{Var}_G(X^2) < 1$ . These observations support the criterion that concordance-inducing distributions G with smaller  $Var_G(X^2)$  are more preferable. Consequently, heavytailed concordance-inducing distributions, such as Student t distributions with small degrees of freedom, are not recommended in comparison to the normal distribution, which leads to van der Waerden's coefficient  $\zeta$ . In addition, we found that the beta distribution-based transformed rank correlations can be good alternatives to Spearman's rho  $\rho_{\rm S}$ .

Below we list limitations, discussion and future research on this work.

• Although Blomqvist's beta  $\beta$  can be optimal in terms of stability of its statistical estimation, this measure has some limitations. For example,  $|\beta(C)| = 1$  does not imply C = M, W; see [9, Proposition 1].

- Given the limitations of Fréchet copulas in practice, it may be interesting to explore optimal concordanceinducing distributions under more practical choices of sets of the underlying copulas  $\mathcal{D}$ , such as a set of parametric copulas or a ball of copulas around a given reference copula.
- In our work we assumed for simplicity that i.i.d. samples from the underlying copula are available. However, this may be unrealistic in practice, and it is therefore of interest whether and how results in this paper change if pseudo-samples from the underlying copula are used in the comparison of measures of concordance in terms of their asymptotic variances.
- Besides Kendall's tau, there are still important measures of concordance, such as Gini's gamma, which are not transformed rank correlations. It is thus of interest to study a broader framework that allows one to include such measures of concordance in comparison. In particular, for a comparison, it may be better to consider other estimators of Kendall's tau than the canonical one considered in this paper due to the required sample size.

# 8. Proofs

## **Proof of Proposition 2.**

1) For  $(U, V) \sim C$ , write  $X = G^{-1}(U)$  and  $Y = G^{-1}(V)$ . Since  $G \in \mathcal{G}_4$ , we have that

$$\operatorname{Cov}(X^2, Y^2) = \mathbb{E}[(XY)^2] - \mathbb{E}[X^2]\mathbb{E}[Y^2] = \mathbb{E}[(XY)^2] - 1, \quad \operatorname{Cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y] = \mathbb{E}[XY].$$

Therefore,

$$\sigma_G^2(C) = \operatorname{Var}(XY) = \mathbb{E}[(XY)^2] - \mathbb{E}[XY]^2 = \operatorname{Cov}(X^2, Y^2) + 1 - \operatorname{Cov}(X, Y)^2.$$

2) If  $(U, V) \sim \Pi$ , then  $(X^2, Y^2)$  and (X, Y) are both independent random vectors, and thus  $\operatorname{Cov}(X^2, Y^2) = \operatorname{Cov}(X, Y) = 0$ . Therefore  $\sigma_G^2(\Pi) = 0 + 1 - 1^2 = 1$  by (3). If C = M, then we have that  $\operatorname{Cov}(X, Y) = \kappa_G(M) = 1$  and that  $(X^2, Y^2) \stackrel{d}{=} (G^{-1}(U)^2, G^{-1}(U)^2)$  for  $U \sim \operatorname{Unif}(0, 1)$ . When C = W, we have that  $\operatorname{Cov}(X, Y) = \kappa_G(W) = -1$  and that

$$(X^{2}, Y^{2}) \stackrel{\text{\tiny d}}{=} (G^{-1}(U)^{2}, G^{-1}(1-U)^{2}) = (G^{-1}(U)^{2}, (-G^{-1}(U))^{2}) = (G^{-1}(U)^{2}, G^{-1}(U)^{2}), \quad U \sim \text{Unif}(0, 1),$$

by radial symmetry of  $G \in \mathcal{G}_4$ . Therefore, in either case, we have  $\rho(X^2, Y^2) = 1$  and thus

$$Cov(X^2, Y^2) = \rho(X^2, Y^2) SD_G(X^2) SD_G(Y^2) = Var_G(X^2).$$

Therefore, by (3), we have that

$$\sigma_G^2(M) = \operatorname{Var}_G(X^2) + 1 - 1^2 = \operatorname{Var}_G(X^2), \quad \sigma_G^2(W) = \operatorname{Var}_G(X^2) + 1 - (-1)^2 = \operatorname{Var}_G(X^2).$$

3) By Cauchy–Schwarz inequality, it holds that  $Cov(X^2, Y^2) \le SD(X^2) SD(Y^2) = Var(X^2)$ . Since  $Cov(X, Y)^2 \ge 0$ , the desired inequalities hold by (3). When C = (M + W)/2, we have that

$$(X,Y) \stackrel{d}{=} B(G^{-1}(U), G^{-1}(U)) + (1-B)(G^{-1}(U), G^{-1}(1-U)),$$

where  $B \sim \text{Bern}(1/2)$  and  $U \sim \text{Unif}(0, 1)$  independent of B. Therefore,

$$\operatorname{Cov}(X,Y) = \frac{1}{2}\operatorname{Cov}(G^{-1}(U),G^{-1}(U)) + \frac{1}{2}\operatorname{Cov}(G^{-1}(U),G^{-1}(1-U)) = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0,$$

and  $Cov(X^2, Y^2) = Cov(G^{-1}(U)^2, G^{-1}(U)^2) = Var_G(X^2)$ . Using (3), we have that

$$\sigma_G ((M + W)/2) = \operatorname{Var}_G(X^2) + 1 - 0^2 = \operatorname{Var}_G(X^2) + 1.$$

4) Since the fourth moment of *G* is finite, we have that  $\operatorname{Var}_G(X^2) < \infty$ . Therefore  $\sigma_G^2(C) < \infty$  by 3).

5) The desired result follows from (3) since radial symmetry of  $G \in \mathcal{G}$  implies that

$$(G^{-1}(U), G^{-1}(V)) \stackrel{\text{a.s.}}{=} (-G^{-1}(U_{\nu_1}), G^{-1}(V_{\nu_1})) \stackrel{\text{a.s.}}{=} (G^{-1}(U_{\nu_2}), -G^{-1}(V_{\nu_2})) \stackrel{\text{a.s.}}{=} (-G^{-1}(U_{\nu_1 \circ \nu_2}), -G^{-1}(V_{\nu_1 \circ \nu_2})).$$
(12)

**Proof of Proposition 3.** For  $C, C' \in C_2$  and  $p \in [0, 1]$ , define the random vector  $(\tilde{X}, \tilde{Y}) = (G^{-1}(\tilde{U}), G^{-1}(\tilde{V}))$  where  $(\tilde{U}, \tilde{V}) = B(U, V) + (1 - B)(U', V'), (U, V) \sim C, (U', V') \sim C'$  and  $B \sim \text{Bern}(p)$  is independent of (U, V) and (U', V'). Then  $(\tilde{U}, \tilde{V}) \sim \tilde{C}_p$  where  $\tilde{C}_p = pC + (1 - p)C'$ . Moreover, we have that

$$(\tilde{X}, \tilde{Y}) = (G^{-1}(\tilde{U}), G^{-1}(\tilde{V})) = (G^{-1}(BU + (1 - B)U'), G^{-1}(BV + (1 - B)V')) = B(X, Y) + (1 - B)(X', Y'),$$

where  $(X, Y) = (G^{-1}(U), G^{-1}(V))$  and  $(X', Y') = (G^{-1}(U'), G^{-1}(V'))$ . From this representation, it holds that

$$Cov(\tilde{X}^{2}, \tilde{Y}^{2}) = p Cov(X^{2}, Y^{2}) + (1 - p) Cov(X'^{2}, Y'^{2}), \quad Cov(\tilde{X}, \tilde{Y}) = p Cov(X, Y) + (1 - p) Cov(X', Y').$$

Therefore, we have that

$$\begin{split} \sigma_G^2(\tilde{C}_p) &= \operatorname{Var}(\tilde{X}\tilde{Y}) = \operatorname{Cov}(\tilde{X}^2, \tilde{Y}^2) + 1 - \operatorname{Cov}(\tilde{X}, \tilde{Y})^2 \\ &= p \operatorname{Cov}(X^2, Y^2) + (1-p) \operatorname{Cov}(X'^2, Y'^2) + 1 - (p \operatorname{Cov}(X, Y) + (1-p) \operatorname{Cov}(X', Y'))^2 \\ &\geq p \operatorname{Cov}(X^2, Y^2) + (1-p) \operatorname{Cov}(X'^2, Y'^2) + 1 - p \operatorname{Cov}(X, Y)^2 - (1-p) \operatorname{Cov}(X', Y')^2 \\ &= p \operatorname{Var}(XY) + (1-p) \operatorname{Var}(X'Y') = p \sigma_G^2(C) + (1-p) \sigma_G^2(C'), \end{split}$$

where the inequality in the third line holds since

$$p \operatorname{Cov}(X, Y)^{2} + (1 - p) \operatorname{Cov}(X', Y')^{2} - (p \operatorname{Cov}(X, Y) + (1 - p) \operatorname{Cov}(X', Y'))^{2}$$
$$= p(1 - p)(\operatorname{Cov}(X, Y) - \operatorname{Cov}(X', Y'))^{2} \ge 0.$$

Therefore, the map  $C \mapsto \sigma_G^2(C)$  is concave. When  $C, C' \in C_G(k)$  for some  $k \in [-1, 1]$ , we have Cov(X, Y) = Cov(X', Y'), and thus equality holds in the inequality above. Consequently, the map  $C \mapsto \sigma_G^2(C)$  is linear.

**Proof of Proposition 4.** By definition of  $\mu_*$ , it suffices to consider the case when  $Var(X_0 + Y_0) > 0$  for  $X_0 = G_0^{-1}(U)$  and  $Y_0 = G_0^{-1}(V)$  with  $(U, V) \sim C$ .

Since  $\mathbb{E}[X_0 + Y_0] = \mathbb{E}[X_0] + \mathbb{E}[Y_0] = 0$ , we have that

$$\operatorname{Cov}(X_0Y_0, X_0 + Y_0) = \mathbb{E}[X_0Y_0(X_0 + Y_0)] - \mathbb{E}[X_0Y_0]\mathbb{E}[X_0 + Y_0] = \mathbb{E}[X_0Y_0(X_0 + Y_0)].$$

Therefore, it suffices to show that  $\mathbb{E}[X_0Y_0(X_0 + Y_0)] = 0$  when *C* is radially symmetric.

When C is radially symmetric, we have that  $(U, V) \stackrel{d}{=} (U_{\nu_1 \circ \nu_2}, V_{\nu_1 \circ \nu_2})$  and  $(U_{\nu_1}, V_{\nu_1}) \stackrel{d}{=} (U_{\nu_2}, V_{\nu_2})$  for  $(U, V) \sim C$ . Together with the identity

$$1 = \mathbf{1}_{\{U>1/2, V>1/2\}} + \mathbf{1}_{\{U\le1/2, V>1/2\}} + \mathbf{1}_{\{U>1/2, V\le1/2\}} + \mathbf{1}_{\{U>1/2, V\le1/2\}} + \mathbf{1}_{\{U_{v_1}>1/2, V_{v_1}>1/2\}} + \mathbf{1}_{\{U_{v_2}>1/2, V_{v_2}>1/2\}} + \mathbf{1}_{\{U_{v_1}\circ v_2>1/2, V_{v_1}\circ v_2>1/2\}},$$

we have, by (12), that

$$\begin{split} \mathbb{E}[X_0 Y_0(X_0 + Y_0)] &= \sum_{\varphi \in \{l, v_1, v_2, v_1 \circ v_2\}} \mathbb{E}[\mathbf{1}_{\{U_{\varphi} > 1/2, V_{\varphi} > 1/2\}} G_0^{-1}(U) G_0^{-1}(V) (G_0^{-1}(U) + G_0^{-1}(V))] \\ &= \mathbb{E}[\mathbf{1}_{\{U > 1/2, V > 1/2\}} G_0^{-1}(U) G_0^{-1}(V) (G_0^{-1}(U) + G_0^{-1}(V))] \\ &\quad - \mathbb{E}[\mathbf{1}_{\{U_{v_1} \circ v_2 > 1/2, V_{v_1} \circ v_2 > 1/2\}} G_0^{-1}(U_{v_1 \circ v_2}) G_0^{-1}(V_{v_1 \circ v_2}) (G_0^{-1}(U_{v_1 \circ v_2}) + G_0^{-1}(V_{v_1 \circ v_2}))] \\ &\quad + \mathbb{E}[\mathbf{1}_{\{U_{v_1} > 1/2, V_{v_1} > 1/2\}} G_0^{-1}(U_{v_1}) G_0^{-1}(V_{v_1}) (G_0^{-1}(U_{v_1}) - G_0^{-1}(V_{v_1}))] \\ &\quad - \mathbb{E}[\mathbf{1}_{\{U_{v_2} > 1/2, V_{v_2} > 1/2\}} G_0^{-1}(U_{v_2}) G_0^{-1}(V_{v_2}) (G_0^{-1}(U_{v_2}) - G_0^{-1}(V_{v_2}))] = 0, \end{split}$$

which gives the desired result  $\mu_*(G, C) = 0$ .

**Proof of Proposition 5.** Since normal variance mixtures are radially symmetric, the optimal asymptotic variance is obtained when  $\mu^* = 0$ . Since  $\sigma_{G_{\mu^*}}^2(C)$  is invariant under scale transforms of  $G_{\mu_*}$ , we standardize (X, Y) to apply

the covariance formula (3). Namely, we set  $\Sigma = P/\mathbb{E}[W]$  where P is a correlation matrix with off-diagonal entry  $\rho = \rho(X, Y)$ . In this setup, we have that

$$\operatorname{Var}(X) = \operatorname{Var}(Y) = \mathbb{E}[WZ_1^2] - \mathbb{E}\left[\sqrt{W}Z_1\right]^2 = \mathbb{E}[W]\operatorname{Var}(Z_1) - 0 = \mathbb{E}[W]\frac{1}{\mathbb{E}[W]} = 1.$$

Since  $\text{Cov}(Z_1^2, Z_2^2) = 2\rho^2 / \mathbb{E}[W]^2$ , the law of total covariance implies that

$$Cov(X^{2}, Y^{2}) = \mathbb{E}[Cov(X^{2}, Y^{2}) | W] + Cov(\mathbb{E}[X^{2} | W], \mathbb{E}[Y^{2} | W])$$
  
=  $\mathbb{E}[W^{2} Cov(Z_{1}^{2}, Z_{2}^{2})] + Cov(\mathbb{E}[Z_{1}^{2}]W, \mathbb{E}[Z_{2}^{2}]W) = \mathbb{E}[W^{2}] Cov(Z_{1}^{2}, Z_{2}^{2}) + Var(W) Var(Z_{1}) Var(Z_{2})$   
=  $2\rho^{2} \frac{\mathbb{E}[W^{2}]}{\mathbb{E}[W]^{2}} + \frac{Var(W)}{\mathbb{E}[W]^{2}}.$ 

Therefore, we have, by (3), that

$$\sigma_{G_{\mu^*}}^2(C) = \operatorname{Var}(XY) = \operatorname{Cov}(X^2, Y^2) + 1 - \operatorname{Cov}(X, Y)^2 = 2\rho^2 \frac{\mathbb{E}[W^2]}{\mathbb{E}[W]^2} + \frac{\operatorname{Var}(W)}{\mathbb{E}[W]^2} + 1 - \rho^2,$$

and thus the desired result follows.

**Proof of Proposition 7.** Fix  $G \in \mathcal{G}_4$  and  $C_p^F \in C^F$  with  $p = (p_M, p_\Pi, p_W) \in \Delta_3$ . For  $X = G^{-1}(U)$  and  $Y = G^{-1}(V)$  with  $(U, V) \sim C_p^F$ , we have that  $Cov(X^2, Y^2) = (p_M + p_W) Var_G(X^2)$  and that  $Cov(X, Y) = p_M - p_W$ . Therefore, by (3), it holds that

$$\sigma_G^2(C_p^{\rm F}) = (p_M + p_W)v + 1 - (p_M - p_W)^2 =: f(p_M, p_W),$$

where  $v = \operatorname{Var}_{G}(X^{2})$  for notational convenience. Since the Hessian of *f* 

$$H(p_M, p_W) = \begin{pmatrix} \frac{\partial}{\partial p_M^2} f(p_M, p_W) & \frac{\partial}{\partial p_M p_W} f(p_M, p_W) \\ \frac{\partial}{\partial p_W p_M} f(p_M, p_W) & \frac{\partial}{\partial p_W^2} f(p_M, p_W) \end{pmatrix} = \begin{pmatrix} -2 & 2 \\ 2 & -2 \end{pmatrix},$$

is nonpositive definite, f is a concave function.

For  $(p_M, p_W) \in \mathbb{R}^2$  such that  $p_M, p_W \ge 0$  and  $p_M + p_W \le 1$ , consider the reparametrization (p, 0) + r(-1, 1) = (p - r, r) where  $0 \le r \le p \le 1$ . Then

$$f(p-r,r) = pv + 1 - (p-r)^2 - r^2 + 2(p-r)r = -4\left(r - \frac{p}{2}\right)^2 + pv + 1,$$

and thus f represents a parabolic cylinder.

For a fixed  $p \in [0, 1]$ , the function  $r \mapsto f(p - r, r)$  has a maximum  $\overline{f}(p) = pv + 1$  when r = p/2, and a minimum  $\underline{f}(p) = -p^2 + pv + 1$  when r = 0 or r = p. Since  $v \ge 0$ , the maximum of f is given by v + 1 with the maximum attained by p = 1 when v > 0, and by any  $p \in [0, 1]$  when v = 0. Therefore, we have that  $\overline{\sigma}_G^2(C^F) = v + 1 = \sigma_G^2(C)$  with C = (M + W)/2 when v > 0, and with  $C = p(M + W)/2 + (1 - p)\Pi$  for any  $p \in [0, 1]$  when v = 0. For the minimum of f, notice that the function  $f(p) = -p^2 + pv + 1$ ,  $0 \le p \le 1$ , is a concave parabola, and thus the minimum of  $\underline{f}(p)$  is attained at p = 0 or p = 1. With  $\underline{f}(0) = 1$  and  $\underline{f}(1) = v$ , the minimum of f and its attainers are given by  $\underline{\sigma}_G^2(C^F) = 1 \land v = \sigma_G^2(C)$  with C = M or W when  $0 \le v < 1$ , with C = M,  $\Pi$  or W when v = 1 and with  $C = \Pi$  when v > 1.

**Proof of Corollary 4.** For  $G \in \mathcal{G}$ , denote by V(G) (or  $V(\kappa_G)$ ) the variance of  $X^2$  where X follows the standardized distribution of G so that  $\mathbb{E}[X] = 0$  and  $\operatorname{Var}(X) = 1$ . Then the concordance-inducing distributions of  $\rho_S$ ,  $\beta$  and  $\zeta$  are the uniform distribution on  $\left(-\sqrt{3}, \sqrt{3}\right)$ , the symmetric Bernoulli distribution on  $\{-1, 1\}$  and the standard normal distribution N(0, 1). Since

$$V(\rho_{\rm S}) = V({\rm Unif}(0,1)) = 0.8$$
,  $V(\beta) = V({\rm Bern}(1/2)) = 0$  and  $V(\zeta) = V({\rm N}(0,1)) = 2$ 

the result follows from Corollary 3 Part 2.

 $\square$ 

## **Proof of Proposition 8.** Let G be the standardized symmetric Bernoulli distribution.

1) For  $(X, Y) = (G^{-1}(U), G^{-1}(V))$  with  $(U, V) \sim C$ , it suffices to consider the case when Var(X + Y) > 0 by definition of  $\mu_*$ . Since  $G^{-1}(u) = 2\mathbf{1}_{\{u > 1/2\}} - 1$ ,  $u \in (0, 1)$ , we have that

$$X + Y = \begin{cases} 2, & \text{if } \{U > 1/2, \ V > 1/2\}, \\ -2, & \text{if } \{U \le 1/2, \ V \le 1/2\}, \\ 0, & \text{if } \{U > 1/2, \ V \le 1/2\} \cup \{U \le 1/2, \ V > 1/2\}, \\ XY = \begin{cases} 1, & \text{if } \{U \le 1/2, \ V \le 1/2\} \cup \{U > 1/2, \ V > 1/2\}, \\ -1, & \text{if } \{U > 1/2, \ V \le 1/2\} \cup \{U \le 1/2, \ V > 1/2\}. \end{cases}$$

Since  $C(1/2, 1/2) = \overline{C}(1/2, 1/2)$ , we have that  $\mathbb{E}[X + Y] = 2\overline{C}(1/2, 1/2) - 2C(1/2, 1/2) = 0$ , and thus

$$\operatorname{Cov}(X+Y,XY) = \mathbb{E}[(X+Y)XY] - \mathbb{E}[X+Y]\mathbb{E}[XY] = 2\bar{C}(1/2,1/2) - 2C(1/2,1/2) - 0 = 0,$$

which implies that  $\mu_*(\beta, C) = 0$ . 2) Using the notation

$$p(C) = \Pr(\{U \le 1/2, V \le 1/2\} \cup \{U > 1/2, V > 1/2\}) = C(1/2, 1/2) + \overline{C}(1/2, 1/2) = 2C(1/2, 1/2),$$

we have that  $\beta(C) = 4C(1/2, 1/2) - 1 = 2p(C) - 1$  and that

$$\sigma_{\beta}^{2}(C) = \operatorname{Var}(XY) = 4p(C)(1 - p(C)) = 1 - \beta^{2}(C).$$

3) This immediately follows from 2) since  $0 \le p(C) \le 1$ .

4) By 2),  $\sigma_{\beta}^2(C) = 0$  if and only if p(C) = 0 or 1, that is, *C* is a TPI or TNI copula. 5) By 2),  $\sigma_{\beta}^2(C) = 1$  if and only if p(C) = 1/2, that is, *C* is a balanced copula.

$$\underline{\sigma}_{\beta}^{2}(\mathcal{D}) = \sigma_{\beta}^{2}(C_{*}) = 0 \le \underline{\sigma}_{G}^{2}(\mathcal{D}) \quad \text{for all } G \in \mathcal{G}_{4},$$
(13)

and thus  $\sigma^2_*(\mathcal{H}, \mathcal{D}) = 0$  and  $\beta \in G_*(\mathcal{H}, \mathcal{D})$ . 2) By Proposition 2 Part 2 and Proposition 8 Parts 3 and 5, we have that

$$\overline{\sigma}_{\beta}^{2}(\mathcal{D}) = \sigma_{\beta}^{2}(\Pi) = 1 = \sigma_{G}^{2}(\Pi) \le \overline{\sigma}_{G}^{2}(\mathcal{D}) \quad \text{for all } G \in \mathcal{G}_{4}, \tag{14}$$

and thus  $\overline{\sigma}^2_*(\mathcal{H}, \mathcal{D}) = 1$  and  $\beta \in \overline{G}_*(\mathcal{H}, \mathcal{D})$ .

3)  $\beta \in G_*(\mathcal{H}, \mathcal{D})$  immediately follows from 1) and 2).

Finally, all the relations in (13) and (14) remain valid even if  $\sigma_G^2(C)$ ,  $G \in \mathcal{G}_4$ , is replaced by the optimally shifted asymptotic variance  $\sigma_{\mathcal{G}_{\mu_*}}^2(C)$ . Therefore, all the optimality results in 1), 2) and 3) hold for correspondingly modified versions of Definitions 5 and 6.

**Proof of Proposition 10.** By Proposition 9, we have that  $\beta \in G_*(\mathcal{H}, \mathcal{D})$ . Proposition 2 Part 3 also yields  $\overline{\sigma}_G^2(\mathcal{D}) =$ 1 + Var<sub>G</sub>(X<sup>2</sup>) for all  $G \in \mathcal{H}$ . This upper bound remains valid if  $G \in \mathcal{H}$  is optimally shifted since (M + W)/2is radially symmetric. Therefore, regardless of whether the optimal shift is taken into account, the optimal worst asymptotic variance  $\overline{\sigma}^2_*(\mathcal{H}, \mathcal{D}) = 1$  is attained if and only if  $G \in \mathcal{H}$  satisfies  $\operatorname{Var}_G(X^2) = 0$ , that is,  $X \sim G$  is the standardized symmetric Bernoulli distribution. Consequently, we have that  $\overline{G}_*(\mathcal{H}, \mathcal{D}) = \{\beta\}$  and thus  $G_*(\mathcal{H}, \mathcal{D}) = \{\beta\}$ as desired.  **Proof of Proposition 11.** For  $G \in \mathcal{H}$  and  $C \in \mathcal{D}$  in (8), write  $(X, Y) = (G^{-1}(U), G^{-1}(V))$ .

1) In this case, we have that Pr(XY = 0) > 0 and thus  $a \in \mathbb{R}$  in (8) necessarily has to be a = 0. If  $XY \stackrel{a.s.}{=} 0$  holds, then Y = 0 on  $\{X \neq 0\}$ . Together with  $X \stackrel{d}{=} Y$ , we have that  $Pr(X \neq 0) \le Pr(Y = 0) = Pr(X = 0)$ , which leads to the condition  $Pr(X = 0) \ge 1/2$ .

2) In this case, we have that Pr(XY = 0) = 0 since

$$XY \begin{cases} > 0, & \text{if } \{U \le 1/2, \ V \le 1/2\} \cup \{U > 1/2, \ V > 1/2\}, \\ < 0, & \text{if } \{U \le 1/2, \ V > 1/2\} \cup \{U > 1/2, \ V \le 1/2\}. \end{cases}$$

Therefore,  $a \in \mathbb{R}$  in (8) necessarily has to be unequal to 0. Since

$$Pr(XY > 0) = p(\{U \le 1/2, V \le 1/2\} \cup \{U > 1/2, V > 1/2\}) = p(C)$$

and

$$Pr(XY < 0) = p(\{U \le 1/2, V > 1/2\} \cup \{U > 1/2, V \le 1/2\}) = 1 - p(C)$$

the random variable XY can never be almost surely a constant if 0 < p(C) < 1. Therefore, it holds that p(C) = 0 or 1, and thus C is either TPI or TNI.

Assume that *C* is TPI. Then a > 0 since Pr(XY > 0) = 1. By the TPI assumption of *C*, we have that

$$\begin{split} X_+ &= X \mid \{U > 1/2, V > 1/2\} = X \mid \{U > 1/2\} \sim G_+, \\ Y_+ &= Y \mid \{U > 1/2, V > 1/2\} = Y \mid \{V > 1/2\} \sim G_+, \\ X_- &= X \mid \{U \le 1/2, V \le 1/2\} = X \mid \{U \le 1/2\} \sim G_-, \\ Y_- &= X \mid \{U \le 1/2, V \le 1/2\} = Y \mid \{V \le 1/2\} \sim G_-, \end{split}$$

where

$$G_{+}(x) = \begin{cases} 2G(x) - 1, & \text{if } x > 0, \\ 0, & \text{if } x \le 0, \end{cases} \quad G_{-}(x) = \begin{cases} 1, & \text{if } x > 0, \\ 2G(x), & \text{if } x \le 0. \end{cases}$$

In addition to the equalities  $X_+ \stackrel{d}{=} Y_+$  and  $X_- \stackrel{d}{=} Y_-$ , we have that  $X_+ \stackrel{d}{=} -X_-$  and  $Y_+ \stackrel{d}{=} -Y_-$  since the radial symmetry of *G* and the condition  $\Pr(X = 0) = 0$  lead to

$$Pr(-X_{-} \le x) = Pr(X_{-} \ge -x) = 1 - G_{-}((-x) - )$$

$$= \begin{cases} 1 - 1 = 0, & \text{if } x < 0, \\ 1 - 2G((-x) - ) = 1 - 2(1 - G(x)) = 2G(x) - 1, & \text{if } x \ge 0, \\ = G_{+}(x). \end{cases}$$

Moreover, since  $XY \stackrel{\text{a.s.}}{=} a$ , it holds that

$$X_+Y_+ = XY \mid \{U > 1/2, V > 1/2\} \stackrel{\text{a.s.}}{=} a \text{ and } X_-Y_- = XY \mid \{U \le 1/2, V \le 1/2\} \stackrel{\text{a.s.}}{=} a.$$

Since  $X_+Y_+ \stackrel{\text{a.s.}}{=} a$  and  $Y_+ > 0$  a.s., we have that  $X_+ \stackrel{\text{a.s.}}{=} a/Y_+$ . Therefore, Jensen's inequality implies that

$$\mathbb{E}[X_+] = \mathbb{E}\left[\frac{a}{Y_+}\right] = a\mathbb{E}\left[\frac{1}{Y_+}\right] \ge \frac{a}{\mathbb{E}[Y_+]} = \frac{a}{\mathbb{E}[X_+]},$$

which yields  $\mathbb{E}[X_+] \ge \sqrt{a}$ .

Since  $X_+ \stackrel{d}{=} -X_-$  and  $Var(X) = \mathbb{E}[X^2] = 1$ , we have that

$$1 = \mathbb{E}[X^2] = \Pr\left(U > \frac{1}{2}\right) \mathbb{E}\left[X^2 \mid U > \frac{1}{2}\right] + \Pr\left(U \le \frac{1}{2}\right) \mathbb{E}\left[X^2 \mid U \le \frac{1}{2}\right] = \frac{1}{2}\mathbb{E}[X_+^2] + \frac{1}{2}\mathbb{E}[X_-^2] = \mathbb{E}[X_+^2],$$

and thus  $\mathbb{E}[X_+^2] = 1$ . Using  $X_+^2 \stackrel{\text{a.s.}}{=} (a/Y_+)^2 > 0$  a.s. and Jensen's inequality, we have that

$$1 = \mathbb{E}[X_+^2] = \mathbb{E}\left[\left(\frac{a}{Y_+}\right)^2\right] \ge \frac{a^2}{\mathbb{E}[Y_+^2]} = \frac{a^2}{\mathbb{E}[X_+^2]},$$

which yields  $-1 \le a \le 1$ . Together with a > 0, we have the inequalities  $0 < a \le 1$ . Moreover,  $X_+ \stackrel{as.}{=} a/Y_+$  implies that

$$G_{+}(x) = \Pr(X_{+} \le x) = \Pr\left(\frac{a}{Y_{+}} \le x\right) = 1 - \Pr\left(Y_{+} < \frac{a}{x}\right) = 1 - G_{+}\left(\frac{a}{x}\right), \quad x > 0,$$

which leads to the identity (9). The symmetry  $Pr(Z > a^{1/2}) = Pr(Z < a^{1/2})$  for  $Z \sim G_+$  is obtained as a special case by taking  $x = \sqrt{a} > 0$  in (9).

Next assume that C is TNI. Then a < 0 since Pr(XY < 0) = 1. By the TNI assumption, we have that

$$\begin{split} X_+ &= X \mid \{U > 1/2, V \le 1/2\} = X \mid \{U > 1/2\} \sim G_+, \\ Y_+ &= Y \mid \{U \le 1/2, V > 1/2\} = Y \mid \{V > 1/2\} \sim G_+, \\ X_- &= X \mid \{U \le 1/2, V > 1/2\} = X \mid \{U \le 1/2\} \sim G_-, \\ Y_- &= X \mid \{U > 1/2, V \le 1/2\} = Y \mid \{V \le 1/2\} \sim G_-. \end{split}$$

As in the TPI case, it holds that  $X_+ \stackrel{d}{=} Y_+, X_- \stackrel{d}{=} Y_-, X_+ \stackrel{d}{=} -X_+$  and  $Y_+ \stackrel{d}{=} -Y_+$ . Moreover,  $XY \stackrel{as}{=} a$  implies that

$$X_+Y_- = XY \mid \{U > 1/2, V \le 1/2\} \stackrel{\text{as.}}{=} a, \quad X_-Y_+ = XY \mid \{U \le 1/2, V > 1/2\} \stackrel{\text{as.}}{=} a.$$

From these equalities, all the necessary conditions derived in the TPI case hold with  $-Y_-$ ,  $-X_- \sim G_+$  and -a > 0since  $X_+(-Y_-) \stackrel{a.s.}{=} (-X_-)Y_+ \stackrel{a.s.}{=} -a$ .

**Proof of Proposition 12.** For (U, V),  $(\tilde{U}, \tilde{V}) \stackrel{\text{id}}{\sim} C$ , write  $X = g(U, \tilde{U})$  and  $Y = g(V, \tilde{V})$ . 1) The statement holds when Var(X + Y) = 0. Assume that Var(X + Y) > 0. Then we have that

$$\begin{aligned} X+Y &= \begin{cases} 2, & \text{if } \{U > \tilde{U}, \ V > \tilde{V}\}, \\ -2, & \text{if } \{U \leq \tilde{U}, \ V \leq \tilde{V}\}, \\ 0, & \text{if } \{U > \tilde{U}, \ V \leq \tilde{V}\} \cup \{U \leq \tilde{U}, \ V > \tilde{V}\}, \end{cases} \\ XY &= \begin{cases} 1, & \text{if } \{U \leq \tilde{U}, \ V \leq \tilde{V}\} \cup \{U > \tilde{U}, \ V > \tilde{V}\}, \\ -1, & \text{if } \{U > \tilde{U}, \ V \leq \tilde{V}\} \cup \{U \leq \tilde{U}, \ V > \tilde{V}\}. \end{cases} \end{aligned}$$

Then  $\mathbb{E}[X + Y] = 0$  and Cov(X + Y, XY) = 0 by calculation since

$$\Pr(U > \tilde{U}, V > \tilde{V}) = \int_{[0,1]^2} C(u, v) \, \mathrm{d}C(u, v).$$

Therefore, we have that  $\mu_*(\tau, C) = 0$  as desired. 2) By definition of Kendall's tau (10), we have that

$$\Pr(\{U \le \tilde{U}, \ V \le \tilde{V}\} \cup \{U > \tilde{U}, \ V > \tilde{V}\}) = 2 \int_{[0,1]^2} C(u,v) \, \mathrm{d}C(u,v) = p_\tau(C) = \frac{\tau(C) + 1}{2},$$

and thus  $\sigma_{\tau}^{2}(C) = \operatorname{Var}(XY) = 4p_{\tau}(C)(1 - p_{\tau}(C)) = 1 - \tau^{2}(C).$ 

3) This is an immediate consequence from 2) and  $0 \le p_{\tau}(C) \le 1$ .

4)  $\sigma_{\tau}^2(C) = 0$  is attained if and only if  $p_{\tau}(C) = 1$  or 0, that is,  $\tau(C) = 1$  or -1, respectively. By [4, Theorem 3],  $\tau(C) = 1$  or -1 if and only if C = M or W, respectively.

5)  $\sigma_{\tau}^2(C) = 1$  is attained if and only if  $p_{\tau}(C) = 1/2$ , that is,  $\tau(C) = 0$ . When  $C = C_{\nu_1}$  or  $C = C_{\nu_2}$ , the change of sign axiom of measures of concordance in Definition 1 implies that  $\tau(C) = \tau(C_{\nu_1}) = -\tau(C)$  or  $\tau(C) = \tau(C_{\nu_2}) = -\tau(C)$ , either of which yields  $\tau(C) = 0$ .


**Proof of Proposition 13.** 1) and 2) immediately follow from Proposition 12 Parts 3, 4 and 5, and Proposition 9. 3) By Proposition 12 Part 5, it holds that  $\sigma_{\tau}^2(C) = 1$  for  $C \in C_2$  if and only if  $\tau(C) = 0$ . For a Fréchet copula, we have that  $\tau(C_{(p_M,p_\Pi,p_W)}^F) = (p_M - p_W)(p_M + p_W + 2)/3$ ; see [11, Example 5.3]. Therefore,  $\tau(C^F) = 0$  holds if and only if  $p_M = p_W$ , and thus the desired result follows.

## Acknowledgments

We are grateful to Alexander Schied and Ruodu Wang at University of Waterloo for their valuable comments. Takaaki Koike was supported by JSPS KAKENHI Grant Number JP21K13275. Marius Hofert acknowledges financial support from the Natural Sciences and Engineering Research Council of Canada (RGPIN-2020-04897 and RGPAS-2020-00093).

#### References

- S. Abdullah, S. Siddiqua, M. S. H. Siddiquee, N. Hossain, Modeling and forecasting exchange rate volatility in Bangladesh using GARCH models: a comparison based on normal and student's t-error distribution, Financial Innovation 3 (2017) 1–19.
- [2] N. Blomqvist, On a measure of dependence between two random variables, The Annals of Mathematical Statistics (1950) 593-600.
- [3] J. C. De Winter, S. D. Gosling, J. Potter, Comparing the Pearson and Spearman correlation coefficients across distributions and sample sizes: A tutorial using simulations and empirical data., Psychological Methods 21 (2016) 273.
- [4] P. Embrechts, A. McNeil, D. Straumann, Correlation and dependency in risk management: Properties and pitfalls, in: M. Dempster (Ed.), Risk Management: Value at Risk and Beyond, Cambridge University Press, 2002, pp. 176–223.
- [5] M. Epaphra, Modeling exchange rate volatility: Application of the GARCH and EGARCH models, Journal of Mathematical Finance 7 (2016) 121–143.
- [6] C. Genest, F. Verret, Locally most powerful rank tests of independence for copula models, Nonparametric Statistics 17 (2005) 521–539.
- [7] M. Hofert, T. Koike, Compatibility and attainability of matrices of correlation-based measures of concordance, ASTIN Bulletin: The Journal of the IAA 49 (2019) 885–918.
- [8] H. Hult, F. Lindskog, Multivariate extremes, aggregation and dependence in elliptical distributions, Advances in Applied probability (2002) 587–608.
- [9] T. Koike, M. Hofert, Matrix compatibility and correlation mixture representation of generalized Gini's gamma, Canadian Journal of Statistics, doi: 10.1002/cjs.11748 (2022).
- [10] P. Luigi Conti, Y. Nikitin, Asymptotic efficiency of independence tests based on Gini's rank association coefficient, Spearman's footrule and their generalizations, Communications in Statistics-Theory and Methods 28 (1999) 453–465.
- [11] R. B. Nelsen, An introduction to Copulas, Springer, New York, 2006.
- [12] J. Raymaekers, P. J. Rousseeuw, Fast robust correlation for high-dimensional data, Technometrics 63 (2021) 184–198.
- [13] E. Rödel, W. Kössler, Linear rank tests for independence in bivariate distributions-power comparisons by simulation, Computational Statistics & Data Analysis 46 (2004) 645–660.
- [14] M. Scarsini, On measures of concordance, Stochastica 8 (1984) 201-218.
- [15] F. Schmid, R. Schmidt, Nonparametric inference on multivariate versions of Blomqvist's beta and related measures of tail dependence, Metrika 66 (2007) 323–354.
- [16] Z. Sidak, P. K. Sen, J. Hajek, Theory of Rank Tests, Elsevier, Amsterdam, 1999.
- [17] C. Spearman, "general intelligence," objectively determined and measured, The American Journal of Psychology 15 (1904) 201–292.
- [18] A. Tarsitano, et al., Comparing the effectiveness of rank correlation statistics, P: Dip. di Economia e Statistica, University of della Calabria (2009).
- [19] J. Yang, S. Cheng, L. Zhang, Bivariate copula decomposition in terms of comonotonicity, countermonotonicity and independence, Insurance: Mathematics and Economics 39 (2006) 267–284.



**Fig. 3:** Estimates of asymptotic variances  $\sigma_G^2(C)$  and  $\sigma_\tau^2(C)$  against correlation parameters  $\rho \in [-0.99, 0.99]$  of  $C = C_{\rho}^{\text{Ga}}$  (red),  $C_{\rho,\nu}^t$  (blue) with  $\nu = 5$  and  $C_{\theta}^{\text{Cl}}$  (green) with  $\theta = 2\rho/(1-\rho)$  for *G*-transformed rank correlation coefficients  $\kappa_G$  (all except bottom-right) and Kendall's tau  $\tau$  (bottom-right). The concordance-inducing distribution *G* is set to be standardized (solid lines) and optimally shifted (dashed lines) uniform, Beta(0.5, 0.5), normal, t(10) and symmetric Bernoulli distribution. The black dotted lines represent y = 1, V(G) and 1 + V(G) with  $V(\beta) = 0$ ,  $V(\rho_S) = 0.8$ ,  $V(\zeta) = 2$ , V(t(10)) = 3, V(Beta(0.5, 0.5)) = 0.5 and  $V(\tau) = 0$ .